

A dissertation on the food and discharges of human bodies / [Bryan Robinson].

Contributors

Robinson, Bryan, 1680-1754.

Publication/Creation

London : J. Nourse, 1748.

Persistent URL

<https://wellcomecollection.org/works/n56qetxb>

License and attribution


This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>





Memorials of Adair

(Museum of Art and Archaeology)

M. Lenihan

DISSERTATION

ON THE

CAUSE OF

DISCHARGES

OF THE

ATMOSPHERE

BY

M. LENIHAN

OF THE

UNIVERSITY OF

YORK

Handwritten text

A
DISSERTATION
ON THE
F O O D
AND
DISCHARGES
OF
HUMAN BODIES.

By BRYAN ROBINSON, M. D.

L O N D O N,
Printed for J. NOURSE, at the Lamb against
Katherine-Street in the *Strand*.

MDCCLXVIII.

D I S S E R T A T I O N

O N T H E

F L O O D

I N S C H A E R S



By _____
of _____

Printed by _____
at _____



P R E F A C E.

IN the following Dissertation I have given an Account, from Experiments, of the Food and Discharges of human Bodies. By the Statical Tables it appears, that in *England, Ireland, and South Carolina*, Perspiration is considerably greater in the Day than in the Night; whereas, by the Aphorisms of *Sanctorius*, the contrary

obtains in *Italy*; for in seven Hours of quiet Sleep a Person there perspires twice as much, as he does when awake in an equal Time. A firm Belief of this Doctrine of *Sanctorius*, and that it obtains in other Countries as *Sanctorius* affirms it does in *Italy*, has hurt many. For it is natural for Persons thus persuaded, when they awake in the Morning, and find any Moisture on their Skin, to be afraid of rising lest they should check Perspiration, and to continue in Bed till the Moisture goes off of itself; by which erroneous Conduct they relax and weaken their Constitutions, and frequently become Valetudinarians, from being Persons of naturally strong and healthful Bodies.

Sanctorius

Sanctorius seems to have had a different Opinion of the Proportion of Perspiration to Urine at different Seasons of the Year, from what appears by our Tables. He seems not to have much regarded Urine in comparison of Perspiration; and yet they are not far from being equal in the whole Year, taking one Day, one Month, and one Season with another, as appears by Tables 2, 7, 9. And, what is very remarkable, Urine in the whole Year exceeds Perspiration, not only in *England* and *Ireland*, but even in *South Carolina*, a Country much hotter than *Italy*.

But tho' *Sanctorius* may have advanced some Things which are

not true, and some which are true only under certain Circumstances, yet he is highly to be honoured for what he has done ; and had he known the Circulation of the Blood, and been acquainted with true Philosophy, we may easily grant, that he would not have left any thing material on this Subject to be done by others.





A
DISERTATION
ON THE
FOOD and DISCHARGES
OF
HUMAN BODIES.

AS the Discharges of human Bodies depend upon, and are regulated by, the Motion of the Blood; so it may be proper to premise a short Account of that Motion, by which the Nature of the Discharges by Perspiration, Urine, and Stool, will be more clearly understood than they could be without it.

Of the Food and Discharges

The Account here given of the Blood's Motion, is drawn from what I have delivered concerning it in the *Animal Oeconomy*, and from other Experiments and Observations. I shall, by way of Illustration, subjoin some Tables and Observations drawn from Experiments made on other Animals.

Of the Motion of the Blood.

IF a healthful Body be situated in a given manner with respect to the Horizon, the Velocity with which the Blood flows out of the left Ventricle of the Heart into the Aorta, is in the subduplicate Ratio of the Diameter of the Aorta ; and if the Body be perfectly well proportioned, and its Heart be free from the Influences of all disturbing Causes, the Velocity with which the Blood flows out of the Heart into the Aorta, is in the subquadruplicate Ratio of the Length of the Body ; by *Cor. 4. Prop. 12. Anim. Oecon.* And the Morning Number of Pulses in a Minute of a healthful
Body,

of Human Bodies.

3

Body, when it is fitting, and its Heart is free from the Influences of all disturbing Causes, is as the Velocity of the Blood in the Aorta apply'd to the Length of the Body, by *Cor. 1. Prop. 14*; and if the Body be perfectly well proportioned, its Morning Number of Pulses in a Minute is in the subquadruplicate Ratio of the Length of the Body, apply'd to the Length of the Body. To express these Proportions in Symbols, let *D* denote the Diameter of the Aorta in Inches and decimal Parts, when the Aorta is moderately distended by the Systole of the Heart; *L* the Length of the Body in Inches; *P* the Morning Number of Pulses in a Minute, when the Body is fitting, and its Heart is free from the Influences of all disturbing Causes; and *V* the Velocity with which the Blood flows out of the left Ventricle of the Heart into the Aorta, expressed in the Inches it describes in a second Minute of Time; and then *V* will be as $D^{\frac{1}{2}}$, and *P* as $\frac{D^{\frac{1}{2}}}{L}$ in all healthful Bodies in the Morning, when the Bodies are fitting,

4 and

Of the Food and Discharges

and their Hearts are free from the Influences of all disturbing Causes; and V will be as $L^{\frac{1}{4}}$, and P as $\frac{L^{\frac{1}{4}}}{L}$ in all healthful and perfectly well proportioned Bodies under the same Circumstances. Hence, if we can find V and P in a healthful and well proportioned Body of any one length, we may find them by their Measures $L^{\frac{1}{4}}$ and $\frac{L^{\frac{1}{4}}}{L}$, in healthful and well proportioned Bodies of all other Lengths in the Morning, when the Bodies are sitting, and their Hearts are free from the Influences of all disturbing Causes.

The Velocity with which the Blood flows out of the left Ventricle of the Heart into the Aorta, may be found by knowing the cubic Inches of Blood which flow into the Aorta in one Systole, the Orifice of the Aorta in square Inches and decimal Parts of an Inch, and the Time of one Systole in Seconds and decimal Parts of a Second; for, putting K for the cubic Inches of Blood which flow

flow out of the Heart into the Aorta in one Systole, O for the Orifice of the Aorta in square Inches and decimal Parts, and T for the Time of one Systole in Seconds and decimal Parts of a Second, $\frac{K}{O \cdot T}$ will be equal to V. For $\frac{K}{O}$ is the Length described by the Blood with the Velocity V in the Time T, and $\frac{K}{O \cdot T}$ is the Length described with the same Velocity in 1". For the Velocity being given, the Times are as the Spaces described in those Times, that is, $T \cdot 1'' :: \frac{K}{O} \cdot \frac{K}{O \cdot T} = V$. The Time of one Systole of the Heart in Seconds and Parts of a Second, may be found from the Number of Pulses in a Minute, that is, T may be found from P. For allowing, what is generally supposed, the Time of one Systole, to be but half the Time of a Diaστοle, the whole Time of all the Systoles, or all the Pulses, in a Minute, will be only 20 Seconds. But if the Time of P Pulses be 20 Seconds, the Time of one Pulse

Of the Food and Discharges

Pulse will be $\frac{20}{p}$, which Time is equal to T . And therefore, if $\frac{20}{p}$ be substituted instead of T in the foregoing Measure of the Velocity $\frac{K}{O T}$, we shall have $\frac{K P}{20 O} = V$.

If 1500 Grains, or 5.6177 cubic Inches of Blood, flow out of the left Ventricle of the Heart into the Aorta in one Systole, in a strong well proportion'd Man 72 Inches in Length, when his heart is free from the Influences of all disturbing Causes; and if the Periphery of his Aorta when distended by the Force of the Heart pressing out this Quantity of Blood, be 3.5 Inches; all which may be allowed as not exceeding the Truth; then putting D for the Diameter of the Aorta, we shall have $D = 1.114085$, $D^2 = 1.241185$, $D^{\frac{3}{2}} = 1.0555$, and $O = 0.974824$. From some Experiments, the Morning Pulse of such a Man, when he is sitting and his Heart is free from the Influences of all disturbing Causes, beats about 54 times in a Minute.

a Minute. This Number, though considerably less than the Number assigned to strong well proportioned Bodies of that Length in the Table p. 136 of the *Animal Oeconomy*, yet I believe is much nearer the Truth, and may be allowed in Bodies which live regularly, and eat and drink but little at Night; for eating and drinking plentifully at Night commonly makes the Pulse quicker the next Morning. The Number 65 in that Table, was a Mean taken from the Morning Pulses of 120 Soldiers, and on that account might be greater than the Number corresponding to strong well proportioned Men of that Length, who live in a very regular and temperate manner. In such Bodies therefore K is 5,6177, P is 54, $20 \times O$ is 19.496480, and $\frac{K P}{20 O}$ is 15.48 Inches, equal to V. So that the Blood flows out of the Heart into the Aorta of strong well proportioned Men 72 Inches in length, when their Hearts are not influenced by disturbing Causes, with a Velocity that carries it at the rate of 15.48 Inches in 1".

This

This Velocity being found, the Velocity with which the Blood flows into the Aorta of a perfectly well proportioned Body of any other Length in the Morning, when the Body is sitting, and its Heart is free from the Influences of all disturbing Causes, may be found by this Analogy. As 2.913 the biquadrate Root of 72, is to 15.48, the Velocity of the Blood flowing into the Aorta of a healthful well proportioned Body 72 Inches in Length, in the Morning, when the Body is sitting, and its Heart is free from the Influences of all disturbing Causes; so is the biquadrate Root of the Length of any other healthful well proportioned Body, in the same Situation and same undisturbed State of the Heart, to the Velocity with which the Blood flows into the Aorta of that Body; that is, $2.913 : 15.48 :: L^{\frac{1}{4}}. V$; whence $5.314 L^{\frac{1}{4}} = V$. By this Measure of V , the Numbers in the Column V of Table 1. were computed. The Numbers in the Column P of the same Table, were computed by a Measure formed from this Analogy;

Analogy : as 0.04472 the biquadrate Root of 63 divided by 63, is to 60 the Morning Number of Pulses in a Minute of a healthful well proportioned Body 63 Inches in Length, when the Body is fitting and its Heart is free from the Influences of all disturbing Causes ; so is the biquadrate Root of the Length of any other healthful well proportioned Body divided by that Length, to the Morning Number of Pulses in a Minute, when the Body is fitting, and its Heart is free from the Influences of all disturbing Causes ; that is,

$$0.04472 : 60 :: \frac{L^{\frac{3}{4}}}{L} : P.$$

Whence, $1341.7 \times \frac{L^{\frac{3}{4}}}{L} = P.$ And the Numbers in the Column **F** were computed by a Measure formed from this Analogy, $63 \times 63^{\frac{3}{4}} = 177.4899 : 56 :: L \times L^{\frac{3}{4}}. F :$ Whence $0.3155 \times L \times L^{\frac{3}{4}} = F.$ These Measures of **V**, **P** and **F**, may be allowed in healthful well proportioned Bodies, whose Hearts are not influenced by disturbing Causes.

TABLE

Of the Food and Discharges

L	$L\frac{1}{4}$	V	$\frac{L\frac{1}{4}}{L}$	P	$L \times L\frac{1}{4}$	F
72	2.9130	15.4	0.04046	54	209.7360	66.17
70	2.8925	15.37	0.04132	55	202.4750	63.88
68	2.8716	15.26	0.04223	57	195.2688	61.60
66	2.8503	15.15	0.04319	58	188.1198	59.35
64	2.8284	15.03	0.04419	59	181.0176	57.11
63	2.8173	14.97	0.04472	60	177.4899	56.00
62	2.8060	14.91	0.04526	61	173.9782	54.89
60	2.7830	14.79	0.04638	62	166.9860	52.68
58	2.7596	14.66	0.04758	64	160.0568	50.50
56	2.7355	14.54	0.04885	66	153.1880	48.33
54	2.7108	14.41	0.05020	67	146.3832	46.18
52	2.6854	14.27	0.05164	69	139.6408	44.06
50	2.6592	14.13	0.05318	71	132.9550	41.95
48	2.6322	13.99	0.05483	74	126.3456	39.86
46	2.6043	13.84	0.05661	76	119.7978	37.80
44	2.5755	13.69	0.05853	79	113.3220	35.75
42	2.5456	13.53	0.06061	81	106.9194	33.73
40	2.5149	13.36	0.06287	84	100.5960	31.74
38	2.4828	13.19	0.06534	88	94.3464	29.77
36	2.4495	13.02	0.06804	91	88.1820	27.82
34	2.4147	12.83	0.07102	95	82.0998	25.90
32	2.3784	12.64	0.07432	100	76.1088	24.01
30	2.3403	12.44	0.07801	105	70.2090	22.15
28	2.3003	12.22	0.08215	110	64.4084	20.32
26	2.2581	12.00	0.08685	117	58.7106	18.52
24	2.2134	11.76	0.09222	124	53.1192	16.76
22	2.1658	11.51	0.09844	132	47.6454	15.03
20	2.1147	11.24	0.10573	142	42.2940	13.34
18	2.0598	10.95	0.11443	154	37.0746	11.70
16	2.0000	10.63	0.12500	168	32.0000	10.10
14	1.9343	10.28	0.13816	185	27.0800	8.54
12	1.8612	9.89	0.15510	208	22.3344	7.05

The

The Velocity, with which the Blood flows out of the left Ventricle of the Heart into the Aorta of a healthful Body, when it is fitting, and its Heart is free from the Influences of all disturbing Causes, being universally as the square Root of the Diameter of the Aorta; that is, V being universally as $D^{\frac{1}{2}}$; and V and $D^{\frac{1}{2}}$ in a well proportioned Body 72 Inches in Length, being 15.48 and 1.0555; the Velocity, with which the Blood flows into the Aorta of any other healthful Body under the same Circumstances, may be found by this Analogy. $15.48 : 1.0555 :: V : D^{\frac{1}{2}}$. Whence $14.667 D^{\frac{1}{2}} = V$.

The disturbing Causes of the Motion of the Heart are Changes in the sensible Qualities of the Air, Heat and Cold, Dryness and Moisture, Errors in Food, in Exercise of Body, in the Times of sleeping and waking, and the Passions of the Mind; that is, a wrong Use of the *Non-naturals* is the common disturbing Cause of the Motion of the Heart.

B

TABLE

TABLE 2.

W	.	w	:	:	24	.	1
H	.	h	:	:	16	.	1
L	.	l	:	:	4	.	1
D	.	d	:	:	2	.	1
P	.	p	:	:	7	.	20
V	.	v	:	:	7	.	5
Q	.	q	:	:	28	.	5

In *Table 2* are exhibited, the Proportions of the Weights of the Bodies, of a healthful strong well proportioned Man, and a healthful strong well proportioned Child newly born; the Proportions of the Weights of their Hearts, of the Lengths of their Bodies, of the Diameters of their Aortas, of their Pulses in a Minute when their Hearts are free from the Influences of disturbing Causes, of the Velocities of the Blood in their Aortas, and of the Quantities of Blood which in a given Time flow through their Hearts or Lungs, when their Hearts are not affected by disturbing Causes. And from these Proportions compared with one another

another will arise some useful Observations.

Obs. 1. The Weight of the Heart with respect to the Weight of the Body, is greater in Children than in grown Bodies. It was greater in the Child than in the Man, in the Proportion of 3 to 2. For,

$$\frac{h}{w} \cdot \frac{H}{W} :: 1 \cdot \frac{16}{24} :: 3 \cdot 2.$$

Hence the Weight of the Heart with respect to the Weight of the Body, lessens continually from the Birth till Bodies come to their full Growth.

Obs. 2. The Quantity of Blood which flows through the Heart or Lungs in a given Time, in Proportion to the Weight of the Heart, or Quantity of Blood contained in the Body, which Quantity of Blood is proportional to the Weight of the Heart, is greater in Children than in grown Bodies. It was greater in the Child than in the Man, in the Proportion of 20 to 7. For

$$\frac{q}{h} \cdot \frac{Q}{H} :: \frac{5}{1} \cdot \frac{28}{16} :: 20 \cdot 7,$$

B 2 which

Of the Food and Discharges

which is the Proportion of their Pulses in a Minute.

Hence, the Quantity of Blood that flows through the Heart or Lungs in a given Time, in Proportion to the whole Quantity of Blood contained in the Body, lessens continually from the Birth till Bodies arrive at their full Growth.

Obs. 3. The Velocity of the Blood with respect to its Quantity, which Quantity is as the Weight of the Heart, is much greater in Children than in grown Bodies; it was greater in the Child than in the Man, in the Proportion of 80 to 7. For $\frac{v}{h} \cdot \frac{V}{H} :: \frac{5}{1} \cdot \frac{7}{16} :: 80 \cdot 7$.

Hence, though the Blood of Children moves slower than the Blood of grown Bodies, yet for its Quantity it moves much quicker, and much oftener passes through the Lungs. On which account the Blood of Children, notwithstanding the Slowness of its Motion, may by passing

ing oftener through the Lungs, and thereby receiving more of the Acid of the Air in proportion to its Quantity, be more fluid, and of a brighter Colour, than the Blood of grown Persons.

Obs. 4. The Quantity of Blood that flows through the Heart or Lungs in a given Time, in Proportion to the Weight of the Body, is greater in Children than in grown Bodies. It was greater in the Child than in the Man in the Proportion of 30 to 7. For $\frac{q}{w} \cdot \frac{Q}{W} :: \frac{5}{1} \cdot \frac{28}{24} :: 30 \cdot 7$.

Hence, though the Velocity of the Blood is less in Children than in grown Bodies, yet its Motion with respect to the Weight of the Body is greater.

Obs. 5. The Velocity of the Blood with respect to the Length of the Body, is greater in Children than in grown Bodies. It was greater in the Child than in the Man, in the Proportion of 20 to 7,

B 3

which

Of the Food and Discharges

which is the Proportion of their Pulses in a Minute. For $\frac{v}{1} \cdot \frac{V}{L} :: \frac{5}{1} \cdot \frac{7}{4} :: 20 \cdot 7$.

Hence, the Velocity of the Blood, and Number of Pulses in a Minute, with respect to the Length of the Body, lessen continually in growing Bodies till they arrive at their full Growth.

Several other Observations may be made from comparing the Proportions of this Table with one another.

To form some Idea of the flow Motion of the Blood in the capillary Blood-vessels, we must know the Bigness of the red Corpuscles of healthful arterial Blood, and the Order of Colours in the Table p. 88. of the *Dissertation on the Æther of Sir Isaac Newton*, to which Order the red Colour of those Corpuscles belongs. To know the Bigness of the red Corpuscles of healthful arterial Blood, we must allow for their Density, which by Doctor *Jurin's* Experiments, is 1.126, the Density of Water being 1.
For

For were those Corpuscles transparent, their Density is such, that the Sine of Incidence upon them would be to the Sine of Refraction, as $\sqrt{2.126} = 1.4581$, is to 1, by *Cor. Prop. 9.* of the *Dissertation*. And therefore the Thickness of the Corpuscles, that they may exhibit the same red Colour with those of Bubbles of Water of the first Order in the Table, must be less than the Thickness of the Skins of those Bubbles in the Proportion of 1 to 1.4581. And consequently D, which I put for the Thickness of the Corpuscles, will be equal to $\frac{1}{148148} \times \frac{1}{1.4581} = 0.00000046293$ Part of an Inch. The Diameter of the smallest Blood-vessel must be greater than the Thickness of a Corpuscle, to let a Corpuscle pass through it; I shall suppose it to be twice as great, and then the Diameter of the smallest Blood-vessel will be the 0.00000092586 Part of an Inch, and the square Root of this Diameter will be the 0.0009636 Part of an Inch, which being multiply'd into 14.667, gives the 0.014 Part of an Inch

Of the Food and Discharges

for the Space described by the Blood in a capillary Blood-vessel in a Second of Time. With this Velocity the Blood moves at the Rate of 50.4 Inches in an Hour. The Velocity of the Blood in the Aorta of a strong well proportioned Man 72 Inches in length, is above 1100 times greater than the Velocity of the Blood in a capillary Blood-vessel; and it is above 780 times greater in the Aorta of a strong well proportioned Child 18 Inches in length, than it is in a capillary Blood-vessel. This slow Motion of the Blood in the capillary Vessels is what qualifies it for Secretion.

That the Colour of healthful arterial Blood is the Red of the first Order, and that the Colour of healthful venal Blood is composed of the Violet and Indigo of the second Order, appears to me very probable from the following Considerations. The Colour of venal Blood in an inflammatory Fever, when drawn and exposed to the Air, first changes into a very faint Blue, and then into a White,
or

or Yellowish White like the Colour of Buff. These Changes shew, the Colour of this venal Blood to be the Beginning of Black of the first Order, and that its tinging Corpuscles are enlarged by the Acid of the Air. This Effect of the Acid of the Air on this venal Blood, is directly contrary to the Effect it produces in healthful venal Blood; for it changes its dark Colour into a good Red, without making it pass through the intermediate Colours of Blue, Green, Yellow and Orange; and therefore the Red into which it changes it, must be the Red of a superior Order; which shews that its tinging Corpuscles are lessened by the Acid of the Air. The Red into which the Colour of healthful venal Blood changes when exposed to the Air, must be the Red of the first Order; for if it was the Red of the second Order, as I supposed in the *Animal Oeconomy*, then the Change of the Colour of venal Blood, composed of the Purple and Indigo of the third Order, into the Beginning of Black of the first Order the Colour of
inflam-

Of the Food and Discharges

inflammatory venal Blood, would be a Change too great to be admitted. But if the Colour of healthful venal Blood be composed of the Violet and Indigo of the second Order, it may be changed by the Heat and Motion of the Blood in an inflammatory Fever, into the Beginning of Black in the first Order without any Inconvenience, and therefore may be allowed.

Hence, the Acid of the Air has a different Effect on the venal Blood in inflammatory Fevers, from what it has on the venal Blood in Health; it enlarges the tinging Corpuscles of the first, and lessens the tinging Corpuscles of the second. Black being the Colour which Bodies put on, when the Corpuscles on which their Colours depend are divided into smaller Particles by Putrefaction; and the Colour of venal Blood in inflammatory Fevers being much nearer to Black, than the Colour of venal Blood in Health; we may hence conclude, that venal Blood in inflammatory Fevers, is
nearer

nearer to a State of Putrefaction than venal Blood in Health ; and that the Acid of the Air has the same Effect upon venal Blood in inflammatory Fevers, as it has upon the Parts of Bodies dissolved by Putrefaction. All this will be readily conceived from Sir *Isaac Newton's* Account of the Composition of Particles of Salt, which I shall give in his own Words.

“ As Gravity makes the Sea flow
“ round the denser and weightier Parts
“ of the Globe of the Earth, so the At-
“ traction may make the watry Acid
“ flow round the denser and compacter
“ Particles of Earth for composing the
“ Particles of Salt. For otherwise the
“ Acid would not do the Office of a
“ Medium between the Earth and com-
“ mon Water, for making Salts dissolvable
“ in Water ; nor would Salt of Tartar
“ readily draw off the Acid from dis-
“ solved Metals, nor Metals the Acid
“ from Mercury. Now as in the great
“ Globe of the Earth and Sea, the
“ densest Bodies by their Gravity sink
“ down

Of the Food and Discharges

“ down in Water, and always endeavour
 “ to go towards the Center of the Globe;
 “ so in Particles of Salt, the densest Mat-
 “ ter may always endeavour to approach
 “ the Center of the Particle: So that a
 “ Particle of Salt may be compared to a
 “ Chaos; being dense, hard, dry, and
 “ earthy in the Center; and rare, soft,
 “ moist, and watry in the Circumference.
 “ And hence it seems to be that Salts are
 “ of a lasting Nature, being scarce de-
 “ stroyed, unless by drawing away their
 “ watry Parts by Violence, or by letting
 “ them soak into the Pores of the central
 “ Earth by a gentle Heat in Putrefaction,
 “ until the Earth be dissolved by the
 “ Water, and separated into smaller Par-
 “ ticles, which by reason of their Small-
 “ ness make the rotten Compound appear
 “ of a black Colour. Hence also it may
 “ be that the Parts of Animals and Ve-
 “ getables preserve their several Forms,
 “ and assimilate their Nourishment; the
 “ soft and moist Nourishment easily
 “ changing its Texture by a gentle Heat
 “ and Motion, till it becomes like the
 “ dense,

“ dense, hard, dry, and durable Earth
“ in the Center of each Particle. But
“ when the Nourishment grows unfit to
“ be assimilated, or the central Earth
“ grows too feeble to assimilate it, the
“ Motion ends in Confusion, Putrefaction,
“ and Death.”

I now proceed to give an Account of
the Discharges and Food of human Bo-
dies.

*Of the Food and Discharges of Human
Bodies.*

PROPOSITION I.

THE Sum of the Discharges by Per-
spiration, Urine, and Stool in a na-
tural Day or any other Time, is equal to
the Quantity of Food taken in that Time,
lessened by the Difference of the Weights of
the Body at the beginning and end of the
Time, if the Body be heavier at the end

Of the Food and Discharges

of the Time than at the beginning ; and increased by that Difference, if the Body be lighter. If p , u , and s denote the Quantities of Perspiration, Urine, and Stool discharged in a natural Day or any other Time, F the Quantity of Food taken in that Time, and d the Difference of the Weights of the Body at the beginning and end of the Time ; then $p + u + s = F + d$, when the Body is lighter, and $p + u + s = F - d$, when the Body is heavier, at the end of the Time than at the beginning.

For if a Body be of the same Weight at the beginning and end of a natural Day or any other Time, the Sum of the Discharges made by Perspiration, Urine, and Stool, will be equal to the Quantity of Food taken, in that Time. But if the Weight of the Body be greater or lesser at the beginning of the Time than at the end of it, the Sum of the Discharges will fall short of or exceed the Food by the Difference of those Weights. And therefore, that Difference must be added to the Food when the Discharges

exceed the Food, and subducted from the Food when the Food exceeds the Discharges; that is, d must be added to F in the first Case, and subducted from F in the second, to make $p + u + s = F + d$ when the Body is lighter, and $p + u + s = F - d$ when the Body is heavier at the end of the Time than at the beginning. Therefore the Proposition is true.

Cor. I. The Quantity of Perspiration in a natural Day or any other Time, is equal to the Sum of the Food and Difference of the Weights of the Body at the beginning and end of the Time, lessened by the Quantities of Urine and Stool in that Time, when the Body is lighter at the end of the Time than at the beginning; and equal to the Food lessened by the Difference of the Weights of the Body, and by the Quantities of Urine and Stool in that Time, when the Body is heavier at the end of the Time than at the beginning: that is, $p = F + d - u - s$ when the Body is lighter, and $p = F - d - u - s$ when the Body is heavier.

Hence,

Hence, by knowing F , d , u and s , all which may be had by weighing, the Quantity perspired in any Time, whatever that Time be, may be found by this Corollary.

Cor. 2. If a Body be of the same Weight at the End of a natural Day or any other Time, as at the Beginning; the Sum of the Discharges, and Quantity of Food, in that Time, will be equal; and Perspiration will be equal to the Food, lessened by the Quantities of Urine and Stool. If d be 0, then $p + u + s = F$; and $p = F - u - s$.

Cor. 3. If no Food be taken between the two Times of weighing the Body, or, in other Words, if the Person fasts during the intermediate Time; the Sum of the Discharges by Perspiration, Urine, and Stool in that Time, will be equal to the Difference of the Weights of the Body at the Beginning and End of the Time; and Perspiration will be equal to that Difference

ference lessened by the Quantities of Urine and Stool. For F being o , and d being affirmative, because the Body is necessarily lighter at the End of the Time than at the Beginning; $p + u + s = d$, and $p = d - u - s$.

Cor. 4. If the Perspiration and Urine discharged in a natural Day or any other Time be equal, and the Body be of the same Weight at the end of the Time as at the beginning; double the Quantity of Perspiration added to the Stool will be equal to the Food; and the Quantity of Perspiration will be equal to half the Difference of the Food and Stool. If $p = u$, and d be o ; then $2 p + s = F$, and $p = \frac{F - s}{2}$.

Cor. 5. If a Body, by taking a given Quantity of Food in a natural Day, be of the same Weight at the end of the Day as at the beginning; the Sum of Perspiration, Urine, and Stool discharged in that Time, will be given. If d be o , and

C F be

Of the Food and Discharges

F be given, or as 1; $p + u + s$ will be given, or as 1. When this Corollary obtains, if one of the three Discharges be increased or lessened, one or the Sum of the other two will be equally lessened or increased; otherwise the Sum of the three could not be given.

PROPOSITION II.

THE Sum of the Discharges by Perspiration, Urine, and Stool, in a natural Day or any other Time, is nearly proportional to the mean Quantity of Blood, which in that Time flows out of the Heart into the Aorta in one Systole, and the Number of Systoles or Pulses in the same Time, taken together. If q denote the mean Quantity of Blood, which in a natural Day or any other Time, flows out of the Heart into the Aorta in one Systole, and N the Number of Pulses in that Time, then will $p + u + s$ be nearly proportional to $q N$.

For the fecerning Ducts, which draw off the Humours of Perspiration and
Urine

Urine from the Blood, are Continuations of the Blood-veffels, and differ from them only in this, that they are too small to let the red Parts of the Blood, which are its largest Parts, pass through them; and therefore, the Sum of these two Humours flowing through these small Ducts in a natural Day or any other Time, will be proportional to the Quantity of Blood that flows out of the Heart into the Aorta in that Time. But the Quantity of Blood which flows out of the Heart into the Aorta in a natural Day or any other Time, is equal to the mean Quantity thrown out of the left Ventricle in one Systole, and Number of Systoles or Pulses in that Time, taken together. And therefore, the Sum of Perspiration and Urine in a natural Day or any other Time, will be proportional to the mean Quantity of Blood thrown into the Aorta in one Systole, and Number of Systoles or Pulses in that Time, taken together; that is, $p + u$ will be proportional to $q N$. Stool is a very inconsiderable Discharge when compared with Perspiration and Urine; and on that

Of the Food and Discharges

account will but little increase their Sum when added to it. And consequently, $p + u + s$ will be nearly proportional to $q N$.

Cor. If the same Quantity of Blood, flows out of the Heart into the Aorta in a natural Day, or any other Time; the Sum of the Discharges by Perspiration, Urine, and Stool in that Time, will be nearly the same. If $q N$ be given, $p + u + s$ will be nearly given.

$q N$ will be given, when both q and N are given, or when q is as $\frac{1}{N}$. And when $q N$ is given, $p + u + s$ will be nearly given, or nearly as 1. And consequently, when any one of the three Discharges is increased or lessened, one, or the Sum of the other two, will be lessened or increased nearly equally; otherwise their Sum could not be nearly given, or nearly as 1.

PROPOSITION III.

THE Quantity of Food taken in a natural Day or any other Time, increased by the Difference of the Weights of the Body at the beginning and end of the Time, when the Body is lighter at the end of the Time than at the beginning; and lessened by that Difference, when the Body is heavier, is nearly proportional to the Quantity of Blood which flows out of the left Ventricle of the Heart into the Aorta in that Time; that is, $F + d$ is nearly proportional to $q N$, when the Body is lighter at the end of the Time than at the beginning; and $F - d$ is nearly proportional to $q N$, when the Body is heavier.

For $p + u + s = F + d$, when the Body is lighter, and $p + u + s = F - d$, when the Body is heavier, at the end of the Time than at the beginning; by *Prop. 1.* And $p + u + s$, is nearly proportional to $q N$; by *Prop. 2.* And therefore, $F + d$ is nearly proportional to $q N$ when the

Of the Food and Discharges

Body is lighter, and $F-d$ is nearly proportional to qN when the Body is heavier, at the end of the Time than at the beginning.

Cor. 1. If a Body be of the same Weight at the end of a natural Day or any other Time, as at the beginning; the Quantity of Food taken in that Time, will be nearly proportional to the Quantity of Blood, which flows out of the Heart into the Aorta in the same Time. If d be 0, F will be nearly proportional to qN .

Cor. 2. If a Body be of the same Weight at the end of a natural Day or any other Time, as at the beginning; and if during that Time, the mean Quantity of Blood flowing out of the Heart into the Aorta in one Systole be given; the Quantity of Food taken in that Time, will be nearly proportional to the Number of Pulses in the same Time. If d be 0, and q be given; F will be nearly proportional to N .

Cor.

Cor. 3. If the mean Quantity of Blood, which in a natural Day or any other Time flows out of the Heart into the Aorta in one Systole, be reciprocally proportional to the Number of Pulses in that Time, and if the Body be of the same Weight at the end of the Time as at the beginning ; then will the Quantity of Food be nearly given. If q be as $\frac{1}{N}$, and d be 0, F will be nearly as 1.

Cor. 4. If the Heart during a natural Day or any other Time, throws the same Quantity of Blood into the Aorta in each Systole ; if the Number of Pulses in that Time, be proportional to the Morning Number of Pulses in a Minute when the Body is sitting, and its Heart is free from the Influences of all disturbing Causes ; and if the Body be of the same Weight at the end of the time as at the beginning ; then will the Quantity of Food taken in that Time, be nearly proportional to the Morning Number of Pulses

Of the Food and Discharges

in a Minute when the Body is sitting, and its Heart is free from the Influences of all disturbing Causes. If q be given, N be as P , and d be o ; F will be as P .

I now proceed to an Explanation of the following Statical Tables, and Tables of Animals. But before I do this, I shall mention a few Particulars relating to myself.

I am now, in *May* 1747, in the 68th Year of my Age. The Length of my Body is 63 Inches: I am of a sanguin but not robust Constitution, and at present am neither lean nor fat. In the Year 1721, the Morning Weight of my Body without Cloaths was about 131 Averdupois Pounds, the daily Quantity of my Food at a Medium was about 85 Averdupois Ounces, and the Proportion of my Drink to my Meat, I judge was at that Time about 2.5 to 1. At the latter end of *May* 1744, my Weight was above 164 Pounds, and the Proportion of my Drink to my Meat was considerably greater than before, and had been so for some Time.

Time. I was then seized with a paralytick Disorder, which obliged me to make an Alteration in my Diet. In order to settle the Proportion of my Drink to my Meat, I considered what others have said concerning this Proportion. According to *Sanctorius*, though he reckons it a Disproportion, the Drink to the Meat in his Time, was above 10 to 3 in temperate Bodies, *Aph.* 68. *Sect.* 3. *Cornaro's* Drink to his Meat, was as 7 to 6; *Mr. Rye's* in Winter, as 4 to 3; *Dr. Lining's* at a Medium for a whole Year, as 11 to 3; and my Drink to my Meat, as 5 to 2. A Mean taken from all these makes the Drink to the Meat to be as 2176 to 1000, which is above 2 to 1. I kept near to the Proportion of 2 to 1 in the Experiments, from which I composed Table 2. In 17 Months, from the Time I was seized to the End of *October* 1745, I lost above twenty Pounds of my Weight, and above twice as much in the first five Months as I did in the twelve Months after. To this Loss of Weight, caused chiefly, by the Regulation of my Diet, I impute my Recovery from
my

my last Disorder, excepting which, and a Vertigo I had about ten or twelve Years before, I have had no Disorder of Consequence, or that confined me, since I was a Boy.

Explanation of the Tables.

TABLE I is the same with the Table published in the *Animal Oeconomy*, p. 259, and contains the daily Food and daily Discharges taken at a Medium from the Experiments of eight Months, beginning with *April* the last of the Spring Months, and ending with *November* the first of the Winter Months. In this Table the natural Day was divided into three Parts, Morning, Afternoon, and Night; the Morning contained six Hours from eight to two, the Afternoon six Hours from two to eight, and the Night the remaining twelve Hours. The Day made up of Morning and Afternoon, was equal to the Night. I observed the Food and Discharges in these three Parts of the natural Day, every Day for eight Months, and

and with the Means taken from all the Food and all the Discharges in the several Months, I compos'd this Table. The Numbers expressing the Food and the Discharges are Averdupois Ounces. *p* expresses the mean Quantity of Perspiration, and *u* the mean Quantity of Urine, discharged daily in each Month; and consequently, the Numbers in the Column $\frac{p}{u}$, both in this and all the following Tables, express the Proportions of Perspiration to Urine in the several Months, or express Perspiration, Urine being always expressed by 1; for Instance, Perspiration was to Urine in *April*, as 0.904 to 1, or as 904 to 1000; and in *August* Perspiration was to Urine, as 1.713 to 1, or as 1713 to 1000. This Table was made in the Year 1721.

Table 2 was made from the Experiments of a whole Year, beginning the first of *November* 1744, and ending with *October* 1745. In this Table the Day contained the Hours of my being up, and

5 the

Of the Food and Discharges

the Night the Hours of my being in Bed. For the first seven Months to the Beginning of *June*, the Time of my being up was fifteen Hours, and the Time of my being in Bed nine Hours; and for the last five Months to the End of *October*, the Time of my being up was 14.5 Hours, and the Time of my being in Bed 9.5 Hours. The Food and Discharges in this and all the other Tables, are expressed in Averdupois Ounces. For whenever any other Ounce was used, as the *Venetian* Ounce in *Italy* by *Sanctorius*, and the Troy Ounce in *South Carolina* by Dr. *Lining*, I reduced it to the Averdupois Ounce, for the more easy comparing of the Quantities of Food, and Quantities of the Discharges, of Persons living in different Climates. The *Venetian* Pound, according to Mr. *Greaves*, contains 5528 Grains, and the *Venetian* Ounce $460\frac{2}{3}$ Grains; Dr. *Lining's* Pound contains 16 Troy Ounces, or 7680 Grains; and the Averdupois Pound used by Dr. *Kiell*, Mr. *Rye*, and myself, contains 7000 Grains, and the Ounce $437\frac{1}{2}$ Grains.

Table

Table 3 consists of two Parts. The first Part was made from *Table 1*, and the second Part from *Table 2*. Each Part contains the mean Quantities of Food taken daily in the several Months and Seasons of the Year; and the mean Quantities of Urine and Perspiration, which were drawn off from the Blood hourly in the Day and in the Night. In the first Part of the *Table*, the Day was twelve Hours, and the Night twelve Hours; and consequently *T* was 12 both Day and Night. In the second Part of the *Table*, the Day and Night being the Times of my being up and in Bed, which were 15 and 9 in the first seven Months, and 14.5 and 9.5 in the last five Months; *T* was 15 in the Day, and 9 in the Night in the first seven Months; and 14.5 in the Day, and 9.5 in the Night, in the last five Months. For Example, the mean Quantity of Urine discharged daily in *November*, was 20.4 Ounces in the first Part of the *Table*, and 17.96 Ounces in the second Part; and dividing 20.4 by 12, and

17.96

Of the Food and Discharges

17.96 by 15, we have 1.700, and 1,197, the mean Quantities of Urine discharged hourly in the Day in that Month. And the mean Quantity of Perspiration discharged daily in that Month was 17.63 in the first Part of the *Table*, and 20.32 in the second Part; and dividing 17.63 by 12, and 20.32 by 15, the Quotients 1.470 and 1.355 will be the mean Quantities of Perspiration discharged hourly in the Day in that Month. In like manner were got the mean hourly Discharges of Urine and Perspiration in the Night: And thus the whole *Table* was formed.

Table 4 is the same with the *Table* in p. 277. of the *Animal Oeconomy*, and contains the mean Quantities of Perspiration and Urine which were actually discharged hourly by two Persons B and D, myself and another, in four very hot Days in Summer, beginning at six in the Morning, and ending at ten at Night. The Numbers corresponding to six in the Morning, were the Quantities of Perspiration and Urine which were drawn off
from

from the Blood in every Hour of the Night, taking one Hour with another. We both eat our Breakfast at eight in the Morning, dined at two, and supped at eight at Night. The mean Quantities of our daily Food in these four Days, were 86 and 63 Averdupois Ounces. We both kept within, and used no Exercise during that Time.

Table 5 was made from *Table 4*, and contains the mean Quantities of the Food of B and D in a natural Day; and the mean Quantities of Perspiration and Urine, and the Proportion of the mean Quantity of Perspiration to the mean Quantity of Urine, in the three equal Parts of the natural Day, Morning, Afternoon, and Night; supposing the Morning to begin at six and end at two, the Afternoon to begin at two and end at ten, and the Night to begin at ten and end at six next Morning.

Table 6 was made from the particular Tables of the Months, from the Means
of

Of the Food and Discharges

of which Tables I composed Table 2 ; and it contains several of the most remarkable Changes of my Weight, together with my Food and Discharges on those Days in which the Changes happened. The first Column contains the Names of the Months ; the second contains the Days of the Months on which the Changes happened ; the third contains the Quantities of Food taken on those Days ; the fourth contains the Gain or Loss of Weight, Gain being denoted by —, and Loss by + ; the fifth contains the Sum of the Discharges on those Days by Perspiration, Urine, and Stool, which Discharges are set down separately in the sixth, seventh, and eighth Columns ; and the ninth Column contains the Proportion of Perspiration to Urine.

Tables 7, 8, 9, are the yearly Tables, of Dr. Keill at Northampton in England, of Mr. Rye at Cork in Ireland, and of Dr. Lining at Charles-town in South Carolina. In Dr. Lining's Table, d and n express Day and Night ; and p and u Perspiration

tion and Urine as in the other *Tables*; for Instance, $\frac{d u}{n u}$ expresses the Proportion of the Day's Urine to the Night's Urine, and $\frac{d p}{n p}$ the Proportion of the Day's Perspiration to the Night's Perspiration. I have reduced his Degrees of Heat to the Scale of Sir *Isaac Newton*.

Table 10 contains the Quantities of Food in *Averdupois Ounces*, and the Proportions of Perspiration to Urine in the four Seasons of the Year, in *Italy, England, Ireland, and South Carolina*.

Sanctorius tells us, that if the Meat and Drink of one Day be eight Pounds, insensible Perspiration is wont to rise to about five Pounds. *Aph. 6. Sect. 1.* And consequently, there will remain three Pounds to be discharged by Urine and Stool, thirty-two Ounces by Urine, and four Ounces by Stool. *Aph. 59, 60. Sect. 1.* These Quantities of Perspiration and Urine, must be the Quantities discharged

D in

Of the Food and Discharges, etc.

in a natural Day in Summer, by *Aph.* 21.
Seçt. 1. And from these we may get the
 Quantities of Perspiration and Urine in a
 natural Day in Winter, by *Aph.* 41. *Seçt.*
 2. which Quantities will be 48 and 44.
 Hence in *Italy* Perspiration is to Urine in
 a natural Day in Summer as 1.870 is to
 1, or as 1870 is to 1000; and in a natu-
 ral Day in Winter Perspiration is to Urine
 as 1.090 is to 1, or as 1090 is to 1000.
 The Sums of the Proportions of Perspira-
 tion to Urine in Summer and Winter, di-
 vided by 2, will nearly give their Pro-
 portion in Spring and Autumn; which
 therefore will be as 1.480 is to 1, or as
 1480 is to 1000.

STATICAL

STATISTICAL TABLES.

TABLE I.

Dr. ROBINSON, Aged 42, A. D. 1721.

Months	Break		Food		Morning.		Afternoon.		Night.		Whole		Stool	Whole Difch.	$\frac{p}{u}$
	Dinn.	Supp.	Whole	Supp.	Urine	Perfp.	Urine	Perfp.	Urine	Perfp.	Urine	Perfp.			
April	23.37	44.00	16.22	83.59	8.38	11.00	17.40	11.26	16.59	16.05	42.37	38.31	5.27	85.95	0.904
May	23.30	47.35	21.62	92.27	7.18	13.12	17.77	11.62	16.75	19.52	41.70	44.26	7.42	93.38	1.061
June	27.20	48.13	15.10	90.43	7.13	15.68	12.10	12.70	14.53	22.77	33.76	51.15	5.32	90.23	1.515
July	25.98	41.12	15.53	82.63	7.78	14.83	10.30	12.28	12.30	19.32	30.38	46.43	5.85	82.66	1.528
August	22.04	43.10	20.54	85.68	7.35	17.42	10.00	13.15	12.58	20.69	29.93	51.27	4.48	85.68	1.713
Septem.	18.72	48.00	19.74	86.46	7.50	13.48	13.12	11.02	15.50	19.93	36.12	44.43	5.15	85.70	1.230
October	18.85	45.88	15.68	80.42	7.20	9.98	13.72	10.10	16.68	17.52	37.60	37.60	5.05	80.25	1.000
Novem.	22.38	40.92	15.06	78.37	7.65	8.94	12.75	8.69	16.69	18.02	37.09	35.65	4.71	77.45	0.961
Mean	22.73	44.81	17.44	84.98	7.52	13.06	13.39	11.35	15.20	19.22	36.12	43.64	5.41	85.16	1.208
Summer	25.49	45.53	17.42	88.44	7.36	14.54	13.39	12.20	14.53	20.54	35.28	47.28	6.20	88.76	1.340
Autumn	19.87	45.66	18.65	84.18	7.35	13.63	12.28	11.42	14.92	19.38	34.55	44.43	4.89	83.87	1.286
Mean	22.68	45.59	18.03	86.31	7.35	14.08	12.83	11.81	14.72	19.97	34.91	45.86	5.54	86.31	1.315

TABLE 2.
Dr. ROBINSON, Aged 64-5, A. D. 1744-5.

Months.	Food.		± d	Urine		Urine Spec.		Urine		Perspiration.		Stool	Whole Difch.	$\frac{p}{u}$	Pulse in 1'.
	Meat	Drink		Day.	Spec. Grav.	Night	Spec. Grav.	whole	Day.	Night	Whole				
Novem.	18.77	41.73	+	14	17.96	10252	12.22	10273	30.18	20.32	8.03	28.35	60.98	0.939	63
Decem.	20.05	41.98	-	11	18.90	10260	11.76	10277	30.66	19.46	8.49	27.95	61.67	0.911	63
January	19.70	43.34	+	66	17.75	10251	13.43	10255	31.18	18.07	9.22	27.29	65.14	0.875	66
Febr.	18.59	43.10	-	4	14.91	10260	13.03	10266	27.94	17.07	9.55	26.62	61.57	0.953	65
March	20.80	39.90	+	18	16.83	10250	12.13	10269	28.96	19.18	9.44	28.62	61.28	0.988	63
April	19.93	40.10	+	7	17.00	10252	10.89	10275	27.89	21.86	7.86	29.73	60.26	1.066	62
May	19.39	36.93	+	8	19.12	10256	9.05	10291	28.17	19.18	6.71	25.89	56.58	0.919	65
June	22.10	34.23	-	13	16.20	10234	9.07	10256	25.27	19.51	8.12	27.63	55.90	1.093	66
July	22.50	33.79	-	5	17.42	10247	10.37	10271	27.79	17.45	7.63	25.08	56.12	0.902	63.4
August	22.04	34.38	+	5	15.70	10234	10.25	10264	25.95	20.27	7.98	28.25	56.58	1.088	63.8
Septem.	20.53	35.06	+	42	15.61	10236	10.13	10279	25.75	19.80	8.75	28.55	56.99	1.108	63.6
October	19.63	34.32	-	6	15.77	10235	10.42	10272	26.19	18.34	7.16	25.50	53.77	0.973	60.4
Mean	20.33	38.24	+	121	16.93	10247	11.06	10270	27.99	19.21	8.24	27.45	58.89	0.980	63.6
Winter	19.51	42.35	+	69	18.20	10254	12.47	10268	30.67	19.28	8.58	27.86	62.60	0.908	64
Spring	19.77	41.03	+	21	16.25	10254	12.02	10270	28.27	19.37	8.95	28.32	61.04	1.002	63
Summer	21.33	34.98	-	10	17.58	10246	9.50	10273	27.08	18.71	7.49	26.20	56.20	0.967	64.8
Autumn	20.73	34.59	+	41	15.69	10235	10.26	10272	25.96	19.47	7.96	27.43	55.77	1.057	62.6
Mean	20.33	38.24	+	121	16.93	10247	11.06	10270	27.99	19.21	8.24	27.45	58.89	0.980	63.6

T A B L E 3.

Months.	A. D. 1721.			A. D. 1744-5.		
	Food.	$\frac{u}{T}$ Day. Night	$\frac{p}{T}$ Day. Night	Food.	$\frac{u}{T}$ Day. Night	$\frac{p}{T}$ Day. Night
November	78.37	1.700 1.391	1.470 1.502	60.50	1.197 1.358	1.355 0.892
December				62.03	1.260 1.307	1.297 0.943
January				63.04	1.183 1.492	1.205 1.024
February				61.68	0.994 1.448	1.138 1.061
March				60.70	1.122 1.348	1.279 1.049
April	83.59	2.148 1.382	1.855 1.338	60.03	1.133 1.210	1.457 0.873
May	92.27	2.079 1.396	2.062 1.627	56.32	1.274 1.006	1.279 0.745
June	90.43	1.602 1.211	2.365 1.897	56.33	1.117 0.955	1.345 0.855
July	82.63	1.507 1.025	2.260 1.610	56.29	1.201 1.091	1.203 0.803
August	85.68	1.446 1.048	2.547 1.724	56.42	1.083 1.079	1.398 0.840
September	86.46	1.718 1.291	2.042 1.661	55.60	1.076 1.066	1.365 0.921
October	80.42	1.743 1.390	1.673 1.460	53.95	1.088 1.097	1.265 0.753
Mean	84.98	1.742 1.267	2.034 1.602	58.57	1.144 1.202	1.298 0.896
Winter				61.86	1.213 1.385	1.285 0.953
Spring				60.80	1.083 1.335	1.291 0.994
Summer	88.44	1.729 1.211	2.229 1.71	56.31	1.197 1.018	1.273 0.805
Autumn	84.19	1.635 1.243	2.088 1.61	55.32	1.082 1.080	1.343 0.838
Mean	86.31	1.682 1.227	2.158 1.664	58.57	1.144 1.202	1.298 0.896

TABLE 4.

Hours.	B		D	
	Perfsp.	Urine	Perfsp.	Urine
6	1.875	0.937	2.000	1.000
7	1.833	1.000	1.400	1.000
8	2.000	1.000	1.500	1.200
9	2.000	1.166	1.800	1.300
10	2.000	1.500	1.900	1.000
11	1.750	1.000	1.400	1.000
12	2.333	1.000	1.800	1.000
1	2.333	1.250	1.500	1.000
2	2.000	1.000	1.500	1.000
3	3.500	1.500	2.000	1.000
4	2.333	2.000	1.500	1.143
5	2.333	2.000	1.800	1.000
6	2.666	2.000	2.000	1.000
7	2.000	2.000	2.000	1.000
8	2.333	2.333	2.000	1.000
9	2.333	1.666	1.500	1.500
10	2.333	1.666	1.500	1.500

TABLE 5.

Men	Morning.			Afternoon.			Night.			Whole.		
	F	P	$\frac{P}{u}$	F	P	$\frac{P}{u}$	F	P	$\frac{P}{u}$	F	P	$\frac{P}{u}$
B	86	16.249	8.916	1.822	15.165	1.307	15.000	7.496	2.001	51.080	31.577	1.618
D	63	12.800	8.500	1.506	9.143	1.564	16.000	8.000	2.000	43.100	25.643	1.681

T A B L E 6.

Months	Day.	Food	\pm d	Difch	Urine	Perfsp.	Stool.	$\frac{P}{u}$
Nov.	18	61.0	+29	90.0	49.00	38.00	3.00	0.775
Dec.	13	62.0	+11	73.0	41.00	28.05	3.05	0.695
	14	67.0	+6	73.0	41.00	29.00	3.00	0.707
	28	54.0	+17	71.0	34.00	34.00	3.00	1.000
Janu.	2	62.0	+37	99.0	31.25	40.75	27.00	1.304
	25	40.5	+26	66.5	34.05	28.00	4.00	0.811
	30	82.0	+7	89.0	31.75	24.05	32.75	0.771
	31	61.5	+32	93.5	28.05	30.00	35.00	1.052
Febr.	1	64.5	+24	88.5	18.00	23.25	47.25	1.291
	16	58.0	+7	65.0	23.05	28.75	2.75	0.831
	17	81.0	+0	81.0	31.75	28.25	21.00	0.890
March	17	66.0	-15	51.0	26.50	22.00	2.05	0.830
	30	20.0	+28	48.0	21.50	26.25	0.00	1.232
April	24	59.0	+15	74.0	31.05	40.25	2.25	1.262
	25	64.0	-14	50.0	21.05	28.05	0.00	1.325
May	20	23.0	+26	49.0	27.05	20.25	1.25	0.736
	25	23.0	+36	59.0	33.00	23.25	2.75	0.705
June	3	60.0	+12	72.0	27.00	41.75	3.25	1.546
	13	47.0	+21	68.0	24.05	31.50	12.00	1.286
	30	54.0	-17	37.0	22.75	11.05	2.75	0.505
July	12	13.0	+28	41.0	22.00	17.05	1.50	0.795
	28	24.0	+30	54.0	24.05	27.25	2.25	1.112
August	6	77.0	-25	52.0	21.25	24.25	6.50	1.139
	8	34.0	+25	59.0	20.05	31.05	7.00	1.536
Sept.	7	70.0	-20	50.0	22.00	28.00	0.00	1.272
	8	27.0	+26	53.0	24.05	26.00	2.50	1.061
Octob.	30	69.0	-21	48.0	22.25	21.05	4.50	0.955

T A B L E 7.
Dr. KEILL, Aged 39, A. D. 1717-18.

Months.	Weight		Food.		Urine		Perspiration.		Total		$\frac{P}{u}$
	lb.	oz.	oz.	oz.	Day.	Night	Day.	Night	Urine	Perf.	
July	154.09	74.62	18.54	17.98	18.98	15.95	36.52	34.93	0.957		
August	153.75	78.00	17.27	13.76	18.31	13.12	31.03	31.43	1.013		
September	154.60	76.00	16.00	18.00	18.73	13.25	34.00	31.98	0.941		
October	155.99	68.87	21.86	15.55	16.26	11.40	37.41	27.66	0.739		
November	155.85	64.61	22.02	17.17	18.18	10.50	39.19	28.68	0.732		
December	154.84	63.17	18.97	14.42	16.18	11.21	33.39	27.38	0.820		
January	153.77	63.12	23.68	14.51	17.72	11.53	38.19	29.25	0.766		
February	154.80	83.25	20.66	16.34	20.14	10.20	37.00	30.34	0.820		
March	155.93	94.33	28.58	19.07	16.18	10.70	47.65	26.88	0.564		
April	155.38	67.25	25.04	19.99	18.52	11.41	45.03	29.93	0.664		
May	155.96	84.67	16.58	21.10	22.46	13.37	37.68	35.83	0.951		
June	158.76	84.64	19.31	18.41	24.00	13.38	37.72	37.38	0.991		
Mean	155.31	75.21	20.71	17.19	18.80	12.17	37.90	30.97	0.817		
Summer	156.27	81.31	18.14	19.16	21.81	14.23	37.30	36.04	0.966		
Autumn	154.78	74.29	18.38	15.77	17.77	12.59	34.15	30.36	0.889		
Winter	154.82	63.63	21.55	15.37	17.36	11.08	36.92	28.44	0.770		
Spring	155.37	81.61	24.76	18.47	18.28	10.77	43.23	29.05	0.672		
Mean	155.31	75.21	20.71	17.19	18.80	12.17	37.90	30.97	0.817		

TABLE 8.

Mr. RYE, Aged 42, A. D. 1721-2.

Months.	Weight.	Food.		Urine		Total Urine	Total Perfp.	$\frac{P}{u}$
	lb.	oz.	oz.	Day.	Night	oz.	oz.	
May	196.56	102.41	21.96	18.76	40.72	61.10	1.500	
June	197.12	97.91	20.60	16.00	36.60	60.23	1.645	
July	197.62	94.23	19.60	15.73	35.33	58.83	1.665	
August	197.75	99.66	18.15	15.25	33.40	66.60	1.994	
September	196.50	96.54	19.54	15.73	35.27	57.77	1.638	
October	195.81	90.12	21.68	17.19	38.87	49.56	1.275	
November	197.69	87.93	21.87	17.77	39.64	47.37	1.195	
December	199.44	89.12	21.64	19.48	41.12	53.72	1.306	
January	197.25	94.73	20.73	18.80	39.53	50.13	1.268	
February	200.81	94.98	22.48	17.95	40.43	55.14	1.364	
March	200.47	100.53	20.93	19.07	40.00	56.87	1.422	
April	196.53	108.57	22.86	17.96	40.82	62.61	1.534	
Mean	197.80	96.39	21.00	17.47	38.47	56.66	1.472	
Summer	197.10	98.18	20.72	16.83	37.55	60.05	1.600	
Autumn	196.69	95.44	19.79	16.06	35.85	57.98	1.617	
Winter	198.13	90.59	21.41	18.68	40.09	50.41	1.257	
Spring	199.27	101.36	22.09	18.33	40.42	58.21	1.440	
Mean	197.80	96.39	21.00	17.47	38.47	56.66	1.472	

TABLE 9.
Dr. LINING, A. D. 1739-40.

Months.	Meat	Food. Drink.	Whole	+d	Urine.	Perfp.	Stool.	Whole	$\frac{du}{\pi u}$	$\frac{dp}{\pi p}$	$\frac{dp}{du}$	$\frac{\pi p}{\pi u}$	$\frac{p}{u}$	Ba- rom.	Heat by N's Scal	Moist of Air	Rain.
March	27.75	101.03	128.78	- 3.6	77.26	47.48	3.92	128.66	1.215	1.105	0.596	0.655	0.614	30.00	4.69	12	1.141
April	26.51	93.11	119.62	+ 39.6	64.55	52.74	3.65	120.94	1.329	1.275	0.794	0.830	0.817	30.03	6.56	7	1.092
May	28.94	100.43	129.37	+ 12.3	61.69	64.35	3.74	129.78	0.842	1.545	1.252	0.683	1.043	30.10	7.87	9	5.612
June	24.84	112.89	137.73	+ 69.0	57.55	78.33	4.15	140.03	0.789	1.631	1.705	0.825	1.361	30.09	8.81	10	4.648
July	26.80	120.34	147.14	+ 10.8	47.82	95.36	4.32	147.50	0.623	2.151	2.964	0.879	1.994	30.10	9.19	11	3.013
August	27.65	115.44	143.09	+ 4.8	60.79	77.80	4.65	143.25	0.725	2.115	1.717	0.592	1.280	29.60	8.44	12	7.301
Septem.	25.70	104.00	129.70	+ 111.9	43.95	84.58	4.90	133.43	0.862	2.039	2.142	0.906	1.924	30.11	8.06	12	3.200
October	26.31	80.00	106.31	- 72.9	52.34	44.76	6.78	103.88	0.722	1.049	1.003	0.700	0.855	30.22	4.50	12	1.257
Novem.	29.33	90.94	120.27	- 91.8	68.84	44.36	4.00	117.21	1.215	1.072	0.630	0.714	0.644	30.24	3.75	14	1.848
Decem.	29.50	100.44	129.94	- 48.0	77.55	46.69	4.10	128.34	1.310	1.468	0.632	0.563	0.602	30.11	1.87	10	2.736
January	25.94	104.35	130.29	- 58.0	80.18	43.68	4.50	128.36	1.618	1.369	0.532	0.859	0.545	30.09	2.45	18	4.492
Febr.	26.91	103.06	129.97	+ 6.6	85.59	41.08	3.53	130.19	1.723	1.379	0.449	0.568	0.480	30.25	2.62	16	3.135
Mean	27.18	102.17	129.35	- 19.3	64.84	60.10	4.35	129.30	1.081	1.516	1.201	0.731	0.927	30.08	5.73	11.9	3.289
Spring	27.05	99.06	126.11	+ 42.6	75.80	47.10	3.70	126.60	1.422	1.253	0.613	0.684	0.621	30.09	4.62	11.66	1.789
Summer	26.86	111.22	138.08	+ 92.1	55.69	79.35	4.07	139.11	0.751	1.776	1.974	0.796	1.425	30.10	8.62	10.00	4.424
Autumn	26.55	99.81	126.36	+ 43.8	52.36	69.05	5.44	126.85	0.770	1.737	1.621	0.733	1.318	29.97	7.00	12.00	3.919
Winter	28.26	98.58	126.84	- 197.8	75.52	44.91	4.20	124.63	1.381	1.303	0.598	0.712	0.595	30.15	2.69	14.00	3.025
Mean	27.18	102.17	129.35	- 19.3	64.84	60.10	4.35	129.29	1.081	1.517	1.201	0.731	0.927	30.08	5.73	11.91	3.289

T A B L E 10.

Seasons.	Italy.		England.		Cork.		Ireland.		South Carolina.	
	Food.	$\frac{p}{u}$	Food.	$\frac{p}{u}$	Food.	$\frac{p}{u}$	Food.	$\frac{p}{u}$	Food.	$\frac{p}{u}$
Spring	101.08	1.480	81.61	0.672	101.36	1.440	60.80	1.002	126.11	0.621
Summer	101.08	1.870	81.31	0.966	98.18	1.600	88.44	1.340	138.08	1.425
Autumn	101.08	1.480	74.29	0.889	95.44	1.617	84.18	1.286	126.36	1.318
Winter	101.08	1.090	63.63	0.770	90.59	1.257	61.86	0.908	126.84	0.595
Mean	101.08	1.480	75.21	0.817	96.39	1.472	86.31	1.141	129.35	0.927

Observations on the foregoing Tables.

OBS. 1. By Table 1, the Heat and Cold of the Air affect Perspiration and Urine in a different manner; Heat increases Perspiration and lessens Urine, and Cold on the contrary lessens Perspiration, and increases Urine. In *April* and *May* these two Discharges were nearly equal, only Urine exceeded Perspiration a little in *April*, and was exceeded by it a little in *May*. In the three Months *June*, *July*, and *August*, Perspiration exceeded Urine in the Proportion of above 1.5 to 1, or 15 to 10. Perspiration was greatest, and Urine least, in *August*. In *September* Perspiration lessened, and Urine increased. And in *October* and *November* they were nearly equal again, only Urine exceeded Perspiration a little in *November*. At the End of this Month I was interrupted, and hindered from carrying on the Experiments to the End of the Year, as I at first intended; but I repeated them for about ten Days in cold frosty Weather, and found

found that Urine then exceeded Perspiration as much as Perspiration exceeded Urine in Summer. Hence, supposing the Quantity of Food, and the Proportion of the Drink to the Meat, to have been the same in Winter as in Summer; the Proportion of Perspiration to Urine in Summer would have been greater than the Proportion of Perspiration to Urine in Winter, nearly in the Proportion of 18 to 10.

Obs. 2. By comparing the Food and the Discharges of the whole eight Months of Table 1, it appears that they were very nearly equal in a natural Day, taking one Day of that whole Time with another. For taking one Day with another, the daily Discharges exceeded the daily Food by only the 0.18th part of an Ounce, which is not a Quarter of an Ounce Averdupois. This daily Loss of Weight, though small, yet in the whole eight Months, or 244 Days, amounted to 43.92 Ounces; that is, to near two Pounds and a half. It is natural for Bodies to
fall

Of the Food and Discharges

fall away in Summer. *Sanctorius* says, that temperate Bodies are lighter in Summer than in Winter by about three Pounds, that is, by near thirty-eight Averdupois Ounces. A Change in the Weight of a Body at different Seasons of the Year, arises from an Inequality between the Quantity of Food and some of the Discharges in those Seasons. When a Body gains in Weight, the Food exceeds the Discharges; and when it loses in Weight, the Discharges exceed the Food. If therefore from a given Quantity of Food a Body be lighter in Summer than in Winter, the Discharges exceed the Food in Summer, and fall short of the Food in Winter. Consequently, in Summer Perspiration is more increased than Urine is lessened, and Perspiration more lessened than Urine is increased in Winter; whereas to keep a Body always of the same Weight, the Increase or Decrease of Perspiration must be ever equal to the Decrease or Increase of Urine, supposing the Quantity of Stool to be always the same. Hence, the Heat and Cold of the Air
first

first affect the Surface of the Body, and increase or lessen Perspiration; and afterwards, in some little Time, the Increase or Diminution of Perspiration is attended with an equal Diminution or Increase of Urine. Whence, the Body from a given Quantity of Food necessarily becomes lighter in Summer than in Winter, when the Air is hot than when it is cold; supposing the Quantity of Stool, which is a very small Discharge when compared with Perspiration and Urine, to be much the same in Summer and in Winter, as it is found to be in regular and temperate Bodies. This Property of Perspiration and Urine is caused by a Difference in the Motion of the Blood, at the Surface of the Body, and in its inward Parts, at the Skin and in the Kidneys, in these two different Seasons of the Year. The Heat of the Air acting on the Surface of the Body, swells the Vessels of the Skin, Blood-vessels, and secreting Ducts, and increases the Motion of their respective Fluids through them; but when the Motion of the animal Fluids, Blood, and secreted Humours,

Humours, is increased in one Part, it is lessened in the other Parts in healthful Bodies, when it is increased in the Skin, it is lessened in the Kidneys, by *Prop. 19. Anim. Oecon.* and therefore when Perspiration is increased or lessened by the Heat or Cold of the Air, Urine is lessened or increased equally, but not at the very same Time the Change is made in Perspiration; for all Changes of Motion in the animal Fluids are gradual; and consequently, when the Motion of the Blood is increased at the Skin, it will require some Time to be equally lessened in the Kidneys. Hence, from the Nature of the Blood's Motion, the Weight of a healthful Body is less in Summer than in Winter. But though the Weight of the Body, and the Proportion of Perspiration to Urine, be both changed by the Heat and Cold of the Air, yet will these Changes be very inconsiderable in a grown Body which uses but little Exercise, provided the Body be supported with a proper Quantity of good Food, and the Proportion of the Meat to the Drink be not less than

than that of 1 to 2. The Quantity of Food proper to preserve the Weight and Health of a grown Body, which uses but little Exercise, much the same at all Seasons of the Year, may be had from Table 1. p. 10.

Obs. 3. Perspiration and Urine are less affected by the Heat and Cold of the Air, and approach nearer to a Proportion of Equality in Summer and Winter, when the Quantity of the Food is less, and the Proportion of the Meat to the Drink greater, than when the Quantity of the Food is greater, and that Proportion is less. This appears from the Tables 1 and 2. In the Summer 1721, the daily Quantity of my Food at a Medium was 88.44 Ounces, and I judge the Proportion of my Meat to my Drink to have been about $\frac{3}{8}$, my Meat being about 24 Ounces in a Day, and my Drink 64.44. And in the Summer 1745, the daily Quantity of my Food at a Medium was 56.31 Ounces, and the Proportion of my Meat to my Drink was nearly $\frac{3}{5}$, my Meat being 21.33
E Ounces

Of the Food and Discharges

Ounces in a Day, and my Drink 34.98. In the Summer 1721 Perspiration was greater than Urine, in the Proportion of 1340 to 1000; and in the Summer 1745 it was almost equal to Urine, only less in the Proportion of 967 to 1000. And therefore Perspiration and Urine approach nearer to a Proportion of Equality in Summer and Winter, under a lesser Quantity of Food, and a greater Proportion of the Meat to the Drink, than they do under a greater Quantity of Food, and a less Proportion of the Meat to the Drink.

Hence we may conclude, that *Cornaro's* Perspiration and Urine were equal at all Seasons of the Year; for he did not take above half my Quantity of Food, and his Meat to his Drink was almost double of my Meat to my Drink, his being $\frac{6}{7}$, and mine about $\frac{1}{2}$; for $\frac{6}{7}$ is to $\frac{1}{2}$, as 12 to 7. By this small Quantity of Food, and great Proportion of his Meat to his Drink, this noble *Venetian* at the Age of forty, freed himself, by the Advice of his Physicians, from several grievous Disorders contracted
by

by Intemperance, and lived in Health of Body and great Chearfulness of Mind to above an Hundred. And I at the Age of sixty-four, by lessening my Food, and increasing the Proportion of my Meat to my Drink, by lessening my Drink about a third Part, and my Meat about a sixth, of what they were in the Year 1721, have freed myself from the Returns of some slight Disorders, and have greatly for my Age recovered the paralytick Weakness I was seized with about three Years ago. Hence we gather, that good and constant Health consists in a just Quantity of Food, and a just Proportion of the Meat to the Drink ; and that to be freed from chronical Disorders contracted by Intemperance, the Quantity of Food ought to be lessened, and the Proportion of the Meat to the Drink increased more or less according to the Greatness of the Disorders ; and that both the Quantity of Food, and the Proportion of the Meat to the Drink ought to be such as shall make Perspiration and Urine nearly equal at all Seasons of the Year. For Changes of the Seasons, as to

Of the Food and Discharges

Heat and Cold, are the most common Causes of Diseases; *Hipp. Aph. 1. Sect. 3.*

Before I proceed farther in the Observations, it is proper to give an Account of the Quality of my Food, and of the State of my Blood under this Reduction of my Food.

During the Year of Experiments from which I composed Table 2, I commonly eat four Ounces of Bread and Butter, and drank half a Pound of a very weak Infusion of Green Tea for Breakfast. For Dinner I took two Ounces of Bread, and the rest Flesh-meat, Beef, Mutton, Pork, Veal, Hare, Rabbet, Goose, Turkey, Fowl tame and wild, and Fish; and I generally chose the strongest Meats, as fittest, since they agreed well with my Stomach, to keep up the Powers of my Body under this great Diminution of my Food; I seldom took any Garden Stuff with my Meat, finding that it commonly lessened Perspiration, and increased my Weight. I drank four Ounces of Water with my Meat, and a Pound of Claret after I had
done

done eating. And at Night I eat nothing but drank twelve Ounces of Water, with a Pipe of Tobacco. This was my Course of Diet during that Year. Ever since that Time, I have lessened my Food a little; for in the following Year it was only 53 Ounces in a Day at a Medium; and this Year, as far as it is gone, which is seven Months, it is much about the same Quantity. This Diminution of my Food since the first Year, consisted chiefly in lessening the Quantity of Water.

To know the State of my Blood under this Reduction of my Food, I had 1855 Grains of Blood drawn from my Arm the 28th of *May* 1747; which, after standing twenty-four Hours, gave 1205 Grains of *Crassamentum* or red part, and 650 Grains of *Serum*. Let R denote the Weight of the red part, and S the Weight of the *Serum*, and then $\frac{R}{S}$ will be $\frac{1205}{650} = 1.854$. And the specifick Gravity of the *Serum* was 10364, the specifick Gravity of Water being 10000. The mean

Of the Food and Discharges

Proportion of the red part of the Blood to the *Serum*, that is $\frac{R}{S}$, and the specifick Gravity of the *Serum*, are, in the ordinary way of living, 1.400, and 10300 in healthful Men.

Notwithstanding this great Proportion of the red part of my Blood to the *Serum*; it flowed out of the Orifice in a full Stream, and with great Force; which argues that it was very fluid whilst it circulated in the Body. And that this greater than ordinary Proportion of the red part of my Blood to the *Serum*, has been of use to me now I use little Exercise, appears from my having been free two Years from a *sore Throat* and a *Diarrhœa*, Disorders I often had, though they were but slight and never confined me, when I lived more fully and used more Exercise; and it farther appears from my having had no Return of my convulsive and paralytick Disorder.

Now, as the Proportion of the red part of my Blood to the *Serum*, has increased
on

on increasing the Proportion of my Meat to my Drink, the Proportion of the red part of the *Serum* may in some sort be measured by the Proportion of the Meat to the Drink, that is, putting M for the Meat, and D for the Drink, $\frac{R}{S}$ may in some sort be measured by $\frac{M}{D}$. Hence, if the red part of the Blood bears too great a Proportion to the *Serum*, which is the Case of athletick Persons, and others who do not take a sufficient Quantity of Drink with their Meat, that Fault may be corrected, by lessening the Meat or by increasing the Drink. A young Man, who for a considerable Time had not drank with his Meat, and had a very florid Complexion and a scorbutick Eruption all over his Body, Arguments of too great a Proportion of the red part of the Blood to the *Serum*, was freed from the Eruption by drinking with his Meat, without any other Remedy. In the beginning of Fevers, the Proportion of the red part of the Blood to the *Serum* is greater, and at

the end of them lesser, than it is in Health; and the Change of this Proportion in Persons under these Disorders, is owing to their living wholly upon Drink and liquid Nourishment. And Bodies loaded with ferous Moisture, an Argument of too small a Proportion of the red part of the Blood to the *Serum*, have been freed from their Load, by abstaining wholly from Drink,

But though the Proportion of the red part of the Blood to the *Serum*, varies with the Proportion of the Meat to the Drink, yet there are other Things, besides the bare Quantities of Meat and Drink, which have a Share in settling the Proportion of the red part of the Blood to the *Serum*. For this Proportion is greater in Country People than Citizens, in Persons who use Exercise than in Persons who are inactive, and in Persons who live upon Flesh-meats and fermented Liquors, than in Persons who live upon Vegetables and Water. In short, this Proportion is increased by Things which dry
the

the Body and strengthen the Fibres, and lessened by Things of a contrary Nature.

Too great a Proportion of the red part of the Blood to the *Serum*, renders Bodies subject to inflammatory Fevers on taking Cold. For when Blood is drawn, and suffered to stand in the Air till it is cold, its red parts coalesce and form a *Coagulum*, leaving the *Serum* almost as fluid as Water. This shews that the red Corpuscles of the Blood attract one another more strongly, than they do the transparent Corpuscles of the *Serum*; and consequently, when the Blood happens to be greatly stock'd with red Corpuscles, and they are brought still nearer together by Cold, their attractive Forces may make them run together, coalesce, and obstruct the capillary Blood-vessels, in which I have shewn the Motion of the Blood to be exceeding slow. And this they will be the most apt to do in the Blood-vessels of those Parts which are most exposed to the Cold of the Air, as the Blood-vessels of the Throat, Lungs,
and

and *Pleura* are; whence arise Inflammations of those Parts.

Obs. 4. The Quantity and Quality of the Food, and the Proportion of the Meat to the Drink, being given, the Weight of a Body is less, and consequently its Discharges greater in dry Weather than in wet Weather. In the first fifteen Days of *May*, when the Air was dry, the daily Quantities in Ounces of my Perspiration and Urine, at a Medium, were 29.66 and 28.50; and in the nine following Days, when the Air was moist, the mean Quantities of Perspiration and Urine in a Day were 20.39 and 28.94 Ounces. I took above four Ounces of Food more in a Day the first fifteen Days, than I did the nine Days, and yet I gained two Pounds under the lesser Quantity of Food, and should have gained twice as much, had I not one Day lessened my Food above forty Ounces. Hence Perspiration was 9.27 Ounces in a Day less, and Urine only the 0.44 part of an Ounce greater, in

the nine wet Days, than in the fifteen dry Days; and consequently my Perspiration was above twenty times more lessened than my Urine was increased, by this Change of Weather from dry to wet. Hence Bodies, which live much in the same manner, as to the Quantity and Quality of their Food, and the Proportion of their Meat to their Drink, are lighter in dry Weather than in wet Weather, which may be thus accounted for. The Moisture of the Air moistens the Fibres of the Skin, and lessens Perspiration by lessening their vibrating Motion. For the moister the Fibres of the Skin are, the greater is their Density; and the greater their Density is, the less is the Density of the *Æther* lodged within them; and the less the Density of the *Æther* is, the weaker is its vibrating Motion; and the weaker the vibrating Motion of the *Æther* is, the weaker is the vibrating Motion of the Fibres caused by it. The vibrating Motion of the Air, *Æther*, and all elastic Fluids, is made less and slower by Water and watry Moisture. Hence musical

fical Sounds which are caused by a vibrating Motion in the Air, are lower and graver in a moist Atmosphere than in a dry one. The Sound of a musical Catgut String is made weaker and graver by wetting the String. And the Case is the same in the Nerves and Fibres of the Skin, watry Moisture apply'd to them lessens their vibrating Motions, and of consequence lessens Perspiration depending on those Motions. And when Perspiration is thus lessened by the Moisture of the Air, Urine by degrees is increased, but not equally, as appears by the above Instance; whence a Body from a given Quantity of Food, and a given Proportion of the Meat to the Drink, is necessarily heavier in wet Weather than in dry Weather.

Hence we learn, that to keep a Body of the same Weight in wet Weather as in dry, either the Quantity of Food must be lessened, or the Proportion of the Meat to the Drink increased; and both these will

will be done by lessening the Drink, without making any Change in the Meat.

By comparing this Observation with the two first Observations, it appears, that the Heat and Dryness of the Air change the Weight of the Body in like manner, for they both lessen it; and that the Cold and Moistness of the Air likewise change the Weight in the same manner, for they both increase it.

Obs. 5. By the first part of Table 3, when I lived more fully and used more Exercise, both Perspiration and Urine drawn off hourly from the Blood during the whole eight Months, were at a Medium greater in the Day than in the Night; Urine was greater almost in the Proportion of 3 to 2, and Perspiration greater in the Proportion of 5 to 4. And by the second part of the Table, when the Quantity of my Food was considerably less, and the Proportion of my Meat to my Drink greater, the Quantity of Urine drawn off from the Blood hourly
at

Of the Food and Discharges

at a Medium throughout the whole Year, was nearly the same in the Day and in the Night, in the Time of my being up and Time of my being in Bed, only it was something greater in the Night than in the Day; and the hourly Perspiration was remarkably greater in the Day than in the Night, in the Proportion of 1298 to 896. The Cause of the remarkable Inequality between the Quantities of Urine drawn off hourly from the Blood in the Day and in the Night in the first part of the Table, was too great a Quantity of Drink taken at Dinner. For when above a certain Quantity of Drink is taken with the Meat at Dinner, it is apt to run off quick by Urine, and so to make the Day's Urine exceed the Night's Urine more than if less Drink had been taken.

Now as Bodies are apt to be disordered by such a Quantity of fermented Liquors as passes off quick in pale Urine, it is rational to think that the Quantity of Drink ought not to exceed such a Quantity as can be drawn off from the Blood at much the same

same rate in the Day and in the Night ; which Quantity may be known by *Table 1.* p. 10. supposing the Drink to the Meat to be as 2 to 1, and the Body to be grown, and to use but little Exercise ; for the Quantities of Food set down in that Table, and the Proportion of the Drink to the Meat, may not be sufficient for growing Bodies, and Bodies which use much Exercise. And the Cause of the remarkable Inequality between the hourly Perspiration in the Day and in the Night, was a Difference in the Heat and Motion of the Blood in those two times. For the Heat and Motion of the Blood are always greater, from a greater Activity in the Soul, in the Day, than in the Night ; and they are likewise ever greater from the Food taken in the Day-time ; for the Pulse is always quicker after eating than before it, after a full Meal than after a spare one, and after a Meal of drier and stronger Food than after a Meal of Food that is moister and weaker. My Food was remarkably dry and of a strong Nature in the second Table, when compared with what it was in the first. And there-

therefore the remarkable Inequality between the mean hourly Perspiration in the Day and Night might be caused by a Difference in the Motion of the Blood in those Times.

Obs. 6. The mean hourly Perspiration in the Day exceeds the mean hourly Perspiration in the Night, in both parts of Table 3. And the Proportion of the mean hourly Perspiration in the Day to the mean hourly Perspiration in the Night, is greater in the second part of that Table than in the first, when the Food was less in Quantity and drier, than when it was more in Quantity and moister. In Summer and Autumn, the mean hourly Perspiration in the Day in Proportion to the mean hourly Perspiration in the Night, was 1.581 and 1.602 in the second part of the Table, when the Food was less and drier; and 1.302 and 1.292 in the first part of the Table, when the Food was more and moister. To make these Things more evident, I composed the following Tables from Table 3 and Table 7. These Tables, in the Columns $\frac{d u}{n u}$, $\frac{d p}{n p}$, exhibit the mean

mean Quantities of Urine and Perspiration drawn off hourly from the Blood in the Day, in Proportion to their mean Quantities drawn off hourly in the Night; and in the Columns $\frac{dp}{du}$, $\frac{np}{nu}$, exhibit the mean hourly Quantity of Perspiration in Proportion to the mean hourly Quantity of Urine, drawn off from the Blood in the Day and in the Night, in the several Seasons of the Year.

TABLE 11. taken from TABLE 3.

Seasons.	A. D. 1721.				A. D. 1744-5.			
	$\frac{du}{nu}$	$\frac{dp}{np}$	$\frac{dp}{du}$	$\frac{np}{nu}$	$\frac{du}{nu}$	$\frac{dp}{np}$	$\frac{dp}{du}$	$\frac{np}{nu}$
Winter					0.870	1.348	1.059	0.688
Spring					0.811	1.299	1.192	0.744
Summer	1.427	1.302	1.289	1.413	1.175	1.581	1.063	0.790
Autumn	1.315	1.292	1.277	1.290	1.002	1.602	1.241	0.776
Mean	1.371	1.297	1.283	1.356	0.966	1.457	1.139	0.749

TABLE 12. taken from TABLE 7.

Seasons.	$\frac{du}{nu}$	$\frac{dp}{np}$	$\frac{dp}{du}$	$\frac{np}{nu}$
Spring	1.341	1.697	0.748	0.583
Summer	0.947	1.533	1.202	0.742
Autumn	1.165	1.412	0.967	0.798
Winter	1.402	1.567	0.805	0.721
Mean	1.214	1.552	0.928	0.711

From the second part of Table 11, in which the Food was less in Quantity and drier, Perspiration exceeded Urine in the Day, and fell short of it in the Night, at all Seasons of the Year; whereas in the first Part of this Table, where the Food was more in Quantity and moister, Perspiration exceeded Urine both Day and Night, in Summer and Autumn. From the second Part of Table 11 it appears, that Perspiration and Urine are so contrived by the great Author of Nature, that when one increases the other lessens, both in different Parts of the natural Day, and in different Seasons of the Year; by which wise Contrivance, grown Bodies, which do not use much Exercise, and take a proper Quantity of good Food, and rightly proportion the Meat to the Drink, will be little affected with the sensible Qualities of the Air, and will continue of the same Weight, and in good Health, at all Seasons. Other Persons who live irregularly as to Diet, will scarcely be able, as appears from what is said in these Observations,

vations, to pass the four Seasons without Disorders.

Obs. 7. In this Observation I shall shew how Perspiration and Urine are affected by the Passions of the Mind. Anger and Joy increase, and Fear and Sadness lessen, both Perspiration and Urine. The Soul, which has great Power over the Body by virtue of the Æther, when it is made uneasy by the Passion of Anger, raises a strong vibrating Motion in the Æther within its *Sensorium*, which Motion is propagated through the Nerves to all Parts of the Body. This strong Motion in the Membranes of the Heart quicken its Contraction and Dilatation, and thereby quicken the Contraction and Dilatation of the Blood-vessels and secreting Ducts, and of consequence increase the Discharges of Perspiration and Urine, and that more or less in proportion to the Strength and Continuance of the Passion. I had one Instance of this in *November 1744*, in which both Urine and Perspiration were

Of the Food and Discharges

increas'd considerably by this Passion, in the Day and Night, from what they were the Day and Night before. Joy affects these Discharges in like manner as Anger. In the Passions of Fear and Sorrow, Perspiration and Urine are lessened, by a Diminution of the vibrating Motion of the *Æther*, occasioned by a Depression of the Power of the Will and Activity of the Soul, under those Passions. Hence Joy and Anger increase, and Fear and Sorrow lessen, Perspiration and Urine, and the Weight of the Body. And other Passions, as they partake of these, will affect the Discharges and Weight of the Body in like manner. That the Soul has an entire Power over the vibrating Motion of the *Æther*, may be gathered from its moving the Limbs with various Degrees of Velocity, and from its stopping and changing those Motions at pleasure; and likewise from that infinite Variety of musical Sounds caused by an infinite Variety of Vibrations in the Air and *Æther*, which can be produced by a fine Voice.

Consider-

Considering the great Power the Soul has over the Body in causing and regulating its Motions by the Æther, it may be proper to explain the great Law by which the Soul acts in this high Station. The Soul, by its sensitive and intelligent Nature, knows and feels the Wants and Disorders of the Body by the Sensations of Uneasiness and Pain, and perceives the Removal of those Wants and Disorders by the Sensations of Ease and Pleasure. Now since all Uneasiness may be reckoned Pain, and all Ease Pleasure, Pain and Pleasure are the two great Sensations of the Soul, which cause it to excite such Motions in the Æther of the Nerves, and thereby such Motions in the several Parts of the Body, as are fit to remove Pain and give Pleasure. Hunger and Thirst are the painful Sensations whereby the Soul is put upon raising such Motions in the Body as are necessary to procure Meat and Drink. And stretching the Stomach by too full a Meal creates a painful Sensation, whereby the Soul is urged to raise

Of the Food and Discharges

such Motions in the Body as are necessary to free it from the Load ; accordingly, if the Load arise from too much Drink, it often hurries it out of the Body in pale Urine, but if it arises from too much Meat, it commonly throws it off by Stool or Sweat, oftener by Stool in weak Bodies, and by Sweat in strong Bodies.

Obs. 8. The Proportion of Perspiration to Urine is increased by all those Exercises which increase the Motion of the Blood, and warm the Skin ; as is proved by Experiments in the *Animal Oeconomy*, p. 280.

Obs. 9. In *January* and *February* of Table 2, the daily Discharge by Stool at a Medium, was greater than in any of the other Months ; which was owing to a *Diarrhœa* which began the 30th of *January*, and continued four Days to the 3d of *February*. During this Discharge, which at a Medium was above two Pounds a Day, Perspiration and Urine were both lessened, and with respect to
their

their Quantities before this Discharge began, the Decrease of Urine was above double that of Perspiration. For before the *Diarrhæa* began, the daily Quantities of Perspiration and Urine were 27 and 30, and for the four Days of the Discharge they were at a Medium 25 and 24 nearly. The Decrease of Urine during the *Diarrhæa* was 6 Ounces a Day, and the Decrease of Perspiration 2; but $\frac{6}{30}$ is to $\frac{2}{27}$ as 162 to 60, that is as 2.7 to 1. The Matter of a *Diarrhæa* is supply'd by Blood-veffels, which are nearer to the Blood-veffels of the Kidneys than they are to the Blood-veffels of the Skin. On which account, when the Discharge by Stool is much increased by a *Diarrhæa*, both Perspiration and Urine are lessened, and Urine, from its being a nearer Discharge as being supply'd by nearer Blood-veffels, is lessened more than Perspiration. And when a *Diarrhæa* stops, Urine by the Stoppage of that Discharge is more increased than Perspiration. All this necessarily happens from the Nature and Laws of the Blood's Motion. The Truth

Of the Food and Discharges

of this appears, when a *Diarrhœa* or *Dysentery* is stopped by large Quantities of plain Water, Barley Water, Ptisan, or Chicken Water, drank warm; for as soon as the Pain abates, from the acrid Salts being diluted and carry'd off by Stool, the Water turns to the Kidneys, and pass'es off quick by Urine like Spa-water, which soon puts a Stop to the Discharge by Stool. One thing I must observe concerning the Discharge by Stool, that since I lessened my Drink, I have been much more costive than I was before. Costiveness generally attends dry Food in other Animals as well as Men.

Obs. 10. During the Course of Experiments from which I formed Table 2, the Morning Weight of my Body was continually changing, either increasing or decreasing. Sometimes it increased or decreased for some Mornings together, but then the whole Increase or Diminution never amounted to above 63 Ounces. In the beginning of *February* I gained 41 Ounces in six Days, having lost by a *Diarrhœa*

arrhæa 63 Ounces in three Days before. Table 6 shews some of the most remarkable Changes of Weight, whether Increase or Diminution, which happened to me that Year, together with the Months, Days, and Quantities of Food taken on those Days, in which the Changes happened. The Increase and Diminution of my Weight was commonly caused by an Increase and Diminution of my Food; tho' I have known my Weight increased by Fruit and Garden Stuff; when the whole Quantity of what was taken, was less than my ordinary Quantity of Food; which was the Case on the 30th of *June*.

If the Increase of Weight in a small Compass of Time, rise to above a certain Quantity, it will be apt to cause Disorders. I can bear an Increase of above a Pound and a half in one Day, and an Increase of three or four Pounds in six or seven Days, without being disordered; but am apt to think that I should suffer from an Increase of five or six Pounds in that Time. An Increase of Weight may
be

be carry'd off by lessening the Food, or by increasing the Discharges; and the Discharges may be increased either by Exercise, or by Evacuations procured by Art. By lessening the daily Quantity of my Food to 23 Ounces, I have lost 26 Ounces; by fasting a whole Day last Month, I lost 48 Ounces, having gained 27 the Day before; and by fasting a whole Day this Month I have lost 42 Ounces, having gained 7 Ounces the Day before. I am not able to use much Exercise at present, but have shewn from Experiments in the *Animal Oeconomy* how much a Body may lose by Exercise. Mr. Rye, who was a strong well-set corpulent Man of a sanguin Complexion, by a brisk Walk for one Hour before Breakfast, threw off by insensible Perspiration one Pound of increased Weight; and by a Walk of three Hours, he threw off two Pounds of increased Weight. I have lost by a spontaneous *Diarrhæa* two Pounds in twenty-four Hours; and Mr. Rye lost twice that Quantity in the same Time. A strong purging Medicine may lessen the
Weight

Weight of a grown Body about two or three Pounds, as I have shewn in the *Animal Oeconomy*. The best way to take off an Increase of Weight which threatens a Distemper, is either by Fasting or Exercise. But nothing, amidst a Variety of disturbing Causes, will be able so effectually to prevent such an Increase of Weight as shall cause a Distemper, as a very exact and regular Diet which can keep the Body of a right Weight, and prevent the Discharges from running into Irregularities, and Disproportions to one another.

Obs. 11. The specifick Gravity of my Urine in Table 2, was greater in the Night than in the Day, when I was in Bed than when I was up. Hence we learn, that Urine draws off more Contents from the Blood in Sleep than when Bodies are awake; and consequently, that natural Sleep is a very good Sign in Fevers, in which the Blood abounds more with Contents than it does in Health.

Obs.

Obs. 12. According to Dr. *Lining*, the sickly Months in *Charles-Town* at *South Carolina*, are *July*, *August*, and *September*. The Heat of the Air in each of those Months, is greater than its Heat in any other Month except *June*, as may be seen in *Table 9*; and the mean Heat in those three Months, is greater than the Meridian Heat of the Air in *England* in the Month of *July*, which by *Newton's* Scale is 6, in the Proportion of 8.63 to 6, or 863 to 600. And the mean Quantities of Rain, and Moisture in the Air, are each of them greater in those three Months than in the three preceding Months. Hence it seems very reasonable to conclude, that the Sickliness at that Season is caused by the Heat, and Moisture of the Air; for Bodies are more disposed to Putrefaction in a hot and moist Air, than in a cold and dry Air. Perhaps it may be asked, Since the Air in *South Carolina* is much hotter than it is in *England* and *Ireland*, and even in *Italy*, where
it

it is much hotter than in these Countries, why *Lining's* yearly Perspiration fell short of his yearly Urine, taking one Month of the whole Year with another, when in *Sanctorius* and *Rye* Perspiration exceeded Urine at all Seasons of the Year? To this may be answered, That though the Air is hotter in *South Carolina* than in the Countries I have mentioned, and on that account draws the Moisture of the Body more to the Skin, and increases Perspiration; yet both the Quantity and diuretic Quality of the Drink are much greater in *South Carolina* than in the aforesaid Countries, which may make it to pass off in a greater Quantity by Urine than by Perspiration; for a large Quantity of Drink of a diuretic Nature commonly passes off very quick by Urine. The Drink commonly used in Summer in *Charles-town* is a weak Punch thus made: Take of Water 2 Troy Pounds, of Sugar $1\frac{1}{2}$ Ounce, of recent Lime-juice $2\frac{1}{2}$ Ounces, of Rum $3\frac{1}{2}$. M. That which is used in Autumn and Winter is richer, having more Sugar and Rum, and less Acid. And if the
Moisture

Of the Food and Discharges

Moisture of the Air be greater there than in these Countries and in *Italy*, that may likewise contribute to lessen Perspiration, and promote Urine, by *Obs.* 4. *Keill's* Urine exceeded his Perspiration in all Months of the Year except *August*, in which Month Perspiration was nearly equal to Urine, only a little larger, as appears from Table 7. He often drank Punch, and was much on Horseback; both of which promote Urine much more than they do Perspiration, as appears by common Experience. Whether any other Cause contributed to produce this Effect in *Dr. Keill's* Discharges I can not tell, not having known him, and being wholly unacquainted with his Constitution. One Thing indeed, besides those already mentioned, I am satisfied, is much concerned in fixing the Proportion of Perspiration to Urine, and that is, the Force of the Heart in proportion to the attractive Power of the Kidneys. For a weaker Heart, or Kidneys stronger either naturally or by Irritation, may lessen the Proportion
of

of Perspiration to Urine, and on the contrary.

Obs. 13. The Sum of Perspiration and Urine drawn off hourly from the Blood, is greater in the Day than in the Night, in the Time of being up than in the Time of being in Bed, as appears from Table 3, and the Tables of *Keill* and *Lining*. Hence, when the Food is the same, as to Quantity, Quality, and the Proportion of the Meat to the Drink, and the Exercise likewise the same; those Bodies will be apt to gain most in their Weights which get most Sleep; which agrees with Experience. For much Sleep, much Food, and little Exercise, are the principal Things which increase the Weight of the Body, and make Animals grow fat. Consequently, if the Weight of the Body be too great for good and uninterrupted Health, it may be lessened by lessening the Sleep and Food, and by increasing the Exercise. And if the Body, on account of Age or other Infirmities, cannot use any Exercise of Consequence, and takes

Of the Food and Discharges

much the same Quantity of Sleep, its Weight must be lessened by lessening its Food, which may be done by lessening the Drink, without making any Change in the Meat; as I have found by Experience in myself. For my Exercise now being very inconsiderable, and of little Consequence with regard to the Change of my Weight, and the Quantity of sound Sleep being now six or seven Hours, which is much the same as formerly; the Diminution of my Weight ought wholly to be attributed to the Diminution of my Food, which consisted chiefly in lessening the Quantity of my Drink. I go to Bed early, and rise early, and have done so the greatest part of my Life. On the contrary, if the Weight of the Body be too little for good and uninterrupted Health, it may be increased by increasing the Food and Sleep, and by lessening the Exercise; and the Food must be increased chiefly by increasing the Drink and liquid Nourishment. For the Discharges are ordinarily less from Drink and a liquid Nourishment,

ishment, than from dry and solid Food; as I have shewn above.

Obs. 14. By Table 9, Dr. *Lining's* Weight was considerably less in hot Weather than in cold. He fell away in *April, May, June, July, August, and September,* and lost in those six Months above fifteen Averdupois Pounds, which he gained again, and near a Pound and a half more, in *October, November, December, and January.* It is to be observed that the sickly Months in that hot Climate, were some of those in which he fell away. See *Sanct. Aph.* 54, 55, 56. *Sect.* 2.

Obs. 15. There is but one Weight under which a grown Body can enjoy the best and most uninterrupted Health; and that Weight must be such, that Perspiration and Urine may be nearly equal at all Seasons of the Year; for by this Means the Body will be uniformly drain'd of its Moisture, the inward Parts by Urine, and the more superficial Parts by Perspiration, without any irregular and unnatural Dis-

G

charges,

Of the Food and Discharges

charges, and its Morning Weight will continue nearly the same at all Seasons of the Year. This Weight may be settled by the preceding Observations.

Obs. 15. “ In the Winter, when the
 “ Perspiration of an unexercised Person is
 “ only equal to the Urine, the Diet for
 “ twenty-four Hours ought not to exceed
 “ four Pounds or four Pounds and a half.
 “ In Summer the Diet may be six Pounds
 “ and a half, which may be carry’d off
 “ without the Help of Exercise when
 “ the Air is hot and dry. *Rye, Aph.* 25,
 “ 26. If the Meat and Drink of one
 “ Day be four Pounds and a half, the
 “ Perspiration of that Day will be two
 “ Pounds, the Urine two Pounds and
 “ five Ounces, and the Stool three Ounces.
 “ From about four Pounds of Food the
 “ Body daily returns to the same Weight;
 “ but from a lesser Quantity of Food the
 “ Weight of the Body lessens, and from
 “ a greater Quantity it increases.” *Keilley*
Obs. unius anni, p. 17. By comparing
 these Aphorisms and Observations with
 what

what I said above concerning my own Food, we may allow the Quantity of Food that is necessary to preserve a grown Person, who uses but little Exercise, constantly in good Health, to be pretty rightly determined in the Column F of Table 1. p. 10. For that Quantity of good Food will keep the Discharges of Perspiration and Urine equal, and the Morning Weight of the Body the same at all Seasons of the Year. Hence we may judge of the healthful Quantity of Food; for if the Quantity of Food be such as to make the Perspiration and Urine of a natural Day always nearly equal, and the Morning Weight of the Body always nearly the same, that Quantity is the truly healthful Quantity of Food for grown Bodies, who use but little Exercise.

Obs. 16. The Quantity of Food necessary to keep a grown Body in Health, will be better and more easily digested when it is so divided as to make the Meals equal, than when it is so divided as to make them very unequal. For instance; if

three Pounds of Food is to be taken in a Day at three Meals, it will be better to take a Pound at each Meal, than to take two Pounds at one Meal, and half a Pound at each of the other two; and if the same Quantity of Food is to be taken at two Meals, it will be better to take a Pound and a half at each Meal, than to take two Pounds at one Meal, and one Pound at the other. And as to the Distance between one Meal and another, it ought ever to bear some Proportion to the Largeness of the preceding Meal; for instance, if three Meals be taken in a Day, they may be taken at the Distance of eight Hours from each other, and at the Distance of twelve Hours if only two Meals be taken in a Day. But though the Food be equally divided between the Meals, yet there ought to be some Difference in the Distances between them; for Perspiration and Urine being drawn off from the Blood more slowly in the Night than in the Day, the Distance between Supper and Breakfast next Morning, ought to be greater than the Distance between Breakfast and Dinner,

Dinner, or between Dinner and Supper.
N. B. Persons who are subject to Disorders of the Head, ought not to take any Food at Night.

If the Quantity of Food be given, its Quality will cause a Difference in the Time of digesting; for instance, slimy and viscid Meats are longer in digesting in the Stomach than Meats of a contrary Nature; the Flesh of some young Animals is not so soon digested as the Flesh of the same Animals arrived at their full Growth; thus Veal and Lamb are not so soon digested as Beef and Mutton. A Man who took a Vomit every second Night for some Months, observed, that when he had taken Chicken for Dinner, he always threw it up undigested, but never threw up any of his Food undigested when he had made his Dinner on Beef or Mutton.

I now proceed to illustrate what has been said, by Observations drawn from Experiments made on other Animals.

OBSERVATIONS *drawn from Experiments made on other Animals.*

THE following Tables exhibit the mean Weights of the Bodies, Hearts, and Livers, of several Species of Birds and Fishes whose Names are set down in the first Column. The mean Weights of the Bodies, Hearts, and Livers, in Grains, are set down in the Columns W, H, and L; and the Proportion of the Heart and Liver to the Weight of the Body, and of the Liver to the Weight of the Heart, are set down in the Columns $\frac{H}{W}$, $\frac{L}{W}$, and $\frac{L}{H}$. Many of the Numbers are Means taken from ten Experiments, as were most of the Numbers of the Fishes; but some of the Numbers of the Birds were Means taken from a lesser Number of Experiments. The Heart was cleared from all Appendages of Fat and large Vessels, and was only the bare Muscle of the Heart; the Liver was without the Gall-Bladder; and in Birds the Weight of the Body was its Weight without Feathers.

TABLE

T A B L E I.

Wild Birds.	W	H	L	$\frac{H}{W}$	$\frac{L}{W}$	$\frac{L}{H}$
Cock green Linnets	352	5.9	7.9	$\frac{1}{39}$	$\frac{1}{44}$	1.34
Hen green Linnets	396	5.9	11.5	$\frac{1}{67}$	$\frac{1}{34}$	1.95
Cock Sparrows	427	6.3	19.5	$\frac{1}{68}$	$\frac{1}{22}$	3.10
Hen Sparrows	404	5.6	19.0	$\frac{1}{72}$	$\frac{1}{21}$	3.39
Cock Teals	7000	76.0	122.0	$\frac{1}{52}$	$\frac{1}{57}$	1.60
Hen Teals	5080	56.0	144.0	$\frac{1}{91}$	$\frac{1}{35}$	2.57
Mallards	16796	157.0	360.0	$\frac{1}{107}$	$\frac{1}{45}$	2.29
Ducks	15143	120.0	343.0	$\frac{1}{117}$	$\frac{1}{44}$	2.66
Mean of the Male Wild Birds	6144	61.3	127.3	$\frac{1}{100}$	$\frac{1}{48}$	2.08
Mean of the Female Wild Birds	5256	49.1	129.4	$\frac{1}{107}$	$\frac{1}{40}$	2.63
Mean of the whole Wild Birds	5700	55.2	128.3	$\frac{1}{103}$	$\frac{1}{44}$	2.32

T A B L E 2.

Tame Birds.	W	H	L	H W	L H	L H
Cock Pigeons young	3003	29	126	$\frac{1}{103}$	$\frac{1}{24}$	4.34
Hen Pigeons young	2908	27	128	$\frac{1}{108}$	$\frac{1}{23}$	4.74
Cocks, game, $2\frac{1}{4}$ Years old	25674	159	410	$\frac{1}{161}$	$\frac{1}{22}$	2.58
Hen Chickens, game, 9 Weeks old	6064	34	239	$\frac{1}{178}$	$\frac{1}{25}$	7.03
Drakes	17884	117	468	$\frac{1}{153}$	$\frac{1}{38}$	4.00
Ducks	16714	103	426	$\frac{1}{162}$	$\frac{1}{39}$	4.13
Barn-door Cocks, old	17504	84	360	$\frac{1}{208}$	$\frac{1}{49}$	4.28
Barn-door Hens, old	19315	63	508	$\frac{1}{307}$	$\frac{1}{38}$	8.06
Barn-door Cocks, young	21087	105	433	$\frac{1}{204}$	$\frac{1}{49}$	4.12
Barn-door Hens, young	21889	52	454	$\frac{1}{42}$	$\frac{1}{48}$	8.73
Cock Chickens	12380	47	412	$\frac{1}{263}$	$\frac{1}{30}$	8.77
Hen Chickens	10046	34	328	$\frac{1}{293}$	$\frac{1}{31}$	9.65
Cram'd Cocks	21065	92	673	$\frac{1}{227}$	$\frac{1}{31}$	7.31
Cram'd Hens	23990	60	1037	$\frac{1}{400}$	$\frac{1}{23}$	17.28
Mean of the Male Tame Birds	16942	90	412	$\frac{1}{88}$	$\frac{1}{41}$	4.58
Mean of the Female Tame Birds	14418	53	446	$\frac{1}{273}$	$\frac{1}{32}$	8.41
Mean of the whole Tame Birds	15680	72	429	$\frac{1}{218}$	$\frac{1}{36}$	5.96

T A B L E 3.

Round Fish.	W	H	L	$\frac{H}{W}$	$\frac{L}{W}$	$\frac{L}{H}$
Male Mackarels	7054	10.4	125	$\frac{1}{878}$	$\frac{1}{58}$	12.0
Female Mackarels	7579	11.2	223	$\frac{1}{877}$	$\frac{1}{34}$	19.9
Male Salmon Trouts	8925	11.4	164	$\frac{1}{783}$	$\frac{1}{54}$	14.4
Female Salmon Trouts	9074	10.6	131	$\frac{1}{856}$	$\frac{1}{69}$	12.3
Male Salmons	54614	62.5	795	$\frac{1}{873}$	$\frac{1}{89}$	12.7
Female Salmons	72095	70.4	1046	$\frac{1}{1024}$	$\frac{1}{87}$	14.8
Male Herrings	2489	2.2	11.2	$\frac{1}{1131}$	$\frac{1}{222}$	5.1
Female Herrings	2298	2.2	34.6	$\frac{1}{1045}$	$\frac{1}{88}$	15.7
Male River Trouts	4150	3.7	57	$\frac{1}{1122}$	$\frac{1}{73}$	15.4
Female River Trouts	3847	3.1	62	$\frac{1}{1241}$	$\frac{1}{82}$	20.0
Male Cods	90871	67.0	2070	$\frac{1}{1356}$	$\frac{1}{44}$	30.9
Female Cods	112044	84.7	3465	$\frac{1}{1323}$	$\frac{1}{32}$	40.9
Male Whittings	3281	2.3	102	$\frac{1}{1426}$	$\frac{1}{32}$	44.5
Female Whittings	3661	2.4	82	$\frac{1}{1525}$	$\frac{1}{44}$	34.2
Mean of the Male Round Fish	24483	22.8	475	$\frac{1}{1074}$	$\frac{1}{52}$	20.8
Mean of the Female Round Fish	30085	26.4	720	$\frac{1}{1139}$	$\frac{1}{42}$	27.3
Mean of the whole Round Fish	27284	24.6	598	$\frac{1}{1109}$	$\frac{1}{46}$	24.3

T A B L E 4.

Flat Fish.	W	H	L	$\frac{H}{W}$	$\frac{L}{W}$	$\frac{L}{H}$
Male Turbutts	26090	12.00	311	$\frac{1}{2174}$	$\frac{1}{84}$	25.9
Female Turbutts	36400	15.26	460	$\frac{1}{2333}$	$\frac{1}{79}$	30.1
Male Brets	10157	4.37	132	$\frac{1}{2324}$	$\frac{1}{77}$	30.2
Female Brets	14745	5.80	110	$\frac{1}{2542}$	$\frac{1}{134}$	19.0
Male Plaices	16505	6.25	232	$\frac{1}{2641}$	$\frac{1}{71}$	37.1
Female Plaices	20979	9.14	276	$\frac{1}{2295}$	$\frac{1}{76}$	30.2
Male Flounders	2178	0.93	33	$\frac{1}{2342}$	$\frac{1}{66}$	35.5
Female Flounders	4396	2.08	122	$\frac{1}{2113}$	$\frac{1}{56}$	58.6
Male Soles	6557	2.59	51	$\frac{1}{2532}$	$\frac{1}{129}$	19.7
Female Soles	6887	2.79	58	$\frac{1}{2468}$	$\frac{1}{119}$	20.8
Mean of the Male Flat Fish	12297	5.25	152	$\frac{1}{2357}$	$\frac{1}{81}$	29.0
Mean of the Female Flat Fish	16681	7.01	205	$\frac{1}{2380}$	$\frac{1}{81}$	29.2
Mean of the whole Flat Fish	14889	6.12	178	$\frac{1}{2367}$	$\frac{1}{81}$	29.1

OBSERVATIONS *on the Tables of*
Animals.

Obs. 1. **T**HE Weight of the Heart in proportion to the Weight of the Body, is greater in wild Birds than in tame Birds. The former use more Exercise, or move their Muscles more, than the latter ; and from the different Quantities of Motion in the Muscles of wild Birds and tame, we may see the Reason why the Weight of the Heart in proportion to the Weight of the Body, is greater in the former than in the latter. When a Muscle moves, it contracts and dilates alternately, in like manner as the Heart. In its Contraction it empties itself, as the Heart does in its Systole, pressing the Blood forward towards the Heart with greater or lesser Force, in proportion to the Strength of the Contraction. And in its Dilatation the Muscle fills again, as the Heart does in its Diastole, with Blood sent into it by the Force of the Heart.

Now

Now if many and large Muscles be moved, it is evident that much more Blood will flow to the Heart during the Time of their Motion, than did before in an equal Time when the Muscles were at Rest. And the Heart, to prevent an Oppression from an Overload of Blood, is made to quicken its Motion in order to send back the Blood as fast as it is thrown upon it by the Motion of the Muscles. Hence the Motions of the Heart and Blood are much increased by much Motion of the Muscles. On the contrary, when the Quantity of Motion in the muscular System is greatly diminished by the Inactivity of Bodies, the Force of the Heart and Motion of the Blood will likewise be diminished in the same Proportion. But Muscles which are much moved always increase in Magnitude, Weight, and Strength; and yet Bodies which use much muscular Motion, are seldom observed to grow fat or increase in Weight. And therefore the Weight of the Heart in proportion to the Weight of the Body will be greater in wild Animals than in tame, in
Bodies

Bodies which use much Exercise than in
Bodies which are inactive.

Hence Animals which use much Exercise have a greater Appetite, and require more Food than Animals which are inactive. For the Appetite is proportional to the Sum of the Discharges, and the Sum of the Discharges is proportional to the Motion of the Blood, by *Prop. 2.* Which Motion being greater in Animals which use much Exercise than in Animals which are inactive, the Appetite to Food will of Consequence be so too.

Obs. 2. The Weight of the Heart in proportion to the Weight of the Body, is greater in small Birds than in large ones, it is greater in a Sparrow than in a Goose. It is likewise greater in a Mouse than in an Ox, in a Child than in a Man, and probably greater in little Men than in large Men. This Proportion expressed by $\frac{H}{W}$, in a strong healthful Man, in a strong Child newly born, in an Ox, in a Hare,
and

and in a Mouse, has been found by Experiment to be equal to $\frac{1}{240}$, $\frac{1}{160}$, $\frac{1}{248}$, $\frac{1}{110}$, and $\frac{1}{127}$. Now since small Animals are generally more sprightly and active than great Animals, we may allow the Sprightliness and Activity of a Body to depend much upon the Proportion of the Weight of the Heart to the Weight of the Body.

Obs. 3. The Proportion of the Weight of the Heart to the Weight of the Body is less in fat Bodies than in lean. It is less in tame Birds than in wild Birds, and in cram'd Fowl than in Barn-door Fowl; and tame Birds are commonly fatter than wild Birds, and cram'd Fowl fatter than Barn-door Fowl. Butchers observe the same of Beasts, that the fatter a Beast is, the less is its Heart. Now as Animals grow fat, the Weights of their Bodies increase, and their Quantities of Blood and Weights of their Hearts lessen; for the Fat in the Membranes compresses the Blood-vessels, and lessens the Quantity of Blood; and as that lessens, the Weight of
the

the Heart lessens, the Weight of the Heart and Quantity of Blood always decreasing and increasing together. And therefore, the Weight of the Heart in proportion to the Weight of the Body is less in fat Bodies than in lean Bodies. The Motion of the Blood, as well as its Quantity, is less in fat Bodies than in lean, by *Cor. 3. Prop. 12. Anim. Oecon.* And hence we may account for fat Bodies having a lesser Appetite than Bodies which are lean; for the Appetite is regulated by the Sum of the Discharges, and the Sum of the Discharges is regulated by the Motion of the Blood, which Motion being less in fat Bodies than in lean, the Appetite will of consequence be so too.

Obs. 4. The Weight of the Heart in proportion to the Weight of the Body, is greater in the Males than in the Females of Birds both wild and tame; and the Difference of this Proportion in the Males and Females of wild Birds, is less than its Difference in the Males and Females of tame Birds. This Proportion in the Males
and

and Females of wild Birds, is $\frac{1}{100}$ and $\frac{1}{107}$ in Table 1; and in the Males and Females of tame Birds, it is $\frac{1}{138}$ and $\frac{1}{172}$ in Table 2. But $\frac{1}{100}$ is greater than $\frac{1}{107}$, and $\frac{1}{138}$ is greater than $\frac{1}{172}$; and the Difference between $\frac{1}{100}$ and $\frac{1}{107}$ is considerably less than the Difference between $\frac{1}{138}$ and $\frac{1}{172}$. Now by *Obs.* 1. the Weight of the Heart in proportion to the Weight of the Body is increased by Exercise. But the Males of Birds, from their greater Activity and Strength, use more Exercise than the Females. And therefore the Weight of the Heart in proportion to the Weight of the Body, is universally greater in Male Birds than in Female Birds, both wild and tame. This Proportion differs less in the Males and Females of wild Birds, than it does in the Males and Females of tame Birds: For the Males and Females of wild Birds probably exercise much alike, and are nearly equal in Fatness; but the Males and Females of tame Birds differ more both in their Exercise and Fatness; for the Males exercise more, and are remarkably less fat than the Females, as
 appears

appears from Observation. And therefore the Difference of this Proportion is less in the Males and Females of wild Birds, than in the Males and Females of Birds which are tame.

What has been said concerning this Proportion in the Males and Females of wild and tame Birds, holds equally true in the Males and Females of wild and tame Beasts.

Obs. 5. The Weight of the Heart in proportion to the Weight of the Body is much greater in Birds than in Fish. This Proportion at a Medium in the whole of wild and tame Birds, is $\frac{1}{16}$ by Tables 1, 2; and in the whole of round and flat Fish, it is $\frac{1}{32}$ nearly, by Tables 3, 4. Consequently, this Proportion is greater in the whole of the Birds, than in the whole of the Fish, in the Proportion of above 8 to 1. This great Difference in the Proportion of the Weight of the Heart to the Weight of the Body in Birds and Fish, is owing to the great Difference

H

of

of the Heat of the Blood in these two kinds of Animals, the Heat of the Blood being much greater in the former than in the latter. For a greater Degree of Heat in the Blood, will swell and enlarge the Capacity of the System of Blood-vessels, and make it bear a greater Proportion to the Weight of the Body than a lesser Degree of Heat. But the Force of the Heat in Animals is ever proportional to the Capacity of the System of Blood-vessels. And therefore the Force of the Heart with respect to the Weight of the Body will be greater in Birds than in Fish; that is, putting Æ for the Strength of the vibrating Motion of the Æther in the Heart, $\frac{\text{Æ H}}{\text{W}}$ will be greater in Birds than in Fish; by *Cor. 5. Prop. 8. Anim. Oecon.*

The Heat of the Blood of Animals, is caused by the volatile Acid of the Air mixing with it in the Lungs by means of Respiration; and is greater or lesser in proportion to the Quantity of this enlivening Spirit received into the Blood in a

I

given

given Time. And therefore, the Heat of the Blood is much greater in Birds than in Fish, as the former in a given Time receive much more of this enlivening Spirit into their Blood by Respiration than the latter.

Obs. 6. The Proportion of the Weight of the Heart to the Weight of the Body, is greater in round Fish than in flat Fish. The Mean of this Proportion in round Fish and flat Fish was $\frac{1}{1100}$ and $\frac{1}{2307}$ in Tables 3, 4. This Difference in the Proportion of the Weight of the Heart to the Weight of the Body in round Fish and flat Fish, may be owing to a Difference in their Respiration and Exercise. For round Fish often rise to the Surface of the Water to take in Air, a Thing rarely observed of flat Fish, which stay for the most part at the bottom of the Water.

Obs. 7. The Proportion of the Weight of the Liver to the Weight of the Heart is less at a Medium in wild Birds than in tame Birds, and in round Fish than in flat Fish. It is 2.32 and 5.96 in wild and

tame Birds ; and 24.3 and 29.1, in round and flat Fish. Hence we learn, that those Things which increase or lessen the Proportion of the Weight of the Heart to the Weight of the Body, do at the same time lessen or increase the Proportion of the Weight of the Liver to the Weight of the Heart. And consequently, when an Animal grows fat from much Food, much Sleep, and little Exercise, its Heart lessens and its Liver increases ; and on the contrary, when an Animal grows lean from little Food, little Sleep, and much Exercise, its Heart increases, and its Liver lessens. There is but one Weight of the Heart of a grown Body, under which Weight the Body enjoys the best and most uninterrupted Health, and that Weight is such as enables the Heart to supply the several Parts of the Body with just Quantities of Blood, such as can preserve the due Magnitudes of the Parts, and thereby hinder them from disturbing the Offices of one another. When the Weight of the Heart is too little for the Weight of the Body, the Liver enlarges beyond its just Magnitude,

Magnitude, and presses too much on the contiguous Parts, particularly on the Stomach, Intestines, and Diaphragm, and by that Pressure lessens the Capacities of the Stomach, Intestines, and Thorax, whence the Appetite grows less, the peristaltic Motion less, and Respiration more difficult and uneasy.

Obs. 8. The Weight of the Liver in proportion to the Quantity of Blood, is less at a Medium in wild Birds than in tame, and in round Fish than in flat Fish. For the Weight of the Heart and Quantity of Blood in all Animals are proportional to each other; and therefore, since by *Obs.* 7. the Weight of the Liver in proportion to the Weight of the Heart is less in wild Birds than in tame Birds, and in round Fish than in flat Fish; the Weight of the Liver in proportion to the Quantity of Blood, will be less in wild Birds and round Fish, than in tame Birds and flat Fish.

Rickety Children have larger Livers, and less Hearts and less Blood, than Children in Health. They have less Hearts and less Blood than Children in Health, from too

full and too gross a Diet, and too little Exercise; and they have larger Livers than Children in Health, from the Liver's always increasing in Weight, when the Weight of the Heart and Quantity of Blood lessen. In most chronical Diseases, the Liver is larger than in Health, as has been found by dissecting Bodies dying of those Distempers; and it is more than probable, that the Weight of the Heart, and Quantity of Blood in those Distempers, are both less than in Health; for most chronical Diseases arise from too much Food, and too little Exercise, both of which lessen the Weight of the Heart and Quantity of Blood, the first by causing Fatness, and the second by occasioning a Diminution of the Blood's Motion.

Hence when the Liver is grown too large by Intemperance and Inactivity, it may be lessened and brought to a healthful Magnitude by Temperance and Exercise. It may be emptied other ways by Art, but nothing can prevent its filling again, and consequently secure good and constant Health, but an exact Diet and Exercise.

Purging

Purging and Vomiting may lessen the Liver, and reduce it to its just Magnitude; but these Evacuations cannot prevent its increasing again, so long as Persons live too fully and use too little Exercise. This can only be done by lessening the Food and increasing the Exercise. And if sufficient Exercise cannot be used on account of Weakness or some other Cause, frequent Vomits may be substituted in its stead; for frequent Vomits as well as constant Exercise increase the Motion of the Blood. The Effects of Vomits on the Motion of the Blood, appear by the following Experiments. *By observing the Pulse of several Men after taking a Vomit, it has been found, that so soon as a Man begins to grow sick, his Pulse becomes low, quick, and irregular, and in the Action of Vomiting is often so low as not to be felt; that in the Intervals between the Vomits, the Pulse is still low and quick, but not near so low and quick as in the Action of Vomiting; and that after the Operation is over, the Pulse rises gradually, and in the Space of half an Hour or an Hour, becomes fuller than it*

was before the Vomit was taken. Hence we see the Effects of Vomits on the Motion of the Blood, they lessen that Motion during the whole Time of their Operation, and almost quite stop it in the very Action of Vomiting; and after the whole Operation is over, they increase the Motion of the Blood, so as to make it greater than it was before. And from these Effects of Vomits on the Motion of the Blood, we discover their great Usefulness in the Cure of many Diseases.

For Instance, Vomits stop Hæmorrhages from small Vessels. For when a Blood-vessel is opened, the Blood flows faster through that Vessel, and slower through all the rest of the Vessels, than it did before, by *Prop. 19 and 45 Anim. Oecon.* And therefore, all that is necessary to stop a Hæmorrhage from a small Vessel, is to stop the Motion of the Blood in that Vessel, and increase its Motion in all the other Vessels; and both these are done by Vomits, as appears by the foregoing Experiments.

The increasing the Motion of the Blood in all the other Vessels, will lessen the Motion

tion in the Vessel supplying the Hæmorrhage, and thereby effectually prevent a Return of the Discharge. For the same Reason, Vomits lessen immoderate Discharges of the Glands, and Ulcers; for they lessen the Motion of the Blood and Humours in the Parts affected, by increasing their Motion in all the other Parts. By increasing the Blood's Motion, repeated Vomits with a proper Diet I have found to be of great Service in dispersing scrophulous Tumours; which may be allowed, when it is considered that these Tumours are most incident to Children and young Bodies, the Motion of whose Blood is slow; and that they often disappear of themselves when Bodies are grown up, and their Blood has acquired a stronger Motion. When Obstructions arise from too languid a Motion of the Blood, Vomits are generally of use in removing them. And when the Motion of the Blood is too great, and Obstructions are formed by Cold, in the manner explained in p. 67. Vomits after large Bleeding will be of great Service in removing them. In
short

short Vomits, repeated according to the Nature and Obstinacy of the Disorder, are generally of Service in all Irregularities and Disproportions of the Motions of the Blood and other Fluids in different Parts of the Body. The Safety as well as Usefulness of frequently repeated Vomits, is evidently seen in Persons at Sea, and in Women with Child. Persons at Sea who are sick, and vomit much, are commonly the better for it: And frequent Vomiting in Women with Child, is of Service as it prevents Abortion. As all Muscles grow stronger by Exercise, so the muscular Coat of the Stomach grows stronger by Vomiting.

Obs. 9. Since fat Animals have larger Livers, and less Hearts and less Blood, than lean Animals; it is evident, that when a fat Animal grows lean, its Fat and Liver lessen, and its Heart and Blood increase; and on the contrary, that when a lean Animal grows fat, its Fat and Liver increase, and its Heart and Blood lessen. Hence, when a fat Animal grows lean, it
first

first loses its Fat, and afterwards, when its Fat is wasted, if it still goes on to lose in Weight, the Loss then falls on the Heart and Blood; and when a lean Animal grows fat, it first gains Flesh and Blood, and afterwards, when the Flesh and Blood are increased to certain Magnitudes, it then begins to get Fat. So that Animals on the Waste, first lose their Fat, and afterwards their Flesh and Blood; and Animals on the Gain first get Flesh and Blood, and then Fat. Farther, Animals on the Waste, first fall away in their Limbs, and afterwards in their inward Parts; and Animals on the Gain, first gain in their inward Parts, and then in their Limbs.

Obs. 10. The Weight of the Body under which an Animal has the greatest Strength and Activity, which I shall call its *athletic Weight*, is that Weight under which, the Weight of the Heart, and the Proportion of the Weight of the Heart to the Weight of the Body, are greatest. The Strength of an Animal is measured by the Strength of its Muscles, and the Strength
of

of its Muscles is measured by the Strength of its Heart ; and the Activity of an Animal is measured by the Weight of the Heart in proportion to the Weight of the Body ; that is, the Strength and Activity of an Animal are measured by H and $\frac{H}{W}$. And therefore, the Strength and Activity of an Animal will be greatest, when H and $\frac{H}{W}$ are greatest.

If the Weight of the Body of an Animal be greater than its *athletic Weight*, it may be reduced to that Weight, by Evacuations, dry Food, and Exercise. These lessen the Weight of the Body by wasting its Fat, and lessening its Liver ; and they increase the Weight of the Heart by increasing the Quantity and Motion of the Blood. And by lessening W and increasing H , they will soon reduce the Animal to its athletic Weight. Thus a Game Cock in ten Days is reduced to his athletic Weight, and prepared for Fighting. If the Food, which with Evacuations and Exercise, reduced the Cock to his athletic Weight

Weight in ten Days, be continued any longer, the Cock will not have that Strength and Activity which he had before under his athletic Weight; which may be owing to the Loss of Weight going on after he arrives at his athletic Weight, and then falling on the Heart, Blood, and Muscles, which must necessarily occasion a Loss of Activity and Strength. It is known by Experience, that a Cock cannot stand above 24 Hours at his athletic Weight, and that a Cock has changed very much for the worse in 12 Hours. When a Cock is at the Top of his Condition, that is, when he is at his athletic Weight, his Head is of a glowing red Colour, his Neck thick, and his Thigh thick and firm; the Day after, his Complexion is less glowing, his Neck thinner, and his Thigh softer; and the third Day his Thigh will be very soft and flaccid. Four Game Cocks reduced to their athletic Weights were killed, and found to be very full of Blood, with large Hearts, large Muscles, and no Fat. The athletic Weight of an Animal is a very dangerous Weight; by p. 67, and *Hipp. Aph. 3.*

Señ.

Señ. I. Fevers and Apoplexies are the Disorders, which commonly happen to Animals under or near their athletic Weights. Horses fed upon dry Food are much more subject to Fevers and Apoplexies than Horses fed upon Grass; and the former are much nearer their athletic Weights than the latter.

SINCE finishing the above, the following curious and useful Experiment, made on a Cow within eight or ten Days of dropping her Calf, was communicated by a Friend.

	Grains.
Cow alive —————	6461376
Lungs —————	42486
Liver —————	77964
Heart —————	24528

Diameter of the Aorta = 1.9 Inch.

Bag with Calf and Liquor—	735840
Calf —————	427488
Lungs —————	10950
Liver —————	9965
Heart —————	2628

Diameter of the Aorta = .8 Inch.

F I N I S.



BOOKS printed for J. NOURSE, at the
Lamb, against Katherine-Street, in the Strand.

I. **B**OERHAAVE's Medical Correspondence : Containing the various Symptoms of Chronical Distempers, the Professor's Opinion, Method of Cure, and Remedies. To which is added, His Practice in the Hospital of *Leyden*.

N. B. In this Collection are contained many Letters wrote originally in *Engliss*, to the Doctor, by Persons of Distinction, Gentlemen and Ladies, *etc.* with his Answers, such are marked * * * the rest are translated from the *Latin*. Price bound 4s.

II. A General and Compleat Treatise on all the Diseases incident to Children, from their Birth to the Age of Fifteen. By *John Astruc*, M. D. Regius Professor of Medicine at *Paris*, and Chief Physician to his present Majesty the King of *France*, &c. 8vo. Price 3s. 6d.

III. An Essay on Comparative Anatomy : Or, a Summary View of the most material Differences in the Structure of Animals ; in which the Descriptions are all taken from real Dissections, and transferred by Analogy to the Human Body, intermixed with many Practical Observations in Medicine and Surgery. In 8vo. Price 2s. 6d. sewed.

IV. A Discourse of Practical Chemistry. In which are contained all the Operations described in *Wilson's Complete Course of Chemistry*. With many new, and several uncommon Processes. To each Article is given, the Chemical History, and to most, an Account of the Quantities of Oils, Salts, Spirits, yielded in Distillation, &c. From *Lemery*, *Hoffman*, the *French Memoirs*, *Philosophical Transactions*, &c. and from the Author's own Experience. With Copper Plates. By *William Lewis*, M. B. R. S. S.

V. Academical Lectures on Fevers, in which those Disorders are fully treated of, and a Method of Cure subjoined


subjoined to each. Read in the Royal College at *Paris*. By *John Astruc*, M. D. *Regius* Professor of Medicine, and chief Physician to the present King of *France*. In one Volume 8vo. Price bound 5 s.


VI. *Hermippus Redivivus*: or, the Sage's Triumph over Old Age and the Grave. Wherein a Method is laid down for prolonging the Life and Vigour of Man. Including a Commentary upon an Antient Inscription, in which this great Secret is revealed, supported by numerous Authorities. The whole interspersed with a great Variety of remarkable and well attested Relations, in 8vo. Price 2s.


VII. Memoirs of the Royal Society; or, a New Abridgment of the Philosophical Transactions. Giving an Account of the Undertakings, Studies, and Labours of the Learned and Ingenious in many considerable Parts of the World; from the first Institution of that Illustrious Society in 1665, to 1740. In the Course of this Work, every thing is carefully extracted from the Originals, according to the Order of Time; the Latin Tracts Englished; the Terms of Art explained; the Theoretical Parts applied to Practice; and the whole illustrated with a great Number of Copper Plates. A Performance of general Use for the Knowledge and Improvement of Mathematicks, Natural Philosophy, Trades, Manufactures, Arts, &c. By Mr. *Baddam*. The Second Edition. Ten Volumes. 8vo. 1745.


VIII. The Elements of Natural Philosophy: Chiefly intended for the Students in the Universities. By *P. V. Musschenbroek*, M. D. Professor of Mathematicks and Philosophy in the University of *Leyden*. Translated from the Latin, by *John Colson*, M. A. F. R. S. Lucasian Professor of Mathematicks in the University of *Cambridge*. In Two Volumes, 8vo. Price 10 s.


IX. A Course of Lectures in Natural Philosophy. By the late *Richard Helsham*, M. D. Professor of Physick and Natural Philosophy in the University of *Dublin*. Published by *Bryan Robinson*, M. D. The Second Edition. 8vo. To which are added, several Problems by the Editor.











B699
H/5

