

## **Surgical papers / by Alex. Ogston.**

### **Contributors**

Ogston, Alexander, 1844-1929.

Ogston, Alexander, 1844-1929. Contribution to the surgery of bleeding vessels.

Ogston, Alexander, 1844-1929. On a new operation for removal of posterior adhesion of the iris.

### **Publication/Creation**

[Place of publication not identified] : [publisher not identified], [1872?]  
(Aberdeen : A King.)

### **Persistent URL**

<https://wellcomecollection.org/works/x5zv9e9>

### **License and attribution**

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection  
183 Euston Road  
London NW1 2BE UK  
T +44 (0)20 7611 8722  
E [library@wellcomecollection.org](mailto:library@wellcomecollection.org)  
<https://wellcomecollection.org>

*with Alex. Ogston's  
Compliments*

**S**URGICAL **P**APERS

BY

ALEX. OGSTON, M.D.



## A CONTRIBUTION TO THE SURGERY OF BLEEDING VESSELS.

---

IN a former paper (*The Lancet*, April 17th, 1869) I adduced a number of experiments bearing on the subject of the comparative strength of arteries secured by ligature, acupressure, and torsion, from which it appeared that ligature was by far the most secure method, resisting the highest column of mercury (114 inches) brought to bear upon it by the dynamometer, while acupressure and torsion resisted on an average respectively a column of 23·5 and 13 inches. These experiments were criticised at the time by Mr. Lawson Tait, but I still venture to affirm that they give a fair comparison of the value, as far as security goes, of the different methods. Excepting the assumed normal pressure of blood in living vessels, which was, I believe, stated at too high a figure, the only factor omitted by them is that of the elasticity of, and additional security furnished by, the portion of tissue included along with the artery between the pin and wire when acupressure is performed, and which protection accounts for the inner and middle coats of the vessels being, as is alleged, not cut through by this method.

These experiments have not been continued as originally intended. They are sufficient evidence as they stand, and

they gave place to a new line of experimentation undertaken with a view to improve upon, or add to, our present methods of arresting bleeding from cut vessels.


That none of our present modes is entirely satisfactory cannot be denied. Torsion seems, in some hands, to have given marvellously good results; but, having had no opportunity of witnessing it as applied regularly to large vessels, I can only say, under liability to correction by those who have tried and adopted it, that the small amount of internal pressure resisted by it forms a serious, although the only, objection to its use. In Aberdeen, on the contrary, we have abundant opportunity of studying acupressure, and the result of my own observation is to convince me that it cannot retain the place originally claimed for it in operative surgery. The one real advantage that it possesses is that it allows the wound to be freed from foreign bodies one or two days after operation—doubtless a very important matter, but which can be attained without its employment. The asserted frequency of primary union in wounds treated here by it, and the absence of suppuration in such wounds, are to a certain extent correct. In amputations of the mamma it yields first-class results, primary union and absence of pus being generally achieved; but in other amputations suppuration occurs just as usual, the statements to the contrary being due partly to the great enthusiasm of its advocates, and partly to a careful nurse removing, as far as possible, all traces of discharge before the visit of the surgeon.

Acupressure has several disadvantages. The pins frequently prevent complete closure of the wound at the time of the operation, and the edges have to be accurately adjusted at a subsequent period. The removal of the wires requires a

considerable degree of traction, and, in the employment of the ring-loop, is sometimes a matter of difficulty ; while, in at least one recent instance, an acupressed vessel in an amputation wound required, owing to secondary hæmorrhage, the subsequent application of the ligature. The estimate of acupressure I have formed is similar to that of many other surgeons who have given it a trial.

Considering, then, the superior strength of ligatured vessels, it would be a step in advance could the ligatures be applied in such a manner as to retain their superior security, and yet so as to allow of their being removed about the second day after operation, before the tendency to suppuration has commenced in the wound, and also so as to avoid snaring off the ends of the arteries and the tissue grasped by the forceps in the usual mode of applying them.

All these advantages are combined in the method I have now to propose—namely, that the bleeding vessels be secured by the knot depicted in Fig. 1, which if it be run close, and



the short end cut off close to the knot, can be removed by very moderate traction on the long end. The knot, it will be

perceived, is merely the "thumb or over-hand knot," similar to the first part of the knot ordinarily applied to arteries, with the single exception of a loop of the long end being employed instead of the undoubled cord. Its hold is perfectly sufficient when tied, but I shall afterwards come to discuss its security.






FIG. 2

In applying the knot to the flaps of an amputation wound, the following is the method I have adopted:—A piece of *the strongest* ligature silk, well waxed to diminish the risk of slipping after it has been tied, has one of its ends passed an inch or thereby through the eye of a curved needle. (A few such armed needles are prepared before the operation.) The bleeding vessel being seen, and its course above the orifice estimated, the needle is entered from the raw surface of the flap, a quarter of an inch on one side of the course of the vessel, and the same distance above the bleeding orifice; then passed down behind the vessel so as to include it between the needle and the raw surface of the flap; and, having crossed behind the vessel, is brought out a quarter of an inch on the other side of its track. The needle is next drawn

through, bringing a couple of inches of the thread with it, as in Fig. 3, and it may then be detached from the thread. A



loop of the long end, an inch or thereby in length, is then used to form, with the single short end, an ordinary "thumb knot," as in Fig. 1, or, if additional security is required, a



double knot, as in Figs. 2 and 4. The knot is closed upon the vessel and included tissue by traction on the loop and short end; the strands of the loop are placed close to each

other, so as to lie in contact throughout and offer no obstacle to removal when the long end is pulled ; and, finally, the short end is cut off close to the knot.

Such knots do not slip till traction is made on the long end, which ought to be brought out at the side of the wound next the original point of entry of the needle, and not doubled again over the vessel ; and, correspondingly, the loop is laid suitably directed away from the vessel in the direction of the needle when it emerged at the other side of the artery. In small arteries the single knot is amply sufficient ; in large arteries I use the double knot for the sake of security. The ligature is removed, on the second day after the operation, by gentle traction on the long end ; and the amount of force required for its removal, while too great to be readily effected by accident, is, notwithstanding, less, even with the double knot, than that needed to remove the wire where acupressure has been employed.

The advantages of this proceeding consist in the speedy removal of foreign bodies from the wound, and in the fact that no tissue is compressed or strangled by it save the portion lying in the very knot itself. The distal end of the vessel is still in connection with the living parts surrounding it, and its nutrition is not interfered with.

The strength of the method is another of its advantages. Vessels experimented upon in the manner given in detail in my former paper, resist, when secured with the single knot, from 24 to 70 inches of mercury ; and, on an average, 39 inches of mercury, or a pressure of 19 lb. to the square inch, as against 23·5 inches of mercury with acupressure, and 13 inches with torsion. The double knot resists the full column admitted of by my dynamometer—viz., 114 inches of mercury,

or 57 lb. to the square inch. I have tested in animals this application of ligature to cut vessels, and to vessels deligated in continuity, before using it in man, and the results were perfectly satisfactory. In man they have hitherto been equally so. I have at present a stump after Pirogoff's amputation treated in the above way, where the ligatures were removed on the second and third days, and which is healing with as small an amount of discharge as I ever saw obtained, even with acupressure.

The disadvantages are, that, as in acupressure, some tissue is included along with the vessel; and where structures such as important veins and nerves run parallel to the artery, damage might be done by their inclusion. Thus it is best suited for amputations; indeed, without further experience of it, I should hesitate to employ it in the deligation of an artery in continuity upon a human subject. I have had no opportunity of testing it in that most difficult of all vessels to secure, an interosseous artery in the angle of a flap amputation, but believe that, by laying hold of the bleeding orifice and the tissue surrounding it, and then proceeding, as in a flap, to pass the needle behind it through the grasped mass, it would answer as well as the ordinary ligature.—*The Lancet*, May 27th, 1871.



Digitized by the Internet Archive  
in 2019 with funding from  
Wellcome Library

<https://archive.org/details/b30571224>

## ON A NEW OPERATION FOR REMOVAL OF POSTERIOR ADHESION OF THE IRIS.

---

WHEN, in iritis, a single point of adhesion between the pupillary margin of the iris and the lens has been formed, there is great danger that the eye will remain ever after a damaged organ. Should the use of atropine (as is very usual) fail to destroy, while yet soft, this point of adhesion, the disease tends to go on from bad to worse, for the fixed part acts as a continual irritant, causes more or less severe relapses of the iritis, and leads to the increase of breadth of the adhesion, and to atrophic changes in the iris itself. Hence it happens that the usual result of such cases, when left to themselves, is that, finally, the pupil becomes totally adherent to the lens, the anterior capsule in and near the pupil becomes covered with the products of inflammation, and this capsulitis, interfering with the nutrition of the lens, leads to the formation of cataract, while the atrophic iris is bulged out at its periphery by the fluid in the posterior chamber, and forms a convex ring round the occluded pupil. The deeper structures of the eye usually suffer at the same time.

The treatment of adhesion between the iris and lens, posterior synechia, as it is called, is one of the least satisfactory parts of the ophthalmic practice. In iritis and corneitis, where these are so apt to be formed, the use of atropine

furnishes us with a means of preventing, in many cases, this annoying occurrence ; but, in spite of the most careful treatment, adhesions often happen, and constitute a germ of mischief tending continually to progress.

When the adhesions are broad and extensive, no better or surer means could be wished for than the performance of iridectomy, which radically cures the whole mischief at the expense of a slight deformity. When, on the other hand, the adhesions are few, and involve mere points of the pupillary margin, it would be greatly to be desired that we could break down synechiæ without producing deformity of the eye. The use of atropine commonly fails us entirely in such cases, and operation is the only means of attaining this end.

As a substitute for iridectomy in such cases, it has been proposed (Stellwag's *Augenheilkunde*, 1864, p. 179) to make a peripheral incision in the cornea with the usual iridectomy knife, then to introduce a peculiar flat hook, sharp at its concave side, and break down the adhesions by pushing with the convex side, or cutting them with the concave portion. It was hoped that this operation would succeed in an eye under the influence of atropine, but the proposal seems never to have found favour, and I am ignorant of what success, if any, is said to have attended it. The description of it would cause any experienced oculist to hesitate before even giving it a trial.

More recently (*Archiv für Ophthalmologie*, 1869, Bd. xv. Ab. I.) Dr. Passavant, in a letter to Professor von Graefe, suggests that limited adhesions of the iris to the lens should be broken down in the following manner :—The iris being under the influence of atropine, a peripheral incision, as for iridectomy, is to be made in the cornea opposite the site of the

adhesion. The knife is then withdrawn, and an iris forceps is introduced, which, seizing the iris near the pupil, and at a point corresponding to the adhesion, is made to drag the iris outwards towards the wound until the adhesion is seen to rupture. The forceps is then loosed from its hold on the iris by opening the blades, and withdrawn, the operation being thus completed. Dr. Passavant usually contents himself (p. 261) with the destruction of one adhesion at a time, and operates on another after an interval of a few days. He says —“In more than fifty such operations I have not met with a single injurious result.”


This paper of Dr. Passavant appeared so honestly written that a trial of his method was instituted in the next case that presented itself in the Aberdeen Hospital. In this, as in all the cases where I have tried it, the operation was followed by no bad result as regards the iris; but, though the adhesion was seen to tear, the contraction of the pupil, which invariably followed on the escape of the aqueous humour, allowed the two ends of the adhesion to lie so close to each other that they united again in spite of the free use of atropine, and by the time the corneal wound was healed, the same state of matters existed as before the operation, only the adhesion was not so broad as previously.

In both the above-mentioned operations the fault lies chiefly in their necessitating the escape of the aqueous humour, and so permitting the contraction of the pupil which follows to bring the ends of the separated adhesions so close that they reunite. They show, however, with what impunity the iris may be manipulated, and that iritis is not even a usual result of careful and well-judged interference.

The following mode of operating on cases of limited

adhesions, for which iridectomy is too severe a measure, is based upon two facts—first, that the iris is a structure sufficiently tough to resist very decidedly attempts to transfix it, and will be pushed before the point of a sharp instrument so as even to tear from its attachments rather than allow the sharp point to pass through it; and second, that if the aqueous


FIG. 1.



humour be not permitted to escape, an instrument may be introduced into the anterior chamber, and the iris manipulated without causing contraction of an atropised pupil.

The pupil being put well under atropine, and the patient being under chloroform or bichloride of methylene, a spring speculum is fixed on the lids, and the operator fixes the bulbus with a pair of forceps held in the left hand. The situation of the adhesion is now well studied, and, if necessary, concentrated artificial light is employed to render it distinct. This latter, however, is necessary only with feeble daylight or opacities of the cornea. A needle (Fig. 1) is now introduced through the cornea, near its sclerotic margin, at a considerable distance on one side of the adhesion, and in such a direction that, when pushed onwards, it crosses at right angles that radius of the lens in which the adhesion is lying (Fig. 2). The needle is pushed onwards into the anterior chamber, and so guided by the operator that its point becomes engaged in the pupillary margin of the iris at the root of the adhesion. The simple contact of the needle point with the iris is sufficient for this, and no attempt must be made to bury the needle in the tissue. By a gentle lever action of the needle point towards the periphery of the iris, the corneal wound being the fulcrum, the adhesion is broken down by being stretched

FIG. 2.



and ruptured, and the needle being slowly withdrawn from the eye, the operation is completed.

The needle I use is depicted in Fig. 1, and its point is rounded off a little, so as not to be acuminate.

There seems to be no reaction of iris or cornea consequent on this slight interference; on the contrary, I have found the dilatation of the pupil always keep the torn ends of the adhesion apart under the use of atropine until absorption had taken place.

The simplicity of the proceeding obviates the necessity of detailing cases, but the following example will show how readily it succeeds where Passavant's method fails:—

*Case.*—Jas. T., aged 18, admitted into the Aberdeen Hospital on October 25, 1869, with nebulous opacities of both corneæ and synechia posterior iridis on both eyes. The right pupil was nearly totally adherent, the left pupil being fixed

only at two points—viz., above and to the outer side. On October 28, 1869, the external adhesion of the left iris was separated by Passavant's method. There was no subsequent reaction, but the adhesion reunited to nearly its former extent, in spite of the assiduous employment of atropine. On November 23, an iridectomy was performed on the right eye with complete results. On December 16 the external adhesion on the left eye was broken down by the operation above described, and on December 21 the upper adhesion was similarly operated on, a round movable pupil being secured, and the boy was dismissed cured on December 27, with orders to return after six weeks to be treated for the haziness of the corneæ.—*Medical Times and Gazette, May 28th, 1870.*