

An appendix to the Observations upon Mr. Pott's General remarks on fractures / [Thomas Kirkland].

Contributors

Kirkland, Thomas, 1722-1798.

Pott, Percivall, 1714-1788. General remarks on fractures.

Publication/Creation

London : T. Becket & P.A. de Hondt, 1771.

Persistent URL

<https://wellcomecollection.org/works/eb4zz6n5>

License and attribution

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>

3134 5
A N

A P P E N D I X

T O T H E

O B S E R V A T I O N S

U P O N

Mr. POTT'S GENERAL REMARKS ON
FRACTURES.

By THOMAS KIRKLAND, M. D.



Printed for T. BECKET and P. A. DE HONDT,
in the Strand, 1771.

5

PREFACE

TO THE

READER

BY

JOHN R. KELLY

THOMAS KIRKLAND

AND

JOHN

NEW YORK: PUBLISHED BY
JOHN R. KELLY

A N
A P P E N D I X
T O T H E
O B S E R V A T I O N S
U P O N
M^r. POTT'S GENERAL REMARKS ON
FRACTURES.

A N

P R E F A C E

TO THE

READER

BY

THE GENERAL REMARKS ON
THE FUTURE

P R E F A C E.

THE very great ease, with which we reduced the dislocated hip, mentioned in the following Appendix, induced me to re-examine the nature of this injury; and the result of my enquiry is now offered to the consideration of the gentlemen of the faculty, as several circumstances of consequence seem to have escaped the notice of those, who have written on this subject. Even our latest writers have not in this case extended their improvements beyond preserving the muscles in a state of non-resistance; and yet, whoever examines into this matter, by proper experiments, will see whether this is the chief point we ought to have

have in view in the cure of the injury immediately under consideration; whether the bone, being readily replaced, does not often depend upon another cause; and whether *wholly* attending to the best means of making a considerable extension, may not frequently lead us from the principal object to which we should pay attention, and from the true intentions of cure, where considerable extension prevents reduction. In the following case all possible advantage was at first taken of a relaxed state of the muscles, but making the extension as *commonly practised, and as advised by most writers*, probably prevented success.

But no true idea can be formed about this matter, unless recourse be had to experiment upon recent subjects, and the skeleton, from whence a true judgment may be derived concerning the method pointed out, as seemingly the most advisable to be pursued, when called to relieve this accident.—But time, and the experience

ence of many surgeons alone, can finally determine whether the practice recommended is right, because this accident so very rarely happens. And, indeed, practice is always best established by the concurring testimony of different men, if, instead of implicitly copying from other writers, they give us their own experience and observations.—It therefore affords me infinite pleasure to find, that what I have said concerning the cure of compound fractures has met with the approbation of several eminent surgeons; and I have been informed by three gentlemen of great experience and undoubted veracity, who have seen much of this business in the course of many years practice, that they never lost a patient on account of a compound fracture, nor ever amputated on this occasion; which I know nearly corresponds with the success of several other surgeons.—I am certain, within the last fifteen or sixteen years, I have talked with more than fifty surgeons of experience about this matter; and though it be true

that *almost all die* of bad compound fractures, in large and crowded hospitals, when amputation is not performed, yet it is equally certain, that in the country, when the hospital mode of practice is not followed, they *almost all recover*, under the same predicament, without amputation; from which difference of success, obvious and useful conclusions may be drawn, without entering more minutely into the matter. And we shall only observe, that accidents of this kind most commonly happen to people employed in hard labour, who in London are generally sent to an hospital; but in the country (where there is not an hospital) they are all private patients, which gives the country surgeons more frequent opportunities of seeing the event of these cases in a pure air, and *under a different method of proceeding*, than falls to the share of those gentlemen, who are situated where business of this kind is for the most part, carried on in a place set apart for the purpose. Surely then, the united observations of
many

many country surgeons, uninfluenced in their practice by general bad success in these accidents, is worthy of notice ; especially as speedy amputation has been advised in every bad compound fracture without limitation. And do not those then, who throw cold water upon well-intended information from this quarter, without making proper enquiry, seem desirous of having it believed, that they had rather see all his Majesty's subjects hopping about upon one leg, than that their own opinion should in any instance be thought to be erroneous ?

A N

APPENDIX, &c.

SINCE my Observations upon Mr. PORT's General Remarks on Fractures were published, I have had a patient with a dislocated thigh, from which, I think, I have more satisfactorily learned the nature of this accident than from all I have ever read upon the subject; and I am desirous of adding this case as a supplement to my letter on dislocations, in hopes that it will put an end to the general use of violent extension, which has commonly been practised and recom-

B 2 mended

mended indiscriminately, as a dernier resort on this occasion.

A pauper, fifty years of age, of the parish of Ashfordby, in Leicestershire, was thrown from a waggon-load of coals, and part of the coals falling upon his hip, dislocated his right thigh-bone, the head of it passing inward towards the foramen ovale of the *os pubis*.

Mr. PEAK, a sensible surgeon of Melton-Mowbray, was called in to his assistance, who at once discovered the nature of the accident; but the part being swelled, he bled the patient, applied the usual remedies to abate the swelling, and then, after putting the muscles in a relaxed state, with *a force commonly used* on this occasion, he attempted the reduction, but without success; and not having before seen a dislocated thigh, he desired my assistance.

Accordingly, I saw this man five days after the accident, when the tumor had subsided

subsidied in consequence of proper applications, which gave me a better opportunity of examining the nature of this injury, than often happens, because the patient was very lean.—The knee straddled very much outward, the muscles on the inside of the thigh, which arise from the os pubis, were upon the stretch, from the head of the bone lying under them; and this thigh was longer by two inches than the other. The great trochanter did not protuberate in its usual manner, but was upon a level with the outer edge of the acetabulum, and a nick betwixt them might easily be felt with the finger, upon moving the thigh.—From all which it was evident, that the end of the dislocated bone lay upon and across the cavity from whence its head had escaped.

After securing the patient upon a bed on his right side, and putting the thigh in a right angle with the body (the knee being bent, see p. 56.) my servant and
another

another man only, extended it by a towel fixed above the knee, upon which we made a lever of the thigh bone in the manner already directed, (*ibid.*) and then pushed the ankle outward, upon which the head of the bone slipped into its place without noise, and with the *utmost ease*. Whereas, had a greater extension been made, in all probability reduction had not taken place; for though in general, reducing dislocations consists in making a lever of the dislocated bone, *after* it is brought by extension upon a level with the part from whence it has receded, yet, in inward dislocations of the thigh, there is always an exception to this rule, the head of the bone *being already before its cavity*, in which case, much extension is not only unnecessary, but tends to prevent reduction.

Perhaps the plates prefixed may explain this matter, but if any one acquainted with the structure of the joints before dissection,

section, will take into his hands the os innominatum and a thigh bone separately, he will soon have a tolerable knowledge concerning an inward dislocation of the thigh, by applying the upper edge of the great trochanter, (*a*) just within or upon the outer edge of the acetabulum coxendicis, (*b*) so as to make the head of the bone pass half way over the foramen ovale, (*c*) and he will also see, (by placing the thigh bone so as to make about a right angle with the vertebræ, and then making a lever of it) what a little extension must of course be necessary in the reduction. Nay, it may even be doubted, whether any extension will in this case be required, because making a lever of the bone, raises the great trochanter out of the socket, and prevents it from catching upon the cartilage, and being any impediment in restoring the bone to its proper place.

Again, if the great trochanter is pushed into the middle of the socket, so that the

head of the bone entirely covers the foramen ovale, it will be evident by withdrawing it half an inch or an inch, that the head of the bone will be quite clear of the brim of the acetabulum; and that this very small extension is all that is necessary to replace it. Nor can he omit observing, when the foot turns inward, and the head of the bone towards the upper part of the os pubis (*d*), or when the foot turns outwards, and the head of the bone down towards the ischium * (*e*), that it would be best to pull the ankle gently inward or outward, as either case requires so as to place the head of the bone, if possible, over the foramen ovale, in a line crossing the middle of the acetabulum coxendicis (from *b* to *c*) before any pressure is made with the knee, so as to give it an opportunity of passing into its place, over the notch leading into the acetabulum.

* In these three cases, instead of a nick, there will be a cavity felt at the outer edge of the acetabulum, as the great trochanter will be about the middle of the socket.

where

where, from the ligament yielding, it will meet with the least resistance. And even supposing a rent in the burfal ligament, it will not, I imagine, by embracing the neck of the femur, be any impediment to this position taking place, as it is most probable, that the head of the bone escapes out of the socket in this direction, and that it accidentally inclines up or down afterwards.

But my experiments have led me to think, that the head of the bone never passes out of its place *through a narrow opening* made by it in the burfal ligament, but that the ligament is commonly torn off from the greatest part of the neck of the femur, and no ways confines it in recent dislocations; and when the bone is reduced, I find a great part of the ligament is pressed by it into the bottom of the socket, from whence it is soon raised again and resumes its proper place upon

C

moving

moving the limb forward and backward.

I well remember, in endeavouring to dislocate the thigh in a man I dissected some years ago, the burfal ligament began to tear off at the neck of the femur ; but as I could not readily break the round ligament, I finished the displacing of the bone with my knife, and looked at that time upon the nature of the laceration as accidental. Since which, (not having frequent opportunities in the country, of dissecting the human subject) I have had recourse to experiments upon the joints of brutes, in which I find the separation of the burfal ligament, always begins in the same manner before the round ligament tears from its insertion, and when the head of the bone gets at liberty, and rushes out of its place, this separation is enlarged by the capsular ligament tearing off from the whole, or from the greatest part of the

the

the circumference of the neck of the bone.
—I even tried whether I could force the head of the bone through an opening near the size of a sixpence, made in the burſal ligament without tearing it off at the neck, but without effect, and the laceration in this manner has been ſo conſtantly uniform, that I cannot help thinking but it is always the caſe in complete diſlocations; nor does it appear to me that it can be torn in any other manner: and does not the only inſtance we have upon record of the diſſection of a recent diſlocation, ſeem to confirm this opinion? In that caſe “the capſular ligament was completely
“torn off from the whole circumference
“of the neck of the humerus*.”

Now, I have always thought that PETIT and Dr. HUNTER were right in their opinion concerning the rupture of the capſular ligament in diſlocations; but till I had made theſe experiments (ſuppoſing

* Med. Obſ. and Inq. vol. ii.

a rent through the ligament) I was at a loss to account for the ease, with which a shoulder was reduced, after being drawn from under the pectoral muscle into the axilla; but finding all or the greatest part of the ligament torn from the neck of the dislocated bone, it was easy to see how this came to pass, and why the rupture of the ligament does not prevent reduction, when timely and properly attempted*. Nor had I less pleasure in discovering the manner, in which the lacerated ligament was prevented from interrupting the cure, when forced under the bone into the socket; for after removing part of the muscles, and dislocating a thigh, I placed the bone in different positions, to see in which it would most easily return into its cavity, but in all of them, it pressed in a great part of the burfal ligament before it; but, upon moving the bone in a rotatory motion, the ligament instantly be-

* See POTT on Dislocations, p. 115.

gan to rise, and in a few turns resumed its natural situation †; and many experiments have since convinced me, that this was not accidental, but certain, after which,

† The separation of the ligament at the neck of the bone, and resuming its former place, is not to be confined to joints which are composed of ball and socket; but it happens in a similar manner to those which are joined by ginglymus, except (for example) if the lower end of the humerus is pushed forward upon the radius and ulna, the ligament is mostly separated from these bones.—There are Surgeons who believe, that the burfal ligament is not so frequently lacerated in dislocations, because some of them are weak, distractile, and constantly moistened; but I never could dislocate a joint completely without lacerating the ligament in the manner described, whether the dislocation was made gradually, or by the vis percussiois; and if their belief is not founded upon observation made in dissecting recent dislocations, or upon experiments made on dead bodies, it can give no satisfaction in an affair which can only be determined by matter of fact.—BOERHAAVE thought and believed that the head of the femur could not be dislocated by external violence, *because* of the depth of the socket and the strength of the ligaments in this articulation, and he induced others to be of the same opinion; but undeniable testimonies have since evinced that their belief was ill-founded: from whence we may infer how little regard ought to be paid to theories of any kind.—It is therefore desired, that those who have frequent opportunities of dissecting the human subject, will try whether the laceration does not always happen nearly in the same manner.

in all probability, the ligament very readily adheres to the bone again ; for, does not the periosteum readily re-unite to the broken bone in simple fractures ? and we see inflamed membranes soon adhere to the part with which they are in contact. Nay, even where a bone has been exposed to the air, and covered with matter, yet the pericranium (for instance) will readily adhere to it ; and this adhesion is more likely to take place where the air has not admision, and nature meets with no interruption.—Should not we therefore preserve the limb in a state of rest, for some time after reduction to forward this coalition ?—And may not a neglect, in this particular, from imagining the burfal ligament unhurt, have prevented the re-union of the lacerated parts, and rendered the limb liable to be displaced by every slight accident ?

However if my experiment should have deceived me in this matter, and a rupture
in

in the burfal ligament ſhould ever prevent the head of the thigh bone from being placed in the direction I have recommended before reduction, ſhould not the knee be firſt preſſed outwards a little, to raiſe it above the brim of the ſocket, then making a moderate extenſion in the ſame direction, and afterwards a lever of the thigh (ſee p. 56.) will not reduction take place? Whereas, if theſe ſteps are omitted, may not the head of the bone, if it paſſes over, be locked faſt upon the brim of the acetabulum, (*d* or *e*) and either prevent or render reduction very difficult?

But when the diſlocated part lies already upon the notch (as was the caſe of our patient) the knee ſhould not be twiſted any way by moving the ancle, before the head of the bone is pushed oppoſite the acetabulum; otherwiſe it will be preſſed upon the very part we have given a caution to avoid, and make a

greater extension necessary than is consistent in this operation; for I apprehend it will succeed much better if the bone *slips close over* the ligament into its place.

But in all these *inward dislocations*, placing the thigh in a right angle with the body, seems to me from experiment and from the ease with which our patient was cured, to be a material point; not wholly in taking off the resistant state of the muscles (because *very little* extension is only wanting) but principally in placing the bone in the most convenient situation for passing into the middle of the socket whereas, when the extension is made with the thigh in a right line with the body, the head of the bone is not raised or set at liberty from the brim of the acetabulum, but is drawn sliding down in contact with the parts upon which it lies without any possibility of the extension affording the least assistance, unless the head of the femur is situated in the groin obliquely

obliquely above its cavity, and then this kind of extension gives us an opportunity of pushing it rudely into its place at the time it is passing by the breach in the side of the acetabulum.—And what is worse, this is done by guess work, there being no criterion fixed when it is the most proper time to make a lever of the bone. Whereas, pressing the ankle outward, and bringing the thigh in a right angle with the body, places the bone in a proper situation previous to reduction, and at once with certainty determines this matter without the torture consequent on great extension.—And if we are to suppose the burfal ligament embracing the neck of the femur from a rent through it, will not the consequence of extension with the thigh in a right line with the body, be horrible?

But if instead of the bone lying towards the groin, it lies *strait across* the foramen ovale, will not great extension with the

D

thigh

thigh in a right line pull it below its cavity, and frustrate the reduction? For though, when in this situation, it has been reduced, while this kind of extension has been made; yet, if there is any dependence to be had on anatomical enquiries, it appears that the reduction of the bone must have taken place upon making a lever of the thigh, before the extension had done much mischief, (owing probably to the resistant state of the muscles) for it will be evident to every one who makes the experiment, that when the head of the femur lies in the supposed position, the making a lever of the thigh bone is more likely to reduce it *without any*, than *with much* extension.—Nay, even when the thigh makes a right angle with the body, great extension will keep the head of the bone at a distance from the part, into which it should enter, and when it ceases, it is ten to one but the patient will be in statu quo: and upon a fair enquiry into this matter, I am
apt

apt to think it will be found, that all the difficulties which have arisen in reducing inward dislocations of the thigh, have been owing to violent extension with the limb in a wrong position *.

If the head of the thigh bone is placed upon, or near the brim of the outer edge of the acetabulum, I apprehend we shall have a clear idea of an *outward* dislocation of the hip, which sometimes happens; how it came to pass that the reduction was so easily accomplished in the cases I have mentioned, by only moving the thigh, and pressing with my hand

* WISEMAN, who wrote *from practice*, and seems to have reduced dislocations of the thigh with great ease, says, "There is no need of so great extension in an inward luxation of the thigh." And in "tough bodies" where (he thinks) the strongest extension is required, he orders only one man to *pull at the leg*.—Probably, the extension he made obliquely towards the sound leg, with the patient laid upon a table flat on his back, being gentle, did neither good nor harm, but the reduction more likely took place, simply from his making a lever of the bone, when (in pushing the knee up towards the belly) it came into the situation we have recommended. See WISEMAN.

against the trochanter, (see p. 53 and 54.) and that the same ease must always attend this kind of luxation.—But when the head of the bone is dislocated beyond the acetabulum under the glutæi muscles, the general rule of making extension sufficient to bring it upon a level with the brim of its cavity, takes place. Nor, in this case, would the thigh making a right angle with the body, be a proper position, because when the head of the bone is under the glutæus minimus it is above the part into which it must return, and drawing it downward will give a better opportunity of putting it into its place; and accordingly we see Mr. WHITE * reduced this kind of dislocation with the *greatest facility imaginable* by placing the patient upon his back, and making his extension with the thigh in a strait direction; but if a relaxed state of the muscles, and bringing the head of the bone directly across the middle of its cavity should seem

* Cases in Surgery, p. 124.

preferable, I apprehend the patient should be placed on his sound side, and the extension made with the thigh forming an obtuse angle with the body, which is probably the position to which the limb will incline after the accident.

R E C A P I T U L A T I O N.

From the whole of what has been said, does it not appear, that the injury, of which we have been treating, may easily be remedied, when proper steps are taken? For reducing an inward dislocation of the thigh, seems to consist only in placing the patient in the manner we placed the pauper, or (which appears to me a more convenient and eligible method for performing the operation) sitting upright upon the edge of a table, or a bed of a convenient height, on which he must be properly secured.—This being done, if the head of the bone lies directly across the foramen ovale, (which I apprehend
may

may be known by the inner condyle of the femur being neither elevated nor depressed) a gentle extension, and making a lever of the bone seems to be all that is necessary.—But if it does not lie already in this direction, it seems right to push the ancle outward or inward, so as to bring the knee into its natural position, or (if this cannot be done) to press the knee a little outward before these steps are taken.—And lastly, when the great trochanter is felt with its natural protuberance, the ancle may be pushed outward, to direct the head of the bone into the bottom of the socket, if it is not immediately forced in by the action of the muscles.—In the same situation too, when the thigh is dislocated *outward*, but not beyond the brim of its acetabulum, we see (p. 56) it may easily be reduced, by only moving it, and pressing upon the great trochanter.—But when the head of the bone is under the glutæus minimus, a greater extension must be made with the

thigh in a different direction *, before the great trochanter can be pushed inward, or the knee outward with advantage; but in each kind of dislocation it seems necessary to move the thigh about immediately after reduction, the sooner to restore the ligament to its proper place, and hasten the complete recovery of the patient: and to this we may add, that it is not necessary that all the different movements we have recommended should be performed in the same instant of time, as if we were playing legerdemain, which has been advised, but they may regularly succeed each other, with considerable advantage both to the operator and the patient.

* See above.

F I N I S.

A P P E N D I X

though in a different direction, the great trochanter can be pushed in-
ward, or the knee outward with either
tact; but in each kind of dislocation it
seems necessary to move the thigh
immediately after reduction, the object
to restore the ligament to its proper place
and hasten the complete recovery of the
patient; and to this we may add, that it
is not necessary that all the dislocations
movements we have recommended should
be performed in the same instant of time,
as it were playing legions, which
has been advised, but they may regularly
succeed each other with considerable ad-
vantage both to the operator and the pa-
tient.

See above.

F I N I S