

Hand atlas of human anatomy / by Werner Spalteholz.

Contributors

Spalteholz, Werner, 1861-1940.
Barker, Lewellys F. 1867-1943

Publication/Creation

Philadelphia ; London : Lippincott, [1930?]

Persistent URL

<https://wellcomecollection.org/works/d8quuwxc>

License and attribution

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>

WERNER SPALTEHOLZ
HAND ATLAS OF
HUMAN ANATOMY

VOL. II.

D. IX


20/5

DA, C(2)



22500302927

Med
K8227



51608

HAND-ATLAS

OF

HUMAN ANATOMY

BY

WERNER SPALTEHOLZ

PROFESSOR OF ANATOMY IN THE UNIVERSITY AT LEIPZIG

SIXTH EDITION IN ENGLISH

VOL. II

REGIONS, MUSCLES, FASCIÆ, HEART, BLOOD-VESSELS

PHILADELPHIA AND LONDON

J. B. LIPPINCOTT COMPANY

2 015 659
ANATOMY, Texts : 20 cent

DA, C (2)




TRANSLATIONS OF SPALTEHOLZ' HAND-ATLAS OF HUMAN ANATOMY
HAVE APPEARED IN ENGLISH, ITALIAN, AND RUSSIAN
AND A SPANISH EDITION IS IN PREPARATION


ALL RIGHTS RESERVED

WELLCOME INSTITUTE LIBRARY	
Coll.	WelMOMec
Coll.	
No.	05


PRINTED IN GERMANY




332. Regions of the head and neck.




333. Regions of the body, viewed from in front.




334. Regions of the body, viewed from behind.




335. Regions of the male perineum.



336. Regions of the female perineum.



337. Muscles of the head and face,
viewed somewhat from the right.



338. Muscles of the head and face, viewed from in front.

Musculus epicranii. Form: flat, broad. Position: covers the roof of the skull.

Posterior part, fleshy: **mm. occipitales** (see Fig. 347). Origin: lineae nuchae supremae oss. occipitalis. Insertion: galea aponeurotica. Action: draw the galea backward.


Middle portion, tendinous: **galea aponeurotica**, large, flat tendon, inserted between mm. occipitales and frontales; loosely and movably united with the periosteum, very firmly connected with the cutis.

Anterior portion, fleshy: **mm. frontales** (see also Fig. 337). Origin: galea aponeurotica. Insertion: skin of the eyebrows crossed by fibres of mm. orbiculares oculi. Action: draws eyebrows upward, wrinkles forehead transversely; move the galea aponeurotica.


Innervation: nn. faciales.

M. procerus, unpaired. Origin: ossa nasalia. Insertion: skin between and above the eyebrows. Action: draws skin at the root of the nose downwards, causes transverse folds at the root of the nose. Innervation: nn. faciales.

M. quadratus labii superioris (see also Fig. 337). Form: flat, triangular. Position: lateral from the nose; origins partly covered by m. orbicularis oculi. *Caput angulare*. Origin: proc. frontalis and margo infraorbitalis maxillae. Insertion: skin of the ala nasi and of the sulcus nasolabialis. *Caput infraorbitale*. Origin: near under the margo infraorbitalis. Insertion: sulcus nasolabialis. *Caput zygomaticum*. Origin: facies malaris oss. zygomatici. Insertion: sulcus nasolabialis. Action: draws ala nasi upwards and upper lip upwards and outwards, widens nasal opening and deepens the sulcus nasolabialis. Innervation: n. facialis.




339. Muscles of the right eyelid, viewed from in front.




340. Muscles of the right eyelid, viewed from behind.

M. orbicularis oculi (see also Figs. 337 and 338). **Form:** flat, elliptical. **Position:** in the eyelids and at their periphery. **Pars orbitalis.** **Origin:** crista lacrimalis anterior maxillae, anterior limb of the lig. palpebrale mediale (upward and downward). The fibres of the upper medial portion run to the skin of the medial part of the eyebrow. The rest of the fibres surround the eyelids in flat, concentric arches, intercross above with the fibres of the m. frontalis and interweave with one another lateralward from the external angle of the eye. Fibres running in a vertical direction join up laterally, and below fibres which radiate from the medial and lateral part into the skin of the cheek. **Action:** draws the eyebrow medianward and downward, the skin of the cheek upward and medianward; helps to close the lids; forms radial folds in the skin of the lateral angle of the eye. **Pars lacrimalis [Horneri].** **Origin:** crista lacrimalis posterior oss. lacrimalis; it runs, partially crossed, behind the lig. palpebrale mediale and the tear-sac to the medial angle of the eye, spreads out thinly within the lids (corresponding in extent to about that of the tarsi) and go to a thin, horizontal band of connective tissue, **raphe palpebralis lateralis** (see also Figs. 925 and 929), which extends from the external angle of the eye to the lateral margin of the orbit (see p. 817). In the lids it forms the **pars palpebralis**. **Action:** widens tear-sac, closes eyelids. **Innervation:** n. facialis.



341. Muscles of the region of the mouth, viewed somewhat from the right side.

(Platysma, mm. risorius, quadrati and orbicularis oculi have been removed.)



342. Muscles of the region of the mouth, viewed from behind.

(The tissues about the mouth have been loosened from their bony substratum and the muscles have been dissected out from the posterior surface.)

M. corrugator (see Figs. 339 and 340). **Form**: flat, narrow. **Position**: near the root of the nose. **Origin**: pars nasalis oss. frontalis. **Insertion**: extending lateralward and a trifle upward, skin over the middle of the eyebrow. **Action**: causes vertical folds at the side of the root of the nose. **Innervation**: n. facialis.

M. nasalis. **Form**: flat, triangular. **Position**: lateralward from and below the ala nasi, partly covered by the m. quadratus labii superioris. **Pars transversa**. **Origin**: jugum alveolare of the upper canine tooth, a few strands also in the skin at the furrow of the nasal ala. **Insertion**: to the dorsum of the nasal cartilage spreading out by a thin tendon to the muscle of the other side. **Pars alaris**. **Origin**: jugum alveolare of the upper lateral incisor tooth. **Insertion**: lateral and lower margin of the ala nasi. **Action**: draws ala nasi downward, narrows nasal opening. **Innervation**: n. facialis.

M. depressor septi. **Form**: flat, quadrangular. **Position**: below the nose, just medianward from the preceding muscle. **Origin**: jugum alveolare of the upper lateral incisor tooth. **Insertion**: septum mobile nasi and posterior part of the nostril. **Action**: draws the septum mobile downward, narrows nasal opening. **Innervation**: n. facialis.

M. risorius (see Fig. 337). **Form**: flat, triangular; very variable. **Position**: lateral from the angle of the mouth, upon the platysma. **Origin**: fascia parotideomasseterica. **Insertion**: unites at the angle of the mouth, or somewhat below it, with the m. triangularis. **Action**: helps to draw the angle of the mouth lateralward; causes dimple. **Innervation**: n. facialis.

M. zygomaticus (see also Fig. 337). **Form**: oblong, flat-cylindrical. **Position**: lateralward from and above the angle of the mouth, separated from caput zygomat. m. quadrati labii sup. by the v. facialis anterior. **Origin**: facies malaris oss. zygomatici. **Insertion**: skin of the angle of the mouth; a great deal of the fibres go over into the m. orbicularis oris. **Action**: draws the angle of the mouth lateralward and upward. **Innervation**: n. facialis.

M. caninus (see also Fig. 337). **Form**: flat, triangular. **Position**: in the fossa canina, covered by the m. quadratus labii superioris; comes to the surface between the latter and the m. zygomaticus. **Origin**: fossa canina. **Insertion**: with converging fibres, partly to the skin of the angle of the mouth; partly the fibres go over arch-like into the lower lip. **Action**: draws the angle of the mouth upward; the muscles of the two sides together lift the lower lip and help to close the mouth. **Innervation**: n. facialis.

M. triangularis (see also Figs. 337 and 348). **Form**: flat, triangular. **Position**: below the angle of the mouth; covers the foramen mentale and partly the m. quadratus labii inferioris. **Origin**: just above the basis mandibulae below the foramen mentale. **Insertion**: with converging fibres to the angle of the mouth and, in largest part, arch-like into the upper lip. Single bundles sometimes connect the muscles of the two sides below the chin, upon the platysma, as the m. transversus menti (see Fig. 348). **Action**: draws the angle of the mouth downward; the muscles of the two sides together draw the upper lip down and help to close the mouth. **Innervation**: n. facialis.


M. quadratus labii inferioris (see also Figs. 337 and 348). **Form**: flat, quadrangular. **Position**: below the lateral part of the mouth, for the most part covered by the m. triangularis; it covers the foramen mentale. **Origin**: the upper part is a direct continuation of the platysma, the lower part arises from the basis mandibulae. **Insertion**: skin of lower lip. **Action**: draws the lower lip lateralward and downward. **Innervation**: n. facialis.

M. incisivus labii superioris (see also Fig. 344). **Form**: flat, triangular, narrow. **Position**: above the angle of the mouth, covered by the m. quadratus labii superioris, medially of the m. caninus, at the upper margin of the m. orbicularis oris. **Origin**: jugum alveolare of the upper lateral incisor tooth. **Insertion**: arch-like to the angle of the mouth, interwoven there with the other muscles. **Action**: draws the angle of the mouth medianward and upward. **Innervation**: n. facialis.

M. incisivus labii inferioris (see also Fig. 344). **Form**: flat, quadrangular, narrow. **Position**: below the angle of the mouth, covered by the m. quadratus labii inferioris, laterally of the m. mentalis, at the lower margin of the m. orbicularis oris. **Origin**: jugum alveolare of the lower canine tooth, connecting with the m. mentalis. **Insertion**: to the lower margin of the m. orbicularis oris, interweaving with it. **Action**: draws the angle of the mouth medianward and a trifle downward. **Innervation**: n. facialis.

M. orbicularis oris, a muscle-plate situated in the lips which owes its formation to the fact that the muscles going to the mouth opening, at the angles of the mouth, completely or partly radiate out into the upper or lower lip. The fibres run in the lips partly intercrossing, for the most part transversely, at the angles of the mouth arch-like between the two lips and are attached in part to the external skin and to the mucous membrane. In this way a sort of ring muscle is formed. **Action**: if the latter contracts alone it closes the mouth and points it. **Innervation**: nn. faciales.

M. mentalis. **Form**: thick, cylindrical, short. **Position**: on the chin. **Origin**: jugum alveolare of the lower medial incisor tooth. **Insertion**: with diverging bundles in front and obliquely downward into the skin of the chin, interweaving there partly with fibres of the other side. **Action**: lifts the skin of the chin and wrinkles it. **Innervation**: n. facialis.




343. Fascia temporalis and m. masseter,
viewed from the right side.

Fascia parotideomasseterica (see Fig. 337) covers the lateral (and medial) surface of the glandula parotis as a thick membrane, the m. masseter as a thinner membrane, and becomes lost medianward below the m. zygomaticus, platysma and m. risorius.

Fascia temporalis, very dense, covers the fossa temporalis externally, arises from the linea temporalis superior, divides below into two layers, is separated usually by fat from the m. temporalis and goes to the medial and lateral surface of the arcus zygomaticus and the os zygomaticum.

M. masseter. *Form:* flat, quadrangular, thick. *Position:* immediately upon the lateral surface of the ramus mandibulae; behind partly covered over by the glandula parotis, in front and below by the platysma. It consists of two layers which are separated from one another distinctly behind only; the external layer is the longer and broader. *Origin:* external layer: anterior and middle part of the arcus zygomaticus (tendon of origin jagged, reaching far downward); internal layer: middle and posterior part of the arcus zygomaticus; both layers also from its inner surface. *Insertion:* external layer (multipinnate): obliquely backward and downward to the lateral surface of the angulus mandibulae and its periphery; internal layer: downward and somewhat forward to the lateral surface of the ramus mandibulae. *Action:* lifts the lower jaw. *Innervation:* n. massetericus (V. 3).



344.

M. buccinator and m. temporalis, viewed from the right side.


M. incisivus labii superioris

Mm. zygomaticus, caninus, triangularis
(cut through)M. incisivus
labii inferioris

(The arcus zygomaticus has been sawed away at its origins, the joint of the lower jaw having been opened at the same time. An outer layer of the m. temporalis is cut off.)

M. temporalis. *Form:* flat, triangular, broad and thin above, narrow and thick below, bipinnate. *Position:* covered by the fascia temporalis and the arcus zygomaticus; it fills up the fossa temporalis completely. *Origin:* nearly the whole extent of the fossa temporalis and medial surface of the fascia temporalis and (a few strands) the arcus zygomaticus. *Insertion:* the anterior fibres run vertically, the posterior at first almost horizontally, than markedly curved; all the fibres converge to the processus coronoideus mandibulae and descend deep on its medial surface. The tendon reaches inside in a jagged form as far as half the height of the muscle. *Action:* lifts the lower jaw and draws the processus coronoideus and thus also the processus condyloideus somewhat backward. *Innervation:* nn. temporales profundi (V. 3).


Fascia buccopharyngea, very thin, covers the lateral surface of the m. buccinator; it becomes lost in front at the angle of the mouth, is stretched out as a denser cord between the hamulus pterygoideus and the posterior end of the crista buccinatoria [mandibulae] as the *raphe pterygomandibularis* (see Fig. 345) and goes over behind upon the outer surface of the muscles of the throat.



345. Mm. pterygoidei of the right side, external view.

(The arcus zygomaticus has been removed as in Fig. 344; the processus coronoideus and an adjacent piece of bone have been chiselled out of the ramus mandibulae.)

M. buccinator (see also Figs. 341, 342 and 344). **Form:** flat, oblong-quadrangular, thin. **Position:** beneath all the other muscles of the mouth like a transverse, curved band, close to the mucous membrane of the cheeks and lips, along the rows of teeth. **Origin:** raphe pterygomandibularis (see p. 265), lateral surface of the processus alveolaris maxillae and crista buccinatoria (or corresponding portion) of the lower jaw bone. The fibres run partly parallel, partly they decussate into the upper and lower lip at the angles of the mouth and there go over into the fibres of the muscles of the other side. Part of the fibres are attached laterally of the angle of the mouth to the mucous membrane. In the lips it forms the deepest layer of the m. orbicularis oris. In the region of the crown of the second (or first) upper molar it is pierced by the ductus parotideus. **Action:** it draws the angle of the mouth lateralward, closes the mouth opening, presses the lips and cheeks upon the teeth. **Innervation:** n. facialis.




346. Muscles of mastication on right side, viewed from behind and somewhat medianward.

(The posterior half of the skull has been removed.)

M. pterygoideus externus (see also Fig. 345). *Form*: triangular, thick, inferior head bipinnate. *Position*: in the fossa infratemporalis, covered lateralward by the m. temporalis, arcus zygomaticus and m. masseter. *Origin*: by two heads; *upper* and lesser *head*: flat, from the crista infratemporalis and the facies infratemporalis of the great wing of the sphenoid; *lower head*: rounded, partly from the lateral surface of the lamina lateralis proc. pterygoidei and from the processus pyramidalis oss. palatini, partly from the facies infratemporalis maxillae. *Insertion*: the fibres run converging backward and lateralward to the fovea pterygoidea proc. condyloidei mandibulae, the fibres of the upper head partly also to the anterior surface of the capsula articularis and of the discus articularis of the joint of the lower jaw. *Action*: draws the mandibula forward, the proc. condyloideus and discus articularis under the tuberculum articulare; when it acts on one side alone, it draws the jaw somewhat toward the other half of the body. *Innervation*: n. pterygoideus externus (V. 3).

M. pterygoideus internus (see also Fig. 345). *Form*: oblong-quadrangular, thick. *Position*: on the medial side of the lower jaw, in a direction similar to that of the m. masseter; it is covered lateralward partly by the m. pterygoideus externus, m. temporalis, ramus mandibulae and m. masseter. *Origin*: tendinously fossa pterygoidea, by a small part also (lateralward from the m. pterygoideus externus) from proc. pyramidalis oss. palatini and facies infratemporalis maxillae. *Insertion*: by parallel fibres downward, backward and somewhat lateralward to the medial surface of the angulus mandibulae and its neighborhood. *Action*: lifts the lower jaw; when it acts on one side only, it draws the jaw toward the opposite side of the body. *Innervation*: n. pterygoideus internus (V. 3).




347. Right occipital and cervical muscles, viewed from behind.

M. auricularis anterior (O. T. attrahens aurem) (see Figs. 337 and 929). Form: flat, triangular. Position: in front of and above the ear. Origin: galea aponeurotica. Insertion: spina helices auriculare. Action: it draws the auricle forward and upward. Innervation: n. facialis.

M. auricularis superior (O. T. attollens aurem) (see also Figs. 337 and 929). Form: flat, triangular. Position: above the ear. Origin: galea aponeurotica. Insertion: narrow, to the eminentia fossae triangularis auriculæ. Action: it draws the auricle upward. Innervation: n. facialis.


M. auricularis posterior (O. T. retrahens aurem) (see also Fig. 929). Form: oblong-quadrangular, single or divided. Position: behind the ear. Origin: pars mastoidea oss. temporalis, upon the tendon of the m. sternocleidomastoideus. Insertion: with parallel fibres to the eminentia conchæ auriculæ. Action: it draws the auricle backward. Innervation: n. auricularis posterior (VII).

M. transversus nuchæ, inconstant. Form: oblong-quadrangular. Position: it lies superficially upon the linea nuchæ superior, in the same layer as the mm. trapezius and sternocleidomastoideus. Origin: protuberantia occipitalis externa. Insertion: tendon of the m. sternocleidomastoideus. Innervation: n. auricularis posterior (VII).



348. Right platysma, viewed from in front.


Platysma (see also Figs. 337 and 342). **F o r m**: oblong-quadrangular, thin, very large. **P o s i t i o n**: just beneath the skin in the clavicular region, in the neck and in the lower part of the face. **O r i g i n**: fascia pectoralis in region of breast and shoulder. **I n s e r t i o n**: the fibres run parallel, medianward and upward; the posterior fibres radiate out upon the fascia parotideomasseterica, the anterior go partly to the angle of the mouth, partly into the m. quadratus labii inferioris, partly (from the region of the first molar tooth till to the median plane) to the basis mandibulae. Below the chin the fibres of the two sides often decussate. **A c t i o n**: it lifts the skin of the neck from the subjacent tissue; it draws the angle of the mouth to the side and downward. **I n n e r v a t i o n**: ramus colli n. facialis.



349. Muscles of the neck (2nd layer), viewed from in front.

(The head has been bent backward; the platysma has been removed.)

M. sternocleidomastoideus (see also Figs. 347 and 350). **Form**: long, flat-cylindrical, strong. **Position**: on the lateral and anterior surface of the neck, covered in its lower half by the platysma. **Origin**: by two heads; *medial head*: by a long tendon from the anterior surface of the manubrium sterni, partly covering the articulation sternoclavicularis; *lateral head*: by a short tendon from the upper surface of the sternal extremity of the clavicle. Between the two heads there is usually a triangular space, which is broader below, and to which corresponds the fossa supraclavicularis minor (see Fig. 332). **Insertion**: the lateral head runs a little steeper and passes under the medial; the two unite at about the middle of the neck and go obliquely upward, backward and lateralward to become attached by a short tendon to the lateral surface of the processus mastoideus and to the linea nuchae suprema. **Action**: the muscle of one side acting alone turns the head to the opposite side and the face at the same time upward; the muscles of both sides acting together lift the face a little and draw the head a trifle forward, or, if the head be fixed, they elevate the clavicle (in the articulation sternoclavicularis) and the sternum (and with them also the thorax). **Innervation**: n. accessorius and a branch of the n. occipitalis minor (C₂).




350. Muscles of the neck (2nd layer), viewed from the right side.

(The platysma has been removed.)

M. digastricus (see also Figs. 349, 352 and 568). *Form*: two rounded bellies, *venter posterior* and *venter anterior*, with a long intermediate tendon. *Position*: below, behind and medianward from the mandibula, covered behind by the glandula parotis, in front partly by the glandula submaxillaris. The *venter posterior* (bipinnate) arises in the incisura mastoidea oss. temporalis and goes over in front and below into an intermediate tendon, which is fastened by a loop-like strip of the fascia colli to the corpus oss. hyoidei; the *venter anterior* arises from this intermediate tendon and usually also from the corpus oss. hyoidei and goes to the fossa digastrica mandibulae. *Action*: when the hyoid bone is fixed, it depresses the lower jaw; when the lower jaw is fixed, it draws the hyoid bone upward. *Innervation*: *venter posterior*: r. digastricus n. facialis; *venter anterior*: n. mylohyoideus (V.3).


M. stylohyoideus (see also Figs. 349, 351 and 352). *Form*: flat, oblong, thin. *Position*: behind and below the mandibula, just in front of and above the posterior belly of the m. digastricus (vide supra). *Origin*: processus styloideus. *Insertion*: forward, downward and medianward to the corpus oss. hyoidei; in the neighborhood of its insertion it surrounds the m. digastricus by two limbs, a lateral, stronger and a medial, feebler. *Action*: it draws the hyoid bone backward and upward. *Innervation*: r. stylohyoideus n. facialis.



351. Muscles of the neck (3rd layer), viewed from in front.

(Like Fig. 349; in addition however the mm. sternocleidomastoidei, trapezii and digastrici have been removed; the right clavicle has been sawed through close to the sternum and removed.)


M. omohyoideus (see also Figs. 349, 350 and 352). **Form**: flat narrow, long; divided in the centre by an intermediate tendon into two bellies, *venter inferior* and *venter superior*. **Position**: on the lateral and anterior surface of the neck, covered behind and below by the clavicle and the m. trapezius, in the middle by the m. sternocleidomastoideus. The intermediate tendon is ensheathed in the fascia colli and united with the anterior wall of the v. jugularis interna; the venter inferior is attached by stiff connective tissue to the clavicle. **Origin**: margo superior scapulae and lig. transversum scapulae superius, extending to the proc. coracoideus. **Insertion**: runs in front of the mm. scaleni forward and upward to the intermediate tendon and thence, bends upward to the corpus oss. hyoidel, where it lies laterally from the m. sternohyoideus and in front of the m. thyrohyoideus. **Action**: it draws the hyoid bone downward, makes the fascia colli tense, and somewhat extends the v. jugularis interna. **Innervation**: r. descendens n. hypoglossi (fibres from C₁, C₂, C₃).



352. Muscles of the neck (3rd layer), viewed from the right side.

The platysma, m. sternocleidomastoideus and the anterior part of the m. trapezius have been removed.)

M. sternohyoideus (see also Figs. 349—351). *Form*: flat, oblong-quadrangular, thin, mostly having an inscriptio tendinea near its origin. *Position*: on the anterior surface of the neck in front of the larynx, the trachea and the glandula thyroidea, next to the mm. sternothyroideus and thyrohyoideus, covered above by the platysma, below by the m. sternocleidomastoideus; the muscles of the two sides approach very close to each other a little above their origin and sometimes even touch in the median line. *Origin*: posterior surface of the manubrium sterni, of the articulatio sternoclavicularis and of the clavicle (see Fig. 364). *Insertion*: the fibres run parallel to one another upward, at first also somewhat medianward, and go to the corpus oss. hyoidei (*bursa m. sternohyoidei*, see p. 275). *Action*: it draws the hyoid bone downward. *Innervation*: r. descendens n. hypoglossi (fibres from C₁, C₂, C₃).



353. Ligamentum sternoclaviculare


Ligamentum interclaviculare

Muscles of the hyoid bone (deep layer), viewed from in front.

(The mm. sternohyoidei, omohyoidei, digastrici and mylohyoidei have been removed.)

M. sternothyroideus (see also Figs. 349, 351 and 352). **Form**: flat, oblong-quadrangular, very thin, often having an inscriptio tendinea. **Position**: on the anterior surface of the neck, close to the glandula thyroidea, covered by the mm. omohyoideus and sternohyoideus. **Origin**: posterior surface of the manubrium sterni and of the first costal cartilage (see Fig. 364). **Insertion**: upward and at the same time slightly lateralward to the lamina cartilaginis thyroideae (linea obliqua); single fibres go over immediately into the mm. thyrohyoideus and thyropharyngeus. **Action**: it draws the cartilage thyroidea downward. **Innervation**: r. descendens n. hypoglossi (fibres from C₁, C₂, C₃ [C₄]).

M. thyrohyoideus (see also Figs. 349—352). **Form**: flat, quadrangular, very thin. **Position**: on the anterior surface of the neck, close to the cartilago thyroidea, covered by the mm. omohyoideus and sternohyoideus. **Origin**: lamina cartilaginis thyroideae (linea obliqua) (see also above). **Insertion**: corpus and cornu majus oss. hyoidei (*bursa m. thyrohyoidei*, see p. 275). **Action**: draws the hoid bone closer to the cartilago thyroidea and vice versa. **Innervation**: r. thyrohyoideus n. hypoglossi (fibres from C₁ [C₂]).



354. Muscles of the floor of the mouth, viewed from behind and above.

(The muscles of the tongue have been completely removed.)


M. mylohyoideus (see also Figs. 349, 351 and 568). *Form*: flat, triangular, broad; the right and the left muscle together are hollowed above out like an trough. *Position*: the right and the left commonly stretch between the lower jaw and the hyoid bone and thus form the floor of the mouth; immediately adjacent to its lower surface lies on each side the anterior belly of the m. digastricus and the glandula submaxillaris. *Origin*: linea mylohyoidea mandibulae. *Insertion*: the fibres run, parallel to one another, backward, downward and medianward and are attached for the most part to a connective tissue raphe situated in the median plane; the posterior fibres go to the anterior surface of the corpus oss. hyoidei. *Action*: when the lower jaw is fixed, it draws the hyoid bone forward and upward and raises the tongue, when the hyoid bone is fixed, it depresses the lower jaw. *Innervation*: n. mylohyoideus (V. 3).

M. geniohyoideus (see also Figs. 353, 568 and 582). *Form*: flat-cylindrical, oblong. *Position*: on the upper (buccal) side of the m. mylohyoideus, completely hidden by it from below; the mm. geniohyoidei of the two sides are immediately adjacent to one another in the median plane. *Origin*: by a short tendon from the spina mentalis mandibulae. *Insertion*: the fibres, diverging a little, extend to the anterior surface of the corpus oss. hyoidei. *Action*: when the lower jaw is fixed, it draws the hyoid bone forward and upward; when the hyoid bone is fixed, it depresses the lower jaw. *Innervation*: C₁, C₂ and very probably also n. hypoglossus.

Bursa [mucosa] subcutanea prominentiae laryngeae (see Fig. 353), often absent, usually found only in men and only in older individuals; it is unpaired and lies in the subcutaneous tissue in front of the upper part of the thyroid cartilage.

Bursa m. sternohyoidei (see Fig. 353), almost constant, is better developed in the male than in the female and lies, if unpaired, in the median plane or, if paired, close to the median plane upon the lig. hyothyreoideum medium, between it on the one hand and the m. sternohyoideus and the fascia colli on the other; it extends upward behind the hyoid bone.

Bursa m. thyreochoidei (not shown in the drawing), still more constant than the preceding, lies below the cornu majus oss. hyoidei, upon the membrana hyothyreoidea, between it and the upper end of the m. thyreochoideus.




355. Deep muscles of the neck, viewed from in front.

(The sternum and the anterior ends of the ribs with their muscles have been removed.)

M. rectus capitis lateralis. Form: flat, short. Position: lateral from the articulo atlantooccipitalis. Origin: proc. transversus atlantis. Insertion: inferior surface of the proc. jugularis oss. occipit.. Action: bends the head to the side. Innervation: direct branch of the plexus cervicalis (C_1).

M. rectus capitis anterior (O. T. rectus capitis anticus minor). Form: flat, short. Position: in front of the articulo atlantooccipitalis. Origin: massa lateralis and root of the proc. transversus atlantis. Insertion: pars basilaris oss. occipit.. Action: bends the head forward. Innervation: direct branch of the plexus cervicalis (C_1 , C_2).




356. Anterior muscles of the cervical spine on the right side.

M. longus capitis (O. T. rectus capitis anticus major) (see also Figs. 351, 352 and 355). **Form**: flat, oblong-triangular; has in the middle part on the anterior surface an oblong field of tendons.

Position: just in front of the upper end of the cervical spine and in front of the membrana atlantooccipitalis anterior. **Origin**: tubercula anteriora of the transverse processes of the 3rd—6th cervical vertebra. **Insertion**: upward and medianward to the pars basilaris oss. occipitalis. **Action**: bends the head and the upper cervical vertebrae forward and the cervical spine lateralward. **Innervation**: direct branch of the plexus cervicalis (C₁, C₂, C₃).

M. longus colli (see also Fig. 355). **Form**: flat, oblong-triangular, broadest in the middle. **Position**: close to the anterior surface of the cervical spine and of the upper thoracic vertebrae, in the groove between the bodies and transverse processes of the vertebrae; it is partially covered above by the m. longus capitis. Two portions. Inferior, medial portion: from the bodies of the 3rd thoracic to the 5th cervical vertebra, goes to the bodies of the 4th—2nd cervical vertebra and to the anterior tubercles of the transverse processes of the 6th and 5th cervical vertebra; superior, lateral portion: from the anterior tubercles of the transverse processes of the 6th—3rd cervical vertebra, goes to the tuberculum anterius atlantis. **Action**: bends the cervical spine forward and lateralward. **Innervation**: direct branches of the cervical and brachial plexuses (C₂, C₃, C₄, C₅, C₆).



357. Mm. scaleni of the right side, viewed from the right.


M. scalenus anterior (O. T. *scalenus anticus*) (see also Figs. 350 to 352 and 355). **Form:** flat, oblong-quadrangular. **Position:** close to the inferior part of the cervical spine. **Origin:** by four tendinous slips from the anterior tubercles of the transverse processes of the 3rd to the 6th cervical vertebra. **Insertion:** downward and forward by means of a short tendon to the tuberculum scalenii [Lisfranci] of the 1st rib. **Action:** when the cervical spine is fixed it elevates the 1st rib; when the 1st rib is fixed it bends the cervical spine forward and lateralward; and when acting with its fellow of the opposite side the two bend the cervical spine directly forward. **Innervation:** direct branches of the cervical and brachial plexuses ([C₄], C₅, C₆, C₇).

M. scalenus medius (see also Figs. 350—352 and 355). **Form:** flat, oblong-triangular. **Position:** close to the inferior part of the cervical spine, behind the m. scalenus anterior, separated from it at the 1st rib by a triangular space (*scalenus fissure*). **Origin:** by 6 (7) slips from and beside the posterior tubercles of the transverse processes of the 7th—2nd (1st) cervical vertebra. **Insertion:** downward and lateralward to the lateral surface of the 1st rib (behind the sulcus subclaviae). **Action:** when the cervical spine is fixed it elevates the 1st rib; when the 1st rib is fixed it bends the cervical spine lateralward. **Innervation:** direct branches of the cervical and brachial plexuses ([C₄], C₅, C₆, C₇, C₈).

M. scalenus posterior (O. T. *scalenus posticus*) (see also Figs. 350—352 and 355). **Form:** flat, oblong-quadrangular. **Position:**

close to the inferior part of the cervical spine, bounded in front by the m. scalenus medius, behind by the m. iliocostalis cervicis, lateralward by the m. levator scapulae; usually fused with the m. scalenus medius. **Origin:** by 2 slips from the posterior points of the transverse processes of the 5th and 6th cervical vertebrae. **Insertion:** downward and lateralward to the lateral surface of the 2nd rib. **Action:** is related to the 2nd rib as is that of the preceding muscle to the 1st rib. **Innervation:** direct branches of the brachial plexus (C₇ or C₈).


M. scalenus minimus (inconstant). **Form:** flatly-rounded, narrow. **Position:** in the slit between the mm. scalenus anterior and medius; it separates the brachial plexus from the a. subclavia. **Origin:** by 1—2 slips from the transverse processes of the 7th or of the 6th and 7th cervical vertebra. **Insertion:** downward and forward to the lateral surface of the 1st rib and to the pleural cupola. **Action:** makes the pleural cupola tense. **Innervation:** direct branches of the brachial plexus (C₈).



358. Transverse section through the neck, at the level of the uppermost cartilages of the trachea, viewed from above (after Fr. Merkel).

The **fascia praevertebralis** is a thin but strong membrane which covers the anterior surface of the cervical spine and the deep muscles of the neck; part of it descends with these into the thorax, part extends upon the outer surface of the mm. scaleni and of the thorax: it covers directly all the muscles of the neck represented in Fig. 355.


The **fascia colli** extends as a strong membrane from the hyoid bone to the sternum and clavicle in front of the larynx, laryngeal muscles and trachea. It is attached to the sternum by two leaves, which enclose the *spatium suprasternale* and its fatty contents. Laterally it extends above in front of the m. omohyoideus and past it as far as the anterior surface of the bundle of blood vessels, where it becomes lost; it forms a sheath for the intermediate tendon and the venter inferior of the m. omohyoideus. Behind, it becomes thinner and passes below the m. sternocleidomastoideus, upon the fat which helps to fill up the lateral cervical triangle, and below the m. trapezius to the lig. nuchae; the external surface of the mm. sternocleidomastoideus and trapezius is covered only by delicate muscle-fascia.



359. Muscles of the right side of the chest, viewed from in front.

(The platysma has been removed.)


M. pectoralis major (see also Figs. 360, 367, 368 and 394). **Form**: flat, quadrangular when the arm is hanging down, broad, thick. **Position**: on each side it passes out from the median plane to the anterior surface of the thorax and of the axilla, above it is partly covered by the platysma; separated from the muscle of the other side by a groove of varying depth. **Origin**: medial half up to two third of the clavicle (*pars clavicularis*), anterior surface of the sternum and of the second to the sixth costal cartilage (*pars sternocostalis*), and the anterior leaf of the vagina m. recti abdominis (*pars abdominalis*). The *pars clavicularis* is as a rule distinctly separated at the sternoclavicular joint from the *pars sternocostalis* by an open space, and remains separated from it in its entire length. **Insertion**: see p. 281.



360. Right m. pectoralis major, viewed from in front, semi-schematically.

(Two strips have been cut out of the muscle. The dotted black lines indicate the course of the upper, posterior margins of the muscle bundles, the dotted red lines that of the lower, anterior margins.)

M. pectoralis major (continued). **Insertion:** The fibres converge lateralward to the crista tuberculi majoris of the humerus and there become attached by tendons. The pars clavicularis runs in parallel fibres to the anterior part of the terminal tendon; a lateral triangular field of the terminal tendon is here covered by the anterior part of the m. deltoideus. The pars sternocostalis usually converges toward a thick, tendinous plate in the posterior section of the tendon of insertion, and can be divided into several large compartments which run fan-shaped after one another toward the humerus. The lowest part of the pars sternocostalis and the pars abdominalis run slightly twisted without deviating to any marked degree from their path to a tendinous leaf on the dorsal surface of the tendon of insertion, this leaf being connected by loose tissue with the rest of the tendinous mass; the same leaf may rise above the proximal margin of the latter. The tendon of insertion, accordingly, consists of two layers, one anterior, thick (formed by the pars clavicularis and the main mass of the pars sternocostalis), and one posterior, thin (pertaining to the lower part of the pars sternocostalis and to the pars abdominalis); these layers are continuous below but can be separated from above (see Fig. 361). The tendon of insertion is connected below with that of the m. deltoideus. Its posterior layer sometimes divides before it is attached and encloses the tendon of the caput longum m. bicipitis; it also lines the groove between the cristae tuberculi majoris and tuberculi minoris, and is there united with the tendinous fibres of the m. latissimus dorsi. Between the posterior surface of the terminal tendon and the anterior surface of the caput longum m. bicipitis brachii there often lies an oblong bursal sack, **bursa m. pectoralis majoris** (see Fig. 397). **Action:** it moves the scapula forward, draws the upper arm medianward and forward, and rotates it inward; it draws the vertically elevated upper arm downward; the upper part presses the arm somewhat upward against the scapula, and can also raise the cingulum extremitatis superioris a trifle; the lower part can raise the ribs when the shoulder girdle is fixed. **Innervation:** nn. thoracales anteriores ([C5], C6, C7, C8, Th1).




361. Muscles of the right side of the chest (2nd layer), viewed from in front.

(The platysma and the m. pectoralis major have been removed.)

M. pectoralis minor. Form: flat, oblong-triangular. Position: on the anterior surface of the thorax and of the axilla, covered by the m. pectoralis major; it may project somewhat beyond the lateral margin of the m. pectoralis major. Origin: by (four or) three digitations from the anterior surface of the (2nd or) 3rd—5th rib. Insertion: the fibres converge upward and lateralward to the proc. coracoideus scapulae; there is a short tendon which may be united with the m. coracobrachialis. Action: it moves the scapula (and the clavicle) downward and turns its angulus inferior backward and medianward; when the shoulder blade is fixed, it elevates the ribs to which it is attached. Innervation: nn. thoracales anteriores (C₇, C₈).


M. subclavius. Form: flat-cylindrical, narrow, short, pinnated. Position: hidden between the clavicle and the 1st rib, covered by the platysma, the clavicle and the m. pectoralis major. Origin: with a strong tendon, from the anterior surface of the 1st costal cartilage and 1st rib, just lateralward from the lig. costoclaviculare. Insertion: obliquely lateralward and upward to the inferior surface of the clavicle as far as the lig. coracoclaviculare and between its two parts. Action: it draws the clavicle (and the scapula) downward and forward. Innervation: n. subclavius (C₅ [C₆]).



362. Right m. serratus anterior,

viewed from the side and somewhat from in front.

(The mm. pectorales as well as the muscles of the anterior and posterior scapular fossae have been removed; the clavicle has also been removed and the scapula drawn forcibly backward.)



363. Right posterior wall of the thorax with muscles, viewed from in front.

(Diaphragm, m. psoas and m. quadratus lumborum have been removed. The ligamenta intercostalia interna are cut off in the 5th—9th intercostal space.)

M. serratus anterior (O. T. serratus magnus) (see Figs. 359, 361, 362, 367, 368 and 394). **Form:** flat, quadrangular. **Position:** on the lateral and posterior wall of the thorax, close to the ribs and mm. intercostales, covered in front and above by the mm. pectorales, behind by the m. subscapularis, by the scapula and its posterior muscles as well as by the m. latissimus dorsi and m. trapezius; lateralward and below, with its 4–5 inferior digitations, which penetrate between the digitations of the m. obliquus externus abdominis, just beneath the skin. **Origin:** by 9–10 fleshy digitations from the 8–9 uppermost ribs (by 2 digitations from the 2nd rib [tuberositas costae II]); also from a tendinous arch between 1st and 2nd rib. **Insertion:** uppermost 2 digitations (from 1st and 2nd rib) converging to the angulus medialis scapulae; 3rd digitation (from 2nd rib) (occasionally also the 4th digitation, from the 3rd rib) runs diverging to the margo vertebralis (the thinnest part); 4th–10th digitations (from 3rd–9th rib) extend converging to the angulus inferior; the latter portion is the strongest. The fibres run as a whole curved so as to be convex, follow exactly the curvature of the wall of the thorax and increase in length below; the fibres from the 5th–8th rib are the longest. **Action:** it presses the scapula to the thorax, draws it forward and lateralward, acting with especial power upon the angulus inferior; at the same time it rotates the scapula so that the angulus medialis is depressed and the acromion slightly elevated. **Innervation:** n. thoracalis longus (C₅, C₆, C₇ [C₈]).


Mm. intercostales externi (see also Figs. 362, 370 and 387). **Form:** flat, short, thin; they contain a great deal of tendinous tissue. **Position:** in the intercostal spaces, from the tubercles of the ribs on, to about the lateral ends of the costal cartilages; in the first intercostal space sometimes to the sternum; behind they are continuous with the mm. levatores costarum. **Origin:** lower margin of each rib, external from the sulcus costae. **Insertion:** the fibres run parallel downward and sternalward to the upper margin of the next rib below. Medianward from the anterior end running in the same direction are glistening tendinous band of connective tissue (*ligamenta intercostalia externa*, not shown in the figure). **Action:** muscles contract on inspiration. **Innervation:** nn. intercostales I–XI.

Mm. intercostales interni (see also Fig. 361, 362, 364 and 372). **Form:** flat, short thin. **Position:** in the intercostal spaces from the region of the angles of the ribs to the sternum; they lie more internally than the mm. intercostales externi, and are accordingly covered by the latter externally as far as the lateral ends of the costal cartilages; between the costal cartilages they are covered only by the ligg. intercostalia externa; behind they leave the anterior surface of the mm. intercostales externi free medianward from the angles of the ribs. **Origin:** inferior margin of each rib; more exactly, inwards from the sulcus costae from the anguli costarum almost to the margin of the costal cartilage, and outwards from the sulcus costae in the middle of the rib to the sternum, so that the intercostal vessels and nerves run behind between the mm. intercostales externi and interni, then between the two layers of the mm. intercostales interni, and in front, on the inner side of the latter. **Insertion:** the fibres run parallel downward and vertebralward to the upper margin of the next rib below. **Action:** the portions extending between the ossa costalia contract on expiration; those bundles running between the costal cartilages, *mm. intercartilaginei*, act like the mm. intercostales externi. **Innervation:** nn. intercostales I–XI.

The *ligamenta intercostalia interna* are thin bands of connective tissue which are situated, partly in front on the inner side of the mm. intercostales interni in connection with the m. transversus thoracis (not shown in figure), partly behind on the anterior surface of the mm. intercostales externi in connection with the mm. intercostales interni and the mm. subcostales (see Fig. 363).


Mm. subcostales, very variable. **Form:** flat, triangular, very thin. **Position:** on the internal surface of the ribs, usually only of the lower ribs, in the region of the anguli costarum. **Origin:** from the inner surface of the ribs. **Insertion:** the fibres run like those of the mm. intercostales interni, but jump over one or two ribs. **Action:** contract on expiration. **Innervation:** corresponding nn. intercostales.

M. transversus thoracis (see Fig. 364). **Form:** flat, quadrangular, very thin. **Position:** directly upon the posterior surface of the anterior thoracic wall, below close to the diaphragma and bounding the m. transversus abdominis. **Origin:** posterior surface of the proc. xiphoideus and of the lower half of the corpus sterni. **Insertion:** lower bundle transverse, the upper obliquely upward and lateralward to the 6th to the 3rd (or 2nd) rib (costo-cartilaginous junction); the upper ones usually pass over two intercostal spaces. **Action:** it draws the ribs downward on expiration. **Innervation:** nn. intercostales III (or II)–IV.



364. Anterior wall of the thorax with muscles,
viewed from behind.


(On the right side the fibres of origin of the diaphragma have been removed.)



365. Diaphragm, viewed from below and in front.

(The muscles of the chest have been removed, the abdominal muscles have been cut through transversely.)


Diaphragma (*diaphragm*) (see also Figs. 364 and 366). **Form and position:** flat, broad, kidney-shaped, tendinous in the middle (*centrum tendineum*), otherwise fleshy; it is stretched out transversely between the abdominal and thoracic cavities; it bulges into the thorax, being convex above, markedly so lateralward, less so in the middle, so that to the right and to the left there arises a cupola-like elevation, those of the two sides being separated by a shallow median depression; in the adult on moderate expiration the right cupola reaches to close below the costocartilaginous junction of the 4th rib, the left cupola to just above the costocartilaginous junction of the 5th rib (see Fig. 461). **Origin:** from the posterior surface of the proc. xiphoideus sterni (*pars sternalis*), from the internal surface of the anterior ends of the 7th—12th rib (*pars costalis*) and from the lumbar vertebrae (*pars lumbalis*).




366. Diaphragm, pars lumbalis, viewed from below and in front.

(The diaphragm has been separated from the upper ribs and from the sternum and drawn somewhat upward; on the right the mm. psoas and quadratus lumborum have been removed.)


Diaphragma (continued). The pars lumbalis arises from the lig. longitudinale anterius and is divided on each side by perforating vessels and nerves (somewhat variably) into three *crura* (the right ones reaching a little farther down): *crus laterale*, on the right thicker, from the *crura* (the right reaching a little farther down); *crus mediale*, on the right thicker, from the anterior surface of the 3rd and 4th, *crus intermedium*, narrow, from the anterior surface of the 2nd and 3rd, *crus laterale*, from the lateral surface of the 2nd (or 1st) lumbar vertebral body and from a strip of fascia which extends thence in front of the m. psoas to the transverse process of the 1st lumbar vertebra and is fused with the fascia iliaca (see p. 343) (*arcus lumbocostalis medialis* [Halleri]). A second strip of fascia, between the point last mentioned and the 12th rib, (*arcus lumbocostalis lateralis* [Halleri]), forms a bridge over the m. quadratus lumborum and serves chiefly to give origin also to muscle fibres; a triangular space is frequently found over it in the muscle, *trigonum lumbocostale* [Bochdaleki]. The *crura* medialis of the two sides diverge somewhat above and in front and leave between them and the spine an oblong space, the *hiatus aorticus* (for the aorta and ductus thoracicus), which is closed before the centre of the 12th thoracic vertebral body by a small tendinous arch. Over this, the right *crus mediale* divides into two bundles which enclose a slit which reaches almost to the centrum tendineum, the *hiatus oesophageus* (for the oesophagus and the nn. vagi); the left *crus mediale* sends only a few corresponding fibres to the right. Between the *crus mediale* and the *crus intermedium* run usually the v. azygos (or hemiazygos) and the nn. splanchnici; between the *crus intermedium* and the *crus laterale* runs the truncus sympathicus. **Insertion:** on all sides radially to the *centrum tendineum* (O. T. central tendon), which has the shape of a clover-leaf or sometimes of a kidney and consists of interwoven tendon bundles. In its right half lies the *foramen venae cavae*, quadrangularly rounded (for the v. cava inferior). **Action:** it enlarges the thoracic cavity; the cupolae descend markedly, the centrum tendineum descends less. **Innervation:** nn. phrenici (C3, C4, C5).



367. Muscles of the abdomen,
viewed from the right side and the front.



368. Muscles of the abdomen, viewed from in front.




369. Mm. pyramidales, viewed from in front.


(The anterior leaf of the sheath of the rectus has been partially dissected off and reflected.)

M. obliquus externus abdominis (see also Figs. 367, 368, 380 and 382). **Form**: flat, quadrangular, broad, much longer in front than behind. **Position**: on the anterior and lateral surface of the abdominal and inferior thoracic region. **Origin**: by 8 fleshy digitations from the outer surface of the 12th—5th rib; the uppermost digitation is usually covered by the pars abdominalis m. pectoralis majoris (see Fig. 370); the 4 upper digitations are inserted between the digitations of the m. serratus anterior; the 3 lower between those of the m. latissimus dorsi. **Insertion**: the upper fibres run more transversely, forward, the lower more obliquely, forward and downward. The largest, upper part goes over medianward into a broad, tendinous sheath which runs in front of the m. rectus abdominis and helps to form the anterior leaf of the vagina m. recti abdominis (see p. 279). The next lower muscle bundles pass obliquely to the *ligamentum inguinale [Poupart]* (O. T. Poupart's ligament) (see p. 298) and leave medianward an open space (*annulus inguinalis subcutaneus*) (see p. 298) for the passage of the funiculus spermaticus. The muscle fibres arising farthest back are attached in fleshy masses to the external lip of the iliac crest; they form, when the muscles are relaxed, a ridge (see Fig. 374) running in a horizontal direction (*flank pad*), projecting slightly beyond the crista iliaca. **Action**: it draws the thorax downward (expiration); bends the spinal column forward and toward the same side and rotates it toward the other side; when the thorax is fixed it elevates the pelvis. **Innervation**: nn. intercostales V—TII and the nn. iliohypogastricus and ilio-inguinalis (total Th₅—Th₁₂ [L₁]).

M. pyramidalis, of variable degree of development, sometimes absent altogether. **Form**: flat, oblong-triangular. **Position**: close to the median plane above the symphysis pubis, in front of the m. rectus abdominis, in a duplicature of the anterior leaf of the vagina m. recti abdominis. **Origin**: anterior surface of the ramus superior ossis pubis and of the symphysis ossium pubis. **Insertion**: the fibres run upward and medianward to the linea alba. **Action**: it tightens the linea alba in the upright position and gives rise to a median groove below the navel when the abdominal muscles are relaxed. **Innervation**: n. intercostalis XII; possibly also fibres from L₁ and L₂ or only from these.




370. Muscles of the abdomen (2nd layer), viewed from in front.




371. Transition of the tendon of the right m. obliquus internus abdominis into the sheath of the rectus.

(The m. obliquus externus abdominis has been completely removed above; below, it has been cut through and reflected; the m. obliquus internus abdominis has been separated at its upper border from the costal cartilages; the anterior leaf of the sheath of the rectus has been removed in the upper portion.)


M. obliquus internus abdominis (see also Figs. 370 and 382—384). **Form**: flat, quadrangular, broad, in front considerably longer than behind. **Position**: on the anterior and lateral surface of the abdomen, completely covered by the m. obliquus externus abdominis. **Origin**: posterior surface of the combined leaves of the fascia lumbodorsalis (see p. 306), linea intermedia of the iliac crest, lateral two thirds of the lig. inguinale (see p. 298). **Insertion**: the most posterior fibres pass upward and forward to the inferior margins of the 12th—10th rib. The other fibres coming from the iliac crest diverge and pass medianward so that the uppermost run parallel to the ends of the 10th—8th costal cartilages (connected with them only by loose tissue), the lowermost run transversely; the fibres from the lig. inguinale are somewhat curved as they run downward. All the muscle bundles not fastened to the ribs go over, lateralward from the m. rectus, along a curved line, into a tendinous plate. This divides in its upper two thirds into two lamellae, of which one passes in front of, the other behind, the m. rectus abdominis to go to the median plane and there form the basis for the corresponding leaf of the *vagina m. recti abdominis* (O. T. sheath of the rectus) (see also Fig. 373 and p. 297). In the lower third, all the fibres run in front of the m. rectus abdominis (see also Fig. 374 and p. 297); the posterior leaf of the rectal sheath is lacking here generally (see p. 297); its lower margin is curved so as to be concave and is called the *linea semicircularis [Douglassi]* (see Fig. 375 and p. 297). The lower margin of the muscle forms in part the upper wall of the inguinal canal (see p. 299); in addition the m. cremaster goes off from it (see p. 299). **Action**: the upper part draws the thorax downward (expiration), bends the spinal column forward and toward the same side, rotates it toward the same side; when the thorax is fixed it elevates the pelvis; the lower part acts in a similar manner to the m. transversus abdominis. **Innervation**: nn. intercostales X—XII and the nn. iliohypogastricus, ilioinguinalis and spermaticus externus (total [Th₁₀] Th₁₁, Th₁₂, L₁ [L₂]).



372. Mm. recti abdominis, viewed from in front.




373. Transverse section through the abdominal wall above the linea semicircularis [Douglassi] (after W. Braune).



374. Transverse section through the abdominal wall below the linea semicircularis [Douglassi] (after W. Braune).

M. rectus abdominis (see Figs. 372 and 375). **Form**: flat, oblong-quadrangular, above broader and thinner, below narrower and thicker. **Position**: on the anterior surface of the abdomen close to the linea alba. In front it is covered by the anterior leaf of the vagina m. recti abdominis; behind it lies above immediately on the 6th–8th rib cartilage, below on the m. transversus abdominis and on the posterior leaf of the vagina m. recti, in the lower third where the latter is lacking, directly upon the fascia transversalis. **Origin**: anterior surface of the 5th–7th costal cartilage, of the proc. xiphoideus and of the lig. costoxiphoideum. **Insertion**: its fibres run downward, parallel to one another. In its upper half it is interrupted by three intercalated tendons, *inscriptiones tendineae*; the uppermost, usually incomplete, goes slightly arch-shaped at the level of the horizontal part of the 8th costal cartilage, the lowermost runs transversely at the level of the navel, and the middle one, between these two, in a similar direction; these intermediate tendons often do not go entirely through the thickness of the muscle; they are very intimately connected with the anterior leaf, but loosely with the posterior leaf of the rectal sheath. The number and extent of these tendons varies; usually there is an incomplete one in the lateral half of the muscle below the navel at the level of the linea semicircularis [Douglassi]. Below, the muscle fibres converge and are attached by a strong flat tendon to the ramus superior oss. pubis and to the anterior surface of the symphysis ossium pubis; the fibres of the tendons are there very much interwoven. **Action**: it draws the thorax downward (expiration) and bends the spinal column forward; when the thorax is fixed it elevates the pelvis. Single sections of the muscle can contract separately. **Innervation**: nn. intercostales VI–XII.



375. Muscles of the abdomen (3rd layer), viewed from in front.

(Mm. obliqui abdominis have been completely removed; furthermore on each side the m. rectus abdominis and the anterior leaf of its sheath have been removed on each side in their upper two thirds.)


M. transversus abdominis (see also Figs. 364, 372 and 383). **Form**: flat, quadrangular, broad, much longer in front than behind. **Position**: on the anterior and lateral surface of the abdomen, completely covered by the mm. obliqui abdominis, partly also by the m. rectus abdominis. The muscle fibres which arise from the 7th and 8th rib lie entirely behind the m. rectus abdominis. **Origin**: from the internal surfaces of the 7th—12th costal cartilage, from the union of the two leaves of the fascia lumbodorsalis (see Figs. 383 and 384), from the internal lip of the iliac crest and from the lateral two thirds of the lig. inguinale (see p. 298). **Insertion**: the fibres are parallel, run transversely forward and go over into a tendon plate along a line which is concave medianward (*linea semilunaris* [Spigeli]). This tendon plate (see also Figs. 373 and 374) runs, in its upper two thirds, behind the m. rectus abdominis and strengthens the posterior leaf of the sheath of the rectus; in the lower third, where there is no posterior leaf of the rectal sheath (see below), it goes in front of the m. rectus abdominis and fuses with the anterior leaf of the rectal sheath; some bundles may run there also behind the m. rectus. The lowermost fibres arch downward and help to form the posterior wall of the inguinal canal (see pag. 300); they are attached partly to the upper surface of the os pubis, partly to the lig. inguinale. **Action**: it pulls the ribs downward (expiration) and diminishes the size of the abdominal cavity. **Innervation**: nn. intercostales V—XII, the nn. iliohypogastricus, ilioinguinalis and spermaticus externus (total [Th₅] Th₁₂, L₁, L₂).

Vagina m. recti abdominis (*sheath of the rectus*) (see also Figs. 368, 370, 371, 373 and 374) consists on each side of two broad tendinous leaves, which cover directly the anterior and posterior surfaces of the m. rectus abdominis and which are formed by the tendons of the mm. obliqui et transversus abdominis; medianward both leaves meet those of the other side in the linea alba. The anterior leaf covers the whole anterior surface of the m. rectus and is firmly fused with its inscriptiones tendineae. It is very thin above and there serves to give origin to the pars abdominalis of the m. pectoralis major; it becomes stronger lower down and consists of interwoven fibres. In its upper two thirds it is formed by the anterior leaf of the tendon of the m. obliquus internus and the tendon of the m. obliquus externus (see Fig. 373) fused with it, in its lower third by the union of the tendons of the mm. obliqui and of the m. transversus abdominis (see Figs. 374 and 375). The posterior leaf is much shorter, begins on the costal cartilages at the angulus infrasternalis and ends as a stiff membrane, more or less sharply, somewhat below the navel in a margin, the *linea semicircularis* [Douglasi], which is concave below and stretches out lateralward from the linea alba; it consists of the transverse tendinous fibres of the posterior leaf of the tendon of the m. obliquus internus abdominis and those of the m. transversus abdominis. The posterior leaf more or less thinned and formed only by fibres of the m. transversus, extends frequently further downwards and rarely reaches to the upper edge of the symphysis, leaving a gap only for the passage of the vasa epigastrica inferiora.

Linea alba (see also Figs. 368, 369 and 372—374) is a tough connective tissue strip which extends in the median plane between the mm. recti abdominis from the proc. xyphoideus to the symphysis ossium pubis. Below the proc. xyphoideus it gradually broadens, being about twice as wide at the level of the navel as it is higher up, and again narrows below the navel to a thin cord which is attached behind the tendons of the mm. recti abdominis to the lig. pubicum superius by means of a triangular expansion (*adminiculum lineae albae*) (see Fig. 379). The sheath of the rectus is attached to the linea alba; indeed it forms it, in a way, through the union of the two leaves of the right side with the two leaves of the left. Approximately below its middle and about at the level of the body of the 4th lumbar vertebra the linea alba contains the navel, *umbilicus*; individually its position changes to no small degree.

Ligamentum fundiforme penis (O. T. suspensory ligament) (see Figs. 367 and 368) (in the male) is a broad band, consisting essentially of elastic fibres; it arises, above the symphysis ossium pubis, from the anterior surface of the vagina m. recti and the linea alba as well as from the fascia superficialis abdominis. It surrounds the penis like a loop by means of its two limbs and often radiates out into the scrotum.

Ligamentum fundiforme clitoridis (O. T. suspensory ligament) (see also Fig. 728) (in the female) correspond to the ligamentum fundiforme penis.




376. Right anterior abdominal ring and fossa ovalis in the male.

Ligamentum inguinale [Pouparti] (O. T. Poupart's ligament) (see also Figs. 367, 368, 375, 379, 421 and 427) is a powerful, flatly rounded tendon, which extends from the spina iliaca anterior superior to the tuberculum pubicum and can be palpated through the skin as a hard cord. It is not an independent ligament but formed medially by the tendinous fibres of the m. obliquus externus, laterally by that of both mm. obliqui and the m. transversus abdominis; to its lower margin the fascia lata is attached and this causes a convex downward bulging of the ligament. Medianward the ligament does not end at the tuberculum pubicum alone but also sends diverging fibres to the pecten ossis pubis; the latter insertion is a small triangular ligament, *ligamentum lacunare [Gimbernati]* (O. T. Gimbernat's ligament) (see also Fig. 379 and 421), which is fused with the fascia pectinea, extends obliquely backward and upward and with its lateral concave margin bounds the annulus femoralis (see p. 343).

Canalis inguinalis (inguinal canal) (see also Figs. 369 and 377—379) is a cylindrical space filled up completely by the ductus deferens (in the male, by the lig. teres uteri in the female) with the accompanying vessels (see p. 607 and 619); it perforates the lowermost portion of the abdominal muscles obliquely from behind and above, forward, downward and medianward. The canal has two openings, one posterior, *annulus inguinalis abdominalis* (O. T. internal abdominal ring), and one anterior, *annulus inguinalis subcutaneus* (O. T. external abdominal ring), and (somewhat schematically) four walls, one anterior, one inferior, one superior, one posterior. It is narrower in the female, but somewhat longer than in the male.

The **annulus inguinalis subcutaneus** (see also Fig. 369) lies somewhat laterally from and a little above the tuberculum pubicum. It is formed by the separation of the lowermost fibres of the tendon of the m. obliquus externus from one another giving rise to an acute-angled space, which is bounded below by fibres which go to the tuberculum pubicum (*crus inferius*, O. T. external pillar), above by fibres which run to the anterior surface of the symphysis ossium pubis (*crus superius*, O. T. internal pillar); the larger lateral portion of the space is filled up by curved fibres which run transversely, *fibrae intercrurales* (O. T. intercolumnar fascia, external spermatic fascia.) These fibres leave as a free opening only a small space, convex laterally, which will permit the passage of the tip of the index finger.



377. Right inguinal canal in the male (2nd layer), viewed from in front.

(The m. obliquus externus abdominis has been cut through above the anterior abdominal ring and reflected partly medianward and partly downward. The spermatic cord has been pressed somewhat downward within the inguinal canal.)


Canalis inguinalis (continued) (see also Figs. 369, 376, 378 and 379).

Anterior wall: formed by the tendon plate of the m. obliquus externus abdominis, especially also by the fibrae intercrurales.

Lower wall: m. obliquus externus; the lig. inguinale [Pouparti] and the fibrous bands of the crus inferius immediately adjacent do not extend in their medial portion in a straight line to the tuberculum pubicum but are curved somewhat downward so that they form a groove open above for the spermatic cord.

Upper wall: m. obliquus internus; the lowermost fibres of this muscle, arising from the lig. inguinale, extend in curves medianward and downward, forming lateralward the muscular, narrow, upper wall of the inguinal canal and going over medianward into the anterior layer of the sheath of the rectus; this medial part has its lower insertion sometimes farther lateralward and then passes close in front of and along with the falx inguinalis to the upper surface of the os pubis.

M. cremaster (see also Figs. 376 and 705) (in the male). It arises by a thick lateral bundle in continuation of the lowermost fibres of the mm. obliquus internus and transversus abdominis from the upper margin of the lig. inguinale and by a weak, medial bundle from the anterior layer of the sheath of the rectus; the former lies on the lateral, the latter on the medial surface of the spermatic cord; both pass with the spermatic cord through the subcutaneous opening of the inguinal canal, spread out in common upon the tunica vaginalis communis of the testis and epididymis, and surround it like bows, forming also a long-meshed net. **Action:** it elevates the testis. **Innervation:** n. spermaticus externus (L_2 or L_1 , L_2). — In the female this muscle consists of only a few fibres which go to the lig. teres uteri.




378. Right inguinal canal in the male (3rd layer), viewed from in front.

(The mm. obliqui abdominis have been cut through above the inguinal canal and reflected medianward and lateralward. The spermatic cord has been pressed somewhat downward within the inguinal canal.)

Canalis inguinalis (continued). **Posterior wall** (see also Fig. 379): m. transversus; the lowermost fibres of the m. transversus arch downward on the median side of the posterior abdominal ring, spread out in a divergent manner and become attached medianward to the upper surface of the os pubis, lateralward to the upper margin of the lig. inguinale. The approximately triangular plate thus arising forms the posterior wall of the inguinal canal and varies in strength at different places. Medianward it contains a strip of strong tendinous fibres, *falx [aponeurotica] inguinalis*, which extends from above downward and medianward to the upper surface of the os pubis, and becomes firmly attached by its medial border to the tendon of the m. rectus; in front it is not rarely covered and strengthened by tendon fibres of the m. obliquus internus (see p. 299). In the lowermost portion of this strip there are also tendon fibres which extend from the lateral margin of the m. rectus abdominis, in curves, downward and lateralward (see Fig. 379); not seldom they are more powerful than the others. The lateral part of the posterior wall is strengthened by the lig. interfoveolare (see p. 301).


The *ligamentum inguinale reflexum [Collesi]* (see also Fig. 369) is a thin band of fibres, one finger broad, which comes into view medial from the annulus inguinalis subcutaneus, between the two crura. It extends from the medial end of the lig. inguinale and lig. lacunare obliquely upward and medianward behind the crus superius to the anterior layer of the sheath of the rectus and in so doing lies partly in front of the lowermost portion of the falx inguinalis, partly it connects with fibres of the m. obliquus externus of the other side.




379. Posterior wall of the right inguinal canal in the male, viewed from behind.

Canalis inguinalis (continued). **Posterior wall** (continued): The lateral part of the plate connected with the *m. transversus*, which forms the posterior wall (see p. 300), is strengthened laterally by a band of fibres markedly concave lateralward and upward, the *ligamentum interfoveolare* [Hesselbachi]. This lies just in front of the *a. epigastrica inferior*. It consists of fibres which bend directly downward from the *m. transversus* and of those which go off behind the same farther up; both kinds of fibres extend downward, bend around chiefly lateralward and upward, partly also medianward and downward to the inner surface of the *lig. inguinale* and fuse with it. The *falx inguinalis* and the *lig. interfoveolare* vary greatly in strength and breadth. The thinner spot, situated between the two, which lies directly behind the subcutaneous abdominal ring, is broad, when they are narrow, and vice versa; it contains always single tendon fibres of the *m. transversus*, consists otherwise of irregularly and loosely arranged fibres, and is perforated by the *a. spermatica externa* (see Fig. 519) and sometimes the *n. spermaticus externus*. On the posterior surface of the posterior wall there often run a few bundles of muscle fibres, *m. interfoveolaris*, either directly behind the *lig. interfoveolare*, or somewhat more medianward; they arise like a fan behind the *m. transversus* (sometimes also from it) and extend downward to the posterior surface of the *lig. inguinale* and the *lig. lacunare*.

The *annulus inguinalis abdominalis* lies somewhat medianward above the middle of the *lig. inguinale*, is bounded medianward and below by the *lig. interfoveolare* and flattens out gradually lateralward and upward.




380. Broad muscles of the back, viewed from behind.



381. Broad muscles of the back (2nd layer), viewed from behind.

(The mm. trapezii and sternocleidomastoidei have been removed.)

M. trapezius (see Fig. 380). **Form**: flat, triangular, broad. **Position**: in the neck and upper part of the back, just beneath the skin. **Origin** (see also Fig. 347): tendinous from the linea nuchae superior, protuberantia occipitalis externa, lig. nuchae und lig. supraspinale as far as the 12th thoracic vertebra; the tendon fibres are connected with those of the other side, are longer in the neighborhood of the vertebra prominens, and there form, on each side, a large triangular area, on the 11th and 12th thoracic vertebra frequently a small area. **Insertion**: the upper fibres extend obliquely downward and forward to the acromial extremity of the clavicle; the middle fibres transversely to the acromion and to the spine of the scapula; the lower fibres obliquely upward and lateralward to become attached by a small, triangular, flat tendon to the medial part of the spine of the scapula. **Action**: it draws on contraction of all the bundles the shoulder-blade toward the spine and rotates it so that the angulus medialis is displaced medianward, the acromion upward and medianward, the angulus inferior slightly lateralward and forward; the upper portion alone elevates, the lower portion alone depresses the shoulder-blade and turns it somewhat, causing the lower angle to move lateralward; when the shoulder-blade is fixed, it draws the head backward and rotates it, along with the face, somewhat toward the other side. **Innervation**: n. accessorius and a branch of the plexus cervicalis (C₂, C₃, C₄).




382. Muscles of the lumbar region (1st layer), viewed from behind.

(On the left side the *m. latissimus dorsi* has been removed.)

M. latissimus dorsi (see also Figs. 367, 380, 381 and 394). **Form**: flat, triangular, thin. **Position**: in the lower part of the back and lateral region of the thorax, being partly covered above by the *m. trapezius*. **Origin**: tendinous from the posterior layer of the fascia lumbodorsalis, from the *proc. spinosus* of the 7th thoracic vertebra on downward, from the *crista iliaca* and, by three or four fleshy digitations, from the external surface of the three or four lowermost ribs, not infrequently also from the *angulus inferior*. **Insertion**: the fibres run, converging, lateralward, the upper fibres more transversely, the lower almost straight upward; they lie on the lateral wall of the thorax close to the *m. serratus anterior*, cover in part the *angulus inferior scapulae*, join closely to the lower margin of the *m. teres major*, wind around it to its anterior surface and become attached by a flat tendon to the *crista tuberculi minoris humeri*. Between the two muscles there lies the *bursa m. latissimi dorsi* (see Fig. 396). **Action**: it draws the arm medianward and backward and rolls it inward; it draws the vertically elevated arm downward; it draws the shoulder-blade medianward and downward. **Innervation**: *n. thoracodorsalis* ([C6] C7, C8).

M. rhomboideus major (see Figs. 380 and 381). **Form**: flat, quadrangular. **Position**: upper part of the back, covered by the *m. trapezius*; under it a triangular area immediately underneath the skin. **Origin**: tendinous from spinous processes and interspinous ligaments of the 1st—4th thoracic vertebra. **Insertion**: the fibres run parallel downward and lateralward to the *margo vertebralis scapulae* (lower half). **Action**: it draws the shoulder-blade medianward and upward and rotates it at the same time so that the *angulus inferior scapulae* is more markedly turned medianward and upward than the *acromion*. **Innervation**: *n. dorsalis scapulae* (C5).

M. rhomboideus minor (see Fig. 381), often only indistinctly separable from the preceding muscle. **Form**: flat, oblong-quadrangular. **Position**: directly on the upper margin of the preceding muscle. **Origin**: tendinous from the *lig. nuchae* and the spinous processes of the two lower cervical vertebrae. **Insertion**: like that of the preceding muscle, just above the same, including the *trigonum spinae scapulae*. **Action and innervation**: same as for the preceding muscle.




383. Muscles of the lumbar region (2nd layer), viewed from behind.

(The mm. latissimus dorsi and obliquus externus abdominis have been removed on both sides; on the left side the mm. obliquus internus abdominis and serratus posterior inferior have also been removed).

M. levator scapulae (O. T. levator anguli scapulae) (see Figs. 350—352 and 381). **Form:** flatly rounded, oblong. **Position:** on the lateral surface of the neck, covered above by the m. sternocleidomastoideus and below by the m. trapezius, in front coming in contact with the m. scalenus posterior, behind with the mm. splenii cervicis et capitis. **Origin:** by four tendinous processes becoming weaker from above downward, from the transverse processes (posterior tubercles) of the 1st—4th cervical vertebra (see Figs. 357 and 386). **Insertion:** the bundles run slightly twisted downward, backward and lateralward to the margo vertebralis scapulae above the trigonum spinae scapulae. **Action:** it draws the shoulder-blade medianward and upward and rotates it so that the angulus inferior is turned strongly medianward and upward, and the acromion less markedly upward; when the shoulder-blade is fixed, it bends the cervical portion of the spine and rotates it somewhat to the same side. **Innervation:** n. dorsalis scapulae ([C₂] C₃, C₄, C₅).

Trigonum lumbale [Petiti] (O. T. Petit's triangle) (see Fig. 382), a triangular thinner spot of the posterior abdominal wall, varying in size; below it is bounded by the crista iliaca, lateralward by the posterior margin of the m. obliquus externus abdominis, medianward by the lateral margin of the m. latissimus dorsi. In this triangle the muscle fibres of the m. obliquus internus are visible; in front of them lies the tendon of origin of the m. transversus abdominis and of the m. quadratus lumborum.

Fascia nuchae (not represented in the figure) is a thin membrane of connective tissue which lies just beneath the mm. trapezius and rhomboidei; it goes off from the inferior surface of the squama occipitalis, covers the mm. splenii capitis et cervicis and semispinalis capitis, i. e. the muscles depicted in Fig 392, is connected medianward with the lig. nuchae, is continued in front into the fascia colli and goes over below at the m. serratus posterior superior into the fascia lumbodorsalis.



384. Transverse section through the muscles of the lumbar region (after W. Braune).

The **fascia lumbodorsalis** consists of two layers, the posterior and the anterior layer.

The posterior layer (see also Figs. 380, 382, 383 and 385) lies just below the mm. trapezius, rhomboidei and latissimus dorsi on the posterior surface of the m. sacrospinalis, is thin at its beginning at the m. serratus posterior superior, where it is continuous with the fascia nuchae (see p. 305), forms a sheath below for the m. serratus posterior inferior and, becoming stronger, descends as far as the coccygeal region; on its way it is fused medianward with the spinous processes and supraspinal ligament of the thoracic and lumbar vertebrae and with the crista sacralis media; below, it fuses with the tendons of origin of the m. sacrospinalis, with the posterior ligaments of the sacrum and with the crista iliaca, lateralward with the anguli costarum and, below the ribs, between them and the crista iliaca, with the anterior layer. It consists above of transverse, below of oblique crossing fibres, which represent in part the immediate continuation of the mm. serratus posterior inferior and latissimus dorsi arising from the fascia.

The anterior layer (see also Figs. 363, 366, 383 and 391) is thinner, lies in front of the m. sacrospinalis, behind the mm. quadratus lumborum and psoas major and extends only from the 12th rib to the crista iliaca, being fastened to both. It arises by four processes from the transverse processes of the 1st—4th lumbar vertebra, is connected below with the lig. iliolumbale and contains, above, powerful, curved fibrous bands which extend from the 1st and 2nd lumbar vertebra to the 12th rib, the *ligamentum lumbocostale* (see Figs. 363, 366 and 391). The fibres of the anterior layer run essentially transversely, unite lateral from the m. sacrospinalis with the fibres of the posterior layer and thus together form the tendon of origin of the m. transversus abdominis; the m. obliquus internus abdominis arises in part from the posterior surface of the combined layers.

385. Broad muscles of the back (3rd layer), viewed from behind.


(The shoulder-blades and the muscles going to them have been removed.)


M. serratus posterior superior (O. T. serratus posticus superior). Form: flat, quadrangular, very thin. Position: upper part of the back, covered by the mm. rhomboidei. Origin: by a flat tendon from the lig. nuchae or the spinous processes of the 6th and 7th cervical, as well as from the 1st and 2nd thoracic vertebra. Insertion: the fibres run parallel, obliquely lateralward and downward and terminate by four fleshy digitations on the 2nd to 5th rib. Action: it elevates these ribs. Innervation: nn. intercostales I—IV, often also a branch of the plexus brachialis (C₅).

M. serratus posterior inferior (O. T. serratus posticus inferior) (see also Fig. 382). Form: flat, quadrangular, very thin. Position: lower part of the back, covered by the m. latissimus dorsi and partly also by the m. trapezius. Origin: posterior layer of the fascia lumbodorsalis in the region of the 10th to 12th thoracic and the 1st and 2nd lumbar vertebra. Insertion: the fibres run parallel obliquely lateralward and upward and terminate by four fleshy digitations on the 9th—12th rib. Action: it draws these ribs downward and outward; it contracts on inspiration. Innervation: nn. intercostales IX—XII.

M. splenius capitis (see also Figs. 347, 352, 381 and 386). Form: flat, oblong-quadrangular, thick. Position: in the neck, partly covered by the mm. trapezius, sternocleidomastoideus, serratus posterior superior and rhomboidei. It embraces on both sides the mm. semispinalis capitis, longissimus capitis et cervicis. Origin: lig. nuchae at the level of the 3rd—7th cervical vertebra as well as from the spinous processes of the 1st and 2nd thoracic vertebra. Insertion: the fibres converge somewhat and pass upward and lateralward by a thick, short tendon to the proc. mastoideus and to the linea nuchae superior oss. occipitalis. Action: it rotates the head so that the face is turned upward and toward the same side; the muscles of the two sides together bend the head backward. Innervation: rami laterales of the rami posteriores of the nn. cervicales I—IV (V).

M. splenius cervicis (O. T. splenius colli) (see Figs. 347, 381 and 386). Form: flat, oblong-quadrangular, narrow. Position: it forms in a way a continuation downward of the preceding muscle and is often indistinctly separable from it. Origin: procc. spinosi of the 3rd—6th thoracic vertebra. Insertion: by 2—3 processes to the tubercula posteriora of the procc. transversi of the 2—3 uppermost cervical vertebrae. Action: it bends the neck backward and lateralward and rotates the 2—3 upper cervical vertebrae like the preceding muscle rotates the head. Innervation: like that of the preceding muscle.





386. Muscles of the neck (left: 2nd layer, right: 3rd layer), viewed from behind.

(On both sides the mm. trapezius, sternocleidomastoideus and levator scapulae have been removed, on the right side the mm. splenii have also been removed.)

387. Long muscles of the back (1st layer), viewed from behind

(On the right side the m. iliocostalis has been pulled lateralward, on the left side it has been reflected.)

M. sacrospinalis (O. T. *erector spinae*) (see also Figs. 383, 384 and 388—391). **Form:** triangular-prismatic, long. **Position:** lateralward from the m. transversospinalis, from the sacral region to the neck. **Origin:** fleshy in the depth, otherwise by long tendons from the crista iliaca, ligg. sacroiliaca posteriora, crista sacralis media and spinous processes of the lumbar vertebrae and of the 2 (or 3) lowermost thoracic vertebrae. The mass of fibres is divisible in the lumbar region into three separate muscles: m. iliocostalis, m. longissimus and m. spinalis.


M. iliocostalis (O. T. *external division of erector spinae*), farthest lateralward, extends between the ribs or their homologues.

M. iliocostalis lumborum (O. T. *sacrolumbalis*). **Insertion:** by nine processes, the lower ones fleshy, the upper ones thin and tendinous, to the angles of the 12th—4th rib. **Action:** it can depress the ribs.

M. iliocostalis dorsi (O. T. *accessory to the iliocostal*). **Origin:** covered by the preceding, it arises by six fleshy digitations from the angles of the 12th—7th rib. **Insertion:** by eight thin tendons to the angles of the 7th to 1st rib and the transverse process of the 7th cervical vertebra.


M. iliocostalis cervicis (O. T. *cervicalis ascendens*) (see also Fig. 386). **Position:** it is bounded in front by the m. scalenus posterior, lateralward and behind by the m. levator scapulae, medianward by the mm. splenius and longissimus cervicis. **Origin:** covered by the preceding muscle, it arises by 4—6 fleshy processes from the angles of the 6th—3rd (2nd or 1st) rib. **Insertion:** by three narrow tendons to the transverse processes of the 6th—4th cervical vertebra. **Action:** it bends the neck backward and toward the same side; it can elevate the ribs.

Action of the whole m. iliocostalis: it bends the spinal column backward and toward the same side. **Innervation:** rami laterales of the rami posteriores of the nn. thoracales I to XII and lumbalis I.



388. Long muscles of the back (2nd layer), viewed from behind.

(The mm. iliocostales have been removed; the mm. longissimi dorsi have been reflected outward; the m. longissimus cervicis has been, on the right side, drawn lateralward, on the left side, medianward; the m. longissimus capitis has been drawn medianward on the right side.)



M. longissimus (O. T. middle division of erector spinae) (see also Fig. 387) is the middle bundle of the m. sacrospinalis; it extends (in the thorax only by its medial digitations) between the transverse processes or their homologues.

M. longissimus dorsi. Origin: it receives very frequently medianward several short, accessory bundles from the procc. mammillares of the 1st and 2nd lumbar vertebrae and the transverse processes of the lower 6–7 thoracic vertebrae. Insertion: 1st, by thin, lateral processes, fleshy below, tendinous above (often partly absent), to the transverse processes of the 5th to 1st lumbar vertebra and the angles of the 12th–2nd rib (see Fig. 387); 2nd, by medial processes, below fleshy, above tendinous (always present), to the procc. accessorii of the 5th to 1st lumbar vertebra and the procc. transversi of all the thoracic vertebrae. Action: it bends the spinal column backward and toward the same side.

M. longissimus cervicis (see also Fig. 386). Form: flat, thin. Position: medianward from the m. iliocostalis cervicis, lateralward from the m. longissimus capitis covered by mm. splenius cervicis and trapezius. Origin: uppermost 4–6 thoracic vertebrae. Insertion: the middle cervical vertebrae. Action: it bends the neck backward and toward the same side.

M. longissimus capitis (O. T. trachleomastoid) (see also Fig. 386). Form: flat, oblong-quadrangular, thin; usually fused with the preceding muscle. Position: medianward from the m. longissimus cervicis, lateralward from the m. semispinalis capitis covered by mm. splenii and trapezius. Origin: procc. transversi of the 3rd–1st thoracic vertebra and procc. articulares of the 7th–3rd cervical vertebra. Insertion: thickened and tendinous to the proc. mastoideus. Action: it draws the head backward, bends and turns it toward the same side.

Innervation: rami laterales of the rami posteriores of the nn. cervicales I–VIII, thoracales I–XII, lumbales I–V.

389. Short muscles of the back (1st layer), viewed from behind.

(The *m. semispinalis capitis* has been removed on the left side.)

M. spinalis (O. T. inner division of erector spinae) (see Fig. 387), very thin, forms the medial part of the *m. sacrospinalis* and extends between spinous processes.

M. spinalis dorsi. Origin: fleshy from the bundles of the *m. longissimus*, which arise from the two uppermost lumbar and the two (or three) lowermost thoracic vertebrae. Insertion: grown together with *mm. semispinalis* and *multifidus*, from the 9th (or 8th) to the 2nd thoracic vertebra inclusive; one vertebra (10th or 9th) is passed over without attachment.

M. spinalis cervicis (O. T. *spinalis colli*), very variable. Origin: two uppermost thoracic and two lowermost cervical vertebrae. Insertion: 4th to 2nd cervical vertebra. Action: they extend the spinal column. Innervation: rami mediales of the rami posteriores of the nn. cervicales III—VIII and thoracales II, VI, VII, VIII.

M. spinalis capitis see under *m. semispinalis capitis*.

M. transversospinalis is the name applied to the mass of fibres running obliquely upward from the procc. transversi to the procc. spinosi and which like a three-sided prismatic cord fills the groove between the procc. spinosi and transversi. On the surface its fibres run more vertically than deep down. It is subdivided into *m. semispinalis*, *m. multifidus* and *mm. rotatores*.

M. semispinalis (see also Fig. 388) is the superficial layer of the *m. transversospinalis*; its fibres pass over more than four vertebral bodies before becoming attached. The muscle is absent in the lumbar region.


M. semispinalis dorsi. Origin: procc. transversi of the six lower thoracic vertebrae. Insertion: procc. spinosi of the six upper thoracic and two lower cervical vertebrae.

M. semispinalis cervicis (O. T. *semispinalis colli*). Origin: procc. transversi of the six upper thoracic vertebrae. Insertion: procc. spinosi of the 5th—2nd cervical vertebra.


Action: they bend the spinal column backward and toward the same side, and rotate it toward the opposite side.

Innervation: rami mediales of the rami posteriores of the nn. cervicales III—VI (VII) and thoracales (III) IV—VI, XI, XII.

M. semispinalis capitis (O. T. *complexus*) (see also Figs. 347 and 386). Form: flat, oblong-quadrangular, thick. Position: bounded medianward by the lig. nuchae, in front by *mm. semispinalis cervicis*, *recti capitis posteriores* and *obliqui capitis*, lateralward by the *m. longissimus capitis*, covered in large part by the *mm. splenii* and *trapezius*. Origin: procc. transversi of the six upper thoracic vertebrae, procc. transversi et articulares of the 3—4 lower cervical vertebrae and (sometimes absent) procc. spinosi of the uppermost thoracic and lowermost cervical vertebrae. The fibres coming from the procc. spinosi are sometimes separated so as to form an independent muscle, the *m. spinalis capitis*. Insertion: the fibres run parallel upward and somewhat medianward to the under surface of the squama occipitalis; they are interrupted, especially medianward by one or several jagged intermediate tendons. Action: it draws the head backward and raises the face toward the other side. Innervation: rami mediales et laterales of the rami posteriores of the nn. cervicales I—IV.



390. Short muscles of the back (2nd layer), viewed from behind.



M. multifidus (see also Fig. 388) is the second layer of the muscular mass extending from the procc. transversi to the procc. spinosi obliquely upward; it lies in the thorax and neck just below the m. semispinalis, in the lumbar region just below the m. longissimus dorsi; its fibres bridge over from two to four vertebral bodies. **Origins:** tuberositas iliaca, ligg. sacroiliaca posteriora, posterior surface of the sacrum, procc. mamillares of the lumbar vertebrae, procc. transversi of the thoracic vertebrae and procc. articulares of the 7th—5th cervical vertebra. **Insertions:** procc. spinosi of the lumbar, thoracic and cervical vertebrae (with the exception of the atlas). **Action:** it bends the whole spinal column backward and a trifle toward the same side and rotates the cervical portion and the thoracic portion of the spine toward the opposite side (but not the lumbar portion of the spine). **Innervation:** rami mediales of the rami posteriores of the nn. cervicales III to VIII, thoracales I—XII and lumbales I—V.

The **mm. rotatores** (see Fig. 391) form the third, deepest layer of the fibre masses going obliquely upward from the procc. transversi to the procc. spinosi; they are completely hidden by the m. multifidus. They consist of single, narrow, thin muscle bands.

Mm. rotatores longi bridge over one vertebra.

Mm. rotatores breves go to the next vertebra above.

Both go on the lumbar vertebrae from the proc. mamillaris, on the thoracic vertebrae from the root of the proc. transversus and on the cervical vertebrae from the proc. articularis inferior to the root of the proc. spinosus or to the arcus vertebrae, are present from the sacrum as far as the 2nd cervical vertebra and, below, run nearly vertically, above, more transversely. **Action and Innervation:** same as for the m. multifidus.


391. Short muscles of the back (3rd layer), viewed from behind.


Mm. levatores costarum.

Form: flat, triangular, thin.
Position: behind the articulationes costotransversariae, covered by the *m. longissimus*.
Origin: procc. transversi of the 7th cervical and 1st–11th thoracic vertebra. **Insertion:** lateralward and downward to the next rib below (*mm. levatores costarum breves*); in the 3–4 lower intercostal spaces also to the next rib but one below (*mm. levatores costarum longi*). They are directly adjacent to the *mm. intercostales externi*. **Action:** elevate the ribs. **Innervation:** ramus anterior of the *n. cervicalis VIII* and *nn. intercostales I–XI*.

Mm. interspinales, short bundles between every two neighboring spinous processes; from the epistropheus to the sacrum. They are often double on the cervical vertebrae, are usually absent from the thoracic vertebrae but very powerful on the lumbar vertebrae. **Action:** they bend the spinal column backward. **Innervation:** rami mediales of the rami posteriores of the *nn. cervicales III–VIII*, *thoracales I–XII* and *lumbales I–V*.

Mm. intertransversarii (*O. T. intertransversales*), short bundles between every two neighboring transverse processes. They are usually absent from the thoracic vertebrae. They are double on the cervical vertebrae, extending between adjacent tubercula anteriora (*mm. intertransversarii anteriores*) and also between adjacent tubercula posterora (*mm. intertransversarii posteriores*). (See Fig. 392.) They are double also on the lumbar vertebrae extending between the procc. mamillares et accessorii (*mm. intertransversarii mediales*) and between the procc. transversi (*mm. intertransversarii laterales*). **Action:** they bend the spinal column to the side. **Innervation:** rami anteriores et posteriores of the *nn. cervicales II–VIII*; rami posteriores of the *nn. thoracales*; rami anteriores et posteriores of the *nn. lumbales I–V*.





392. Short muscles of the neck, viewed from behind and below.

M. rectus capitis posterior major (O.T. rectus capitis posticus major). **Form:** flat, oblong-triangular. **Position:** immediately upon the vertebrae. **Origin:** proc. spinosus epistrophei. **Insertion:** slightly diverging, upward and lateralward to the planum nuchale of the squama occipitalis. **Action:** it draws the head backward, and rotates it (with the atlas) toward the same side. **Innervation:** n. suboccipitalis (C_1), possibly also ramus posterior n. cervicalis II.

M. rectus capitis posterior minor (O.T. rectus capitis posticus minor) (see also Fig. 391). **Form:** flat, triangular. **Position:** immediately upon the membrana atlanto-occipitalis posterior, reaching as far as the median plane; lateralward it is partially covered by the preceding muscle. **Origin:** tuberculum posterius atlantis. **Insertion:** slightly diverging, upward to the planum nuchale of the squama occipitalis. **Action:** it draws the head backward. **Innervation:** n. suboccipitalis (C_1).

M. obliquus capitis superior. **Form:** flat, oblong-triangular. **Position:** lateralward and backward from the articulatio atlantooccipitalis. **Origin:** proc. transversus atlantis. **Insertion:** slightly diverging, obliquely upward and medianward to the planum nuchale of the squama occipitalis and attaches there behind the m. rectus capitis posterior major. **Action:** it draws the head backward and bends it slightly to the same side. **Innervation:** n. suboccipitalis (C_1).

M. obliquus capitis inferior. **Form:** flat, oblong. **Position:** immediately upon the vertebrae. **Origin:** proc. spinosus epistrophei. **Insertion:** lateralward and upward to the proc. transversus atlantis. **Action:** it rotates the atlas (with the head) toward the same side. **Innervation:** n. suboccipitalis and ramus posterior n. cervicalis II (total C_1 , C_2).


393. Muscles of the right upper arm, viewed from the lateral side.

M. deltoideus (see also Figs. 359, 361, 380, 381, 394, 398 and 400). Form: flat, triangular, bent so as to be markedly through-like, thicker in the middle than at the borders. Position: just beneath the skin; it forms the fleshy projection of the shoulder. Origin: fleshy from the extremitas acromialis clavicularae and the acromion, tendinous from the spina scapulae connected with the fascia infrapinnata. Insertion: the fibres of the anterior and posterior part run in themselves converging in almost a straight line downward and lateralward, the middle fibres curve downward over the tuberculum majus humeri (*bursa subdeltoidea*, see Fig. 318); the middle part of the muscle presents a markedly multipinnate appearance; all the fibres converge to the tuberositas deltoidea humeri, lateralward and below the insertion of the m. pectoralis major; they fasten partly to a tendinous plate covering the inner surface. Action: the middle portion lifts the upper arm lateralward in the shoulder joint; the anterior portion draws the arm forward and rolls it inward, the posterior portion draws the arm backward and rolls it outward; these both portions acting together draw the arm hanging in the vertical plane medianward, but the more abducted arm lateralward. Innervation: n. axillaris (C5, C6).

Bursa subcutanea

acromialis (see Fig. 400), often present, just beneath the skin, on the upper surface of the acromion.






394. Muscles of the right axilla, viewed from below.

(The upper arm has been lifted to the horizontal plane.)


The **axilla**, when the upper arm is maintained in the horizontal plane, has the shape of a three-sided pyramid, the apex of which lies above, below the clavicle, and the base of which corresponds to the lower wall, covered only by skin and fascia. This inferior wall is bounded in front by the *m. pectoralis major* and sometimes by the *m. pectoralis minor*, behind by the *mm. latissimus dorsi* and *teres major*, medianward by the *m. serratus anterior*. Its anterior and posterior margins in the uninjured body stand out prominently laterally as the *anterior* and *posterior axillary folds*. The anterior fold is formed chiefly by the *m. pectoralis major* and sometimes the *m. pectoralis minor*, the posterior fold by the *mm. latissimus dorsi* and *teres major*. The anterior wall is formed by the *mm. pectoralis major* and *pectoralis minor*, the posterior wall by the *mm. latissimus dorsi*, *teres major* and *m. subscapularis*, the medial wall by the *m. serratus anterior*. Along the angle between the anterior and posterior wall lies the *m. coracobrachialis*; in the groove behind the latter, between it and the muscles of the posterior wall, run the large arm vessels and nerves. The axilla is filled with blood vessels, lymph vessels, lymph glands, nerves, and masses of fat. From the axilla two fissures or spaces filled with connective tissue, known as the two *axillary fissures* or *gaps* (see Fig. 401), and separated from each other by the *caput longum m. tricipitis*, lead dorsalward. The *medial axillary fissure* (for the *a. and vv. circumflexae scapulae*), is triangular and lies medially from the *caput longum m. tricipitis*, between it, the *mm. subscapularis* and *teres minor* and *mm. longissimus dorsi* and *teres major*. The *lateral axillary fissure* (for the *a. and vv. circumflexae humeri posteriores* and *n. axillaris*), is square-shaped and lies laterally from the *caput longum m. tricipitis*, between it, the *mm. subscapularis* and *teres minor*, *collum chirurgicum humeri* and *mm. latissimus dorsi* and *teres major*.



395. Fasciae of the right axilla, viewed from below.

(The drawing corresponds to Fig. 394; the upper arm is in the horizontal plane.)

Axilla (continued). The anterior surface of the *m. pectoralis major* is covered over by the thin *fascia pectoralis*. This is more powerful lateralward and extends, from the lower margin of the muscle to the lower margin of the *mm. latissimus dorsi* and *teres major*, transversely across the axilla to form the *fascia axillaris*. The latter (see also Fig. 857) is multiply perforated and presents usually a more powerful, curved, fibrous band, the (*fascia*-) *axillary arch*, which varies much in form and curvature, and extends from the region of the tendon of the *m. pectoralis major* to that of the *m. latissimus dorsi*; it is manifoldly connected with these tendons. The fascia or the axillary arch presents not infrequently bundles of muscles (*muscular axillary arch*), running from the *m. latissimus dorsi* to the insertion of the *m. pectoralis major*, less frequently from the *m. pectoralis major* to the insertion of the *m. latissimus dorsi*. The fascia axillaris goes over behind into the thin fascia of the *mm. latissimus dorsi* and *teres major*, in front into the thin covering of the *m. deltoideus* and lateralward into the somewhat stronger *fascia brachii*. This forms a sheath for the whole muscular mass of the upper arm (see also Figs. 857–860) and sends on each side a stronger strip to the *margo medialis* and the *margo lateralis humeri*, the *septum intermusculare mediale* (O. T. internal intermuscular septum) and the *septum intermusculare laterale* (O. T. external intermuscular septum), the former passing medianward and the latter lateralward between the *m. triceps brachii* on the one hand and all the other muscles of the upper arm on the other. The medial septum (see Figs. 399 and 404) lies immediately behind the insertion of the *m. coracobrachialis*, behind the *mm. brachialis* and *pronator teres* and in front of the *caput mediale m. tricipitis brachii*. The lateral septum (see Figs. 393 and 402) begins behind the insertion of the *m. deltoideus* and runs behind the *mm. brachialis*, *brachioradialis*, *extensor carpi radialis longus*, in front of the *caput laterale* and *caput mediale* of the *m. triceps brachii*. The muscles mentioned arise in part from these septa. The septa contain a great deal of fibres running along the bone, reach until to the *epicondyli humeri* and are there broadest and strongest.



396.
Muscles of
the right
shoulder-
blade,
viewed from in front.

M. subscapularis (see also Figs. 394 and 398). **Form**, flat, triangular, fan-shaped, multipinnate. **Position**: it fills up almost entirely the fossa subscapularis and passes immediately in front of the shoulder joint. **Origin**: facies costalis scapulae and fascia subscapularis (see below). **Insertion**: the fibres converge upward and lateralward and are attached by a short powerful tendon to the tuberculum minus and crista tuberculi minoris of the humerus; on the posterior surface above and laterally they are fused with the capsula articularis humeri; there the *bursa m. subscapularis* is found (see p. 198 and Fig. 397). **Action**: it rolls the arm strongly inward, abducts the arm hanging in the vertical plane, adducts the arm lifted 60° lateralward; the upper portion draws the arm slightly forward, the lower portion draws the arm from the vertical position backward, and draws the lateral lifted arm also forward. **Innervation**: nn. subscapulares (C_5 , C_6 [C_7]).

The **fascia subscapularis** (not shown in drawing) covers, as a thin membrane, the anterior surface of the m. subscapularis and is connected lateralward with the continuation of the fascia brachii, which covers the m. coracobrachialis (see also Fig. 395).

M. teres major (see also Figs. 380, 381, 394 and 397—402). **Form**: flat-cylindrical. **Position**: at the margo axillaris of the shoulder-blade, in front covered by the tendon of the m. latissimus dorsi, behind and lateralward partly by the m. latissimus dorsi, partly by the m. deltoideus and caput longum m. tricipitis brachii. **Origin**: facies dorsalis of the angulus inferior scapulae. **Insertion**: the fibres run in slight spirals lateralward and upward and are attached by a strong tendon to the crista tuberculi minoris. There, between two layers of the tendon, lies the *bursa m. teretis majoris* (not shown in drawing) which connects sometimes with the bursa m. latissimi dorsi. **Action**: it draws the arm medianward and backward and rolls it inward; it draws the vertically elevated arm downward. **Innervation**: nn. subscapulares (C_6 , C_7).


Bursa ligamenti coracoclavicularis, small, often present, between the lig. trapezoideum and the lig. conoideum of the lig. coracoclaviculare.

Bursa subacromialis, large, almost constant, between the acromion and the lig. coracoacromiale on the one hand and the tendon of the m. supraspinatus on the other. It often communicates with the following bursa. It is seldom united before the tendon of the m. supraspinatus with the joint cleft.

Bursa subdeltoidea, large, almost constant, between the m. deltoideus and the tuberculum majus humeri, often communicating with the preceding bursa.

Bursa m. coracobrachialis, of moderate size, almost constant, situated in front of the upper part of the tendon of the m. subscapularis, between it, the proc. coracoideus and the tendon of the m. coracobrachialis.


Bursa m. latissimi dorsi, fairly large, oblong, constant, between the tendons of the nn. latissimus dorsi and teres major; sometimes communicating with the bursa m. teretis majoris.



397. Muscles on the anterior surface of the right shoulder joint, viewed from in front.

(The clavícula, mm. deltoideus, pectoralis major and subscapularis have been removed.)


M. biceps brachii (see also Figs. 359, 361, 393, 394, 398 and 399). **Form:** rounded, spindle-shaped, thick. **Position:** on the anterior surface of the upper arm; above, at its origins, it is covered by the mm. pectoralis major and deltoideus. **Origin:** by two heads. **Caput longum:** from the tuberositas supraglenoidalis scapulae and from the labrum glenoidale by a long, thin tendon, which extends first lateralward within the cavity of the shoulder joint, then curves, markedly, downward in the sulcus intertubercularis (see also Fig. 282) and is accompanied in its course by a prolongation of the joint-capsule, the *vagina mucosa intertubercularis* (see also p. 198). Below the sulcus intertubercularis the tendon is fixed in its position by the m. pectoralis major (often by a tube-like duplicature of its tendon, see p. 281). **Caput breve:** by a narrow, flat, shorter tendon from the proc. coracoideus, at first fused with the m. coracobrachialis. **Insertion:** both heads extend downward and unite to form a spindle-shaped muscle-belly, the fibres of which, a little above the elbow joint, go over for the most part into a flatly rounded, thick tendon, partly, also into a superficial thin tendinous plate. The latter, *lacertus fibrosus* (O. T. semilunar fascia), runs obliquely medianward and downward in front of the m. pronator teres and becomes lost in the fascia antebrachii; the tendon proper, bending into the depth, becomes attached to the tuberositas radii (*bursa bicipitoradialis* and *bursa cubitalis interossea* see p. 321). **Action:** it lifts the upper arm forward and rolls it inward; the caput longum abducts the upper arm, the caput breve adducts it; both flex the forearm and supinate it, especially, when it is flexed. **Innervation:** n. musculocutaneus (C₅, C₆ [C₇]).



398. Muscles of the right upper arm, viewed from in front.

M. coracobrachialis (see Figs. 361, 394, 397 and 399). **Form:** flat, oblong-quadrangular. **Position:** in the axilla, covered in front by the m. pectoralis major, lateralward and in front adjoining the caput breve m. bicipitis, behind adjoining the m. subscapularis and the tendons of the mm. latissimus dorsi and teres major; below on the medial surface of the upper arm, between the mm. biceps and brachialis on the one side the caput mediale of the m. triceps on the other. **Origin:** tendinous from the proc. coracoideus scapulae, being fused there with the m. pectoralis minor and the caput breve m. bicipitis (*bursa m. coracobrachialis* see p. 318). **Insertion:** the fibres run distalward to the margo medialis humeri

and are attached just below its middle, being often partially fused there with the m. brachialis. **Action:** it lifts the upper arm forward and adducts it; it rolls the arm outward when it is lifted 60° lateralward. **Innervation:** n. musculocutaneus (C6, C7).




399. Muscles of the right upper arm

(2nd layer), viewed from in front.

M. brachialis (O. T. brachialis anticus) (see also Figs. 393 and 398). Form: flat, spindle-shaped; the muscle projects on both sides of the humerus. Position: on the anterior surface of the upper arm, covered for the most part by the m. biceps; the grooves left free in front of it, medianward and lateralward, by the side of the m. biceps, are called the *sulcus bicipitalis medialis* and *lateralis* (see Fig. 398). Origin: fleshy from the septa intermuscularia and from the facies anteriores humeri, surrounding, above, the insertion of the m. deltoideus like a horse-shoe and often fused with the insertion of the m. coracobrachialis. Insertion: the fibres converge immediately before the elbow (united there with the capsule) and go by a short tendon somewhat backward to the proc. coronoideus and to the tuberositas ulnae. Action: it flexes the forearm. Innervation: n. musculocutaneus; the lateral part is frequently supplied by a fine branch of the n. radialis (altogether C₅, C₆ [C₇]).

Bursa bicipitoradialis, constant; between the tuberositas radii and the tendon of the m. biceps.


Bursa cubitalis interossea, often present in later life; adjoining, lateralward, the tendon of the m. biceps, medianward, the ulna and chorda obliqua, behind, the m. supinator.



400.
Muscles of the
right upper arm,
viewed from behind.

M. supraspinatus (see also Figs. 381, 401 and 402). **Form:** oblong-triangular, thick, bipinnate. **Position:** in the fossa supraspinata scapulae, covered medianward by the m. trapezius, lateralward by the acromion, lig. coracoacromiale and m. deltoideus. **Origin:** fossa supraspinata and fascia supraspinata. **Insertion:** the fibres converge lateralward and, becoming tendinous, extend to the tuberculum majus humeri; there they become very closely fused laterally with the capsule of the joint. **Action:** it lifts the upper arm considerably lateralward, rolls it slightly outward and elevates it somewhat forward. **Innervation:** n. suprascapularis ([C₄] C₅ [C₆]).

M. infraspinatus (see also Figs. 380, 381 and 401). **Form:** flat, oblong-triangular, thick, bipinnate. **Position:** in the fossa infraspinata scapulae, partially covered medianward by the m. trapezius, lateralward by the m. deltoideus. **Origin:** fossa infraspinata and fascia infraspinata. **Insertion:** the fibres converge as they pass lateralward to become inserted by a strong tendon at the tuberculum majus humeri (*bursa m. infraspinati* see p. 324); they are there closely fused, particularly laterally, with the capsule of the joint. **Action:** it rolls the upper arm outward, and draws it slightly forward; the vertically hanging arm abducts the upper, adducts slightly the lower part; when the arm is lifted 60° lateralward, the entire muscle abducts. **Innervation:** n. suprascapularis ([C₄] C₅, C₆).




401. Muscles of the right upper arm, viewed from behind, the m. deltoideus having been removed.

M. teres minor (see also Fig. 400). **Form:** flat, oblong-quadrangular. **Position:** just below the m. infraspinatus, often fused with it; covered behind and lateralward by the m. deltoideus. **Origin:** facies dorsalis scapulae, close to the margo axillaris, and fascia infraspinata. **Insertion:** the fibres run lateralward and parallel to one another, are fused (particularly closely lateral) with the capsule of the joint and are attached by a short tendon to the tuberculum majus humeri. **Action:** it rolls the upper arm outward, adducts it and lifts slightly forward the arm hanging down, a little backward the laterally elevated arm. **Innervation:** n. axillaris (C5, C6 [C7]).

Fascia supraspinata (see Fig. 400) is a powerful fascia, distinctly striated medianward, which covers the m. supraspinatus; it is fused medianward with the margo superior, the margo vertebralis and the spina scapulae and lateralward becomes lost.

The **fascia infraspinata** (see Figs. 380 and 381) is attached to the margo vertebralis and to the spina scapulae, is stronger medianward, covers the mm. infraspinatus and teres minor and reaches lateralward the humerus.

M. triceps brachii (see also Figs. 393, 394, 400 and 402). **Form:** flat, oblong-quadrangular. **Position:** it occupies the whole posterior surface of the upper arm. **Origin:** by three heads. **Caput longum:** by a flat tendon from the tuberositas infraglenoidalis scapulae; it runs downward between the mm. teres minor and teres major, the fibres being parallel, is partly covered above by the m. deltoideus and overlaps the caput mediale, in part, from behind and medianward; below it continues in a tendon plate attached to the tendon area. **Action:** adducts the upper arm and draws it somewhat backward; it extends the forearm. **Caput mediale** (see Fig. 402): from the facies posterior humeri, distal from the sulcus n. radialis, as well as from the septa intermuscularia mediale et laterale; the fibres reach below close to the m. anconaeus and run, converging somewhat, to a large quadrangular flat tendon, which begins in the middle of the upper arm (continued on p. 324).



402. Muscles of the right upper arm (deep layer), viewed from behind.

(The acromion has been sawed through, the mm. infraspinatus and teres minor have been completely, the caput laterale m. tricipitis partially, removed.)

M. triceps brachii (continued).
Origin: *Caput laterale:* from the facies posterior humeri, proximal to the sulcus n. radialis, as well as from the septum intermusculare laterale; the fibres run, parallel, downward, bridging over the sulcus n. radialis (which the a. profunda brachii and the n. radialis) and extend partly over the posterior surface of the caput mediale to the upper and lateral margin of the tendon of the same. **Insertion:** fused with the joint capsule, olecranon ulnae (*bursae olecrani* see p. 333). **Action** of the entire m. triceps: it extends the forearm. **Innervation:** n. radialis (C⁶, C⁷, C⁸, [Th¹]).


Bursa m. infraspinati, small, frequently present, between the tendon of insertion of the m. infraspinatus and the capsule of the joint.


403. Muscles of the right forearm, viewed from in front.

M. brachioradialis (O. T. supinator longus) (see also Figs. 404, 408 and 409). **Form**: flat, broad above, pointed below. **Position**: superficial on the lateral surface of the upper arm and the volar surface of the forearm. **Origin**: margo lateralis humeri and septum intermusculare laterale. **Insertion**: above, it runs between the m. triceps and the m. brachialis; below, it is bounded behind by the m. extensor carpi radialis longus, ulnarward by the mm. pronator teres and flexor carpi radialis and becomes attached by a long, thin, flat tendon, beginning in the middle of the forearm, close above the proc. styloideus radii. **Action**: it flexes the forearm powerfully; when it is flexed at a right angle, it brings the hand in an intermediate position which is neighboring the extreme pronation; when the forearm is greatly flexed, it pronates powerfully; when the forearm is extended, it supinates the same. **Innervation**: n. radialis (C₅, C₆).

M. palmaris longus (see also Fig. 413) (is frequently absent). **Form**: spindle-shaped, narrow. **Position**: superficial on the volar surface of the forearm. **Origin**: epicondylus medialis humeri and fascia antebrachii, fused with its neighbors. **Insertion**: it extends above between the mm. flexores carpi radialis and ulnaris on the m. flexor digitorum sublimis, then downward and radialward upon the latter and goes over by a long, thin tendon, in front of the lig. carpi transversum, into the aponeurosis palmaris. **Action**: it pronates the forearm; it flexes the hand and makes the aponeurosis palmaris tense. **Innervation**: n. medianus (C₇, C₈, Th₁).

M. pronator teres (O. T. pronator radii teres) (see also Figs. 404 and 406). **Form**: oblong-quadrangular. **Position**: flexure of the elbow and volar surface of the forearm. **Origin**: by two heads: partly from the septum intermusculare mediale and the epicondylus medialis humeri (*caput humerale*), partly from the proc. coronoideus ulnae (*caput ulnare*) (commonly weaker); the two heads are usually separated by the n. medianus. **Insertion**: the fibres run, parallel, obliquely radialward and downward to be come attached by a short tendon to the facies volaris and lateralis radii. **Action**: it pronates the forearm and flexes it. **Innervation**: n. medianus (C₆, C₇).






404. Muscles of the right forearm, viewed from in front.

(The mm. biceps, brachialis, palmaris longus and flexor digitorum sublimis have been removed.)

M. flexor carpi radialis (see also Fig. 403). **Form**: flat, oblong-spindle-shaped, bipinnate in its lower half. **Position**: superficial on the volar surface of the forearm. **Origin**: epicondylus medialis humeri and fascia antebrachii, fused with its neighbors. **Insertion**: the fibres go, at first diverging, between mm. pronator teres and palmaris longus, radialward and downward, and then become attached bipinnate to a powerful tendon running from the middle of the forearm between the mm. brachioradialis and palmaris longus to the hand, where the tendon, covered by the lig. carpi transversum (in a separate canal of it, see p. 335), lies in the groove of the os multangulum majus; finally, covered by the tendon of the m. flexor pollicis longus, it goes to the basis oss. metacarpalis II and III (see Fig. 406). In the hand the **bursa m. flexoris carpi radialis** (see Fig. 417) lies between the bone and the tendon. **Action**: it pronates the forearm, flexes and abducts the hand. **Innervation**: n. medianus (C₆, C₇ [C₈]).

M. flexor carpi ulnaris (see also Figs. 403, 406, 409 and 411). **Form**: flat, oblong-triangular; above it forms, with its radial, anterior surface, a groove, in which lies the ulnar part of the m. flexor digitorum profundus. **Position**: superficial on the ulnar margin of the forearm. **Origin**: partly from the epicondylus medialis humeri and from the fascia antebrachii, fused with its neighbors (*caput humerale*), partly by a long, flat, thin tendon-plate which is fused with the fascia antebrachii, from the posterior surface of the olecranon and from the margo dorsalis ulnae (*caput ulnare*). **Insertion**: it runs distalward, coming in contact radialward with the m. flexor digitorum sublimis and is attached by a long, powerful tendon to the os pisiforme; the **bursa m. flexoris carpi ulnaris** (not shown in figure) is frequently to be found there. **Action**: it flexes and adducts the hand. **Innervation**: n. ulnaris (C₈, Th₁).



406. Muscles of the right forearm

(3rd layer),

viewed from in front.

(The mm. biceps, brachialis, brachioradialis, flexor carpi radialis, palmaris longus and flexor digitorum sublimis have been removed; the lig. carpi transversum has also been removed.)

M. flexor digitorum profundus (see also Figs. 404, 405 and 407). **Form:** flat, spindle-shaped. **Position:** on the volar surface of the forearm, directly upon the ulna and m. pronator quadratus; completely covered by the mm. pronator teres, flexor carpi radialis, palmaris longus, flexor digitorum sublimis and flexor carpi ulnaris. **Origin:** facies volaris and ulnaris ulnae and membrana interossea antebrachii and fascia antebrachii (inner surface of the sheet of origin of the caput ulnare of the m. flexor carpi ulnaris). **Insertion:** the fibres run almost parallel downward; from the muscle belly, from the middle of the forearm on, go off four long, flat tendons, which go beneath and along with the tendons of the m. flexor sublimis, under the lig. carpi transversum to the hollow of the hand and thence to the 2nd–5th finger; each tendon runs, at the level of the 1st phalanx, between the halves of the corresponding tendon of the m. flexor sublimis to the base of the 3rd phalanx (see Fig. 415). **Action:** it flexes the hand and adducts it slightly, flexes the 1st–3rd phalanx of the 2nd–5th finger. **Innervation:** for the 3rd, 4th and 5th finger: n. ulnaris; for the 2nd, 3rd and 4th finger: n. interosseus volaris of the n. medianus (total C₆, C₇, C₈, Th₁).

M. flexor pollicis longus (see also Figs. 405 and 407). **Form:** flat, oblong-triangular, unipinnate. **Position:** on the volar surface of the forearm, immediately upon the radius and the m. pronator quadratus; covered by the mm. pronator teres, brachioradialis, flexor carpi radialis and flexor digitorum sublimis. **Origin:** facies volaris radii and usually, by a thin bundle, from the epicondylus medialis humeri. **Insertion:** the fibres run to a long tendon beginning ulnarward; this extends, along with the tendons of the mm. flexores digitorum, below the lig. carpi transversum to the hand, then in the groove of the m. flexor pollicis brevis to the base of the 2nd phalanx of the thumb (see Fig. 415). **Action:** it flexes the hand and abducts it slightly, turns the thumb dorsalward and flexes its 2nd phalanx. **Innervation:** n. interosseus volaris of the n. medianus (C₆, C₇, C₈).

407. Muscles of the right forearm


(4th layer), viewed from in front.


(The mm. flexor digitorum profundus and flexor pollicis longus have been partially, all the more superficial muscles completely, removed.)

M. pronator quadratus (see also Fig. 405). Form: flat, quadrangular. Position: on the volar surface of the forearm near the wrist, immediately upon the radius, the ulna and the membrana interossea; covered by all the flexor muscles going to the hand. Origin: facies volaris ulnae. Insertion: the fibres run parallel, radialward and somewhat downward to the facies volaris radii. Action: it pronates the forearm. Innervation: n. interosseus volaris of the n. medianus (C6, C7, C8, Th1).

M. supinator (O. T. supinator brevis) (see also Figs. 405, 406, 410 and 411). Form: flat, bent as a hollow groove, quadrangular; it is obliquely pierced by the r. profundus n. radialis (see Fig. 877). Position: close below the elbow joint on the volar, radial and dorsal surface of the forearm, immediately upon the radius, covered by the mm. brachioradialis, extensores carpi et digitorum communis and anconaeus. Origin: epicondylus lateralis humeri, lig. collaterale radiale of the elbow joint and crista m. supinatoris ulnae. Insertion: the fibres run obliquely downward to the facies volaris, radialis and dorsalis of the radius, partially surrounding this bone in a curve. Action: it supinates the forearm. Innervation: ramus profundus n. radialis ([C5] C6, C7, C8).

M. extensor carpi radialis longus (O. T. extensor carpi radialis longior) (see Figs. 403—406, 408—410 and 412). Form: flat, oblong-spindle-shaped. Position: usually superficial on the radial side of the forearm, covered in front by the m. brachioradialis, bounded behind by the m. extensor carpi radialis brevis. Origin: septum intermusculare laterale, margo lateralis and epicondylus lateralis humeri. Insertion: the fibres run downward and converge in the middle of the forearm to form a long, flat tendon, which extends upon the facies radialis radii (there partly crossed by the mm. abductor pollicis longus and extensor pollicis brevis) under the lig. carpi dorsale (through its second compartment) to the dorsal surface of the basis oss. metacarpalis II; at the wrist the tendon of the m. extensor pollicis longus crosses over it. Action: it flexes the forearm, supinates the extended and pronates the flexed forearm; it abducts and extends the hand. Innervation: ramus profundus n. radialis ([C5] C6, C7, C8).





408. Muscles of the right forearm, viewed from the radial side.

M. extensor carpi radialis brevis (see also Figs. 403, 409 and 410). Form: spindle-shaped. Position: usually superficial on the radial side of the forearm, coming in contact in front with the m. extensor carpi radialis longus, behind with the m. extensor digitorum communis. Origin: epicondylus lateralis humeri, joint capsule and fascia antebrachii, fused with its neighbors. Insertion: the fibres go downward and converge, becoming attached in the middle of the forearm to a flat, long tendon which goes close behind that of the m. extensor carpi radialis longus through the second compartment of the lig. carpi dorsale (see p. 334) to the proc. styloideus oss. metacarpalis III; there, between the tendon and the bone, lies the *bursa m. extensoris carpi radialis brevis* (see Fig. 411), rarely absent. The tendon is crossed over by the same muscles as is that of the m. extensor carpi radialis longus (see also Fig. 412). Action: it extends the hand and abducts it. Innervation: ramus profundus n. radialis (C₆, C₇ [C₈]).

M. extensor digitorum communis (see also Figs. 409 and 412). Form: oblong-spindle-shaped. Position: superficial on the back of the forearm, radialward bounded by the m. extensor carpi radialis brevis, ulnarward by the m. extensor carpi ulnaris. Origin: epicondylus lateralis humeri, joint capsule and fascia antebrachii, fused with its neighbors. Insertion: the fibres run parallel downward and in the middle of the forearm become separated into four single bellies. Each of these goes unipinnated over into a long, flat tendon; these pass together through the fourth compartment of the lig. carpi dorsale and then diverge to go to the dorsal surface of the first phalanx of the 2nd—5th finger; each tendon sends a branch to the base of the first phalanx and each becomes attached by three processes, one in the middle going to the base of the second phalanx and two at the side converging in front and going to the base of the third. The

four tendons are connected with one another upon the ossa metacarpalia by means of three oblique bands, *juncturae tendinum* (see Fig. 412), that be-


tween the 2nd and 3rd finger being weaker, or missing. Each tendon presents a triangular expansion upon the first phalanx to which become attached the mm. lumbricales and interossei (see Fig. 419). Action: it extends the hand and adducts it slightly, extends the 1st (only slightly the 2nd and 3rd) phalanx of the 2nd—5th finger; spreads the fingers apart. Innervation: ramus profundus n. radialis ([C₆] C₇, C₈).

409. Muscles of the right forearm, viewed from behind.

M. extensor digiti quinti proprius (O. T. extensor minimi digiti). **Form**: oblong-spindle-shaped, narrow. **Position**: immediately on the ulnar side of the preceding muscle, fused with it above. **Origin**: in common with the preceding muscle. **Insertion**: the long, thin tendon runs through the fifth compartment of the lig. carpi dorsale and fuses on the back of the hand with the tendon going to the little finger from the m. extensor digitorum communis (see also Fig. 412). **Action**: it extends the 1st phalanx of the 5th finger and removes it slightly from the 4th. **Innervation**: ramus profundus n. radialis (C₇, C₈).

M. anconaeus (see also Figs. 402, 408, 410 and 411). **Form**: flat, triangular. **Position**: superficial on the back of the forearm, hidden by the fascia antebrachii, above immediately adjacent to the caput mediale of the m. triceps, radialward bounded by the m. extensor carpi ulnaris, ulnarward by the ulna. **Origin**: by a short, strong tendon from the epicondylus lateralis humeri as well as from the lig. collaterale radiale. **Insertion**: the fibres diverge to the facies dorsalis ulnae. **Action**: it extends the forearm. **Innervation**: n. radialis ([C₆] C₇, C₈ [Th₁]).

M. extensor carpi ulnaris (see also Figs. 408, 410 and 412). **Form**: flat, oblong-spindle-shaped. **Position**: superficial on the back of the forearm, radialward bounded by the m. extensor digitorum communis resp. the m. extensor digiti V proprius, ulnarward above by the m. anconaeus, below by the ulna. **Origin**: epicondylus lateralis humeri, lig. collaterale radiale of the elbow joint and fascia antebrachii, fused with its neighbors; also by a thin, tendinous leaf (fused with the fascia antebrachii) from the facies dorsalis ulnae. **Insertion**: the fibres run ulnarward and downward to the facies dorsalis ulnae and upon this go over pinnated into a long tendon which runs through the sixth compartment of the lig. carpi dorsale, radially of the proc. styloideus ulnae (see Fig. 419), to the basis oss. metacarpalis V. **Action**: it adducts the hand and extends it a little. **Innervation**: ramus profundus n. radialis ([C₆] C₇, C₈).




410. Muscles of the right forearm, viewed from behind.

(The m. extensor digitorum communis has been removed.)

M. abductor pollicis longus (O. T. extensor ossis metacarpi pollicis) (see also Figs. 403, 408, 409, 411, 412 and 414). Form: flat, spindle-shaped, bipinnated. Position: on the posterior and radial side of the forearm; behind immediately upon the membrana interossea and the radius, covered by the mm. extensor carpi ulnaris and extensor digitorum communis, bordering proximally on the m. supinator, distally on mm. extensores pollicis; radialward superficially passing over the mm. extensores carpi radiales. Origin: facies dorsalis ulnae, membrana interossea and facies dorsalis radii. Insertion: the fibres converge radialward and downward to form a flat tendon, which, at the radial side of the forearm, goes between the mm. extensores carpi radiales and flexor carpi radialis through the first compartment of the lig. carpi dorsale, partly to the m. abductor pollicis brevis, partly to the basis oss. metacarpalis I. Action: it supinates slightly the forearm; it flexes the hand slightly volarward and abducts it; it abducts the thumb and draws it dorsalward. Innervation: ramus profundus n. radialis ([C₆] C₇, C₈).

M. extensor pollicis brevis (O. T. extensor primi internodii pollicis) (see also Figs. 406, 408, 409, 411 and 412). Form: spindle-shaped. Position: just distalward from the preceding muscle. Origin: membrana interossea and facies dorsalis radii. Insertion: it runs exactly as does the preceding muscle (often inseparably fused with it) and, close to it, goes upon the back of the os metacarpale I to the base of the first phalanx of the thumb. Action: it supinates slightly the forearm; flexes the hand a trifle volarward and abducts it; it abducts the thumb and draws it dorsalward; it extends the first phalanx of the same. Innervation: ramus profundus n. radialis ([C₆] C₇, C₈).



411. Muscles of the right forearm (deep layer), viewed from behind.

(The mm. extensores carpi, digitorum communis and digiti V proprius have been removed.)

M. extensor pollicis longus (O. T. extensor secundi internodii pollicis) (see also Figs. 408—410 and 412). Form: long-spindle-shaped, pinnated. Position: ulnarward from and dorsally to the two preceding muscles, immediately upon the ulna and radius, covered by the same muscles as are they. Origin: facies dorsalis ulnae and membrana interossea. Insertion: the fibres go over into a long tendon, which passes obliquely or in a bow (around a ridge of the radius) behind the tendons of the mm. extensores carpi radiales through the third compartment of the lig. carpi dorsale and then to the base of the second phalanx of the thumb. Action: it supinates the forearm slightly; it extends the hand and abducts it; it draws the thumb dorsalward and adducts the same; it extends its phalanges. Innervation: ramus profundus n. radialis ([C₆] C₇, C₈).


M. extensor indicis proprius (O. T. extensor indicis) (see also Fig. 410). Form: oblong-spindle-shaped. Position: ulnarward from and dorsally to the preceding muscle, immediately upon the ulna, covered by the mm. extensor carpi ulnaris, extensor digiti V proprius and extensor digitorum communis. Origin: facies dorsalis ulnae and membrana interossea. Insertion: the fibres converge to form a flat tendon which goes in common with the tendons of the m. extensor digitorum communis through the fourth compartment of the lig. carpi dorsale and then fuses with the ulnar margin of the tendon of this muscle that goes to the index finger. Action: it supinates the forearm a trifle; it extends the hand and abducts it slightly; it extends the index finger and draws it ulnarward. Innervation: ramus profundus n. radialis ([C₆] C₇, C₈).

Bursa subcutanea olecrani, very frequent, upon the olecranon.

Bursa infratendinea olecrani, very frequent, in the tendon of the m. triceps.

Bursa subtendinea olecrani, occasional, in front of the tendon of the m. triceps.

Bursae subcutaneae epicondyli [humeri] lateralis et epicondyli [humeri] medialis (not shown in figure), the former occasional, the latter more frequent in the adult, upon the corresponding epicondylus humeri.




413. Fascia of the right palm.

The **ligamentum carpi volare** (see also Figs. 414 and 415) is a band of fibres running transversely in the fascia antebrachii (see Fig. 851) just over the wrist; it covers the flexor muscles and also the a., vv., and n. ulnares. Its radial portion is connected with the ligamentum carpi transversum.

The **aponeurosis palmaris** (O. T. palmar fascia) lies in the palm just beneath the skin connected with it firmly by short fibres. It is triangular and consists of two layers. Their superficial fibres running longitudinally are the continuation of the tendon of the m. palmaris longus (when it is absent they begin before the lig. carpi transversum) and go by five diverging processes partly to the skin of the fingers at the level of the capitula oss. metacarpalium. The deep transversely running layer is the continuation of the fibres of the lig. carpi transversum; near the free margin of the interdigital folds sharply limited bands, *fasciculi transversi*, are stretched out just beneath the skin between the 2nd—5th finger; they are also partially connected with one another. From the deep layer sagittal partition-walls go into the depth to the ossa metacarpalia which bound one from another canal-shaped spaces for the tendons of the mm. flexores digitorum and for the vessels and nerves; distally they connect with the ligg. vaginalia. The *thenar* and *hypothenar* eminences are covered only by thin fascia. Upon the hypothenar eminence there lies in addition the

M. palmaris brevis, flat, quadrangular. Origin: ulnar margin of the aponeurosis palmaris. Insertion: to the skin at the ulnar margin of the hand. Action: it wrinkles the skin, and deepens the hollow of the hand. Innervation: ramus superficialis of the ramus volaris manus n. ulnaris ([C₇] C₈ [Th₁]).

Lig. carpi transversum (O. T. anterior annular ligament) (see also Figs. 414—416) is a tough fibrous band which extends transversely between the eminentiae carpi and the adjacent ligaments; it bridges over the sulcus carpi (see p. 119) and sends out on the ulnar side of the tendon of the m. flexor carpi radialis, tendinous branches deeply to the ossa multangula and capitatum, so that there is a separate canal formed for this tendon.



M. palmaris longus

Ligamentum
carpi volare

Os pisiforme

M. palmaris
brevisAponeurosis
palmarisVessels and
nervesFasciculi
transversi


414. Muscles of the right palm.


M. abductor pollicis brevis (O. T. abductor pollicis). Form: flat. Position: superficial upon the ball of the thumb. Origin: lig. carpi transversum and tuberculum oss. navicularis. Insertion: radial sesamoid bone, base of the first phalanx of the thumb and tendon of the m. extensor pollicis longus. Action: it abducts the thumb and draws it volarward; it flexes its first and extends its second phalanx. Innervation: n. medianus (C6, C7 [C8]).

M. abductor digiti quinti (O. T. abductor minimi digiti) (see also Fig. 412).

Form: flat, oblong. Position: superficial upon the ball of the little finger. Origin: lig. carpi transversum and os pisiforme. Insertion: base of the first phalanx of the 5th finger and to its extensor tendons. Action: it draws the 5th finger volarward and abducts it; it flexes its 1st and extends its 2nd and 3rd phalanx. Innervation: ramus profundus of the ramus volaris manus n. ulnaris ([C7] C8 [Th1]).

M. flexor digiti quinti brevis (O. T. flexor brevis minimi digiti), frequently absent. Form: flat, oblong. Position: superficial, on the radial side of the preceding muscle. Orig.: lig. carpi transversum and hamulus oss. hamati. Insertion: base of the first phalanx of the 5th finger. Action: it flexes the first phalanx of the little finger. Innervat.: like that of the preceding muscle.






415. Muscles of the right palm (2nd layer).

(The mm. abductor pollicis brevis, abductor digiti V and flexor digiti V brevis have been removed; a piece has been cut out of the tendons of the m. flexor digitorum sublimis.)

Mm. lumbricales, four flat, narrow muscles. **Position**: in the palm between the tendons of the m. flexor digitorum profundus. **Origin**: two radial from the radial side of the corresponding tendons, two ulnar (bipinnated) each from two adjacent tendons of the m. flexor digitorum profundus. **Insertion**: the fibres converge to form narrow tendons which run, volarward from the lig. capitulum transversum (see Fig. 292), to the posterior surface of the first phalanx and to the triangular expansions of the tendons of the m. extensor digitorum communis (see Fig. 412). The 1st, 2nd and 4th m. lumbricalis goes to the radial side of the 2nd, 3rd and 5th finger. The 3rd either runs to the radial side of the 4th finger, or forks and goes to the radial side of the 4th and to the ulnar side of the 3rd finger; both kinds of insertion are equally frequent. **Action**: they flex the first and extend the second and third phalanx of the 2nd–5th finger. **Innervation**: 2(–3) radial from the n. medianus, 2(–1) ulnar from the ramus profundus of the ramus volaris manus n. ulnaris, the 3rd often from both (C₇, C₈, Th₁).

416. Deep muscles of the ball of the right thumb.

(The long flexor muscles and the superficial muscles of both eminences have been removed.)




M. opponens pollicis (see also Fig. 415). Form: flat, oblong-quadrangular. Position: in the ball of the thumb, covered by the m. abductor pollicis brevis. Origin: ligam. carpi transversum and tuberculum ossis multanguli majoris. Insertion: corpus and capitulum ossis metacarpalis I. Action: it adducts the thumb and draws it volarward. Innervation: n. medianus (C₆, C₇ [C₈, Th₁]).

M. flexor pollicis brevis (see also Figs. 414 and

415). Form: oblong-quadrangular. Position: ulnar margin of the ball of the thumb and in the depth, bounded radialward by the m. opponens pollicis, ulnarward by the m. adductor pollicis. Origin: by two heads; *superficial head*: lig. carpi transversum; *deep head*: ossa multangula, capitatum, metacarpale II. Insertion: *superficial head* to the radial sesamoid bone, *deep head* by two divisions (one narrower, radial and ohne thicker, ulnar) to the radial and ulnar sesamoid bones, both heads also to the base of the first phalanx of the thumb and partly to the tendon of the m. extensor pollicis longus. Between the two heads and upon the radial compartment of the deep head is a groove for the tendon of the m. flexor pollicis longus. Action: it draws the thumb volarward; bundle of fibres to the radial sesamoid bone abduct, those to the ulnar sesamoid bone adduct the thumb; it flexes the first and extends the second phalanx of the thumb. Innervation: superficial portions by the n. medianus, deep by the ramus profundus of the ramus volaris manus n. ulnaris (C₆, C₇, C₈).

M. adductor pollicis (see also Figs. 414 and 415). Form: flat, triangular. Position in the depth of the palm, immediately upon the bones and the mm. interossei; covered by the tendons of the mm. flexores digitorum, being bounded radialward by the preceding muscle, and frequently fused with it. Origin: os metacarpale III. Insertion: ulnar sesamoid bone, base of the first phalanx of the thumb and tendon of the m. extensor pollicis longus. Action: it adducts the thumb and draws it volarward; it flexes the first and extends the second phalanx of the thumb. Innervation: ramus profundus of the ramus volaris manus n. ulnaris (C₇, C₈, Th₁).


M. opponens digiti quinti (O. T. opponens minimi digiti) (see also Figs. 414 and 415). Form: flat, oblong-quadrangular. Position: in the ball of the little finger, almost completely covered by the mm. abductor and flexor digiti V brevis. Origin: lig. carpi transversum and hamulus oss. hamati, pierced by an opening for the rami profundus a. ulnaris and rami volaris manus n. ulnaris. Insertion: corpus and capitulum oss. metacarpalis V. Action: it draws the fifth finger volarward. Innervation: ramus profundus of the ramus volaris manus n. ulnaris ([C₇] C₈ [Th₁]).



417.
Sheaths of
the tendons
in the right
palm,
injected and colored
red.

Vagina [mucosa] tendinum mm. flexorum communium forms a common sheath for the tendons of the mm. flexores digitorum communes; it begins a little above the lig. carpi transversum and ends in the middle of the palm; very often a special sheath entirely or partially separated from the main sheath is to be found, radialward, for the tendons of the index finger (see also p. 340).


Vagina tendinis m. flexoris pollicis longi accompanies the tendon of the m. flexor pollicis longus from the upper margin of the lig. carpi transversum as far as the second phalanx; it communicates with the preceding sheath in half the cases.



418. Mm. interossei volares manus dextrae.

The *vaginae tendinum digitales* (see Fig. 417) are four long sheaths, which on the volar surface of the 2nd—5th finger from the region of the capitulum oss. metacarpalis on, surround the tendons of the mm. flexores digitorum communes in common as far as the third phalanx; that of the 5th finger is connected in the adult very often with the vagina tendinum mm. flexorum communium. The tendons are bridged over in the region of each of these tendinous sheaths, in certain places, by powerful horse-shoe-shaped connective tissue bands, *ligamenta vaginalia*, attached beside them on both sides to the bone (see p. 208), and are thereby prevented from coming away from the underlayer; the bands run, partly transversely (on the level of the joints), partly obliquely or crossed (between the joints), *ligamenta annularia* and *ligamenta cruciata manus digitorum* (see Fig. 414).

Mm. interossei volares (O. T. palmar interossei), three flat, triangular, small muscles. **Position:** in the depth of the palm, in the interspaces between the ossa metacarpalia II—V, covered by all the other muscles of the palm. **Origin:** on the lateral surfaces of the ossa metacarpalia — the first on the ulnar surface of the 2nd finger, the second and third on the radial surfaces of the 4th and 5th (see also Fig. 184). **Insertion:** each of the narrow tendons goes dorsalward from the lig. capitulum transversum (see Fig. 292) on the same side of the finger to the back of the first phalanx and to the triangular expansion of the corresponding tendon of the m. extensor digitorum communis (see Fig. 412). **Action:** they draw the first phalanx of the 2nd, 4th and 5th finger toward the middle finger, flex the first phalanx and extend the second and third phalanx of the 2nd, 4th and 5th finger. **Innervation:** ramus profundus of the ramus volaris manus n. ulnaris ([C₇] C₈, Th₁).



419. *Mm. interossei dorsales manus dextrae.*

Mm. interossei dorsales (see also Figs. 412 and 418), four flat, triangular, bipinnate, small muscles. **Position:** in the interspaces between the ossa metacarpalia, nearer the back of the hand. **Origin:** each comes from the neighboring surfaces of the ossa metacarpalia (see also Fig. 185). **Insertion:** each of the narrow tendons runs dorsalward from the lig. capitulorum transversum (see Fig. 292) — the first and second each on the radial side of the 2nd and 3rd finger, the third and fourth each on the ulnar side of the 3rd and 4th finger, to the back of the first phalanx and to the triangular expansion of the corresponding tendon of the *m. extensor digitorum communis* (see also Fig. 412). **Action:** the first adducts the thumb; the first and second draw the 1st phalanx of the 2nd and 3rd finger radialward, the third and fourth draw that of the 3rd and 4th finger ulnarward; all flex the first phalanx and extend the second and third phalanx of the 2nd—4th finger. **Innervation:** ramus profundus of the ramus volaris manus n. ulnaris ([C₇] Cs, Th₁).

Vagina tendinum mm. abductoris longi et extensoris brevis pollicis, vagina tendinum mm. extensorum carpi radialis (at the newborn one for each tendon), **vagina tendinis m. extensoris pollicis longi** (communicating constantly with the preceding), **vagina tendinis m. extensoris digiti quinti** and **vagina tendinis m. extensoris carpi ulnaris** (see Fig. 420) surround their corresponding tendons at the lig. carpi dorsale and for varying distances distal to it.

Vagina tendinum mm. extensoris digitorum communis et extensoris indicis (see Fig. 420) forms a sheath for the tendons of these muscles in common from the proximal margin of the lig. carpi dorsale as far as the middle of the ossa metacarpalia.




420. Sheaths
of the tendons
of the back of
the right hand,
injected and colored
red.

**Bursae intermeta-
carpophalangeae**, fre-
quent, one to three, are
small, lie between two adja-
cent fingers at the level of
the capitula oss. metacarpa-
lium, dorsalward from the
ligg. capitulorum transversa,
between the 2nd and the 5th
finger.


**Bursae subcutaneae
metacarpophalangeae dorsales**
occur on the back of the articulationes metacarpophalangeae, usually only on the 5th finger.

Bursae subcutaneae digitorum dorsales (see Fig. 412), just beneath the skin on
the back of the articulationes digitorum, usually small, occurring constantly or almost con-
stantly on the first finger joints (between the first and the second phalanx), occasionally on
the second joint of the 2nd and 4th finger.



421. Section through the muscles of the right inguinal region, parallel to and directly under the lig. inguinale [Poupart].

Fascia iliaca (see also Fig. 423). It covers the entire free surface of the m. iliopsoas, begins above very thinly on the arcus lumbocostalis medialis [Halleri], on the bodies of the lumbar vertebrae and on the crista iliaca, is strongest below, and passes over medianward into the fascia pelvis. Below the lateral portion of the lig. inguinale [Poupart] it extends downward with the m. iliopsoas, is attached lateralward to the spina iliaca anterior superior and to the lig. inguinale, medianward to the eminentia iliopectinea and is named, from this point on, the *fascia iliopectinea*. This divides the space under the lig. inguinale into two compartments, a lateral, *lacuna musculorum*, for the m. iliopsoas and n. femoralis, and a medial, *lacuna vasorum*; the latter contains lateralward the a. femoralis (in front of it the n. lumboinguinalis), medianward the v. femoralis and, between the latter and the lateral concave edge of lig. lacunare [Gimbernati], the *annulus femoralis*, which is closed by the fascia transversalis (*septum femorale* [Cloqueti]) and serves for the passage of lymph vessels. Below the lig. inguinale the fascia iliopectinea covers the floor of the fossa iliopectinea and forms at the same time the deep layer of the fascia lata (see also p. 348).




422. Lumbar muscles,

viewed from in front. (The largest part of the left m. psoas major has been removed.)

M. quadratus lumborum (see also Figs. 365, 366 and 384). *Form*: flat, oblong-quadrangular. *Position*: in the posterior wall of the peritoneal cavity, directly in front of the fascia lumbodorsalis, covered partly in front by the m. psoas major. Two, often slightly separated, very variable layers. *Anterior layer*. *Origin*: labium internum cristae iliacae, lig. iliolumbale, procc. transversi of the 5th—2nd lumbar vertebra. *Insertion*: inferior border of the 12th rib and the body of the 12th thoracic vertebra. *Posterior layer*. *Origin*: labium internum cristae iliacae and lig. iliolumbale. *Insertion*: procc. transversi of the 4th—1st lumbar vertebra and inferior margin of 12th rib. *Action*: it draws the last rib downward; it bends the lumbar spine lateralward and dorsalward. *Innervation*: rami musculares plexus lumbalis (Th₁₂, L₁, L₂, L₃).

M. iliopsoas consists of *m. psoas minor*, *m. psoas major* and *m. iliacus*.


M. psoas minor (see also Figs. 365, 366 and 384), inconstant. *Form*: flat, long, narrow. *Position*: directly in front of the m. psoas major. *Origin*: lateral surface of the bodies of the 12th thoracic and 1st lumbar vertebra and intervertebral disc between them. *Insertion*: fascia iliaca. *Action*: it bends the lumbar spine lateralward and forward and makes the fascia iliaca tense. *Innervation*: rami musculares plexus lumbalis (L₁, L₂).



423. Fascia iliaca dextra.

M. psoas major (see Figs. 365, 366, 384, 421, 422 and 427). **Form**: round, oblong-triangular. **Position**: in the posterior wall of the peritoneal cavity, directly in front of the mm. quadratus lumborum and iliacus, just lateral from the lumbar vertebrae and the apertura pelvis superior, the latter projecting a trifle. **Origin**: lateral surfaces of the bodies and intervertebral discs of the 12th thoracic to the 4th lumbar vertebra, small tendinous arches bridging over the aa. and vv. lumbales, as well as the procc. transversi of the lumbar vertebrae. **Insertion**: converging downward, forward and lateralward and then, fused with the m. iliacus, with a narrow tendon to the trochanter minor femoris also dorsalward. **Action**: it bends the lumbar spine lateralward and forward; it flexes the thigh considerably and rotates it slightly outward or, when the latter is fixed, it bends the pelvis and the body forward. **Innervation**: rami musculares plexus lumbalis and n. femoralis (L₂, L₃, L₄).


M. iliacus (see also Figs. 365, 421, 422 and 427). **Form**: flat, triangular. **Position**: in the fossa iliaca and anterior to the hip-joint, covered medianward by the m. psoas major. **Origin**: fossa iliaca, inferior circumference of the spina iliaca anterior inferior and capsule of the hip-joint; the latter origin (see Fig. 428) is also designated as the third head of the m. iliopsoas. **Insertion**: the fibres converge downward in front of the hip-joint (*bursa iliopsoa* see p. 350), then backward in common with the m. psoas major to the trochanter minor femoris (*bursa iliaca subtendinea* see p. 351). **Action**: it flexes considerably the thigh and rolls it a trifle outward or, when this is fixed, it draws the pelvis, together with the body, forward. **Innervation**: rami musculares n. femoralis (L₃, L₄).



424. Muscles of the true pelvis on the right side, viewed from within.

M. obturator internus (see also Figs. 425 and 436). **Form:** flat, triangular. **Position:** in the interior of the true pelvis, on its lateral wall and close behind the hip-joint. **Origin:** medial surface of the membrana obturatoria and of the os coxae in the region of the foramen obturatum. **Insertion:** the fibres converge to the incisura ischiadica minor; there they bend lateralward (*bursa m. obturatoris interni* see p. 361) at a right angle and, becoming tendinous, pass between the two mm. gemelli to the fossa trochanterica femoris. **Action:** it rotates the thigh outward, adducts and extends it somewhat. **Innervation:** rami musculares of the plexus sacralis (L5, S1, S2 [S3]).

M. coccygeus (see also Figs. 425, 723 and 724). **Form:** quadrangular, thin. **Position:** on the anterior surface of the lig. sacrospinous, forming with it a common mass of very variable composition (see p. 212). **Origin:** spina ischiadica. **Insertion:** lateral edge of lower sacral and upper coccygeal vertebrae. **Innervation:** ramus muscularis of the plexus pudendus (S3, S4 [S5]).




425. Muscles of the true pelvis

on the right side,
viewed from without and below.

M. obturator externus (see also Figs. 429—431). **Form:** triangular, thick. **Position:** anterior surface of the small pelvis and beneath the hip-joint; greatly concealed; covered in front by the mm. iliopsoas, pectineus and adductores brevis et minimus, behind of the mm. quadratus femoris and gluteus maximus. **Origin:** anterior surface of the membrana obturatoria and of the os coxae in the region of the foramen obturatum. **Insertion:** converging lateralward to the fossa trochanterica femoris. **Action:** it rolls the thigh outward; it flexes it and adducts it. **Innervation:** n. obturatorius (L_3 , L_4).

M. piriformis (see also Figs. 424, 436 and 437). **Form:** flatly-rounded, triangular. **Position:** behind the hip-joint bordering above upon the m. gluteus minimus, below upon the gemellus superior. **Origin:** facies pelvina oss. sacri, the 2nd—4th foramen sacrale anterius embracing laterally; the upper part of the incisura ischiadica major. **Insertion:** the fibres run, converging, through the foramen ischiadicum majus (see also p. 212) lateralward to the tip of the trochanter major femoris (*bursa m. piriformis* see p. 361). **Action:** it abducts the thigh outward, rolls it and extends it somewhat backward. **Innervation:** rami musculares plexus sacralis (S_1 , S_2 , S_3).



426. Fascia of the right thigh, viewed from in front.

Fascia lata [anterior half] (see also Figs. 376, 423 and 874) covers the free surface of the muscles of the thigh and is stronger lateralward and above. From near the knee a strong fibrous process runs from the region of the epicondylus medialis femoris divergingly upward and lateralward; a furrow in the skin corresponds to this when the knee is passively over extended and the muscles are relaxed; below this furrow the lowermost part of the m. vastus medialis projects as an oblong bulge running obliquely upward and lateralward (*suprapatellar bulge*). The fascia also forms a sheath for the m. sartorius behind and divides above into a deep and a superficial layer. The deep layer covers the *fossa ilipectinea*. This (see Figs. 423 and 427) possesses a triangular entrance, comes in relation above with the lig. inguinale, laterally with the m. sartorius and ends below under the latter; lateralward it is bounded by the m. iliopsoas, medianward by the m. pectineus; above it is continued into the lacuna vasorum. It is the lateral part of the larger, triangular *trigonum femorale* [*fossa Scarpa major*] (O. T. Scarpa's triangle), which is limited above by the lig. inguinale, laterally by the m. sartorius, and medially by the medial edge of the m. adductor longus. Lateralward the deep layer is identical with the *fascia ilipectinea* (see p. 343), and medianward, upon the m. pectineus, it is called the *fascia pectinea*. The superficial layer passes from the upper edge of the m. sartorius to the lower edge of the lig. inguinale and partially covers the fossa ilipectinea. It ends medianward with a concave, spiral-shaped edge, *margo falciformis* (O. T. falciform ligament) which arises below from the fascia pectinea (*cornu inferius*), is often indistinctly outlined in its middle portion and, above (*cornu superius*), runs medianward to the lig. inguinale and the lig. lacunare.

427.

Muscles of the
right thigh,

viewed from in front.


Fascia lata (continued). The fossa bounded by the margo falciformis is called the *fossa ovalis*; it is covered by the *fascia cribrosa*, a deep layer of the fascia superficialis which extends from the margo falciformis to the fascia pectinea and presents openings for lymph glands and vessels. The space from the annulus femoralis as far as the entrance to the fossa ovalis is called also the *canalis femoralis*.


Bursae praepatellares see p. 232.

Bursa infrapatellaris subcutanea (see Fig. 426), frequently present, just beneath the skin, in front of the lig. patellae.

Bursa subcutanea tuberositas tibiae (see Fig. 426), frequently present, in front of the tuberositas tibiae.

M. sartorius (see also Fig. 439). Form: flat, narrow, very long. Position: superficial, in a long spiral above on the anterior, below on the medial surface of the thigh. Origin: by a short tendon from the spina iliaca anterior superior. Insertion: the fibres run parallel, at first medianward and downward in front of the m. iliopsoas and in front of the groove between the m. quadriceps and the mm. adductores, then downward medial from the m. vastus medialis in front of the m. gracilis, then behind the epicondylus medialis, and thence obliquely forward with a flat, diverging tendon to the tuberositas tibiae and fascia cruris; the tendon covers the insertions of the mm. gracilis and semitendinosus and fuses with them (*bursa m. sartorii propria* see p. 351). Action: it flexes the thigh, abducts it and rotates it somewhat outward; it flexes the leg and rotates it inward. Innervation: n. femoralis (L_2, L_3).





428.

Muscles of the right thigh, viewed from in front.

(The mm. sartorius and tensor fasciae latae have been completely, the m. iliopsoas for the most part removed.)

Bursa iliopectinea (see also Fig. 421), constant, large, between the m. iliopsoas and the lig. iliofemorale; it is sometimes connected with the cavity of the hip-joint.

M. pectineus (see also Figs. 421 and 427). **Form:** flat, oblong-quadrangular. **Position:** medial from and below the hip-joint, bounded lateralward by the m. iliopsoas, medianward by the m. adductor longus, behind by the mm. adductor brevis and obturator externus, covered at the insertion by the m. sartorius. **Origin:** fleshy from the ramus superior oss. pubis, pecten oss. pubis and ligamentum pubicum superius (see p. 215). **Insertion:** parallel downward, lateralward and backward to the linea pectinea femoris (*bursa m. pectinei* see p. 351). **Action:** it adducts and flexes the thigh, and rotates it slightly outward. **Innervation:** n. femoralis, rarely also ramus anterior n. obturatorii (L₂, L₃).

M. gracilis (see also Figs. 427, 435, 437 and 439). **Form:** flat, long, thin. **Position:** superficial, on the medial surface of the thigh, medianward from the mm. adductores; behind it is bounded by the m. semimembranosus. **Origin:** by a broad, very thin tendon from the ramus inferior oss. pubis and oss. ischii. **Insertion:** parallel downward, behind the epicondylus medialis femoris and thence forward by a thin tendon to the tuberositas tibiae, just behind the tendon of the m. sartorius (*bursa anserina* see p. 351). **Action:** it adducts the thigh and flexes it somewhat; it flexes the leg and rotates it inward. **Innervation:** ramus anterior n. obturatorii (L₂, L₃, L₄).

429. Muscles of the right thigh, viewed from in front.

(The mm. sartorius, tensor fasciae latae, pectineus and gracilis have been removed; the knee-joint has been opened on the medial side.)

M. quadriceps femoris consists of four muscles, which are partly separated above, but inseparably united at their insertion: *m. rectus femoris*, *m. vastus lateralis*, *m. vastus intermedius* and *m. vastus medialis*. Innervation: n. femoralis (L_2, L_3, L_4).

M. rectus femoris (see also Figs. 427 and 428). Form: spindle-shaped, thick, bipinnate. Position: on the anterior surface of the thigh, covered above by the m. sartorius, otherwise superficial. Origin: tendinous, partly from the spina iliaca anterior inferior, partly arch-like above the acetabulum (*bursa m. recti femoris* see p. 352). Insertion: the fibres diverge from a middle tendon and run distalward to a tendon beginning on the dorsal surface where it goes over above the patella into the common tendon. Action: it flexes the thigh and abducts it; it extends the leg.

M. adductor longus (see also Figs. 427 and 428). Form: flat, oblong-triangular, thick. Position: on the anterior surface of the thigh, bounded behind by the mm. adductores brevis et magnus, medially by the m. gracilis, laterally above by the m. pectineus, separated from it below by a triangular hollow which varies in size; behind it is covered by the m. sartorius. Origin: by a thick, long tendon from the anterior surface of the ramus superior oss. pubis. Insertion: diverging downward and lateralward to the labium mediale of the linea aspera femoris. Action: it adducts and flexes the thigh. Innervation: ramus anterior n. obturatorii (L_2, L_3).


Bursa iliaca subtendinea, very frequently present, between the m. iliopsoas and the trochanter minor; it communicates often with the hip joint.

Bursa m. pectinei, very frequently present, between the m. pectineus and m. iliopsoas, just below the trochanter minor.


Bursa m. sartorii propria (see also Fig. 428), between the tendon of the m. sartorius and the tendons of the mm. gracilis and semitendinosus; not rarely connected with the

Bursa anserina, constant, large, between the tendons of the mm. gracilis and semitendinosus and the tibia.

Spalteholz, Atlas. 6th ed.



430. Muscles of the right thigh, viewed from in front.



(Like Fig. 429); in addition the *m. adductor longus* has been entirely, the *mm. vasti medialis* and *lateralis* have been partially removed; the knee-joint has been opened.)

M. adductor brevis. Form: triangular, thick. Position: in front covered by the *mm. pectineus* and *adductor longus*, above adjoining the *m. obturator externus*, behind the *m. adductor minimus*. Origin: fleshy from the anterior surface of the ramus inferior ossis pubis. Insertion: diverging downward and lateralward to the labium mediale of the linea aspera femoris. Action: it adducts and flexes the thigh and rotates it somewhat outward. Innervation: ramus anterior et posterior n. obturatorii (L₂, L₃, L₄).

M. adductor minimus (O. T. upper portion of adductor magnus) (see Figs. 431 and 436—438). Form: triangular, thick. Position: behind the preceding, bounded above by the *mm. obturator externus* and *quadratus femoris*, below by the *m. adductor magnus*, behind by the latter and the *m. gluteus maximus*. Origin: anterior surface of the rami inferiores ossis pubis and ossis ischii. Insertion: diverging lateralward and downward to the labium mediale of the linea aspera femoris. Action: it adducts the thigh and flexes it somewhat. Innervation: ramus posterior n. obturatorii (L₂, L₃, L₄).

M. adductor magnus (see Figs. 427, 429, 431, 435, 437 and 438). Form: triangular, thick. Position: on the medial side of the thigh, in front covered by the *mm. adductores brevis et longus* and *sartorius*, bounded above and in front by the *m. adductor minimus*, medianward by the *m. gracilis* and posteriorly by the *mm. semimembranosus*, *semitendinosus* and *biceps*; it lies above between the *m. gracilis* and the *mm. semitendinosus* and *semimembranosus* over a certain distance superficially (see Fig. 442). Origin: fleshy from the anterior surface of the ramus inferior oss. ischii and tuber ischiadicum. Insertion: the upper, anterior fibres diverge lateralward and downward to the labium mediale of the linea aspera femoris (see also p. 354), the lower, posterior fibres

converge downward to a thick tendon (see Figs. 431 and 438) which runs to the epicondylus medialis; together with the femur it forms the boundaries of the *hiatus tendineus [adductorius]* through which the vessels of the thigh pass (see also p. 354). Action: it adducts and extends the thigh. Innervation:

ramus posterior n. obturatorii and n. tibialis (total L₂, L₃, L₄, L₅, S₁).

Bursa m. recti femoris, frequently present, between the tendon of origin of the *m. rectus femoris* and the acetabulum.

431. Muscles of the right thigh, viewed from in front.


(The muscles have been removed with the exception of the *mm. obturator externus, adductores minimus et magnus* and *vastus intermedius*.)


M. vastus intermedius (O. T. *crureus*) (see also Figs. 429 and 430). **Form:** flat, oblong. **Position:** on the anterior side of the thigh, directly upon the bone; in front covered by the *m. rectus femoris*; at the sides covered over by the *mm. vasti medialis et lateralis*, being fused with them partly. **Origin:** anterior surface of the *corpus femoris*. **Insertion:** the fibres run parallel downward, covered in front by a long, thin, flat tendon, and go, above the patella, into the common tendon (see p. 354).

M. articularis genus (O. T. *subcrureus*) (see Figs. 430, 441 and 443) forms, in a sense, the deepest layer of the preceding muscle. **Form:** flat, thin. **Position:** directly upon the lower third of the femur. **Origin:** anterior surface of the *corpus femoris*. **Insertion:** upper and lateral surfaces of the capsule of the knee-joint. **Action:** it makes this capsule tense. **Innervation:** *n. femoralis* (L_3, L_4).

M. vastus lateralis (O. T. *vastus externus*) (see Figs. 427—430, 438 and 440). **Form:** flat, bent furrow-like. **Position:** on the lateral surface of the thigh, chiefly superficial, covered only by the *m. tensor fasciae latae* and the *tractus iliotibialis*, bounded behind by the *m. biceps femoris*, separated from the *m. vastus intermedius* by a connective tissue layer, which contains the *r. descendens a. circumflexae femoris lateralis*. **Origin:** anterior and lateral surface of the *trochanter major* and *labium laterale* of the *linea aspera femoris*. **Insertion:** the fibres run parallel, in a curve, medianward and downward and are attached partly to a tendinous sheet lying on the medial surface of the muscle, partly, above and lateralward from the patella to the common tendon (see p. 354).

Bursa trochanterica m. glutaiei minimi, almost constant, fairly large, between the tendon of the *m. glutaiei minimus* and the *trochanter major femoris*.






432. External muscles of the right hip, viewed from the lateral surface.

M. vastus medialis (O. T. *vastus internus*) (see Figs. 427—430, 438 and 439). **F o r m**: flat, bent furrow-like. **P o s i t i o n**: on the medial and anterior surface of the thigh; covered at its origin by the m. sartorius; otherwise superficial; comes into contact behind with the mm. adductores longus et magnus, anteriorly commonly grown together with the m. vastus intermedius. The muscle layer of the m. vastus medialis reaches further distalwards, than that of the m. vastus lateralis, often below the upper border of the patella. **O r i g i n**: labium mediale lineae asperae femoris. **I n s e r t i o n**: the fibres go, parallel, in a curve, lateralward and downward, partly to the anterior tendinous plate of the m. vastus intermedius, partly, above and medianward from the patella, to the common tendon. From the anterior surface of the m. adductor magnus, approximately from the lower margin of the m. adductor longus on, a tendinous plate, the *lamina vastoadductoria*, goes to the medial surface of the m. vastus medialis; this bridges over a triangular vascular canal, which extends obliquely backward and downward, the *canalis adductorius* [*Hunteri*] (O. T. Hunter's canal), whose distal aperture, the *hiatus tendineus* [*adductorius*], opens into the ham (see also p. 352).

The common tendon of the m. quadriceps femoris is attached to the upper and to the lateral margins of the patella and partly goes over, in front of it, directly into the lig. patellae. Some of the tendon fibres of the mm. vasti medialis and lateralis run however downward beside the patella to the anterior surface of the condyli tibiae and thus form the *retinacula patellae mediale et laterale* (see Figs. 427 and 439), of which the medial one is the broader and the more distinct, and the outer attached to the tractus iliotibialis; to these are also to be added other ligamentous bands which extend from the epicondylus forward to the patella and into which also bands of muscle are inserted.

A c t i o n of the mm. vasti: they extend the leg. **I n n e r v a t i o n** of the same: n. femoralis (L_2, L_3, L_4).




433. External muscles of the right hip, viewed from the lateral surface.

(In order to show the varying mode of attachment of the m. gluteus maximus, a piece has been cut out of the superficial layer of the tractus iliotibialis and the deeper layer has been dissected out.)

M. tensor fasciae latae (O. T. tensor vaginae femoralis) (see also Figs. 427 and 432). **Form:** flat, oblong-quadrangular, thick. **Position:** on the lateral surface of the hip, just beneath the skin; bounded in front by the mm. sartorius and rectus femoris, medianward and behind by the m. gluteus medius, and is fused with the latter at the origin. **Origin:** tendinous from the spina iliaca anterior superior and from the fascia lata. **Insertion:** the fibres run parallel downward and go over, at the junction of the upper with the middle third of the thigh, into the tractus iliotibialis [Maissiati] (see p. 356). The muscle is covered on its lateral surface by a thin fascia, on its medial surface by a thicker layer of fascia; the latter is attached above at the spina iliaca anterior inferior. **Action:** it makes the fascia lata tense, especially the tractus iliotibialis; it flexes and abducts the thigh and rotates it somewhat inward; it causes slight extension and outward rotation of the leg. **Innervation:** n. gluteus superior, often also n. femoralis (total L4, L5, S1 [S2]).

M. gluteus maximus (see also Figs. 380, 432, 435, 436 and 438). **Form:** flat, quadrangular, very thick. **Position:** just beneath the skin of the buttock, the fleshy basis of which it forms. **Origin:** lateral surface of the ala oss. ilium behind the linea glutea posterior, fascia lumbodorsalis, the posterior surface of the sacrum and coccyx and their ligaments, as well as the lig. sacrotuberosum. **Insertion:** the fibres run, nearly parallel, lateralward and downward; those of the upper portion go, in the form of a broad tendon, over the lateral surface of the trochanter major femoris (*b. trochanterica m. glutei maximi* see p. 361), curve downward and pass over into the deep layer of the tractus iliotibialis [Maissiati], hidden by the superficial layers of the same; the fibres of the lower portion are attached by a thick, flat tendon to the tuberositas glutea femoris (*bursae gluteofemorales* see p. 361). **Action:** it extends the thigh backward, rotates it outward and adducts it; it makes the fascia lata tense, especially the tractus iliotibialis; the muscles of both sides press, at fixed standing thighs, the buttocks together. **Innervation:** n. gluteus inferior ([L4] L5, S1, S2).

434. Fascia of the right thigh, viewed from behind.



Fascia lata (posterior half) (see also Figs. 432, 433 and 440) is, upon the posterior surface of the m. gluteus maximus, in general thin; distalward it is somewhat thicker. Interwoven in the fascia along the lateral surface, from the lower end of the m. tensor fasciae latae on, and from the m. gluteus maximus is a broad, thick, glistening, tendinous strip of parallel fibres, the *tractus iliotibialis* [*Maissiat*] (O. T. iliotibial band), which is attached below to the condylus lateralis tibiae. This strip consists above (see Fig. 433) of a deep layer, which is the direct continuation of tendinous fibres of the m. gluteus maximus, and of a superficial layer, which consists partly of tendinous fibres of the m. tensor fasciae latae, partly of fibres which descend from the thick fascial covering of the m. gluteus medius. From the latter region comes also a thicker, flat bundle which crosses the insertion of the m. gluteus maximus and extends in a curve from above downward and medianward to the tuber ischiadicum. This strand, cutting deep into the muscle, corresponds to the *sulcus gluteus* (O. T. gluteal groove), which is visible as a furrow, running transversely in the skin when the m. gluteus maximus is contracted (at the standing leg and, when standing on both feet, at both legs). A less marked strengthening band, in the popliteal space, extends from above and lateralward down-

ward and medianward and corresponds also to a similarly directed furrow in the skin.


Arising from the fascia lata are septa of connective tissue which extend between the individual muscles and groups of muscles; the strongest of these pass behind the origins of the mm. vastus medialis and vastus lateralis to the labium mediale and labium laterale of the linea aspera femoris, forming the *septum intermusculare* [*femoris*] *mediale* and *septum intermusculare* [*femoris*] *laterale* (see Fig. 438); they serve partly for the attachment of the neighboring muscles.


Bursa trochanterica subcutanea, frequently present, small, just beneath the skin, upon the lateral surface of the trochanter major.

435. Muscles of the right thigh, viewed from behind.

M. semitendinosus (see also Figs. 436 and 439). **Form:** triangular, long, tapering below, pinnate; it has in the middle of its dorsal surface an *inscriptio tendinea*. **Position:** on the posterior surface of the thigh; behind, in large part, superficial, covered only above by the m. gluteus maximus; lateralward it is bounded by the caput longum m. bicipitis, medianward and in front by the mm. adductor magnus and semimembranosus. **Origin:** tendinous from the tuber ischiadicum, in common with the caput longum m. bicipitis (*bursa m. bicipitis femoris superior*, see p. 361). **Insertion:** it runs converging downward, then lies with a long tendon which begins on the median side in a furrow of the m. semimembranosus and extends from there behind the tendon of the m. gracilis and in common with it, in a curve, forward to the tuberositas tibiae (see m. sartorius p. 349) (*bursa anserina* see p. 351). **Action:** it extends the thigh backward, adducts it and rotates it somewhat inward; it flexes the leg and rotates it inward. **Innervation:** n. tibialis (L4, L5, S1, S2).

M. semimembranosus (see also Figs. 430, 436, 437 and 439). **Form:** flat, oblong-quadrangular, pinnate. **Position:** on the posterior surface of the thigh; below, partly just beneath the skin, otherwise covered by the m. semitendinosus, above also by the m. gluteus maximus; lateralward below it is bounded by the m. biceps, in front by the mm. quadratus femoris, adductor magnus and gracilis. **Origin:** by a flat, long, strong tendon from the tuber ischiadicum (*bursa m. bicipitis femoris superior* see p. 361). (Continued on p. 358.)






436. Muscles of the right buttock (2nd layer), viewed from behind.

(The *m. gluteus maximus* has been cut through and reflected lateralward.)

M. semimembranosus (continued). *Insertion*: the tendon reaches downward on the lateral surface of the muscle as far as its lower half; from its medial surface fibres go medianward and downward to a tendinous strip, which begins in the upper half, runs downward on the medial surface and goes, behind the epicondylus medialis femoris between the tendons of the *mm. gracilis* and *semitendinosus* (see Fig. 439), to the condylus medialis tibiae; it is there, partly covered by the ligamentum collaterale tibiale, (bent forward) attached to the bone; in part it bends around into the lig. popliteum obliquum and in part it goes over into that portion of the fascia cruris which covers the posterior surface of the *m. popliteus* (see Fig. 309) (*bursa m. semimembranosi* see p. 366). *Action*: it extends the thigh backward, adducts it and rotates it somewhat inward; it flexes the leg and rotates it inward. *Innervation*: n. tibialis ([L₄] L₅, S₁).

M. gluteus medius (see also Figs. 380, 427, 428, 433 and 435). *Form*: flat, triangular, thick, bipinnate. *Position*: on the lateral surface of the hip; in its upper and lateral parts superficial, covered only by the fascia lata, which is here very thick; in front it is overlapped by the *m. tensor fasciae latae*, behind by the *m. gluteus maximus*, being bounded medianward by the *m. gluteus minimus* and in front very frequently fused with it. *Origin*: lateral surface of the ala oss. ilium, between the lineae gluteae anterior and posterior; fascia lata. *Insertion*: the fibres converge downward and become attached by a short tendon to the lateral surface of the trochanter major, near its apex (*bursae trochantericae m. glutei medii* see p. 361). *Action*: it abducts the thigh; the anterior portion flexes it and rotates it somewhat inward, the posterior, larger portion extends it and rotates it somewhat outward. *Innervation*: n. gluteus superior (L₄, L₅, S₁ [S₂]).




437. Muscles of the right thigh, viewed from behind.

(The mm. gluteus maximus, gluteus medius and semitendinosus have been largely, the m. obturator internus has been completely removed.)

Mm. gemelli superior et inferior (see also Fig. 436).

Form: flat, quadrangular, narrow. **Position:** on the posterior surface of the os coxae and the hip joint, bounded above by the m. piriformis, below by the m. quadratus femoris, behind covered by the m. gluteus maximus. **Origin:** *superior*: from the spina ischiadica; *inferior*: from the tuber ischiadicum. **Insertion:** the two muscles form together a groove in which runs the tendon of the m. obturator internus (*b.m. obturatoris interni* see p. 361); the fibres run, parallel, lateralward

mostly to the tendon of the m. obturator internus, partly beside this to the fossa trochanterica femoris. **Action:** they rotate the thigh outward, adduct it and extend it slightly. **Innervation:** rami musculares of the plexus sacralis (superior: L5, S1, S2 [S3]; inferior: L4, L5, S1).



438. Muscles of the right thigh, viewed from behind.

(The mm. glutei maximus et medius, piriformis, obturator internus, gemelli, semitendinosus and semimembranosus, as well as the caput longum m. bicipitis, have been removed.)

M. gluteus minimus (see also Figs. 431 and 437). **Form:** flat, triangular. **Position:** on the lateral surface of the hip, just above the hip-joint partly on the capsule of same; covered by the m. gluteus medius, in front partly fused with it; bounded in front by the mm. rectus femoris and tensor fasciae latae, behind by the m. piriformis. **Origin:** lateral surface of the ala oss. ilium between the lineae gluteae anterior and inferior. **Insertion:** the fibres converge downward, lateralward and forward, go with a strong tendon to the anterior surface of the trochanter major femoris (*b. trochanterica m. glutei minimi* see p. 353). **Action:** it abducts the thigh, the much larger anterior portion flexes and rotates it somewhat inward, the posterior portion extends it and rotates it somewhat outward. **Innervation:** n. gluteus superior (L4, L5, S1 [S2]).

M. quadratus femoris (see also Figs. 430, 436 and 437). **Form:** flat, quadrangular. **Position:** behind and below the hip-joint, bounded in front by the m. obturator externus, below by the m. adductor minimus, behind by the mm. semimembranosus, biceps and gluteus maximus, above by the m. gemellus inferior. **Origin:** tuber ischiadicum. **Insertion:** the fibres run parallel lateralward to the crista intertrochanterica femoris. **Action:** it rotates the thigh outward, adducts it and flexes it somewhat from the extended position; it draws the flexed leg

downward. **Innervation:** rami musculares of the plexus sacralis (L4, L5, S1).

Bursa ischiadica m. glutei maximi (see also Figs. 436 and 437), very frequently present, upon the posterior surface of the tuber ischiadicum and of the origins of the mm. biceps and semitendinosus, between these and the m. gluteus maximus.



439. Muscles of the right knee, viewed from the medial surface.

Bursa trochanterica m. glutei maximi (see Figs. 436—438), constant, very large, upon the posterior and lateral surface of the trochanter major as well as upon the origin of the m. vastus lateralis, between these and the upper portion of the m. gluteus maximus.

Bursae gluteofemorales (see Figs. 437 and 438), constant, two or three small bursae below the preceding; partly lateral, partly medial from the attachment of the m. gluteus maximus to the femur.


Bursa trochanterica m. glutei medii anterior (see Fig. 437), constant, sometimes double; between the tendon of the m. gluteus medius and the apex of the trochanter major.

Bursa trochanterica m. glutei medii posterior (see Fig. 437), very frequently present, between the tendon of the m. gluteus medius and the tendon of the m. piriformis.

Bursa m. piriformis (see Fig. 437), frequently present, in front of and below the tendon of the m. piriformis, between it, the bone and the m. gemellus superior.

Bursa m. obturatoris interni (see Fig. 437), constant, between the incisura ischiadica minor and the m. obturator internus; it is very often continued into the groove formed by the mm. gemelli, for which it forms a lining; the latter part is very rarely independent.

Bursa m. bicipitis femoris superior (see Fig. 437), inconstant, between the tendons of the mm. biceps and semitendinosus on the one hand and the tendon of the m. semimembranosus on the other.




440. Muscles of the right knee, viewed from the lateral surface.

(The fascia lata with the exception of the tractus iliotibialis has been removed.)

M. biceps femoris (see also Figs. 435—438). **Form**: triangular-prismatic, long, with two heads. **Position**: on the posterior surface of the thigh, chiefly superficial, covered only above by the m. gluteus maximus; bounded medianward by the mm. semitendinosus and semimembranosus, lateralward by the m. vastus lateralis, in front by the mm. quadratus femoris and adductor magnus. **Origin**: *caput longum*: by a short, strong tendon from the tuber ischiadicum, in common with the m. semitendinosus (*bursa m. bicipitis femoris superior* see p. 361); the tendon is continued on the medial surface of the muscle as far as its inferior half; from it muscle fibres run out lateralward and downward to a tendon which begins on the lateral surface in the upper half of the muscle; *caput breve*: fleshy, from the labium laterale of the linea aspera femoris; the fibres go backward and downward to the tendon of the caput longum. **Insertion**: the lowermost, common part extends downward behind the epicondylus lateralis femoris (*b. bicipitogastrocnemialis* see p. 366) and goes by a strong, rounded tendon to the capitulum fibulae; in so doing it spans the lig. collaterale fibulare from behind and lateralward by two limbs (*bursa m. bicipitis femoris inferior* see p. 363). **Action**: it extends the thigh backward, adducts it and rotates it outward; it flexes the leg and rotates it outward. **Innervation**: *caput longum*: n. tibialis ([L5] S1, S2, S3); *caput breve*: n. peroneus communis (L4, L5, S1).

441. Muscles of the right leg,


viewed from the lateral surface.



The **retinacula mm. peroneorum** are strong bands of the fascia cruris and fascia dorsalis pedis which hold the mm. peronei firmly in their bony grooves. The **R. superius** (see Figs. 444, 446 and 460) goes, from the malleolus lateralis, partly into the deep layer of the fascia cruris, partly to the lateral surface of the calcaneus. The **R. inferius** (see also Figs. 315 and 460) forms a bridge over the mm. peronei on the lateral surface of the calcaneus, is fastened to the bone in front and behind them and sends a septum between the two tendons to the proc. trochlearis calcanei. At the front above it is connected with the superficial layer of the lig. cruciatum.

Bursa m. bicipitis femoris inferior (see also Fig. 442), almost constant, on the caputulum fibulae between the tendon of the m. biceps femoris and the lig. collaterale fibulare.

M. tibialis anterior (O. T. tibialis anticus) (see also Figs. 442, 452, 453 and 456). Form: triangular-prismatic, oblong. Position: superficial on the anterior surface of the leg; bounded medianward by the tibia, lateralward above by the m. extensor digitorum longus, below by the m. extensor hallucis longus. Origin: condylus lateralis and facies lateralis tibiae, membrana interossea cruris and fascia cruris. Insertion: the fibres converge downward to a flat tendon, which extends medianward, from the lower third of the leg on, in front of the tibia and articulo talocruralis through the medial compartment of the lig. transversum and of the lig. cruciatum cruris, to the plantar surface of the os cuneiforme I (*b. subtendinea m. tibialis anterioris* see p. 375) and mainly to the plantar surface of the basis oss. metatarsalis I (see also Figs. 319 and 456). Action: it flexes the foot dorsalward and can supinate it somewhat, as well as pronate it a trifle form an intermediate position. Innervation: n. peroneus profundus (L₄, L₅ S₁).



442. Muscles of the right leg, viewed from in front.

M. extensor digitorum longus

(see also Figs. 441, 452 and 453).

Form: flat, oblong, pinnate.

Position: just beneath the

skin, on the lateral surface of

the leg; bounded medianward

above by the m. tibialis anterior,

below by the m. extensor hal-

lucis longus, lateralward above by

the m. peronaeus longus, below by

the m. peronaeus brevis. Origin:

fleshy, from the condylus lateralis

tibiae, capitulum and crista anterior

fibulae, septum intermusculare [fi-

bulare] anterius, membrana inter-

ossea cruris and the inner surface

of the fascia cruris. Insertion:

the fibres extend downward and for-

ward to a tendon beginning median-

ward in the middle of the leg, which

runs, through the lateral compart-

ment of the lig. transversum and of

the lig. cruciatum cruris, forward

and downward and becomes subdivi-

ded below into five tendons. Of these,

the four medial tendons diverge to

the heads of the ossa metatarsalia II

to V, unite at the base of the first

phalanx of the 2nd—4th toe with

the corresponding tendons of the m.

extensor digitorum brevis, pass for-

ward upon the dorsal surface of the

phalanges of the toes, and each be-

comes attached by a middle process

to the second, and by two lateral

processes to the third phalanx.

Upon the first phalanx each tendon

forms, as in the fingers (see p. 330),

a triangular expansion, into the late-

ral processes of which the mm. lum-

bricales and interossei may be inser-

ted. The fifth tendon, that situated

farthest lateralward, and the muscle

belonging to it, usually incomplete-

ly separated above, namely the m.

peronaeus tertius (see also Figs.

441, 452 and 460), extends to the

dorsal surface, the base of the os

metatarsale V. Action: it flexes

the foot dorsalward, pronates and

abducts it; it draws the 2nd—5th toe some-

what dorsalward. Innervation: nerv.


peronaeus profundus (L₄, L₅, S₁).


443. Muscles of the right leg (2nd layer), viewed from in front.

(The mm. tibialis anterior, extensor digitorum longus and peroneus longus have been removed; the lig. cruciatum cruris has been partially cut open.)

M. extensor hallucis longus (O. T. extensor proprius hallucis) (see also Figs. 441, 442, 452, 453 and 456). **Form:** flat, oblong, pinnate. **Position:** on the anterior surface of the leg, below just beneath the skin, above overlapped by its two neighbors; bounded lateralward by the m. extensor digitorum longus, medianward by the m. tibialis anterior. **Origin:** membrana interossea cruris and facies medialis fibulae. **Insertion:** the fibres run parallel forward and downward to a flat tendon beginning medianward in the inferior third; this extends, through the lateral compartment of the lig. transversum and through the middle compartment of the lig. cruciatum cruris, forward and medianward on the dorsal surface of the os metatarsale I to the base of the second phalanx of the great toe; with the first phalanx it is connected often by thin bands (see also Fig. 459). **Action:** it flexes the foot dorsalward, pronates and abducts it somewhat; it extends the first toe dorsalward. **Innervation:** n. peroneus profundus (L₄, L₅, S₁).

The **fascia cruris** (see also p. 363 and Figs. 875 and 880) covers in the front the free surface of the mm. tibialis anterior, extensores and peronei. It is connected at the knee with the fascia lata, is stronger above than below and, near the ankle-joint, is strengthened by a broad band of transverse fibres, the **ligamentum transversum cruris** (O. T. broad part of anterior annular ligament) (see Figs. 442, 452 and 453). This band sharply limited generally only below, is attached at its ends to the fibula and tibia and is divided, by a septum going to the tibia, into two compartments: one medial for the m. tibialis anterior; one lateral for the mm. extensor hallucis longus and extensor digitorum longus; the lower margin of the ligament is connected medianward with the upper medial limb of the lig. cruciatum cruris. *On the back of the leg* the fascia forms a superficial and a deep layer which unite below behind the two bones (see also p. 378). The superficial layer covers the dorsal surface of the m. triceps surae, the deeper layer covers the mm. flexores longi and tibialis posterior, the aa. and vv. tibiales posteriores and peroneae and the n. tibialis. Medianward, upon the facies medialis tibiae, the fascia is fused with the periosteum. Lateralward it sends a strip to the fibula, in front of and behind the mm. peronei, **septum intermusculare [fibulare] anterius** and **septum intermusculare [fibulare] posterius** (see also Fig. 447) from which the muscles arise in part.





444. Muscles of the right leg, viewed from behind.

M. triceps surae, consists of two muscles, *m. gastrocnemius* and *m. soleus*, which are separated at their origins, but are inserted by means of a common tendon, *tendo calcaneus* [Achillis], into the middle of the posterior surface of the tuber calcanei (*b. tendinis calcanei* [Achillis] see p. 369).

M. gastrocnemius (see also Figs. 309, 435 and 439—442). **Form**: flat, long, with two heads. **Position**: superficial on the back of the leg; it forms the main mass of the calf. **Origin**: tendinous, from the planum popliteum femoris, fused with the capsule of the joint; *caput laterale*: above the condylus lateralis (*b. m. gastrocnemii lateralis* see p. 368) (it frequently contains a small sesamoid bone, *fabella*), *caput mediale*: above the condylus medialis (*b. m. gastrocnemii medialis* see p. 367) (it contains very seldom a small sesamoid bone). **Insertion**: fibres for each head arise also from a tendinous strip lying on the surface; they diverge below to form a broad tendon, which begins high up on the anterior surface of the muscle, becomes narrower and thicker below and goes over into the tendo calcaneus; the two heads converge, from their origins on, downward and are separated below only by a narrow groove; the belly of the medial head extends somewhat farther distalward. **Action**: it flexes the leg; it flexes the foot plantarward, supinates and adducts it. **Innervation**: n. tibialis (S₁, S₂).

Bursa m. semimembranosi (see also Figs. 438 and 445), constant, double; one between the *m. semimembranosus* and *caput mediale m. gastrocnemii*, in front coming in contact with the capsule of the joint (it may communicate with the joint cavity); the other, between the tendon of the *m. semimembranosus* and the condylus medialis tibiae (see Fig. 447).

Bursa bicipitogastrocnemialis, very rare, between the tendon of the *m. biceps* on the one side, the *caput laterale m. gastrocnemii* and *m. plantaris* on the other side; it is bounded medianward and behind by the n. peroneus communis.

Bursa subcutanea calcanea (not illustrated), constant, in the sole of the foot, just beneath the skin on the inferior surface of the tuber calcanei.

Bursa subcutanea tendinis calcanei [Achillis] (see also Fig. 460), frequently present, on the posterior surface of the tuber calcanei.

445. Muscles of the right leg (2nd layer), viewed from behind.

(The m. gastrocnemius has been removed.)


M. soleus (see also Figs. 440 to 442 and 444). Form: flat, spindle-shaped, thick. Position: on the posterior side of the leg, almost completely covered by the m. gastrocnemius, bounded above by the m. popliteus. Origin: from the capitulum and facies posterior fibulae, linea poplitea tibiae and from a tendinous arch between the two. Insertion: backward and downward to a broad superficial tendinous sheet, which becomes narrower and thicker below and goes over into the tendo calcaneus. Action: it flexes the foot plantarward, supinates and adducts it. Innervation: n. tibialis (L_5, S_1, S_2).


M. plantaris (see also Figs. 435, 441 and 444). Form: flat, short, spindle-shaped. Position: on the back of the leg, bounded above lateralward by the caput laterale m. gastrocnemii, in front by the capsule of the joint and the m. popliteus. Origin: planum popliteum femoris above the condylus lateralis, fused with the capsule of the joint. Insertion: medianward and downward to a long, narrow tendon which passes upon the posterior surface of the m. soleus, in front of the m. gastrocnemius, to the medial side of the tendon of the latter and goes over into the tendo calcaneus; in addition some fibres go to the fascia. Action: like the m. gastrocnemius; it makes the fascia tense. Innervation: n. tibialis (L_5, S_1, S_2).

M. popliteus (see also Figs. 309, 312, 443, 446 and 447). Form: flat, triangular. Position: on the back of the knee-joint and of the tibia, covered by the mm. plantaris and gastrocnemius, bounded below by the m. soleus. Origin: by a narrow tendon from the epicondylus lateralis femoris (*b. m. poplitei* see Figg. 312 and 447), in addition by a fleshy origin from the lig. popliteum arcuatum. Insertion: diverging medianward and downward to the facies posterior tibiae. Action: it flexes the leg and rotates it inward. Innervation: n. tibialis (L_4, L_5, S_1).

Bursa m. gastrocnemii medialis, almost constant, between the uppermost part of the capsule of the knee-joint, the femur and the caput mediale m. gastrocnemii; it very frequently communicates with the cavity of the knee-joint.

Spalteholz, Atlas. 6th ed.





446. Muscles of the right leg (3rd layer), viewed from behind.

(The mm. gastrocnemius, plantaris and soleus have been removed.)

M. flexor digitorum longus (see also Figs. 444, 450, 451, 456 and 457). **Form:** flat, oblong, bipinnate. **Position:** on the back of the tibia and of the m. tibialis posterior (crossing it obliquely), bounded laterally by the m. flexor hallucis longus and often united with it, covered by the m. soleus. **Origin:** fascies posterior tibiae and the deep layer of the fascia cruris (see p. 365). **Insertion:** obliquely downward to a tendinous strip, which begins at the medial side of the muscle and goes over above the malleolus into a rounded tendon; this extends, behind and below the tendon of the m. tibialis posterior, on the medial surface of the lig. talotibiale posterius and at the medial margin of the sustentaculum tali of the calcaneus (see p. 155) to the sole of the foot, being held fast by the deep layer of the lig. laciniatum (see also Figs. 319 and 320). It then crosses, plantarward, the tendon of the m. flexor hallucis longus, receives from it a tendinous bundle (seldom gives off a similar bundle) and breaks up into four diverging tendons; these go to the heads of the 2nd—5th os metatarsale, perforate the corresponding tendons of the m. flexor digitorum brevis and always become fastened to the base of the 3rd phalanx. The tendon is covered on the medial border of the foot by the m. abductor hallucis, in the sole plantarward by the m. flexor digitorum brevis. **Action:** it flexes the foot plantarward, supinates and adducts it; it draws the 3rd phalanx of the 2nd—5th toe plantarward. **Innervation:** n. tibialis (L₅, S₁).

M. flexor hallucis longus (see also Figs. 445, 450, 451 and 456—458). **Form:** above flat, below rounded, oblong, bipinnate. **Position:** on the posterior surface of the fibula and of the m. tibialis posterior; it is bounded medianward by the m. flexor digitorum longus (s. above), lateralward by the mm. peronei, covered by the m. soleus. **Origin:** facies posterior fibulae, septum intermusculare posterius and the deep layer of the fascia cruris (see p. 365). **Insertion:** the fibres converge downward to a tendinous strip. Behind the malleolus medialis begins at the medial side of the muscle the rounded tendon, which goes in the sulcus m. flexoris hallucis longi of the talus and calcaneus, held in position by the deep layer of the lig. laciniatum (see also Figs. 319 and 320), behind and under the tendon of the m. flexor digitorum longus, to the sole of the foot. There it crosses dor-

salward the tendon of the latter muscle, gives off a process to it (see above), which goes especially to the tendon bundle for the 2nd toe (often also for the 3rd toe) and extends between the bellies of the m. flexor hallucis brevis and the two sesamoid bones to the base of the 2nd phalanx of the 1st toe. At the margin of the foot it is covered by the m. abductor hallucis, in the sole plantarward in part by the m. flexor digitorum brevis. **Action:** it flexes the foot plantarward, supinates and adducts it; it draws the 1st and 2nd (and 3rd) toe plantarward. **Innervation:** n. tibialis (L₅, S₁, S₂).

Bursa m. gastrocnemii lateralis, often present, between the uppermost part of the capsule of the knee-joint and the caput laterale m. gastrocnemii; it frequently communicates with the cavity of the joint.

447. Muscles of the right leg

(4th layer), viewed from behind.


(All the muscles have been removed with the exception of the mm. tibialis posterior and peronaeus brevis.)

M. tibialis posterior (see also Figs. 446 and 456—458). **Form**: flat, oblong, penniform. **Position**: directly upon the posterior surface of the bones of the leg and of the membrana interossea cruris, above between the mm. flexor digitorum longus and flexor hallucis longus, covered below by these, above and below also by the m. soleus. **Origin**: facies posterior tibiae, membrana interossea cruris and facies medialis fibulae. **Insertion**: the fibres run obliquely downward to a tendinous strip situated on the medial margin; the round tendon, going out of it above the malleolus medialis, extends directly upon the tibia, in front of the m. flexor digitorum longus, downward and medianward to the sulcus malleolaris tibiae. It is there held in place by the lig. laciniatum, goes in a curve to the medial side of the ligg. talotibiale posterius and calcaneotibiale, thence on the medial and inferior surface of the lig. calcaneonaviculare plantare to the sole of the foot and, lying directly upon the bones and ligaments, it sends strong processes to the tuberositas oss. navicularis (there often a sesamoid cartilage, sometimes a sesamoid bone, see p. 160) and to the os cuneiforme I, and feebler processes to the ossa cuneiformia II and III, cuboideum and metatarsalia II, III, IV (see also Figs. 245 and 319). **Action**: it flexes the foot plantarward, supinates and adducts it. **Innervation**: n. tibialis (L_5, S_1).


M. peronaeus longus (see Figs. 441, 442, 444—446, 451, 458 and 460). **Form**: flat, oblong, bipinnate. **Position**: superficial on the lateral surface of the leg, above upon the fibula, below upon the m. peronaeus brevis; bounded in front by the m. extensor digitorum longus, behind by the m. soleus. **Origin**: capitulum and facies lateralis fibulae, fascia cruris and septa intermuscularia [fibularia]. **Insertion**: the fibres go obliquely downward to a tendinous strip; the flat tendon extends above on the lateral, below on the posterior surface of the m. peronaeus brevis downward, then behind and below the malleolus lateralis, upon the lig. calcaneofibulare in a curve to the sulcus m. peronaei [longi] of the calcaneus and is held in place by the retinacula mm. peronaeorum (see p. 363); thence it runs in the sulcus m. peronaei [longi] upon the tuberositas oss. cuboidei to the sole of the foot and there, covered plantarward by the mm. adductor hallucis (caput obliquum) and flexores digitorum as well as by the lig. plantare longum, it goes obliquely forward to the medial border of the foot; it is attached to the os cuneiforme I and tuberositas oss. metatarsalis I (see Fig. 319). In front of the tuberositas oss. cuboidei it often contains a sesamoid cartilage, rarely a sesamoid bone. **Action**: it flexes the foot plantarward, pronates and abducts it. **Innervation**: n. peronaeus superficialis (L_4, L_5, S_1).

Bursa m. poplitei (see also Fig. 312) see p. 232.

Bursa tendinis calcanei [Achillis] (see Fig. 446), constantly present, large, between the posterior surface of the tuber calcanei and the tendo calcaneus [Achillis].



448. Fascia of the sole of the right foot.



M. peroneus brevis (see Figs. 441—447, 452, 453 and 460).
Form: flat, oblong, bipinnate.
Position: on the lateral side of the leg upon the fibula, partly superficial, partly covered by the m. peroneus longus; bounded in front by the mm. extensor digitorum longus and peroneus tertius, behind by the m. flexor hallucis longus. **Origin:** facies lateralis fibulae and septa intermuscularia [fibularia]. **Insertion:** the fibres converge downward to a tendinous strip; the muscle extends medianward and in front of the tendon of the m. peroneus longus, in a curve behind and below the malleolus lateralis (in a groove on the same) and then forward, directly upon the lig. calcaneofibulare and above the proc. trochlearis calcanei; it is held in place by the retinacula mm. peroneorum (see p. 363); thence the rounded tendon goes downward and forward to the tuberositas oss. metatarsalis V; it gives off, usually, also a process to the tendon of the m. extensor digitorum longus going to the little toe (see Fig. 460). **Action:** it flexes the foot plantarward, pronates and abducts it. **Innervation:** n. peroneus superficialis (L₄, L₅, S₁).

The **aponeurosis plantaris** is a thick layer of glistening tendinous fibres just beneath the skin; it covers the inferior surface of the muscles of the sole of the foot. It begins at the tuber calcanei, extends forward, diverging with five processes, and there becomes attached partly to the skin beneath the heads of the ossa metatarsalia, but especially by means of bundles branching off into the depth to go to the ligg. vaginalia. It is strongest in the middle and feebler over the muscles of the great toe and of the little toe; it sends tough septa into the depth near the m. flexor digitorum brevis; otherwise it is connected behind with the fascia


cruris and at the sides with the fascia dorsalis pedis. Between its cutaneous insertions, there are stretched out, as in the hand (see p. 335), three bands of fibres which run transversely, the *fasciculi transversi*.

Bursa subcutanea calcanea see p. 366.

449. Muscles of the sole of the right foot.


M. flexor digitorum brevis (see also Figs. 450 and 451). Form: oblong-quadrangular, thick. Position: in the sole of the foot, covered only by the aponeurosis plantaris; bounded medianward by the m. abductor hallucis, lateralward by the m. abductor digiti V, dorsalward by the m. quadratus plantae and tendo m. flexoris digitorum longi. Origin: processus medialis tuberculi calcanei and aponeurosis plantaris. Insertion: the fibres diverge in front into four muscular bellies and flat tendons which go to the 2nd—5th toe; each tendon subdivides at the level of the first phalanx into two processes, permitting the corresponding tendons of the m. flexor digitorum longus to pass through between them, and become attached in the same manner as the corresponding tendons of the hand (see p. 360) to the second phalanx. The muscle belly and the tendon of the 5th toe are usually very weak, and may even be missing. Action: it draws the second phalanx of the 2nd to 5th toe plantarward. Innervation: for 2nd—4th toe: n. plantaris medialis (L₅, S₁); for 5th toe: n. plantaris lateralis (S₁, S₂).

M. abductor digiti quinti (O. T. abductor minimi digiti) (see also Figs. 450, 453 and 460). Form: flatly rounded, oblong. Position: on the lateral margin of the foot, covered only by the aponeurosis plantaris, bounded medianward by the mm. flexor digitorum brevis and quadratus plantae. Origin: inferior surface of the calcaneus, aponeurosis plantaris. Insertion: forward and lateralward partly to the tuberositas oss. metatarsalis V, partly by a half-tendinous insertion to the base of the first phalanx of the 5th toe. Action: it draws the first phalanx of the little toe plantarward and eventually lateralward. Innervation: n. plantaris lateralis (S₁, S₂).



450. Muscles of the sole of the right foot (2nd layer).

(The m. flexor digitorum brevis has been removed down to the tendons.)




M. quadratus plantae (O. T. accessorius or flexor accessorius digitorum pedis) (see also Fig. 451). Form: flat, quadrangular. Position: in the sole of the foot, bounded plantarward by the m. flexor digitorum brevis, dorsalward by the lig. plantare longum and caput obliquum m. adductoris hallucis, lateralward by the m. abductor digiti V. Origin: by two processes from the inferior and medial surfaces of the calcaneus. Insertion: forward to the lateral margin of the tendon of the m. flexor digitorum longus. Action: it draws the third phalanx of the 2nd to 5th toe plantarward. Innervation: n. plantaris lateralis (S₁, S₂).

Mm. lumbricales, four. Form: spindle-shaped, narrow. Position: between the terminal tendons of the m. flexor digitorum longus, bounded plantarward by the m. flexor digitorum brevis, dorsalward by the mm. adductor hallucis and interossei. Origin: partly from the medial margin, partly from the opposed margins of the tendons of the m. flexor digitorum longus. Insertion: small tendons extend forward, plantarward from the ligg. capitulum [oss. metatars.] transversa, to the medial surface of the first phalanx of the 2nd—5th toe and sometimes go over in the triangular expansions of the tendons of the mm. extensores digitorum. Action: they

generally only flex the first phalanx of the 2nd to the 5th toe. Innervation: one medial muscle: n. plantaris medialis; three lateral: n. plantaris lateralis (total L₅, S₁, S₂).

452. Muscles of the back of the right foot.



The *fascia dorsalis pedis* (see also p. 363) is stretched out in the form of a membrane, for the most part thin, over the tendons and the muscles of the back of the foot; above it is connected with the fascia cruris, at the sides with the aponeurosis plantaris. Interwoven in it, at the junction of the foot with the leg, is situated the powerful *ligamentum cruciatum cruris* (O.T. lower part of anterior annular ligament) (see also Figs. 443, 453, 456 and 460). This usually has the shape of a Y. Its lateral, strongest limb consists of two layers, of which the more superficial arises on the lateral surface of the calcaneus (connected with the retinaculum mm. peroneorum inferius), the deep one arising in the sinus tarsi; both extend medianward, the former in front of, the latter behind the tendons of the *m. extensor digitorum longus* and unite on the medial side of the same, then bifurcates into two medial limbs. The inferior medial, more delicate, limb runs in front of the tendons of the mm. *extensor hallucis longus* and *tibialis anterior* to the os naviculare and os cuneiforme I (see also Fig. 456). The superior medial, stronger limb goes to the anterior surface of the malleolus medialis (see also: fascia cruris p. 365); it consists likewise of two layers, which run in front of and behind the mm. *extensor hallucis longus* and *tibialis anterior*; the main mass of fibres usually runs in front of the *m. extensor hallucis longus* and behind the *m. tibialis anterior*. Going off from the medial surface of the os cuneiforme I and os metatarsale I is a curved strengthening band of fibres which passes in front of the mm. *tibialis anterior* and *extensores hallucis* to the basis oss. metatarsalis II.

453. Muscles of the back of the right foot (2nd layer).


(The mm. tibialis anterior and extensores longi as well as the inferior limb of the lig. cruciatum cruris have been removed.)

M. extensor digitorum brevis (see also Figs. 441, 452 and 460). **Form:** flat, triangular. **Position:** upon the back of the foot, directly upon the bones, partly covered by the tendons of the mm. extensor digitorum longus and peroneus III. **Origin:** lateral and superior surface of the corpus calcanei, in front of the entrance into the sinus tarsi, covered by the lateral limb of the lig. cruciatum cruris. **Insertion:** the fibres diverge forward and form four small muscular bellies, which, upon the ossa metatarsalia, go over into four flat, delicate tendons. Of these, the three lateral run to the first phalanx of the 2nd–4th toe, fuse with the corresponding tendons of the m. extensor digitorum longus and go with the latter to the second and third phalanx. The medial, strongest tendon, along with the muscle belonging to it, which is mostly distinctly separated is given a special name, being called the **m. extensor hallucis brevis**; it extends obliquely medianward and becomes attached to the base of the first phalanx of the great toe (see also Fig. 454). **Action:** it draws the 1st–4th toe dorsalward and lateralward. **Innervation:** n. peroneus profundus (L₄, L₅, S₁).

Bursa subtendinea m. tibialis anterioris (see also Fig. 457), very frequently present, between the medial surface of the os cuneiforme I and the tendon of the m. tibialis anterior.

Bursae intermetatarsophalangeae (see also Figs. 458 and 459), four, between the heads of the ossa metatarsalia; the three medial almost constant, the fourth only occasionally present.






454. *Mm. interossei dorsales pedis dextri*,

viewed from the dorsal surface of the foot.

Mm. interossei dorsales (see also Fig. 455), four. **F o r m**: oblong-triangular, bipinnate, small. **P o s i t i o n**: in the interspaces between the ossa metatarsalia, projecting somewhat plantarward; covered dorsalward by all the muscles of the back of the foot, plantarward by all the muscles of the sole of the foot. **O r i g i n**: from the adjacent sides of each pair of ossa metatarsalia. **I n s e r t i o n**: the fibres of each muscle run converging forward to a small tendon; this extends dorsalward from the lig. capitulum [oss. metatars.] transversum to the lateral surface of the base of the first phalanx and also may go to the triangular expansions of the tendons of the mm. extensores digitorum. The tendon of the first (most medially situated) muscle runs on the medial side of the 2nd toe, while the tendons of the second to the fourth muscle lie on the lateral surface of the 2nd—4th toe. **A c t i o n**: the first draws the first os metatarsale toward the second; the first draws the first phalanx of the 2nd toe medianward, the others draw the first phalanx of the 2nd—4th toe lateralward; all usually only flex the first phalanx of these toes, without extending the 2nd and 3rd phalanx as in the hand. **I n n e r v a t i o n**: n. plantaris lateralis (S₁, S₂).




455. Mm. interossei pedis dextri, viewed from the plantar surface.

M. flexor digiti quinti brevis (O. T. flexor brevis minimi digiti pedis) (see also Figs. 449—451). **Form:** rounded, narrow, small. **Position:** in the sole of the foot, on the plantar surface of the os metatarsale V, partly covered plantarward by the m. abductor digiti V, otherwise only by the aponeurosis plantaris. **Origin:** basis oss. metatarsalis V and lig. plantare longum. **Insertion:** the fibres run converging forward and, becoming tendinous, go to the base of the first phalanx of the little toe. **Action:** it draws the 1st phalanx of the little toe plantarward and lateralward. **Innervation:** n. plantaris lateralis (S_1 , S_2).

M. opponens digiti quinti (see Figs. 451 and 458), often absent. **Form:** flat, triangular, small. **Position:** directly upon the plantar surface of the os metatarsale V, covered plantarward by the m. abductor digiti V, bounded medianward by the m. flexor digiti V. **Origin:** lig. plantare longum, fused with the m. flexor digiti V brevis. **Insertion:** the fibres run forward and somewhat lateralward to the os metatarsale V. **Action:** it draws the little toe plantarward and medianward. **Innervation:** n. plantaris lateralis (S_1 , S_2).


Mm. interossei plantares, three. **Form:** oblong-spindle-shaped, small. **Position:** in the interspaces between the ossa metatarsalia II—V, plantarward from the mm. interossei dorsales; projecting somewhat toward the sole of the foot and covered there by all the muscles of the sole of the foot. **Origin:** from the medial surface of the ossa metatarsalia III—V. **Insertion:** the fibres of each muscle go, converging, forward to a narrow tendon, which extends dorsally from the lig. capitulorum [oss. metatarsalium] transversum to the medial surface of the basis of the first phalanx and also may go to the triangular expansions of the tendons of the mm. extensores digitorum. The three muscles run on the medial side of the 3rd—5th toe. **Action:** they draw the first phalanx of the 3rd—5th toe medianward and generally only flex their first phalanx, without extending the 2nd and 3rd phalanx as in the hand. **Innervation:** n. plantaris lateralis (S_1 , S_2).



456. Muscles of the right foot, viewed from the medial surface.

M. abductor hallucis (see also Figs. 449—452 and 455). **Form:** flat, oblong-triangular, bipinnate. **Position:** superficial on the medial margin of the sole of the foot; bounded lateralward by the mm. flexor digitorum brevis and flexor hallucis brevis. **Origin:** medial surface of the tuber calcanei, superficial layer of the lig. laciniatum and malleolus medialis; also from the tuberositas oss. navicularis and the plantar surface of the os cuneiforme I. **Insertion:** the fibres converge to a tendinous strip, lying in the muscle, which goes over into a flat, strong tendon; this is fused in front with the medial belly of the m. flexor hallucis brevis and goes to the medial sesamoid bone and to the base of the first phalanx of the great toe. **Action:** it draws the first phalanx of the great toe medianward and plantarward. **Innervation:** n. plantaris medialis (L₅, S₁).

Ligamentum laciniatum (O. T. internal annular ligament) (see also Figs. 319, 445, 446, 451, 457 and 458) is a strengthening band in the lower part of the fascia cruris, below and behind the malleolus medialis. It goes off from the posterior and inferior margins of the malleolus medialis, covers the tendon of the m. tibialis posterior and holds it fast on the subjacent bone; it then divides into two layers. The superficial layer is stretched out from there to the medial surface of the tuber calcanei and goes over above into the two united layers of the fascia cruris (see p. 365); proximally it frequently has a sharp concave margin, sometimes without sharp limit; distally it serves as an origin to the fibres of the m. abductor hallucis. (Continued on p. 379.)



457. Bursae and sheaths of tendons of the right foot, injected (sheaths of tendons stained red), viewed from the medial surface.

(The mm. abductor hallucis and flexor hallucis brevis have been removed.)

Ligamentum laciniatum (continued). The deep layer passes just behind the tendons of the mm. flexor digitorum longus and flexor hallucis longus to the medial surface of the calcaneus, is firmly attached to the margins of the corresponding bony grooves and holds the tendons firmly upon the bones; it reaches above as far as the malleolus and is continuous with the vascular sheath. Between the two layers is a canal for the aa., vv. and nn. plantares; a septum, rarely absent, separates the medial branches from the lateral.

Bursa subcutanea malleoli medialis (see also Fig. 459), frequently present, upon the malleolus medialis.


Vagina tendinis m. tibialis posterioris (see also Fig. 458) begins over the medial malleolus and ends at the os naviculare; it frequently communicates above with the following.

Vagina tendinis m. flexoris digitorum pedis longi (see also Fig. 458) begins below the preceding and ends in the sole of the foot at the point where its tendon crosses that of the m. flexor hallucis longus, below the os naviculare; above it communicates sometimes with the preceding sheath, below sometimes with the following.

Vagina tendinis m. flexoris hallucis longi (see also Fig. 458) begins below the preceding and ends at or in front of the point where the tendon of the m. flexor digitorum longus is crossed, below the os naviculare; there it sometimes communicates with the preceding.

458. Bursae and sheaths of tendons of the sole of the right foot, injected; sheaths of tendons stained red.

(The short muscles of the 1st and 2nd layer have been for the most part removed.)



Bursa subtendinea m. tibialis posterioris, frequently present, oblong-trough-shaped, between the lateral tendinous process of the m. tibialis posterior (see Fig. 319) on one side, the ossa naviculare and cuneiforme II on the other side.

Vagina tendinis m. peronei longi plantaris begins at the sulcus m. peronei [longi] of the os cuboideum and ends at about the medial margin of the lig. plantare longum; it communicates only rarely with the vagina tendinum mm. peroneorum communis.

Vaginae tendinum digitales pedis, five, one on the volar surface of each toe; they begin behind, at the great toe in the neighborhood of the base, on the other toes in the neighborhood of the heads of the ossa metatarsalia, and terminate near the terminal phalanx; they are surrounded like those in the fingers (see p. 340) by the ligg. vaginalia with the ligg. cruciata and annularia; they never communicate with the sheaths which lie more proximalward.

Bursae mm. lumbricalium pedis, four, lying on the anterior extremities of the mm. lumbricales, between them and the ligg. capitulorum


[oss. metatars.] transversa; the three medial bursae are almost constant, the fourth occurs in half the cases.


459. Bursae and sheaths of tendons of the back of the right foot, injected; sheaths of tendons stained red.

Vagina tendinis m. tibialis anterioris (see also Fig. 457) begins somewhat above the lig. transversum cruris and ends usually somewhat above the articulation talonavicularis.

Vagina tendinis m. extensoris hallucis longi (see also Fig. 457) begins as the most deeply situated of the anterior tendon sheaths, below the lig. transversum cruris and ends farthest forward in the region of the articulation tarsometatarsae of the great toe, distalward from the strengthening band situated there (see p. 374); the distal end sometimes has the appearance of being pinched off.

Vagina tendinum m. extensoris digitorum pedis longi (see also Fig. 460) forms a sheath for the tendons of the m. extensor digitorum longus and of the m. peroneus III; it begins somewhat above the preceding below the lig. transversum cruris and ends over the middle of the os cuneiforme III.






460. Bursae and sheaths of tendons of the right foot,

injected (sheaths of tendons stained red), viewed from the lateral surface.

Bursa subcutanea malleoli lateralis (see also Figs. 441 and 459), very frequently present, upon the malleolus lateralis.

Vagina tendinum mm. peroneorum communis (see also Fig. 458) surrounds the tendons of the mm. peronei and bifurcates, at its proximal end for a shorter, at its distal end for a longer, distance into two parts, one for each tendon. It begins above the malleolus lateralis and ends at the beginning of the tuberositas oss. cuboidei; it communicates there only rarely with the vagina tendinis m. peronei longi plantaris. Behind and below the malleolus it may communicate with the joint cavity of the articulatio talocrualis.




461. Position
of the heart
in the thorax,
viewed from in front.

(The level of the diaphragm on moderate expiration is shown by the black dotted line; the outlines of the heart are indicated in red.) — (In part after W. Braune.)


The **heart (cor)** is an almost wedge-shaped, hollow, muscular body, intercalated as a pressure-pump in the circulatory system. It consists of two *fore-chambers* (O. T. auricles), the *atrium dextrum* and *atrium sinistrum*, and of two ventricles, the *ventriculus dexter* and *ventriculus sinister*, of which, after birth, the right auricle communicates only with the right ventricle, and the left auricle only with the left ventricle. The blood flows from the auricle into the ventricle through its *ostium venosum* and leaves the ventricle through its *ostium arteriosum*. The heart is situated asymmetrically in the middle of the thorax, so that the smaller part is on the right, the greater part on the left side (see also Fig. 683). The base of the heart (*basis cordis*) formed by the auricles is directed somewhat backwards and to the right, the apex of the heart, *apex cordis* (formed by the left ventricle alone), looks forward and to the left and comes into direct contact with the chest-wall in the 5th intercostal space, somewhat medial from the costocartilaginous junction. The *ostium arteriosum dextrum* (O. T. pulmonary orifice) lies usually at the sternal end of the left 3rd intercostal space, or behind the left 3rd costal cartilage; the *ostium arteriosum sinistrum* (O. T. aortic orifice) lies somewhat more medianward and downward, close below the middle of the left half of the sternum at the level of the 3rd intercostal space. The middle point of the *ostium venosum dextrum* (O. T. right auriculoventricular opening) lies behind the right half of the sternum at the level of the sternal end of the right 4th intercostal space; the *ostium venosum sinistrum* (O. T. left auriculoventricular opening) lies behind the sternal end of the 3rd left intercostal space.

The Heart.



462. The heart, moderately distended, viewed from above and in front.
(*Facies sternocostalis.*)


The heart presents for examination an inferior, almost plane surface, lying upon the diaphragm, the *facies diaphragmatica* (O. T. posterior surface), and a markedly curved surface looking upward and forward, the *facies sternocostalis* (O. T. anterior surface); the latter lies partly just behind the sternum and behind the anterior ends of the 3rd to the 6th pair of ribs, in part it is overlapped by the margins of the lungs. The two corresponding chamber-surfaces pass on the right rather suddenly into one another, thus forming a sort of margin, *margo acutus*, whilst on the left the transition takes place gradually, and is termed *margo obtusus*; these two „margins“ converge at the apex of the heart.



463. The heart, moderately distended, viewed from below.


(*Facies diaphragmatica.*)

The **heart**: the fore-chambers are separated from the ventricles by a groove, the *sulcus coronarius* (O. T. auriculoventricular groove), which is usually filled up with bloodvessels and fat; it runs nearly perpendicular to the long axis drawn from the apex to the middle of the base and is on the sternocostal surface in part hidden by the beginning of the a. pulmonalis and aorta. Corresponding to the septum between the two ventricles, on the inferior surface runs fairly straight the *sulcus longitudinalis posterior* (O. T. posterior interventricular groove); on the upper surface runs twisted so as to be somewhat S-shaped from the base toward the apex the *sulcus longitudinalis anterior* (O. T. anterior interventricular groove). The latter sulcus begins, covered by the left auricular appendix, on the left side of the a. pulmonalis and meets the sulcus longitudinalis posterior at the right of the apex of the heart in a shallow groove, the *incisura [apicis] cordis*.




464. Superficial muscle layer of a maximally contracted heart, viewed from above and in front.

(The fore-chambers have been dissected off. The outlines of the dilated ventricles of the heart are indicated in red, the position of the ostium arteriosum sinistrum being assumed to be fixed.)



465. Superficial muscle layer of a maximally contracted heart, viewed from below.

(The fore-chambers have been dissected off. Two windows have been cut into the superficial muscle layer in order to show the deeper bands of fibres. The outlines of the dilated ventricles of the heart are indicated in red, the position of the ostium arteriosum sinistrum being assumed to be fixed.)



466. The musculature of the ventricles of the heart, dissected free from one another; drawn apart; viewed from below. Somewhat schematically represented.


(On the right and left the superficial muscle is shown.)

The **heart** is covered on its outer surface by the visceral layer of the pericardium, the *epicardium*, on its inner surface by the *endocardium*; the main mass of the wall is formed of muscle, the *myocardium* (see Fig. 472).

Musculature of the heart. Owing to the network-like arrangement, the elements of the heart muscle bundles cannot be as sharply delimited as in the skeletal musculature. Besides, the bundles undergo manifold anastomosis with one another. The anastomoses appear as flat layers in the superficial portions so that there one can recognize a distinct lamellation, while in the depth, where the anastomoses are more evenly developed in all directions, the layers are far less marked. In spite of this there is a dominant direction of the course of the fibres; it is this only that will be described in what follows. The reader should notice that the direction of the fibres in the directly successive layers changes only gradually.

The *musculature* *) of the *fore-chambers* (see Figs. 464 and 465) is very thin and partly little regular. On the posterior and especially on the anterior surface fibres go transversely from the wall of one atrium to that of the other; otherwise the fibres run arch-like around the mouths of the veins and continue as rings on the termination of the vv. pulmonales as far as to the reflection of the pericardium; the point of origin of the left auricular appendix is also surrounded by nearly ring-like fibres. On the inner surface of the right fore-chamber strong muscle fibres form the substratum of the crista terminalis and of the limbus fossae ovalis; those of the latter also send a strong bundle into the tuberculum intervenosum; the muscle bands arise in the region of the trigonum fibrosum dextrum. From the muscle band of the crista terminalis the mm. pectinati take their origin (see p. 394). In addition, some fibres go over directly into the radial fibre-layer of the valvulae tricuspidalis and bicuspidalis (see page 393). The lumen of the auricular appendices is traversed by numerous irregularly arranged muscular columns (*trabeculae carneae*).

*) In the description of the musculature (pp. 388—389^b) the heart is thought as being placed with the base upward, the facies diaphragmatica backward and the apex downward.




467. Course of the principal layers of muscle of the left ventricle of the heart, viewed from below. Partly schematic.

(Parts of the wall of the heart have been removed above, on the left side below.)

The *musculature of the ventricles* (see also Figs. 464—466 and 468—470) is in its superficial layer common to both ventricles, but otherwise partly separated. The most superficial muscle layer (*vortex fibres*, see Figs. 464, 465 and 470) arises in general from both annuli fibrosi and runs in slightly (left-)interwoven spiral turns (the fibres of the left ventricle usually more steeply than the fibres of the right ventricle) in front from the right and above downward and to the left, behind from the left and above downward and to the right. The layer is in general easily separable from its substratum. At the apex of the left ventricle the fibres converge and come to lie in layers (*vortex cordis*, see also Fig. 464), bend sharply around into the interior of the left ventricle and become lost on the septal wall, in the papillary muscles and in the trabeculae carneae of the left chamber.


The *musculature proper of the right ventricle* (see also Figs. 464—466 and 468—470) is thin and shows in general two layers. The portion containing the ostium venosum shows, especially well marked in the dilated heart, an external, thin, continuous layer and an inner layer arranged more or less like a network. The fibres of the *external* layer arise from the posterior portions of both annuli fibrosi and between them, and run in the wall of the right ventricle forward in a direction similar to that taken by the vortex fibres, though less perpendicularly. Then a part of the fibres bends around sharply into the interior and mixes with the fibres of the inner layer (see below). Another part enters in front into the septal wall; these fibres are joined by fibres which surround the conus arteriosus in a ring-shaped manner. The *inner* layer is formed of short bands of fibres which arise partly at the upper margin of the septal wall, go toward the apex, turn at varying levels toward the external wall and run upward in this again, in part to end as the muscoli papillares, in part to reach the annulus fibrosus as the *trabeculae carneae*; between these columns are others which are stretched out transversely. The conus arteriosus possesses on the inside of the mentioned layer of circular muscle a longitudinal layer at right angles to it; the latter forms longitudinal ridges in the contracted heart.



467^a. The atrioventricular bundle and its distribution in the right ventricle.

(The wall of the atrium and of the ventricle has been partly removed. The atrioventricular bundle has been exposed by removal of the endocardium and of the muscle layer covering it.)


The *musculature of the left ventricle* is distinctly thicker than that of the right. Its fibres arise from the annulus fibrosus sinister and run downward (in layers and interwoven in slightly spiral turns) on the anterior, left and posterior surface of the left ventricle in a direction similar to that taken by the vortex fibres, though always less perpendicularly. Behind, the fibres enter into the septal wall and run forward in it. Then a part of them turns into the depth and bend around into the anterior papillary muscle; another part runs farther in the anterior and left ventricular wall and ascends partly in the posterior papillary muscle, partly alongside of it to the region of the trigonum fibrosum dextrum. Next to the external oblique fibres in the middle are those which are arranged approximately horizontally, that is, at right angles to the axis of the heart, and, next to these in the inner layer, are the fibres that run again obliquely, but in an opposite direction to the external (in right handed screw-like turns, that is in front from the right and below to the left and upward) (see Fig. 467). Thus, in general, the fibres leave the heart's apex free; this apex is thin-walled, and is formed only by the vortex fibres. The fibres, that run toward the base at the internal surface of the ventricle join the continuations of the vortex fibres (see page 389) and are arranged in gentle screw-like turns that run, however, nearly parallel to the long axis of the heart. Along with the papillary muscles they form as the *trabeculae carneae*, numerous long thick longitudinal ridges, which are united by thinner cross ridges to form a network. The septal wall below the ostium arteriosum is the only part that is completely smooth; upon it the left branch of the atrioventricular bundle (see p. 389^b) runs downward.



467^b. The atrioventricular bundle and its distribution in the left ventricle.


(The left atrium and the left ventricle have been cut open and spread apart. The atrioventricular bundle has been exposed by removal of the endocardium.)

The musculature of the atria is connected with that of the ventricles only by means of a narrow bundle situated in the septal wall. This bundle is known as the *atrioventricular bundle* (see also Fig. 467^a); it is a part of the *conduction system of the heart* and is composed of muscle fibres of a somewhat lighter color and of a slight yellowish tint. It begins a little in front of the opening of the sinus coronarius, close behind the septum membranaceum ventriculorum (see page 392), just above the upper, sharp margin of the septum musculare ventriculorum usually in a small thickening (*Tawara's node*), into which radiate certain bands of uncertain origin derived from the atrial musculature (especially from the region below the opening of the sinus coronarius). The *trunk*, 1–2 mm. in thickness, of the bundle runs out of this node for a distance of about 1 cm. on the lower margin of the septum musculare (displaced somewhat to the right); it runs forward and undergoes subdivision into a right and a left branch. The *right branch* runs, as a rounded bundle, in a curve that is slightly convex forward, on the right side of the ventricular septum. It extends downward, more or less hidden in the muscle mass, just behind the origin of the anterior medial papillary muscle (see p. 396) and enters the large trabecula carnea, which forms a bridge between the septal wall and the large papillary muscle. From this point on it cannot be followed further with the naked eye. The *left branch* extends downward and a little to the left, and, at the lower margin of the septum membranaceum beneath the point of attachment of the valvula semilunaris posterior aortae, reaches the left surface of the ventricular septum, passes obliquely downward and forward upon it as a broad but very thin band of fine fibres, and then divides, while still in the upper third of the ventricular septum, into two branches. Of these two branches, the anterior splits up and passes to the trabeculae, which extend to the base of the anterior papillary muscle; the posterior, bending a little downward, similarly reaches the base of the posterior papillary muscle; a few fibres are split off and become lost in the rest of the trabecular network. The left branch with all its visible branches lies close beneath the endocardium and is partly visible through it.



468 and 469. Cross-sections of two hearts between the upper and middle third of the ventricles, perpendicular to the long axis, seen from the apical side.

The upper heart has been fixed in systole, the lower in diastole. (After L. v. Krehl.)



470. Base of the contracted ventricles, after removal of the fore-chambers. The valves are closed.


(The outlines of the distended ventricles are indicated in red, the position of the ostium arteriosum sinistrum being assumed to be fixed.)

The **annuli fibrosi** are dense, narrow bands of connective tissue, which surround the ostia venosa at the base of the ventricles. They serve for the attachment of the muscular fibres of the atria and of the ventricles, and from them arise in largest part the valvulae tricuspidalis and bicuspidalis. The right surrounds the ostium venosum dextrum in the form of an oval closed ring. The left is horse-shoe-shaped surrounding the right, inferior and left sides of the ostium venosum sinistrum; it begins right and left at the root of the aorta in a nodular thickening on either side, the *trigonum fibrosum*; between these the anterior cusp of the valvula bicuspidalis arises directly from the membranous wall of the root of the aorta.

The **differences between the contracted and dilated ventricles** are as follows (see also Figs. 462—465, 468 and 469):

In the *dilated ventricles* the diameter of the base is nearly the same as that of the long axis; the form approaches that of a hemisphere. The conus arteriosus projects somewhat, the lower surface in its left section projects distinctly convex and shows in the middle and in its right section a furrow-shaped shallow depression, particularly marked toward the apex; the ostia venosa and arteriosa are large, the cavities of the ventricles are wide open; the trabeculae carneae extend partly through the lumen of the ventricles, and the muscoli papillares stand out partly free in the ventricular cavities.


On maximal *contraction of the ventricles*, the diameters of the base are markedly shortened, while that of the long axis diminishes only slightly; the form is wedge-shaped. The conus arteriosus shortens and sinks in; the facies diaphragmatica bulges throughout slightly convexly, and the furrow-shaped depression in the dilated heart is still perceptible only at the level of the apex slightly bent downward; the sulcus longitudinalis anterior has a more marked spiral twist, the ostia venosa are markedly diminished in size, the ostia arteriosa less markedly, the cavities of the ventricles are represented only by a slit (with the exception of a space situated above the apices of the papillary muscles); the trabeculae carneae and the muscoli papillares lie close to the walls and form ridge-like projections.



471. Transverse section through a distended heart approximately parallel to the facies diaphragmatica.

(Inferior half of the section, viewed from above.)

The **septum ventriculorum** (*interventricular septum*) (see also Figs. 466, 468 and 469) separates the two ventricles from one another completely. It is muscular in by far its largest part (*septum musculare ventriculorum*), is formed by a corresponding portion of the musculature of each of the two ventricles and can be divided into its two components only by artificial means. On contraction it thickens and the distance between the two sulci longitudinales diminishes. The *septum membranaceum ventriculorum* (O. T. undefended space) is a small area of pure connective tissue situated in the region of the atria. Its left surface is situated just between and in front of the opposed margins of the valvulae semilunares dextra et posterior aortae (see also Figs. 467^a, 467^b and 477); its right surface is divided into two parts by the cuspis medialis valvulae tricuspidalis arising from it; the posterior part looks into the right atrium, the anterior into the right ventricle. About the relations of the septum membranaceum to the atrioventricular bundle see p. 389^b.




472. Transverse section through the valvula bicuspidalis [mitralis]. The valve is set.

(Inferior half, viewed from above.)

The **musculi papillares** (*papillary muscles*) (see also Figs. 468, 469, 471, 475 and 476) are more or less isolated, bluntly wedge-shaped elevations of the internal muscular layer of the ventricles. They vary much in size and in different hearts present considerable differences in appearance. From the tip of each muscle there go off one or several tendinous threads (*chordae tendineae*) of variable strength, which become attached partly to the free margins (*chordae tend. of the I order*), partly to the ventricular surface (*chordae tend. of the II order*) of the cusps of the valves guarding the ostia venosa; the insertion is in the form of a triangular flat expansion. Each papillary muscle gives off chordae usually to two adjacent cusps. In addition other tendinous threads (*chordae tend. of the III order*) arise directly from the musculature of the ventricles near to the peripheral margin of the valves and become attached in the same way also to the ventricular surface of the cusps. The chordae tendineae are of unequal length and are so arranged that the valve is set when they are all put upon the stretch, and vice versa.


The *valvula tricuspidalis* and *valvula bicuspidalis [mitralis]* (see also Figs. 470 and 471) consists each of a tube-like membrane, which is attached to the annulus fibrosus (also partly to the wall of the root of the aorta on the left side) and is divided at its free margin by deep incisures into large *sails* or *cusps* (*cuspides*) and by less deep incisures into variably developed, smaller, intermediary cusps. They are covered on both sides by endocardium, contain in their peripheral third external circular and internal radial muscle fibres which are connected with the musculature of the fore-chambers (see p. 388) and possess in this part also vessels. At the end of the diastole of the ventricles, the central portions of the atrial surfaces of the cusps become applied to one another (the valves become set); the closure is water-tight and during the systole becomes gradually ever more complete, since the ostia venosa are markedly diminished in size, and thus ever larger sections of the cusps of the valves are applied to one another.



473. Right fore-chamber (*atrium dextrum*) of a markedly distended adult heart, viewed from the right.

(A part of the wall of the atrium has been removed.)

The **atrium dextrum** (*right fore-chamber*, O. T. right auricle) (see also Figs. 462–465 and 471) is of the shape of an irregular short cylinder. A groove, variably developed, running perpendicularly over the posterior external surface, the *sulcus terminalis atrii dextri*, and a ridge on the corresponding spot of the inner wall, the *crista terminalis*, mark off a posterior, medial portion, the *sinus venarum [cavarum]*, which is derived from the right horn of the sinus reunions of the embryonic heart (see also p. 402). It contains in its lower and upper walls the mouths of the vv. cavae superior and inferior and in general has smooth walls. Between the opening of the v. cava inferior and the sulcus coronarius to the left of the sulcus terminalis there is often visible on its outer surface an irregular lumpy projection (*appendix auricularis posterior*) (see Fig. 463). The posterior wall of the sinus is bounded partly by the atrium sinistrum, being slightly shoved forward by the same and presents there a transverse ridge, the *tuberculum intervenosum [Loweri]* (O. T. tubercle of Lower). The left wall is formed by the *septum atriorum* (O. T. interauricular septum), which consists partly of muscle, partly (*pars membranacea septi atriorum*) of pure connective tissue. The latter spot lies in the lower part of the wall, is somewhat deepened (*fossa ovalis*) and is surrounded in front, above and behind by a strong ridge of muscle (*limbus fossae ovalis [Vieussenii]*). Starting at the anterior limb of the limbus, and arising from the inferior wall, is a thin, often partially perforated, fold of connective tissue which surrounds the mouth of the v. cava inferior from in front and sometimes also from the right, *valvula venae cavae [inferioris, Eustachii]* (O. T. Eustachian valve). Just in front of it, in the angle between the inferior, left and anterior walls, lies the mouth of the sinus coronarius; it is only imperfectly closed from below by the sickle-shaped *valvula sinus coronarii [Thebesii]* (O. T. coronary valve or valve of Thebesius), which is often perforated. The anterior wall of the atrium contains the ostium venosum of the right ventricle. The right wall presents numerous muscular ridges projecting on the inner surface (*musculi pectinati*) (see also Fig. 471); these arise from the crista terminalis and extend approximately in the direction of the long axis of the heart as far as the sulcus coronarius; between them at the dilated heart the wall is translucent.




474. Right fore-chamber (*atrium dextrum*) of a fetal heart (8th month), viewed from the right.

(A part of the wall of the atrium has been removed. Magnification: 2 : 1.)

Atrium dextrum (continued). From the upper wall projects the bluntly wedge-shaped *auricula dextra* (right auricle of the heart) (O. T. right auricular appendix). It curves around the first portion of the aorta ascendens to the left, is slightly notched above and below but otherwise smooth; it presents inside numerous irregular crossing muscular columns, *trabeculae carneae*.

During fetal life the two atria communicate with one another through an opening, the *foramen ovale*, situated at the spot which later is the fossa ovalis. At its margin in front and above it is surrounded by a horse-shoe-shaped, powerful band of muscle, which later becomes the limbus fossae ovalis, and behind and below it is bounded by the *valvula foraminis ovalis* which rises as a thin connective tissue plate from behind and below and projects, groove-like, into the left atrium. Just after birth, as soon as the pressure in the left atrium becomes higher than in the right, this valve is pressed upon the limbus and fuses with it almost completely, occasionally, however, leaving a small capillary slit open. The *valvula foraminis ovalis* becomes accordingly later the floor of the fossa ovalis; the anterior concave margin of the valve is often retained and is visible inside the left atrium on the septal surface (see p. 397).

The **ventriculus dexter** (right ventricle) (see Figs. 462—466, 468—471 and 475) has the shape of a wedge; its left concave surface is attached to the right side of the left ventricle; its apex does not reach the apex of the heart. The base of the wedge is hidden by the attachment of the right atrium; the inferior, the right and the superior anterior surfaces lie free; the latter goes very gradually over into the *arteria pulmonalis*, and in the distended heart projects markedly convexly forward (*conus arteriosus*). On the inside, marking off the *conus arteriosus* from the rest (the venous portion) of the ventricle, is a muscular projection *crista supraventricularis* which extends, arch-like, from the chamber-base, to the anterior medial papillary muscle (see p. 396) or passes a little before it; a slight ridge, which extends from the region immediately behind this papillary muscle on the septum in the direction to the apex, to the roots of the great papillary muscle, contains the continuation of the right limb of the atrioventricular bundle (see p. 389^b). The venous portion presents at the septum a fairly smooth wall, but elsewhere very many *trabeculae carneae* (see p. 389). The *ostium venosum* can be closed by the *valvula tricuspidalis* (tricuspid valve); this presents (see Fig. 470) a large cusp adjoining the interventricular septum (*cusps medialis*), further a small *cusps anterior* corresponding about to the *conus arteriosus*, a large *cusps posterior* turned toward the right free wall, and, usually, between the *cusps medialis* and *cusps posterior* an additional small intermediary cusp.




475. Right ventricle (*ventriculus dexter*)

of a markedly distended heart, viewed from the right and above.

(The largest part of the right and anterior wall has been removed; the atria have been removed.)


Ventriculus dexter (continued). The chordae tendineae of the anterior and posterior cusps come partly from a large papillary muscle which stands up free from among the trabeculae carneae below the junction of the conus arteriosus with the venous portion; besides this there is usually one smaller (posterior medial) papillary muscle between the septum and the inferior (diaphragmatic) wall, as well as one very small (anterior medial) between the anterior and medial cusp on the septum just below the crista supraventricularis (see p. 395), in whose place several papillary muscles always can be found; the medial cusp receives many chordae directly from the septum itself. The papillary muscles of the right ventricle present marked individual variations. The conus arteriosus is more tubular, has a fairly smooth wall and goes over through the *ostium arteriosum* into the a. pulmonalis; at the entrance to the latter are situated the *valvulae semilunares arteriae pulmonalis anterior, dextra, sinistra* (semilunar valves of the pulmonary artery) (see p. 398).



476. Left ventricle (*ventriculus sinister*)

of a distended heart, viewed from in front and somewhat from the left. (The anterior wall of the left ventricle has been almost entirely removed, the left wall partially and with them also the anterior papillary muscle; the atria have been cut away. A arrow has been introduced into the ostium arteriosum.)

The **atrium sinistrum** (*left fore-chamber*) (O. T. left auricle) (see Figs. 462—465 and 478), is wedge-shaped, broad behind, narrower in front. At the junction of the posterior with the lateral surfaces, right and left, lie the openings of the *venae pulmonales*. The right wall, formed partly by the *septum atriorum*, shows frequently the concave margin of the *valvula foraminis ovalis* (see p. 395). The anterior wall contains the *ostium venosum ventriculi sinistri*. From the junction of the right with the superior wall goes off the zigzag-curved *auricula sinistra* (*left auricle of the heart*) (O. T. left auricular appendix) which curves around the *a. pulmonalis* from the left and reaches as far as its anterior surface. This left auricular appendix contains inside it numerous interlacing *trabeculae carneae*; otherwise the wall of the atrium is smooth inside.



477. Valvulae semilunares aortae, opened up.

(The left ventricle and the aorta have been opened by a cut passing between the right and the left valve; the walls have been spread apart.)


The **ventriculus sinister** (*left ventricle*) (see Figs. 462—472 and 476) has the shape of an egg, the large end of which is directed toward the right and backward and contains the ostia venosum et arteriosum; the other end forms the apex of the heart. As regards the inner surface see also p. 389^a. The *ostium venosum* is guarded by the *valvula bicuspidalis* [*mitralis*] (*bicuspid* or *mitral valve*). This consists of two large cusps and usually two small intermediary cusps; of the larger cusps, the *cuspis anterior*, situated to the left and behind, arises from the annulus fibrosus sinister; the *cuspis posterior*, situated to the right and in front, comes however only partially from the annulus fibrosus sinister, having its origin chiefly from that connective tissue portion of the root of the aorta which extends between the trigona fibrosa, so that the ventricular surface of this cusp goes over directly into this part of the wall of the aorta (see also Figs. 471 and 476). The chordae tendineae come from two papillary muscles, an anterior and a posterior one, each of which can in turn be divided into several wedges; both muscles are developed from the left wall and correspond in their position approximately to the *sinistra, posterior*; the posterior portion of the left valve and the left portion of the posterior valve arise from the membranous part, the other parts of these valves and the right valve from the muscular part of the root of the aorta.

Each *valvula semilunaris* presents just below the middle of its concave, free margin a small nodular thickening, *nodulus valvulae semilunaris* [Arantii], and on both sides of this, near the margin, a sickle-shaped, thinner portion, *lunula valvulae semilunaris*, by means of which it lies against the neighboring valve.

The small or **pulmonary circulation** consists of the a. pulmonalis, the vv. pulmonales and their branches.


The **a. pulmonalis** (see Figs. 462—464, 478, 680 and 681) extends on the left side of the aorta ascendens obliquely to the left and backward and divides below the arcus aortae into a *ramus dexter* and a *ramus sinister*. The former goes below the arcus aortae behind the aorta ascendens and v. cava superior, in front of the bronchus dexter to the hilus of the right lung; the r. sinister runs in front of the bronchus sinister and the aorta thoracalis to the left lung. In the prolongation of the main stem a band of elastic-connective tissue, the *ligamentum arteriosum*, the remains of the obliterated ductus arteriosus [Botalli], extends to the concave, anterior wall of the end of the aorta (see p. 491).

The **vv. pulmonales** (see Figs. 478, 671, 672, 680 and 681) come from the hilus pulmonalis on each side and open into the atrium sinistrum. They run on both sides in front of the main branches of the a. pulmonalis and in front of the bronchi; in addition the right veins go behind the v. cava superior and the atrium dextrum, the left in front of the aorta thoracalis.



**478. Position of the great vessels,
viewed from in front.**


(Pieces have been cut out of the v. cava superior, aorta and a. pulmonalis; the heart has been turned downward, so that the apex looks downward.)



479. The arteries of the heart, viewed from above and in front.

(The ostium arteriosum dextrum and the a. pulmonalis have been removed.)

The *a. coronaria [cordis] dextra* (right coronary artery) (see also Fig. 480) arises in the right sinus aortae [Valsalvae] (see Figs. 477 and 484), runs at first between the right auricular appendix and the conus arteriosus, then in the sulcus coronarius first to the right and then upon the facies diaphragmatica to the left; finally it bends around into the sulcus longitudinalis posterior and, as the *ramus descendens posterior*, arrives close to the apex of the heart. It gives off large branches to the right ventricle, small branches to the right atrium and (from the r. descendens posterior) to the left ventricle and to the septum ventriculorum.




480. The arteries of the heart, viewed from below.

The *a. coronaria [cordis] sinistra* (*left coronary artery*) (see also Fig. 479) arises from the left sinus aorta [Valsalvae] (see Figs. 477 and 484), is usually larger than the right and divides just after its origin into two branches. The *ramus descendens anterior* extends at first behind the a. pulmonalis, then to the left from it, covered by the left auricular appendix, to the sulcus longitudinalis anterior, in which it runs as far as the incisura [apicis] cordis, usually going over upon the facies diaphragmatica; it gives off larger branches to the septum ventriculorum and the left ventricle and smaller branches to the right ventricle. The *ramus circumflexus*, hidden below the left auricular appendix, goes in the sulcus coronarius first to the left, then upon the facies diaphragmatica to the right and does not reach the sulcus longitudinalis posterior; it gives off larger branches to the left ventricle and smaller ones to the left atrium.

Both aa. coronariae vary not rarely in as much that the ramus circumflexus of the left reaches on the facies diaphragmatica as far as to its right margin (and thereby gives off the r. descendens posterior), or that the right may run as far as the left margin; in the first case the right, in the last case the ramus circumflexus of the left is shortened accordingly.

Both aa. coronariae anastomose with one another by means of innumerable minute branches, usually inside the muscle, more rarely on the outer surface just beneath the epicardium.




481. The veins of the heart, viewed from below.

Most of the **veins of the heart** open into the *sinus coronarius*. This lies on the *facies diaphragmatica* in the *sulcus coronarius*, extends to the left as far as the opening of the *v. obliqua atrii sinistri* [Marshalli] and opens at the right into the right atrium (see Fig. 473); it arises from the cross-piece of the embryonic *sinus reuniens* (see also p. 394) and, as such, possesses cross-striped (cardiac) muscle fibres. The vein openings into the sinus are partly closed at their mouths by valves, which, elsewhere, are absent. All the veins of the heart anastomose with one another.

The *v. cordis parva* opens from the right into the sinus; it runs in the *sulcus coronarius* and comes partly from the right ventricle, partly from the right atrium.

The *v. cordis media* extends in the *sulcus longitudinalis posterior* from the apex of the heart to the base; it receives its branches from both ventricles and from the *septum ventriculorum* and opens from the right into the sinus.

The *v. obliqua atrii sinistri* [Marshalli] (O. T. oblique vein of Marshall) runs downward as a small branch on the left side of the left atrium in front of the pulmonary veins and opens from the left into the sinus. It connects above with the fine, fibrous *ligamentum venae cavae sinistralae* (see Fig. 463) which runs downward from the *v. anonyma sinistra* in front of the left pulmonary vessels and gives rise to a low fold of the epicardium; together with this ligament the *v. obliqua atrii sinistri* is a remnant of the embryonic *v. cava sinistra*.



482. The veins of the heart, viewed from above and in front.


The **veins of the heart** (continued) (see also Fig. 481).

The *v. posterior ventriculi sinistri*, on the left side of the heart, receives branches from the left ventricle and opens into the *v. cordis magna* or to the right into the sinus coronarius.

The *v. cordis magna* arises at the apex of the heart, runs at first in the sulcus longitudinalis anterior toward the base, then in the sulcus coronarius around the left surface to the sinus coronarius; it collects its blood partly from the right, but chiefly from the left ventricle, from the septum ventriculorum and from the left fore-chamber.

The *vv. cordis anteriores*, varying in number, upon the right ventricle, open at the sulcus coronarius directly into the right atrium.

The *vv. cordis minimae* open directly into the cavities of the right and left heart by means of the *foramina venarum minimarum* [Thebesii] (see Fig. 471).




483. The heart, viewed from the right and in front, showing the place where the pericardium is reflected.

(The large arteries have been drawn away somewhat from the atria.)

The **pericardium** (see also Figs. 678—683) is a serous sack, closed on all sides, which presents a visceral layer, *epicardium*, firmly attached to the heart muscle, and a parietal layer. Between the two layers is a small slit-like space in which are contained several grams of a clear serous fluid, the *liquor pericardii*.

The *epicardium* (see also Figs. 462 and 463) covers completely the free surface of the ventricles of the heart and contains deposits of fat of variable size along the vessels. The atria are also covered on their free surface in large part by the epicardium, only the posterior surface of the left and a narrow, vertical strip of the right atrium remaining free from it; the v. cava superior is covered by it for a distance of one little finger's breadth; the v. cava inferior and vv. pulmonales are covered by it only at the point where they open into the heart. The a. pulmonalis and aorta ascendens are connected with one another by connective tissue, are covered in common by the epicardium and are separated from the anterior wall of the atria by a slit, the *sinus transversus pericardii*, which is closed above and below but open to the right and to the left. The epicardium is reflected into the parietal layer of the pericardium along the junction between the aorta ascendens and the arcus aortae, as well as over the bifurcation of the a. pulmonalis.




484. Aortic arch with its branches, viewed from the left and in front.

(After a plaster mould.)

Pericardium (continued) (see Figs. 483, 678—683). The *parietal layer* is fused, on its lower surface, firmly with the diaphragm, on the posterior surface loosely with the oesophagus and the aorta thoracalis; to the right and left it is intimately united with the pleura pericardiaca. In front it is partly covered by the thymus and loosely connected with it; it lies in part just behind the corpus sterni and is fastened to it by several powerful, tendinous bands of fibres, the *ligamenta sternopericardiaca* (not illustrated), as well as by loose connective tissue.

The **aorta** runs, from the ostium arteriosum sinistrum, at first, upward (*aorta ascendens*); it then bends around backward (*arcus aortae*), and goes finally downward (*aorta descendens*), above, in front of the thoracic spine (*aorta thoracalis*), below, in front of the lumbar spine (*aorta abdominalis*). It ends in front of the lower margin of the 4th lumbar vertebra, a little below the navel.

The beginning of the **aorta ascendens** (see also Figs. 461, 462, 478 and 479) lies behind that of the a. pulmonalis and presents a flask-like swelling (*bulbus aortae*) with three smaller bulgings (*sinus aortae [Valsalvae]*), of which each corresponds to one valvula semilunaris aortae. Thence the aorta passes upward and turns only a little to the right and forward; it lies behind the right auricular appendix of the heart and ascends in the median plane behind the corpus sterni, on the right side of the a. pulmonalis and on the left side of the v. cava superior; behind, it adjoins the left atrium and the ramus dexter a. pulmonalis. The only larger branches given off by it are the two *aa. coronariae [cordis]* (see pp. 400 and 401).



485. Superficial arteries of the neck, viewed from the right and somewhat from in front.

(The head is bent somewhat backward; the parotid gland has been completely removed; the mm. pectoralis major, sternocleidomastoideus and the muscles of the face have been partially removed.)

The **arcus aortae** (*aortic arch*) (see Figs. 478, 484, 607, 681 and 682) arches upward, being convex, from before backward, and at the same time somewhat to the left, crossing the point of division of the a. pulmonalis and the bronchus sinister; on the right it lies upon the trachea just above its bifurcation; its highest convexity reaches to the level of the breast-bone junction of the first rib; below it ends on the left side of the body of the 4th thoracic vertebra.

From the upper wall of the aortic arch go off: *a. anonyma*, *a. carotis communis sinistra* and *a. subclavia sinistra*.

At the junction of the arcus aortae with the aorta thoracalis there is frequently a constriction, the *isthmus aortae*, and, just beyond this, an oblong expansion, the *aortic spindle* (see Fig. 484). Both formations are especially well marked when the vessel is much curved.

A. anonyma (innominate artery) (see Figs. 478, 500 and 607) runs obliquely in front of the trachea from the left and below to the right and upward, covered in front by the venae anonymae; it divides behind the articulatio sternoclavicularis dextra and somewhat above it into the *a. carotis communis dextra* and the *a. subclavia dextra*.

A. carotis communis (common carotid artery) (see also Figs. 358, 486, 492, 500 537 and 540) runs (left behind the articulatio sternoclavicularis), at first on the right somewhat farther forward, on the lateral side of the trachea and oesophagus, then behind the lobe of the thyroid gland, in front of the mm. scaleni and longus colli, upward and somewhat backward and divides near the cornu superius of the cartilago thyreoidea into the *a. carotis externa* and the *a. carotis interna*. It is accompanied in front and lateralward by the v. jugularis interna, in front by the ramus descendens n. hypoglossi, behind by the n. vagus, and is partially covered by the mm. sternocleidomastoideus, sternohyoideus, sternothyreoideus and omohyoideus.

A. carotis externa (external carotid artery) (see also Figs. 489, 490 and 492) passes almost straight upward, partly covered by the venter posterior of the m. digastricus and by the m. stylohyoideus, runs along the posterior margin of the ramus mandibulae on its medial surface (surrounded generally on all sides by the glandula parotis) and divides at the collum mandibulae into its two terminal branches: *a. temporalis superficialis* and *a. maxillaris interna*. It is partly covered by the v. facialis communis and by the n. hypoglossus. Branches:

1. **A. thyreoidea superior** (superior thyroid artery) (see also Figs. 486, 492 and 500) arises from the anterior wall of the point of division, sometimes even somewhat more deeply; runs in a curve forward and downward to the upper end of the lateral lobe of the thyroid gland, covered partly by the venter superior of the m. omohyoideus. Branches:


- a) *Ramus hyoideus* upward to the hyoid bone.
- b) *Ramus sternocleidomastoideus* lateralward and downward to the m. sternocleidomast.
- c) *A. laryngea superior* perforates the membrana hyothyreoidea, being covered by the m. thyrohyoideus, and anastomoses on the lateral surface of the laryngeal muscles with the *a. laryngea inferior* (from the *a. thyreoidea inferior*) and with a branch of d).
- d) *Ramus cricothyreoideus* (see Figs. 486 and 500), in front of the m. cricothyreoideus and the lig. cricothyreoideum medianward; under the cartilago thyreoidea it gives a branch to the laryngeal muscles (see c) and anastomoses with that of the other side.
- e) and f) *Ramus anterior* and *ramus posterior*, to the anterior and posterior surface of the thyroid gland; both break up into numerous *rami glandulares* (to the gland).

2. **A. lingualis** (lingual artery) (see also Figs. 486 and 492) goes off, at the level of the greater cornu of the hyoid bone, from the anterior wall, extends in a curve upward upon the m. constrictor pharyngis medius, then, parallel to and above the hyoid bone, medially to the m. hyoglossus forward upon the m. longitudinalis inferior and the m. genioglossus. Branches:

- a) *Ramus hyoideus* downward to the lateral surface of the hyoid bone.
- b) *Rami dorsales linguae*, double or single, on the medial surface of the m. hyoglossus, upward to the region of the root of the tongue and the tonsil.
- c) *A. sublingualis* arises at the anterior border of the m. hyoglossus, runs between the m. mylohyoideus and the m. genioglossus below the glandula sublingualis forward, supplies the floor of the mouth and the gums of the anterior teeth; it anastomoses through the m. mylohyoideus with the *a. submentalis*.
- d) *A. profunda linguae* (O. T. ranine artery), very tortuous, runs upward and forward between the m. longitudinalis inferior and the m. genioglossus; it anastomoses above the frenulum linguae with that of the other side.

3. **A. maxillaris externa** (external maxillary artery) (O. T. facial artery) (see also Figs. 487 and 492) arises, close above the *a. lingualis* (sometimes in common with it), from the anterior wall, covered by the venter posterior of the m. digastricus and by the m. stylohyoideus; it runs forward in a S-shaped curve on the medial and upper surface of the glandula submaxillaris, between it and the lower jaw, then bends around the basis mandibulae to the anterior margin of the m. masseter and passes, very tortuous, upon the mm. buccinator and caninus to the side of the nose. In the face it is covered by the platysma, mm. risorius, zygomaticus and quadratus labii superioris, separated from the latter by the v. facialis anterior. Branches:

- a) *A. palatina ascendens* (O. T. ascending or anterior palatine artery) (see Figs. 486 and 492) arises near the origin (frequently from the *a. carotis externa* or *a. pharyngea ascendens*), runs upward upon the m. stylopharyngeus, then between this muscle and the m. styloglossus, then upon the m. constrictor pharyngis superior to the medial surface of the m. tensor veli palatini; it supplies the soft palate and the pharynx.
- b) *Ramus tonsillaris* (see Fig. 492), frequently a branch of a), runs upon the m. constrictor pharyngis superior to the tonsilla palatina and to the soft palate.



486.

The arteries of the larynx and of the tongue, viewed from the right.

(The lamina dextra of the cartilago thyroidea, the right half of the lower jaw and the m. hyoglossus have been partially removed.)


3. A. maxillaris externa (continued):

- c) *A. submental* (see Fig. 485) runs under the m. mylohyoideus, between it and the glandula submaxillaris, forward to the region of the chin; it anastomoses with the a. sublingualis.

- d) *Rami glandulares*, arising partly from c), to the glandula submaxillaris.
- e) *A. labialis inferior* (O. T. inferior coronary artery) (see Fig. 487) arises below the angle of the mouth, runs medianward in the lower lip in the m. orbicularis oris, close to the mucous membrane, and forms a broad anastomosis with that of the other side.
- f) *A. labialis superior* (O. T. superior coronary artery) (see Fig. 487) arises at the angle of the mouth, runs in the upper lip like e) in the lower; branches to the nasal septum.
- g) *A. angularis* (see Fig. 487), the terminal branch, ascends almost vertically along the lateral margin of the nose and anastomoses with the a. dorsalis nasi (from the a. ophthalmica).

4. *A. temporalis superficialis* (superficial temporal artery) (see Figs. 485, 487—489) ascends behind the proc. condyloideus mandibulae, in front of the tragus, over the root of the zygoma; it is covered at first by the glandula parotis, and then, very tortuous, lies upon the fascia temporalis and galea aponeurotica; it divides above the ear into its two terminal branches, *ramus frontalis* and *ramus parietalis*. Branches:

- a) *Rami parotidei*, in part also directly from the a. carotis externa, to the parotid gland.
- b) *A. transversa faciei* (O. T. transverse facial artery), runs forward to the cheek directly upon the m. masseter, just below the zygoma, covered behind by the parotid gland.
- c) *Rami auriculares anteriores*, 3—4, to the auricle and to the external auditory canal.
- d) *A. temporalis media* (medial temporal artery) (see Fig. 489) perforates the fascia temporalis just above the root of the zygoma, runs upward in the sulcus a. temporalis mediae of the temporal bone and is distributed in the m. temporalis.
- e) *A. zygomaticoorbitalis*, above the zygoma, to the upper margin of the orbit.
- f) *Ramus frontalis* (O. T. anterior terminal branch), is tortuous and extends first forward and upward, then backward upon the m. epicranii.
- g) *Ramus parietalis* (O. T. posterior terminal branch), tortuous, backward and upward.
- f) and g) form a rich anastomotic network with one another, with the aa. frontales and occipitales.



487. Superficial arteries of the face, viewed from the right.


(The platysma, m. risorius and m. quadratus labii superioris have been removed.)

5. **A. sternocleidomastoidea** (see Fig. 485) arises, at the same level as the a. maxillaris externa (often from 6.), from the outer wall and goes in a curve, in front of the n. hypoglossus and the v. jugularis interna, lateral- and downward to the m. sternocleidomastoideus.

6. **A. occipitalis** (see Fig. 489), somewhat above the previous artery, from the posterior wall, runs at first upward upon the v. jugularis interna, covered by the venter posterior of the m. digastricus; then it goes backward in the sulcus a. occipitalis between the m. rectus capitis lateralis and the venter posterior m. digastrici, then between the mm. splenius capitis and semispinalis capitis; finally it bends upward and arrives between the insertions of the mm. trapezius and sternocleidomastoideus at the linea nuchae superior upon the m. epicranii. Branches:

- a) *Rami musculares* to the m. sternocleidomast. and neighboring muscles; among these
- aa) *Ramus descendens*, downward between the mm. splenius capitis and semispinalis capitis.
- b) *Ramus mastoideus* through the foramen mastoideum to the dura mater (see also Fig. 490).
- c) *Ramus auricularis* obliquely upward and forward toward the ear.
- d) *Rami occipitales*, very tortuous, upward; those of the two sides anastomose with one another and with the aa. temporales superficiales. A small branch, the *ramus meningeus*, goes through the foramen parietale to the dura mater.


7. **A. auricularis posterior** (posterior auricular artery) (see Figs. 485 and 489), above the preceding from the posterior wall, runs, covered by the glandula parotis, on the upper margin of the venter post. m. digastrici, upward and in front of the proc. mastoideus behind the ear. Branches:



488. Arteries of the roof of the skull, viewed from above.

7. *A. auricularis posterior* (continued):

- a) *A. stylomastoidea* (see Figs. 492 and 493) goes through the foramen stylomastoideum into the canalis facialis, there gives off branches backward to the cellulae mastoideae (*rami mastoidei*), forward to the m. stapedius (*ramus stapedius*) and through the canaliculus chordae tympani into the middle ear (*a. tympanica posterior*); it anastomoses with the ramus petrosus superficialis (from the a. meningea media) (see p. 412).
- b) *Ramus auricularis* (O. T. anterior terminal branch) to the medial surface of the external ear, and partly, perforating it, to the lateral surface of the same.
- c) *Ramus occipitalis* (O. T. posterior terminal branch) behind the ear, upward; it anastomoses with the a. occipitalis and the a. temporalis superficialis.




489. Deep arteries of the face, viewed from the right.

(The parotid gland has been entirely removed, the zygoma, the ramus of the lower jaw and the muscles of the lower jaw have been partially removed [see Fig. 345]; the canalis mandibulae has been chiselled open in a part of its course.)

8. **A. maxillaris interna** (internal maxillary artery) (see also Figs. 490—492) runs medially forward from the collum mandibulae (1st portion), then, tortuous, forward, medianward and upward, usually on the lateral surface of the m. pterygoideus externus between it and the m. temporalis (2nd portion) to the fossa pterygopalatina, in which or before which (3rd portion) it divides into its terminal branches. Nearly in half the cases it lies on the medial surface of the m. pterygoideus externus and arrives between the two heads of this muscle into the fossa pterygopalatina. Branches:

1st portion (four branches):

- a) *A. auricularis profunda* backward to the joint of the jaw, auditory canal and tympanic membrane.
- b) *A. tympanica anterior* upward through the fissura petrotympanica [Glaseri] into the middle ear.
- c) *A. alveolaris inferior* (O. T. mandibular or inferior dental artery) runs downward and forward between the ramus mandibulae and the m. pterygoideus internus to the foramen mandibulare; it passes through the canalis mandibulae, giving off branches to the roots of the teeth, gums and bone. Branches:
 - ca) *Ramus mylohyoideus* goes off before the entrance of the artery into the canalis mandibulae and runs, in the sulcus mylohyoideus, forward and downward.
 - cb) *A. mentalis* leaves the canal through the foramen mentale and goes to the chin.



490. Arteries of the skull and of the nasal cavity, right half, viewed from within.


(The canalis pterygoideus and the canales palatini have been partly chiselled open.)

8. **A. maxillaris interna** (see also Figs. 489, 491 and 492) (continued):

- d) *A. meningea media* (middle meningeal artery) bends under the m. pterygoideus externus upon its medial surface, passes upward to the foramen spinosum and is distributed inside the skull, at first imbedded in the sulci arteriosi of the sphenoid bone and the parietal bone, to the dura mater of the frontal, temporal and parietal region. Through the fissura orbitalis superior it anastomoses with the a. lacrimalis. Branches:
 - da) *Ramus meningeus accessorius* goes off outside the skull, often directly from the a. maxillaris interna, to the mm. pterygoidei, tuba auditiva and its neighborhood.
 - db) *Ramus petrosus superficialis* (O. T. petrosal branch) (see Fig. 493) arises just above the foramen spinosum, runs lateralward and backward to the hiatus canalis facialis and anastomoses there with the a. stylomastoidea (from the a. auricularis posterior).
 - dc) *A. tympanica superior* (see Fig. 493) arises like the preceding branch, goes through the apertura superior canaliculi tympanici and through the canaliculus tympanicus into the middle ear and upon the promontorium.

2nd portion (four [muscular] branches):

- e) *A. masseterica* goes lateralward through the incisura mandibulae to the m. masseter.



491. Arteries of the nasal septum, viewed from the left.

8. *A. maxillaris interna* (see also Figs. 489, 490 and 492) (continued):


- f) *Aa. temporales profundae posterior et anterior* (posterior and anterior deep temporal arteries) upward in the m. temporalis, the posterior close upon the squama temporalis, the anterior upon the ala magna ossis sphenoidalis.
- g) *Rami pterygoidei*, varying in number, often from f) or h), to the mm. pterygoidei.
- h) *A. buccinatoria* (O. T. buccal branch), downward and forward to the m. buccinator, for muscles of the face, mucous membrane of the mouth and gums of the upper jaw.

3rd portion (four branches):

- i) *A. alveolaris superior posterior* (posterior dental or alveolar branch) arises near the fossa pterygopalatina, passes forward and downward through the foramina alveolaria maxillae in the canales alveolares to the sinus maxillaris, upper molar teeth and their gums.
- k) *A. infraorbitalis*, forward through the fissura orbitalis inf. to the sulcus and canalis infraorbitalis, gives off branches to the orbital contents, and also the *aa. alveolares superiores anteriores* (O. T. anterior dental branch) in the canales alveolares to the anterior teeth and their gums; terminal branches through the foramen infraorbitale to the face.
- l) *A. palatina descendens* (O. T. posterior or descending palatine branch) passes downward to the canalis pterygopalatinus. Above it gives off the *a. canalis pterygoidei* [Vidii] (O. T. Vidian artery) which goes through the canalis pterygoideus backward to the roof of the pharynx. Below it divides into the *aa. palatinae minores et major*, which run through the canales palatini and emerge from the foramina palatina minora et majus; the minor branches go to the soft palate and its surroundings, the major branch goes to the inferior surface of the hard palate as far forward as the gums.
- m) *A. sphenopalatina* (O. T. sphenopalatine or nasopalatine artery), through the foramen sphenopalatinum into the nose; it supplies the lateral wall of the nose and the pharynx by means of the *aa. nasales posteriores laterales*, runs at the inferior surface of the body of the sphenoid to the nasal septum and breaks up into the *aa. nasales posteriores septi* which anastomose in the canalis incisivus with the a. palatina major.

9. *A. pharyngea ascendens* (ascending pharyngeal artery) (see Figs. 492 and 493) from the posterior surface, near the a. lingualis, ascends, at first between the a. carotis externa and interna, then in front of the latter between it and the lateral wall of the pharynx, medial from the mm. styloglossus and stylopharyngeus, as far as the base of the skull. Branches:

- a) *Rami pharyngei*, 2-3, partly descending, partly ascending, to the pharynx and palate.




492. Deep arteries of the neck, viewed from the right.

(The muscles of the neck and face have been in large part removed; the bones of the face have been partially removed.)

9. A. pharyngea ascendens (continued).

- b) *A. meningea posterior* ascends upon the lateral surface of the *a. carotis interna* and goes through the foramen jugulare to the dura mater (see Fig. 498).
- c) *A. tympanica inferior* (see Fig. 493) passes through the fossula petrosa and the canaliculus tympanicus into the middle ear and upon the promontorium.




493. A. carotis interna and the middle ear, viewed from the right.

(The temporal bone has been cut open and drawn as in Fig. 13; the plexus venosus caroticus internus has been introduced after N. Rüdinger. Magnification: 3:2.)

A. carotis interna (internal carotid artery) (see also Figs. 485, 492, 494, 499 and 537), curved slightly like the letter S, runs in front of the mm. longus colli and longus capitis and in front of the tubercula anteriora of the procc. transversi of the 4th—2nd cervical vertebra to the base of the skull, bounded medianward by the lateral wall of the pharynx; lateralward and behind it adjoins the v. jugularis interna, in front and lateralward below the a. carotis externa, above the glandula parotis and the mm. styloglossus and stylopharyngeus; the latter muscles separate it from the a. carotis externa. It then enters the canalis caroticus, and passes through it going first upward, then horizontally medianward and forward, runs through the foramen lacerum in a groove of the fibrocartilago basalis, turns upward and goes, slightly curved, in the sulcus caroticus of the body of the sphenoid bone to a position below the processus clinoides anterior; there it bends sharply backward and upward, medial from the proc. clinoides anterior and behind the foramen opticum, perforates the dura mater and breaks up into its terminal branches lateral from the chiasma opticum, below the substantia perforata anterior, *a. cerebri anterior* and *a. cerebri media*. Inside the canalis caroticus it is surrounded by the plexus venosus caroticus internus, in the sulcus caroticus by the sinus cavernosus (see also Fig. 533). In the neck it gives off no branches, in the canalis caroticus the *ramus caroticotympanicus*, in the sinus cavernosus small branches to the surrounding parts and during its course behind the foramen opticum, from the convexity of its last curve, the *a. ophthalmica*.

1. **Ramus caroticotympanicus** (O. T. tympanic branch), a fine branch which passes through one of the canaliculi caroticotympanici to the mucous membrane of the middle ear.

2. **A. ophthalmica** (see Figs. 494, 915, 916, 919—921 and 924), at the lateral inferior surface of the n. opticus, goes through the foramen opticum into the orbit, then bends somewhat upward and passes transversely over the n. opticus, between it and the m. rectus superior, to the junction of the medial with the upper wall of the orbit; there it passes, below the m. obliquus superior, forward and divides between the trochlea and the ligamentum palpebrale mediale into its terminal branches: *aa. palpebrales mediales*, *dorsalis nasi* and *frontalis*. Branches:




494. Arteries of the right orbit, viewed from above.

Magnification: 10 : 7.

(The bony roof and the contents of the orbit have been partially removed.)

2. A. ophthalmica (continued):

- a) *A. centralis retinae* (central artery of the retina) (see Fig. 919 and p. 804) generally arises in common with c), perforates the medial inferior surface of the sheath of the optic nerve and passes in the axis of the same to the retina.
- b) *A. lacrimalis* (lacrimal artery) runs forward over the m. rectus lateralis at the lateral wall of the orbit; it sends several aa. ciliares posteriores and small branches into the foramen zygomaticoorbitale, perforates the lacrimal gland, supplies it and ends at the lateral angle of the eye in aa. palpebrales laterales (one for each eyelid, see p. 417) which send fine branches, aa. conjunctivales posteriores, to the conjunctiva. The a. lacrimalis anastomoses with the a. meningea media through the fissura orbitalis superior.
- c) *Aa. ciliares posteriores breves et longae* (short and long posterior ciliary arteries), 4—6, partly from neighboring branches go, manifoldly divided, to the posterior section of the bulbus (see pp. 801 and 802).
- d) *Rami musculares*, partly from neighboring branches, several in number, among them frequently one large, lateral, superior and one medial, inferior to the eye muscles.
- e) *Aa. ciliares anteriores* (see p. 802) arise usually from the anterior ends of the rami musculares, run forward in the tendons of the straight right eye muscles (see p. 802).
- f) *A. supraorbitalis* (see also Figs. 487, 488 and 492) passes just below the roof of the orbit to the incisura supraorbitalis and through this to the forehead.
- g) *A. ethmoidalis posterior* (see also Figs. 490 and 491) through the foramen ethmoidale post. to the posterior ethmoidal cells and to the posterior superior part of the nasal cavity.




495. Arteries of the base of the brain.

(The pole of the right temporal lobe and the right side of the cerebellum have been cut away.)

2. *A. ophthalmica* (continued):

- h) *A. ethmoidalis anterior* (see Figs. 490, 491 and 494) passes through the foramen ethmoidale anterius upon the lamina cribrosa, there gives off the *a. meningea anterior* upward to the dura mater, then perforates the lamina cribrosa and supplies the anterior part of the wall of the nasal cavity (*aa. nasales anteriores*).
- i) *Aa. palpebrales mediales* (not illustrated), one in each eyelid lateralward; they form by anastomosis of each medial artery with one *a. palpebralis lateralis* (from the *a. lacrimalis*) the *arcus tarseus superior* and *inferior*, situated near the free margin of the eyelid, in front of the tarsus. They give off fine branches to the conjunctiva (*aa. conjunctivales posteriores*) (see also p. 802).
- k) *A. dorsalis nasi* (O. T. nasal branch) (see Figs. 487 and 492) perforates the m. orbicularis oculi above the ligamentum palpebrale mediale, passes downward and anastomoses with the *a. angularis* (from the *a. maxillaris externa*).
- l) *A. frontalis* (see Figs. 487, 488, 492 and 494) goes through the incisura frontalis to the forehead and supplies the muscles and skin in that situation.




496. A. cerebri media and a. chorioidea of the right side, viewed from below.

(The temporal lobe and the occipital lobe have been removed to open up the cornu inferius ventriculi lateralis and to permit the fossa cerebri lateralis [Sylvii] to come into view; the cerebellum and the pons have been cut off just in front of their anterior margin.)

3. **A. communicans posterior** (posterior communicating artery) (see also Figs. 495, 497—499) passes backward over the sinus cavernosus, below the tractus opticus, lateralward from the tuber cinereum and the corpus mamillare, gives off small branches to the parts named and opens into the a. cerebri posterior (from the a. vertebralis).

4. **A. chorioidea** (O. T. anterior choroid artery) (see also Fig. 495), delicate, goes between the tractus opticus and the gyrus hippocampi backward and lateralward to the cornu inferius of the lateral ventricle and breaks up in the plexus chorioideus ventriculi lateralis into branches; it sends besides small branches to the surrounding parts.


5. **A. cerebri media** (middle cerebral artery) (see also Figs. 495, 498 and 499) runs lateralward to the fossa cerebri lateralis [Sylvii] and in this to the lateral surface of the frontal, parietal and temporal lobe; it supplies the substantia perforata anterior, the lateral portion of the gyri orbitales, the island and the gyri frontalis inferior, centrales, the lobuli parietales, the gyri supramarginalis, angularis and temporalis superior; it is connected by numerous anastomoses with the aa. cerebri anterior and posterior.



497. Arteries of the medial surface of the right cerebral hemisphere.

6. **A. cerebri anterior** (see also Figs. 495, 496, 498 and 499) runs medianward above the n. opticus, sends branches to the cerebral ganglia through the substantia perforata anterior and to the chiasma opticum, approaches closely the vessel of the same name of the other side and forms a broad anastomosis with it by means of the *a. communicans anterior*. It next goes into the fissura longitudinalis cerebri around the genu corporis callosi, lying directly upon it, and passes backward upon the corpus callosum. It supplies the corpus callosum, the medial surface of the frontal and parietal lobes almost as far as the fissura parietooccipitalis, the gyrus rectus, the lobus olfactorius, the gyri orbitales as far as the medial limb of the sulci orbitales and the gyri frontales superior et medius and sends a branch to the falx cerebri. It communicates by numerous anastomoses with the aa. cerebri media and posterior.

A. subclavia (see Figs. 478, 485, 492 and 500) arises on the right side from the a. anonyma, on the left side directly from the arch of the aorta (see p. 407); the right is therefore shorter than the left, which ascends at first behind the v. anonyma sinistra, on the left side of the trachea. The artery runs on each side lateralward in a curve behind the m. sternocleidomastoideus over the pleural cupola and over the first rib, being convex upward. In front it is bounded at first by the v. subclavia (crossed in front by the ansa subclavia and the nn. vagus and phrenicus), then by the m. scalenus anterior, behind by the mm. scaleni minimus and medius and comes to lie in the scalenus fissure in the sulcus subclaviae of the first rib. From there on it goes, in the depth of the fossa supraclavicularis major, downward and lateralward to the axilla, behind and below the middle of the clavicle; it is bounded behind and above by the plexus brachialis and is covered by lymph glands and adipose tissue; the a. transversa scapulae crosses its anterior surface and the v. subclavia leans against it in front and below. From the lower margin of the m. subclavius (at the surgeons: of the m. pectoralis minor) on, its continuation is called the *a. axillaris*. Arising from it medianward from the m. scalenus anterior are the following branches: *a. vertebralis*, *a. mammaria interna* and the *truncus thyrocervicalis*, which again divides in a variable manner into the *a. thyroidea inferior*, *a. cervicalis ascendens*, *a. cervicalis superficialis* and *a. transversa scapulae*; medially or dorsally from the m. scalenus anterior the *truncus costocervicalis* is given off, which divides into the *a. intercostalis suprema* and the *a. cervicalis profunda*; in the space between the mm. scaleni (sometimes a little lateralward from it) the *a. transversa colli* arises.



498. Arteries of the base of the brain in their relation to the base of the skull


after removal of the brain; right half of the skull, viewed from the left. (The falx cerebri has been completely removed, the tentorium cerebelli partially. The a. meningea posterior and the ramus meningeus a. vertebralis have been drawn upon the dura mater.)

1. **A. vertebralis** (see Figs. 492, 495, 499 and 500), from the upper circumference of the a. subclavia, passes, at the lateral margin of the m. longus colli, lateralward and behind the a. carotis communis, curved slightly backward to the foramen transversarium of the 6th (sometimes the 5th, 4th, 3rd or 2nd, very seldom the 7th) cervical vertebra, then through the foramina transversaria of the 6th—2nd cervical vertebra, in front of the emerging cervical nerves, upward; it is next curved markedly lateralward to the foramen transversarium of the atlas and goes medianward in the sulcus arteriae vertebralis (see Figs. 267, 273 and 392) in a curve behind the massa lateralis atlantis, covered directly by the mm. obliquus capitis superior and rectus capitis posterior major. Between the atlas and os occipitale it then perforates the dura mater, bends upward, forward and medianward in front of the medulla oblongata to the clivus and unites on it acute angled with the vessel of the same name of the other side, approximately in front of the lower, posterior margin of the pons, to form the unpaired **a. basilaris**. This goes upward and forward upon the clivus in the sulcus basilaris of the pons and, at the upper, anterior margin of the pons, below the substantia perforata posterior and at the upper margin of the dorsum sellae, divides into its two terminal branches, the *aa. cerebri posteriores*. —

Branches of the a. vertebralis:

- a) *Rami spinales*, from the cervical portion, go through the foramina intervertebralia to the canalis vertebralis and its contents. They give off branches to the neighboring muscles.
- b) *Ramus meningeus* arises a little in front of the point where the vessel passes through the dura mater and extends, through the foramen magnum, upward to the dura mater of the posterior fossa of the skull.

A. communicans anterior



499. Arteries of the base of the brain in their relation to the skull after removal of the brain; viewed from above.

(On the right side the a. carotis interna has been exposed in its course in the sinus cavernosus.)

1. **A. vertebralis** (continued) (see also Figs. 495—497) (branches inside the dura mater):
 - c) *A. spinalis posterior*, delicate, bends around the lateral margin of the medulla oblongata and runs downward, forming anastomoses with that of the other side, in the sulcus lateralis posterior of the spinal cord, in front of and lateralward from the posterior roots; it unites with branches of the rami spinales of the a. vertebralis, aa. intercostales, lumbales and sacrales laterales and supplies the spinal cord and its membranes.
 - d) *A. spinalis anterior*, delicate, unites with the artery of the opposite side at the upper end of the spinal cord and runs downward, partly unpaired, partly plexiform being composed of one to four elongated meshes, in front of the fissura mediana anterior as far as the filum terminale; like the preceding artery it receives a variable number of branches of the rami spinales, and supplies the spinal cord and its membranes.
 - e) *A. cerebelli inferior posterior* (posterior inferior cerebellar artery) turns backward to the posterior part of the inferior surface of the cerebellum; it gives off branches to the plexus chorioideus ventriculi quarti.

Branches of the **a. basilaris** (see Figs. 495—499):

- f) *A. cerebelli inferior anterior*, paired, runs lateralward to the anterior part of the inferior surface of the cerebellum.
- g) *A. auditiva interna* (O. T. internal auditory artery) goes lateralward to the nn. facialis and acusticus and with these into the meatus acusticus internus (see p. 854).
- h) *Rami ad pontem* (O. T. transverse or pontal arteries), numerous small branches, to the substance of the pons.
- i) *A. cerebelli superior*, paired, arises shortly before the terminal subdivision, extends at the anterior, upper margin of the pons, lateral- and backward, forms a loop around the pedunculus cerebri so as to arrive upon the upper surface of the cerebellum and supplies mainly this region; it gives off branches to the plexus chorioideus ventriculi tertii.
- k) *A. cerebri posterior* (posterior cerebral artery), paired, at first parallel to the preceding artery, separated from it by the n. oculomotorius and further lateralward by the n. trochlearis and by the tentorium cerebelli; it goes at the anterior margin of the pons around the pedunculus cerebri to the inferior surface of the occipital lobe. It gives off branches through the substantia perforata posterior to the cerebral ganglia, branches to the cerebral penduncles and to the plexus chorioideus ventriculi lateralis and supplies the whole occipital lobe and the temporal lobe with the exception of the gyrus temporalis superior. It communicates by numerous anastomoses with the aa. cerebri anterior and media.


The a. cerebri posterior receives near its origin the a. communicans posterior (from the a. carotis interna); in this way arises the *circulus arteriosus [Willisi]* (O. T. circle of Willis) which lies over the sella turcica and the diaphragma sellae and surrounds the chiasma opticum, the tuber cinereum and the corpora mamillaria.

2. **Truncus thyrocervicalis** (O. T. the thyroid axis) (see also Figs. 485 and 500) arises at the medial margin of the m. scalenus anterior from the anterior wall. Branches:

- a) *A. thyroidea inferior* runs, in front of the m. scalenus anterior, at first upward and then behind the v. jugularis interna, the n. vagus and the a. carotis communis in front of the a. vertebralis medianward and forward to the posterior surface of the glandula thyroidea (see also Fig. 685), into which it penetrates (*rami glandulares*); it gives off on its way the *rami pharyngei, oesophagei, tracheales*, as well as the *a. laryngea inferior* (see Fig. 486), which goes below the m. circopharyngeus into the larynx and anastomoses upon the lateral surface of the internal laryngeal muscles with the a. laryngea superior (from the a. thyroidea superior).
- b) *A. cervicalis ascendens*, usually a branch of a), ascends upon the origins of the mm. scaleni and levator scapulae, at first medianward from the n. phrenicus, behind the v. jugularis interna often going as far upward as the base of the skull; it gives off below several *rami spinales* into the foramina intervertebralia as well as *rami musculares* to the surrounding muscles; one of the latter, *ramus profundus* (see also Fig. 492), runs between the transverse processes of the 4th and 5th cervical vertebra backward to the deep muscles of the neck; it can take the place of the a. cervicalis profunda.
- c) *A. cervicalis superficialis* goes behind the m. omohyoideus transversely through the fossa supraclavicularis major in front of the plexus brachialis and of the mm. scaleni and levator scapulae to the inner surface of the m. trapezius and supplies this and adjacent muscles. It alternates in its formation with the a. transversa colli.
- d) *A. transversa scapulae* (O. T. suprascapular or transversalis humeri) (see also Fig. 503) bends downward and forward, in front of the m. scalenus anterior, to the posterior surface of the clavicle and runs along this bone over the lig. transversum scapulae superius to the fossa supraspinata; it then runs behind the collum scapulae in front of the lig. transversum scapulae inferius to the fossa infraspinata, and there forms a broad anastomosis with the a. circumflexa scapulae. Near the incisura scapulae arises the *ramus acromialis*, which goes through the m. trapezius to the rete acromiale.

3. **A. mammaria interna** (internal mammary artery) (see also Fig. 501) goes from the inferior wall of the a. subclavia medianward and downward behind the v. subclavia directly upon the pleura and then vertically downward just behind the cartilages of the 1st—7th rib. It runs nearly parallel to the lateral margin of the sternum, is separated from it, increasing from above downward, one to two cm. and is covered behind by the pleura and by the m. transversus thoracis. In the region of the 6th intercostal space it divides into its two terminal branches: *a. musculophrenica* and *a. epigastrica superior*. Branches:

- a) *Aa. mediastinales anteriores*, delicate branches, backward to the structures lying in the spatium mediastinale anterius.
- b) *Aa. thymicae* (not illustrated), partly from neighboring vessels, backward to the thymus.
- c) *Rami bronchiales* (not illustrated) to the lower end of the trachea and the bronchi; often absent.




500. A. subclavia dextra, viewed from the right.

(Most of the clavicle has been removed, as have been the muscles attached to the clavicle, the m. scalenus anterior and the mm. pectorales and deltoideus.)

3. A. mammaria interna (continued) (see also Fig. 501):

- d) *A. pericardiophrenica* with the n. phrenicus in front of the root of the lung on the pericardium downward to the diaphragm; it supplies the pericardium and the diaphragm.
- e) *Rami sternales*, medianward to the posterior surface of the sternum; there they form a network of anastomoses with the vessels of the same name of the other side.
- f) *Rami perforantes*, 6—7, the uppermost between the two heads of the m. sternocleidomastoideus (see Fig. 485) or between the clavicle and the first rib, the others forward through the uppermost 6 intercostal spaces. They supply the m. pectoralis major (*rami musculares*) and the skin of the front of the chest (*rami cutanei*). Among the latter that which comes from the 2nd intercostal space is usually very large, bends downward to the mamma and supplies it by means of the *rami mammarii*. The branch perforating the 6th intercostal space is also large and supplies the skin in front of the m. rectus abdominis as far as the navel.
- g) *Rami intercostales*, two for each intercostal space, arise separately or two each from a little trunk. They run lateralward in the six upper intercostal spaces close to the ribs, supply the muscles and the pleura and anastomose with the aa. intercostales. The branch running at the lower margin of each rib is the larger one.


- h) *A. musculophrenica* extends behind the cartilages of the 7th—10th (or 11th) rib, just over the origins of the pars costalis of the diaphragm, lateralward and downward; it gives off *rami intercostales* for the 7th—10th (or 11th) intercostal space and is distributed also in the diaphragm and in the abdominal muscles.
- i) *A. epigastrica superior* passes between the processus xiphoideus and the 7th costal cartilage forward to the m. rectus abdominis, runs downward in it and anastomoses in the region of the navel with the *a. epigastrica inferior*; it gives off several small branches which go forward to the skin in the neighborhood of the navel.
4. **Truncus costocervicalis** (see also Figs. 492 and 512), from the posterior wall, divides after a short, upward course into two terminal branches:
- a) *A. cervicalis profunda* (deep cervical artery), backward between the 1st rib and the proc. transversus of the 7th cervical vertebra, runs upon the mm. semispinalis cervicis and multifidus cervicis as far as the epistropheus. It supplies the muscles of the neck and, by a descending branch, the long muscles of the back; several branches into the intervertebral foramina.
- b) *A. intercostalis suprema* (first intercostal artery) bends downward and backward in front of the neck of the first rib to the 1st intercostal space and also usually in front of the neck of the 2nd rib to the 2nd intercostal space. It gives off *rami posteriores* to the muscles and skin of the back and *rami spinales* into the foramina intervertebralia; in these branches and in its further course it behaves quite like the aa. intercostales (see p. 437).
5. **A. transversa colli** (O. T. transverse cervical artery or transversalis colli) (see Figs. 500 and 503), from the upper wall of the *a. subclavia*, runs among the branches of the plexus brachialis, usually between C₆ and C₇, directly upon the lateral surface of the m. scalenus medius, lateralward and backward and divides beneath the m. levator scapulae into its two terminal branches. It gives off a branch to the fossa supraspinata and supplies the muscles in the neighborhood of the same (see also p. 422).
- a) *Ramus ascendens* extends upward under the m. levator scapulae and the mm. splenii in the neck to supply these muscles.
- b) *Ramus descendens* passes downward between the mm. rhomboidei and the m. serratus posterior superior along the margo vertebralis of the scapula; it supplies these and the adjacent muscles and sends branches to the skin.
- A. axillaris** (*axillary artery*) (see Figs. 485, 500 and 502) runs distalward from the lower margin of the m. subclavius along the lateral angle of the axilla; it is covered at first by the proc. coracoideus, then lies in the groove behind the m. coracobrachialis and is bounded behind by the mm. subscapularis, latissimus dorsi and teres major, medianward by the m. serratus anterior. Superficially (medianward) it is covered by the v. axillaris; on the dorsal, lateral and ventral and partially also on the medial surface it is directly surrounded by the long branches of the plexus brachialis. From the lower margin of the m. pectoralis major on, its continuation is called the *a. brachialis*. Besides small branches to the m. subscapularis, *rami subscapulares*, it gives off the following branches:
1. **A. thoracalis suprema** (O. T. superior thoracic artery) (see Fig. 502) arises behind the m. subclavius and runs downward behind the m. pectoralis minor to the muscles of the 1st and 2nd intercostal space; it may be substituted by a branch of the *a. thoracoacromialis*.
2. **A. thoracoacromialis** (O. T. acromiothoracic or thoracic axis) (see Figs. 485, 500 and 502) arises at the upper margin of the m. pectoralis minor and ends in *rami pectorales* between the mm. pectorales. Branches:
- a) *Ramus acromialis*, transversely behind the mm. pectoralis major and deltoideus, in front of the proc. coracoideus, lateralward, supplies these muscles, perforates the m. deltoideus and ends in the *rete acromiale*, situated upon the upper surface of the acromion.
- b) *Ramus deltoideus* goes downward in the slit between the mm. pectoralis major and deltoideus; it supplies especially the latter muscle and the skin over it.
3. **A. thoracalis lateralis** (O. T. long thoracic artery) (see Figs. 500 and 502) arises behind the m. pectoralis minor and runs upon the m. serratus anterior downward as far as the 5th or 6th intercostal space; it supplies especially the m. serratus anterior. Several branches (*rami mammarii externi*) go through the m. pectoralis major to the skin of the mammary region. It is often a branch of the *a. subscapularis*.
4. **A. subscapularis** (see Figs. 502 and 503), arises at the lower margin of the m. subscapularis (see also p. 426), goes downward and soon divides into two terminal branches:
- a) *A. circumflexa scapulae* (O. T. dorsalis scapulae) bends backward between the mm. subscapularis and teres major through the medial axillary space and then, medial from the caput longum of the m. triceps, goes upward, partly in front of, partly behind the m. teres minor, into the fossa infrapinnata. It supplies the neighboring muscles and skin and anastomoses freely with the *a. transversa scapulae*.



501. Arteries of the anterior wall of the trunk, viewed from behind.

On the left the mm. transversi thoracis and abdominis as well as the posterior leaf of the sheath of the m. rectus have been removed.)

(The aa. epigastricae are drawn upon the surface of the mm. recti abdominis.)



502. Arteries of the right axilla, viewed from in front.


(The arm is lifted, the mm. pectorales have been partially removed.)

4. A. subscapularis (continued):

b) *A. thoracodorsalis*, between the mm. latissimus dorsi and serratus anterior along the margo axillaris scapulae downward and backward, supplies the neighboring muscles and skin.

5. *A. circumflexa humeri anterior* (O. T. anterior circumflex artery) (see Fig. 500) (generally slighter than the following) goes, directly on the anterior surface of the collum chirurgicum humeri, covered by all the muscles, first lateralward, then backward. It supplies these muscles, the shoulder joint and by means of 1—2 branches (*aa. nutriciae humeri*) the superior epiphysis.

6. *A. circumflexa humeri posterior* (O. T. posterior circumflex artery) (see Figs. 500, 503 and 505) arises at the same level as the preceding artery, sometimes in common with it and is not rarely a branch of the a. subscapularis (occasionally arising from a trunk from which originate also the a. subscapularis and a. profunda brachii), passes with the n. axillaris backward through the lateral axillary space between the humerus, mm. subscapularis and teres minor, caput longum m. tricipitis as well as mm. latissimus dorsi and teres major and surrounds the collum chirurgicum from behind and lateralward, lying always upon the bone and covered by all the muscles. It supplies these muscles, the shoulder joint and the skin over the m. deltoideus; it usually gives off two branches (*aa. nutriciae humeri*) to the superior epiphysis of the humerus.




503. Arteries of the right shoulder-blade, viewed from behind.

(The mm. supraspinatus, infraspinatus, teres minor and deltoideus have been partially removed.)

A. brachialis (*brachial artery*) (see Figs. 502, 504, 506 and 507) extends from the inferior margin of the m. pectoralis major onward, at first just behind the m. coracobrachialis, in front of the caput longum and caput mediale of the m. triceps, then in the sulcus bicipitalis medialis almost straight downward; it then passes upon the m. brachialis, covered by the lacertus fibrosus, to the depth of the elbow, there to divide at the level of the proc. coronoideus ulnae into the *a. radialis* and the *a. ulnaris*. Superficial to it medianward and in front of it run the vv. brachiales; on its lateral anterior surface above descends the n. medianus, which, however, above the elbow passes in front of the artery to its medial side; on its medial surface lies above the n. ulnaris, on its posterior surface the n. radialis; these, however, soon diverge from it. Besides several small branches to the neighboring muscles and skin it gives off:

1. **A. profunda brachii** (O. T. superior profunda artery) (see Figs. 504 and 505) arises from the upper portion of the vessel, bends backward and downward and follows a spiral course along with the n. radialis in the sulcus n. radialis around the posterior surface of the humerus downward and lateralward; in its course it runs between the origins of the caput laterale and caput mediale m. tricipitis and is covered by the former. Branches:

504. Arteries of the right upper arm, viewed from in front.



1. *A. profunda brachii* (continued) (see also Fig. 505):

a) *Ramus deltoideus* (see also Fig. 502) goes off immediately below the origin of the artery or arises from the a. brachialis or from the a. collateralis ulnaris superior; it runs lateralward, transversely over the anterior surface of the humerus, covered by the mm. coracobrachialis and biceps, to the m. deltoideus and gives off branches to this and to the m. brachialis.

b) *A. collateralis media* runs downward in the substance of the caput mediale of the m. triceps and sometimes reaches the rete articulare cubiti, which it then helps to form.

c) *A. collateralis radialis* (O. T. articular branch of the superior profunda) appears somewhat below the middle of the upper arm between the caput laterale and the caput mediale of the m. triceps (with the n. cutaneus antebrachii dorsalis of the n. radialis), runs just behind the septum intermusculare laterale as far as the epicondylus lateralis and there opens into the rete articulare cubiti. It supplies the neighboring muscles, the skin over the m. triceps and partly also that over the mm. brachialis and biceps.

d) *A. nutricia humeri* (not illustrated) arises sometimes also from a muscular branch or from the a. brachialis itself; it passes through the canalis nutricius to the bone and marrow of the diaphysis.

2. *A. collateralis ulnaris superior* (O. T. inferior profunda artery) (see also Figs. 505—507) arises usually a little below the preceding artery, runs downward with the n. ulnaris on the medial surface of the caput mediale m. tricipitis (behind the septum intermusculare mediale) and enters into the rete articulare cubiti near the olecranon. It supplies the neighboring muscles and skin.

3. *A. collateralis ulnaris inferior* (O. T. anastomotica magna artery) (see also Figs. 505—508) goes off somewhat above the epicondylus medialis humeri, runs medianward in front of the m. brachialis, perforates the septum intermusculare mediale, bends directly upon the bone backward and lateralward and enters into the rete articulare cubiti; it gives off branches to the adjacent muscles.

505. Arteries of the right upper arm, viewed from behind.

(The m. triceps has been partially removed, the other muscles of the upper arm have been completely removed.)

A. radialis (radial artery)


(see also Figs. 504, 506—511) extends radialward, slightly curved, upon the m. supinator and then runs almost straight to the wrist between the m. brachioradialis on one side, the mm. pronator teres and flexor carpi radialis on the other side; it is covered over in its upper portion by the margin of the m. brachioradialis and, below, lies free, covered only by the fascia antebrachii; above it lies, in its course, upon the insertion of the m. pronator teres, below first upon the mm. flexor digitorum sublimis and flexor pollicis longus; then upon the m. pronator quadratus or the radius and is accompanied in the upper half at its radial side by the ramus superficialis n. radialis. Distalward from the proc. styloideus radii it bends to the back of the hand, covered by the tendons of the mm. abductor pollicis longus and extensor pollicis brevis, then runs distalward through the tabatière and beneath the tendon of the m. extensor pollicis longus, passes between the bases of the ossa metacarpalia I and II through the m. interosseus dorsalis I into the palm and ends in the arcus volaris profundus. Branches:

1. **A. recurrens radialis** (O. T. radial recurrent artery) (see Fig. 507) goes off directly after the artery arises, bends upward and backward like a hook along the n. radialis upon the m. supinator; it supplies the adjacent muscles and the skin over them and sends off a branch beneath the origin of the m. brachioradialis through the septum intermusculare laterale into the rete articulare cubiti (see Fig. 508).


2. **Rami musculares** (see Fig. 506) in the forearm, numerous, to the adjacent muscles and to the skin.

3. **Ramus volaris superficialis** (O. T. superficial volar branch) (see Fig. 510) arises at the level of the proc. styloideus radii, runs distalward upon or partly between the bundles of the m. abductor pollicis brevis, otherwise covered only by the thin fascia of the thenar, and goes over usually in the form of a curve into the *arcus volaris superficialis*. It gives off branches to the muscles and skin of the thumb.

Spalteholz, Atlas. 6th ed.



506. Arteries of the right forearm, viewed from in front, 1st layer.



(The mm. pronator teres, palmaris longus, flexor carpi radialis and flexor digitorum sublimis have been partially removed from their origins.)

4. **Ramus carpeus volaris** (O. T. anterior radial carpal) (see Figs. 507 and 511), delicate, goes ulnarward at the distal margin of the m. pronator quadratus, directly upon the bone; it helps to form the *rete carpi volare*.

5. **Ramus carpeus dorsalis** (O. T. posterior radial carpal) (see Figs. 508 and 509) arises from the portion of the artery situated in the back of the hand, extends ulnarward directly upon the bones and ligaments and helps to form the *rete carpi dorsale*.

6. **A. metacarpea dorsalis I** (O. T. dorsalis indicis or dorsal artery of the first finger) (see Fig. 509) goes off shortly before the passage of the artery through the m. interosseus dorsalis I, divides soon afterward and supplies the adjacent margins of the index finger (as far as the first phalanx) and thumb. A small branch to the radial margin of the thumb usually arises separately.

A. ulnaris (*ulnar artery*) (see also Figs. 504, 507—511) goes into the depth at the upper margin of the m. pronator teres and then turns ulnarward beneath this as well as beneath the mm. flexor carpi radialis, palmaris longus and flexor digitorum sublimis, directly upon the m. flexor digitorum profundus; it then runs along the volar margin of the m. flexor carpi ulnaris, between it and the m. flexor digitorum sublimis to the wrist. It is hidden in its course beneath the margins of the neighboring muscles, is accompanied ulnarward for part of the way by the n. ulnaris and rests upon the m. flexor digitorum profundus. In the hand it runs beneath the lig. carpi volare, then upon the lig. carpi transversum on the radial side of the os pisiforme and ends in the arcus volaris superficialis. Branches:

1. **Aa. recurrentes ulnares** (O. T. ulnar recurrent arteries) (see also Fig. 505), single or in a common trunk from the beginning of the artery; they run in a re-

current direction, that is, proximalward; one runs usually upon the anterior surface of the m. brachialis toward the a. collateralis ulnaris inferior, the other runs backward beneath the origins of the mm. flexor carpi radialis and flexor digitorum sublimis to the sulcus n. ulnaris of the epicondylus medialis humeri and there enters into the rete articulare cubiti. It supplies the surrounding muscles and skin.

2. **Rami musculares**, in the forearm, numerous branches, to the surrounding muscles and to the skin.

507. Arteries of the right forearm,

viewed from in front, 2nd layer. (The mm. brachioradialis, pronator teres, flexor carpi radialis and flexores digitorum sublimis et profundus have been removed.)

3. **Ramus carpeus volaris** (O. T. anterior ulnar carpal) (see also Fig. 511), usually double, delicate, goes in the neighborhood of the wrist, upon the bone, to the *rete carpi volare*.


4. **Ramus carpeus dorsalis** (O. T. posterior ulnar carpal) (see Figs. 508, 509 and 511) runs, directly upon the bone, around the ulna to the back of the hand and to the *rete carpi dorsale*.


5. **Ramus volaris profundus** (O. T. deep ulnar artery) (see Figs. 510 and 511) arises close to the os pisiforme, goes into the depth between the origins of the mm. flexor digiti V brevis and abductor digiti V through the origin of the m. opponeus digiti V and helps to form the *arcus volaris profundus*; it gives off small branches to the muscles of the hypothenar. From this branch, or from the stem of the a. ulnaris itself, an artery (*a. digitalis volaris*) passes beneath the m. palmaris brevis upon the muscles of the hypothenar (giving branches to these and to the skin) to the ulnar margin of the 5th finger.

6. **A. interossea communis** (O. T. common interosseous artery) (see also Fig. 506) goes off a little below the tuberositas radii and divides immediately into two branches:

a) **A. interossea volaris** (O. T. anterior interosseus artery) runs distalward directly upon the volar surface of the membrana interossea, covered by the m. flexor digitorum profundus; it supplies the muscles of the volar surface and sends a series of branches through the membrane to the muscles of the posterior surface; it also sends branches (*aa. nutritiae*) to the radius and ulna. It reaches the dorsal surface of the membrana interossea at the upper margin of the m. pronator quadratus and runs distalward upon it to the *rete carpi dorsale* (see Figs. 508 and 509); a small branch remains upon the volar surface and goes to the *rete carpi volare* (see also Fig. 511). Branch:

aa) **A. mediana** (O. T. median artery or arteria comes nervi mediani) (see Fig. 506) from the first portion of the artery, often directly from the a. ulnaris, slender runs at the ulnar side of the n. medianus distalward, finally to reach the palm.





508. Arteries of the right forearm, viewed from behind.

(The m. extensor digitorum communis has been completely removed, the mm. extensores carpi, indicis proprius, digiti V proprius partially.)


6. A. interossea communis (continued):


b) *A. interossea dorsalis* (O. T. posterior interosseous artery) perforates the membrana interossea a little below the chorda obliqua, appears at the lower margin of the origin of the m. supinator and then runs almost straight distalward upon the ulnar origins of the mm. abductor pollicis longus, extensor pollicis longus and extensor indicis proprius. It is distributed to the adjacent muscles and skin and ends in the rete carpi dorsale. Branch:

ba) *A. interossea recurrens* (O. T. interosseous recurrent or posterior interosseous recurrent) runs upward upon or near the m. supinator (covered by the m. anconaeus), gives off branches to both muscles and to the skin covering them and goes to the rete articulare cubiti.

Rete articulare cubiti (see also Fig. 505) is a wide vascular network, situated especially on the dorsal surface of the elbow joint, in part more superficially upon the olecranon and the tendon of the m. triceps, in part deeper between this tendon, the humerus and the elbow joint. Opening into it from above are the aa. collaterales ulnares superior et inferior, collateralis media and collateralis radialis, from below large branches from the posterior a. recurrens ulnaris and the a. interossea recurrens and small branches from the a. recurrens radialis. A little above the capsule of the joint there is usually a large, transverse anastomosis, which is fed especially by the a. collateralis ulnaris inferior. Branches go off from the network to the surrounding bones, joints, muscles and skin.

Rete carpi dorsale (O. T. posterior carpal rete) (see also Fig. 509) lies upon the back of the wrist and is arranged in two layers: one more superficial, consisting of finer vessels, upon the lig. carpi dorsale and one deeper, directly upon the ligaments and bones of the wrist. The rete is formed by the rami carpei dorsales of the a. radialis and of the a. ulnaris, as well as by the terminal branches of the aa. interossee volaris et dorsalis. From the deep part of the network arise the aa. metacarpeae dorsales (O. T. dorsal interosseous arteries) II, III and IV, which run distalward upon the mm. interossei dorsales II, III and IV, one upon each, each dividing at the level of the heads of the ossa metacarpalia into 2 aa. digitales dorsales (O. T. dorsal digital arteries) for the adjacent margins of the 2nd to the 5th finger. The latter vessels are short and usually end at the first finger-joints. A vessel arises usually directly from the network for the ulnar margin of the little finger. The aa. metacarpeae dorsales receive the rami perforantes of the aa. metacarpeae volares.






510. Arteries of the right palm, superficial layer.

(The aponeurosis palmaris has been removed.)

The **arcus volaris superficialis** (O. T. superficial palmar arch) lies just beneath the aponeurosis palmaris upon the tendons of the m. flexor digitorum sublimis; its convexity is directed distalward. It is formed chiefly by the termination of the a. ulnaris, which unites usually with the delicate ramus volaris superficialis of the a. radialis. It gives off small branches to the surrounding muscles and skin as well as three *aa. digitales volares communes* (O. T. palmar digital arteries), which extend distalward upon the 2nd—4th m. lumbricalis, receive the *aa. metacarpeae volares* II—IV (see also Fig. 511) and divide at the metacarpophalangeal joints, each into two *aa. digitales volares propriae* (O. T. collateral digital arteries) for the adjacent surfaces of the 2nd—5th finger.




511. Arteries of the right palm, deep layer.

(All the structures have been removed with the exception of the mm. interossei, the muscles of the ball of the little finger and the insertions of the mm. opponens pollicis and flexor pollicis brevis.)

The *aa. digitales volares proprias* (see also Fig. 510) supply the volar surfaces and, from the region of the first joint of the finger on, also the dorsal surfaces of the fingers.

The **arcus volaris profundus** (O. T. deep palmar arch) lies directly upon the proximal ends of the ossa metacarpalia and the mm. interossei, covered by the tendons of all the flexors of the fingers and by the m. adductor pollicis. It forms a flatter curve than the superficial arch; its convexity looks distalward. In its formation is chiefly concerned the termination of the a. radialis, which anastomoses with the ramus volaris profundus of the a. ulnaris. Proximalward it gives off branches to the rete carpi volare, distalward four *aa. metacarpeae volares* (O. T. princeps pollicis, radialis indicis and palmar interosseous arteries) which run upon the mm. interossei. The first goes to the two margins of the thumb and to the radial margin of the index finger. The 2nd, 3rd and 4th artery gives off, close to its origin, a *ramus perforans*, which passes through the mm. interossei to the corresponding a. metacarpea dorsalis; the arteries finally anastomose, close to the heads of the ossa metacarpalia, with the *aa. digitales volares communes*.


The **rete carpi volare** (O. T. anterior carpal rete) (see also Fig. 507) lies directly upon the volar surface of the wrist and consists of several fine branches from the a. interossea volaris, from the rami carpei volares of the *aa. radialis* and *ulnaris* and from the arcus volaris profundus.



512. Aorta thoracalis, viewed from in front.

(On the left side the mm. intercostales interni and subcostales have been removed; the diaphragm has been cut away at its origins. A part of the a. subclavia dextra has been cut off.)

The **aorta thoracalis** (*thoracic aorta*) (see also Figs. 607, 681—692) runs, from the left side of the body of the 4th thoracic vertebra on, downward and somewhat to the right and passes in front of the lower margin of the 12th thoracic vertebra, a little to the left of the median plane, through the hiatus aorticus of the diaphragm and there goes over into the aorta abdominalis. On the left side it is directly adjacent to the v. hemiazygos and is covered by the pleura mediastinalis (and is accordingly in direct contact with the cavum pleurae); on the right side it adjoins the ductus thoracicus and the v. azygos, above also the oesophagus, which further down lies on its anterior surface; in front it comes into relation also with the bronchus sinister and with the pericardium in the region of the left atrium. It gives off branches to the thoracic viscera, *rami viscerales*, and to the walls of the thorax, *rami parietales*.



513. Branches of the arteriae lumbales dextrae.

(A piece of the lumbar spine with its muscles, viewed from the right and somewhat from above.)

A. Rami viscerales: 1. **Aa. bronchiales** (see Fig. 512), 2—3, from the anterior wall of the first part of the vessel, very frequently (especially on the right side) from the a. intercostalis III. They go forward to the right and left bronchus and along with these into the lungs.

2. **Aa. oesophageae** (not illustrated), 3—7, from the anterior wall at different levels, forward and to the right to the oesophagus; the lowermost anastomoses with the a. gastrica sinistra.


3. **Rami pericardiaci** (not illustrated), delicate, to the posterior wall of the pericardium.

4. **Rami mediastinales** (not illustrated), arising partly in common with the preceding, slender, going to the contents of the spatium mediastinale posterius; the lowermost spread out also upon the pars lumbalis of the diaphragm as the aa. phrenicae superiores (not illustrated).

B. Rami parietales: **Aa. intercostales** (see also Fig. 512), from the posterior wall, on each side (11) 10 for the (2nd) 3rd—11th intercostal space and the lower margin of the 12th rib. The upper arteries run upward in a recurrent direction, the lower more horizontally; the right are the longer. All of them lie upon the lig. longitudinale anterius, behind the truncus sympathicus and the v. azygos or hemiazygos, the right also behind the oesophagus and the ductus thoracicus. Each arrives at the lower margin of a capitulum costae and there divides.

a) **Ramus posterior** gives off a **ramus spinalis**, which goes through the foramen intervertebrale into the canalis vertebralis to the spinal cord and its surroundings. It then goes backward, medianward from the lig. costotransversarium anterius, gives off **rami musculares**, and divides into two branches. One of these runs between the mm. semispinalis and longissimus, appears in the vicinity of the median plane and turns lateralward to the skin (**ramus cutaneus medialis**); the other lies between the mm. longissimus and iliocostalis and goes often (especially in the lower segments) also to the skin (**ramus cutaneus lateralis**).

b) **Ramus anterior** runs forward in the spatium intercostale beneath the v. intercostalis internally of the mm. intercostales externi, at first covered only by the fascia endothoracica and the pleura, then between the mm. intercostales externi and interni, in the anterior half within the latter or at their inner side (see p. 285); it gives off a branch to the upper margin of the next lower rib and also many **rami musculares**. In front it anastomoses with the corresponding ramus intercostalis of the a. mammaria interna or of the a. musculophrenica. Besides the digitations of origin of the m. obliquus abdominis externus **rami cutanei laterales** [**pectorales et abdominales**] (not illustrated) go, some backward (**ramus posterior**) to the lateral, some forward (**ramus anterior**) to the anterior wall of the trunk; **rami mammarii laterales** go from the latter to the mammary gland. Only in the 4th—6th intercostal space, as a rule, small branches pass to the skin medial from the nipple (**rami cutanei anteriores** [**pectorales et abdominales**]) (not illustrated) and some to the mammary gland (**rami mammarii mediales**).



514. Aorta abdominalis in the male, viewed from in front.

(On the right side the kidney and the mm. psoas and quadratus lumborum have been removed.)

The **aorta abdominalis** (see also Figs. 550 and 686) runs downward on the anterior surface of the lumbar vertebrae, a little to the left of the median plane. It is directly covered in front by the corpus pancreatis, the pars inferior duodeni and the peritoneum, lies on the left side of the v. cava inferior and divides in front of the lower margin of the 4th lumbar vertebra into the *aa. iliacae communes dextra et sinistra*. Its branches are divisible into *rami parietales* (to the walls of the abdomen) and *rami viscerales* (to the abdominal viscera).

A. Rami parietales:

1. **A. phrenica inferior**, paired, arises in front of the 12th thoracic vertebra, separated from, or in common with that of the other side, goes upward, forward and lateralward on the inferior surface of the pars lumbalis diaphragmatis (on the right side behind the foramen venae cavae) and helps to supply the diaphragm. Branches:

Rami suprarenales superiores, from the first part of the artery, to the adrenal.

2. **Aa. lumbales I—IV** (O. T. lumbar arteries) (see also Fig. 513), paired, go off, usually at a right angle, from the posterior wall, one in front of the body of each of the 1st—4th lumbar vertebra; they run backward on the lateral surface of the same, covered by the crura diaphragmatis and the origins of the m. psoas major, and pass through the m. quadratus lumborum to the muscles and to the skin of the lateral wall of the abdomen. Each gives off one branch:

Ramus posterior, corresponding exactly to the r. posterior of an a. intercostalis (see p. 437), gives off a *ramus spinalis* and goes to the muscles and skin of the lumbar region.

3. **A. sacralis media** (O. T. middle sacral artery) (see also Figs. 521 and 522), unpaired, from the posterior wall of the aorta a little above the bifurcation, extends downward, somewhat tortuous, in the median plane in front of the 5th lumbar vertebra, sacrum and coccyx. It gives off branches to the neighboring bones, muscles, nerves and to the rectum and ends in a vascular skein, the *glomus coccygeum* (not illustrated), situated beneath the tip of the coccyx. Branch:

A. lumbalis ima runs lateralward and downward, one on each side, upon the body of the 5th lumbar vertebra to the mm. psoas major and iliacus.

B. Rami viscerales: 3 paired and 3 unpaired; sometimes a branch to the ureter.


1. **A. suprarenalis media** (O. T. middle suprarenal artery), paired, arises a little below the a. coeliaca and runs directly in front of the crus mediale diaphragmatis to the adrenal.

2. **A. renalis** (O. T. renal artery) (see Fig. 550), paired, arises nearly at a right angle at the level of the 1st lumbar vertebra (or somewhat lower), runs transversely in front of the crus mediale diaphragmatis (on the right behind the v. cava inferior) behind the v. renalis and sends its branches in front of and behind the pelvis of the kidney to the substance of the kidney (see also Fig. 691). Variations in the origin and number are frequent. Each, by means of small branches to the tunica adiposa of the kidney, anastomoses with the aa. phrenica inferior, lumbales, suprarenales, spermatica interna and mesentericae. Each gives off also the *a. suprarenalis inferior* to the adrenal and two small branches to the ureter.

3. **A. spermatica interna** (O. T. spermatic artery) (see also Figs. 515, 516 and 550), paired, arises, at an acute angle, at the level of the 2nd lumbar vertebra, runs downward, lateralward and forward (on the right in front of the v. cava inferior) upon the mm. psoas major and minor, in front of the ureter, behind the peritoneum, gives off branches to the tunica adiposa of the kidney, to lymph glands and seldom to the ureter and approaches the lateral margin of the pelvic inlet. Thence it is distributed differently in the two sexes. In the male, as the *a. testicularis* (see also Figs. 518, 521, 550 and 703), it goes upon the m. psoas major, in front of the a. and v. iliaca externa to the annulus inguinalis abdominalis and, surrounded by the plexus pampiniformis, in the spermatic cord to the epididymis and testicle; it anastomoses there with the a. deferentialis. In the female, as the *a. ovarica* (see Figs. 522, 523 and 554), it crosses the a. and v. iliaca externa, nearly in front of the ureter, enters into the lig. suspensorium ovarii, surrounded by the plexus pampiniformis, runs along the margo mesovaricus of the ovary, gives off branches to it and forms a broad anastomosis with the ramus ovarii of the a. uterina.

4. **A. coeliaca** (O. T. coeliac axis) (see also Figs. 515 and 549), unpaired, from the anterior wall between the 12th thoracic and the 1st lumbar vertebra, is short and divides at the upper margin of the corpus pancreatis into three branches:

- a) **A. gastrica sinistra** (O. T. gastric or coronary artery) (see Figs. 515 and 516) bends forward and upward in the plica gastropancreatica to the right side of the cardia and runs in the omentum minus, along the curvatura minor of the stomach, to the right and downward. Small branches to the oesophagus (*rami oesophagei*, see p. 437), larger ones to the anterior and posterior wall of the stomach and to the omentum minus; all the arteries supplying the stomach freely anastomose with each other.
- b) **A. hepatica** (O. T. hepatic artery) (see Figs. 515 and 516) runs upon the right crus mediale diaphragmatis at the upper margin of the pancreas behind and below the vestibulum bursae omentalis to the left, posterior surface of the pylorus. Branches:
 - ba) **A. gastrica dextra** (O. T. pyloric artery) arises at the pylorus and runs in the omentum minus along the curvatura minor to the left; it unites with the a. gastrica sinistra (see above) and gives off branches to the anterior and posterior wall of the stomach.




515. Arteries of the abdominal viscera, deepest layer, viewed from in front.

(The left lobe of the liver, the largest part of the stomach, the small intestines and the colon transversum have been removed; the peritoneum has been, in large extent, dissected off.)

4. *A. coeliaca* (continued):

- bb) *A. hepatica propria* ascends between the layers of the lig. hepatoduodenale, to the left of the ductus choledochus and in front of the v. portae (see Figs. 548 and 635), to the porta hepatis and there divides into a *ramus dexter* (O. T. right terminal branch) to the right half and a *ramus sinister* (O. T. left terminal branch) to the left half of the liver (see also Fig. 632). The former gives off the *a. cystica* to the gall bladder.
- bc) *A. gastroduodenalis* (gastroduodenal artery) runs downward on the left and dorsal surface of the pars superior duodeni and divides into two branches:
 - α) *A. pancreaticoduodenalis superior* extends downward and to the left on the concave side of the pars descendens and pars inferior duodeni, between it and the caput pancreatis, unites with the *a. pancreaticoduodenalis inferior* and supplies, with its *rami pancreatici*, the caput pancreatis, and, with its *rami duodenales*, the duodenum.
 - β) *A. gastroepiploica dextra* (right gastroepiploic artery) (see Fig. 516) arrives at the lower surface of the pars superior duodeni between the two anterior layers of the omentum majus, runs to the left along the curvatura major of the stomach and unites with the *a. gastroepiploica sinistra*. Branches to the anterior and posterior wall of the stomach (see p. 439), as well as *rami epiploici* to the great omentum.
- c) *A. lienalis* (O. T. splenic artery) (see Figs. 549, 637 and 638) bends to the left and extends to the left above the v. lienalis at the upper margin of the pancreas, as well as behind the bursa omentalis, in front of the left adrenal and kidney, in a tortuous course to the hilus of the spleen. Branches (see p. 442):



516. Arteries of the abdominal viscera, more superficial layer, viewed from in front.

(The left lobe of the liver, the small intestines and the colon transversum have been removed;
 the mesenterium and mesocolon have been dissected off.)

4. **A. coeliaca** (continued):

- ca) *Rami pancreatici*, numerous, to the corpus and to the cauda pancreatis.
- cb) *A. gastroepiploica sinistra* (left gastroepiploic artery) (see also Fig. 516) runs forward in the lig. gastrolienale, goes to the right, between the two anterior layers of the omentum majus, along the curvatura major of the stomach, and unites with the *a. gastroepiploica dextra*; its branches are like those of the latter artery (see also p. 439).
- cc) *Aa. gastricae breves* (see Fig. 515) in the lig. gastrolienale to the fundus ventriculi.
- cd) *Rami lienales* from the hilus to the substance of the spleen (see Fig. 637).

5. **A. mesenterica superior** (superior mesenteric artery) (see also Figs. 514—516, 549 and 635), unpaired, arises just below the *a. coeliaca*, separated from it by the *v. lienalis*, in front of the 1st lumbar vertebra; it goes downward at first behind the caput pancreatis, then passes through the incisura pancreatis in the groove on the anterior surface of the processus uncinatus pancreatis and in front of the pars inferior duodeni between the layers of the mesenterium and runs in a curve, slightly convex to the left and forward, as far as the end of the small intestine.


- a) *A. pancreaticoduodenalis inferior* (see Figs. 515 and 516) goes off from the right side, runs to the right on the concave surface of the pars inferior duodeni between it and the pancreas, supplies both organs and unites with the *a. pancreaticoduodenalis superior*.
- b) *Aa. intestinales* (O. T. vasa intestini tenius), 12—20, from the convexity of the arch on the left side, pass between the layers of the mesenterium to the loops of the jejunum (*aa. jejunales*) and ileum (*aa. ileae*). Each artery bifurcates and forms arch-like anastomoses with the branches of neighboring vessels; the branches arising from these form further anastomoses and so on until two to five series of anastomoses follow upon one another (see Fig. 547). From the last numerous branches of nearly equal size and almost equidistant from one another go to the small intestines. Small branches go also to the contents of the mesentery. The last *a. ilea* unites with the *a. ileocolica*.
- c) *A. ileocolica*, from the right side, runs behind the peritoneum to the right and downward to the end of the small and beginning of the large intestine, anastomoses with the last *a. ilea* and with the *a. colica dextra*. A small branch, the *a. appendicularis* (not indicated), goes behind the ileum to the vermiform process.
- d) *A. colica dextra* (right colic artery), from the right side, very often a branch of the preceding artery, goes to the right behind the peritoneum to the colon ascendens; it anastomoses with the *a. ileocolica* and with the *a. colica media*.
- e) *A. colica media* (middle colic artery), from the right side, in front of the pancreas; it extends to the right and forward in the mesocolon transversum to the colon transversum; it anastomoses with the *a. colica dextra* and with the *a. colica sinistra*.

6. **A. mesenterica inferior** (see Figs. 513—516), unpaired, arises between the 3rd and 4th lumbar vertebra and runs downward and to the left, beneath the peritoneum. Branches:

- a) *A. colica sinistra* (left colic artery) beneath the peritoneum to the left to the colon descendens; it anastomoses with the *a. colica media* and with the *aa. sigmoideae*.
- b) *Aa. sigmoideae*, often single, run downward and to the left to the colon sigmoideum; they anastomose with the *a. colica sinistra* and with the *a. haemorrhoidalis superior*.
- c) *A. haemorrhoidalis superior* (see also Figs. 516, 521 and 522), the terminal branch of the *a. mesenterica inf.*, bends downward in front of the *a. and v. iliaca communis sinistra* and in front of the promontorium and goes downward in the mesorectum, in front of the sacrum and behind the rectum, and then divides into two branches which run right and left to the intestinum rectum; it anastomoses above with the *aa. sigmoideae*, below with the *aa. haemorrhoidales mediae* (from the *aa. hypogastricae*).


A. iliaca communis (iliac artery) (see also Figs. 514, 518, 519, 521, 522 and 550), paired, goes downward, lateralward and forward, slightly curved, in front of the bodies of the 4th and 5th lumbar vertebra at the medial margin of the *m. psoas major*; it runs on the left behind the *a. haemorrhoidalis superior* and, oftener on the left, behind the ureter. It lies in front of and somewhat to the left of the *v. iliaca communis*. It divides in front of the articulatio sacroiliaca into the *a. hypogastrica* and the *a. iliaca externa*; a small branch to the ureter.

A. hypogastrica (see Figs. 514, 518, 519, 521, 522 and 551) turns downward and somewhat backward into the small pelvis and divides, after a short course, usually into a posterior and an anterior branch. The posterior branch bends markedly backward to the upper part of the foramen ischiadicum majus (above the *m. piriformis*), gives off the *aa. iliolumbalis* and *sacralis lateralis* and ends as the *a. glutea superior*. The anterior branch runs in front of the *m. piriformis* to the inferior portion of the foramen ischiadicum majus, gives off the *aa. umbilicalis* (or *aa. vesicales superiores*), *obturatoria*, *vesicalis inferior*, *deferentialis* (in the male), *uterina* (in the female) and *haemorrhoidalis media* and bifurcates into the *aa. glutea inferior* and *pudenda interna*. The subdivisions of the posterior branch (also the *a. obturatoria* and the *a. glutea inferior*) go chiefly to the walls of the pelvis as *rami parietales*, those of the anterior branch (with the exception of the *a. obturatoria* and the *a. glutea inferior*) chiefly to the viscera as *rami viscerales*. In addition to these a small branch to the ureter.



517. Branches of the a mesenterica superior, viewed from in front.

(The small intestine has been cut away at the mesentery with the exception of two loops, the transverse colon has been drawn upward, the great omentum cut off, and the mesentery reflected to the left. The mesentery and mesocolon have been partially dissected away.)




518. Right a. obturatoria and a. epigastrica inferior, in the male, viewed from the left.

A. *Rami parietales*:

1. *A. iliolumbalis* (O. T. iliolumbar artery) (see also Figs. 514 and 519) turns lateralward and upward behind the n. obturatorius, between the basis ossis sacri and the m. psoas major and divides into two branches:

- a) *Ramus lumbalis* corresponds to the ramus posterior of an a. lumbalis (see Fig. 513), runs dorsally between the proc. transversus of the 5th lumbar vertebra and the basis of the sacrum and gives off a *ramus spinalis* to the spinal canal; it supplies the neighboring muscles and skin.
- b) *Ramus iliacus* runs behind the m. psoas major along the crista iliaca, gives off branches to the bones and muscles; it unites with the a. circumflexa ilium profunda.




519. Abnormal a. obturatoria dextra in the male (*Var.*)

(A window has been cut out of the mm. obturator internus and levator ani.)

2. **A. sacralis lateralis** (lateral sacral artery) (see Figs. 514 and 521) downward, single or double, medially from the foramina sacralia, upon the fascia pelvina of the sacrum, gives off *rami spinales* through the foramina sacralia anteriora into the canalis sacralis, and branches to the muscles and to the skin of the back.

3. **A. obturatoria** (obturator artery) (see Figs. 518, 520—522 and 527), frequently from the posterior branch, extends, covered by peritoneum, forward and downward beneath the linea terminalis pelvis and beneath the n. obturatorius to the canalis obturatorius, inside which it divides into its terminal branches. In the pelvis, small branches to its surroundings, as well as:

- a) **Ramus pubicus.** This runs medianward on the posterior surface of the ramus superior ossis pubis and anastomoses on the upper surface of the lig. lacunare [Gimbernati] (see Fig. 525) or laterally from it with the ramus obturatorius of the a. epigastrica inferior. By this anastomosis the a. obturatoria arises in about one third of the cases from the a. epigastrica, it lies more often laterally from the annulus femoralis, more rarely on the septum femorale or on the lig. lacunare.
- b) **Ramus internus,** medial from the membrana obturatoria, downward to the ischium.



520. Terminal branches of the a. obturatoria dextra (male pelvis), viewed from in front and from the right.

(The muscles of the thigh have been cut off short at their origin.)

3. A. obturatoria (continued): Terminal branches:


- c) *Ramus posterior* goes downward in the membrana obturatoria to the anterior surface of the ramus superior ossis ischii, and gives off, as it goes, the *a. acetabuli*, which passes through the incisura acetabuli to the hip-joint and through the lig. teres to the caput femoris; it also gives off branches which pass through the membrane to the m. obturator internus, and sends its terminal branches in part forward upon the origin of the m. obturator externus, in part backward, especially to the ischium.
- d) *Ramus anterior* forward upon the external surface of the membrane and then, partly through, partly in front of the m. obturator externus, toward the preceding artery.

4. **A. glutea superior** (gluteal artery) (see Figs. 518, 521 and 528), between L_5 and S_1 backward through the foramen suprapiriforme, gives small branches to the surrounding muscles and bones and divides into a superficial branch, which goes lateralward and downward between the mm. gluteus maximus and medius, and into a deep branch, which runs lateralward between the mm. gluteus medius and minimus. This latter branch divides into a *ramus superior* along the upper margin of the m. gluteus minimus and into a *ramus inferior* lower down, on the m. gluteus minimus. It gives off, also, branches to the skin of the region.

5. **A. glutea inferior** (see Figs. 518, 521 and 528) runs between S_2 and S_3 or S_3 and S_4 through the foramen infrapiriforme, between the lower margin of the m. piriformis and the lig. sacrospinous backward and downward to the anterior surface of the m. gluteus maximus. It supplies especially this muscle, its neighbors and the skin covering that region and gives off a fine *a. comitans n. ischiadici* which goes downward to the n. ischiadicus.

B. Rami viscerales.

1. **A. umbilicalis** (O. T. hypogastric artery) (see Figs. 521, 522, 561 and 562), present in its full development only before birth, runs in a curve, on the inner wall of the small pelvis, forward and upward to the anterior abdominal wall, arrives on this behind the lateral margin of the tendon of the m. rectus abdominis and then goes, beneath the peritoneum in the plica umbilicalis lateralis, to the navel (see also Figs. 379 and 650). It gives off the *aa. vesicales superiores* to the upper part of the bladder; from these go small branches to the ureter. After birth only the beginning of the artery and the branches to the bladder remain open, the largest part of the vessel becomes obliterated to form the *ligamentum umbilicale laterale* (O. T. lateral false ligament) (see Fig. 650).




521. Arteries of the male pelvis, viewed from the left.

(The rectum has been drawn somewhat to the left, the peritoneum has been removed.)

2. **A. deferentialis** (O. T. artery of the vas deferens) (present only in the male), frequently arising from the preceding artery or from the following, passes forward and downward to the ductus deferens, there divides, one part going downward as far as the vesicula seminalis, the other upward to the inguinal canal and thence to the epididymis; it anastomoses there with the a. testicularis.

3. **A. vesicalis inferior** (see also Figs. 522 and 523) goes downward to the fundus vesicae, in the male also to the vesicula seminalis and the prostata, in the female to the vagina.


4. **A. haemorrhoidalis media** (middle haemorrhoidal artery) (not constant) (see also Fig. 518) comes frequently from the a. pudenda interna and runs over the m. coccygeus and m. levator ani, downward and medianward to the rectum and to the surrounding muscles; it gives off branches to the vesiculae seminales and to the prostata in the male, to the vagina in the female. It anastomoses also with the aa. haemorrhoidales superior and inferior.



522. Arteries of the female pelvis, viewed somewhat from the left.

(The organs have been partly drawn out of their position; the peritoneum has been removed.)

5. **A. uterina** (uterine artery) (present only in the female) (see also Fig. 523) goes forward beneath the peritoneum at first on the inner wall of the small pelvis, then turns medianward, runs through the parametrium obliquely forward in front of the ureter to the cervix uteri and there gives off a small branch to the ureter as well as the *a. vaginalis*, which runs downward on the lateral wall of the vagina. The uterine artery then runs along the margo lateralis uteri, following a very tortuous course between the layers of the ligamentum latum uteri, as far as the fundus uteri and sends off numerous corkscrew-like, tortuous branches to the anterior and posterior surface of the uterus, anastomosing with one another and with those of the other side; it ends in a branch which runs in the ligamentum latum partly to the ligamentum teres uteri and to the tuba uterina (*ramus tubarius*, anastomosis with *a. ovarica*), partly to the ovary (*ramus ovarii*); the latter branch runs along the margo mesovaricus of the ovary and forms a broad anastomosis with the *a. ovarica*.




523. Arteries of the uterus and surrounding structures, viewed from behind and somewhat from the left.

(On the left side the ovary, the Fallopian tube and the ligamentum teres uteri have been displaced from their normal position.)

6. **A. pudenda interna** (see Figs. 519, 524 and 525) goes downward at the lower margin of the m. piriformis behind the lig. sacrospinous, encompasses the spina ischiadica and passes forward, in the form of a curve, along the medial surface of the ramus superior, then along that of the ramus inferior ossis ischii; in its course it runs somewhat (about the width of the little finger) above the processus falciformis and is covered by the fascia obturatoria; close behind the m. transversus perinei superficialis it divides into the *a. perinei* and the *a. penis* (or *a. clitoridis*). It frequently gives off above the lig. sacrospinous small branches to the surrounding muscles and nerves, and often, also, the *a. haemorrhoidalis media*. Other branches:


- a) *A. haemorrhoidalis inferior* (O. T. external or inferior haemorrhoidal artery) (see Figs. 524 and 525), usually double or triple. It runs, transversely medianward through the fat of the fossa ischiorectalis, being fairly superficial, to the fat, muscles and skin about the anus. It anastomoses with the *aa. haemorrhoidalis media* and *sacralis media*.
- b) *A. perinei* (O. T. superficial perineal artery) (see Figs. 524 and 525) extends, over or under the m. transversus perinei superficialis, forward and medianward; it is superficial, being covered only by the fascia superficialis perinei; it goes in the male to the posterior surface and to the septum of the scrotum by means of the *aa. scrotales posteriores*, in the female into the labium majus and labium minus by means of the *aa. labiales posteriores*. It also gives off small branches to the superficial muscles of the perineum.
- ca) *A. penis* (in the male only) (see Figs. 519, 524, 711, 726 and 730) runs forward, in the direction of the main stem, close to the ramus inferior ossis pubis, within the trigonum urogenitale, leaves it behind the lig. transversum pelvis, passes in front of and below the latter until it lies below the lig. arcuatum pubis where it is continued directly into the *a. dorsalis penis*. Branches:
 - a) *A. bulbi urethrae* (O. T. artery of the bulb) (see Fig. 524) goes off at the posterior margin of the trigonum urogenitale and runs medianward and forward to the bulbus urethrae and to the neighboring muscles.
 - β) *A. urethralis* (s. Fig. 524), forward and medianward to the corpus cavernosum urethrae.



524. Arteries of the male perineum.

(On the left side the dissection shows the *m. transversus perinei profundus*; the *crus penis* has been drawn aside.)

γ) *A. profunda penis* arises close under the *ligamentum arcuatum pubis*, penetrates the *crus penis* on its medial surface and runs in it partly backward, partly forward close to the *septum penis* (see also Fig. 712) as far as its tip; its branches anastomose with one another and with those of the opposite side.



525. Arteries of the female perineum.

(Dissection similar that of Fig. 524; the crus clitoridis has been drawn aside.)

- d) *A. dorsalis penis* (see Figs. 521, 524, 526 and 712) goes near the lig. suspensorium penis to the dorsum penis and in its medial groove, on each side near the unpaired v. dorsalis penis, as far as the glans; it gives branches to the skin, to the corpora cavernosa penis et urethrae and especially to the glans penis.
- cb) *A. clitoridis* (present only in the female) is comparable in its course to the a. penis; it gives off the *a. bulbi vestibuli [vaginae]* to the perineal muscles, the vagina and the bulbus vestibuli, the *a. urethralis* to the latter and to the vestibulum vaginae, and the *a. profunda clitoridis* to the crus clitoridis; the *a. dorsalis clitoridis* runs upon the clitoris to the glans clitoridis, praeputium clitoridis and labia minora.

A. iliaca externa (external iliac artery) (see Figs. 514, 518, 519, 521, 522, 527, 550, 551 and 554) goes downward, lateralward and forward on the medial, anterior surface of the m. psoas major; it runs at first to the lateral side of the v. iliaca externa, in front of the fascia iliaca and behind the peritonaeum; on the right side oftener than on the left the anterior and medial surface of its initial portion is crossed by the ureter. It goes through the lacuna vasorum beneath the lig. inguinale (see Fig. 421) and from there on is called the *a. femoralis*. Branches:

526. Arteries of the right thigh,

viewed from in front, superficial layer.


(The m. sartorius has been for the most part removed and a piece has been cut out of the m. rectus femoris.)

1. **A. epigastrica inferior** (O. T. deep epigastric artery) (see Figs. 501, 518, 519 and 521) arises just above the lig. inguinale and runs in a curve, first obliquely medianward, then vertically upward. In its course it lies at first behind the lig. inguinale and lig. interfoveolare, medial from the annulus inguinalis abdominalis and then,

passing behind the tendon of the m. transversus abdominis, arrives at the posterior surface of the m. rectus abdominis. It causes the peritoneum to bulge forward to form the plica epigastrica (see also Figs. 379 and 650) and thus separates the fovea inguinalis medialis and the fovea inguinalis lateralis from each other. Finally it sinks into the substance of the m. rectus abdominis, supplies this muscle and a part of the skin covering it, and anastomoses freely with the a. epigastrica superior. Branches:

a) **Ramus pubicus** (see Fig. 518) goes medianward behind the pelvic attachment of the m. transversus abdominis just above the ramus superior ossis pubis and gives off the **ramus obturatorius**, which runs downward behind the lig. lacunare [Gimbernati] to the ramus pubicus of the a. obturatoria (see p. 445).

b) **A. spermatica externa** (O. T. cremasteric branch) (in the male) (see Fig. 519), **a. ligamenti teretis uteri** (in the female), perforates the posterior wall of the inguinal canal, enters the spermatic cord or the lig. teres uteri and spreads out in the tunics of the testicle, or in the lig. teres and in the labia majora.



527. Arteries of the right thigh, viewed from in front, deep layer.

(The mm. sartorius, tensor fasciae latae, rectus femoris, pectineus, adductor longus and gracialis have been removed. A large portion of the a. femoralis has been cut off.)

2. **A. circumflexa ilium profunda** (deep circumflex iliac artery) (see also Figs. 514, 518 and 519) arises behind or below the lig. inguinale, passes (at first in a duplication of the fascia iliaca, then upon the latter) lateralward and upward in the groove between the mm. transversus abdominis and iliacus, gives off small branches to the neighboring muscles, as well as one larger branch upward between the mm. transversus and obliquus internus (see Fig. 501) and extends backward along the crista iliaca; anastomosis with the r. iliacus a. iliolumbalis.

A. femoralis (femoral artery) (see also Fig. 526) follows a fairly straight course, in the groove between the mm. pectineus and adductores on the one hand, the m. iliopsoas and m. vastus medialis on the other, downward and medianward and finally through the canalis adductorius. It lies above upon the deep layer of the fascia lata and is covered, from the inferior angle of the trigonum femorale on, by the m. sartorius. Above, it lies lateralward from the v. femoralis, farther down, it lies in front of the same (see also Fig. 556). Branches:


1) **A. epigastrica superficialis** (superficial epigastric artery) (see Fig. 526) arises just below the lig. inguinale, perforates the margo falciformis and runs upward as far as the region of the navel; it gives off branches to the skin and to the m. obliquus externus abdominis.

2) **A. circumflexa ilium superficialis** (superficial circumflex iliac artery) (see Fig. 526), frequently a branch of the preceding, goes lateralward, usually superficially upon the fascia lata, to the spina iliaca anterior superior and supplies the skin and muscles.

3) **Aa. pudendae externae** (O. T. superficial and deep external pudic arteries) (see Fig. 526), usually two in number, run medianward, in front of and behind the v. femoralis to the anterior surface of the scrotum and penis (*aa. scrotales anteriores*) or of the labia majora (*aa. labiales anteriores*).

4) **Rami inguinales** (see Fig. 526) to the lymph glands and muscles of the fossa iliopectinea.

5) **Rami musculares** (see Fig. 526) to the neighboring muscles and skin.



528.

Arteries of the right thigh, viewed from behind.

(The m. gluteus maximus has been reflected inward, the mm. gluteus medius, quadratus femoris and caput longum m. bicipitis have been partially removed.)

6) **A. genus suprema** (O. T. anastomotica magna) (see Fig. 527) goes off in the canalis adductorius, perforates the lamina vastoadductoria (see also p. 772), runs downward between it and the m. sartorius behind the epicondylus medialis femoris and finally accompanies the n. saphenus for a variable distance (*ramus saphenus*). It gives off *rami musculares*, as well as a *ramus musculoarticularis* (O. T. deep branch) in the m. vastus medialis and *rami articulares* (O. T. superficial branch) below the same to the *rete articulare genus*.

7) **A. profunda femoris** (O. T. profunda or deep femoral artery) (see also Figs. 526 and 527) arises in the fossa iliopectinea and descends behind the a. femoralis and a little lateral from it. It runs in front of the mm. iliopsoas, pectineus and adductor brevis, behind the m. adductor longus, and ends as the a. perforans tertia. Branches:

a) **A. circumflexa femoris medialis** (O. T. internal circumflex artery) goes off near the origin and runs medianward behind the a. and v. femoralis. Branches:

aa) **Ramus superficialis**, in front of the m. pectineus, medianward between the mm. adductores longus and brevis.

ab) **Ramus profundus**, a larger branch, bends upward, passes backward between the mm. iliopsoas and pectineus and, below the m. obturator externus, to the anterior surface of the m. quadratus femoris. It sends branches to the hip-joint and divides into a branch ascending to the fossa trochanterica, and a branch descending behind the adductor muscles; it supplies the neighboring muscles and skin.


b) **A. circumflexa femoris lateralis** (O. T. external circumflex artery) lateralward in front of the m. iliacus. Branches:

ba) **Ramus ascendens**, goes upward and lateralward beneath the m. rectus femoris to the mm. glutei and tensor fasciae and to the fossa trochanterica.

bb) **Ramus descendens**, a larger branch, downward and lateralward beneath the m. rectus femoris between the mm.

vasti intermedius and lateralis almost to the knee; it also gives off branches to the skin.

c) **A. perforans prima** (O. T. superior or first perforating artery) perforates the adductors at the lower margin of the m. pectineus.



529. Arteries of the right leg, viewed from behind.

(The muscles of the thigh with the exception of the m. adductor magnus have been completely removed; the mm. gastrocnemius, plantaris, soleus and flexor hallucis longus have been partially removed.)

d) *A. perforans secunda* (see Figs. 527 and 528) perforates the adductors at the lower margin of the m. adductor brevis.

e) *A. perforans tertia* (see Figs. 527 and 528) perforates the adductors somewhat above the canalis adductorius.

All three aa. perforantes supply the adductors and the muscles and skin of the posterior surface.

The *a. nutricia femoris* is usually single and given off by the *a. perforans II*; more rarely it is double (*superior* and *inferior*) and then comes from the aa. perforantes I and III.

A. poplitea (popliteal artery) (see also Fig. 528) runs downward, from the hiatus tendineus [adductorius], behind the planum popliteum femoris, upon the capsula articularis of the knee-joint, upon the posterior surface of the m. popliteus and in front of the tendinous arch between the origins of the m. soleus and divides just below the latter into the *a. tibialis posterior* and the *a. tibialis anterior*. In its course it is covered above by the m. semimembranosus, in the popliteal space, behind and somewhat lateralward, by the v. poplitea and (behind this) is overlapped by the n. tibialis; below, it is hidden beneath the mm. gastrocnemius and plantaris. Branches:

1) **A. genus superior lateralis** (O. T. superior external articular artery) (see also Fig. 530) goes, above the condylus lateralis femoris, directly upon the bone to the *rete articulare genus*.

2) **A. genus superior medialis** (O. T. superior internal articular artery) like the preceding, above the condylus medialis femoris.


3) **A. genus media** (O. T. azygos articular artery) perforates the capsule above the lig. popliteum obliquum and goes to the ligg. cruciata and to the synovial folds.

4) **A. genus inferior lateralis** (O. T. inferior external articular artery) (see also Fig. 530) goes upon the m. popliteus, along the upper margin of the condylus lateralis tibiae, covered by the lig. collaterale fibulare, to the *rete articulare genus*.

5) **A. genus inferior medialis** (O. T. inferior internal articular artery) (see also Fig. 527) goes, below the condylus medialis tibiae, covered by the lig. collaterale tibiale, to the *rete articulare genus*.


6) **Aa. surales**, several, sometimes from one trunk, to the m. triceps surae and to the skin.

A. tibialis posterior (posterior tibial artery) extends downward and medianward in front of the m. triceps surae upon the posterior surface of the m. tibialis posterior in the groove between the mm. flexores digitorum longus and hallucis longus, above in front of the n. tibialis, elsewhere medianly of it, covered by the deep layer of the fascia cruris. Below, it runs in front of the medial margin of the tendon of Achilles in a curve to the regio retromalleolaris medialis lying there nearly in the middle between the malleolus medialis and the tendo calcaneus and there divides into the aa. plantares medialis and lateralis. It gives off branches, to the muscles and to the skin and farther:



530. Arteries of the right leg, viewed from in front.

(The mm. tibialis anterior and extensor hallucis longus have been partially removed.)



1. **Ramus fibularis** (see Fig. 529) extends lateralward toward the collum fibulae to the mm. soleus and peroneus longus.

2. **A. peronaea** (peroneal artery) (see Fig. 529) arises just below the collum fibulae and runs downward on the posterior surface of the m. tibialis posterior, between it and the m. flexor hallucis longus, near the fibula; below, it lies upon the posterior surface of the membrana interossea cruris or of the tibia and breaks up behind the ankle-joint into the *rami calcanei laterales* (O. T. external calcaneal arteries) which go to the *rete calcaneum* and its neighborhood. Numerous branches to the muscles, and further:

- A. nutricia fibulae** (see Fig. 529) into the canalis nutricius fibulae.
- Ramus perforans** (O. T. anterior peroneal artery) (see also Figs. 529 and 531) perforates the membrana interossea just above the syndesmosis tibiofibularis, unites with the a. malleolaris anterior lateralis and helps to form the *rete malleolare laterale* and the *rete calcaneum*.
- A. malleolaris posterior lateralis** (see Fig. 529), directly upon the fibula, to the *rete malleolare laterale*.
- Ramus communicans** (see Fig. 529), upon the posterior surface of the tibia, just above the ankle joint, to the a. tibialis posterior.

3. **A. nutricia tibiae** (see Fig. 529) into the canalis nutricius tibiae.

4. **A. malleolaris posterior medialis** (O. T. internal malleolar branch) (see Fig. 529), directly upon the tibia to the *rete malleolare mediale*.

5. **Rami calcanei mediales** (O. T. internal calcaneal branches) (see Figs. 529 and 532) form, with the rami calcanei laterales (see above), the *rete calcaneum*, which is situated superficially upon the tuber calcanei, as well as upon the surrounding muscles and tendons.

A. tibialis anterior (anterior tibial artery) goes forward above the membrana interossea cruris and directly downward upon its anterior surface. Above, it lies between the m. tibialis anterior and the m. extensor digitorum longus, below, between the former muscle and the m. extensor hallucis longus. Farther down, it is situated upon the anterior surface of the tibia and of the capsule of the upper ankle-joint, as well as behind the m. extensor hallucis longus and the deep layer of the lig. cruciatum cruris and there becomes continuous with the *a. dorsalis pedis*. It gives off numerous branches to the muscles and skin, and farther:

1. **A. recurrens tibialis posterior** (posterior tibial recurrent artery) (see Fig. 529), which comes off at the beginning of the artery, extends, behind the fibula, to the articulatio tibiofibularis. It is often absent.

2. **A. recurrens tibialis anterior** (anterior tibial recurrent artery) upward, upon the anterior surface of the condylus lateralis tibiae, to the *rete articulare genus*, which is spread out partly upon the bones and ligaments of the knee joint, partly upon the tendon of the m. quadriceps femoris, upon the lig. patellae and upon the patella itself (*rete patellae*).

531. Arteries of the back of the right foot.

3) **A. malleolaris anterior lateralis** (O. T. external malleolar artery) to the malleolus lateralis, unites with the ramus perforans of the a. peronaea and helps to form the *rete malleolare laterale* upon the lateral malleolus and its neighborhood.

4) **A. malleolaris anterior medialis** (O. T. internal malleolar artery) goes to the malleolus medialis and helps to form the *rete malleolare mediale*, situated there.

A. dorsalis pedis (see also Fig. 530) extends forward, upon the ligaments and bones of the back of the foot, between the mm. extensores hallucis longus and brevis and breaks up between the proximal ends of the ossa metatarsalia I and II into the *ramus plantaris profundus* and the a. *metatarsae dorsalis I*. Branches:

1. **Aa. tarseae mediales** (O. T. internal tarsal artery), 2—3, run upon the bones and ligaments to the medial border of the foot and help to form the *rete malleolare mediale*.


2. **A. tarsea lateralis** (O. T. external tarsal artery) lateralward and forward upon the bones and ligaments, covered by the m. extensor digitorum brevis, toward the basis ossis metatarsalis V, unites there with the lateral end of the a. arcuata. Branches which anastomose with one another and, with branches of the a. dorsalis pedis and a. arcuata, form the *rete dorsale pedis*.

3. **A. arcuata** (O. T. metatarsal artery) lateralward directly upon the proximal ends of the ossa metatarsalia unites with the a. tarsea lateralis. Branches:

a) **Aa. metatarsae dorsales II to IV** (O. T. tarsal interosseous arteries) run forward upon the 2nd—4th m. interosseus dorsalis and each divides into two aa. *digitales dorsales* (dorsal digital arteries) behaving just as do the arteries to the fingers. Larger *rami perforantes posteriores* and smaller, inconstant *anteriores* between the proximal or between the distal side of the ossa metatarsalia go to the aa. *metatarsae plantares*.


4. **Ramus plantaris profundus** (O. T. communicating, first interosseous perforating or plantar artery) between the proximal ends of the ossa metatarsalia I and II, into the sole of the foot, to the arcus plantaris.

5. **A. metatarsae dorsalis I** forward like the other aa. *metatarsae dorsales* to the great toe and the medial surface of the 2nd toe; it gives off a *ramus perforans anterior* between the heads of the ossa metatarsalia.



532. Arteries of the sole of the right foot.

(The muscles have, for the most part, been removed.)




A. plantaris medialis (see p. 455) (O. T. internal plantar artery) goes forward, between the two layers of the lig. laciniatum (through its upper compartment, see p. 379), and there gives off the *ramus superficialis*, which passes through the m. abductor hallucis, giving off branches to it, arrives at the medial margin of the foot and runs as far as the great toe. It then enters the groove between the mm. abductor hallucis and flexor digitorum brevis (*ramus profundus*), supplies these muscles and the skin and goes, between the two bellies of the m. flexor hallucis brevis, into the a. metatarsae plantaris I or into the medial a. digitalis plantaris of the great toe.

A. plantaris lateralis (see p. 455) (O. T. external plantar artery) runs, between the two layers of the lig. laciniatum (through its lower compartment see p. 379), then between the mm. flexor digitorum brevis and quadratus plantae, forward and lateralward toward the basis ossis metatarsalis V; there it lies more superficially, between the mm. quadratus plantae and flexor digiti V and bends around medianward into the arcus plantaris. It gives off branches to the neighboring bones, ligaments, muscles and skin.

The **arcus plantaris** lies directly beneath the proximal ends of the 2nd to the 4th metatarsal bone and beneath the mm. interossei, between them and the caput obliquum of the m. adductor hallucis, being more superficial lateralward than medianward. It diminishes in caliber from the lateral toward the medial margin of the foot, arises, lateralward, directly from the a. plantaris lateralis and unites medianward, between the ossa metatarsalia I and II, with the ramus plantaris profundus of the a. dorsalis pedis. It gives off recurrent branches to the bones and muscles; running forward from it are the 4 *aa. metatarsae plantares* (O. T. digital or plantar digital arteries). Each of these passes forward between two metatarsal bones, receives the rami perforantes of the aa. metatarsae dorsales and divides at the bases of the first phalanges into the *aa. digitales plantares* (O. T. collateral digital arteries), which behave just as do those to the fingers. The artery for the medial margin of the great toe is usually given off by the a. metatarsae plantaris I; that for the lateral margin of the little toe arises usually as a separate artery from the end of the a. plantaris lateralis.

533. Frontal section through the right sinus cavernosus, viewed from behind. Magnification: 5:2.

The **sinus durae matris** (*sinuses of the dura mater*) are venous canals which run, chiefly, between the divergent layers of the dura mater and lie at the same time in corresponding bony grooves. They stand open on section and are devoid of valves; the sinus sagittalis superior and cavernosus are crossed in a network-like manner by bands of connective tissue. They receive the *vv. cerebri, meningeae, diploicae* (partially), *auditivae internae* and the *vv. ophthalmicae*.



1. Sinus transversus (O. T. lateral sinus)

(see Figs. 534 and 535), paired, collects most of the blood from the skull cavity. It begins on each side at the protuberantia occipitalis interna; it is usually larger on the right than on the left side and runs curved lateralward in the sulcus transversus of the occipital bone and of the parietal bone, in the attached margin of the tentorium cerebelli. Thence, it goes tortuous medianward and downward, in the sulcus sigmoideus of the temporal bone and in the sulcus transversus of the pars lateralis oss. occipitalis, to the foramen jugulare to become continuous there with the upper end of the v. jugularis interna. In the region of the tentorium cerebelli it is triangular-prismatic in shape, elsewhere semi-cylindrical.


2. **Sinus sagittalis superior** (O. T. superior longitudinal sinus) (see Figs. 534, 535, 815 and 816), unpaired, extends from before backward, in a curve, in the attached margin of the falx cerebri near the crista frontalis and in the sulcus sagittalis of the frontal bone, of the parietal bones and of the occipital bone. It opens very often with a right, larger branch into the right and with a left, smaller branch into the left sinus transversus; nearly as often it opens undivided into the right, seldom undivided into the left sinus transversus. It is triangular in transverse section. The sinus anastomoses with the sinus sagittalis inferior.

3. **Sinus sagittalis inferior** (O. T. inferior longitudinal sinus) (see Figs. 534 and 535), unpaired, from before backward in the free margin of the falx cerebri, into the sinus rectus.

4. **Sinus rectus** (straight sinus) (see Figs. 534, 535 and 815), unpaired, goes as continuation of the preceding and after picking up the v. cerebri magna [Galen], as a triangular canal, at the point of attachment of the falx cerebri to the tentorium cerebelli, from in front and above, backward and downward; it opens sometimes undivided into the left sinus transversus, but usually divided with a left, larger branch into the left and with a right, smaller branch into the right sinus transversus. Sometimes in front of the protuberantia occipitalis interna exists a cavity (*confluens sinuum*), into which the sinus sagittalis superior and the sinus rectus open undivided and from which go off both sinus transversi.

5. **Sinus occipitalis** (see Figs. 534 and 535), unpaired or paired, begins at the protuberantia occipitalis interna and goes downward, close to the median plane in the attached margin of the falx cerebelli. Above, it begins oftener divided than single at the beginning of the sinus transversi. Below, it bifurcates, turns lateralward and forward on both sides above the foramen occipitale magnum and enters the sinus transversus, close to the foramen jugulare. It anastomoses with the plexus venosi vertebrales interni and varies much in the degree of its development.


6. **Sinus cavernosus** (see also Figs. 493, 534 and 535), paired, is an irregular space, which is situated, on each side, near the sella turcica and the corpus oss. sphenoidalis. It is bounded above and lateralward by that part of the dura mater which is stretched out from the procc. clinoides anterior and posterior to the angulus superior pyramidis and to the middle skull fossa (see also p. 709); it surrounds, in addition, the a. carotis interna, the n. abducens and the plexus cavernosus. Tributaries:



534. Sinus durae matris, viewed from above.

(The right orbit and sinus cavernosus have been opened; the tentorium has been cut away.)

- a) *V. ophthalmica superior* (superior ophthalmic vein) (see also Figs. 539, 915, 916 and 919), devoid of valves, corresponds only approximately to the a. ophthalmica, which possesses, in addition, two delicate accompanying veins. It anastomoses freely above the lig. palpebrale mediale by its anterior extremity (*v. nasofrontalis*) (see Figs. 538 and 539) with the v. angularis (of the v. facialis anterior), runs backward near the m. obliquus superior, then lateralward between the n. opticus and the m. rectus superior, finally backward through the fissura orbitalis superior into the sinus cavernosus, laterally and above the annulus tendineus communis; it sends an anastomosis to the v. ophthalmica inferior. It receives the *vv. ethmoidales anterior et posterior*, *v. lacrimalis*, *vv. musculares* and the veins of the bulbus oculi and of the eyelids (see p. 802; the *v. centralis retinae* opens into the sinus cavernosus).




535. Sinus durae matris of the right half of the skull,
viewed from the left.

b) *V. ophthalmica inferior* (see Figs. 539, 915 and 916), devoid of valves, arises from the veins of the tear-sac, of the eyelids, &c., in the anterior part of the orbit, between the medial and inferior wall; it runs in the inferior part of the orbit and divides into two branches. One branch goes backward through the fissura orbitalis superior (below the annulus tendineus communis [Zinni]) into the sinus cavernosus; the other passes lateralward through the fissura orbitalis inferior and opens into the ramus profundus of the *v. facialis anterior*. It receives *vv. musculares* and veins of the bulbus oculi and anastomoses medianward with the *v. ophthalmica superior*.

c) *Sinus sphenoparietalis* (see Fig. 534), paired, begins on each side a little behind the sutura coronalis on the parietal bone in a bony groove or in a bony canal, turns medianward beneath the ala parva ossis sphenoidalis and enters the anterior end of the sinus cavernosus. It receives usually one *v. meningea media*, *vv. cerebri inferiores* (from the fissura cerebri lateralis [Sylvii]), as well as the *v. ophthalmomeningea*, and also the *v. diploica temporalis anterior*.

7. and 8. **Sinus intercavernosus anterior** and **sinus intercavernosus posterior** (see Fig. 534), unpaired, run in front of, behind and partly also beneath, the hypophysis and unite the two sinus cavernosi with each other. They form, with these, the **sinus circularis**.

9. **Sinus petrosus superior** (superior petrosal sinus) (see Fig. 534), paired, goes lateralward and backward on each side, from the posterior part of the sinus cavernosus in the sulcus petrosus superior of the temporal bone and in the attached margin of the tentorium cerebelli, to the sulcus sigmoideus and into the sinus transversus.



536. Venae diploicae, viewed from the right.

(The lamina externa of the roof of the skull has been filed off.)

10. **Sinus petrosus inferior** (inferior petrosal sinus) (see Figs. 534 and 535) paired, extends lateralward and downward on each side from the posterior, inferior portion of the sinus cavernosus in the sulcus petrosus inferior of the occipital and temporal bone to the foramen jugulare and runs downward through its most anterior subdivision; it opens, either in the foramen jugulare or below it, into the bulbus v. jugularis superior.

11. **Plexus basilaris** (see Figs. 534 and 535), unpaired, lies as a plexus upon the clivus and is connected above, on both sides, with the sinus cavernosus and the sinus petrosus inferior, below, at the foramen occipitale magnum, with the plexus venosi vertebrales interni.

Vv. cerebri collect the blood partly from the surface, partly from the interior of the brain (see also p. 674) and open particularly into the sinus sagittalis superior, transversus, sphenoparietalis, petrosus superior and rectus; they possess no valves.

Vv. meningeae (see Figs. 534 and 535) stand open, are devoid of valves, are numerous in the dura mater encephali, are connected above with the sinus sagittalis superior and anastomose manifoldly with one another. They accompany the stems of the corresponding arteries, there being usually two veins to one artery, and open into a neighboring sinus. The *vv. meningeae mediae*, usually two in number, run near the a. meningeae media, are connected with the sinus sphenoparietalis and sometimes with the rete foraminis ovalis, extend downward through the foramen spinosum and open into the plexus pterygoideus.

Vv. auditivae internae (see p. 856) (not illustrated) come from the internal ear through the meatus acusticus internus and open into the sinus transversus or the sinus petrosus inferior.

Venae diploicae (see Fig. 536) are thin walled tubes, which lie in the flat *canales diploici* [Brescheti] in the substance of the bones of the roof of the skull. They have no valves; they are arranged in the form of a network and are very variable. The main stems run from above downward and are connected at their extremities by openings (*emissaria*) with the external veins of the skull or with the sinus durae matris or (usually) with both at the same time; they possess valves alone at their mouths which generally allow a flowing off outward only.

- 1) *V. diploica frontalis* opens into the v. supraorbitalis and the sinus sagittalis superior.
- 2) *V. diploica temporalis anterior* into a v. temporalis profunda and the sinus sphenoparietalis.
- 3) *V. diploica temporalis posterior* from the emissarium pariet. to the emissarium mastoid..
- 4) *V. diploica occipitalis* opens into the emissarium occipitale.

Emissarium parietale, in the foramen parietale, unites the sinus sagittalis superior with the superficial veins of the skull; its valves permit a flow in an outward direction only.

Emissarium mastoideum, in the foramen mastoideum, between the sinus transversus and the v. auricularis posterior or the v. occipitalis or the beginning of the v. jugularis externa.

Emissarium occipitale, at the level of the protuberantiae occipitales, opens usually either internally into a sinus transversus or into the confluent sinuum, externally into the v. occipitalis, often into both.

Emissarium condyloideum (not illustrated), in the canalis condyloideus, unites the sinus transversus and the plexus venosi vertebrales externi.

Plexus venosus caroticus internus (see Fig. 493) surrounds the a. carotis interna in the canalis caroticus in the form of a network, is connected above with the sinus cavernosus, receives veins from the cavum tympani and opens below into the v. jugularis interna.

Rete foraminis ovalis (not illustrated) in the foramen ovale from the sinus cavernosus to the plexus pterygoideus; it is often connected above with the vv. meningae mediae.

Rete canalis hypoglossi (not illustrated) in the canalis hypoglossi from the sinus occipitalis and the plexus venosi vertebrales interni to the end of the sinus petrosus inferior.

Vena cava superior (see Figs. 462, 478, 537, 546 and 680), unpaired, free from valves, arises behind the first right costal cartilage by the union of the *vv. anonymae dextra et sinistra* and extends downward and curved moderately convex toward the right, in front of the root of the right lung and at the right side of the aorta ascendens, to the right atrium. It is bounded in front by the thymus gland and on the right by the right lung; it is covered above on its right side by the pleura mediastinalis, below by the visceral layer of the pericardium. In the upper part of its course the v. azygos (see p. 474) opens into it from behind.

V. anonyma (O. T. innominate vein) (see also Figs. 478, 537 and 546) is formed behind the articulatio sternoclavicularis and in front of the a. subclavia by the union of the v. jugularis interna, v. subclavia and v. jugularis externa; the latter opens usually into the v. subclavia. The right vein is bounded on the right by the right lung and runs vertically downward; the left, distinctly longer, extends to the right and downward, behind the manubrium sterni and the thymus, directly over the arcus aortae, in front of the aa. subclavia sinistra, carotis communis sinistra and anonyma. Both veins are devoid of valves. Branches:

1. **Vv. thyreoideae inferiores** (inferior thyroid veins) (see Fig. 537) from the *plexus thyreoideus impar*. This lies partly in front of the lower margin of the isthmus glandulae thyreoideae, in part it surrounds the beginning of the trachea like a ring; it anastomoses with the vv. thyreoideae superiores of both sides and receives, behind, the *vv. tracheales*, the *vv. oesophageae* and the v. laryngea inferior (corresponding to the arteries of the same names). Three veins usually go off from the plexus: a right vein opens at the angle of junction of the vv. anonymae or into the v. anonyma dextra or into the v. jugularis interna dextra, a left vein into the v. anonyma sinistra and a middle, v. thyreoidea ima, running downward in front of the trachea, into the v. anonyma sinistra.

2. **V. mammaria interna** (internal mammary vein) (see Figs. 364 and 537) corresponds almost exactly to the artery of the same name. Arising from the union of the double *vv. epigastricae superiores* (rich in valves) and the *vv. musculophrenicae* it is double in the region of the m. transversus thoracis; above the latter it is single and situated medial from the artery. It receives (with the vv. musculophrenicae) the anterior extremities of the 9(—10) superior *vv. intercostales* (see p. 473), which have valves at their mouths, farther the *rami perforantes* (especially marked in the 1st or 2nd intercostal space), which have valves, as well as *rami sternaes* from the networks which are situated upon both surfaces of the sternum, and in which there are several large transverse connections between the trunks of the two sides, sometimes especially one such on the anterior surface of the processus xiphoideus.

3. **Vv. phrenicae superiores** (not illustrated) possess valves, are double and correspond to the a. pericardiophrenica.

4. **Vv. thymicae, pericardicae, mediastinales anteriores, bronchiales anteriores** (not illustrated) from the viscera of the thorax, open separately or partially in common.

5. **V. vertebralis** (see Fig. 539) corresponds to the cervical part of the a. vertebralis, on the lateral side of which it extends downward from the foramen magnum through the foramina transversaria of the 1st—6th (often —7th) cervical vertebra. It begins in the plexus venosi vertebrales, anastomoses with the v. occipitalis and receives branches from the spinal canal, from the deep muscles of the neck and from the plexus vertebrales externi. It stands open and has valves at its mouth. Branch:

V. cervicalis profunda (see Fig. 537) corresponds approximately to the a. cervicalis profunda, begins near the back of the head in the plexus vertebralis posterior und runs downward behind or in front of the m. semispinalis cervicis; it goes forward beneath the proc. transversus of the 7th cervical vertebra into the v. vertebralis. There are valves at its mouth.

V. jugularis interna (internal jugular vein) (see also Fig. 540) corresponds together with the vv. jugulares externa and anterior to the a. carotis communis. It arises as the direct continuation of the sinus transversus in the lateral compartment of the foramen jugulare with a swelling filling up the fossa jugularis, the *bulbus v. jugularis superior*, above runs downward behind the a. carotis interna on the lateral wall of the pharynx, becomes displaced gradually to the lateral side first of the a. carotis interna, then of the a. carotis communis and extends downward, close to the latter, to the v. anonyma. A little above its termination it presents a spindle-shaped dilatation, the *bulbus v. jugularis inferior*, which is closed above by a single or double valve. Otherwise it is devoid of valves. Branches:

1. **Sinus durae matris** see pp. 459—462.

2. **V. canaliculi cochleae** (not illustrated) comes from the cochlea (see p. 856) through the canaliculus cochleae and opens into the bulbus v. jugularis superior.

3. **V. occipitalis** (see p. 468).


4. **Vv. pharyngeae** vary in number and come at different levels from the *plexus pharyngeus* which covers the outer surface of the pharynx and receives branches from the deep muscles of the pharynx, from the tuba auditiva, the palate and dura mater (*v. meningea*, corresponding to the a. meningea posterior), as well as the *v. canalis pterygoidei* [*Vidii*] (from the canalis pterygoideus); the plexus connect with the neighboring veins. The inferior vv. pharyngeae may open also into other branches of the v. jugularis interna. They are all devoid of valves.

5. **V. lingualis** (lingual vein) (see also Fig. 540) runs backward at first exactly as does the a. lingualis, later, however, on the lateral surface of the m. hypoglossus; it receives a large *v. sublingualis* and the *vv. dorsales linguae*. The a. lingualis is accompanied medially from the m. hypoglossus by small veins, which together with the *v. comitans n. hypoglossi* open into the v. lingualis. All branches possess valves; the vein communicates with the vv. pharyngeae and the v. thyreoidea superior; it opens sometimes into a v. facialis or a v. thyreoidea superior.

6. **Vv. thyreoideae superiores** (superior thyroid veins) (see also Fig. 540), usually two, accompanying the artery only in part; one opens far below, the other above, near the a. thyreoidea superior. The latter receives a *v. sternocleidomastoidea* and the *v. laryngea superior* (corresponding to the a. laryngea superior), which may also open into the v. facialis communis, and frequently opens into the v. facialis communis. These veins possess valves.


7. **V. facialis communis** (see also Figs. 539 and 540) arises beneath the angle of the lower jaw, from the *vv. faciales anterior* and *posterior*, which may also open separately into the v. jugularis interna. It runs downward and backward, covered only by the platysma, crosses the a. carotis externa and terminates at the level of the hyoid bone. It has no valves. Branches:

a) *V. facialis anterior* (see also Figs. 538—540) corresponds in its course to that of the a. maxillaris externa, which itself possesses in addition two small accompanying veins; it is a little tortuous and runs usually behind the artery and in part more superficially, covered only by the mm. zygomaticus, risorius and platysma, from the root of the nose to the angle of the lower jaw. It arises superficially at the medial angle of the eye as the *v. angularis*, formed by the *vv. frontalis* (usually unpaired, and then very frequently opening into the left v. angularis), *supraorbitalis* (runs transversely medianward partly beneath the m. orbicularis oculi over the margo supraorbitalis, anastomoses with the vv. temporales superficialis and media) and *nasofrontalis* (see p. 460); in front of the root of the nose it forms a broad union with the vein of the other side. Above, it receives further: the *vv. palpebrales superiores*, *nasales externae*, *palpebrales inferiores* and *labialis superior*, besides the *ramus profundus*; this is formed in the fossa infratemporalis by the *vv. alveolares superiores* (corresponding to the arteries of the same name) and by branches of the plexus pterygoideus, and receives one branch of the v. ophthalmica inferior; it runs forward upon the facies infratemporalis maxillae and terminates below the arcus zygomaticus. Farther down enter the *v. labialis inferior*, *vv. massetericae* and *vv. parotideae anteriores*. Beneath the lower jaw it receives further the large *v. submental*, which is connected with the v. sublingualis and v. jugularis anterior, as well as one *v. palatina*, which corresponds to the a. palatina ascendens. All the branches of the vein possess efficient valves at their mouths; the main trunk also contains such valves at the level of the ductus parotidis.



537. Deep veins of the neck, viewed from the right.

(The muscles of the neck and face have been in large part removed; the bones of the face have been partially removed.)




538. Superficial veins of the face, viewed from the right.

(With partial use of a preparation by W. Braune.)

(Below the zygoma, the subcutaneous fat and a part of the muscles have been removed.)


- b) *V. facialis posterior* (see also Figs. 537 and 539) arises superficially in front of the ear through the union of the vv. temporales superficialis et media, runs downward in the substance of the glandula parotis behind the ramus mandibulae and lateral from the carotis externa and bends obliquely forward below the angulus mandibulae, over or under the m. digastricus, to the v. facialis communis. It gives off a large branch at a variable level to the v. jugularis externa. It receives:
- ba) *V. temporalis superficialis*, from the superficial plexus of the roof of the skull, runs downward behind the trunk of the artery. It possesses a valve in front of the ear.
- bb) *V. temporalis media* runs transversely backward, deviating from the artery, in the m. temporalis, anastomoses at the lateral angle of the eye with the superficial veins, in the muscle with the vv. temporales profundae and perforates the fascia temporalis over the root of the zygoma. It possesses valves.
- bc) *Vv. auriculares anteriores*, from the lateral surface of the external ear.
- bd) *Vv. parotidae posteriores*, from the glandula parotis.



539. Deep veins of the face, viewed from the right.

(The bones and muscles have been removed as in Fig. 489.)

- be) *V. articulares mandibulae*, from a plexus surrounding the joint of the jaw, which also receives the *vv. tympanicae* (not illustrated) (through the fissura petrotympanica [Glaseri] from the middle ear) and veins from the external auditory passage and eardrum membrane and is connected with the plexus pterygoideus.
- bf) *V. stylomastoidea* (not illustrated), corresponding to the artery of the same name, anastomoses at the hiatus canalis facialis with the *vv. meningae mediae*.
- bg) *V. transversa faciei* (transverse facial vein), frequently double, corresponds to the artery of the same name.
- bh) *Plexus pterygoideus* extends from the fossa pterygopalatina as far as the collum mandibulae and lies, with wider vessels, on both sides of the m. pterygoideus externus, with narrower vessels, also on the medial surface of the m. pterygoideus internus and on the membranous wall of the tuba auditiva; it forms a sheath for the a. maxillaris interna, whose branching area corresponds to its inlets. It receives: the *v. sphenopalatina* (from the nasal cavity through the foramen sphenopalatinum), the *vv. meningae mediae* (see p. 462), the *vv. temporales profundae* (from the m. temporalis, in which it forms a plexus with branches of the v. temporalis media), branches from the maxilla, the *vv. massetericae*, *v. alveolaris inferior* (the latter two corresponding to the arteries of the same names) and the *rete foraminis ovalis* (see p. 463). It anastomoses with the *vv. articulares mandibulae* and with the plexus pharyngeus and empties in part through the ramus profundus v. facialis anterioris, especially however through one or two valve-bearing vessels, which accompany the a. maxillaris interna, into the v. facialis posterior.



540. Veins of the neck, viewed from in front.

(On the right the m. sternocleidomastoideus, on the left the m. pectoralis major, has been removed.)

V. jugularis externa (external jugular vein) (see also Figs. 537—539) arises below the ear by two roots: the anterior, frequently the larger, comes from the v. facialis posterior (see p. 466), the posterior arises behind the ear. It crosses the m. sternocleidomastoideus at a sharp angle, covered by the platysma, goes above the clavícula into the depth and opens usually into the v. subclavia, generally with the v. jugularis anterior. Valves at its mouth and in its middle. Branches:

1. **V. occipitalis** corresponds to the a. occipitalis. It opens partly into the v. jugularis externa, partly, doubled, accompanies the a. occipitalis to the v. jugularis interna; it anastomoses with the v. vertebralis and the plexus vertebralis posterior.

2. **V. auricularis posterior** from the superficial plexus behind the ear.

3. **V. jugularis anterior**, downward from the chin upon the mm. mylohyoideus and sternohyoideus, extends lateralward below the origins of the m. sternocleidomastoideus to the v. jugularis externa or the v. subclavia. It anastomoses with the adjacent veins, and, by the large *arcus venosus juguli*, below the glandula thyroidea with the vein of the opposite side. More rarely instead of this vein only an unpaired median trunk (*v. mediana colli*) is present.


541. Superficial veins of the back of the right hand.

(Partly after W. Braune.)

The **veins of the upper extremity** run partly in the depth between the muscles, partly in plexuses, superficially, upon the fascia superficialis. Both possess numerous valves, the deep veins more than the superficial, and are manifoldly connected with one another by anastomoses, which are devoid of valves.


The **deep veins** (not illustrated) follow almost exclusively the corresponding arteries and are double, with the exception of those in the fingers and in the axilla. In the *palms* they begin as single *vv. digitales volares propriae* (see Fig. 542), which unite to form the *vv. digitales volares communes* and go into the double *arcus volaris venosus superficialis*. In the depth run the *vv. metacarpeae volares*, which open into the double *arcus volaris venosus profundus*; they send in addition several *rami perforantes* through the spatia interossea metacarpi to the deep veins of the back of the hand, several smaller and regularly two larger veins into the 1st metacarpal space and through the muscles of the little finger to the rete venosum dorsale manus. From the palmar arches arise the *vv. ulnares* and *vv. radiales* and from these again the *vv. brachiales* with their tributaries, corresponding exactly to the arteries. On the **back of the hand** deep, small veins accompany the deep layer of the rete (arteriosum) carpi dorsale (see p. 432) and empty partly into the *vv. radiales*, partly into the rete venosum dorsale manus.

The **superficial veins of the back of the hand** represent the chief canals of outflow for the skin. They begin upon the phalanges of the fingers in a plexus, which permits of the recognition of longitudinal trunks, *vv. digitales dorsales propriae*, and each terminates on the first phalanx in the *arcus venosus digitalis* which is concave proximalward. From the union of every pair of these arches, arise larger *vv. metacarpeae dorsales* (originally four); these open in a very variable manner in a distal convex *arcus venosus dorsalis manus*, the position of which corresponds to about the middle of the ossa metacarpalia, or they form the *rete venosum dorsale manus*, the meshes of which are elongated in the direction of the long axis of the limb.



542. Superficial veins of the right forearm, viewed from in front.


(Palm after W. Braune.)



The **superficial veins of the palm** are much finer than those of the back of the hand. They begin upon the phalanges in a plexus in which one can make out more or less distinctly the *vv. digitales volares propriae* and out of which most of the blood is conducted to the posterior surface through several vessels at the lateral margins of the fingers. Similarly the *vv. intercapitulares* (see also Figs. 541) conduct blood in the spaces between the fingers from the volar to the dorsal vessels. The superficial network of the palm lies upon the aponeurosis palmaris and upon the fascia of the thenar and hypothenar eminences; in the latter two places the network is somewhat closer. Proximalward it goes into the beginning of the *v. mediana antebrachii*.

The **superficial veins of the forearm** are also arranged in the form of a plexus and very variable. Among them three are usually very distinct (see also Figs. 864 and 865):

1. **V. basilica** (see also Fig. 541) begins upon the back of the hand, at the ulnar end of the *arcus venosus dorsalis* or at the *rete venosum dorsale manus*, loops around the ulnar side of the forearm to its volar surface and runs on this as far as the elbow, then in the *sulcus bicipitalis medialis* along the upper arm as far as its middle third. There it perforates the fascia, goes still some distance upward and unites sooner or later with one of the (more delicate) *vv. brachiales*. On the forearm it is thinner, on the upper arm it is thicker than the *v. cephalica*.
2. **V. cephalica** (see also Fig. 541) arises on the back of the hand, mostly from the radial end of the *arcus venosus dorsalis* or from the *rete venosum dorsale manus*, crosses the *tabatière*, winds around the radial side of the forearm to its volar surface and extends on this as far as the elbow; thence it goes upward in the *sulcus bicipitalis lateralis* and (in a duplicature of the *fascia brachii*) in the groove between the *mm. pectoralis major* and *deltoideus*, sinks into the *fossa infraclavicularis* and after union with a *v. thoracoacromialis* empties into the *v. axillaris* (see also Fig. 543). Usually a second longitudinal vessel is situated radially on the forearm, the *v. cephalica accessoria*, which opens into the main vein.




543. Veins of the right axilla, viewed from in front.

(Arrangement as in Fig. 502.)

3. *V. mediana antebrachii* (see Fig. 542), smaller than the vv. basilica and cephalica, originates from the superficial network of the hollow of the hand, extends on the ulnar side of the volar surface of the lower arm to the region of the elbow and opens for the most part into the v. basilica, more rarely into the v. mediana cubiti. Occasionally it bifurcates and runs partly to the v. basilica as *v. mediana basilica*, and partly to the v. cephalica as *v. mediana cephalica*. In the flexure of the elbow more frequently the *v. mediana cubiti* (see Fig. 542) leads as a connecting branch, from the v. cephalica in front of the lacertus fibrosus obliquely proximalward to the v. basilica; not seldom it conducts the main mass of the blood of the v. cephalica to the v. basilica. In both cases the superficial vessels laterally from the lacertus fibrosus anastomose freely through a valveless vessel with the deep veins.

V. axillaris (*axillary vein*) (see also Fig. 537) arises usually at the lower margin of the m. pectoralis major by the union of the two vv. brachiales and extends as far as the clavicle on the medial and anterior (superficial) surface of the artery, which is also accompanied directly by small veins. Its branches correspond to those of the artery. The *v. thoracalis lateralis* receives the *v. thoracoepigastrica* (see Fig. 555 and p. 484), which extends subcutaneously upward from the v. femoralis or from the v. epigastrica superficialis on the anterior and lateral surface of the trunk. Into this or into the neighboring veins empty the *vv. costoaxillares* which, in the upper 6—7 intercostal spaces, carry the blood from the middle portions of the vv. intercostales (see p. 473) to the axilla. The v. axillaris and its branches are richly provided with valves.




544. Spinal veins, seen in a median section, viewed from the left.

V. subclavia (*subclavian vein*) (see Fig. 537) is the name of the continuation of the v. axillaris from the lower margin of the m. subclavius on. It accompanies the a. subclavia on its anterior and inferior surface and crosses the first rib in front of the m. scalenus anterior. Behind the articulatio sternoclavicularis it helps to form the v. anonyma. It has valves at its beginning and at its termination. In addition to it some small veins directly accompany the a. subclavia. Tributaries (besides the v. jugularis externa):

1. and 2. **V. transversa scapulae** and **v. transversa colli** (see Figs. 537 and 540), near the corresponding arteries, unite usually to form one stem, which frequently opens into the v. jugularis externa. They carry valves at their mouths.

Veins of the spinal column. In the canalis vertebralis, between the dura mater spinalis, the periosteum and the ligaments of the canal, lie close venous plexuses, **plexus venosi vertebrales interni**, the tributaries of which come partly from the vertebrae, partly from the spinal cord. These plexuses have no valves and (see also Fig. 545), extend chiefly in the longitudinal direction from the foramen occipitale magnum to almost as far as the tip of the sacrum and permit the recognition of four trunks, two anterior and two posterior. The *anterior* lie upon the posterior surface of the bodies of the vertebrae and of the intervertebral disks on each side lateral from the lig. longitudinale posterius; they consist of several large veins, running very close together, which anastomose manifoldly with one another, possess a lumen which stands open especially in the cervical and thoracic part and are accordingly also called the *sinus vertebrales longitudinales*. The *posterior* plexuses are situated on both sides on the anterior surface of the arcus vertebrarum and of the ligamenta flava and anastomose through spaces in the latter with the plexus (externi) posteriores; they are more delicate and less numerous and form more irregular, wider networks. The anterior plexuses are united with one another on the posterior surface of each vertebral body by transverse trunks, which run between the periosteum and the lig. longitudinale posterius and there receive the *vv. basivertebrales*. The latter (see also Fig. 545) extend radially in the vertebral body toward the middle of the posterior surface and open in common or separately, being provided at the mouth with valves; in front they anastomose through the bones with the plexus (externi) anteriores. The posterior plexuses are connected with one another by branches arranged in the form of a network; with the anterior they are connected by numerous shorter vessels which are present especially on the medial side of each radix arcus vertebrae and leave free only the upper part of the foramen intervertebrale. Through the connecting branches of the different plexuses with one another arise the more or less distinct vascular rings, *retia venosa vertebrarum*, one at the level of each vertebra. The upper end of the plexuses surrounds the foramen occipitale magnum like a wreath (see Fig. 535), is connected with the sinus occipitalis, plexus basilaris and rete canalis hypoglossi and empties into the vv. vertebrales. In the canalis sacralis the vessels become smaller, the plexuses become looser and empty through the foramina sacralia into the vv. sacrales.




545. Spinal veins in a horizontal section, viewed from above.

(Only the body of the vertebra has been cut through.)

Plexus venosi vertebrales externi (see also Fig. 544) lie on the external surface of the spinal column, have no valves and are divisible into anterior, smaller plexuses and posterior, larger plexuses. The *plexus venosi vertebrales anteriores* are best developed in the cervical spine and there lie upon the bodies of the vertebrae and upon the mm. longi capitis et colli; they receive tributaries from the muscles and from the bodies of the vertebrae and are connected lateralward with the vv. intervertebrales and with the following plexuses. The *plexus venosi vertebrales posteriores* lie in part directly upon the posterior surface of the vertebral arches, processes and ligaments, in part between the short muscles of the back. They are best developed in the cervical portion (see also Figs. 537 and 539) and are connected there especially with the vv. occipitales, vv. vertebrales and vv. cervicales profundae; besides, they anastomose through spaces in the ligamenta flava with the plexus interni and unite in front with the plexus (externi) anteriores, with the vv. intervertebrales and with the vv. intercostales.

Vv. intervertebrales accompany the nn. spinales in the foramina intervertebralia and lie chiefly in plexuses on their anterior surfaces. They receive the vv. spinales of the spinal cord, which bear valves at their openings, and are connected with the plexus vertebrales interni and externi representing their channels of outflow; they have valves where they terminate in the neck in the v. vertebralis, or in the trunk in the vv. intercostales (vide infra), lumbales or sacrales laterales.

The **vv. intercostales** (see Fig. 546), situated usually singly at the lower margin of a rib in the sulcus costae, close to and above the artery, increase evenly in diameter from above downward. The upper 9 (—10) form venous circles which empty their contents partly sternalward into the vv. mammaria interna and musculophrenicae (see p. 463), partly vertebralward into the system of the v. azygos; at their mouths they carry valves correspondingly directed; from their middle portions they send off branches in the 1st—6th (or 7th) intercostal space to the axilla (*vv. costoaxillares*, see p. 471). The lower 2—3 vv. intercostales have no outflow forward and receive tributaries, supplied with valves, from the diaphragm and from the abdominal muscles. Veins of smaller diameter and inconstant in length are to be found also at the upper margin of each rib; they are manifoldly connected with the others. Each v. intercostalis unites with a *ramus posterior* near the bodies of the vertebrae, which is larger than it is itself and which runs near the ramus posterior of the a. intercostalis; into this flow tributary vessels from the skin and from the muscles of the back, as well as the *ramus spinalis*, which, as a continuation of the vv. intervertebrales (vide supra), collects the blood from the spinal column and its contents.



546. Vena azygos, viewed from in front.

(The diaphragm, the right mm. intercostales interni and the lumbar muscles have been removed.)

V. azygos, unpaired, begins as a continuation of the *v. lumbalis ascendens dextra* (see p. 478) between the right *crus mediale* and *crus intermedium* of the diaphragm, ascends in the *spatium mediastinale posterius* on the right or anterior side of the vertebral bodies, and in its course lies in front of the *aa. intercostales dextrae*, to the right of the *aorta thoracalis* and of the *ductus thoracicus* and behind the *oesophagus*. At the level of the 5th or 4th thoracic vertebra it bends forward over the root of the right lung and opens into the *v. cava superior* (see also Figs. 478 and 680). In this latter portion of its course it usually possesses valves, which, however, are not efficient. It anastomoses sometimes at the beginning with the *v. renalis dextra*, receives the nine lowermost *vv. intercostales dextrae*, as well as the *vv. oesophageae*, *vv. bronchiales posteriores* and *vv. mediastinales posteriores* (from the corresponding organs) (not illustrated), besides the *v. hemiazygos*. The latter (see also Fig.

681) begins, unpaired, between the left *crus mediale* and *crus intermedium* of the diaphragm as the continuation of the *v. lumbalis ascendens sinistra*, but is in most cases connected with the *v. renalis sinistra*. Devoid of valves, it extends upward, behind the *aorta thoracalis*, on the left side of the bodies of the vertebrae, receives several *vv. intercostales sinistrae*, *vv. oesophageae* and *vv. mediastinales posteriores*, and bends to the right in front of one of the inferior or middle thoracic vertebrae behind the *ductus thoracicus* to go to the *v. azygos*. The intercostal veins, next higher up on the left side, empty into a *v. hemiazygos accessoria*, which goes downward on the left side of the thoracic spine and empties either into the *v. hemiazygos* or directly into the *v. azygos*. The 3—4 upper intercostal veins open on each side into a *v. intercostalis suprema*, which on the right side usually opens into the *v. azygos* (supplied with valves), on the left side into the *v. hemiazygos accessoria*, but in addition also communicates above on each side with the *v. anonyma* or with the *v. vertebralis*.

547.

Arteries and veins of a piece of small intestine.


The **vena portae** (*portal vein*) (see Fig. 548 and 549) collects the blood from the whole digestive tube, from the cardia on as far as the m. levator ani, from the pancreas, the gall bladder and the spleen. Its *trunk* arises behind the caput pancreatis, on the left and posterior surface of the pars superior duodeni, from the union of the v. mesenterica superior and the v. lienalis; it is 2–3 cm long and runs upward in the lig. hepatoduodenale, behind the a. hepatica and the hepatic and biliary ducts, to the porta hepatis; there it divides into a right and a left branch one for each half of the liver. Its *roots* follow, in general, the branches of the arteries of the same name and are supplied with valves only in their finer subdivisions immediately after their exit from the wall of the stomach and intestine.

1. **V. mesenterica superior** (see Figs. 548 and 549) runs on the right side of the a. mesenterica superior, arises from numerous *vv. intestinales* (from the small intestines), from the v. ileocolica, v. colicae dextrae, v. pancreaticoduodenalis, v. duodenales, v. pancreaticeae, v. colica media and v. gastroepiploica dextra. The intestinal veins anastomose with one another very much as the arteries do but more frequently.

2. **V. mesenterica inferior** (see Figs. 548 and 549) collects its blood from the v. haemorrhoidalis superior (see also p. 482), the vv. sigmoideae and the v. colica sinistra. Its peripheral course agrees with that of the a. mesenterica inferior; its trunk however separates from this and goes in a curved direction upward and to the right behind the peritoneum (frequently in the plica duodenojejunalis) and behind the pancreas; it usually opens into the v. lienalis, frequently into the v. mesenterica superior, sometimes at the point of junction of the two.


3. **V. lienalis** (see Figs. 548 and 549) is formed by the union of the veins of the spleen with the v. gastroepiploica sinistra and with the vv. gastricae breves; it runs from the left to the right beneath the a. lienalis and behind the pancreas, and in its course receives several pancreatic veins. It crosses the aorta between the a. coeliaca and the a. mesenterica superior.





548. Tributaries of the portal vein.

(The transverse colon and the small intestines have been partially removed; the liver has been drawn upward.)



549. Trunk of the portal vein.

(The stomach, liver and pancreas have been partially, the small intestines and the transverse colon completely, removed.)

4. **V. coronaria ventriculi** (see also Fig. 548) runs from the right to the left along the *curvatura minor* of the stomach, anastomoses near the pylorus with the *v. pylorica*, at the cardia with the oesophageal veins and opens near the *a. gastrica sinistra* into the trunk of the *v. portae* or the *v. lienalis*.

5. **V. pylorica** (see also Fig. 548), more delicate than the preceding, extends from the left to the right near the *a. gastrica* at the *curvatura minor* of the stomach, anastomoses with the preceding vein and opens at the pylorus into the stem of *v. portae*.

6. **V. cystica** (see Fig. 548) from the gall bladder, opens into the stem or the right branch of the portal vein.

The **accessory portal veins** (not illustrated) are small vessels, which arise in the folds of the peritoneum connecting with the liver, or which come especially from the stomach and pass either into the stem of the portal vein or directly into the substance of the liver.

The **anastomoses of the portal** with the *vv. cavae* are regularly present as follows:

- 1) At the cardia of the stomach between the *v. coronaria ventriculi* (*v. portae*) and the oesophageal veins (*v. azygos*);
- 2) at the rectum in the plexus haemorrhoidales between the *v. haemorrhoidalis superior* (*v. mesenterica inferior*) and the *vv. haemorrhoidales mediae et inferiores* (*vv. hypogastricae*). Valves usually prevent a flow back from the *v. hypogastrica* to the *v. portae*.
- 3) *vv. parumbilicales* [*Sappey*] (not illustrated), 4—5 small vessels, which are connected in the neighborhood of the navel with the superficial veins and with the *vv. epigastricae inferiores*, extend near the *lig. teres hepatis* to the liver and empty usually directly into its substance. The largest anastomoses also with the left *v. epigastrica superior*. One of these veins is said to be the remains of the *v. umbilicalis*. They are supplied with valves; the current is directed toward the liver.

V. cava inferior (see also Fig. 686) arises by the union of the vv. iliacae communes in front of the intervertebral disk between the 4th and 5th lumbar vertebra behind the first portion of the a. iliaca communis dextra somewhat to the right of the median plane. From there on, it ascends at first in front of the bodies of the lumbar vertebrae and in front of the origins of the right m. psoas major, then in front of the a. renalis dextra and in front of the right crura diaphragmatis and at the same time somewhat to the right and forward, goes through the foramen venae cavae of the diaphragm and, curved somewhat forward, opens into the right atrium of the heart. In its course it lies on the right side of the aorta abdominalis and is covered in front by the peritoneum, by the pars inferior duodeni and by the caput pancreatis; above, it is embedded in the fossa venae cavae of the liver and firmly fused with it, so that only its posterior wall lies partly free (see Fig. 632). It is devoid of valves. Its branches are divisible into *radices parietales* (from the walls of the peritoneal cavity) and *radices viscerales* (from the abdominal viscera).

A. Radices parietales;

1. **V. phrenica inferior**, paired, partly double, corresponds to the a. phrenica inferior and opens at the foramen venae cavae.

2. **Vv. lumbales** (see also Figs. 546 and 551), on each side 4—5, run close to and over the aa. lumbales, correspond to the vv. intercostales (see p. 473) and receive tributaries from the muscles and from the skin of the back, from the plexuses of the spine and from the abdominal muscles. They possess (insufficient) valves, run medianward upon the lateral and anterior surface of the bodies of the vertebrae (those of the left side behind the aorta) and open into the posterior wall of the v. cava. The vv. lumbales are united on each side with one another by means of longitudinal anastomoses which are situated behind the m. psoas major in front of the roots of the procc. transversi. The sum of these anastomoses forms on each side a vessel running vertically, the *v. lumbalis ascendens*, which is connected below with the v. iliolumbalis and frequently with the v. iliaca communis, above however is continued on the right side directly into the v. azygos, on the left side into the v. hemiazygos (see p. 474).

B. Radices viscerales:

1. **Vv. hepaticae** (see also Fig. 631), 2—3 large trunks, collect the blood which is led to the capillary system of the liver through the a. hepatica propria and v. portae; they leave the liver in the uppermost part of the fossa venae cavae and open into the v. cava just below the diaphragm. The right v. hepatica, largest, carries the blood from the right lobe of the liver, the middle, smallest, vein from the lobi caudatus [Spigeli] and quadratus, the left from the left lobe of the liver. Several small vv. hepaticae open in the fossa venae cavae, below the large ones, directly into the v. cava inferior. Valves are only rarely present at the mouths of the large veins.


2. **V. renalis** (renal veins) (see also Fig. 686), paired, arises by the fusion of from three to five branches near the hilus renalis, runs medianward and at the same time somewhat upward at the level of the 1st lumbar vertebra or a little deeper in front of the a. renalis and opens into the lateral wall of the v. cava. The left v. renalis is somewhat the larger and longer; it lies usually in front of the aorta just beneath the origin of the a. mesenterica superior. It receives tributaries from the capsula adiposa of the kidney and from the ureter; the left v. renalis receives also usually the v. spermatica sinistra. In addition the left vein anastomoses very frequently with the v. hemiazygos, the right sometimes with the v. azygos. The vv. renalis rarely possess valves at their mouths.

3. **V. suprarenalis** (suprarenal veins) (see also Fig. 692), paired, comes from the anterior surface of the adrenal and opens on the right side usually into the v. cava, on the left side into the v. renalis. It is devoid of valves.

4. **V. spermatica interna** (spermatic vein) (see also Figs. 551, 553 and 554), paired, accompanies the a. spermatica interna directly and is separated from it only at its termination. It consists of from 2 to 3 anastomosing vessels which surround the artery like a plexus, receives branches from the ureter, from the peritoneum and from the capsula adiposa of the kidney and unites finally to form a single trunk. The right vein opens obliquely into the anterior wall of the v. cava and there possesses efficient valves, the left goes almost at a right angle into the inferior wall of the v. renalis sinistra. The beginning of the vein is different in the two sexes.


In the male, the *v. testicularis*, begins at the posterior margin of the testicle and epididymis with numerous fine and few thick branches (see also Fig. 703), which form a close plexus (*plexus pampiniformis*) around the artery inside the funiculus spermaticus, and extends with this through the canalis inguinalis. It anastomoses extensively on the testicle and epididymis with the v. spermatica externa. It possesses valves outside the canalis inguinalis, in the abdominal cavity only one near the annulus inguinalis abdominalis.

In the female, *v. ovarica*, it arises partly from numerous vessels, which leave the ovary at its hilus, partly from broad anastomoses with the upper part of the plexus uterovaginalis; in addition it receives some branches from the lig. teres uteri and from the tuba uterina. All these branches, anastomosing, form a close plexus (*plexus pampiniformis*) around the artery, inside the lig. suspensorium ovarii and are devoid of valves.




550. Vena cava inferior in the male, viewed from in front.

(On the right side the kidney has been removed, a piece of the m. psoas and of the a. iliaca communis has been cut out; on the left side the abdominal wall has been reflected downward.)



551. Veins of the male pelvis, right half, viewed from the left.

(The left m. psoas major has been removed, the rectum has been drawn somewhat to the side.)




552. Veins of the female perineum (see Fig. 525).

V. iliaca communis (*common iliac vein*) (see Figs. 550 and 558), paired, devoid of valves, behind and to the right of artery, arises from the *v. hypogastrica* and the *v. iliaca externa*. It frequently anastomoses with the *v. lumbalis ascendens*. Branch of the left vein:

V. sacralis media (middle sacral vein), unpaired, often double below, near the *a. sacralis media*, forms with branches of the *vv. sacrales laterales* the *plexus sacralis anterior* (see p. 484); it is connected with the *plexus haemorrhoidalis externus* and *vesicalis*.

V. hypogastrica (*hypogastric vein*) (see Figs. 550 and 551) lies directly behind the *a. hypogastrica*. Its branches possess valves only outside the pelvis. Branches:

1. **V. pudenda interna** (*O. T. internal pudic vein*) (see also Figs. 551 and 554) begins below the symphysis ossium pubis with an anastomosis from the *v. dorsalis penis* (*clitoridis*) and receives the *vv. profundae penis* (*clitoridis*), *vv. urethrales*, *vv. bulbi urethrae* (*bulbi vestibuli* [*vaginae*]), *vv. scrotales* (*labiales*) *posteriores* and *vv. haemorrhoidales inferiores*; the latter are connected with the *plexus haemorrhoidales*. All the branches correspond to the arteries of the same name. The vein usually runs first beside the *a. penis* (*clitoridis*) in the *trigonum urogenitale* and then (for the most part double) accompanies the *a. pudenda interna*.



553. Veins of the female pelvic organs, viewed from the left.


(The superficial layers of the venous plexuses [see Fig. 554] have been removed; the peritoneum has been for the most part removed.)

2. **Plexus haemorrhoidalis internus** (see Figs. 552 and 627) in the tela submucosa of the rectum, with ampulla-like dilatations in the region of the annulus haemorrhoidalis (see also p. 547) empties partly into a delicate, superficial network at the anus, partly by means of numerous perforating branches into the **plexus haemorrhoidalis externus** (see also Fig. 551) situated upon the external surface of the tunica muscularis. From the latter arise the *v. haemorrhoidalis superior*, devoid of valves (to the *v. mesenterica inferior* of the *v. portae*), the *vv. haemorrhoidales mediae*, with valves, paired (to the *v. hypogastrica*), which receive branches from the bladder, prostata and seminal vesicles in the male (uterus and vagina in the female) and the *vv. haemorrhoidales inferiores*, with valves, paired, from the neighborhood of the anus (to the *v. pudenda interna*); in addition branches go to the *v. sacralis media*.

3. **Plexus vesicalis** (see also Figs. 551, 554 and 730) surrounds the lower part of the bladder and in the male the basis prostatae lateralward and behind; it is, with valves in almost all its branches, connected with the following plexus, receives *vv. deferentiales* (from the testicle, along the ductus deferens) and is emptied on each side by means of several *vv. vesicales*.

4. **Plexus pudendalis** (see also Figs. 551 and 554) lies behind and over the lig. arcuatum pubis, in front of the prostata and bladder (in the male, the bladder and urethra in the female), receives the main trunk of the unpaired *v. dorsalis penis (clitoridis)* (see also Fig. 709), which runs between the lig. arcuatum pubis and the lig. transversum pelvis, is connected with the *v. pudenda interna* and with the plexus vesicalis and empties partly into the *vv. vesicales*, partly directly into the *v. hypogastrica*.

5. **Plexus uterovaginalis** (in the female) extends over the walls of the vagina and as an especially close plexus along the margo lateralis of the uterus, where it partly fills up the parametrium and surrounds the ureter. It is connected with the plexus pudendalis and haemorrhoidalis externus and empties on each side chiefly through the large *vv. uterinae*, which surround the *a. uterina* and form broad anastomoses with the plexus pampiniformis of the *v. ovarica*. There are almost no valves in the uterine veins, and those which are found can not be closed.



**554. Veins of the female pelvis, right half,
viewed from the left and somewhat in front.**

(The ovaries and uterus have been drawn out of their position; the peritoneum has been largely removed; the ureters have been removed.)

6. **V. iliolumbalis** (see Figs. 546, 550 and 551), in part double, accompanies the a. iliolumbalis and often opens into the v. iliaca communis. It receives the vv. intervertebrales (see p. 473), muscular veins and sometimes the v. lumbalis V, anastomoses with the vv. lumbalis ascendens, circumflexa ilium profunda and sacrales laterales and possesses many valves.

7. **Vv. sacrales laterales** (see Fig. 550), usually double, run near the a. sacralis lateralis and open into the v. hypogastrica or into a branch of the same. Their branches form, with those of the v. sacralis media, the *plexus sacralis anterior* upon the facies pelvina of the sacrum.

8. **V. obturatoria** (see Figs. 551 and 554) near the a. obturatoria; its branches are usually doubled; it forms a broad anastomosis in the pelvis with the v. iliaca externa, and sometimes opens into this. Its branches anastomose freely with the v. circumflexa femoris medialis.

9. **V. glutaea superior** (see Fig. 557) accompanies the a. glutaea superior; its branches are usually doubled. At its entrance into the pelvis its walls stand open and it possesses valves. Behind the m. piriformis it anastomoses with the v. glutaea inferior.

10. **V. glutaea inferior** (see Fig. 557), near the a. glutaea inferior; its branches, usually doubled, accompany the branches of the latter. It receives a large branch from the n. ischiadicus and anastomoses on the m. quadratus femoris by means of very wide vessels with the v. circumflexa femoris medialis and with the uppermost v. perforans of the vv. profundae femoris.

V. iliaca externa (*external iliac vein*) (see Figs. 550, 551 and 556) begins as the continuation of the v. femoralis behind the lig. inguinale, on the medial side of the a. iliaca externa and, as it ascends, becomes displaced behind the artery. It receives a broad anastomosis from the v. obturatoria and frequently possesses one (usually insufficient) valve. Branches:

1. **V. epigastrica inferior** (see Figs. 550, 551 and 556), usually doubled, accompanies the a. epigastrica inferior. It anastomoses with the vv. epigastricae superiores and behind the ramus superior ossis pubis with the v. obturatoria; in addition it receives the vv. parumbilicales, as well as the v. spermatica externa (see also p. 478) from the testicle and epididymis and branches from the vv. cutaneae abdominis. Below the navel it possesses numerous valves.

2. **V. circumflexa ilium profunda** (see Figs. 550, 551, 554 and 556), double in most of its course, surrounds the a. circumflexa ilium profunda like a plexus and often opens into the v. epigastrica inferior; in so doing it crosses the a. iliaca externa on its anterior or posterior surface. It anastomoses with the v. iliolumbalis and is supplied with valves.

V. femoralis (*femoral vein*) (see Fig. 556) is the continuation of the v. poplitea from the hiatus tendineus [adductorius] on. In the canalis adductorius [Hunteri] it lies behind the a. femoralis and somewhat lateralward from it; further up it runs for the most part exactly behind the artery, and arrives finally on the medial side of the artery in the fossa iliopectinea. In its lower portion two vv. comitantes also run close to the artery. It possesses usually 4–5 pairs of valves. Branches:

1. **V. epigastrica superficialis** (see also Fig. 559) is situated in the subcutaneous tissue and carries the blood downward from the lower half of the anterior abdominal wall. It opens in the fossa ovalis separately or, united with neighboring veins, into the v. femoralis or into the v. saphena magna and possesses valves which conduct downward.

2. **V. circumflexa ilium superficialis** (see Fig. 559), sometimes doubled for part of its course, close to the corresponding artery in the subcutaneous tissue, opening either independently or in common with the v. epigastrica superficialis. It is supplied with valves.


3. **V. thoracoepigastrica**, single or doubled for long distances, extends, on the anterior or lateral wall of the trunk, in the subcutaneous tissue, from the axilla as far as the fossa ovalis. Its middle portion is devoid of valves, in its upper part valves direct the blood upward to the axilla, in its lower part downward to the fossa ovalis. Above, the vein opens into the v. thoracalis lateralis (see p. 471), below it goes either separately or united with the v. epigastrica superficialis into the v. femoralis.

4. **Vv. pudendae externae** (see also Fig. 559), near the aa. pudendae externae, frequently flow into the v. saphena magna; they receive branches from the skin of the abdomen, from the scrotum (or the labia majora), vv. scrotales (labiales) anteriores, and from the penis (or the clitoris), v. dorsalis penis (clitoridis) cutanea; the latter runs unpaired or doubled upon the fascia penis (clitoridis) as far as the symphysis pubis and there bends lateralward.

(Continuation see p. 486.)


Vv. cutaneae abdominis et pectoris form, in the subcutaneous tissue of the abdominal and thoracic wall, a network of vessels, supplied with valves, from which the blood is led away on each side 1st by superficial vessels to the neck, to the axilla (v. thoracoepigastrica) and to the fossa ovalis (vide supra), 2nd by deep, perforating vessels to the thoracic cavity (to the vv. mammaria interna and intercostales), to the peritoneal cavity (vv. parumbilicales) and to the veins of the anterior abdominal wall (vv. epigastricae superiores et inferiores).

In the neighborhood of the papilla mammae the vv. cutaneae form a wreath of vessels of uneven caliber, the *plexus venosus mamillae*.



555. Superficial veins of the trunk,
viewed from in front (after W. Braune).

(The arrows indicate the way in which the current is directed by the valves.)



556. V. femoralis dextra, viewed from in front.

(The lig. inguinale and the mm. sartorius, rectus femoris, pectineus and adductor longus have been entirely or partially removed; a piece has been cut out of the a. femoralis.)

V. femoralis (continued):

5. V. saphena magna (see p. 489).

6. V. profunda femoris accompanies the a. profunda femoris and possesses several pairs of valves. Branches:

- a) *Vv. perforantes* (see also Fig. 557), partly doubled, arise on the posterior surface of the m. adductor magnus and communicate freely there by longitudinal anastomoses with one another, with the vv. gluteae inferior, circumflexa femoris medialis and with the v. poplitea; they then perforate the mm. adductores near the aa. perforantes.
- b) *Vv. circumflexae femoris mediales* (see also Fig. 557) run near the corresponding arteries and anastomose freely behind, on the m. quadratus femoris, with the vv. gluteae inferior and perforans I, in front with the v. obturatoria. They are supplied with valves.
- c) *Vv. circumflexae femoris laterales*, supplied with valves, accompany the a. circumflexa femoris lateralis and anastomose with the preceding and with the v. gluteae inferior.

V. poplitea (*popliteal vein*) (see Figs. 557 and 895) arises from the union of the vv. tibiales and runs just behind the a. poplitea, in its distal part at the same time somewhat medianward, in its proximal part somewhat lateralward from it. Besides, near the artery there is below a v. comitans lateralis, above a v. comitans medialis. It is supplied with valves, receives the v. saphena parva (see p. 490) and the vv. articulares genus (double near the corresponding arteries, not illustrated) and anastomoses with the lower vv. perforantes.


557. Deep veins of the right thigh, viewed from behind.

(The *m. gluteus maximus* has been reflected lateralward; otherwise like Fig. 528.)

The deep veins of the foot and leg (not illustrated) accompany the arteries, usually two to each artery, and are plentifully supplied with valves. Numerous anastomoses with the superficial veins lead the blood in the back of the foot from the depth to the surface, in the sole of the foot, in the leg and in the thigh from the surface to the depth.

In the sole of the foot they begin as the *vv. digitales plantares* and then form the *vv. metatarsae plantares*, which empty partly through several *rami perforantes* into the deep and the superficial veins of the back of the foot, partly into the *arcus venosus plantaris profundus*. From the latter go off the *vv. plantares laterales*, which by union with the finer *vv. plantares mediales* form the *vv. tibiales posteriores*; in the leg the latter receive the *vv. peronaeae*. The *vv. plantares laterales* also send an anastomosis to the *v. saphena parva*, the *vv. plantares mediales* to the *v. saphena magna*.

On the back of the foot the deep veins are relatively delicate; they begin as the *vv. metatarsae dorsales pedis* and are continued into the *vv. tibiales anteriores*.



558. Superficial veins of the right leg,


viewed from in front (after W. Braune.)

(The arrows indicate the way the stream is directed by the valves.)

The **superficial veins of the leg** form a continuous network in the subcutaneous tissue, varying in complexity, rich in valves, and permitting the recognition in it of single larger vessels which act as the principal canals of exit. They are connected with the deep veins by numerous anastomoses (for direction of current see p. 487).

In the **sole of the foot** (not illustrated) there is a very close network (*rete venosum plantare cutaneum*), which possesses different channels of outflow. In the first place branches go to the deep veins, secondly, however, and chiefly, the blood is led away through superficial trunks around the borders of the foot to the dorsum of the foot. These pass at the medial and lateral margins of the foot into the vv. marginales medialis and lateralis of the dorsum of the foot, dorsalward over the margin of the heel toward the leg, and forward into a larger vessel (*arcus venosus plantaris cutaneus*), which runs in the toe-ball furrow at the distal margin of the fasciculi transversi aponeurosis plantaris and is continued over the medial and lateral margins of the foot into the beginnings of the vv. marginales (vide infra). This vascular arch receives superficial branches from the toes and sends off into the space between each two toes single or double vv. intercapitulares to the vv. digitales dorsales pedis.

On the **back of the foot** the veins begin at the toes as the vv. digitales pedis dorsales, which unite after receiving the vv. intercapitulares to form the short vv. digitales communes pedis and flow into the *arcus venosus dorsalis pedis [cutaneus]*; this is an irregular chain of anastomoses between the vv. digitales communes and the vv. marginales and lies upon the distal portion of the ossametatarsalia. Proximalward from it lies the irregularly shaped *rete venosum dorsale pedis cutaneum*, which receives many tributaries from the depth and is continued into the anterior network of the leg. In the network, near the lateral and medial margins of the foot, can be made out on each side a larger vessel, the v. marginalis lateralis (see Fig. 560) and the v. marginalis medialis, which chiefly receive tributaries from the rete venosum plantare cutaneum.





559. Superficial veins of the right thigh, viewed from in front.

(The arrows indicate the way the current is directed by the valves.)

In the plexus of the superficial veins of the leg two large venous trunks stand out above the foot, the *v. saphena magna* and the *v. saphena parva*, of which the latter belongs essentially to the leg alone.

V. saphena magna (see also Figs. 556 and 558) begins in front of the medial malleolus as the continuation of the *v. marginalis medialis* and there receives superficial branches from the rete venosum plantare cutaneum and deep branches from the *vv. plantares mediales*. It then runs almost vertically upward upon the *facies medialis tibiae*, makes a slight curve behind the *condyli mediales* of the tibia and femur and then goes fairly straight but obliquely upward on the medial and anterior surface of the thigh, and enters the anterior medial wall of the *v. femoralis* in the *fossa ovalis*. In its entire course it is situated subcutaneously upon the *fascia cruris* and *fascia lata*. On its way it receives numerous branches from the superficial networks of the heel, of the leg and of the thigh, as well as veins from the tibia, and anastomoses manifoldly with the *v. saphena parva*. The superficial veins coming from the back of the thigh often form a separate trunk, the *v. saphena accessoria* (not illustrated), which accompanies the *v. saphena magna* medianward at a variable distance from it and empties into it at an inconstant level.





560. Superficial veins of the right leg,

viewed from behind (after
W. Braune).

(The arrows indicate the way the current is directed by the valves.)

V. saphena parva (see also Fig. 557) begins behind the lateral malleolus as the continuation of the *v. marginalis lateralis* of the foot; in its formation there takes part, besides superficial branches of the *rete venosum plantare cutaneum*, also an anastomosis from the deep *vv. plantares laterales*. It runs at first on the lateral margin, then on the posterior surface of the *tendo calcaneus* [Achillis]; farther up ensheathed in a duplicature of the superficial layer of the *fascia cruris*, it lies in the groove between the two heads of the *m. gastrocnemius* (see also Fig. 880). The mouth of the *v. saphena parva* arises sometimes in the leg in the *v. saphena magna* or in deep veins; most frequently it embouches undivided into the hollow of the knee in the *v. poplitea*, sometimes also divided into the latter as well as into the *v. saphena magna* or into deeper thigh veins; sometimes the mouth arises above the hollow of the knee into deep, posterior veins or the *v. saphena magna*. The *v. femoropoplitea* runs as a small vessel superficially along the thigh and empties its contents downward into the *v. poplitea* and into deep thigh veins or only into the latter, upward under the *m. glutaeus maximus* into the *vv. glutaeae*. The *v. saphena parva* receives in its course numerous branches from the superficial network of the heel and of the leg and anastomoses usually manifoldly with the *v. saphena magna*.


The fetal circulation.

During embryonic life the apparatus concerned in the circulation of the blood undergoes various transformations.

The arrangement which exists in the second half of pregnancy up to birth is designated the **placental circulation**. This presents the following peculiarities (see Figs. 561 and 562).

The **v. umbilicalis** (*umbilical vein*), devoid of valves, leads the blood, which has become *arterial* in the placenta, through the *umbilical cord* (*junculus umbilicalis*) to the navel and thence in the free margin of the lig. falciforme hepatis to the fossa venae umbilicalis of the liver. There a part of the blood goes through direct branches to the capillary system of the liver; another part becomes mixed, in that the v. umbilicalis connects with the left branch of the v. portae, with the venous contents of the portal vein (from the stomach, intestines, pancreas and spleen) and also arrives, along with this, in the capillary system of the liver; finally the third, smaller part runs directly to the left v. hepatica and thus to the v. cava inferior through the *ductus venosus* [*Arantii*], which is situated in the fossa ductus venosi of the liver (see also Fig. 632) and connects the v. umbilicalis with the left v. hepatica. The v. cava inferior carries mixed blood therefore on its passage through the foramen venae cavae of the diaphragm. This is composed of 1st the pure arterial contents of the ductus venosus [*Arantii*], 2nd the contents of the vv. hepaticae, which carry the blood away from the capillary system of the liver, and 3rd the venous contents of the v. cava inferior (especially from the renal and suprarenal veins, pelvic veins and those of the lower extremities). Passing through the v. cava inferior, this blood reaches the right atrium, which, in the fetus, is freely connected with the left atrium through the foramen ovale (see Fig. 474). Since the foramen ovale is situated in the direction of the current of blood arriving through the v. cava inferior, and since, in addition, the valvula venae cavae [*inferioris*, *Eustachii*] to a certain extent hinders an alteration of this direction of the current, the main mass of blood streams through the foramen ovale into the left atrium and in so doing receives and mixes with only a part of the venous blood which arrives in the heart through the v. cava superior. Into the left atrium open the vv. pulmonales, and their blood, which in the fetus is venous, mixes with the other contents. The so-called *mixed arterial* blood thus arising goes especially to the upper half of the body through the left ventricle, the aorta ascendens and the arcus aortae and its large branches, the aa. anonyma, carotis sinistra and subclavia sinistra; only a part flows into the aorta thoracalis.


The v. cava superior carries the venous blood from the upper half of the body and (through the v. azygos which is markedly developed in the fetus) from the wall of the trunk, to the right atrium. Thence a small part passes through the foramen ovale into the left atrium, the larger part, however, mixed with a part of the blood from the v. cava inferior and hence called *mixed venous* blood, enters the right ventricle and passes into the a. pulmonalis. The a. pulmonalis divides, in the fetus, into three branches: the rami dexter et sinister, which go to the lungs (as yet not functioning) and the *ductus arteriosus* [*Botalli*], which runs in the direction of the main trunk and enters, at the end of the arcus aortae, into the concave anterior wall of the latter. The greater part of the mixed venous blood now runs through this ductus arteriosus as far as the beginning of the aorta thoracalis; there it mixes with the residue of the mixed arterial blood coming from the end of the arcus aortae and passes through the branches of the aorta thoracalis and aorta abdominalis to the intestinal canal, to the genito-urinary organs, to the inferior portion of the wall of the trunk and to the lower extremities, chiefly, however, through the two aa. umbilicales (see p. 446) into the umbilical cord and to the placenta, where the blood again becomes arterialized.



561. Bloodvessels of the fetus,

viewed from in front and somewhat

from the left. (The abdominal and thoracic cavities have been widely opened, the lungs, the whole alimentary canal, the pancreas, the spleen, the left lobe of the liver and the serous membranes have been removed and the heart has been drawn aside to the right.)



562. Scheme of the circulation of the blood in the fetus,

viewed from in front and somewhat from the left (see Fig. 561).

(The heart is in its correct position and has been partly opened. The arrows indicate the direction of the blood current.)

Printed by Fischer & Wittig, Leipzig.







