Annual report of the Medical Department / Federated Malay States.

Contributors

Federated Malay States. Medical Department.

Publication/Creation

Kuala Lumpur : Government Press, [1926]

Persistent URL

https://wellcomecollection.org/works/ccw5wwvg

License and attribution

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



FEDERATED MALAY STATES.

ANNUAL REPORT

OF THE

MEDICAL DEPARTMENT

FOR THE YEAR

1926



BY

Dr. R. DOWDEN.

Principal Medical Officer, F.M.S.

KUALA LUMPUR:

PRINTED AT THE FEDERATED MALAY STATES GOVERNMENT PRESS.

1927.

Digitized by the Internet Archive in 2019 with funding from Wellcome Library

CONTENTS.

THE RESIDENCE OF THE PARTY OF T		
ADMINISTRATION.—		PAGE
STAFF		1
Financial		8
PUBLIC HEALTH.—		
(a) General Remarks		3
(b) Communicable Diseases	THE THE	4
(c) VITAL STATISTICS		7
THE SHALKS AT BROKEN O VEHICLE AND THE	MAIN MAIN	8
SANITATION		9
INSPECTION OF SCHOOLS		9
PUBLIC HEALTH EDUCATION		9
QUARANTINE AND PORT HEALTH WORK	TRIE TA	10
INFANT WELFARE WORK	***	10
HOSPITALS, DISPENSARIES AND VENEREAL DISEASE CL	INICS	11
CURE FOR OPIUM HABIT		12
PRISONS		14
LEPER ASYLUMS	***	14
TEMPORARY CHINESE DECREPIT ASYLUM, PORT SWETTE	NHAM	14
VETERINARY BRANCH		15
METEOROLOGY		16
SCIENTIFIC.—		
COMMENTS ON THE REPORT OF THE DIRECTOR, INSTITUTE FO		3.0
Research	*** ***	
COMMENTS ON THE REPORT OF THE MEDICAL SUPERINTENDENT MENTAL HOSPITAL	r, Central	970
TABLE 1.—RETURN OF DISEASES AND DEATHS (IN-P	ATIENTS)	19
TABLE 2.—MEDICAL STAFF		29
ANNUAL REPORT OF THE DIRECTOR, INSTITUTE FOR RESEARCH, FEDERATED MALAY STATES		
ANNUAL REPORT OF THE MALARIA RESEARCH LAB INSTITUTE FOR MEDICAL RESEARCH		
ANNUAL REPORT OF THE CHEMICAL LABORATORIES, I FOR MEDICAL RESEARCH		
ANNUAL REPORT OF THE CHIEF HEALTH OFFICER		66
ANNUAL REPORT OF THE REGISTRAR-GENERAL OF BIR		
DEATHS, FEDERATED MALAY STATES		
ANNUAL REPORT OF THE CHIEF SURGEON, PERAK		
ANNUAL REPORT OF THE CHIEF SURGEON, SELANGOR		86

	PAGE
ANNUAL REPORT OF THE MEDICAL SUPERINTENDENT, CENTRAL MENTAL HOSPITAL, TANJONG RAMBUTAN	
ANNUAL REPORT OF THE SPECIALIST, VENEREAL DISEASES FEDERATED MALAY STATES	
ANNUAL REPORT OF THE OPHTHALMOLOGIST SURGEON, IPOH	. 125
ANNUAL REPORT OF THE DEPUTY MEDICAL OFFICER IN CHARGI	
ANNUAL REPORT OF THE RADIOLOGIST, FEDERATED MALAY STATES	
ANNUAL REPORT OF THE MEDICAL OFFICER IN CHARGE OF LEPER ASYLUM, KUALA LUMPUR	
ANNUAL REPORT OF THE LADY MEDICAL OFFICER, INFAN' WELFARE CENTRE, KUALA LUMPUR	
ANNUAL REPORT OF THE LADY MEDICAL OFFICER, INFANT WELFARE CENTRE, IPOH	
ANNUAL REPORT OF THE LADY MEDICAL OFFICER, INFAN. WELFARE CENTRE, TAIPING	

FEDERATED MALAY STATES.

ANNUAL REPORT OF THE MEDICAL DEPARTMENT FOR-THE YEAR ENDING 31st DECEMBER, 1926.

ADMINISTRATION.

STAFF.

 The total authorized staff of the Medical Department, Federated Malay States, including all branches on the 31st December, 1926, was 1,049. It is divided as follows:

				Total		1,049
Radiological Branch	41 - 255		1.335	***		6
Venereal Disease Branch		***				5
Veterinary Branch	***	***				82
Central Mental Hospital			***		***	34
Institute for Medical Research	ch			***		37
Health Branch						82
Hospital Branch			***		***	853

The following were the principal changes which took place during the year:

Dr. R. Dowden, Principal Medical Officer, proceeded on leave on 10th April, 1926, and Dr. A. R. Wellington, Senior Health Officer, acted for him until 14th November, 1926, when he went to Australia to attend the Conference on Health Conditions in the Pacific.

Dr. A. K. Cosgrave, Senior Medical Officer, Selangor, acted as Senior Health Officer from 10th April, 1926, until the end of the year.

Dr. W. H. Hart, Acting Senior Medical Officer, Pahang, was transferred to Selangor and acted as Senior Medical Officer, Selangor, from 10th April, 1926, until the end of the year.

Dr. H. R. Dive, Medical Officer, acted as Senior Medical Officer, Pahang, from 10th April, 1926, until the end of the year.

Dr. William Fletcher, Bacteriologist, proceeded on leave on 21st January, 1926, and Dr. A. N. Kingsbury, Pathologist, acted for him until 7th June, 1926.

Dr. A. T. Stanton, Director of Government Laboratories, proceeded on one month's leave on 2nd July, 1926, and was appointed Chief Medical Adviser to the Colonial Office, with effect from 11th August, 1926, and Dr. Fletcher, Bacteriologist, acted for him during the period and was appointed Director of Government Laboratories on 11th August, 1926.

APPOINTMENTS.

During the year, the following officers were appointed:

Dr. W. H. Hart as Senior Medical Officer, Pahang, on 31st August, 1926.

Dr. R. M. Dannatt as Acting Chief Surgeon, Selangor, on 5th March, 1926.

Dr. G. D. Gordon as Medical Officer on 5th March, 1926.

Dr. E. C. Chitty as Medical Officer on 6th March, 1926.

Dr. J. J. O'Grady as Medical Officer on 11th March, 1926.

Dr. F. G. Greenwood as Medical Officer on 25th March, 1926.

Dr. D. W. G. Faris as Medical Officer on 25th March, 1926.

Dr. E. A. Struthers as Medical Officer on 25th March, 1926.

Dr. P. J. Mulcahy as Medical Officer on 25th March, 1926.

Dr. J. G. Castellain as Health Officer on 17th May, 1926.

Mr. K. D. Turner as Veterinary Surgeon on 24th April, 1926.

Dr. (Miss) Ivy Collier as Lady Medical Officer on 22nd May, 1926.

Dr. R. T. B. Green was appointed to act as Assistant Bacteriologist on 27th March, 1926.

Dr. A. N. Kingsbury was appointed Professor of Bacteriology, College of Medicine, Singapore, on 30th May, 1926.

Capt. K. B. Williamson was appointed Professor of Biology, College of Medicine, Singapore, on 24th September, 1926.

Dr. C. C. Taffs as Medical Officer on 13th August, 1926.

Dr. T. C. Wakefield as Medical Officer on 14th August, 1926.

Dr. B. D. Merrin as Medical Officer on 14th August, 1926.

Mr. B. A. R. Gater of the Agricultural Department was appointed as Malaria Research Officer on 1st November, 1926.

Dr. (Mrs.) Whyte was appointed as Lady Medical Officer on 1st December, 1926.

RESIGNATIONS AND RETIREMENTS.

Dr. W. F. Macdonald retired on 21st April, 1926, on medical grounds.

Dr. B. Cross engagement expired on 25th May, 1926.

Dr. V. M. Matthews engagement expired on 3rd July, 1926.

Dr. (Miss) Ivy Collier retired on 20th November, 1926, on medical grounds.

Dr. P. J. Mulcahy resigned on 27th December, 1926.

Dr. I. P. Masters, Senior Medical Officer, Pahang, retired on medical grounds on 30th August, 1926.

Capt. F. Golding, Financial Secretary to the Senior Medical Officer, Perak, was granted eight months' leave on 10th October, 1926, prior to retirement.

DEATH.

I regret to record with deep regret the death of Dr. J. G. Dunlea, Medical Officer, on 14th September, 1926.

ASSISTANT SURGEONS.

APPOINTMENTS.

The following Assistant Surgeons were appointed during 1926:

Mr. J. Samuel, Assistant Surgeon, on 9th April, 1926.

Mr. Thiam Hock Wee, Assistant Surgeon, on 1st June, 1926.

Mr. K. Natarajan, Assistant Surgeon, on 1st July, 1926.

Mr. V. M. Balakrishna Pannikar, Assistant Surgeon, on 11th July, 1926.

Mr. G. S. Venkatesan, Assistant Surgeon, on 11th July, 1926.

Mr. Ooi Kheng Seng, Assistant Surgeon, on 15th September, 1926.

Mr. P. Kumara Menon, Assistant Surgeon, on 29th October, 1926.

PROMOTIONS.

The following were promoted to Deputy Medical Officers during 1926:

Mr. G. Abraham as Deputy Medical Officer on 1st November, 1926.

Mr. E. J. de Cruz as Deputy Medical Officer on 1st November, 1926.

Mr. K. N. Ghosh as Deputy Medical Officer on 1st November, 1926.

Mr. M. Gupta as Deputy Medical Officer on 1st November, 1926.

Mr. A. Ponniah as Deputy Medical Officer on 1st November, 1926.

Mr. S. Danasamy as Deputy Medical Officer on 1st November, 1926.

Mr. A. Saravanamuthu as Deputy Medical Officer on 1st November, 1926.

Mr. R. Vythilingam as Deputy Medical Officer on 1st November, 1926.

Mr. Abdul Latiff as Deputy Medical Officer on 1st November, 1926.

Mr. V. Supramaniam as Deputy Medical Officer on 1st November, 1926.

Mr. A. E. Doraisamy as Deputy Medical Officer on 1st November, 1926.

Mr. A. Viswalingam as Deputy Medical Officer on 1st November, 1926.

RESIGNATION.

Mr. P. Nadarajah, Assistant Surgeon, resigned on 17th June, 1926.

NURSING SISTERS.

APPOINTMENTS.

- Miss E. Augarde, Nursing Sister, on 19th June, 1926.
- Miss E. G. Hill, Nursing Sister, on 10th July, 1926.
- Miss D. Matthews, Nursing Sister, on 16th July, 1926.
- Miss M. T. O'Sullivan, Nursing Sister, on 29th July, 1926.
- Miss D. M. Hansford, Nursing Sister, on 7th August, 1926.
- Miss M. C. Cox, Nursing Sister, on 26th August, 1926.
- Miss M. J. Teale, Nursing Sister, on 26th August, 1926.
- Miss M. Duncan, Nursing Sister, on 25th September, 1926.
- Miss C. M. Saunders, Nursing Sister, on 7th October, 1926.
- Miss M. Grice, Nursing Sister, on 16th October, 1926.
- Mrs. C. Wilson, Nursing Sister, on 16th October, 1926.
- Miss R. E. Yeomans, Nursing Sister, on 16th December, 1926.
- Miss F. L. Webb, Nursing Sister, on 31st December, 1926.

PROMOTIONS.

Miss M. E. Fisher, promoted to Matron, Grade II, on 23rd December, 1926.

RETIREMENT.

Miss L. M. Jacobs, Matron, Grade II, retired on pension on 23rd December, 1926.

RESIGNATIONS.

- Miss E. H. McIlrath, Nursing Sister, engagement expired on 6th January, 1926.
- Miss S. Smith, Nursing Sister, engagement expired on 17th April, 1926.
- Miss E. Lampard, Nursing Sister, resigned on 18th May, 1926.
- Miss E. M. McDougall, Nursing Sister, resigned on 10th June, 1926.
- Miss D. M. Hodgson, Nursing Sister, resigned on 19th June, 1926.
- Miss M. Kaylor, Nursing Sister, resigned on 24th November, 1926.
- Miss E. L. Crocker, Nursing Sister, resigned on 30th November, 1926.
- 2. The Health and Hospital Branches were both understaffed by some fifty per cent. during the year. Taking this fact into consideration the staff are to be congratulated on the amount of work done, and as the duties of the department have increased in every direction, and 1926 was an exceptionally unhealthy year, the work of the department was carried out under great difficulties.

It is hoped that the new appointments in the Health Branch and the special allowance to those who hold the Diploma in Public Health qualification, together with the increased initial pay in the Medical Branch, a greater number of recruits will be forthcoming in 1927.

ENACTMENTS.

3. List of Enactments affecting Public Health enacted during the year:

"The Health Boards Enactment, 1926," was passed during the year and comes into force on 1st January, 1927. This Bill, which is the outcome of the Estates Health Commission, is an attempt, made for the first time, to organise the health work on large and small estates on a sound basis. It ought to result in a higher standard of both preventive and curative work and better estate hospitals.

FINANCIAL.

- (a) Statement of Revenue for the year 1926-
 - Revenue (Hospital fees, licences, etc.) ... \$ 374,080
- (b) Statement of Expenditure for the year 1926-
 - Personal Emoluments and Other Charges ... 4,115,518
- (c) The Estimated Revenue for the Federated Malay
 - States for the year 1926 is 81,043,259

PUBLIC HEALTH. (a).—GENERAL REMARKS.

4. The general health of the country was not satisfactory during the year under review, the death-rate being 29.22 per mille as against 23.60 for the previous year. This was due to a very great extent to the epidemic of malaria which swept over the country during the months of April, May, June, July and August. In some districts it was much more severe than in others. An increased virulence in the type of the disease was observed. Exceptionally heavy rains at the end of 1925 followed by a prolonged drought, and the increased amoust of clearing jungle land which followed the rise in the price of rubber may have been important contributing factors.

Various theories have been advanced by Government and privately employed medical men to account for this.

One group holds that waves of virulent malaria are to be expected every ten years or so, but this is also more or less true of exceptional rains. Others consider that the great increase in the number of imported Tamil labourers from India introduced a great number of and a more virulent type of parasites. The majority of the cases were P. falcparum and P. vivax infections, and no differences in the forms seen under the microscope are reported. It seems probable that the main factors were the floods and heavy rains, the silting of the water courses throughout the country which prevents flood water flowing off, and causes a rise in the sub-soil water level which lasts for a long period. The sub-soil water forming many new seepage areas and breeding places, while the increase in the Indian labouring population may have increased the supply of gametocytes. However the incidence of malaria was universal throughout the Federated Malay States and occurred in many places where there had been no influx of new labourers.

- 6. None of these theories are capable of absolute proof but many new breeding places were found in Kuala Lumpur by the Health Officer, Dr. Black and his staff, after the rains, and it is not unreasonable to suppose that there were similar happenings elsewhere.
- 7. During the year the anti-malarial measures were in no way relaxed and the distribution of quinine was carried out as usual.
- 8. Tablets of quinine to the number of 1,043,200 were issued to the various Health Officers for ultimate free distribution to the public through the Police, the Education Department and the District Officers, also to the Senior Medical Officers for distribution through the various travelling dispensaries.
- Lectures, lantern demonstrations, notices in all languages, and anti-malaria propaganda was vigorously carried out by the Public Health Education Committee.
- 10. Much remains to be done, as an example of this one may quote the case where after a lecture, the lecturer having displayed various forms of quinine, one of his audience a Malay said to him:

"What you say may be all very well but we know that the quinine you have shown us is made from the ground up bones and livers of dead men."

11. The end of the year was remarkable for rains and floods more universal and heavier than those occurring at the end of 1925 and beginning of 1926, and 1927 may show little or no improvement so far as malaria is concerned.

(b).-COMMUNICABLE DISEASES.

- 12. Malaria.—There were 38,633 cases treated in hospitals with 1,792 deaths and a death-rate of 4.64 as compared with 23,056 cases with 993 deaths and a death-rate of 4.03 for the year 1925.
- 13. Blackwater Fever.—There were five cases of blackwater fever with one death. They occurred in the State of Negri Sembilan.
- 14. Dysentery and Diarrhoea.—The total number of cases treated in hospitals was 6,254 with 1,149 deaths and a death-rate of 18.37 per mille as against 4,081 with 638 deaths and a death-rate of 15.63 for 1925. The types were as follows:

States.	Amoebic.	Deaths.	Bacillary.	Deaths.	Diarrhoea.	Deaths.
Perak	 899	137	915	218	1,389	190
Selangor	 354	68	674	180	796	113
Negri Sembilan	 318	75	278	82	228	23
Pahang	 71	14	179	31	153	18
Total	 1,642	294	2,046	511	2,566	344

^{15.} The year under review was a bad one for bowel complaints. The increase was due probably to chills and privations due to the floods and heavy rains and possibly to contamination of rivers and wells from the same cause. The increase of Indian immigrants bringing women and children of poor stamina with them was also a possible factor.

16. Venereal Diseases.—The total number of cases treated in hospitals during the year 1926 was 4,428 cases with 51 deaths and a death-rate of 1.15 per cent. as compared with 3,688 cases with 91 deaths and a death-rate of 2.46 for the previous year.

The campaign against venereal diseases continues with marked success. Persons of all nationalities seek advice and treatment at the Clinics, which are being increased in number.

- 17. The figures given in the attached report by the Venereal Diseases Specialist speak for themselves, but his most remarkable achievement has been winning the confidence of the three hundred and sixty Chinese known prostitutes living in Kuala Lumpur. They attend regularly at the Clinic and seek advice on other conditions besides venereal diseases. It is impossible for anyone not knowing the Chinese to realize what the Venereal Diseases Specialist has accomplished.
- 18. Injections of arsenical compounds are given at all hospitals and dispensaries. The following are the figures for the different States:

States.			1924.		1925.		1926.
Perak	***		14,217		15,913	***	11,863
Selangor			10,525		9,581		4,845
Negri Sembilan		***	3,244		3,091		2,790
Pahang			2,030	***	2,068	***	2,984
	Total		30,016		30,653		22,482
					-		

- 19. Pulmonary Tuberculosis.—The number of cases treated in hospitals during the year under review was 2,329 cases with 995 deaths and a death-rate of 42.72 per mille as compared with 2,571 cases with 1,051 deaths and a death-rate of 40.88 per mille for the year 1925.
- Extensive propaganda in the form of pamphlets was continued by the Public Health Education Committee.
- 21. There is no doubt that influenza leaves Asiatics very liable to infection by the tubercle bacillus and as influenza is more common that it was, so tubercle bacillus infections increase. The floods with the exposure and privations were responsible to some extent for the increase death-rate and number of cases. Overcrowding in the towns is also largely a factor. A Commission has recently sat on the question of housing—'the report is not yet available. It is impossible to persuade early cases to come into or to remain in hospital. They do not feel seriously ill and all they wish for is "a bottle of medicine". Asiatics will not sacrifice their chances of money making for the possible cure of a disease which they do not believe they have. The Government Asiatic Staff are all periodically examined for symptoms of pulmonary or other tubercular infection.
- 22. Yaws.—The treatment of yaws by arsenical compounds was continued. A total number of 30,233 injections were given during the period under review. The number of cases treated in each State in comparison with the number treated in 1923, 1924 and 1925 is shown in the following table:

States.		1923.	1924.	1925.		1926.
Perak		 14,149	 9,884	 9,155		9,358
Selangor		 4,381	 2,231	 1,305		1,680
Negri Sembilan		 9,358	 7,267	 6,426		4,543
Pahang		 3,247	 3,997	 9,403	***	7,893
	Total	 31,135	 23,389	 26,289		23,747

- 23. The difficulty experienced in persuading patients, who are mostly Malays, to continue injections is still very great, but by making facilities for injections as easy as possible it is hoped that yaws will be eventually eradicated. Serious cases so common a few years ago are rarely seen and the milder cases are becoming less frequent. In some districts where some years ago the disease was common it is no longer seen, but fresh centres of infection crop up from time to time. There is also danger of infectious cases coming in from the neighbouring States of Trengganu and Kedah over the extensive land frontiers against which it is impossible to guard. In the course of time yaws may disappear from the Federated Malay States but unceasing vigilance will be needed.
- 24.—Leprosy.—There were 480 fresh admissions and 66 deaths from this disease during the year.

- Infectious Diseases.—There were no epidemics during the year, except one small outbreak of cholera and some cases of measles.
- 26. Typhus Fever.—Sixty cases of tropical typhus were diagnosed at the Institute for Medical Research during the year, thirty-two belonging to the K. group and twenty-eight belonging to the W. group. A special resume of the Institute Report is given.
- 27. Cholera.—A small outbreak of this disease occurred in the Kuala Langat district of Selangor, there being 13 cases and six deaths. A possible cause of infection was from one of a family who had arrived from India two months previously on an infected ship. These coolies, the parents of one child, had been inmates of the non-infectious hospital of the Quarantine Camp. It may be possible that a member of this family was an intermittent carrier.

Three cholera contacts escaped from the Quarantine Camp led by persons who had been previously in this country. They more probably carried the disease, which was further spread by labourers absconding from Permatang Estate lines.

28. Enteric Fever.—There were 147 cases treated in hospitals with 26 deaths during the year as compared with 116 cases and 34 deaths for the corresponding period of 1925.

States.					Cases.	Deaths.
Perak	 			***	37	 7
Selangor .	 				55	 7
Negri Sembilan	 	***	***		39	 7
Pahang	 				16	 5
			Total		147	 26
						1

The remarks of the Director of Government Laboratories are quoted elsewhere.

- Diptheria.—There were 35 cases diagnosed at the Institute for Medical Research during the year.
- 30. Influenza.—Influenza has been more prevalent during the year under review. There were 5,405 cases with 74 deaths as against 4,437 cases and 44 deaths in 1925.
- 31. Pneumonia.—The number of cases treated was 2,066 with 1,202 deaths and a death-rate of 45.09 as against 1,922 cases with 779 deaths and a death-rate of 40.52 for the year 1925. Here again the rains, floods and increase in the labouring Indian population are responsible for the increase in the cases and death-rates.

Influenza is also a factor, and the Indian labourer is very prone to contract pneumonia and has a low degree of resisting power.

- 32. As soon as conditions permit it is hoped to introduce female nursing in the acute male wards of the Asiatic hospitals. This should reduce the hospital death-rate from pneumonia but the difficulty is to find suitable recruits, and sooner or later nearly all the Asiatic Nurses marry and we lose them just as they complete their training.
- 33. Smallpox.—There has been no outbreak of smallpox during the year under review, and vaccination work was mostly confined to children. The number of vaccinations performed was 113,914 as compared with 126,305 for the corresponding period of last year. They were distributed as follows:

States,		1923.	1924.		1925.		1926.
Perak	10.	 53,926	 54,278		86,125		88,589
Selangor	***	 11,075	 11,745		26,369		14,256
Negri Sembilan		 6,366	 6,568		8,268		5,892
Pahang		 6,357	 5,821		5,543	***	5,727
	Total	 77,724	 78,407		126,305		113,914
				-		_	

34. Beri-Beri.—The number of cases treated for this disease was 1,075 with 140 deaths and a death-rate of 13.02 as compared with 906 with 98 deaths and a death-rate of 10.81 for the corresponding period of 1925.

Very active propaganda work in the form of pamphlets and advertisements in the vernacular press warning the public of the danger of using polished rice were issued by the Committee for Public Health Education. 35. There was a marked increase in number of cases treated in Pahang during the year, most of them came from the Railway Construction area north of Chegah Perah and from mining areas near Bentong and Raub. Many Chinese coolies seem to keep themselves on the border line of health and any serious interference with their food supply soon brings on an attack of beri-beri; there was a large increase in the number of cases as a result of the floods at the end of the year.

They are also very unwilling to go into hospital, even when they recognise that they are ill, when money is to be made and the price of tin is high. In spite of all efforts the use of polished rice is increasing and the population will not use undermilled rice.

Rice polishing extract still holds its own against marmite and other treatments, but all are useful.

HELMINTHIC DISEASES.

36. Ankylostomiasis.—A total of 2,493 cases were treated in 1926 as compared with 3,309 cases in 1925. These figures are for cases who were treated for these diseases only. In various districts, campaigns against this and other intestinal parasites are conducted and efforts are made to improve rural sanitation generally. As soon as staff are available a more general effort throughout the Federated Malay States will be made.

The Public Health Education Committee has carried on propaganda by means of pamphlets.

- 37. Ascaris lumbricoides is almost a universal infection. It has caused death in adults, and in children of all ages is a serious condition untreated, to quote one other form of disease alone.
- 38. The demands made for medicine for worms at the out-door centres and travelling dispensaries are very numerous and refer almost entirely to Ascaris infections as other worms if passed are seldom noticed.

(c).-VITAL STATISTICS.

- 39. Births and deaths figures are obtained from notifications compulsory under the registration of Births and Deaths Enactment which is everywhere in force. The total number of births and deaths is approximately correct. The accuracy of diagnosis as to cause of deaths is in the majority of cases open to question, for few of the cases were seen by a qualified medical man previous to decease. In each of the four large towns, every uncertified body is viewed by the Assistant Health Officer who interrogates the friends and forms a diagnosis so far as possible. In rural districts, these duties are carried out by the Police.
- 40. Deaths in towns are debited against the town only if the deceased was resident there for three months or more previous to death. The towns contain hospitals which cater both for the town and the district surrounding it. It is a well-known fact that chronic cases from the rural areas drift to the towns in the hope of getting more skilled treatment usually at the last moment. Taking all things into consideration even with a qualifying period of three months a number of deaths are debited against the towns which should be debited against the rural areas where the diseases were contracted.
- 41. Because of the peculiar age and sex distribution, and the fact that the labour of this country is largely made up of persons aged 20 to 45, who passed the doctor before embarking for Malaya, the death figures cannot be compared with countries where the labour is indigenous and where age and sex distribution are normal.
- 42. In the case of Chinese immigrants no medical examination is made before they leave China but they are usually able-bodied young bachelors.
- 43. Population.—The population of the Federated Malay States as estimated was at the end of June, 1926, 1,476,032, distributed as follows:

Perak	***	 			***	 654,179
Selangor		 				 457,170
Negri Sembilan		 	***	***		 204,257
Pahang		 				 160,426

Assuming that the races remain in the same proportion as in the Census year the race distribution is as follows:

European	s and	Americ	cans					 6,947
Eurasians					***			 3,495
Malays a	nd oth	er nati	ves of	the Ar	chipels	igo		 558,060
Chinese		***					200	 526,733
Indians			***		***	***		 374,915
Others		M				1000		 5.882

- 44. Owing to the unprecedented severity of the floods which occurred and which were particularly severe in Pahang resulting in the complete destruction of the records of the Registrar of Births and Deaths in that State, the figures given for births and deaths are for the States of Perak, Selangor and Negri Sembilan only.
- 45. Births.—The number of births registered was 39,834 for the whole year, giving a rate of 30.28 (excluding Pahang) per mille of population as against a birth-rate of 28.89 per mille for 1925.

The following table shows the number of births and birth-rates according to races:

	Race.					Number of	t	Birth-rate.
Europeans a	nd Ame	ricans				131		19.86
Eurasians	***	***	444			127		87.76
Malays and	other rac	es of th	he Arc	hipelag	0	17,522		39.12
Chinese		***				12,973		26.61
Indians						9,010		24.68
Others						71		13.67

- 46. Deaths.—Thirty-eight thousand four hundred and forty-five deaths were registered, giving a death-rate of 29.22 (excluding Pahang) per mille. The number of deaths in 1925 was 34,158 and the rate was 23.60 for the whole Federated Malay States.
- 47. It is probable that the introduction of more female Tamil labourers bringing with them children of poor physique, and malaria, dysentery and diarrhoea following the heavy rains and floods may account for the increase.

The distribution of deaths among the several races was as follows:

	R	ace.					Number of deaths.	Death-rate.
European	ns and	Ame	ricans			***	36	 5.46
Eurasian	S					***	46	 13.68
Malays a	and oth	er ra	ces of t	he Arc	hipelage	0	11,657	 26.03
Chinese							14,730	 80.22
Indians	4	***	***				11,920	 32.64
Others				W			56	 10.78

INFANT MORTALITY.

48. The number of deaths of children under one year was 7,718 or an infantile mortality rate per thousand births of 193.75 against a mortality rate of 177.17 in 1925. This death-rate which it is hoped will be reduced in the course of time, compares favourably with those of other tropical countries. The death-rates for the three States were:

		States.					ns of child er one yea	Death-rate per 1,000 births.
Perak							3,459	 172.18
Selangor	***				110	***	2,888	 207.56
Negri Se	mbila	n					1,371	 225.86
Pahang	(Figu	ires not	ave	ilable	owing	to fl	oods)	

49. The infantile mortality rate in Negri Sembilan still remains the highest, but it is hoped that with the further extension of Infant Welfare work to reduce this in the near future.

(d).—GENERAL EUROPEAN POPULATION.

- 50. The general health of the European population continues to be good. There was little sickness and invaliding. The total European and American population as estimated at the end of June, 1926, was 6,947. There were 131 births, giving a birth-rate of 19.86 per mille and 36 deaths with a death-rate of 5.46 per mille as compared with a birth-rate of 20.72 per mille and a death-rate of 4.03 per mille in the previous year.
- 51. It is necessary to remark upon the increasing number of cases of neuresthenia, Europeans of both sexes being affected. There is no doubt that in the case of many men the cause is the increase in the amount of work to be done, the monotony of the climate and of the life, in spite of recreations unknown in the past, and perhaps the tendency to burn the candle at both ends, stay up half the night and work all day. In most stations the daily routine is the same and the members of the clubs do not change. Small worries become great ones only too easily, often

with the most disastrous results. In the case of European women lack of definite occupation and the straining after amusement and excitement are often to blame, but the climate does not suit a good many ladies. It is possible that were the fourteen days casual leave now available yearly extended to a month things might improve. It is to be hoped that when the Cameron's Highlands Hill Station is an accomplished fact conditions will be altered.

SANITATION.

- 52. The Health Branch is still understaffed; of the authorised senior staff of one Senior Health Officer, 15 Health Officers and eight Chief Sanitary Inspectors, only six Health Officers and six Medical Officers seconded to the Health Branch, and two Chief Sanitary Inspectors were available. One of the latter on his return from leave at the end of February, 1926, was stationed at Tapah to supervise the anti-malarial works in connection with the road to Cameron's Highlands which is to be developed into a Hill Station.
- 53. Of the 28 Sanitary Inspectors eight of them were absent in Singapore for six months undergoing a course preparatory to taking the examination for the Royal Sanitary Institute certificate which certificate must be held before an Inspector can be promoted to the higher grades. Their services were not available to the branch from May until November.

INSPECTION OF SCHOOLS.

- 54. The inspection of schools is still being carried out by both Medical and Health Branches of the Department. Frequent visits are being paid to the schools near the main roads by the travelling dispensaries. All schools are supplied with a stock of quinine tablets.
- 55. Dental caries is largely prevalent and more attention should be paid to the feeding of children. The food stalls in schools are often untidy and the food unwholesome. Food vendors and hawkers should be prohibited as far as possible. As previously reported latrine accommodation and water supply are very unsatisfactory in a number of schools. School inspection, especially the medical inspection of school children, will never be satisfactory until a staff of medical school inspectors, male and female, are appointed. A whole time school dentist has also been asked for.
- 56. It is necessary to note that in the matter of sanitation all this department can do is to recommend improvements to the Education authorities, and it is for the Education and the Public Works Departments to carry out the necessary alterations. The staff have done their best to carry out these inspections but both branches are much under strength and many schools are far from the beaten tract and the time to visit them cannot be found.
- 57. The work under the Labour Code was satisfactory in view of the number of staff. Of 1,450 estates, 725 were visited by Health Officers and of 156 estate hospitals 154 were visited. With the improved conditions in this branch recruits cught to be forthcoming. Further details of the work of the Health Branch will be found in the Senior Health Officer's report annexed.

PUBLIC HEALTH EDUCATION.

- 58. The Public Health Education Committee met on several occasions during the year; at Horticuitural Shows exhibitions were put up which were well attended, lectures were given and pamphlets which had been prepared by the Committee were distributed in large numbers in various languages.
- 59. Popular lectures on malaria were given in Malay and Chinese in various parts of the country, these were illustrated by lantern slides and were well attended.
- 60. A cinematograph film of the work done at the Infant Welfare Centre, Kuala Lumpur, was also exhibited on several occasions to large and interested audiences. The film was photographed and produced by Mr. Hoflin, Secretary of the Infant Welfare Advisory Board.
- 61. Lectures on venereal diseases, illustrated by lantern slides, were given in Chinese, Tamil and other Clubs and Institutions.

QUARANTINE AND PORT HEALTH WORK.

- 62. A very marked increase of immigrants, 99,066 for 1926 compared with 48,748 for 1925, taxed the accommodation at Port Swettenham Quarantine Camp somewhat severely, in view of the fact that several successive steamers arrived infected with cholera. The situation was satisfactorily dealt with by the Quarantine Camp Staff. Improvements in the Camp will be effected in 1927 and absconding made more difficult.
- 63. During the year 75 ships with immigrant labourers were boarded and inspected. The labourers were landed at the Quarantine Camp. Of the 75 ships, 21 were infected—eight with cholera, two with cholera and smallpox, four with smallpox, one with cerebro-spinal meningitis, one with cholera and cerebro-spinal meningitis and five with chickenpox.
- 64. The number of immigrants who entered the Quarantine Station, Port Swettenham, was 99,089, the number remaining on 31st December, 1925, was 1,408, making a total of 100,497. The largest number on any one day was 7,713 on the 20th of June. The following table shows how these immigrants were distributed:

Remaining on 31st December, 1926	***	m 4 -1	1,697
Died in Hospital and Cholera Wards	11.20		 457
Absconded from the Quarantine Camp	1	00000	 55
Transferred to Klang Hospital	2.0	100.77	 104
Discharged to Depôt	1440	012 0 19	

The total number of persons removed to the Camp from cholera infected ships was 15,061, of these 157 either developed cholera or had the disease on admission to the Camp, 39 cases remained on 31st December, 1926. Of the 196 treated 79 died giving a percentage of 40.31.

- 65. Every precaution was taken to cut short the course of the epidemics. Prophylaxis by anti-cholera inoculations, and essential oils were adopted, and I think with success, as shown by the figures, 157 cases with 15,061 contacts.
- 66. Pneumonia are not uncommon amongst the immigrants. They seem to be largely due to defective ventilation arrangements between decks. Unfortunately two new ships, in which special precautions were adopted as regards ventilation and sanitation, having been taken by the Government of India as transports for troops to China.

INFANT WELFARE WORK.

67. Infant Welfare work continues with great success and the attendance at the Clinics have greatly increased. The number attending Clinics in Kuala Lumpur, Ipoh, Taiping and Seremban for 1925 and 1926 were as follows:

Place. Kuala Lumpur				1925. 23,134		1926. 29,831
Kuala Lumpur Ipoh	,	***		15,528		14,080
Taiping			***	18,259		13,559
Seremban (opened	in August,	1926)	000	100	***	3,895
		Total		56,916		61,365

68. It is interesting to note the change that has taken place in the attitude of the parents towards the aim and object of the Centre. They were originally full of fears and prejudices and attended only after much persuasion and in a very hesitating manner; they vouchsafed no signs of approval or otherwise at the instructions given them and appeared completely mystified and far from happy. To-day on visiting the Centre one can see for oneself that these same people are now thoroughly at home and have lost their fears, their attendance weekly is regular and they bring not only their own babies but persuade their neighbours to bring theirs; they demand instructions saying "please tell me about the feeding", they occasionally even ask for the instructions to be written in case they forget them. Such change of attitude is due in a large measure to time, helped no doubt by the fact that they have grasped a clearer idea of the aim and object of the work, and thus realise their responsibility with regard to the welfare of their children.

District visiting is being carried out as much as possible. It is recognised as the most important part of child welfare service and the work done in this direction has been exceptionally good through the year.

69. The reports of the Lady Medical Officers, Infant Welfare Centres, are appended.

HOSPITALS, DISPENSARIES AND VENEREAL DISEASES CLINICS.

70. Hospitals.—The number of in-patients treated in hospitals was 127,332 with 9,178 deaths and a death-rate of 7.21 as compared with 103,762 with 6,964 deaths and a death-rate of 6.71 in 1925. The distribution of patients in the different States was as shewn below:

			1925.				1926.						
States.		Cases.	Deaths.		Death- rate.	Cases.		Deaths.		Death-			
Perak		46,551	 3,198	***	6.87	56,909		3,970		6.98			
Selangor		34,768	 2,172		6.24	40,464		2,904		7.17			
Negri Sembilar	1	14,684	 1,090		7.04	19,400		1,615		8.32			
Pahang	***	7,759	 504	***	6.49	10,559		689		6.52			
Total		103,762	 6,964		6.71	127,332		9,178		7.21			
	-					-		-		Commence of the last			

 The principal diseases commonly treated in hospitals were malaria, venereal diseases, ankylostomiasis, dysentery, diarrhoea, beri-beri, pneumonia and pulmonary tuberculosis.

The following table shows the number of cases and deaths during the years 1925 and 1926.

		1925.			1931,				
Diseases,	No. of cases,	No, of deaths,	Percentage of deaths.	No. of cases,	No, of deaths,	Percentage of deaths.			
Malaria	23,056	993	4.03	38,633	1,792	4.64			
Venereal diseases	3,688	91	2.46	4,428	51	1.15			
Ankylostomiasis	3,309	190	5.74	2,493	120	4.81			
Dysentery	2,694	549	20.38	3,848	852	22.14			
Diarrhoea	1,387	89	6.41	2,566	344	13.41			
Beri-beri	906	98	10.81	1,075	140	13.02			
Pneumonia	1,922	779	40.52	2,666	1,202	45.09			
Pulmonary tuber-					10.00				
culosis	2,571	1,051	40.88	2,329	995	42.72			

72. Womens' Hospitals and Wards.—The work in womens' hospitals and wards is increasing and there has been a gratifying increase in the number attending for confinement. The number of patients treated in Kuala Kangsar Hospital during the year was 1,776 an increase of 516 of the previous year. It is very unfortunate that the records and books were lost or damaged in the flood at the end of the year in view of the considerable increase in the number of both in-door and out-door patients that demonstrated the growing readiness of Malays to make use of this hospital.

The hospital was flooded on December 30th, some of the houses being completely submerged. The water rose to a height of ten feet in the wards. The consequent damage and loss in respect of hospital property especially in the case of books and papers was considerable. All the patients had to be evacuated and were removed eventually to Taiping.

- 78. The Maternity Ward.—The success of the new maternity ward in its first full year is the most interesting feature of the report of the Lady Medical Officer, Kuala Kangsar. It has been rare for any of the six beds to remain long unoccupied, the number of labour cases treated being 86 an increase of 24 on the number for 1925. This does not include premature labours or cases admitted for complication after the birth of the child.
- 74. The Malay Ward, Kuala Kangsar.—The popularity of the hospital with Malays has doubtless been raised owing to the confidence shewn by His Highness the Sultan of Perak in allowing his wife the Raja Perempuan to be admitted as a patient for operation. She made an excellent recovery.
- 75. There was a decided increase in the number of children and babies treated. It is very satisfactory that, in many cases, mothers shewed their confidence in the treatment by leaving their children entirely to the care of the hospital. The fact that mothers seem no ionger insistent on accompanying their children has made the Lady Medical Officer consider the establishment of a children's ward to be a practical possibility in the future. The addition of a European Sister to the staff has noticeably raised the standard of nursing in this hospital.

76. In other hospitals where Lady Medical Officers are stationed the work has been most satisfactory and these ladies have added greatly to the numbers of Asiatic women admitted for confinements. The Lady Medical Officers are excellent at general work as well as at maternity.

Practically every hospital now has its labour ward and room and Asiatic midwife to assist the qualified staff.

SURGERY.

77. Operations.—The number of operations undertaken during the year is shewn below:

		Major.				antida	
States.	1925.		1926.		1925.		1926.
Perak	344		448	***	2,229	***	2,691
Selangor	334		311		1,579		1,257
Negri Sembilan	86	11.00	98		1,618		1,680
Pahang	12		17		296	***	314
Total	776		869		5,722	do]ud	5,942

The reports of the Chief Surgeons, Perak and Selangor, are attached as appendices.

There are now three highly trained and skilful surgeons available in the Federated Malay States and there are also promising juniors.

78. A glance at the lists of operations performed will show that Federated Malay States surgery is at last on a sure footing. Patients no longer go to Java and Singapore and when new operation rooms and hospitals are erected a great increase in surgical work may be anticipated. High class Asiatics are beginning to avail themselves of the surgeons knowledge.

EYE DISEASES.

79. Out-Patients.—The number of new cases during the year at the Eye Clinics was as follows:

				1925.	1926.
Kuala Lumpur	44-		 ***	1,369	 2,112
Ipoh and Taiping	4.40	4	 	3,097	 3,740

There were a total number of 13,329 attendances as compared with 23,220 for the year 1925.

80. In-Patients.—One thousand three hundred and forty-one cases were admitted into hospital as compared with 1,071 cases during the year 1925. The chief diseases met with were: conjunctivitis, trachoma, gonorrhoeal-ophthalmia, ulcers and opacities of cornea, cataracts, neuroretinitis, glaucoma and injuries to cornea and eye-ball, resulting chiefly from accidents chiefly amongst the workmen in factories and metal quarries, and burns from acids and corrosives.

There are three treatment Centres, Taiping, Ipoh and Kuala Lumpur and the work is increasingly popular. The reports are attached as appendices.

CURE FOR OPIUM HABIT.

81. There is little to report this year except that the number of admissions gradually declined throughout the year until at the end there were no patients in any of the wards.

Many absconded and the great majority was discharged at their own request, of those who underwent the full course of treatment it is impossible to say how many were benefited permanently as these cases cannot be followed up once they leave hospital. The opinion is that the percentage is almost negligible.

It would appear that some came in for rest and food. Many were caught with opium in their possession and some went out to secretly smoke opium during the night.

82. I think it may now be taken that the opening of the opium wards has been a failure.

Actually the sales of opium increased every month after the opening of these wards. Six thousand patients were treated and if the cure was a success this number ought to have materially affected the consumption as far fewer Chinese smoke opium than was formerly the case.

X-RAY AND ELECTRICAL TREATMENT.

83. Dr. C. F. Constant was the Radiologist throughout the year. The figures given below show the actual number of patients treated or examined:

X-ray cases	 	 	 	722	714
Electrical treatment	 	 	 	***	209

- 84. A well organised X-Ray Department has been opened in Kuala Lumpur. New X-ray equipment has been installed in Ipoh and an X-ray set will be installed in Seremban in 1927.
- 85. The Radiological Building took a long time to mature and was very badly required. As none of the old buildings could support the new X-ray apparatus and as the old apparatus broke down beyond repair for a time the Federated Malay States had no X-rays available. The Radiologist visits Ipoh at regular intervals, and a Medical Officer with radiological experience is available for the minor examinations. Electrical treatment on modern lines is available in Kuala Lumpur and will be installed in Ipoh and Seremban as soon as apparatus is to hand. The Radiologist has been greatly hindered by breakages in transit due to bad packing in England and also to lack of instructions from the English makers as to assembling plant, a sad contrast to the business-like and thorough methods obtaining in America. A word must be said concerning the Radiologist's energy and skill in assembling and putting up the apparatus and his great knowledge as a practical electrician. His report is attached.

HOSPITALS AND DISPENSARIES.

86. Out-Patients.—The number of out-patients treated in all hospitals, dispensaries, travelling dispensaries and dispensary boats for 1926 was 630,052 as compared with 621,793 in 1925. The following figures show the number treated during the past four years:

States.	1923.		1924.		1925.		1926.
Perak	190,990		221,096		216,282		210,511
Selangor	159,402		177,896		219,739		227,864
Negri Sembilan	92,692	100	96,432		99,047	10110	101,652
Pahang	84,329		88,837		86,725		90,025
Total	527,413		584,261	od mo	621,798	I on't	630,052

- 87. Of the total of 630,052 cases 156,652 were treated by the travelling dispensaries and 16,817 by dispensary boats on the Pahang River. In Selangor a system of house to house visits was inaugurated. The travelling dispensaries staff visited Malay kampongs which were some distance off the road and whose inhabitants would not trouble to attend at the usual stopping places of the travelling dispensary. A total of 460 patients out of a population of 3,416 was treated by these means. It is intended to extend this system as more senior Malay dressers become available.
- 88. In 1927 a number of new and improved type of larger travelling dispensaries are to be supplied. They are being built on a one ton Morris Chasis. They will gradually replace the fords which are becoming worn out after many years of excellent service.
- 89. The motor boats dispensaries are under construction one for the Perak River and one for the Pahang River. The former has now, at the time of writing, been supplied and it promises to be a great success. The design produced by the Medical Department with the aid of the Radiologist, who was for some years in the Royal Navy, has been adopted by Messrs. Thornycroft as a standard for river ambulance launches. Little was forgotten and as an ambulance launch these ten knot ships have no rivals. They will be of immense value to the Malay riverine population.

PRISONS.

90. During 1926 the general health of the prisoners has continued satisfactory. The total number treated during the year in the different gaol hospitals was 635 with 10 deaths and a death-rate of 1.57 as against 958 with 16 deaths and a death-rate of 1.69 for the year 1925. They were distributed as in the following table:

o-1980 tower tild so and				1925.						1926.		
Place.		Cases.		Death	8.	Per- centage.		Cases.	I	eaths.	ce	Per- ntage.
Pudu Gaol, Kuala Lumpu	r	158		1		0.57	***	125	***	1		0.80
Gaol, Taiping												4.96
						1.10						144
Seremban		255		6		2.35		119	***	2		1.68
, Kuala Lipis	***	54		-		-	***	55		-		-
Kuantan	911	20		-		-		17		-		-
of believed of live to Tot	al	958	111	16		1.69		635		10		1.57
				1000								

As is usual, all new prisoners are kept under observation. Their blood and faeces are examined and they are vaccinated before being passed to labour.

Dysentery which was at one time very prevalent has wholly disappeared as a result of measures taken in Taiping in 1917.

LEPER ASYLUMS.

91. Kuala Lumpur.—The total number of lepers treated during the year was 704 with 42 deaths and a death-rate of 5.96 per cent. as compared with 635 cases and a death-rate of 4.88 per cent. for the year 1925. A report on the working of this institution is attached as an appendix.

A feature for the work is the number of voluntary admissions and the return of escaped patients. Old and unsuitable as this settlement is, it is a credit to the officers and patients in charge.

- 92. Pulau Pangkor Laut.—In this asylum which is for Malays only 68 cases of leprosy were treated during the year. There were six deaths. The Medical Officer in charge reports that the hoped for results when the "Tai Foong Chee" treatment was introduced in 1925 were not obtained. The inmates were soon tired of this form of treatment and it was not given a fair chance. The failure, perhaps, are attributable to the extraordinary mentality of the Malay.
- 93. Once admitted to the asylum he makes up his mind to stay there. The Medical Officer states that during the last three years he can only point to one case, a young female, who is anxious to get well and join her people again. She alone persevered with Tai Foong Chee and E.C.C.O. and improvement in her case is marked.
- 94. The District Officer, Lower Perak, recently visited and remarked in the Visitors' Book that the patients are as happy as it is possible for them to be. More accommodation will be required before long in spite of that supplied already, and the island is rather too small, or shortly will be, for requirements. Marriages amongst the patients are now permitted by Government.
- 95. Taiping Leper Wards.—Eighty-eight cases were admitted during the year. Sixty-six remained at the end of 1925. There were 14 deaths in 1926. It is hoped to shortly transfer all the Taiping lepers to the Kuala Lumpur Asylum which is being extended. Only a small reception ward will remain in Taiping.

TEMPORARY CHINESE DECREPIT ASYLUM, PORT SWETTENHAM.

96. The policy is to allow this camp to decline in numbers until it is empty or transferred elsewhere. No new decrepits were admitted into this home since July, 1925, owing to want of accommodation. The number of decrepits remained on 1st January was 441 and the number remaining at the end of the year was 229. Sixteen were discharged, sixty-seven absconded, sixty-three died and sixty transferred. Chronic patients should not be in the hospitals which should be reserved for cases of acute disease. A large institution for such patients will be available in the course of time twelve miles from Kuala Lumpur where plenty of agricultural land is at disposal and where these people can be taught trades and earn some money. A site for a temporary camp near Port Swettenham is under consideration.

VETERINARY BRANCH.

		RIN	DERPEST.				
States.			Cases.		Deaths.		Destroyed.
Perak	 ***		850		642		
Selangor	 ***		134	***	1		133
Negri Sembilan	***		1,327		1,221		5
Pahang	 		2,811		2,021	***	664
	70-4-1		F 100		0.005		
	Total		5,122	***	3,885	***	797

97. In Negri Sembilan, the seriousness and extent of the outbreak was due to the fact that the areas infected were mostly Malay kampong areas where buffaloes are plentiful and predominated over cattle, with their great susceptibility and high mortality. In most cases the disease was held in check by simple quarantine measures, but success depended on the strictness with which these measures were enforced, and were obeyed. Great trouble was experienced in some districts in this direction and it was due to the lack of co-operation on the part of the public that the disease persisted for over four months.

98. In Pahang outbreaks occurred in Temerloh, Kuala Lipis, Bentong, Raub and Kuantan. The one in Raub district was most severe. The outbreak commenced in November, 1925, and lasted until November, 1926.

FOOT-AND-MOUTH DISEASE.

St	ates.			Cases.		Deaths.	Destroyed.
Perak			***	 85	***	4	
Selangor		***		 285		1	
Negri Sem	bilan			 2		-	
Pahang				 -		-	 -
			Total	 320		5	 _
						The same	100

99. Rabies.—Fourteen cases of rabies occurred in Negri Sembilan. The first case was at Johol, and cases following in the Kuala Pilah and Tampin districts. One of these cases occurring in a monkey.

Two hundred and seventy contact dogs were vaccinated with special canine rabies vaccine supplied by the Institute for Medical Research. Two vaccinated dogs developed the disease, one three days after vaccination, the other about a month later.

No cases occurred in the other States during the year.

100. Swine Fever.—There were no cases of swine fever reported in the Federated Malay States during the period under review.

. 101. Surra.—There were only four cases of this disease. They occurred in dogs admitted for treatment in the Animal Infirmary, Taiping, Perak, all of them were destroyed.

VETERINARY QUARANTINE STATIONS.

102. Port Swettenham.—Fourteen thousand one hundred and forty-nine animals were imported via Port Swettenham, of which number 4,224 cattle were quarantined for 10 days. Four outbreaks of rinderpest and two of foot-and-mouth disease were detected in newly imported animals. The new cattle Quarantine Station built by the Railway Department was repaired, a water supply was laid on and it was used during the latter half of the year.

103. Kuala Lumpur.—One thousand and five cattle were quarantined, these were chiefly animals imported from Singapore. The new cattle Quarantine Station at Bungsar Road was taken over during September, the old one at Bukit Sentul being given up. The former is more convenient as it contains a large grazing area and it is closer to the railway and abattoirs.

104. Kuantan.—One thousand one hundred and ninety-four animals passed through the pig Quarantine Station, Kuantan, no disease occurring.

105. Perak.-The following cattle were quarantined at:

Port Weld	***	 		***		 3,297
Parit Buntar		 				 112
Selama		 	***			 307
Teluk Anson		 	***		***	 859

106. Negri Sembilan.—Animals were detained in the cattle Quarantine Stations as follows: Seremban 123 cattle and 189 buffaloes, Tampin 41 cattle and 58 buffaloes and Port Dickson 163 cattle.

VETERINARY PROSECUTIONS.

107. There were a total number of 1,171 prosecutions resulting in 1,126 convictions. The fines imposed amounted to a total of \$12,796. They were distributed as follows:

States.		Prosecution	8.	Conviction	8.	Fines.
Perak	***	 469	***	457		\$5,668
Selangor		 395	***	390		3,279
Negri Sembilan	711	 109		93		2,623
Pahang		 198		186		1,230
	Total	 1,171		1,126	Ingres.	\$12,796

METEOROLOGY.

RESUME OF METEOROLOGICAL CONDITIONS IN 1926.

- 108. During the first half of the year the rainfall was generally below the normal, especially during January and February. The temperature was considerably above normal for the same period. A maximum temperature of 97°F was recorded in Kuala Lumpur on April 3rd and again on April 4th. This is the highest shade temperature recorded in Kuala Lumpur, the highest previously experienced being 96°F.
- 109. The rainfall during the period December 17th-31st was general throughout the Peninsula, and was the heaviest of which there are any records. 26th to 29th were the days of heaviest falls, especially in Pahang. At Kuantan 20 inches were recorded on the 27th, and 54 inches in the four days 26th-29th.

SCIENTIFIC.

COMMENTS ON THE ANNUAL REPORT OF THE DIRECTOR, GOVERNMENT LABORATORIES, F.M.S.

- 110. This report is full of matters of the greatest interest and a few words drawing attention to the most salient features may be of interest.
- 111. "Plasmochin".—Experiments were conducted with a quinine substitute named "Plasmochin" a German preparation manufactured by the Farbenindustrie Aktiengesellschaft at Leverkusen. It was found to be effective in benign tertian and quartan infections and destructive to all the sexual forms of parasite in the blood, but to have very little effect on subtertian parasites. In using it very careful microscopical work and frequent examinations are necessary. There is also some question as to its toxinity not yet settled. It is therefore not suitable for general use and distribution. In fact Messrs. Bayer recommend the use of quinine in subtertian infections in addition to "Plasmochin".
- 112. Tsutsugamushi Disease, Kedani or Japanese River Fever.—Four cases occurred amongst Europeans, of whom three were certainly infected in the Federated Malay States. The disease is not uncommon in Sumatra but only one previous case was known in the Federated Malay States. The disease is conveyed by the bite of small acarines, rather similar in form but smaller than the Autumn mite seen in England. Field rats act as the intermediate hosts. Rats make their nests in lallang and blukar and these places harbour the mite which feeds on the rat's blood but attacks man also, and many other animals. The mite is microscopic in size. One European case was fatal. A pamphlet on the prophylaxis for planters and their labourers is being prepared for distribution.
- 113. Tropical Typhus.—Sixty cases were diagnosed and of these thirty-two belonged to the "K" group or strain and twenty-eight to the "W" group. Laboratory names to distinguish slightly differing reactions obtained from otherwise apparently similar organisms. The disease is moderately common in the Federated Malay States though only recently recognised.
- 114. Melioidosis.—This fatal disease was recognised in 1926 in Indo-China, a case was found as a natural infection in a dog in the Federated Malay States, and a man also in the Federated Malay States was infected through a superficial ulcer on the leg. Until 1926 this disease had not been found outside Burma and the Malay Peninsula. In all there were four cases occurring in Kuala Lumpur. The disease is so rapidly fatal that it is impossible to say how many occurred elsewhere.

- 115. Leptospirosis, Infective Jaundice.—The organism has been passed through guinea pigs and definitely recognised. The staff have been investigating how many diseases may be caused in the Federated Malay States by leptospiras it is suspected that forms of this type of organisms are responsible for several.
- 116. Measles.—Dr. Neave Kingsbury conducted experiments with serum taken from the blood of convalescent measles cases. It was found that injections of this serum modified attacks of measles and conferred immunity lasting for some time.
- 117. Pneumonia.—Overcrowding amongst Police recruits was found to be responsible for an attack of pneumonia.

Overcrowding in the same barracks had previously caused outbreaks of cerebro-spinal meningitis in other years.

- 118. Bacteriophage.—The Bacteriophage supplied by Dr. F. d'Herelle from Egypt was found to be non-effective against the Flexner group of dysenteries common here, though it was most effective in Rio de Janeiro with Shiga infections. Enquiries are being made, and Dr. d'Herelle will visit the Federated Malay States in 1927.
- 119. Diptheria.—Dr. Fletcher decides amongst other observations that carriers are common and the population highly immune.
- 120. Enteric (Typhoid Fever).—The evidence goes to show that this disease is unusually uncommon in the Federated Malay States. Barbadoes claims to be very healthy but had for five years an average of 1,332 cases yearly which were notified, as against the Federated Malay States hundred or so notified cases. The population of Barbadoes is 159,499 only, that of the Federated Malay States over 1,500,000.
- 121. Anti-rabic Treatment.—The Pasteur Institute has been a success 183 persons were given anti-rabic treatment in Kuala Lumpur. A branch treatment Centre was opened in Malacca the material being supplied from Kuala Lumpur.

Prophylactic inoculations of dogs against rabies were successfully carried out.

- 122. Morbid Growths.—In all 195 specimens of morbid growths were examined and reported on for various hospitals.
- 123. Chemical Division.—The Chemical Division carried out numerous investigations for the medical and other departments. Amongst others a laborious investigation into the effluents from small experimental sewage installations which is still going on.

Rice polishing extract for use in beri-beri was manufactured on a large scale by the Chemists.

124. Malaria Bureau.—Nine special reports were issued during 1926. Exhibits were shown at Agricultural Exhibitions. The investigation in malaria in rice-fields was continued in Perak and Negri Sembilan.

Type collections of Malayan mosquitoes were sent to fifteen persons.

The larvicidal properties of rubber oil were enquired into. It is efficient only when heavily applied, and it must be used fresh.

- 125. Malaria Carriers.—Experiments made on A. separatus tended to prove that it was not a carrier.
- 126. During the year the Institute for Medical Research and the Malaria Bureau have been short of staff and overworked, but the results are excellent.

COMMENTS ON THE REPORT OF THE MEDICAL SUPERINTENDENT, CENTRAL MENTAL HOSPITAL, TANJONG RAMBUTAN.

127. There were at the end of 1926, 1,212 male and 338 female patients at the Central Mental Hospital. They were made up as follows:

Incircui AAVA						Males.		Females.
Remained	on 31st Dece	mber,	1925		***	1,145		306
Admitted d			***			489		155
	-Recovered		***			151		56
,,	Relieved					56		20
,,	Not improv	ed		***		54		22
"	Not insane			***		1	***	
Absconded			****		***	78	***	-
Died						82	***	25
				Total		1,212	***	338

128. The following admissions were made during the year from places outside the Federated Malay States:

				Males.	Females.
Singapore	 	 ***		 1	 -
Kedah	 	 ***	***	 40	 11
Kelantan	 	 		 2	

- 129. Admissions.—The total admissions during the year was 735 which shows an increase of 69 on last year, when the numbers were 666. In 1924 the admissions were 614.
- Discharges.—These numbered 360 of whom 207 were discharged recovered and 76 relieved.
- 131. The Medical Superintendent comments on the danger of alcohol taking the place of opium amongst the Chinese.

APPENDICES.

- 132. The following are attached as appendices:
 - A .- Report of the Director, Institute for Medical Research.
 - B.—Report of the Malaria Research Officer, Institute for Medical Research.
 - C .- Report of the Chemist, Institute for Medical Research.
 - D .- Report of the Chief Health Officer.
 - E .- Report of the Registrar-General of Births and Deaths.
 - F .- Report of the Chief Surgeon, Perak.
 - G .- Report of the Chief Surgeon, Selangor.
 - H .- Report of the Medical Superintendent, Central Mental Hospital.
 - I .- Report of the Specialist, Venereal Diseases.
 - J .- Report of the Ophthalmologist Surgeon, Ipoh.
 - K.—Report of the Deputy Medical Officer in Charge of Ophthalmic Clinic, Kuala Lumpur.
 - L.—Report of the Radiologist, Federated Malay States.
 - M.—Report of the Medical Officer in Charge of the Leper Asylum, Kuala Lumpur.
 - N.—Report of the Lady Medical Officer, Infant Welfare Centre, Kuala Lumpur.
 - O .- Report of the Lady Medical Officer, Infant Welfare Centre, Ipoh.
 - P.—Report of the Lady Medical Officer, Infant Welfare Centre, Taiping.

KUALA LUMPUR, 20th April, 1927. R. DOWDEN,
Principal Medical Officer, F.M.S.

TABLE 1.

RETURN OF DISEASES AND DEATHS (IN-PATIENTS) FOR THE YEAR 1926.

The form shows in the main the arrangement of diseases in the International Nomenclature, 1921 Edition. To save space the unimportant diseases of any class can be grouped in their places as "Other Diseases" of the class.

- * i.e. the year previous to that for which the return is made.
- † "Total cases treated" will, of course, include those remaining in hospital at the end of the previous year.
- ‡ The figures in this column to be carried on to the next year's return.

						og in 925.	Yearly	y total.	8 0 8	g in 926.	
	Diseas	es.				Remaining in Hospital at end of 1925.	Admis- sions.	Deaths.	+ Total cas treated.	‡ Remaining in Hospital at end of 1926.	Remarks,
	I.—EPIDEMIC, ENDEM		o In	FECTIO	US					TAKE OF THE PARTY	
1.	Enteric group—							1			
2. 3.	(a) Typhoid fever (b) Paratyphoid A. (c) Paratyphoid B. (d) Type not define Typhus Relapsing fever Undulant fever	***				13 1 3 65	115 4 15 9 56 	23 3 3	128 4 15 10 59 606	13 ₂ ₃	
	Other types	***				9	482	13	491	2	
5.	Malaria— (a) Tertian (b) Quartan (c) Aestivo—autum (d) Cachexia (e) Black-water (f) Mixed infection					123 17 320 157 	9,224 557 18,624 9,264 28 216	194 12 1,146 407 5 22	9,847 574 18,944 9,421 28 222	277 7 389 197 	
6.	Small-pox-				3 1			100	1	CLEUMD O	
8. 9. 10. 11.	Alastrim Measles Scarlet fever Whooping cough Diphtheria Influenza					27 5 125	588 58 45 5,405	3 5 11 74	560 58 45 5,530		Solding to be seen to
	Miliary fever Mumps		***			2	96	2	98	"1	
14.	Cholera Epidemic diarrhœa					1			1		
16.	Dysentery—				min!		1000	100	- Intel	1 mg 200 h	
17.	(a) Amæbic (b) Bacillary (c) Undefined or du Plague—	 e to ot	her c	auses		36 71 5	1,606 1,975 204	294 511 47	1,642 2,046 209	55 84 23	
	(a) Bubonic (b) Pneumonic (c) Septicæmic (d) Undefined						0				
		***	***	***	***	***	1		Inner I		
20. 21. 22.	Spirochætosis— Ictero-hæmorrhagica Leprosy Erysipelas Acute poliomyelitis Ergenhalitis lethaggies		,	· · · · · · · · · · · · · · · · · · ·	3	658 2	6 483 46 1	1 66 15 1	6 1,141 48 1	719 	
	Encephalitis lethargica Epidemic cerebro-spina					2	21	13	23	3	

TABLE 1-(cont.)

			ng is al at 1927	Yearly	total.	1868	ng is 1 a t 1926	
	Diseases.		*Remaining in Hospital at end of 1925.	Admis- sions.	Deaths.	† Total ca treated.	#Remaining in Hospital a t end of 1926.	Remarks.
	1885	a gland	345					
	I.—EPIDEMIC, ENDEMIC, AND INFECTI DISEASES—(cont.)	ous					and the same	
5.	Other epidemic diseases—							
	(a) Rubeola (German measles)			7	***	7		
	(b) Varicella (chicken-pox)	***	4	88		92	1	
	(c) Kala-azar (d) Phlebotomus fever			***				TBeneral
	(e) Dengue		1	66	***	67	0.222	
	(f) Epidemic dropsy (q) Yaws		10	228		238	9	
	(q) Yaws (k) Trypanosomiasis			220	***	200		army arminist.
	Glanders				***			
	Anthrax				***			
	Tetanus			76	57	76	1	TO (6)
	Mycosis				111		222	and and the
	Tuberculosis, pulmonary and laryngeal Tuberculosis of the meninges or	oontrol	183	2,139	992	2,322	177	
•	nervous system	centrar		10	8	10	***	
	Tuberculosis of the intestines or peri	toneum	1	22	14	23	***	- Sendar
	Tuberculosis of the vertebral column Tuberculosis of bones and joints	883	1 3	30	11	5 33	2 4	
		***		00	**	00	100	
i.	Tuberculosis of other organs—						- Im-	1-40
	(a) Skin or subcutaneous tissue (lup (b) Bones	1215	***	2		2	1	
	(c) Lymphatic system	0.83		23	1	23	1	
	(d) Genito-urinary			5	1	5		Don Hank
	(e) Other organs			5	3	5	***	
7.	Tuberculosis disseminated—							
	(a) Acute			3	3 22	3		
	(b) Chronic		2	39	22	41	1	
3.	Syphilis—	1.304						
	(a) Primary		40	722 781	15	762	51	
	(b) Secondary (c) Tertiary	1000	56 33	308	15 19	837 341	48	
	(d) Hereditary	1000	1	28	. 8	29	1	Epidemia in
	(e) Period not indicated			49	2	49	1	Transact C.
	A.—Gonorrhœa and its complications		54 71	648 1,435	4	702 1,506	12 54	
	B.—Gonorrheal ophthalmia		1	42	1	43	4	
	C.—Gonorrhœal arthritis		6	146	2	152	11	
	D.—Granuloma venereum			137	103	7	2	
	Septicæmia Other infectious diseases		3	417	22	137 420	5	
	Trypanosomiasis						2.2	(b) Part
	II.—General Diseases not Mention	NED					"Simula"	1000
	ABOVE.		13					Tellow Tier
3.	Cancer or other malignant tumours	of the	200.0					and the
	buccal cavity		2	26	11	28	3	Towns of
4.	Cancer or other malignant tumours stomach or liver		- XX	52	41	52	The state of the s	Talenda To
5.	Cancer or other malignant tumours			02	-	02	***	Meralpolan .
	peritoneum intestines, rectum			21	11	21	-	saling stands
6.	Cancer or other malignant tumours			99	8	24		The state of the state of
	female genital organs		1	23	0	24		and the same

Table 1—(cont.)

	BEET E THE STATE OF	g in	Yearly	total.	80 8	at at 26.	
H.—General Diseases not Mentioned Arove—(cont.)	Pi	pital of 19		1	ca, ted.	ining sital of 19	Dame of a
H.—General Diseases not Mentioned Arove—(cont.)	Diseases.	Host	nis.	ths.	otal	Hosp end	Remarks.
Anove—(cont.)		* 18	Adr	Dea	+ 1	++	
Anove—(cont.) 47. Cancer or other malignant tumours of the breast							
Anove—(cont.) 47. Cancer or other malignant tumours of the breast	II.—General Diseases not Mentioned						
As Cancer or other malignant tumours of the skin				TANKS			AND WALL
48. Cancer or other malignant tumours of the skin	47. Cancer or other malignant tumours of the						
Skin			9	1	9		- Harris I am
not specified	skin	2	25	8	27	4	186 BEL
50. Tumours non-malignant		7	41	14	48	3	Company of the Co
12 296 2 308 13	50. Tumours non-malignant		-				- Topologia in
53. Seurcy (including Barlow's disease)	FO CI : 1			2			California Cl. 115
127	53. Scurvy (including Barlow's disease)		7				on elstochal (49
56. Rickets 3 6 5 9 57. Diabetes (not including insipidus) 6 92 11 98 5 58. Anæmia—	FF Dani Lani		2010		100000000000000000000000000000000000000		an annuta 18
57. Diabetes (not including insipidus) 6 92 11 98 5 58. Anæmia — (a) Pernicious							District of the last
(a) Pernicious		6	92	11	98	5	manual D
(b) Other ansemias and chlorosis	58. Anæmia—	1		The last	HIPERIN .	In the last	College and the last
59. Diseases of the pituitary body 60. Diseases of the thyroid gland— (a) Exophthalmic goitre				1111111111			Vietni ne
60. Diseases of the thyroid gland— (a) Exophthalmic goitre (b) Other diseases of the thyroid gland, myxodema 61. Diseases of the para-thyroid glands 62. Diseases of the supra-renal glands 63. Diseases of the supra-renal glands 64. Diseases of the supra-renal glands 65. Leuksemia— (a) Leuksemia (b) Hodgkin's disease (b) Hodgkin's disease (lead, mercury, etc.)				10000	12000		Damient 2
(a) Exophthalmic goitre (b) Other diseases of the thyroid gland, myxodema 61. Diseases of the para-thyroid glands 62. Diseases of the supra-renal glands 63. Diseases of the supra-renal glands 64. Diseases of the supra-renal glands 65. Leukæmia (a) Leukæmia (b) Hodgkin's disease (b) Hodgkin's disease (lead, mercury, etc.)		100			100		sold (a)
(b) Other diseases of the thyroid gland, myxedema			1		1	MIN THE	100.00
61. Diseases of the para-thyroid glands 62. Diseases of the thymus	(b) Other diseases of the thyroid gland,				- Alle	de la com	100 W
62. Diseases of the thymus		1000				***	6800 TO
64. Diseases of the spleen		0.00					Samuel Superior
65. Leukæmia— (a) Leukæmia (b) Hodgkin's disease							
(a) Leukaemia (b) Hodgkin's disease (b) Hodgkin's disease (c) 1	A STATE OF THE PARTY OF THE PAR		102	The state of	102	Section 2	wake-M.
(b) Hodgkin's disease			9	1	9	1442	
67. Chronic poisoning by mineral substances							dillogard of
(lead, mercury, etc.) 13 4 13 68. Chronic poisoning by organic substances (morphia, cocaine, etc.) 35 388 1 423 4 69. Other general diseases <td< td=""><td></td><td>1</td><td>40</td><td>1</td><td>41</td><td>1</td><td>total salas A 49</td></td<>		1	40	1	41	1	total salas A 49
68. Chronic poisoning by organic substances			13	4	13	(md) 70 at	min min ou
69. Other general diseases	68. Chronic poisoning by organic substances		000	-	100	1111111111	7117 (0)
Auto-intoxication	69. Other general diseases			82		I Total	6
Hæmophilia	Auto-intoxication		1	7999	1	bigge	12
Diabetes insipidus							wat (e)
AND ORGANS OF THE SENSES. 70. Encephalitis (not including encephalitis Lethargica)							In amount 10
AND ORGANS OF THE SENSES. 70. Encephalitis (not including encephalitis Lethargica)		10 /		1		Ball	1947 (a) Ang
70. Encephalitis (not including encephalitis Lethargica)	III.—Affections of the Nervous System						1013. (0)
Lethargica)	AND ORGANS OF THE SENSES.	110		The said	All The	State of the	a mella land 30
Lethargica)	70. Encephalitis (not including encephalitis	2000	8000		7000	and the section and	
meningitis or cerebro-spinal meningitis) 74 62 74 1 72. Locomotor ataxia 4 8 3 12 3 73. Other affections of the spinal cord 40 13 40 74. Apoplexy 7 5 7 (a) Hæmorrhage 2 34 17 36 3 (b) Embolism 1 1 1	Lethargica)	2	20	6	22		
72. Locomotor ataxia	meningitis or cerebro-spinal meningitis)		74	62		1	
74. Apoplexy	72. Locomotor ataxia		8	3	12	3	
(a) Hæmorrhage 2 34 17 36 3 (b) Embolism 1 1 1		1839		7-20	100	A COLUMN	and a more
	(a) Hæmorrhage		34	17	36		and property of
		10000		Jan St.	de la contra de		Tollies He . S.
	(c) Antoniossis in in in in						

TABLE 1-(cont.)

	ng ii	1925	Yearly	total.	nso	ng i	
Diseases.	temaini	Hospital at end of 1925.	Admis- sions.	Deaths.	† Total cases treated.	# Remaining in Hospital at end of 1926.	Remarks.
	-		Ac	Ď	+	#	4
					T TON IN		
III.—Affections of the Nervous System					Clerch		
AND ORGANS OF THE SENSES-(cont.)					or Internal		
. Paralysis—							
		86	170	35	206	36	
C. I C. I C. I		19	109	17	128	33	
Other forms of mantal alienation		7	13 539	7	13 546	16	
Dailana .		7	101	5	108	5	
. Eclampsia, convulsions (non-puerperal) 5 year		201				- 113 114	
T 0 .15 1.			6	2	6	1	
CAS		1	41	27	42	1	
A Transaction			17	100	17		
D Namitia		12	156	2	168	10	
		2	44	***	46	1	
. Cerebral softening			4	3	4	1	
Other affections of the nervous system, sue as paralysis agitans		9	420	17	429	8	
			420	1,	120		
. Affections of the organs of vision—						Charles and	
	***	94	340	3	434	61	
7.5 m. Shama		13 2	764 116	***	777 118	16 15	
(1) Thursday of the one			22		22	10	
(a) Other effections of the one		96	611	2	707	123	
5. Affections of the ear or mastoid sinus		6	293	4	299	6	
						and the state of	
IV Approximate or any Comment	101	111					
IV.—Affections of the Circulatory							
System.							
			24	. 21	24	Territorial Control	
		5	22	9	27	1	
Other diseases of the beaut	***	2	5 67	30	5 69		
(a) Valvular		6	68	30	74	3	
Mitral		18	227	92	245	14	
Aortic			35	12	35	1	
Pulmonavy	8500 DE					0.00	
(h) Myocarditie	***	2	28	3 22	7 30	of the same of	
	***	-	20	20	90	water and	Diabetes
. Diseases of the arteries—							
(b) Autonio colomocio		2	14	9	16	***	
(c) Other diseases	2000	1	12	3 4	12	2	
Embolism or thrombosis (non corobrel)			9	2	6 9	THE REAL PROPERTY.	988
. Diseases of the veins—			HITTE	Agrony	miller	1000	
H	12	10	222	100	****	and the	
Varionee voine		13	211	3	224	8	
Phlebitis			3 2	1	3 2.		STREET, S
Diseases of the lymphatic system	100		36	3	36	1	to the address
Lymphangitis		7	132		139.	2	
Lymphadenitis, bubo (non-specific)		44	560	- 4	604.	42	
fi. Other affections of the circulatory eveters			4		4.		
other amountains of the circulatory system	***	5	103	54	108.	144	

TABLE 1-(cont.)

					See 2	ng in l at 925.	Yearly	total.	cases d.	ng in 926.	
		Diseases.				* Remaining in Hospital at end of 1925.	Admis- sions,	Deaths.	+ Total ca treated.	‡ Remaining in Hospital at end of 1926.	Remarks.
W	A	on man Danner		· Cuon							
	-AFFECTIONS						1			and the same of	CT 170
31.	Diseases of th Adenoids			***	•••	***	1 9		1 9		and the same of th
	Polypus					"1	16		17		
	Rhinitis					î	17		18		street discrep
	Coryza		177			2	133		135		1000 (B)
98.	Affections of	he larynx—								edismaq !	100 00
	Laryngitis		0		20.3	1	48	2	49		Selbuonid T
99.	Bronchitis				100	7	180	8	187	5	ABREST AT
	(a) Acute					86	2,237	54	2,323	61	partition I at
	(b) Chronic					41	944	59	985	35	modificación de la constante d
	Broncho-pneu		***		***	40	579	307	619	17	appointed.
01.	Pneumonia (a) Lobar	***	***	***	***	50	85 1,739	54 766	1,789	41	AND THE PARTY OF T
	(b) Unclass	ified	***	***	***	31	484	245	515	13	The state of the s
02.	Pleurisy, emp					18	344	50	362	14	
	Congestion of						2		2		12 (c)
	Gangrene of t						36	27	36	1	SE 10 161
	Asthma		***	***		14	592	18	606	26	to Dilate Cal
	Pulmonary en		***	***			- 7		7		in the training
07.	Other affectio			***		14	557	15	571	8	
	Fullmonary :	spirochaetosis	***	****		***		1	1	***	
7	IDISEASES	OF THE DIGES	STIVE	SYSTE	M.						e boult
08.	A.—Diseases	of teeth or gu	ms		Sec.	***	3		3	navi sets	Germaniti III
	Caries, pyor				***	1	130	1	131	2	MALIEUTINES NO
		ections of the				***	2	2	2		Date Lant -17
	Stomatitis					2	122	7	124		
	Glossitis, et	Z		***		***	10		10	2	soff-file
09.	Affections of	the pharynx or	tonsil	s—					sancey.	OK) EN	178 · ·
	Tonsillitis					3	207	2	210	7	
	Pharyngitis						104		104	1	Sport Steelers 19
	Affections of										Division A 300
11.	A.—Ulcer of				***	4	60	14	64	2	MINE-II
10	Other affection	he duodenum				***	14	3	14	1	STREET, LI
12.	Gastritis			***	***	7	47 397	10	47	9	amiles S
	Dyspepsia, e	te	***	***		14	474	9	488	10	
				***	***	11	100		300	100	D PRODUCTION OF THE PARTY OF TH
13.	Diarrhœa and					-			. 37		and the same of th
	Under two y			***	***	17	424	69	441	8	to represent the
14.	Diarrhoea and		***	***	***	1	68	5	64	1	
	Two years a Colitis		***	***	***	60	1,998	270	2,053	64	
	Ulceration			***		5	16	8	124 16	- States	
14	Sprue				***	3	83	20	86	3	
	Ankylostomia					105	2,388	120	2,493	55	
	Diseases due t						FIRM			- 132	American de la companya del companya del companya de la companya d
	(a) Cestoda	DATE OF THE PARTY				1	-		0	1	
	(b) Tremate			***	***		5		6		DANIESTE.
		da (other than	ankyl	ostom	a)		2		2	***	30) mol 13
	Asca	ris				60	2,652	21	2,712	36	TO STREET OF
		ocephalus disp	ar	***			1		1	1	an pull
	Trieb	ina				***					CALCULATION OF
		anculus				1	17		18	1	
	Stron	gylus			***	***	2		2		
	Oxyu	ris									

TABLE 1 .- (cont.)

			Remaining in Hospital at end of 1925.	Yearly	y total.	808	#Remaining in Hospital at end of 1926.	
	Diseases.		spita d of	- 8	*	Total car treated.	pital of	Remarks.
			Rem Ho en	Admis- sions.	Deaths	Tota	Rem Hos	
		-		Y	Á	+	77	
							CHT 90 80	MITTERS A.
VI.—Dr	SEASES OF THE DIGESTIVE						Comments.	to many the t
	System-(cont.)	9.1						
iseases due to	intestinal parasites—(cont.)							and and
(d) Cocci		1						Corpen
(e) Other	parasites			15		15		
(f) Uncla		46.7	1 3	145		1		A CONTRACT
 Appendicit Hernia 		1690	10	145 208	19	148 218	6 24	Company of
	ons of the anus, fistula, etc.		4	97	1	101	4	100 E (10)
	affections of the intestines			79	7	79	***	(d)
Enteropt Constipat			9	628		687		al-dimension of
	w atrophy of the liver			020			6	
21. Hydatid of	the liver			1	1	1		Cart (A)
22. Cirrhosis o	100	***	4	124	41	128	11	S. Sentines 2
(a) Alcol (b) Othe			1 20	37 255	22	38		HOLIMAN DESCRIPTION OF THE PERSON OF THE PER
23. Biliary Cal				200	119	275	15	andre a
24. Other affec	tions of the liver		***	4	1	4	a olijon	Communical h
Abscess			5	117	27	122	7	THE PART T
Hepatitis Cholecyst			4	125 21	4	129	7	Stanisland.
Jaundice	itis		6	145	10	21 151	2	CANCEL TO
 Diseases of 	the pancreas							
	(of unknown cause)		4	69	52	73	1	Carrier 1
27. Other affect	tions of the digestive system	0	20	577	40	597	15	- IN-N
					199	111	1	Ditmooks
VII.—DISE	ses of the Genito-Urin.	ARY	16					- Indiana
Syst	EM (NON-VENEREAL).				allend	I WE THEN	edg side by	nerstadt 0
28. Acute nepl	ritis		31	433	123	464	20	indicate P
29. Chronic			20	405	161	425	26	CONTRACTOR OF
 A.—Chylu B.—Schist 		***	***			***		Total Land
	tions of the bidness		4	51		1122		700 M
Pyelitis,				50	9 8	55	2 2	to Dilleg affer to
32. Urinary ca	lculus		1	41	2	42	2	Description
33. Diseases of			3	13		16	1	
Cystitis			2	83	13	85	2	- marriage L
 Diseases of (a) Strice 	trivo		0	100		143	THE REAL PROPERTY.	The world of the Party
(b) Othe			2 5	102 195	3 7	104 200	3	ment of
 Diseases of 	the prostate			2		200	13	
Hypertro			***	1		1		Directority.
Prostatit	is		4	16		20		
of man		No. of the last of	21	228	7	249	-	
Epididyn			2	89		91	7	Demount of
Orchitis		5000	2	131		133	3	
Hydrocel Ulcer of	Ponis		1	84		85	5	mar fall
	other non-malignant tumo	ours of	4	22		26	1	
the ova	ries		2	16	1	18	1	
38. Salpingitis		2000		15	*****	15	settle.	
39. Uterine to	of the pelvis nours (non-malignant)			1	1	1	***	
o corine tu	morrhage (non-puerperal)			10	1	10		
 Uterine has 	morrange (non-buerberal)		111		****			

Table 1-(cont.)

			og in 1 at 1925.	Yearly	y total.	1808	ng in 1 at 926.	
Disea	ises.		Remaining in Hospital at end of 1925.	Admis- sions.	Deaths.	+Total cas treated.	## Hospital at end of 1926.	Remarks.
				4	A	+	++	
VII.—DISEASES OF T	HE GENITO-U	RINARY					MARKET START	
System (Non-Vi	ENEREAL)-(ce	mt.)	22	la late			To a married	
141. A.—Metritis	81 12			- 8		8	***	
B.—Other affections	of the fema	ale genita		100				
organs		*** **	. 17	278	14	295	5	
Displacements of u				19		19	***	
Amenorrhœa Dysmenorrhœa		***		19 28	***	19	1	
Leucorrhea				83	***	88	3	
				00	***	00	1.00	
142. Diseases of the breas	st (non-puerpe	ral)—	1 1910	Ul war	Side attack		But said	and the State of the
Mastitis		77	. 2	25	1	27	1	- (30m ·
Abscess of breast			. 1	15	417	16	***	
77777 75					mah u		MITORITA	
VIII.—PUERF	ERAL STATE.		1 1					
143. A.—Normal labour		***	89	1,915	5	2,004	78	
B.—Accidents of Pro	The state of the s	444 11		11	3	11	***	
(a) Abortion (b) Ectopic gestati		***		176	2	179	***	
(c) Other accident	on			5 96	1 22	5 98	2	
144. Puerperal hæmorrha		444 44		4	1	4	I selmonths	
45. Other accidents of pa				75	14	79	2	
 Puerperal septicæmia 				86	40	90	Military Inc.	
147. Phlegmasia dolens		***		4	12.	4	1	
148. Puerperal eclampsia				21	5	21	1	
149. Sequelæ of labour	of the booset	****		72	9	72		
150. Puerperal affections Pregnancy		***	1	10	***	10	9	
riegiandy			. 1	48	***	49	9	
IX.—Affections	OF THE SKIN	AND						
CELLULAR	Tissues.				ana sa -			
151. Gangrene	*** ***			41	21	48	4	
152. Boil				133		136		
Carbuncle 153. Abscess				1 200	0.77	1.000		And Assessed
Whitlow		***		1,899	37	1,980	84	
Cellulitis			00	686	62	725	46	S Herrie Chy
154. A.—Tinea	11 2			133		134	6	dio) accust of
B.—Scabies			. 28	1,101	1	1,129	41	boldsrolled 10
155. Other diseases of the	skin		. 245	4,368	52	4,613	266	Summan, 7
Brythema Urticaria	***	**** **	0	4	***	4		National Property
Eczema		***	3.01	66 493	1	68	1	Fi shamay 1
Herpes		:		79	***	509 79	12	Il abelia W. o
Psoriasis	021.			10		10	no sonociati	
Elephantiasis				28		30	2	(I) abono W. V
Myiasis				3	THE STATE OF	8	988	b) alumin V/
Chigos						****		DI SOLUCIONE AND
Cutaneous leishma Cutaneous leishma				285	1	243	7	and toward it
Outsideous ieisnima	masis dicers		. 88	1910	16	1,998	135	Daniel A
X.—Diseases of Bo	NES AND OPG	ANS OF					molt mi	mmH-H
				1	34	to resid to	MI INOL	
	LUDER		1				0	O STONES OF THE PARTY OF
LOCOMOTION (OTHER		***	. 1	5 32	1	6 32	3 3	Contribulg.
LOCOMOTION (OTHER 156. Diseases of bones								
LOCOMOTION (OTHER 156. Diseases of bones 'Osteitis	1 1							dortesti
LOCOMOTION (OTHER 156. Diseases of bones				2		2	1000	dorientili o
LOCOMOTION (OTHER 156. Diseases of bones Osteitis 157. Diseases of joints			35					dortestis company), to do strough

Table 1—(cont).


RETURN OF DISEASES AND DEATHS (IN-PATIENTS) FOR THE YEAR 1926—(cont.)

						-11	ining in pital at of 1925.	Yearly	total.	808	Remaining in Hospital at end of 1926.	
		n:					Remaining i Hospital a end of 192	1000	-	otal castreated.	inin sita of 1	D1
		Disea	ISOS.				na osp	Imis- sions.	i i	Total	nd nd	Remarks.
						- 28	Ber E	Admissions	Deaths	Tol	Re He	
		1						¥	ď	+	1 7	
											1 8	
		IMALF	ORMATIC	ONS.					MINUS	1000	N 10 1119	Augh- He
59.	Malforma						25	16	1	41	21	P. STATE
	Hydroce		***					2	1	2	***	intelled to
	Hypospa		***		***		***		***			10/02/09/50
	Spina bi	nda, etc.	***		***	****	***	3		3		- negro
	XII	-DISEASE	S OF IN	FANCY							L'ON STREET	Lancon of the land
160	Congenita						1	44	26	45	4	Designation
	Premature					***	7	177	146	177	2	dringali-
	Other affe							32	20	32	3	In Inches let
	Infant ne				months	000000000000000000000000000000000000000						
	over)	and the second second						10	6	10		THE REAL PROPERTY.
	*****							750			1	
		-AFFECTIO	NS OF	OLD A	GE.		100	0.50	00			HIN -
164.	Senility		***	***			68	356	99	424	107	minutes and
	Senile de	ementia						39	6	39	1	Winds - St.
XI	VAFFE	CTIONS PR	ODUCED	BY E	XTERNA	L			-		2.00%	COLD COL
		CAU										200
165	Suicide by	noisoning										The same of
	Corrosive			(lane		***	***	***		***		Superify (
	Suicide by					***	***	***		***		Lawrence Co.
	Suicide by				n			4	4	4		Hermond of
	Suicide by							1	1	1		Considerate .
	Suicide by		***					***		***		Li winners
	Suicide by					8		3		3		Consequents.
	Suicide by					***			***			maring.
173.	Suicide by	crushing	***	***		***		***	***			
	Other suic					***	***	111				
175.	Food pois		***		***	***	***	23	4	23		
170	Botulisn		***	***	***		***		***		***	
170.	Attacks of Snake bi					***	***	1	***	10		
	Insect b		***	***	***	***		12 14	1	12	2	
177	Other acci		nnings		***	***	1	43	6	44	1	
	Burns (by	44	omingo				i	99	4	100	8	
	Burns (ot		v fire)				2	110	12	112	4	
180.	Suffocation	a (accident	al)									
181.	Poisoning	by gas (ac	cidental)				1	***	1		
	Drowning					***						
183.	Wounds (by firearms	s, war e	xcepted)	***	1	26	2	27	1	
	Wounds (or stabb	ing ins	trumer	nts)	63	2,199	37	2,262	85	
	Wounds (***	***	35	1,270	15	1,305	34	
186.	Wounds (in mines or	r quarri	es)			14	612	4	626	11	
	Wounds (1	201	1	202	2	
100.	Wounds (flisted by	g., ranw	ay accie	lents, e	tc.)	5	100	3 3	105	4	
	Injuries in Wounds in				KICKS,		6	343		349	15	
-	Execution			the second second	nts	***	***	***		***		
	A.—Over				110	***		***				
	B.—Hung							2	1	2		
	Exposure		the state of the s						***			
	Exposure											
	Heatstro	ke					***	1	1	1	***	
	Sunstrol		***	***			***					
	Lightning							1		1	***	
	Electric sh		***	***	***	***		1		1	***	
197.	Murder by	nrearms	***	***					***	***		


Table 1—(cont.)

		Yearly Total.		90 90	ng in Il at 1926.	
Diseases.	Remaining in Hospital at end of 1925.	Admis- sions.	Deaths.	+ Total cases treated.	# Remaining in Hospital at end of 1926.	Remarks.
XIV.—Affections Produced by External Causes—(cont.) 198. Murder by cutting or stabbing instruments 199. Murder by other means	 2 43 119 	80 219 795 2,018	7 2 72 23 1	80 221 838 2,137 1	 5 2 70 65	
XV.—Ill-Defined Diseases. 204. Sudden death (cause unknown) 205. A.—Diseases not already specified or ill-defined Ascites Edema Asthenia Shock Hyperpyrexia B.—Malingering C.—Filariasis	 171 2 4 	4 4,027 61 22 104 14 39 14 1	4 18 15 27 8 12 	4 4,198 63 22 108 14 39 14	126 1 2 	
XVI.—DISEASES, THE TOTAL OF WHICH HAVE NOT CAUSED 10 DEATHS. Pyrexia of unknown origin Paraphinasis Toxaemia Opium habit N. D. D. and under observation Accompanying Rupture of spleen Sciatica	6 17 100 56 19 	466 482 3 1 1,481 1,743 1,600 1	1 26 1 7 1 	472 499 3 1 1,874 1,799 1,619	62 6 1 62 48 	
Total	5,064	122,268	9,178	127,332	5,116	

GENERAL SYSTEMIC & PREVENTABLE DISEASES TOTAL CASES - 130251



TOTAL DEATHS - 8512




ENERAL SYSTEMS OF THE PROPERTY OF THE LAST


TOTALOGESATEDES



INFECTIVE DISEASES TOTAL CASES - 84468



TOTAL DEATHS - 5591



LIBRARY 3 LIBRARY 1 4 NOV. 1927 E

TABLE 2

MEDICAL STAFF ON 31st DECEMBER, 1926.

- 1 Principal Medical Officer
- 4 Senior Medical Officers
- 3 Chief Surgeons
- 1 Ophthalmologist Surgeon
- 23 Medical Officers
- 8 Lady Medical Officers (including 2 Infant Welfare)
- 1 Personal Assistant to the Principal Medical Officer
- 12 Deputy Medical Officers
- 11 Assistant Medical Officers
- 38 Assistant Surgeons
- 19 Dressers, Special Grade
- 66 Dressers, Grade I
- 251 Dressers, Grade II
- 85 Dressers, Grade III
- 99 Probationers
- 3 Matrons, Grade I
- 4 Matrons, Grade II
- 40 European Sisters (including 6 Infant Welfare)
- 78 Asiatic Nurses
- 17 Native Midwives.

HEALTH BRANCH.

- 1 Senior Health Officer
- 11 Health Officers
- 2 Chief Sanitary Inspectors
- 4 Assistant Surgeons
- 5 Health Inspectors, Grade I
- 23 Health Inspectors, Grade II
- 1 Malaria Inspector.

INSTITUTE FOR MEDICAL RESEARCH.

- 1 Director
- 1 Bacteriologist
- 1 Pathologist
- 1 Assistant Pathologist
- 1 Chemist
- 3 Assistant Chemists
- 1 Malaria Research Officer
- 1 Research Student in Tropical Medicine
- 1 Assistant Surgeon
- 1 Librarian
- 1 Laboratory Assistant, Special Grade
- 4 Laboratory Assistants, Grade II
- 4 Laboratory Assistants, Grade III
- 4 Probationers.

CENTRAL MENTAL HOSPITAL.

- 1 Medical Superintendent
- J Assistant Medical Superintendent
- 1 Agricultural Officer
- 1 Senior Assistant Physician
- 1 Assistant Physician
- 1 Second Assistant Physician
- 1 Inspector
- 1 Assistant Inspector
- 2 Dressers, Grade II
- 2 Probationers
- 1 Matron
- 1 Work Mistress
- 2 Nurses
- 1 Steward
- 1 Storekeeper.

VETERINARY BRANCH.

- 5 Veterinary Surgeons
- 3 Assistant Veterinary Surgeons
- 10 Veterinary Inspectors
- 7 Veterinary Assistants

VENEREAL DISEASE BRANCH

- 1 Venereal Disease Specialist
- 1 Assistant Surgeon
- 1 Nurse
- 2 Dressers, Grade II (Dispensary)
- 1 Dresser, Grade III (Laboratory)

RADIOLOGICAL BRANCH.

- 1 Radiologist
- 1 Assistant Surgeon
- 1 Dresser, Grade II
- 1 Probationer.

ANNUAL REPORT OF THE INSTITUTE FOR MEDICAL RESEARCH FOR THE YEAR, 1926.

An attempt has been made during the year to improve the routine laboratory diagnosis of diphtheria and enteric fevers by the elaboration of the tests applied and by the correlation of the results with the clinical histories. This has been possible because the number of cases is not large; it has rendered the diagnosis more accurate and, at the same time, it has made the routine work more interesting and profitable. The system might be extended to the serum diagnosis of syphilis with advantage, but the number of Wassermann tests is large and it would entail an increase of staff.

Research has been undertaken in many directions. Among the problems immediately before us are the discovery of the vectors of the two forms of tropical typhus: the aetiology of the local form of the tsutsugamushi disease: the classification of the pathogenic leptospirae and the nosology of the leptospiral diseases of Malaya: the preparation of a bacteriophage active against dysentery bacilli of the Flexner group: and more important than all, comes the problem of malaria, which is the country's greatest scourge.

Details of the work which has been done during the year are given under the headings of the more important diseases.

MALARIA.

The price of quinine places it beyond the reach of most of the inhabitants of the vast malaria-stricken regions of the world. The places in which it can be produced successfully are limited, and, even were it sold at cost price many of those who need it would not have the money to buy it. An efficient and trustworthy substitute for quinine which could be produced cheaply in any well-equipped chemical factory would be of enormous benefit to mankind. Already, several preparations have been put on the market as substitutes for the cinchona alkaloids but they have proved useless. Chemists have tried for years to make a synthetic quinine, and recently a substance, closely allied to quinine, and which is said to be a powerful remedy for malaria, has been produced in the Bayer Works at Elberfeld. This substance is called "Plasmochin". In July, the makers were good enough to send us a consignment of the drug and, at the end of the year, Sir Ronald Ross gave us a further supply. We are indebted to Professor W. Nocht for putting us in communication with the Farbenindustrie Aktiengesellschaft at Leverkusen.

Plasmochin is described in the papers given to us by Sir Ronald Ross as an alkylamino-6-methoxy-quinolin salt, obtained by synthetic methods. A tasteless, light yellow, finely granular powder, fairly easily soluble in alcohol, soluble in water to 0.03 per cent. at 20°C. Readily converted by the hydrochloric acid of the stomach into the hydrochloride of plasmochin. It is put up in 0.02 gramme tablets (approximately one-third of a grain) each with a little groove across the middle so that it can be accurately broken in half, when one wishes to give doses such as 0.01 and 0.03 grammes.

The following doses are recommended in the treatment of benign tertain and quartan fevers: One tablet should be given three times a day; that is, 0.06 grammes or about one grain daily. After five days' treatment, there should be an interval of three days and then four days further treatment, and so on. "With due observance of these intervals the treatment may be continued for four or six weeks."

In the treatment of subtertian malaria, the administration of quinine and plasmochin together is recommended, because quinine acts more promptly on the schizonts while the plasmochin attacks the crescents. For this purpose the makers supply a sugar-coated tablet containing 0.01 gramme of plasmochin and 0.125 grammes of quinine sulphate (about one-sixth of a grain of plasmochin with two grains of quinine). The dose recommended is two of these tablets three times a day; that is, about one grain of plasmochin and twelve grains of quinine daily. This dose should be continued for one month without interruption; in the second month it should be reduced to one tablet three times a day.

Plasmochin is more poisonous than quinine. It should not be given on an empty stomach nor should it be administered to persons suffering from diseases of the heart or liver. "It should not be left to the patient to take the medicine himself but its administration should be always under the control of the nursing staff." Cyanosis occasionally appears in patients who are undergoing treatment. When it does so, treatment should be stopped until it has disappeared, which it does in a day or two without leaving any ill effects. Large doses may cause cramps and pains in the stomach. The maximum daily amount for an adult is three doses of two-and-a-half tablets: that is, 0.15 grammes or two-and-a-half grains in a day. Even 0.1 gramme

daily, occasionally causes cyanosis, and it is advisable not to exceed that amount. If convulsive pains of the stomach or cyanosis of the lips are noticed the plasmochin should be stopped, and not given again until the symptoms have completely disappeared.

Dr. W. Röhl, chief of the chemotherapeutic division of the Elberfeld Works, has carried out comparative experiments with quinine and plasmochin, respectively, in the treatment of bird-malaria due to infection with plasmodium relictum. He found that a dose of 1 c.cm. of a 1 in 800 solution of quinine was just sufficient to control the parasites, and that a stronger dose than 1 c.cm. of a 1 in 200 solution was poisonous for a bird weighing 20 grammes. When plasmochin was employed in place of quinine, Dr. Röhl found that 1 c.cm. of a 1 in 50,000 solution controlled the parasites and that the dose could be increased to 1 c.cm. of a 1 in 1,500 solution before symptoms of poisoning appeared. That is to say, plasmochin proved thirty times as powerful as quinine, and the effective dose was only one-thirtieth of the poisonous dose. The "working margin" of plasmochin being therefore 1 to 30 as compared with 1 to 4 in the case of quinine.

Professor P. Mühlens, of Hamburg, administered plasmochin to 134 patients who were suffering from chronic malaria. The parasites disappeared and relapses were rarer in tertian and quartan malaria than they are after treatment with quinine. In subtertian malaria, plasmochin proved less powerful than quinine in its action upon the schizonts, but Professor Mühlens states that it has a unique and most valuable property in this form of the disease, namely, that it destroys the gametocytes, which quinine cannot do. Professor Mühlens therefore recommends that quinine and plasmochin should be administered together in the treatment of subtertain malaria, because the quinine acts upon the schizonts while the plasmochin destroys the crescents and thus prevents the patient from infecting mosquitoes and so from being a source of danger to those around him. Plasmochin has also been used in the Balkans, in Italy and in Spain, where it was found as effective in the treatment of acute malaria as it had proved successful in the treatment of chronic relapsing cases in Hamburg.

The Treatment of Malaria in Kuala Lumpur with Plasmochin.—This inquiry is not yet complete; the action of the drug in subtertian malaria must be followed in a much larger number of cases before we can reach trustworthy conclusions in that respect, and the combined treatment with plasmochin and quinine remains to be tested, but a sufficient number of benign tertian cases has been treated to make it worth while presenting this interim report.

Our observations were made on forty-six patients who had been admitted, on account of malaria, to the District Hospital at Kuala Lumpur, and for this privilege we are indebted to Dr. J. P. Fitzpatrick, the Medical Officer in charge of that institution. Thirty-three of the patients were Tamils and thirteen were Chinese. They were all immigrant labourers, thirty had been less than one year in the Malay States and almost all of them were suffering from acute malaria. This series of forty-six patients comprised thirty-one cases of benign tertain fever, four subtertian, eight mixed tertian and three quartan. Neither the patients with benign tertian nor those with quartan malaria were selected in any way; but, in the subtertian and mixed-tertian groups, only mild cases were chosen for treatment.

It is not practicable to keep the ordinary Asiatic hospital patient under observation, and shielded from re-infection with malaria, for a period sufficiently long to gauge the efficiency of drugs in preventing relapses. The patients at the District Hospital in Kuala Lumpur are drawn from a wide area, only six of those treated with plasmochin came from the town or its suburbs. As soon as they considered themselves well enough, they insisted on leaving the hospital, there was no opportunity for further observation and, as most of them returned to the places where they had contracted malaria, there was every chance of re-infection. We were therefore obliged to limit our inquiry to the immediate effects of plasmochin upon the patient, the parasites and the fever.

Each patient was given a dose of one-and-a-half tablets, or 0.03 grammes, twice a day. That is, 0.06 grammes, or about one grain daily. The average weight of the patients was 48.2 kilogrammes (106 lbs.). This is about 70 per cent. of the average weight of a European. The dose of 0.06 grammes, which was given to our Asiatic patients, is equivalent to a dose of 0.085 grammes given to man weighing 70 kilogrammes.

Blood-films were stained with Leishman's stain and examined daily. The number of microscopic fields examined and the number of parasites were noted in each case; in the patients' records the result of each examination was written as a vulgar fraction with the number of parasites as the numerator and the number of fields

examined as the denominator. This is admittedly a rough method; but, as all the slides were prepared by the same hand, it is useful for purposes of comparison and for estimating the progress of the case.

A sample of urine was examined every day, and, at the same time an inquiry was made as to the presence of any toxic symptoms produced by the drug. In most instances the period of special observation and examination was limited to five days, but, in ten cases, this was extended to eight days or longer.

Plasmochin in Benign Tertian Malaria.—Thirty-one men with benign tertian infections were given one-and-a-half tablets of plasmochin, twice a day, with the result that none of them had a temperature over 100°F. after the third day of treatment, and, in every case, with the exception of a man who was suffering from bronchitis, it was normal after the fourth day.

The effect of plasmochin upon the parasites was equally striking. No parasites were found, later than the fourth day of treatment, in any of the thirty-one patients with the single exception of a man in whose blood one trophozoi'e was discovered in a thousand fields examined on the morning of the fifth day. In twenty-two of these men, no parasites were seen later than the third day.

It is evident from these results that the immediate action of plasmochin is quite equal to that of quinine.

Plasmochin in Quartan Malaria.—Only three cases of quartan malaria have been available for treatment up to the present time. Here again the action of plasmochin proved equal to that of quinine. The parasites disappeared after two days treatment, in one case; and after five days, in the other two. The temperature was normal after the fourth day in two instances; in the third case, there was slight fever until the sixth day, but this was not due to malaria.

Mixed-tertian Malaria.—Plasmochin was administered to eight men who were suffering from mixed infections with P. vivax and P. falciparum. The benign tertian parasites quickly disappeared but the subtertian parasites, on the contrary, did not. They were found in seven of the patients later than the fourth day, and, in three cases, as they were still present at the end of a week's treatment, we were obliged to give quinine.

Subtertian Malaria.—Only four cases of subtertian malaria have been treated with plasmochin. In one, there were only crescents; in the other three, trophozoites were present, and, in two of them, it became necessary to give quinine at the end of a week's treatment, because the fever continued the patients were getting worse, and parasites were still present in large numbers.

It is evident from these results that plasmochin, given alone, is not suitable for the treatment of subtertian and mixed tertian infections.

Subtertian Gametocytes.—Plasmochin was given to four patients with crescents in their blood. The first had an average of five crescents in a hundred field; none was found after three days' treatment. The second had an average of six crescents; none was found after five days. The third had twenty-five crescents; none was found after four days. The fourth had thirty-five crescents in a hundred fields; they gradually disappeared and none could be found after the seventh day of treatment, though thick and thin films were searched for five days.

It would be wrong to base conclusions upon so small a number of cases; but the results, such as they are, support the claim that plasmochin destroys the gametocytes of P. falciparum.

Toxic Symptoms Due to Plasmochin.—Plasmochin is not unpleasant to take because it is tasteless and the dose is small. No toxic symptoms or by-effects were observed. There was no instance of nausea, vomiting, diarrhoea, deafness, vertigo, amaurosis, tremor or headache due to the drug. The urine was examined daily. Albumen was present in the urine of six patients when they were admitted to hospital; this cleared up as the malaria parasites disappeared, except in two of the quartan cases. The plasmochin did not cause albuminuria. It is difficult to detect a slight degree of cyanosis in people with very dark skins. One man became slightly cyanosed; but this was not necessarily due to the plasmochin, as he was very ill with subtertian malaria. Records of blood-pressure were made during the course of treatment in thirty-seven cases but there was no evidence that the plasmochin had any effect upon it.

Objections to Plasmochin.—The makers state that plasmochin is more poisonous than quinine, consequently it would not be safe to supply it (as quinine is supplied in the Malay States) to police stations and to Malay head-men for free distribution.

Plasmochin is as powerful as quinine, in its action upon the parasites of benign tertian and quartan malaria. It is far more pleasant to take than quinine; but there is the drawback that its administration must be controlled by microscopical examination during treatment. Many malarial fevers are due to mixed benign and subtertian infections. In these cases the subtertian parasites may escape notice at the beginning of treatment and there is the risk that they may multiply without detection, while the patient is taking plasmochin, unless the blood is examined repeatedly during the course of treatment. It is therefore not a suitable remedy for outpatients who live at a distance from a hospital where their blood can be examined.

TABLE I.
BENIGN TERTIAN MALARIA.

Showing the Last Day of Treatment on which Fever and Parasites were Present in Each Case.

Case number.	wh	st day nich fev oresent	er wl	Last day on nich parasites present.		Case number.		ast day which few present	er whi	ast day ch para presen	sites
1		3	1000	4		26		1	1000	4	
2		2		2		27		4		1	
3		- 4		4		28		2		2	
7	9.95	1		2	b Galler	29		1	***	1	
8	· Leve	1		2	20 100	31		3		4	
9		2		1	September 1	32		8	1449.	3	
10	1100	3	***	2	30 000	34		1		1	
11		1		2	od Hira	35		8	***	4	
14		4		4		36		5		5	
15	***	2		3		40		1	***	2	
16		1		3		45		1		4	
18		1		4		46		1		1	
19		2		2	19 1111	48		1		3	
28		4		3		49		2	444	3	
25		1		2	O. Invited	50	***	2		2	

TABLE II.

QUARTAN, SUBTERTIAN AND MIXED TERTIAN MALARIA.

Showing the Last Day of Treatment on which Fever and Parasites were Present in Each Case.

Case number.		Type of disease.		st day fever.	Last day parasites found.
4	***	Quartan	- **	6	2
42	257	THOU THE PARTY IN		2	5
44				4	5
5	***	Subtertian	1	6	6 Quinine
13		**	and di	1	2
17			414	6	6 ,,
20		,,	***	0	7 Crescents only
12		Mixed tertian		3	7 Quinine
21		**	***	8	7 ,,
24		"		3	5
33		"	1000	8	8 ,,
88		"		2	4
39			***	9	9 ,,
48		"		4	6
47	***		***	2	5

(The word "Quinine" after a number, means that plasmochin was discontinued and quinine was given instead.)

THE TSUTSUGAMUSHI DISEASE.

The tsutsugamushi disease or kedani fever, is a continued fever which follows the bite of trombicula akamushi, and some closely allied species of mites which convey the virus from rodents to man. It is characterized by a small ulcer covered with a black crust, which appears at the place where the mite has bitten; this is followed by painful swelling of the nearest group of lymphatic glands, a typhus like cruption and high continued fever of two or three weeks duration. The disease is endemic in Japan and Formosa, and also in Sumatra. Dowden has recorded a suspected case in the Federated Malay States, Lagrange has described a typical case

in Annam, and Faust has reported two cases from the central Yangtse Valley. With the exception of the case reported by Dowden, it has not been recognised, hitherto, in any British Colony or Protectorate.

The presence of tsutsugamushi or kedani fever on the east coast of Sumatra is of particular interest to us in the Malay Peninsula, because it is only about twelve hours distant across the Straits of Malacca. The disease was first recognised in Sumatra by Schüffner, who described it under the title of "Pseudotyphoid. A variety of Japanese kedani fever". During the year 1908 he saw 158 cases on the estates of the Senembah Company; and Walch reports that more than three hundred cases were diagnosed in the Company's hospital between 1908 and 1923.

In 1923 there was an outbreak of the disease on a group of rubber estates belonging to the Goodyear Tyre Company, and Keukenschrijver reported that it was particularly prevalent on an estate which was not properly weeded, during the slump in the price of rubber, but was allowed to become over-run with tall grass (lalang) and undergrowth, which served as a home for rats. Many of the coolies who were employed in clearing away the undergrowth, became infected. As the clearing progressed the number of casualties declined; in 1923 there were ninety-three cases, in 1924 there were sixty-four, and during the first half of 1925 there were only twelve. These estates are now clean-weeded and Keukenschrijver is confident that the disease will disappear as the hiding places of the rats have been destroyed. Walch is of opinion that the ordinary cover-crops on rubber estates are not likely to be dangerous, because they do not provide suitable accommodation for the rats which build their nests in undergrowth and lalang.

The Actiology.—The actiology of the disease has been investigated with great thoroughness by Japanese workers. Miyijima and Kawamura demonstrated some years ago, that the reservoirs of infection in the tsutsugamushi of Japan are certain rodents, particularly a vole, microtus montebelli. Hatori has found that, in Formosa, the common house-rat of the country, M. rattus rufescens, is the home of the virus; while the small Malay house-rat, M. concolor, and a field-rat M. diardii, have been incriminated by Walch and Keukenschrijver in Sumatra. The infected rat does not become ill, as a rat with plague does. It remains apparently healthy but its spleen is enlarged. A rat inoculated with the blood of a man suffering from tsutsugamushi shows no symptoms of illness, but its spleen increases to nearly double its original size.

Large numbers of minute mites infect the rodents which inhabit those spots where the tsutsugamushi disease is endemic; it is not uncommon to find fifty or more clustered together within a rat's ear. These mites are the larvae of trombiculae and closely resemble leptus autumnalis, the European harvest-mite. Many of the mites, such as trombicula pseudo-akamushi, are harmless to man and the dangerous species are not the same in Sumatra as in Japan. In Sumatra the chief vector of the disease is T. deliensis; in Japan and Formosa it is T. akamushi. On an estate in Sumatra, where cases of tsutsugamushi were occurring, Walch found that 50 per cent. of the rats harboured the mite, T. deliensis, but on healthy estates, less than 3 per cent. were infested. The adult and nymphal forms of these small acarines do not attack animals, but feed on the juices of reeds and plants. They are found under fallen leaves and decayed vegetation. In Japan, they burrow into the earth, in cold weather, and are captured by digging up the earth and mixing it with water, when they float to the surface. The larvae have been found on almost all the animals which go into the places where they are lurking, rodents, shrews, dogs, cats, goats and buffaloes. Birds possibly play on important part in disseminating the disease by transporting the mites over long distances. The trombiculae have been found on the breast, round the eyes and at the base of the beak, in birds of many species. Keukenschrijver examined a number of bubuts, or crow-pheasants (centropus javanicus), which frequent lalang and undergrowth in Malaya, with the result that he found the majority of them infested with T. deliensis. According to Walch the mites have been found on acrocephalus orientalis, a bird which migrates southward from Japan in the winter, and many possibly carry the disease to distant countries.

The trombiculae are of microscopic size; the adult form of the Japanese T. akamushi measures one millimetre (1/25 inch) in length, and the larva, which attacks man and animals, is only 0.2 mm. (1/125 inch) long, which is less than the ordinary itch-mite, acarus scabei. The T. deliensis of Sumatra is smaller still.

The larval mites feed only once; when they have fed on an infected host they do not carry the infection to a fresh animal as, fleas carry plague or mosquitoes carry malaria; but while they go through their metamorphosis, as nymphs and as egg-laying adults, they still contain the virus, as Nagayo has demonstrated by injecting their crushed bodies into monkeys. The adult mites do not feed on animals, but they pass on the virus, through their eggs, to the next generation of larvae and these transmit it to the hosts on which they fed.

The virus of the tsutsugamushi disease will not pass through an N. or a V. Berkefeld filter, but its nature has not been finally determined. The resemblance of the disease to typhus and to the spotted fever of the Rocky Mountains has led many observers to search for Rickettsia, and Nagayo with his colleagues at the Government Institute for Infectious Diseases in Tokyo, has demonstrated Rikettsia-like bodies, not only in the skin and lymphatic glands of men and monkeys, but also within the larval mites themselves.

We have had but little opportunity, so far, of investigating the aetiology of the disease in the Federated Malay States. Mr. Lesslar, the Assistant Pathologist, has examined a number of local rats (M. griseventer and M. concolor) and has found that a large proportion of them harbour clusters of trombiculae within their ears. The species of these mites has not yet been determined; they are probably T. pseudo-akamushi. The examination of the mites carried by animals and rodents in country districts will be undertaken when circumstances permit; and also an investigation of the tungau, a small mite which frequently attacks Malay children by burrowing into their skin.

Tsutsugamushi in Selangor.—During the year 1926, four European planters, who were suffering from tsutsugamushi, have been seen in consultation with Dr. J. W. Field at the European Hospital in Kuala Lumpur. Three of these men came from an oil-palm plantation where they had been working in an area overgrown with weeds and long grass, on the edge of virgin jungle. The fourth patient was a rubber planter. He was taken ill on his return from Sumatra where he had been inspecting some estates on the east coast, but it is probable that he became infected in the Malay States prior to his visit to the Dutch Colony.

One of these men died on the fifteenth day of his illness, the other three recovered, but they were all seriously ill. The first patient to be admitted was not seen by us until he was convalescent and no primary ulcer was noticed in his case. The symptoms in the other three cases were typical of the disease. The onset was gradual, with two or three days prodromal weariness. In each of them, there was a characteristic primary ulcer covered with a black scab and surrounded by a dusky areola. In two cases, the ulcer was situated on the back of the leg; in one, it was on the front of the arm. The nearest lymphatic glands were much enlarged and tender, forming the primary bubo. The bubo subsided as the temperature rose, but the black scab did not separate until the end of the fever. Discomfort in the lower part of the abdomen was a prominent early symptom. A typhus-like rash appeared on the trunk about the fifth day and spread to the limbs later on. There was no rash on the face. The face was congested and the eyes were injected. The temperature rose rapidly but the pulse remained slow until the beginning of the second week. Nervous symptoms, such as headache, restlessness, sleeplessness, delirium and incontinence, were prominent in all cases.

In one instance, the fever lasted for twenty-six days, with a pseudo-crisis at the end of the second week; in another, the temperature became normal on the twentieth day, and, in a third, on the twenty-first day; the fourth patient died from heart failure on the fifteenth day.

The results of inoculating guinea-pigs, a gibbon, and a Macacus monkey were negative. There was leucopenia at the commencement of the fever. The Widal and Weil-Felix reactions were negative.

A full description of these four cases together with temperature charts and photographs will be published in Bulletin No. 1 of the Institute for Medical Research in 1927. Details of one case are given here as an illustration. The patient in question was a muscular European, thirty-six years of age. On July 23rd, he noticed a small insect-bite, a little blister, on the back of his right calf. He suspected what was wrong because he had been reading about tsutsugamushi in a medical book, while his assistant was in hospital suffering from it. Three days later, when a bubo appeared and he felt out of sorts on getting up in the morning, he had little doubt that he had become infected. On July 27th, he had headache with rigors, fever and vomiting, and he was admitted to hospital on the following day with a temperature of 103°F, and a pulse rate of only 76. His eyes were injected and his face was very flushed. There was no rash, no general enlargement of the spleen. The chief complaint at this stage, as in the other cases, was discomfort and pain across the lower part of the abdomen.

When he was admitted to hospital there was a typical primary ulcer, in the middle of the right calf, covered with a black scab and surrounded with a red areola. The primary bubo was situated in the right groin and consisted of the enlarged and tender inguinal and femoral glands. The black scab increased a little in size, up to the tenth day, and did not separate until the twenty-sixth day of illness, when it fell off and left a small, deep pit with a base covered by dried serum.

There was no rash on admission; it began to come out on July 29th, the fourth day. There were then about thirty slightly-raised, discrete, rose-spots on the chest and abdomen, and the same number on the back. They measured about 5 mm. in diameter and looked like rather large mosquito bites. By the next morning, the rash had become quite thick; the rose-macules of the preceding day were darker in colour and, between them, there were great numbers of small spots, more deeply situated, some of which had coalesced to form a subcuticular mottling. The rash was thickest on the trunk, and more scattered, on the arms and legs. At this time, there was no rash on the hands and feet but they were affected later, and, on the eighth day, the eruption was thicker on the knees and on the dorsal surface of the feet than on any part of the body. The rash had almost disappeared from the trunk by the twelfth day, but it was still prominent on the arms and legs, and in the palms of the hands. There were large irregular blotches on the knees, with smaller macules between them. The eruption had entirely disappeared by the sixteenth day.

The pain in the lower part of the abdomen, to which we have referred, did not last more than forty-eight hours after the patient came into hospital; but he was greatly distressed by frequent retching and vomiting, which began on the sixth day and lasted until the fourteenth. The constant retching, and the discomfort in the epigastrium prevented his sleeping and caused him great distress.

His pulse did not reach 100, until the seventh day, subsequently it became more rapid, and it was generally between 100 and 120 until the sixteenth day. Headache was severe during the second week, and there was slight delirium on the thirteenth and fourteenth days. The temperature was never very high except on the sixth day, when it rose to 104.2°F, otherwise it was not above 103°F. There was a kind of pseudo-crisis at the end of the second week; the temperature came down and the patient was much better, but on the sixteenth day it suddenly went up again and oscillated between 100°F, and 102°F, for four days. Finally it became normal on the twenty-seventh day of illness, and convalescence was uninterrupted. There was a characteristic leucopenia at the onset, with only 4,000 leucocytes in a cubic millimetre on the fourth day. A subsequent count, made on the eleventh day of illness, gave 9,000 cells, of which 85 per cent. were polymorphonuclear and 1 per cent. eosinophiles. The Weil-Felix reaction was negative with the indologenic strains of X 19 (No. 67 and Warshaw) of the National Collection of Type Cultures. The non-indologenic strain, Kingsbury, was not agglutinated on the tenth or fifteenth days, but on the twentieth and twenty-fourth days it was agglutinated in low dilutions. On the thirty-fourth and forty-first days the result of this reaction was negative again.

Prophylaxis and Treatment.—The most effective means of dealing with a nidus of infection on an estate appears to be by clearing the area of the weeds and undergrowth in which the rats live. The labourers who work at the clearing should have their bodies smeared with some repellant such as citronella oil and sulphur ointment or with naphthalene paste. Keukenschrijver has found a mixture of cajaput and tobacco-juice to be a good preventive. Japanese authors recommend that miteproof cotton clothing should be worn, but this is out of question in a hot country like Malaya. The two patients who were bitten on the leg were in the habit of wearing stockings and shorts. The loose texture of woollen stockings offers no obstacle to the passage of the mites and the fibres of wool are ideal things for the mites to cling to. Possibly the most practical prophylactic measure is to wash the body and the clothes thoroughly, immediately on leaving the infected area; in this way the mites which have attached themselves to the body or clothing may be got rid of before they fix themselves in the skin. The clothes should be soaked in 2 per cent. lysol or izal for thirty minutes. Carbolic soap may be employed to advantage. Nagayo states that, when a person is bitten, the onset of the disease may be checked by excising the part of the skin where the mite is found or by extracting the larva with a needle. After the ulcer has appeared and the fever has set in, treatment must be directed to the alleviation of symptoms, there is no specific remedy. Hatori recommends that the ulcer should be excised or cauterized.

TROPICAL TYPHUS.

Sixty cases of tropical typhus were diagnosed during the year; thirty-two belonging to the K. group and twenty-eight belonging to the W. group. Owing to the lack of staff it has not been possible to make any extensive investigations into the aetiology of this disease, though the occurrence of a number of cases on an estate twenty miles from Kuala Lumpur afforded a rare opportunity. Guinea-pigs were inoculated with blood drawn from three cases, on the fifth, the sixth and the eighth days, respectively; but in no instance was there any reaction.

Notes on tropical typhus have appeared in our annual report for the last two years, and three monographs on the same subject have been printed in the series of Bulletins from the Institute for Medical Research. It has been shown in these

publications that there are two groups of tropical typhus, distinguished from one another by the agglutination reaction of the patients' blood-serum with cultures of B. proteus X 19. In one group which we call the K. group, the serum agglutinates a non-indol-producing strain "Kingsbury"; in the other group, called the W. group, the serum agglutinates the ordinary strains of B. proteus X 19, such as "Warsaw" and "No. 67" of the National Collection of Type Cultures, which are producers of indol.

The two serological groups are epidemiologically distinct. Though tropical typhus is not contagious and does not occur in epidemics, it nevertheless has a patchy distribution and is prevalent in certain places. When a crop of cases occurs in a district they are all of one sort; that is to say, they belong to one group. In 1924 there was a small outbreak in a military camp at Sungei Besi, in which all the cases belonged to the K. group. Again during 1926, a number of cases occurred on an estate in Selangor and the examination of the blood of six of these patients showed that all belonged to the K. group. On the other hand, seven cases which occurred during the same year in the suburbs of Kuala Lumpur all belonged to the W. group. The W. group does not show the same tendency as the K. group, to limitation within certain areas and districts. The different distribution of the two groups indicates a distinct aetiology for each, and one may conjecture that they are not both conveyed to man by the same agent.

The tsutsugamushi disease bears a close resemblance to typhus, and in some cases of this disease there is a slight rise in the titre of the K. group agglutinins during convalescence; moreover, we have seen the two diseases occur together on the same plantation. In view of the prevalence of a form of tsutsugamushi (the "pseudotyphoid" of Schuffner) on the east coast of Sumatra, we have made a detailed search in all the cases of tropical typhus, which we have seen ourselves for the primary ulcer and bubo which are characteristic of the Sumatra disease, but they have not been present in a single instance.

During the past year we have seen tropical typhus and tsutsugamushi side by side in Kuala Lumpur, and we have no hesitation in saying that they are distinct and separate diseases. It is true that they have many points in common; for example, the rapid development of the fever, the prominence of nervous symptoms and the multiform, typhus-like rash; but, on the other hand, each disease has some peculiar characteristic features which distinguish it from the other, notably the primary ulcer in tsutsugamushi and the strongly positive Weil-Felix reaction in tropical typhus. The primary sore and the primary bubo are almost invariably present in the tsutsugamushi disease. Nagayo goes so far as to say that "these two symptoms are constantly present and can be regarded as pathognomonic". In the tsutsugamushi or pseudo-typhoid of Sumatra, Schuffner was able to find the primary sore in 79 per cent. of the native cases and in 100 per cent. of the Europeans. Hatori states that it can always be found if it is looked for carefully. In tropical typhus, on the contrary, there is neither ulcer nor bubo. Kenneth Maxy has recently published clinical observations on 209 cases of non-contagious endemic typhus in Alabama and Georgia, but in the none of these was a primary ulcer noticed.

The confusion in nosology is largely a confusion in language. In German-speaking countries the word typhus is employed in the designation of typhoid and relapsing fever as well as for typhus: a qualifying adjective is added to distinguish one from another. Thus, typhus abdominalis corresponds to typhoid, typhus recurrens to relapsing fever, and typhus exanthematicus to typhus. Schuffner who first described the tsutsugamushi disease of Sumatra, called it, when he was writing in English, "Pseudotyphoid, a variety of Japanese Kedani fever". He was careful to distinguish it from typhus and he wrote, that "its course corresponds in all respects to that seen in enteric fever.......This course sharply distinguishes the disease from typhus with its brusque onset and termination by crisis". In Dutch and German journals however, in which most of Schuffner's work was published, the name pseudotyphoid became pseudotyphus; and in English translations from these papers the word pseudotyphus has been retained and has led to the confusion of this disease with the mild, non-contagious tropical typhus.

A case of tropical typhus belonging to the W. group, which we saw in consultation with Dr. Field in September, 1926, at the European Hospital in Kuala Lumpur, is worthy of record because of its pronounced features. The patient was a merchant who lived in a large brick-built house on the outskirts of Kuala Lumpur. His illness began with headache and shivering, but he did not take to his bed until two days later. A comparatively gradual onset, such as this, is more common in the W. group than it is in the K. group where the invasion is usually rapid. The patient was admitted to hospital on the fourth day of his illness with headache, pain in the back and a temperature which rose to 104°F. in the evening. A macular rash appeared over the upper part of the chest on the sixth day, and subsequently spread to the arms and legs.

The most striking thing about the macules, when they first appeared, was their even spacing; they were not grouped or confluent. They were like mosquito-bites, and, on the eighth day, the patient looked as though he had been bitten all over the chest and back with regular intervals of about one inch between the bites. Subsequently, smaller spots appeared between the larger macules, and there was some coalescence, particularly on the back. The eruption was most prominent on the ninth day. There was never any rash on the face, but the whole of the rest of the body was invaded, including the palms of the hands and the soles of the feet. Fading was slow and there were still some dusky spots on the backs of the hands as late as the twenty-first day. Nervous symptoms were prominent. The mental condition of the patient, a man of naturally bright disposition, was much changed, he became querulous and resented all attention. He was sleepless and troubled by bad dreams in the earlier part of his illness and on the eleventh and twelfth days there was slight delirium. His lips, his hands and his voice were tremulous as early as the first week; in the third week he became very deaf; in the fourth week the right knee-jerk was lost while the left was slightly increased. At about the same time, there was a transient paresis of the left arm; he was unable to raise it from the bed for a day or two, but the power quickly returned during convalescence. There was profuse epistaxis on the tenth, eleventh, twelfth and thirteenth days, and a rather severe bronchopneumonia. The respirations were over forty for a week, and the patient's condition appeared grave. For the first two weeks the temperature reached 104°F. nearly every day; it then subsided gradually and by the twenty-fifth day it had become normal. Convalescence was rapid. The Weil-Felix reaction was made with B. proteus X 19, strain "Warsaw" of the National Collection of Type Cultures. The reaction was negative on the eighth day; positive at 1 in 250 on the twelfth day; at 1 in 2,500 on the twenty-first day and at 1 in 1,000 on the thirty-third day. There was no agglutination with the K. strain of B. proteus.

Bacillus Agglutinabilis. Strain U2.—Professor James Wilson of Belfast very kindly sent us a culture of a coliform organism which he has found to be agglutinated by the serum of people with typhus fever. He has named it bacillus agglutinabilis. Strain U2. The bacillus is a gram-negative, coliform organism, which grows well on ordinary media, without spreading in a proteus-like film. It is non-motile; it does not produce indol in peptone water. It forms acid within twenty-four hours in glucose and mannite, and a small bubble of gas in the latter after four days. Acid is formed in saccharose in ten days. Lactose and dulcite are not fermented.

An emulsion of B. agglutinabilis was put up with the blood-serum of twenty-one controls in Kuala Lumpur who were not suffering from typhus. Agglutination did not take place in dilutions above 1 in 60 except in one instance when it occurred at 1 in 120; this was a case of tsutsugamushi, a disease which is closely allied to typhus.

B. agglutinabilis is agglutinated by the blood of people with tropical typhus of the W. group, but not as strongly as B. proteus X 19. It was agglutinated by a dilution of 1 in 4,000 of a patient's serum collected on the twelfth day of illness. B. proteus X 19; strain Warsaw, was agglutinated at twice this titre.

B. agglutinabilis is not agglutinated by the blood of people with tropical typhus of the K. group. It was not agglutinated above a titre of 1 in 60 by a patient's serum which agglutinated B. proteus, strain Kingsbury, at 1 in 30,000 on the nineteenth day of illness.

The tests were made with emulsions of living organisms in Wright's pipettes, and the results were read after two hours at 37°C.

TABLE SHOWING AGGLUTINATION OF B. AGGLUTINABILIS IN TROPICAL TYPHUS.

				I.—Y	ISOP.	K. GRO	UP.			
Day of il B. proteus	lness. Kingsbur	(6) v 30		(10) 120		(13) 2,000		(16) 8,000	 30,000	 60,000
B. proteus										10
B. agglutina									 60	 60
			п.–	-PONN	AMPA	LAM W.	GRO	UP.		

 Day of illness.
 (7)
 (12)
 (15)
 (21)

 B. proteus Kingsbury ...
 10
 ...
 80
 ...
 ...

 B. proteus Warsaw ...
 1,000
 ...
 8,000
 ...
 15,000
 ...
 4,000

 B. agglutinabilis ...
 ...
 4,000
 ...
 4,000
 ...
 480

(The figures show the highest dilution at which agglutination occurred; e.g., the figure 8,000 means that agglutination occurred in a dilution of 1 in 8,000.)

Bacillus Proteus X 19. Strains Metz and Syrie.—Dr. D. J. Babtet of the Institut Pasteur de Hanoi, kindly sent us two strains of B. proteus X 19, named "Metz" and "Syrie", respectively. They had been supplied to him from the Pasteur Institut in Paris.

These two strains grow on agar as a spreading film. The organisms are gram-negative and motile. They produce indol in peptone water. They ferment saccharose, glucose and maltose with the production of acid and a bubble of gas. They are clearly members of van Loghem's indologenes group of B. proteus.

They are agglutinated by the blood of persons with tropical typhus of the W. group. Metz was agglutinated at a titre of 1 in 4,000 and Syrie at 1 in 15,000, in a case where the strain "Warsaw" of the National Collection, was agglutinated at 1 in 8,000 on the twelfth day.

The strains Metz and Syrie are not agglutinated by the blood serum in tropical typhus of the K. group. The blood of a patient which agglutinated the strain B. proteus Kingsbury at 1 in 30,000, on the nineteenth day, did not agglutinate the strains Metz and Syrie, which are apparently identical with the strains Warsaw, No 67 and Jerusalem of the National Collection of Type Cultures at the Lister Institute in London.

MELIOIDOSIS.

There remains much to be learnt about melioidosis, but each year sees some small addition to our knowledge of this fatal malady. In this report we have to record the discovery of its existence in Indo-China; its natural occurrence in a dog; and a case of infection through a superficial ulcer in a man.

Until this year, the disease has not been described as occurring outside the comparatively narrow limits of Burma and the Malay Peninsula. Recently it has been discovered, however, that melioidosis occurs in French Indo-China. In January, 1926, Dr. F. H. Guerin, Director of the Pasteur Institute at Saigon, very kindly sent us a culture of B. whitmori, the causative organism of this fatal malady, which had been isolated by Dr. R. Pons from the blood of an Annamite woman who died on the fourteenth of her illness. The biological characters and the pathological effects of this Saigon strain of B. whitmori were investigated in this laboratory, where it was found to be the most virulent strain of the organism from a human source, which has been encountered. In September we received another strain of B. whitmori from Saigon; it had been cultivated by Dr. Henry G. S. Morin from the blood of a European, who died shortly afterwards. The disease, in this instance, was so acute that there was little time for clinical observation. This is the second case, on record, of infection in a European.

In addition to human cases of the disease melioidosis has been found occurring naturally in the rat, rabbit, guinea-pig, cat and horse. During the current year, one more species was added to the list by the isolation of B. whitmori from the organs of a dog, dead of a distemper, which had been sent to the Institute for examination by Major S. L. Symonds, B.V.Sc.

Bacillus whitmori was isolated from four cases, all of them fatal, admitted to hospitals in Kuala Lumpur during the year. An account of one of these is given here because it is the only clear case of infection through a skin wound which we have met with. The patient, in question, was a Tamil labourer, thirty-six years of age. He worked on the railway and lived in a hut near the Batu River in Kuala Lumpur, in a district called Kampong Semarang, where there are many rats. His illness was said to have begun with fever and headache on November 20th, 1926, and he was admitted to the District Hospital four days later. He had an ulcer on the right leg and a suppurating bubo, in the right groin, which was opened. He was constipated, on admission to hospital, but for five days before he died he had diarrhoea.

We did not see this patient until after his death, which occurred on December, 9th. There was an ulcer on the right leg which was healing at the edges, and a bubo, in the right groin which, had been opened. The iliac glands on the same side were enlarged and suppurating. The spleen was enormous and full of large and small abscesses. There were minute haemorrhagic areas in the lungs. The liver was enlarged and contained innumerable small abscesses. B. whitmori was isolated from the ulcer on the right leg and from the pus in the femoral and iliac lymphatic glands on the same side. It was also cultivated from the abscesses in the liver and the spleen.

The main point of interest in this case was the demonstration of the bacillus in the ulcer on the leg which was evidently the point of inoculation; the infection having spread by way of the lymphatics to the femoral and lymphatic glands. In Bulletin No. 5 of this Institute, published in 1924, we stated that we had not met with a case in which there was any history or evidence of a primary local lesion. On the other hand, infection appears to have gained entry through a skin lesion in a large proportion of the cases of melioidosis observed by Whitmore, Knapp and Krishnaswamy in Rangoon. Many of their patients were morphia-injectors, and they suspected that the virus was carried by the syringe.

LEPTOSPIROSIS.

Leptospiras belong to the group of spiral organisms which includes spirochaeta and treponema. They are accepted as the cause of the following diseases, infectious jaundice (spirochaetosis icterohaemorrhagica), the seven-day fever of Japan (nanukayami), the autumn fever of Japan (akiyami A) and as the probable cause of yellow fever.

In the report of this Institute for the first six months of the year 1925, we announced that the presence of leptospirosis in the Malay Peninsula, which had been suspected for a long time, had at length, been proved by the cultivation of leptospiras from the blood of a patient in the Malay Hospital at Kuala Lumpur, in the month of April, and by the passage of the organism through a series of guinea-pigs. Three additional cases were met with before the close of the year.

During 1926 the energy of the staff has been directed, so far as routine duties have permitted, to the investigation of this disease and more especially to determine if leptospiras are the cause of one disease in this country or of several.

The organisms have been isolated from nine patients in Kuala Lumpur during the year 1926, and in addition there has been a number of cases of severe jaundice, many of them fatal, from which we have not been successful in cultivating them. Several similar cases have occurred recently in the neighbourhood of Ipoh and Dr. P. H. Hennessy has sent us guinea-pigs inoculated with the blood of some of them but none of the animals succumbed to infection.

Dr. H. Noguchi of the Rockefeller Institute, Professor Inada of the Imperial University, Tokio, and Dr. C. C. Okell, of the Wellcome Research Laboratories, have kindly sent us strains of leptospiras for comparison with those isolated locally.

We are still at the beginning of our work, but, from what has been accomplished so far, it appears that there are several diseases in this country which are due to leptospiras, and that infectious jaundice, seven-day fever and autumn fever all occur here in addition to another unclassified group. We have been unable to distinguish, by serological means, the leptospirae of infectious jaundice from the leptospira icteroides which Noguchi considers to be the cause of yellow fever.

MEASLES.

Dr. A. N. Kingsbury has investigated the prophylactic action of convalescent serum in measles and he reports as follows:

The use of convalescent serum as a prophylactic during epidemics of measles was mentioned in the previous report. Groups of children numbering 93, 132, 50 and 120 were given 2.0, 2.5, 3.0 and 3.5 c.cms. of serum, respectively. During the seven weeks following inoculation the incidence of measles in these groups was 15 per cent., 4.5 per cent., 14 per cent. and 1 per cent. Cases continued among adults and uninoculated children after the last cases in the inoculated groups so that opportunity for infection was not lacking.

Prophylactic serum is generally believed to confer a passive immunity for about two to three months. Secondary outbreaks of measles on some of the rubber estates resulted in a much lower incidence among inoculated than among non-inoculated children although an interval of over four months had elapsed from the date of the original treatment. Also the clinical course in inoculated children was milder than in the controls.

PNEUMONIA.

In July and August there was an outbreak of pneumonia in the police-barracks at Kuala Lumpur in consequence of overcrowding. There were six deaths between August 3rd and September 4th. In addition to these fatalities, one or two of the recruits died in their homes, to which they had been taken by their parents who surreptitiously removed them from the hospital in the night, and took them away in bullock-carts under cover of darkness. Those who died were young Malay boys from distant villages and only one had more than four months service.

At the time of the outbreak, the number of recruits in the barracks was considerably in excess of the accommodation provided. In one badly ventilated, low-ceilinged dormitory, on the ground floor, seventy recruits were crowded into a space which should not have held more than forty-four. It was in this room that most of the cases occurred. The epidemic ceased when emergency measures were undertaken to obviate the overcrowding, by housing some of the men in schoolrooms and other buildings. Epidemics due to overcrowding have occurred in these barracks before—there have been at least two outbreaks of cerebro-spinal meningitis—and they will continue to occur until proper accommodation is provided for the recruits.

THE BACTERIOPHAGE IN BACILLARY DYSENTERY.

Dr. F. d'Herelle, Chief of the bacteriological service of the Sanitary Maritime and Quarantine Council of Egypt, has been good enough to send us a supply of his bacteriophage for the treatment of bacillary dysentery. He describes it as an ultravirus parasitic on bacteria. It has been employed with great success in some countries, notably in Brazil, where it is said that there was not a single death in the first ten thousand cases of dysentery treated at the Oswaldo Cruz Institute in Rio de Janeiro.

We were eager to put the bacteriophage to the test because bacillary dysentery is one of the most serious diseases in the Federated Malay States, and because no treatment adopted hitherto has had any apparent influence in curing or mitigating the disease amongst the poorer classes who are its principal victims.

The bulk of the dysentery in this country is due to infection with organisms of the Flexner group and we have investigated the action of the bacteriophage in this type of disease. The result of this investigation will be reported in Bulletin No. 3 of the Institute for Medical Research in 1927. The bacteriophage has not proved satisfactory and Dr. d'Herelle is about to undertake the preparation of a fresh sample from cultures of strains, isolated in Kuala Lumpur, which we have sent to Alexandria where he is working. Dr. d'Herelle is visiting India, shortly, and the Government of the Federated Malay States has invited him to visit this country at the same time.

The following is a summary of the work done in connexion with the bacteriophage during the past year.

Summary.—A supply of the bacteriophage was received from Dr. F. d'Herelle. The majority of the deaths from dysentery in the Malay States are due to organisms of the Flexner group: infections with Shiga's bacillus are uncommon. The bacteriophage was administered to twenty-two men suffering from dysentery caused by bacilli of the Flexner group. A bacteriological examination of their faeces was made every day. Dysentery bacilli persisted in the faeces of six patients for more than ten days after the commencement of treatment. In eleven cases, the bacilli were found up to the sixth day but not after the eighth. In the remaining five cases they were not found after the fourth day. Three patients died during treatment; healthly dysentery bacilli were isolated from their intestines after death. The organisms isolated from the faeces of these twenty-two patients, and from the intestines of the fatal cases, had the usual features of normal dysentery bacilli in appearance, growth, agglutination and carbohydrate reactions. The action of the bacteriophage upon Andrewes's type strains was tested in vitro. It was rather more virulent for these than for freshly isolated strains. When the bacteriophage was added to broth cultures of freshly isolated dysentery bacilli, a few organisms survived in each case, but the resulting "mixed cultures" had little vitality, as a rule; and they were generally non-agglutinable by specific sera. Only a single case of dysentery due to Shiga's bacillus was available for treatment with the bacteriophage. In this instance, no dysentery bacilli were isolated after the second day of treatment. The action of the bacteriophage upon Shiga's bacillus, in vitro, was much stronger than its action upon organisms of the Flexner group.

Conclusions: The sample of bacteriophage employed in the treatment of twenty-two cases of bacillary dysentery due to organisms of the Flexner group had no apparent influence either upon the course of the disease or upon the infecting bacilli.

(Cultures of the dysentery bacilli isolated from these patients have been forwarded to the Curator of the National Collection of Type Cultures at the Lister Institute, London).

DIPHTHERIA.

Diphtheria was almost unknown in the Federated Malay States before the war; but during the last decade it has become comparatively common. Thirty-five cases were diagnosed in the laboratory during the year 1926.

A special investigation of this disease has been made during the last two-and-ahalf years and a report will be issued as Bulletin No. 2 of the Institute for Medical Research in 1927. As the result of this investigation we are of opinion that:

- (1) Diphtheria has been endemic for many years.
- (2) Carriers are common.
- (8) The population is highly immune.
- (4) There is no evidence that the local strains of C. diphtheriae are less virulent than those isolated in other countries.

· The following is a summary of the report:

Summary.—The older practitioners in the Federated Malay States testify to the almost complete absence of diphtheria until recent years. Unconnected sporadic cases occurred at long intervals and four were recorded in Kuala Lumpur between the years 1904 and 1908. The disease has been more common since 1916, but cases are not numerous. It was difficult to explain why the sporadic cases never gave rise to an epidemic, and this investigation was undertaken, in the course of routine work, to determine if the population is as susceptible as it was supposed to be.

It is unusual for more than one case to occur in a family. In eighty-eight cases there were seventy-six instances in which no other member of the household became ill. There were sometimes many carriers in the house where the patient lived, and in the Malay States, as elsewhere, the common origin of the case is undoubtedly the carrier. The proportion of carriers to cases is an index of the immunity of the population. The results of the examination of school children and contacts suggest that diphtheria has been endemic for a long time and that a high degree of immunity is present.

The age incidence of diphtheria affords yet further proof of its endemicity and of the consequent immunity of the population. In an unprotected community all ages suffer, but adult cases are rare in the Asiatic population of the Malay States. They are more common in Europeans.

The bacteriological examination of contacts is of very little practical value, because of the great difficulty of keeping in touch with carriers.

The case mortality and the results of animal inoculations show that the toxigenicity of the local strains of diphtheria is not less than the toxigenicity of those isolated in other countries.

The results of Shick tests clinched the evidence that the population is highly immune to diphtheria. The proportion of susceptible children in two large day-schools attended by the children of better class parents is almost the same as in the mixed schools of New York. Tests which were made in two schools for Malay boys, showed that they were less susceptible than children of the same age in Europe and America.

Diphtheria is almost unknown among the labour forces employed on rubber estates and tin mines. A hundred adults were tested in an institution for decrepit Tamils, mostly from rubber estates, with the result that only two susceptible persons were found among them.

The immunity of the population is evidently of long standing. Possibly diphtheria has always been present among the Malays, but no outbreaks have come to official notice. The immunity may have been produced by diphtheritic inflammation of some part of the body other than the respiratory tract. Bacilli morphologically identical with C. diphtheria and having the same action on carbohydrates, are commonly found in the ulcers to which the labouring classes are susceptible, but the strains which we isolated were not toxigenic and there is no evidence that they were in any way connected with diphtheria.

ENTERIC FEVERS.

The number of blood-specimens sent to the Institute for Medical Research, to be examined for typhoid fever by the agglutination test, grows steadily larger, and the positive results increase year by year. On this account an investigation has been made during the two years 1925 and 1926 to determine if enteric fevers are prevalent in the Malay States and if they are becoming more common. It may be said at once that, judging from the results of this enquiry, typhoid fever is not common in the Federated Malay States and paratyphoid is rare. There has been no epidemic during the twenty-three years which have elapsed since the writer first came to the country, and Dr. G. A. Finlayson, who was Government Pathologist of Singapore, for over twenty years, states that the same is true of the Straits Settlements.

It is probably correct to attribute this freedom to the comparatively even distribution, throughout the year of a heavy rainfall which amounts to more than a hundred inches. The hot sunshine all the year round and the scarcity of flies doubtless play their part and, as far as the rivers are concerned, Dr. A. R. Wellington, the Senior Health Officer, has suggested that the colloidal clay suspended in their waters may be an important factor in their purification. Most of the towns and larger villages are fortunate in having excellent public supplies of water, drawn from uninhabited, jungle-clad hills. There are no large dairy-companies distributing milk collected from many sources and, in consequence, the carrier has less chance of disseminating bacilli here, than in a western city.

The statement that typhoid is uncommon, is justified not only by the absence of the disease in epidemic form, but also by the rarity with which it is found in the post-mortem room and by the scarcity of carriers. Typhoid ulcers have been found only three times in two thousand post-mortem examinations made by members of the Institute staff, during the last six years.

During the same period, that is to say, from 1921 to 1926, a bacteriological examination was made of 4,045 specimens of faeces, but only three carriers were found; giving a rate of 0.74 per cent. The great majority of the specimens were from the dysentery wards of hospitals in Kuala Lumpur and they were plated out on Harris and Teague's eosin methylene-blue agar within a short time after collection.

A carrier rate of less than 0.1 per cent. is extremely low; it may be mentioned, by way of contrast, that the examination of the general population, for typhoid carriers, in various districts of the United States, by means of a single examination of the faeces, gave an incidence of 0.3 to 0.8 per cent. S. W. Walch and his colleagues recently made a bacteriological survey of 1,076 healthy persons connected with the dairy industry in Alabama, and, by means of a single examination of the faeces they found eleven carriers, a rate of 1 per cent.

One of the three carriers found in the Malay States was a European, one a Chinese and one a Tamil. The European's faeces were sent for examination because he was suffering from dysentery. E. histolytica was found by microscopical examination and B. typhosus was isolated by culture. He had no symptoms of typhoid fever, nor had he ever suffered from that disease so far as he knew. Four specimens were examined in March, 1925, one in April, one in September and two in November, with positive results each time, except on one occasion in March. This man was still a carrier of typhoid bacilli at the end of 1926, but he is not known to have infected anyone else. As Walch states, "no outbreak can be traced to the majority of carriers, but it must be assumed for administrative purposes, that every person who harbours typhoid bacilli, is a potential producer of disease." If there were a test to determine the virulence of typhoid cultures it might be found that the organisms in many carrier cases, were harmless. Two urinary carriers in Walch's series, a man and his wife, had owned and worked a dairy for at least five years without, so far as could be ascertained, causing a single case of disease.

The population of the Federated Malay States is about one-and-a-half millions, and nearly one hundred thousand persons are treated in Government hospitals every year. Enteric fever was diagnosed by laboratory methods in eighty-seven cases during 1925 and in one-hundred-and-nine cases during 1926. These figures do not, of course, represent the total number of cases occurring in the whole of the Federated Malay States, but they doubtless include the majority of those admitted to Government hospitals. It is probable that at least one-tenth of the cases in the country come to our notice and that less than a thousand persons contract typhoid fever in a year, throughout the whole of the Federated Malay States.

When one speaks of a disease being common or rare in a country, one means that it is common or rare in comparison with its incidence in other countries, and, with respect to enteric fevers, it is illuminating to compare the Federated Malay States with the colony of Barbados. In the Annual Health Report for the year 1924-25, it is stated that "Barbados enjoys natural and climatic advantages, from a health standpoint, such as can scarcely be found in any other tropical place. The soil is well drained and the water supply is very good". The average number of cases of enteric fever notified during the last five years was 1,332 per annum. The population of the colony is entimated at 159,499 or about one-tenth of that of the Federated Malay States. Consequently, if enteric fever were as prevalent here as it is in the Barbados some thirteen thousand cases would occur every year.

The important question, whether typhoid fever is becoming more common in the Malay States, as indeed some people believe it is, cannot be answered satisfactorily except with the help of accurate statistics, and these are not available. The Senior Medical Officers in charge of the several States were kind enough to supply us with lists, of the cases of typhoid, sent to them from the Government hospitals, but the inaccuracy of these lists made them useless; many patients were omitted who had been shown by bacteriological examination to be infected with typhoid, and others were included who had been found free from the disease by repeated agglutination tests. Typhoid fever is not included in the list of notifiable diseases and consequently some Medical Officers do not notify their cases.

Though trustworthy statistics, giving the incidence of typhoid over a number of years, are not available, there is indirect evidence which shows that it is unlikely that the disease is becoming more prevalent; the bulk of the cases, proved by bacteriological examination, do not come from the large and growing towns, but from the outstations and villages, such as Kuala Pilah, where there is less change and where progress is not so rapid.

The Widal Test .- It was the custom in the past for Medical Officers, who sent specimens of blood to the laboratory for agglutination tests, to receive in return the bald information that the agglutination test was positive or negative, as the result of the examination of a single specimen. This was misleading and dangerous; in some cases where the agglutination was positive, this result was due to prophylactic inoculation or to a previous attack of typhoid; in others, the test was negative because the blood had been collected during the early days of the disease, before the agglutinins had appeared. During the last two years we have tried to make the laboratory diagnosis more accurate by obtaining at least three specimens of blood from each patient. The waxing and waning titre of agglutination obtained by the examination of several samples of blood, collected at intervals during the course of the disease, is a very valuable aid to diagnosis. The trustworthiness of the agglutination curve, obtained in this way, and the fallibility of a single test, are illustrated by the following example. The first specimen of blood, taken from a certain Tamil patient, gave a negative result, it had been collected on the eighth day of illness; the next sample, collected on the twelfth day, agglutinated B. typhosus in dilution of 1 in 240; on the eighteenth day agglutination took place at 1 in 2,000; on the twenty-fifth day at 1 in 3,000, and on the thirty-fourth day at 1 in 250. As a rule, the agglutinins appear towards the end of the second week, reach their maximum in the fourth week and then decline.

The Examination of Excreta.—We tried to obtain three specimens of faeces and urine from each case in which the agglutination test was positive and, in the majority of the cases, the Medical Officers sent the necessary material; but in a few instances repeated requests were unavailing. A sound judgment cannot be given unless the evidence has been carefully collected; the most important part in diagnosis is the provision of suitable specimens for examination and the correlation of the results obtained in the laboratory with the clinical history of the patient. The routine laboratory tests are comparatively simple.

The majority of the specimens were sent to us from a distance, and were not less than twenty-four hours old when they were examined. Experience has shown that it is useless to expect to find typhoid bacilli in material which has been kept for this length of time in a tropical climate, and, in order to overcome the difficulty, small sample of faeces were sent to the laboratory, by post, in tubes containing Teague and Clurman's solution (sodium chloride 0.6 gramme. Glycerine 30 cc. water 70 cc.). The results of this method were most satisfactory. The excreta were obtained, on one or more occasions, in 144 cases where the result of the agglutination test was positive, and in ninety-two, or 63 per cent., the infecting organism was isolated; from the faeces in seventy-six, from the urine in fourteen, and from both urine and faeces in two. All the specimens except thirty-three had been sent from a distance and were one or two days old before they were examined. The bacilli were isolated from the first specimen received in seventy-two cases; from the second specimen in sixteen; from the third, or a later specimen in five.

The Number of Positive Cases.—A positive diagnosis was made on firm grounds, by the procedure which we have described, in 196 cases during the two years 1925 and 1926. One hundred and eighty-two of them were suffering from typhoid fever, nine from paratyphoid A. and five from paratyphoid B. The diagnoses were based on the following results:

B. typhosus was isolated in	and the l	474 1100		83
B. paratyphosus A. was isolated in				
B. paratyphosus B. was isolated in	1			3
Typhoid was diagnosed from the a		curve in	***	56
Typhoid was diagnosed from a sir				
above 1/1,000 in				
Paratyphoid B. was diagnosed from	the agglutin	nation cu	rve in	2
Paratyphoid A. was diagnosed from	the agglutin	nation cu	rve in	3
				100
				196

The Incidence of the Disease.—As we have remarked before, the majority of the cases did not come from the big hospitals in the large towns, but from the smaller agricultural centres.

Number of	cases	from	Perak-						
Taiping	***		***	***	***	***	***	 ***	19
Ipoh			***	***				 ***	14
Rest of P	erak				***			 	37
									70

Number of cases from	Selang	or-						
Kuala Lumpur								33
Rest of Selangor	***			200				37
								70
Number of cases from	Negri	Sembil	an-					
Seremban	****	***	***			***	***	23
Kuala Pilah	n						***	14
Rest of Negri Semb	ilan	***		***				5
								42
Number of cases from	Pahan	g—						
Whole State		***			***	***		8
Number of cases from	Kedah							
Whole State					***			2

Dysentery of the Shiga type is more common in the clusters of native houses, known as kampongs, than it is in the towns, and the same is true of typhoid fever. The inhabitants of these kampongs are accustomed to take their drinking water from a shallow well or from a river flowing past the kampong; and where they drink they generally wash themselves and their clothes. Typhoid is not common among the Chinese who work in the tin mines or among the Tamil labourers on rubber estates.

The distribution of the disease among people of different races was as follows:

83
66
25
10
6
4
2
196

The figure for Malays is rather high in proportion to the number of them admitted to Government hospitals. Probably this is because typhoid is a disease of the kampong rather than the town.

The Mortality.—The case mortality was about the same as in other countries; not more than twenty-six cases or 13 per cent. were fatal, and this is in spite of the fact that nursing, as nursing is understood in Europe, is hardly practicable in the wards of a native hospital. An Assistant Surgeon naively reported concerning one of his typhoid patients "he is delirious and leaves the bed and roams about".

Comparative Table for the Two Yea	RS 19	25-19	26.	110
B. typhosus isolated	***	34		49
Typhoid diagnosed by agglutination		46		58
B. paratyphosus A. isolated		4		2
Paratyphoid A. diagnosed by agglutination	***	-		3
Paratyphosus B. isolated		2	***	1
Paratyphoid B. diagnosed by agglutination		1	***	1
		87		109

PASTEUR INSTITUTE.

A department of the Institute was opened in August, 1924, for the prophylactic treatment of rabies. Carbolized vaccine containing 1 per cent. of nervous tissue in 0.5 per cent. carbolic saline is employed for the treatment of human cases. A full course consists of fourteen daily injections of 5 c.cm. A prophylactic vaccine is prepared for dogs which consists of 20 parts of nervous tissue, 48 parts of glycerine and 32 parts of a 1.25 per cent. solution of phenol in salt-solution. The treatment

consists of a single injection of 6 cubic centimetres. Two strains of fixed virus are employed in the preparation of the vaccine. One of these, the "C" strain, was obtained from the Pasteur Institute at Coonoor and originally came from the Pasteur Institute in Paris; the other, strain "L", was obtained from Colombo.

Dr. A. N. Kingsbury was in charge of the Pasteur Institute for the first four months of the year; since April, Dr. Richard Green has carried on the work with Assistant Surgeon K. Kanagarayer as his assistant. Dr. Green reports as follows:

Treatment of Patients.—During 1926, 183 persons applied for treatment, an increase of sixty over the previous year. One hundred and twenty of them received the full course of fourteen injections; in the remaining cases, nine of whom underwent an incomplete course, it was ascertained that the suspected animals, with which the patients had been in contact, were not suffering from rabies.

Sixty-two, of the 129 cases which were treated, had been bitten by dogs proved by laboratory tests to have been rabid or certified as rabid by Medical or Veterinary Officers; thirty-four had recent abrasions contaminated by saliva from proved rabid animals and thirty-two had been bitten by dogs that were regarded as rabid but which could not be traced. A Malay girl was bitten by a monkey which was killed later. Smears from the monkey's brain showed the presence of Negri bodies, and the girl received a course of treatment. A kitten was diagnosed as suffering from rabies by the Pathologist, Penang, and two Europeans, who were in prolonged and dangerous contact with the kitten, received a full course of treatment.

Rabies in Malacca.—It was decided, early in the year, to establish a branch treatment centre in Malacca, on account of the number of cases of rabies occurring there, and the Acting Pathologist, Malacca, came to the Institute for a course of instruction.

A full course of injections was given to five Europeans, fifteen Chinese and two Indians in Malacca; that is, twenty-two cases apart from those already mentioned as having been treated at the Institute for Medical Research, making a total of 142 cases.

Table I shows the cases grouped according to race and domicile. Table II shows the cases treated at Malacca after the establishment of the local centre there. Fifty-eight per cent. of the patients, who received a full course of treatment, were sent from the State of Kedah; only 22 per cent. belonged to the Federated Malay States and most of these were from Negri Sembilan.

PASTEUR INSTITUTE. TABLE I.

* Cases Grouped According to Domicile and Nationality.

1926 (12 months).	Nationalit	y.	Selangor.	Perak.	Negri Sembilan.	Singapore.	Penang.	Malacea.	Kedah.	Total
Complete	European Eurasian		"	1	3		2	2	4) alaut
course of	Malay	0.0			8	100	1220		16	120
treatment /	Chinese		5		1				24	1
of the same	Indian		4		10	***	***	1	39	1
ncomplete	European Eurasian									Y
course of	Malay	***	***		1		***	***	***	5 9
treatment /	Chinese	***	2		i		***	111	1	(3
	Indian		8				***	***	î	1
rail or how	European Eurasian		1						70 6	1
dvice only	Malay	***	2 3		***	1 60000	***	****		54
dvice only	Chinese	***	17	***	***	-1000	***	***		(39
	Indian		81					***)
otal for 1926	od al.		68	1	24		2	3	85	188

TABLE II.

CASES TREATED IN MALACCA, 1926

	Nationality.			Number ises trea		Total.
	European	 		5		
Complete course of	Eurasian	 				
Complete course of treatment	Malay	 		***		22
treatment	Chinese	 		15		
	Indian	 	111	2	/	

Results of Treatment.-Twenty-nine of the patients were bitten by animals proved to be rabid by laboratory tests, and thirty-four had recent abrasions probably contaminated with the saliva of such animals. None of these patients developed rabies. The close co-operation of the Veterinary Surgeons with the Pasteur Institute has been of the greatest value in tracing cases bitten by dogs, securing material for diagnosis and in keeping records.

TABLE III.

SHOWING RESULTS OF TREATMENT DURING 1926 AT PASTEUR INSTITUTE.

Position of bite.	I,	п.	ш	IV,	v.	VL.	Total.	Number of deaths during or after treatment.
Face Limbs and trunk unclothed Limbs and trunk through	1 20	5 24		1 32	1 22		8 98	in which has
elothing	8	4	***	1	9	1	23	milwesten
Total	29	33		34	32	1	129	Nine received only a partia course of treatment

- Cases bitten by animals proved rabid by laboratory examination.
 Cases bitten by animals certified rabid by Medical or Veterinary Officer. No laboratory tests.
- III. Cases bitten by animals almost certainly rabid according to history. No laboratory tests.
- IV. Cases with recent abrasions contaminated with the saliva of proved rabid animals.
- V. Cases bitten by animals which were untraceable.
- VI. Cases having abrasions contaminated with saliva of animals in which laboratory tests were negative.

TABLE IV.

SHOWING RESULTS OF TREATMENT IN MALACCA CASES.

Position bite.		I.	II.	III.	IV.	Total.	Deaths.
Face Limbs and trunk unclothed Limbs and trunk clothed	 	1 2 2	2	12	3	1 19 2	= = = = = = = = = = = = = = = = = = = =

- I. Cases bitten by dogs proved rabid by laboratory examination.
- II. Cases bitten by dogs almost certainly rabid according to history. No laboratory tests.
- III. Cases which had recent abrasions contaminated with the saliva of dogs proved rabid by laboratory examination.
- VI. Cases which had been in contact with a case of human rabies.

Case of Human Rabies .- Only one case of human rabies came to our notice during the year. This was a Chinese in Malacca who had received no prophylactic treatment. Full details are not available but there seems to be no doubt about the correctness of the diagnosis. The history was briefly as follows: Swee Tan, a male Chinese, age 42, was bitten by a dog in China about November 1st, 1926. The dog was said to have been mad. About twenty-five days later the patient arrived in Singapore and soon afterwards went to Malacca where he became very ill, and was admitted to hospital in Malacca. His temperature on admission was 100°F. and the pulse rate 120. His most striking symptoms were excessive salivation and recurrent spasms of the pharynx. Food, drink or a sudden stimulus excited the pharyageal spasms and a diagnosis of rabies was made on clinical grounds. The patient died on the 1st December, 1926, about one month after having been bitten. A post-mortem examination was not obtainable.

Rabies in Dogs.—The brains of ninety-four dogs were submitted to the Pasteur Institute for diagnosis. Forty-eight contained Negri bodies in fresh smear preparations, and the diagnosis was confirmed by the examination of stained sections. Forty-six of the specimens were negative.

Table V.

Examination of Brains for Negri Bodies, 1926.

Solumen	ocame Born		remk.	N Sambilian		D	T CHANG.		Malseca.	Kadah	Actuals.	Total positive.	Total negative.	Grand total.
+	_ 2	+	- 1	+	- 13	+	- 1	+	13	+ 22	- 16	48	46	94

Prophylactic Inoculation of Dogs.—A vaccine for the prophylactic inoculation of dogs is prepared at the Institute, and four hundred doses were supplied to the Veterinary Officers in various parts of Malaya during 1926.

In Negri Sembilan 270 dogs were inoculated, in Kedah 82, in Selangor 23, in Malacca 16, and in Penang 9. Approximately four hundred dogs thus received protective inoculations. One apparent failure was reported from Negri Sembilan in the case of a dog which was vaccinated and which developed rabies; the diagnosis being confirmed, subsequently, by microscopic examination of the animal's brain. The period from the date of inoculation to the onset of symptoms was about three months. As there was some doubt as to the identification of the dog, the failure of the vaccine is recorded as apparent only. Some of the dogs vaccinated have been quarantined for a period of six months after inoculation, in these no abnormal symptoms have developed.

MORBID HISTOLOGY.

Dr. R. Green who was responsible for this work during the last eight months of the year reports as follows:

Sections of 195 specimens from various hospitals in the Federated Malay States were cut and examined. Details of the benign and malignant tumours are given in the following tables:

TABLE I.
BENIGN TUMOURS

No.	Nationality.	Serial No.	Sex.	Age.	Site of tumour.	Nature of tumour,
1	Chinese	12	F.	40	Nasal cavity	Fibroma
2	,,	156	M.	50	Brain	Glioma
3	,,	51	F.	38	Vulva	Papilloma
4		60	F.	2	Sacrum	War Annua
4 5		69	M.	30	Tongue	Papilloma
	,,	92	F.	33	Uterus	Tillian manage
7		111	F.	37	Breast	Adams Charma
6 7 8	,,	157	M.	_	Paraspinal region	Dermoid cyst
9		160	M.	36	Neck	Soft fibroma
10		169	F.	23	Lip	Cavernous lymph-angioma
11	Tamil	28	M.	40	Peri-articular region	
12		41	M.	33	Skin	37 11
13		173	M.	25	Axilla	TT
14		177	M.	25	Gums	47.7
15		191	M.	18	Tarsal conjunctiva	Polypus
16	Malay	20	M.		Ear	Dilmone
17		77	M.	26	Axilla	Tibuana
18		119	F.	52	Alveolar margin	TT
19	Burmese		F.	20	Ovary	m
20	Eurasian	86	M.	30	Inguinal canal	Dilmone
21	.,	100	F.	23	Alveolar	1001
22	European	24	F.	40	Breast	Distance of January 1
23	n	87	F.	28	,,	
24		192	M.	29	Penis	Davillama

TABLE II.

MALIGNANT TUMOURS

No.	Nationality.	Serial No.	Sex.	Age.	Site of tumour.	Nature of tumour.
1	Chinese	4	M.	61	Liver	. Secondary carcinoma
2	,,	8	M.	50	Chaol	Epithelioma
3		16	M.	26	Anne	STATE OF THE PARTY
4	,,	17	M.	32	Stomach	. Adeno-carcinoma
5	1000	22	M.	24	Livor	Dairouw on wain own
6	" …	26	M.	53	Thyroid	Maliament namillana
7		27	M.	38	Testis	O. marrie
8	,,	81	M.	59	Kidney	Hamanaahaan
9		32	M.	39	Neck	Mined colled someone
10	,,	33	M.	80	No site given	Almoslan samanna
11		37	M.		Femur	Client colled assessment
12	,,	43	M.	40	Rectum	Adone cominens
13	,,	44	M.	54	Stomach	Caminama
14		45	M.	43	Liver	Consulant consistent
15		46	M.	51	Cheek	Daithaliana
16		48	M.	56	Lungs, liver an	
	"				enlaan	Endathaliama
17		54	M.	48	Stomach	Caminomo
18	,,	58	M.	87	Tonsils and cervica	- I Declare the second
10	" "	28			odon d	0.11
19		66	M.	54	Neck	M: 3 11 1
20	,,	70	M.	45	Donatid	Th. 3-41-11
21	"	82	M.	50	Retro-peritonea	
	** ***	0.00		00	olanda	Turninka announce
22		88	M.	50	Andela	Daithallana
23	10	84	M.	52	Ossonhaeme	Commence - 11-3
24	***	89	M	48	Rootum	Carainama
25	,,	65	M.	52	Livor	Drimanus consistences
26		98	M.	57	Orbit	
27	"	106	F.	57	Value	
28	"	110	M.	49	Mosle	
29	,,	115	M.	44	Limon	
30	" …	117	M.	48	Nach	Clause and a second
31	27	118	M.	58	Naca phanna	
32	" …	120	M.	45	Pania	Delate Come
2000		121	M.	44		Epithelioma
33	** ***	97	M.	34	Parotid	. Endothelima
34	"	126	F.	46	Abdominal wall	
35	,,	128	M.	34	T	Fibro-sarcoma .
36	,,	131	F.	11	Liver	
37	" …					
38	" …	135	M.	41	Neels	
39	" …	137	M.	43	Neck	
40	" …	143	M.	60	Perineum	
41	"	150	M.	47	Pleura	
42	" …	153	M.	45	Bowel	
43	"	158	M.	56	Kidney	
44	" …	165	M.	50	Lung	Secondary sarcoma
45	"	167	M.	32	Parotid region	
46		170	F.	52	Cervix uteri	
47		178	M.	46	Cervical glands	
48	,,	181	M.	56	Pylorus	
49	,,	183	M.	45	Liver	
50		184	M.	38	Cervical glands	
51	,,	185	M.	49	Ulcer of neck	
52	,,	190	M.	34	Skin	Epithelioma
53	m ",	195	M.	32	Penis	
	Tamil	7	M.	30	Anus	
55	,,	19	M.	48	Penis	The related to
56	,,	21	M.	28	.,	
57	,,	29	M.	28	.,	
58	,,	39	M.	30	Bones	Periosteal sarcoma
59	,,	55	F.		Lower jaw	Sarcoma
30	,,	57	M.	40	Bowel	Adeno-carcinoma
31	,,	59	M.		Lung	Duimanu accusa
32	,,	67	M.		Lower jaw	Sarcoma
33		71	M.	38	Right foot	Mixed celled sarcoma

TABLE II-(cont.).

			M	LALIG	NANT]	UMOURS-	-(cont.).			
Case No.	Nationality.	Serial No.	Sex.	Age.	Si	te of tumo	ur.		Nature	of tumour.	021
64	Tamil	72	F.	35	Tongu			Epithe	lioma		
65	Tamii	80	F.	55		al region		Epitale	noma		
66	,,	106	M.	55	Penis			,,,			
67	,,	122	M.	35	Cheek			,,			
68 69		127 139	F. M.	50 55	"	***		Saman	one cal	lad as wine	200.0
70	" …	159	M.	45	Liver				ry carci	led carcino)ma.
71	,,	188	M.	30	Rectu			Carcin			
72	Malay	25	F.	50	Lip			Epithe	elioma		
73	"	35 50	F.	40 65	Cervix	on arm		Camain	ama		
75	,,	74	F.	50		axillary		Carcin	-carcine	oma.	
76	European	3	M.	41	Lower			Epithe			
77	,,	101	M.	-	Neck			",			
-			100000	I .				1			
Ver	EREAL DISE	Lepe		ROU	TINE E	EXAMINA	TIONS				
1 53	Wasserman		071.8-					1925.	1	1926.	
	Positiv					A 19111111		2,174		1,717	
	Negati							4.685		4,404	
	Treponema							100			
	Positiv							1		3	
	Negati							5		10	
	Gonococcus										
	Positiv							6	10 000	1	
	Negati							20		12	
Wm	AL REACTIO	* * * * * * * * * * * * * * * * * * * *									
	Positive for		iosus	Son B	el u			302		197	
		B. para						3		4	
		B. para						17		10	
	Negative							814		1,033	
	IL-FELIX R	EACTIONS						-		1,259	
CERI	EBRO-SPINAL	FLUID.	-								
	Positive for			us				9		20	
	,, ,,	D.,						12	****	17	
	,, ,,	D						1		2	
		B. tube						_		2	
	Negative							39		21	
Сно	LERA										
	Positive							176	***	156	
	Negative							518		536	
PLAC	GUE.—										
	Negative	4						12		2	
DIPE	THERIA.										
	Positive	d maseru	1420	114.				42	00220	99	
	Negative							292		401	
LEPI	ROSY.—										
	Positive						***	31	***	41	
	Negative							60		61	
	TERIOLOGICA										
	Positive for					r		106		250	
	" "	-			Shiga			-		8	
		B. typi						34		49	
	11 11							4		2	
	" "				B		***	2	***	1	
	11 11				11			91	***	15	
	, ,, ,,	B. tube	rcule	1818	***	***	***	2	7	-	
	Negative	S ()	***	***	***	***		447	101	830	

BACTERIOLOGICAL E	XAMINATION	OF UR	INE.					
Positive for B	. typhosus			***		10	***	11
,, ,, B	. paratypho.	sus A.	***			-	***	1
Negative		***	***	***		214		308
AUTOGENOUS VACCI	NE	***				-		74
BLOOD-SUGAR ESTI	MATIONS		***	4847		-		170
LEPTOSPIROSIS.—								
Cultures-								
Positive		***	***	***	***	_	244	7
Negative			***	***	***	_	***	16
Urine—								
Positive						-	344	4
Negative	***		***	***		-	***	21
Rats.—								
Twenty-nine e		ad cult	ures	made	with			
Miscellaneous						6,704		7,048
* (The above	figures refe	er to th	he nu	mber	oj spe	cimens	exami	ned not to

* (The above figures refer to the number of specimens examined not to the number of cases.)

CHEMICAL LABORATORIES AND MALARIA BUREAU.

Reports on the work of the Chemical Laboratories by Mr. R. W. Blair, Chemistin charge, and on the work on the Malaria Bureau by Captain H. M. Pendlebury, Acting Malaria Research Officer, are appended.

PUBLICATIONS.

The following publications were issued from the Institute during the year:

(a) Studies from the Institute for Medical Research:

No. 20. "Notes on Malayan Culicidae," by A. T. Stanton.

(b) Bulletins from the Institute for Medical Research, 1925:

No. 1. "The Weil-Felix Reaction in Sporadic Tropical Typhus," by William Fletcher and J. E. Lesslar.

No. 2. "Tropical Typhus and Brill's Disease," by William Fletcher and J. E. Lesslar.

(c) Other Papers:

"On the Carbon Tetrachloride Treatment of Ankylostomiasis," by A. Neave Kingsbury. Transactions of the Royal Society of Tropical Medicine and Hygiene, 1926.

"Some Investigations of Malarial Fevers," by A. Neave Kingsbury.

Transactions of the Royal Society of Tropical Medicine and Hygiene,

"Three cases of Bronchial Spirochaetosis," by J. E. Lesslar and K. Kanagarayer. The Indian Medical Gazette, September, 1926.

"Tropical Typhus," by William Fletcher and J. E. Lesslar. The Indian Medical Gazette, November, 1926.

STAFF CHANGES.

Dr. A. Neave Kingsbury, the Pathologist, left in April to take up the appointment of Professor of Bacteriology in the College of Medicine at Singapore.

Dr. A. T. Stanton, the Director, retired at the end of June on being appointed Chief Medical Adviser to the Secretary of State for the Colonies.

Dr. William Fletcher, formerly the Bacteriologist, was appointed Director in August 11.

Captain K. B. Williamson left in September 24th on his appointment as Professor of Biology at the Medical College in Singapore. Since that time Captain H. M. Pendlebury, Systematic Entomologist in the Museum Department, has been in charge of the Malaria Bureau.

Mr. B. A. R. Gater, Assistant Entomologist, Department of Agriculture, was appointed a Malaria Research Officer on November 1st and went on furlough on December 18th.

Institute for Medical Research, Kuala Lumpur, January 31st, 1927. WILLIAM FLETCHER, Director of Government Laboratories, Federated Malay States.

ANNUAL REPORT OF THE MALARIA BUREAU FOR THE YEAR 1926.

STAFF.

Capt. K. B. Williamson, Malaria Research Officer, Federated Malay States, relinquished his appointment on 24th September, 1926, and assumed duties as Professor of Biology in King Edward VII Medical College at Singapore. Mr. B. A. R. Gater, succeeded him on 1st November, and proceeded on long leave on 17th December, 1926. During the intervening periods of the year Mr. H. M. Pendlebury was in charge of the Bureau.

Mr. Thamboo, the Second Assistant, returned from long leave and resumed duty on 5th October, 1926. Mr. P. Savery and Mr. M. Sewell, protationary assistants, were transferred to the Medical Department, Singapore, and to the Railway Department, Kuala Lumpur, respectively. Mr. P. V. Samuel was appointed probationary assistant on 1st July, 1926.

I have to report, with regret, the death of Mr. Mak Chuen, Third Assistant, which took place in Taiping Hospital on March 7th 1926, whilst he was engaged on rice field investigation work in Perak.

FIGURES RELATING TO FIELD AND LABORATORY WORK.

Breeding places described	1,428
Larvae identified microscopically	36,630
Adults bred out and identified	245
Adults caught in houses	
Adulta examined for malaria parasites	622
Blood specimens examined for malaria parasites	451
Spleens examined	550

REPORTS.

The following is a list of reports issued during the year:

- (i) Interim report on isvestigations in rice fields.
- (ii) Malaria investigations near Kuala Selangor.
- (iii) Investigation on the prevalence of flies in Port Dickson.
 - (iv) Breeding places of mosquitoes in the vicinity of the Residency of the Hon'ble British Resident, Negri Sembilan.
 - (v) Mosquitoes breeding in the Pedas-Rembau valley.
 - (vi) Mosquito survey made in the surroundings of Petaling Hill and Federal Hill, Kuala Lumpur.
 - (vii) Mosquito survey made in the surroundings of the European Hospital, Kuala Lumpur.
 - (viii) Comparative tests on the larvicidal and spreading powers of rubber, solar and crude oils.
 - (ix) Seventeen reports on specimens sent for identification.

EXHIBITIONS.

Anti-malarial exhibits relating to oiling, housing, and drainage, and living aquatic insects and fish, which destroy mosquito larvae, were shown at the Agri-Horticultural shows at Kuala Lumpur from 27-30 August, 1926, and at Teluk Datch, on 18 September, 1926.

INVESTIGATIONS IN RICE FIELDS.

Perak—Bukit Gantang Valley.—The investigation work in rice fields in this valley, which was started during 1925, was stopped during the year, and the temporary laboratory which had been established in the hospital dispensary building, Taiping, was closed.

The mosquito survey in the above valley showed that A. hyrcanus was the predominant species, 88 per cent.; and was associated with A. barbirostris 7.2 per cent. A. fuliginosus 4.7 per cent. and A. aconitus 0.1 per cent.

Adult mosquitoes to the number of 269 were caught in houses situated close to the valley and examined for malaria parasites, but gave negative results.

Kinta District.—At the time of investigation no larvae were found, apparently owing to the fact that all the rice fields were dried up, as it was just after the harvest. A few adults, numbering 90 specimens, were caught in houses in the district. These were examined for malaria parasites, but they all gave negative results.

Negri Sembilan.—The work on rice fields is Negri Sembilan was continued and periodical mosquito surveys were carried out in selected areas, in order to determine the distribution of Anopheles breeding during the different seasons of the year in the district. A party consisting of one probationary assistant and two collectors is stationed at Rembau, the rice field centre of the district, in order to watch any specific changes of Anopheles in a particular valley, and to collect adult mosquitoes from houses in the various kampongs adjacent to the rice fields and examine these for malaria parasites.

It was found, from the mosquito surveys in rice fields at different periods of the year, that A. fuliginosus was always present, often in large numbers, so it would appear that this species breeds in the rice fields of Negri Sembilan throughout the year in spite of the change of conditions which takes place at the various seasons. Then again, A. aconitus, one of the chief associates with A. fuliginosus was found to be prevalent just before, and at the time of, the harvest; but decreases during fallows and entirely disappears by the time the seedlings are transplanted to the rice fields. It is interesting to note that this species begins to reappear when the rice is about four months old, and when the water in rice fields becomes clear owing to fresh supplies either from the irrigation channels, or from adjacent streams.

I have to acknowledge, with many thanks, the assistance rendered by Mr. R. W. Blair, Government Chemist, Institute for Medical Research, and his staff, in analysing the various water samples submitted to him from Rembau-Pedas valley.

Out of 912 adults of various Anopheles species caught in houses in this area, 451 were dissected for malaria parasites, but the results were all negative. This work is being continued.

Of 395 Malay boys from four different schools situated close to the Pedas-Rembau valley, who were examined for spleens, 54 or 13.7 per cent. showed enlarged spleens and of 178 blood slides taken from 178 boys of the above schools 20 or 11.2 per cent. showed infection with quartan parasites including one with benign tertian.

TYPE COLLECTIONS.

Type collections illustrating the commoner species of anophelines have been sent to:

- 1. Health Officer, Pahang.
- 2. Health Officer, Kinta, Ipoh.
- 3. Health Officer, Krian, Parit Buntar.
- 4. Health Officer, Port, Singapore.
- 5. Medical Officer, Estates, Sungei Siput.
- 6. Medical Officer of Health, Department of Home Affairs, Tokyo, Japan.
- 7. Principal Medical Officer, Hongkong.
- 8. Director, Government Institute of Infectious Disease, Tokyo, Japan.
- 9. Director Pasteur Institute, Saigon, Indo-China.
- 10. Director of Public Health, Burma.
- 11. Director, Geneeskundig Laboratorium, Weltevreden, Java.
- 12. Director, Harcourt Butler Institute of Public Health, Rangoon, Burma.
- 13. Dr. Russel, Rural Sanitation Campaign, Malacca.
- 14. Dr. Malcolm Miller, School of Medicine, Edinburgh.
- 15. Sir Ronald Ross, through Sir Malcolm Watson, Klang.

LARVICIDAL PROPERTIES OF RUBBER OIL.

Rubber oil was tested for its spreading power and toxic efficiency on mosquito larvae in comparison to other oils, such as crude and solar.

Laboratory experiments with fresh samples of oils resulted in showing that rubber was the best and solar was better than crude oil. But rubber oil lost a great deal of its efficacy by exposure of a thin layer, for periods of seventeen hours and upwards, to air.

As fresh rubber oil was found to be best, it was decided to make mixtures of rubber and crude or rubber and solar, in certain proportions, and to test them on mosquito larvae in order to see whether the rubber oil would increase the toxicity of the other oils. A series of experiments shewed that rubber oil did not increase the toxicity of the other oils with which it was tested, until an admixture of 50 per cent. was added to each of them.

The tendency of rubber oil to produce very thin inefficient films, probably renders it unsuitable for use on the margin of large ponds, or in other places where the expansion of water is not limited; but it may prove very efficient if sprayed heavily upon pools, or small accumulations of water.

INFECTION EXPERIMENT ON A. SEPARATUS.

Larvae of A. separatus were submitted twice a week through the courtesy of the District Officer, Port Dickson, and adults were bred out in the laboratory for this experiment.

Of 112 adults that were placed to feed upon malaria infectives, 34, 27 and 8 took blood from subtertian, benign and quartan, respectively; and 20, 13 and 8, respectively were dissected, but none of them showed any infection, either in stomach or salivary glands. The average time intervening between feeding and dissection was fifteen days.

VISITORS.

Among the fifteen visitors who signed their names in the book during the year appears the name of Sir Ronald Ross, K.C.M.G., of the Ross Institute for Tropical Diseases, Putney Heath, London.

H. M. PENDLEBURY, Acting Malaria Research Officer.

31st January, 1927.

REPORT OF THE CHEMIST UPON THE WORK OF THE CHEMICAL LABORATORY, INSTITUTE FOR MEDICAL RESEARCH, FOR THE YEAR 1926.

The chemical work of the following departments is performed wholly or in part in the Chemical Laboratory: Medical, Trade and Customs, Police, Railway, Public Works and Forest Department.

The total number of samples examined in the course of the year was 7,756 as compared with 6,021 in the preceding year, an increase of 1,735. There are notable increases in the number of samples of water, toddy, chandu dross and coins. There has been a decrease in the number of samples of milk and liquors.

MEDICAL DEPARTMENT.

Chemical work is performed for the Medical Department in connexion with the Health Branch and the Hospital Branch. The samples submitted may be classified as follows:

- (1) Milk;
- (2) Water;
- (3) Sewage effluents;
- (4) Toddy;
- (5) Exhibits for toxicological analyses;
- (6) Miscellaneous articles.
- (1) Milk.—The following standards are prescribed for milk in the rules of "The Sale of Food and Drugs Enactment, 1913".
 - (a) The quantity of milk fat present in milk must not be less than 3.25 per cent. of the total component parts thereof;
 - (b) The quantity of milk solids, other than milk fat, present in milk must not be less than 8.5 per cent. of the total component parts thereof.

To ascertain whether these provisions are complied with, officers of the Health Branch collect samples for examination in this laboratory.

Five hundred and ninety-seven samples of milk were examined. Of these, 15 (2.5 per cent.) contained less than 3.25 per cent. of milk fat, and 63 (10.5 per cent.) contained less than 8.5 per cent. of milk solids other than milk fat.

The reductase test—a valuable method of determining whether milk is stale or not—was carried out on 64 samples of milk collected in Kuala Lumpur. Of these, 47 completely decolorized the methylene blue in three hours. The reductase test is as follows:

A stock solution is prepared by dissolving one part of methylene blue in 2,000 parts of water. Immediately prior to use one part of this solution is diluted with nine parts of water. One c.c. of the diluted solution is mixed with 10 c.cs. of the milk in a test-tube and then placed in a water bath kept at a temperature of 38°C. The proportion of methylene blue to milk is 1 to 200,000. The test has recently been included in the New Zealand Sale of Food and Drugs Act, and the clause is as follows: "When subjected to the reductase test, it shall not completely decolorize the methylene blue in less than three hours."

In addition, three samples of milk were examined bacteriologically.

(2) Water.—More interest is being taken in the provision and maintenance of pure water supplies, which is becoming one of the most important problems of the day.

Epidemic outbreaks of cholera, typhoid, dysentery, etc., are usually due to an infected water supply, and in the recent outbreak of cholera at Shanghai, suspicion fell upon the waterworks. Cholera Vibrios, the specific cause of cholera, were found to be present in the samples collected from the Intake works, the effluent from the filter beds, and from the taps from the mains. The prevention of cholera consists in municipal, domestic, and personal cleanliness and an uncontaminated water and food supply. Water plays an important part in the prevention of disease, and by providing a pure supply, the public are guarding against the possibility of water-borne diseases.

The jungle-streams of the Federated Malay States are the chief sources of supply, and are, apart from accidental contamination, generally speaking free from pollution. Every effort should be made to protect these supplies, by inspection of the catchment areas, filter beds, etc., and by bacteriological and chemical examinations of the raw and filtered waters.

Chemical examinations were carried out on 528 samples and bacteriological examinations on 99 samples of water as compared with 328 and 36, respectively during the year 1925.

The chemical examinations of the raw and filtered waters of the supplies for Kuala Lumpur numbered 259 and the averages for the year are shown in appendix B.

The averages for the years 1919 to 1926 are shown in appendices C, D, E and F, the highest results are underlined.

The raw waters are filtered through slow sand filters and the filtered waters passed into service reservoirs, the Maxwell's Hill Reservoir being open and the Weld Hill Reservoir covered. The averages differ slightly from those for the year 1925, although the rainfall for the year was lower.

The filtered waters entering the service reservoirs and the waters passing into supply after storage in these service reservoirs were examined chemically and bacteriologically daily during a period of 15 days.

The chlorination treatment of the Klang Water Supply has been continued during the year, and the raw and treated waters were examined daily during a period of 32 days. The quantity of water treated during the period under examination was 9.9 million gallons and the quantity of chlorine added 73.5 lbs. The average dose was 1 in 1.35 millions.

Arrangements were made for regular monthly examinations of the raw and filtered waters of the following supplies:

Parit Buntar, Bagan Serai, Seremban, Tanjong Malim Training College, Tapah, Taiping, Kuala Kubu, Rawang, Kajang, Teluk Anson, Gopeng, Ipoh, Batu Gajah, Kampar, Tanjong Rambutan and Kroh.

Prior to the year under review, these supplies were examined rarely or in most cases not at all.

- (3) Sewage Effluents.—During the year, various types of sewage installations have been erected in Selangor. Samples of the resulting effluents from some of these installations have been examined. The effluents in the majority of cases have failed to conform with the standards prescribed in the Report of the Royal Commission on Sewage Disposal.
- (4) Toddy.—Premises on which toddy is sold are open to inspection by officers of the Health Branch, who are also empowered to take samples for analysis.

The samples were examined to ascertain whether they complied with the standards prescribed in the rules of "The Sale of Food and Drugs Enactment, 1913", viz., (a) "toddy must not contain more than ten per centum of alcohol by volume or have an acidity exceeding 0.8 per centum expressed in terms of acetic acid."

The number of samples examined was 171. All the samples complied with the legal standards.

(5) Toxicological Analyses.—Thirteen exhibits were submitted by the Medical Department. These included viscera, foods and one medicine for suspected poison.

Human Poisoning.—Two viscers were received for examination, one of which contained a very small quantity of an alkaloid, which gave some of the reactions characteristic of nicotine, but was too small for further identification. In one case of opium poisoning, the stomach washings were found to contain opium. Two deaths occurred from gas poisoning in a well. A considerable proportion of carbon dioxide was found in the well. Specimens of blood from the two bodies were examined for carbon monoxide with negative results.

Animal Poisoning.—The viscera from five animals were examined for arsenic—arsenic being found in one case.

Miscellaneous Exhibits for Poison.—One sample of milk and one sample of chloroform were received for examination. No poison was detected in these exhibits.

(6) Miscellaneous Articles.—Five samples of urine were examined for sugar. Five medicines were examined, one of which was found to contain "ganja" (cannabis indica). Seven samples of mineral water were examined bacteriologically. Other samples included butter, dried milk powders, smudge sticks, citronella oil and santonin.

Mention may be made of an interesting sample of heavy red powder resembling mercuric iodide in appearance. This powder on analysis was found to consist of precipitated barium sulphate with a red dye adsorbed, and a small quantity of mercuric iodide.

The total number of articles examined was 37.

VITAMIN B EXTRACT.

The preparation of this extract from rice polishings, for the treatment of beri-beri, was continued throughout the year. During the year 8,606 fluid ounces were prepared and 8,960 fluid ounces were issued to medical practitioners, dispensaries and hospitals as compared with 6,056 fluid ounces prepared and 7,685 fluid ounces issued in 1925. One thousand five hundred and eighty-four fluid ounces were issued free, the remainder was sold at 25 cents per fluid ounce, this being the estimated cost of production.

In the year 1925, experiments with the above-mentioned extract were carried out by Dr. A. Neave Kingsbury, Pathologist, Institute for Medical Research. The results obtained indicate that the method of extraction is a highly efficient one. A new laboratory has been equipped for the preparation of this extract. With the previous apparatus, it was impossible to cope with all the demands.

As the subject of the extraction of vitamins is occupying the attention of many scientists, the methods of extraction of the vitamin B. from rice polishings are detailed below.

The present method differs very slightly from the original method as carried out by Fraser and Stanton in the year 1911.

(1). - ORIGINAL METHOD.

Polishings in quantities of 180 grms., being the amount required by twelve fowls in three days, were mixed with 1,000 c.c. of a 0.3 per cent. hydrochloric acid solution, stirred during the day and the following morning filtered through a Buchner's filter, 100 c.c. of 0.3 per cent. hydrochloric acid solution were used to wash out the vessels. When fluid could no longer be extracted from the mass, it was mixed with 600 c.c. of 0.3 per cent. hydrochloric acid solution stirred during two hours and thereafter filtered as before.

The extracted polishings were mixed with distilled water, nearly neutralized with sodium carbonate, and the volume adjusted to 1,080 c.c., 30 c.c. of this emulsion contained five grms. of polishings less the materials dissolved out by the acidulated water.

The combined filtrates obtained from 180 grms, of polishings were nearly neutralized with sodium carbonate and concentrated at a low temperature to a volume of 1,080 c.c., 30 c.c. of his suspension contained the substances solved out by acidulated water from 5 grms, of polishings.

(2).-PRESENT METHOD.

Rice polishings, preferably fresh from the mill and of the best quality, are sifted to remove portions of husk and broken rice.

The polishings are extracted with four times their weight of 20 per cent. alcohol containing 0.1 per cent. hydrochloric acid.

The extraction is conveniently carried out in upright cylindrical jars of about five litres capacity. The fluid is left in contact with the polishings for one week, the whole being well stirred daily to facilitate extraction.

Six jars are worked up daily, each containing:

lute are normed ab danti's can	 0			
Polishings	 	 	750	grms.
	 ***	 	600	c.cs.
Water	 	 	2,400	**
Pure hydrochloric acid	 	 	3	

The fluid is filtered off through Whatman No. 3 filter paper by means of large Buchner funnels and water suction pumps. The residue is pressed to secure a maximum yield of filtrate.

The filtrate obtained from each jar is about 2,500 c.cs., and the total filtrate from six jars (i.e., from 4,500 grammes of polishings) about 15,000 c.cs.

The filtrate is concentrated under reduced pressure (about 40 mm. of mercury) to about 1,850 c.cs. and made up to 2,025 c.cs. with distilled water, and 225 c.cs. of distilled alcohol are added.

The product obtained from each concentration is allowed to stand two or three days without shaking to enable the precipitate, which is formed on the addition of alcohol to settle. The supernatant liquid is then poured through a Whatman No. 2 folded filter paper and when the liquid has filtered the precipitate is poured on the filter. The precipitate on the paper is washed once or twice with very small quantities of water the washings being added to the main filtrate, and water added to make up to 2,250 c.cs.

One c.c. of this extract represents two grammes of the original polishings. It contains about 9.5 per cent. of alcohol as a preservative.

TRADE AND CUSTOMS DEPARTMENT.

The work for this department consists mainly in the examination of samples in connexion with the assessment of duty, e.g., wines, potable spirits, liquors and denatured spirits, and samples of chandu and chandu dross, submitted by the Chandu Monopoiy Department in connexion with the administration of the Chandu Enactment.

Liquors.

In connexion with the assessment of duty, 91 samples of alcoholic liquors were tested for their spirit strength. In addition, five samples were examined for denaturants, four of which were found to contain the necessary ingredients.

TODDY.

The acidity and alcoholic strength were determined in 171 samples. In eight samples, the acidity exceeded 0.8 per cent. expressed as acetic acid.

It is probable that most of the toddy offered for sale is diluted with water, but no satisfactory method for the detection of this adulteration is at present known. Investigations with this end in view were commenced during the year, and will be continued during 1927.

CHANDU.

Under the provisions of the Chandu Enactment, the importation of chandu of other than Government manufacture is prohibited. The Enactment, further, makes it an offence to be in possession of:

- (a) any chandu reprepared from dross,
- (b) more than seven and a half tahils of Government chandu.
- N.B.—One tahil equals approximately 37.8 grammes.

Two hundred and four samples of chandu and substances suspected to contain chandu were received. Of these, 71 were found to be Government chandu, 118 illicit chandu, four chandu prepared from chandu dross and three imitation chandu containing no opium.

CHANDU DROSS.

The Chandu Monopoly Department purchases chandu dross from licensees at prices varying with the quality of the dross. The chandu dross is collected by the Chandu Monopoly Department and inspections of the dross are carried out by a Chemist from this laboratory.

The number of samples inspected during the year was 4,419. The samples examined were graded as follows:

Grade 1	 	***		***		774.	 4,282
,, II	 **	200	***		***		 94
., III	 					***	 43

DELETERIOUS DRUGS.

Under the provisions of the "The Deleterious Drugs Enactment, 1911", it is an offence to import, sell or use without licence certain drugs (morphine, cocaine, eucaine, etc.) or preparations containing more than a certain percentage of these drugs. In suspected cases, the officers of the Customs Department take samples, which are forwarded to this laboratory for analyses. Two samples were submitted during the year, no deleterious drug was found to be present in the samples.

MISCELLANEOUS SAMPLES.

In connexion with the administration of the Customs and Excise Enactments, twenty-nine samples were received. These included samples of turpentine, Rangoon oil and substances suspected to contain "ganja."

POLICE DEPARTMENT.

One thousand two hundred and fifty-nine exhibits were received from the Police Authorities in connexion with proposed proceedings in the Courts. These may be classified as follows:

- (1) Coins and coining materials;
- (2) Articles for blood stains;
- (3) Toxicological analyses;
- (4) Deleterious Drugs;
- (5) Miscellaneous.
- (1) Coins and Coining Materials.—The number of exhibits examined was 1,077. Of these 1,043 were counterfeit coins, and two moulds. The other exhibits consisted of pieces of metal and chemicals used in the manufacture of counterfeit coins.
- (2) Articles for Blood Stains.—Exhibits, in connexion with charges of murder, were received from the Police Authorities for examination for bloodstains. The number of exhibits received was 96, of which 57 gave positive results. The exhibits, which gave positive reactions for blood were further examined by the precipitin test for human serum, 46 of these gave the reaction characteristic of human serum.

The results of the examinations are tabulated below:

Exhibits.		Number examined.	Number blood-stained	Number human-blood.
Knives, parangs, etc	Service	49	 28	 19
Articles of clothing	***	37	 27	 26
Woods, mats, etc		9	 5	 1
Earth		2	 2	 2

(3) Toxicological Analyses.—Forty exhibits were received for examination. Of these 21 were of human viscera, in 12 of which poison was identified. Morphine was found in 4 cases, arsenic 3, cyanide 2, alcohol 1, oxalic acid 1, santonin 1, and sodium hydroxide in 1.

In the last case, the stomach and contents gave a markedly alkaline reaction. The mucous membrane of the stomach was blackened, hardened and corrugated. Other exhibits received in connexion with human poisoning included samples of urine, vomited matter, foodstuffs, and liquids.

The poisons identified in exhibits other than viscera were opium in five exhibits, arsenic four, datura two, cyanide two, yellow phosphorus one and mercuric chloride in one.

One case of arsenical poisoning deserves mention, where fifty or more coolies suddenly became ill after drinking toddy, without fatal results. Two samples of the toddy received for examination contained a mixture of toddy, sulphur and arsenic. In one sample (770 cubic centimetres) the quantity of arsenic found was equivalent to 6.4 grams. of arsenicus oxide and in the other sample (740 cubic centimetres) the quantity of arsenic found was equivalent to 4.2 grams. of arsenious oxide. The arsenic and sulphur (a mixture used for white ants) had been mixed with the toddy either accidentally or purposely. The complete history of these cases has not yet been obtained.

Case of Datura Poisoning.—The alkaloids of datura stramonium were found in a curry and in two vomits of different individuals who had partaken of this curry.

Case of Mercuric Chloride Poisoning.—A sample of samsu (130 cubic centimetres) was found to contain 2.8 grams, of mercuric chloride. As 0.2 gram, of this substance may be a fatal dose, it follows that less than half an ounce of the samsu contained a fatal dose. Two coolies who tasted the samsu were violently ill. Both subsequently developed symptoms of mercury poisoning but recovered under treatment in hospital.

- (4) Deleterious Drugs.—Two substances suspected to contain one of the deleterious drugs specified in "The Deleterious Drugs Enactment 1925", were examined with negative results.
- (5) Miscellaneous.—There were 35 exhibits submitted. Of these, 11 were found to be "ganja." The other exhibits included some notebooks for finger prints, materials is connexion with explosions and pieces of alleged gold bullion, which were found to contain a core of silver. A liquid, which had been thrown at a person with intent to injure was found on examination to contain caustic soda.

OTHER DEPARTMENTS.

Samples submitted by the Federated Malay States Railway Department included 28 samples of white metal, and two samples of axle oil. Samples of water were also examined as to their suitability for use in boilers and for domestic purposes.

For the Public Works Department samples of water for domestic purposes were analysed.

For the Forest Department samples of the following resins were examined: Damar Penak, Damar Batu, Damar Mata Kuching and dragons blood. A sample of Damar Resak from a species of vatica was found to contain an oil similar to that of Gurjun Balsam

Samples of Minyak Keruing were examined to ascertain whether the oleo-resin of this species of dipterocarpus would be of commercial value.

Two patent specifications is connexion with "The Inventions Enactment, 1914" were examined for the Federal Secretariat.

PRIVATE ANALYSES.

Forty-nine samples were received for examination. Included in this total were samples of water, explosives and kerosine.

The fees for these analyses amounted to \$2,280/-.

LEGAL PROCEEDINGS.

Members of the staff of the Chemical Laboratory gave evidence in legal proceedings on 31 occasions.

STAFF.

The staff of the Chemical Laboratory consists of: Chemist;

Three Assistant Chemists.

In addition there are five laboratory assistants, two attendants for gas plant and one laboratory attendant.

Institute for Medical Research, R. W. Blair,
Federated Malay States, Chemist-in-Charge, Government Laboratories,
Kuala Lumpur, Federated Malay States.

10th February, 1927.

APPENDIX A.

TOTAL NUMBER OF SAMPLES EXAMINED IN THE CHEMICAL LABORATORY DURING THE YEARS 1925 AND 1926.

District and annual District D	URING	THE	YEA	ARS 19	25 ANI) 193	26,		
						,	Number of analyses, 1925.		Number of analyses, 1926.
MEDICAL DEPARTMENT.	Arrila .								
Milk, chemical	1000	***	die	((0.00)			741		597
Milk, bacteriological				The same			12		3
Water, chemical		***	***	***			328		528
Water, bacteriologic	eal			***			36		99
Sewage, effluents				015357			26		10
Toddy							262		171
Toxicological analyse	es						24		13
Miscellaneous	***	111	***	110222		***	27		37
TRADE AND CUSTOMS DEE	ARTME	ENT.—							
Liquors							138		96
Toddy							28		171
Chandu				***			400		204
Chandu dross							3,180		4,419
Deleterious drugs							24		2
Miscellaneous	I /		***		1126		28	0.000	29
POLICE DEPARTMENT									
Coins and coining	materi	als	***	***	odeesun		495		1,077
Articles for blood s	stains			***			107		96
Toxicological analys	es			***			30		49
Liquors							5		-
Deleterious drugs		***					2		2
Miscellaneous							12		35
OTHER DEPARTMENTS									
Miscellaneous		***					87		69
PRIVATE ANALYSES.—									
Water		***	***	***	***		84	***	40
Milk	***	***					12		1
Spirits	***					***	18		-
Toxicological analyse	s		***	***		***	1		-
Miscellaneous	***		***				14		8
					Total		6,021		7,756
							100000		

APPENDIX B.

KUALA LUMPUR WATER SUPPLY.

Chemical Averages for each month of the year 1926. Parts per 100,000 unless otherwise stated.

1	Reinfall in inches.	7.41 7.82 7.82 7.83 7.85 7.85 7.85 7.93 10.00	9.88	Oxidized .msgorita	000 000 000 000 000 000 000 000 000 00	.005
	Oxidized nitrogen.	000000000000000000000000000000000000000	.002	Total solids,	25.55.50.05.75.55.55.55.55.55.55.55.55.55.55.55.55	3.96
(Raw Water.)	.sbilos latoT	00004004404 5555005504	8.83			
(Baw	Chlorine.	2,9,9,9,9,9,9,9,9,9,9,9,9,9,9,9,9,9,9,9	5.86. 86.	(Filtered Water, Oblorine,	AMERICAN SALARY	90.
Ampang.	Oxygen absorbed in 3 hrs.	0956 11578 11578 11578 11517 11517	1795	bedrosda	0731 .0654 .0947 .1270 .1407 .1151 .1151 .1229 .1229 .1228 .1242 .1268	.1087
Intake Works,	Albuminoid nitrogen.	00040 00000 00000 00000 00000 00000 00000 0000		And the state of t	.0036 .0047 .0052 .0052 .0056 .0047 .0036 .0040 .0048 .0048	.0044
Intak	Ammoniacal nitrogen.	00000 00000 00000 00000 00000 00000 0000		Ammoniacal introgen.	000000000000000000000000000000000000000	00000
	нф	666720055005		н	00000000000000000000000000000000000000	7.5
	Colour M. M. Brown.	19888989888	2 2 2 2	Colour M. M. Brown.	77588553884489	22
	Rainfall in inches.	6.24 7.112 14.15 10.62 7.83 4.0 4.81 6.58 11.38	18.82	basibixO .nagordin	000000000000000000000000000000000000000	.005
	Oxidized nitrogen.	424888999999999999999999999999999999999	.003	Total solids.	2.4.2.2.4.2.2.2.2.2.2.2.2.2.2.2.2.2.2.2	3.15
(Raw Water.)	Total solids.	0444 057 057 057 057 057 057 057 057 057 057	8.79	(Filtered Water,	0777770 000 000 000 000 000 000 000 000	20.
(Raw	Chlorine.	8,6,6,8,8,8,6,8,8	20.	(Filter		
servoir.	Oxygen absorbed in 8 hrs.	1135 11279 11279 11680 11680 11690 11690	1634		.0468 .0507 .0507 .0507 .0939 .0946 .0946 .0956 .0966	.0768
Impounding Res	Albaminoid nitrogen.		0144	and an analysis and an analysis and an analysis and an analysis and analysis and analysis and analysis and analysis and analysis	.0032 .0032 .0035 .0035 .0035 .0037 .0037 .0039 .0039	.0041
Impo	Ammoniacal nitrogen.		00012	M liconinoninA mitrogen.	000000000000000000000000000000000000000	00000
	Hd	27.7.0 6.0.0 6.0.0 7.0.0 8.0 8		Hq	0.00 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	6.0
	Colour M. M. Brown.	82123624866566	8 8	Colour M. M. Brown.	0554774458888888888888888888888888888888	16
			111		11/11/11/11/1	:
	ath.	1111111111	 Averages	ath.	111111111111	Averages
	Month	January February March May June July August September October	December	Month	January February March April June July September October November	7

APPENDIX C.

AVERAGES YEARS (1921-1926.)

KUALA LUMPUR WATER SUPPLY: IMPOUNDING RESERVOIR AT AMPANG (RAW WATER)
(High Level Supply).

Parts per 100,000 (unless otherwise stated).

Years 1921-1926.			Ammoniacal nitrogen.	Albuminoid nitrogen.	Oxidized nitrogen.	Chlorides as chlorine.	Oxygen ab- sorbed from permanga- nate 3 hours at 80°F.	Colour M.M. Brown in a 2 feet tube.	Total solids.	Rainfallin inches per mensem.	Reaction pH
1921			.0006	.0124	.002	.06	.1619		4.0		
1922			.0009	.0131	.003	.07	.1641		4.4	8.07	
1923			.0007	.0116	.004	.07	.1736	22	4.0	8.41	
1924			.0006	.0111	.002	.07	.1516	21	4.3	7.61	
1925			.0005	.0102	.004	.07	.1484	26	4.5	11.53	7.5
1926			.0004	.0106	.003	.07	.1521	29	3.8	9.21	6.5
Average	s for 6 y	rears	.0006	.0115	.003	.07	.1586	25	4.2	8.97	7.0

APPENDIX D.

AVERAGES YEARS (1919-1926.)

KUALA LUMPUR WATER SUPPLY: MAXWELL'S HILL (FILTERED WATER) (HIGH LEVEL SUPPLY).

Parts per 100,000 (unless otherwise stated.)

	Years 191	9–1926.		Ammoniacal nitrogen.	Albuminoid nitrogen.	Oxidized nitrogen.	Chlorides as	Oxygen ab- sorbedfrom permanga- nate3hours at 80°F.	Colour M.M. Brown in a 2 feet tube.	Total solids.	Reaction pH
1919		***		.0010	.0094	.010	.08	.0741			
1920			***	.0005	.0074	.010	.06	.0834		5.5	
1921				.0001	.0047	.011	.06	.0828		3.1	
1922		***	***	.0001	.0047	.007	.07	.0835		3.4	
1923	****		***	.0002	.0043	.006	.07	.1049	18	3.5	
1924	***		***	.0008	.0055	.004	.06	.0897	17	3.6	
1925				.0001	.0056	.005	.07	.0850	19	4.0	7.0
1926				.0000	.0041	.005	.07	.0768	16	3.2	6.0
Ay	erages for	8 years		.0004	.0057	.007	.07	.0850	18	3.7	6.5

APPENDIX E.

AVERAGES YEARS (1921-1926).

KUALA LUMPUR WATER SUPPLY: AMPANG INTAKE WORKS (RAW WATER) (LOW LEVEL SUPPLY).

Parts per 100,000 (unless otherwise stated).

Years 1921-1926.		Ammoniacal nitrogen,	Albuminoid nitrogen.	Oxidized nitrogen.	Chlorides as	Oxygen ab- sorbed from pormanga- nate 3hours at 80°F.	Colour M.M. Brown in a 2 feet tube.	Total solids.	Rainfallin inches per mensem.	Reaction pH	
1921			.0005	.0055	.003	.06	.1555		5.1		
1922			.0006	.0067	.004	.06	.1762	***	3.8	9.22	
1923			.0008	.0071	005	.06	.1859	24	4.0	8.74	
1924			.0008	.0051	.005	.05	.1562	24	4.3	8.28	
1925			.0005	.0066	.006	.06	.1709	27	4.2	12.07	7.5
1926			.0003	.0071	.005	.06	.1764	30	3.8	9.88	6.5
Average	es for 6	years	.0006	.0064	.005	.06	.1702	26	4.2	9.64	7.0

APPENDIX F.

AVERAGES YEARS (1919-1926).

KUALA LUMPUR WATER SUPPLY: WELD HILL (FILTERED WATER)
(LOW LEVEL SUPPLY).

Parts per 100,000 (unless otherwise stated).

У	ears 191	9–1926.	Ammoniacal nitrogen.	Albuminoid nitrogen.	Oxidized nitrogen.	Chlorides as	Oxygen absorbed from permanga- nate 3 hours at 80° F.	Colour M.M. Brown in a 2 feet tube.	Total solids.	Reaction pH
1919			 .0010	.0078	.010	.08	.1113			
1920			 .0005	.0064	.009	.07	.1267		6.0	
1921	****		 .0002	.0044	.007	.06	.1206		4.4	
1922			 .0001	.0048	.005	.06	.1329		3.5	
1923			 .0002	.0046	.007	.06	.1465	22	3.6	
1924			 .0003	.0033	.006	.05	.1155	21	3.8	
1925			 .0001	.0044	.006	.06	.1054	23	4.0	8.0
1926			 .0000	.0044	.005	.06	.1087	22	3.9	7.5
Average	es for 8	years	 .0003	.0050	.007	.06	.1210	22	4.2	7.5

REPORT OF THE SENIOR HEALTH OFFICER, FEDERATED MALAY STATES, FOR THE YEAR 1926.

HEALTH STAFF.

1. The sanctioned staff for the year was:

- 2. Of the above, six Health Officers and six Medical Officers, seconded to the Health Branch, and two Chief Sanitary Inspectors were available, one of the latter on his return from leave on 26th February, 1926, being posted to Tapah to supervise the anti-malarial works in connection with the road to Cameron's Plateau (4,750 ft.), which will be a hill station in the future.
- 3. Eight of the Sanitary Inspectors were absent in Singapore for six months, undergoing a course preparatory to taking the examination for the Royal Sanitary Institute certificate, which certificate must be held before an Inspector can be promoted to the higher grades. Their services were not available to the branch from May until November.
 - 4. The following table shows the principal changes among the senior staff:

Date.	Name of officer.	Designation.	From which district.	To which district.	Remarks.
1-2	C.S. Ryles, o.B.E.	Health Officer	Selanger East	Negri Sem- bilan	
21-3	E. R. C. Cooke,	Medical Officer	Railways	vagatta'i	On long leave
10-4	A. R. Wellington	Senior Health Officer	-	To act as Principal	
		Onicer		Medical	
				O ffi c e r F.M.S.	
10-4	A. K. Cosgrave,	Senior Medical Officer, Selan-	100 7 8700	To act as Senior	
		gor		Health Officer	0501
		150		F.M.S.	IDEE
23-4	W. O. Pou	Health Officer	3/0 7 1000	Lower Perak and Batang	Returned from leave
			1000 11000	Padang	
	P. G. Temple	**	Lower Perak	1000	On long leave
1-5	W. J. Vickers	"		Selangor East	Transferred from the Medical Branch
15-5			100, -0000	Perak North	Returned from leave
17-5	J. G. Castellain	Health Officer	-	Railways	On month to month engagement
18-5	A. Reid	**	Perak North	-	Reverted to Medical Branch
19-5	D. W. G. Faris	Medical Officer	-	Port Swet- tenham	Transferred from Medical Branch
13-8	C. C. Taffs	,,	-	Negri Sem- bilan	Recruit
18-12	R. B. Jackson	,,	-	Pahang	Returned from leave
23-12	V. D. Wyborn	"	Pahang	-	Reverted to Medical Branch

DUTIES OF THE HEALTH BRANCH.

- 5. The duties of the members of the Health Branch include:
 - (i) Vital statistics and work under the Registration of Births and Deaths Enactment.
 - (ii) Malaria investigation and mosquito control.
 - (iii) Work under the following Enactments:
 - (a) Quarantine and Prevention of Disease Enactment:
 - (b) Sanitary Boards Enactment;
 - (c) Sale of Food and Drugs Enactment;
 - (d' Labour Code;
 - (iv) General, including inspection of schools, Government lines, water works, etc.
 - (v) Maternity and Infant Welfare.

REVENUE AND EXPENDITURE.

6. The only revenue collected was that for certificates issued under the Registration of Births and Deaths Registration Enactment and for the sale of mosquito identification books. The total amount collected was \$2,887.25.

7. The expenditure debited against the Health Branch was:

Personal Emoluments			\$189,164.87
Temporary Allowances		100	82,079.27
Other Charges, Annually Recurrent		444-	94,453.11
Other Charges, Special Expenditure	due-sold a		6,555.22
Clerical Service	***		\$322,252.47 41,436.60
	Total		\$363,689.07

GENERAL REVIEW OF WORK DONE.

- 8. The Health Branch not yet being up to strength was not able to carry out in full, all the duties pertaining to the branch.
- The Senior Health Officer in addition to his own duties carried out those of the Health Officer, Railways, from 21st March, 1926, to 17th May, 1926.
- 10. Under the Quarantine and Prevention of Disease Enactment work was shared with the Medical Branch. There were no large epidemics though a small outbreak of cholera in the Kuala Langat district in March, caused considerable anxiety.
- 11. Work under the Labour Code was satisfactory in view of the number of the staff. Of 1,450 estates, 725 were visited by Health Officers and of 156 Estate Hospitals 154 were visited.
- School inspection was shared with the Medical Department, 213 visits were made by the Health Staff.
- 13. A very marked increase of immigrants, 99,066 for 1926, compared with 48,748 for 1925, taxed the accommodation at Port Swettenham Quarantine Camp somewhat severely, in view of the fact that several successive steamers arrived infected with cholera, the situation was satisfactorily dealt with by the Quarantine Camp Staff.
- 14. Towards the end of December the rain-fall over parts of the peninsula was unprecedented and on the night of the 27th very severe floods at Kuala Lipis resulted in the Health Office being swept away, all records and papers were totally lost, a very serious matter since the Health Officer is the Registrar of Births and Deaths tor Pahang.

In this report therefore it is impossible to give any figures for Pahang for the year under review.

HEALTH LEGISLATION.

- 15. The following laws affecting the public health were passed during the year:
 - (a) Enactment 10 of 1926. An Enactment to amend the Quarantine and Prevention of Disease Enactments of 1903.
 - (b) Enactment 13 of 1926. An Enactment to provide for the establishment of Health Boards and to prescribe the powers and duties thereof. In force from January 1st, 1927.
 - (c) Enactment 23 of 1926. An Enactment to declare the law relating to lepers.

VITAL STATISTICS.

- 16. Populations are estimated from the Census figures which are believed to be correct. Immigration and emigration have a greater influence on population than have births and deaths. Accurate information concerning imagination and emigration especially by rail are not available and tables cannot be given.
- 17. Increase in population is calculated on the arithmetical increase basis as experience has shown this to be more reliable than that based on geometrical grounds.
- 18. Births and deaths figures are obtained from notifications compulsory under the Registration of Births and Deaths Enactment which is everywhere in force. The tota' number of births and deaths is approximately correct. The accuracy of diagnoses as to causes of deaths is in the majority of cases open to question, for few of the cases were seen by a qualified medical man previous to decase. In each of the four large towns every undercertified body is viewed by the Assistant Health Officer who interrogates the friends and forms a diagnosis. In rural districts these duties are carried out by the Police.
- 19. Deaths in towns are debited against the town only if the deceased was resident there for three months or more previous to death. The towns contain hospitals which cater both for the town and the district surrounding it. It is a well-known fact that chronic cases from the rural areas drift to the towns in the hope of getting more skilled treatment. Taking all things into consideration even with a qualifying period of three months, a number of deaths are debited against the towns which should be debited against the rural areas where the disease was contracted.
- 20. The general death-rate for three States (Pahang excluded) was 29.22 per mille as against 28.60 for the whole Federated Malay States for 1925, this increase is probably due to the wave of malaria which occurred in the first half of the year, the absence of the Pahang figures and the increased immigration which is not reckoned in the population.
- 21. The number of deaths attributable to fevers (most of them probably of malarial origin) was 16,531 or 43 per cent. of the total. Last year the percentage was 42.09.
- 22. Dysentery and diarrhoea accounted for 7.14 per sent. of the total deaths, pulmonary tuberculosis for 4.87 per cent., pneumonia for 6.69 per cent., and convulsion for 11.18 per cent.
- 23. Because of the peculiar age and sex distribution and the fact that the labour of this country is largely males aged 20-45 who passed the doctor before embarking for Malaya, the death figures cannot be compared with countries where the labour is indigenous and where age and sex distribution are normal. With normal age and sex distribution the death-rate would probably be twice what it is.
- 24. There is however no getting away from the fact that the death-rates since 1911, when the Health Branch was first formed have shown a steady decline, with the exception of the year under review.
 - 25. The following figures are put for comparison:

		F	edera	ıl.	1	Kuala Lu	mpui	Town.	E	states		
Year.		General death rate.		Fevers death rate.		General death rate.		Fevers death rate.	General death rate.		Fevers. death rate.	
1911	***	 39.11		17.47	***	39.02		9.87	 62.90	***	Not known	
1926	***	 29.22		12.57	***	21.01		3.74	 14.51		5.89	

26. The figures for estates must be accepted with caution for few Asiatic-owned estates ever admit either the existence of sickness or the occurrence of a death on the estate and nil returns are sent to the Health Officer.

VITAL STATISTICS (FEDERAL).

POPULATION.

27. The population of the Federated Malay States as estimated was at the end of June, 1926, 1,476,082, distributed as follows:

Perak	 	 	 		654,179
Selangor	 	 	 		457,170
Negri Sembilan	 	 	 ***	***	204,257
Pahang	 	 	 		160,426

28. Assuming that the races remain in the same proportion as in the Census year the race distribution is as follows:

State.	Europeans and Americans.	Eurasians.	Malays and other natives of the Archi- pelago.	Chinese.	Indians.	Others.
Selangor Negri Sembilan	2,389 3,054 1,152 352	1,040 1,775 548 132	260,177 105,875 81,797 110,211	228,461 181,071 77,948 39,258	160,136 163,246 41,748 9,785	1,976 2,149 1,069 688
Total .	6,947	3,495	558,060	526,733	374,915	5,882

BIRTHS.

29. Thirty-nine thousand eight hundred and thirty-four births were registered during the year, giving a birth-rate of 30.28 (excluding Pahang) per mille of population. In 1925 the number was 41,818 and the rate was 28.89.

The following table shows the number of births and birth-rates according to races:

		Race.				1	No. of birth	18.	Birth-rate.
Europea	ns an	d Ame	ericans				131		19.86
Eurasian	8						127		37.76
Malays a	nd ot	her rac	es of th	he Arc	hipelag	o	17,522		39.12
Chinese			***				12,973		26.61
Indians							9,010		24.68
Others							71		13.67

DEATHS

30. Thirty-eight thousand four hundred and forty-five deaths were registered, giving a death-rate of 29.22 (excluding Pahang) per mille. The number of deaths in 1925 was 34,153 and the rate was 23.60 for the whole Federated Malay States.

The distribution of deaths among the several races was as follows:

	Rac	ю,				2	No. of death	s.	Death-rate.
European	ns an	d Ame	ericans		***		36		5.46
Eurasian	s						46		13.68
Malays a	and ot	her rac	es of th	e Arc	hipelag	, oz	11,657		26.03
Chinese							14,730		30.22
Indians							11,920		32.64
Others							56		10.78

31. The deaths and death-rates for the total population for the last ten years were as follows:

Year.		Population.	Deaths.	R	ate per mille.
1917	 	 1,244,018	 42,514		34.17
*1918	 	 1,279,859	 67,639		52.85
1919	 	 1,315,700	 38,645	***	29.37
1920	 	 1,851,541	 43,705		32.34
1921	 	 1,304,825	 38,077		29.18
1922	 	 1,360,876	 35,028		25.74
1923	 	 1,389,667	 33,914		24.40
1924	 	 1,418,455	 33,585		23.68
1925	 	 1,447,248	 34,153		23.60
1926	 	 1,315,606†	 38,445†		29.22†

^{*} High figure due to influenza epidemic.

[†] Excluding Pahang.

32. Table showing causes of deaths in 1926:

Diseases.				No	of deaths.	R	ate per mille.
Malaria		***	***		16,531		12.57
Dysentery and diarri	hoea			***	2,744		2.09
Pneumonia		See.	***	***	2,578		1.96
Pulmonary tuberculos	sis	1.00			1,878		1.42
Ankylostomiasis		***		***	228		0.17
Beri-beri					314		0.24
Syphilis	***		***		84	***	0.06
Enteric	***	ine		***	28		0.02
Tetanus					65		0.05
Convulsions					4,297		3.27
Bright's disease			***		195		0.15
Ptomaine poisoning			***	***	126	***	0.10
Influenza	***		***		98		0.07
Other diseases					9,294	***	7.06

33. The following table shows the deaths and death-rates from the principal diseases for the last ten years:

Yea	Year.	Mala	Malaria,		ry and oea.	Pulmo		Beri-	beri.
		Deaths.	Rate.	Deaths.	Rate.	Deaths.	Rate.	Deaths.	Rate.
	F0.385		Ter.						
1917	17.00	18,750	15.07	4,942	3.97	2,446	1.96	1,207	0.97
* 1918		31,515	24.62	4,280	3.34	3,184	2.48	1,277	0.98
1919		16,975	12.90	3,712	2.82	2,445	1.86	930	0.71
1920	10.00	20,595	15.24	3,804	2.81	2,634	1.95	431	0.32
1921	88.3.0	17,168	13.16	2,999	2.30	2,255	1.73	422	0.32
1922		15,570	11.44	2,419	1.78	2,393	1.76	443	0.33
1923	1000	15,516	11.17	2.142	1.55	1,934	1.39	378	0.27
1924		14,283	10.07	1,961	1.38	1,916	1.35	453	0.32
1925		14,377	9.93	1,945	1.34	2,001	1.38	424	0.29
1926		16,531	12.57	2,744	2.09	1,873	1.42	314	0.24

^{*} Influenza epidemic year.

+ Excluding Pahang.

34.---

VITAL STATISTICS (State figures for comparison). Birth Table.

State.		No. of birth.	per	Birth-rate 1,000 livin	g.	1925 birth-rate.
Perak		 20,095		30.72		27.76
Selangor		 13,914		30.44		29.69
Negri Sembilan	2.0 1.1	 5,825	Link	28.51		27.69
Pahang	***	 No figures	N	o figures		32.80

35. Birth statistics of different nationalities.

place and the self-ord		Europeans		Eur	asians.	Malays and other races of the Archipelago.		Chinese.		Indians,		Others.	
State.		Births	Birthrate.	Births,	Birth-rate.	Births.	Birth-rate.	Births.	Birth-rate.	Births.	Birth-rate.	Births.	Birth-rate.
Perak Selangor Negri Sembilan		46 78 7	19.25 25,54 6.08	32 64 31	30.77 .36.06 56.57		39.11 37.82 40.87	6,118 5,332 1,523	26.78 29.45 19.54	3,676 4,418 916	22.96 27.06 £1.94	48 18 5	24,30 8.38 4.68

* No figures.

36.-

DEATH	TABLE	(State	figures	for	com	parison).
-------	-------	--------	---------	-----	-----	---------	----

State.		No.	of deaths.	Death-rate 1926.		Death-rate 1925.
Perak		***	18,506	 28.29	111	23.86
Selangor		***	13,390	 29.29		22.75
Negri Sembilan	***		6,549	 32.08	***	28.51
Pahang (No figur	res).					

37. Deaths and death-rate of different nationalities.

State.		opeans and rienns,	Eurasians,		Malays and other races of the Archipelago,		Chinese,		Indians,		Others.	
State.	Deaths.	Death- rate.	Deaths.	Death-	Deaths.	Desth- rate.	Deaths.	Death- rate.	Deaths.	Death- rate.	Deaths.	Death-
Perak Selangor Negri Sembilan Pahang *	16	6.69 5.24 3.47	8 22 16		6,433 2,667 2,557	24.73 26.13 31.26	7,287 5,239 2,204	31,90 28,93 28,28	4,727 5,431 1,762	29.52 33.27 42.20	35 15 6	17.71 6.98 5.61

^{*} No figures.

38. Table showing deaths and death-rates from principal diseases.

		Malaria.			sentery a liarrhoes			ulmonar berculos	Beri-beri.			
State.	1	R	ate.	i	Ra	te.	by.	Ra	te.	hs.	Rs	te.
	Deaths	1926.	1925.	Deaths	1926,	1925.	Deaths.	1926,	1925.	Deaths.	1926,	1925.
Perak Selangor Negri Sembilan Pahang *	8,73 5,01 2,78	1 10.96	11.11 8.07 8.74	1,027 1,211 506	1.57 2.65 2.48	1.16 1.65 1.25	996 521 356	1.52 1.14 1.74	1.49 1.20 1.65	108 140 66	0.17 0.31 0.32	0.13 0.34 0.39

^{*} No figures.

INFANTILE MORTALITY.

39. There were 7,718 deaths of children under one year of age. The infantile mortality rate or rate per 1,000 births was 193.75; the rate for 1925 was 177.17.

Infantile Mortality Table.

State.				ths of chil der one you of age.	Death-rate per 1,000 births.
Perak	1.12	 		3,459	 172.18
Selangor	***	 		2,888	 207.56
Negri Sembilan		 	1	1,371	 235.36
Pahang (No figures).				

40. Deaths from Zymotic Diseases.

State.			1	Plague.		Cholera.		Smallpox.		ebro-spinal eningitis.
Perak		insti-		-		_	***	-		5
Selangor		500		-	0.4.	6			od. ob	8
Negri Ser	mbilan	11.1		-		-				2
Pahang				-		-	***	-	1	The state of the s

41. Death-rates from principal diseases for the last seven years:

				Pernk.			ielangor.		Neg	ri Sembi	banc,	Pahang.		
Name of the least	Year.		Malaria.	Dysentery and diarrhoes.	Pulmonary taberculosis.	Mahria.	Dysentery and diarrhoes.	Pulmonary taberculosis.	Malaria,	Dysentery and diarrhoea.	Pulmonary tabercalosis.	Malaria.	Dysentery and diarrhoea.	Pulmonary tuberculosis.
1920 1921			15.82 13.32	2.21 1.54	2.11 1.78	13.13 11.80	3.77 3.29	1.89	18.18 13.07	4.31	2.37 1.68	15.24 16.61	1.19	0.94
1922		-	12,29	1.42	1.83	9.96	2.39	2,00	11.51	2,40	1.69	12.10	0.74	0.8
1923			12,46	1.26	1.52	9,48	1.93	1.48	10.53	1.45	1.41	11.36	1.73	0.5
1924			10.66	1.20	1.50	7.40	1.67	1.36	10.50	1.32	1.52	14.59	1.41	0.50
1925	Park.	-	11.11	1.16	1.49	8,07	1:65	1.20	8.74	1.25	1.65	11.91	1.35	1.1
1926		***	13.35	1.57	1.52	10.96	2.65	1.14	13.65	2.48	1.74			

^{*} No figures.

42. Vital statistics for the four large towns, Kuala Lumpur, Ipoh, Taiping and Seremban.

Town.		Estimated	Bir	rths.	Deaths of persons who previous to decease had resided in town three months.			
Town.		population.	Number.	Rate per mille.	Number.	Rate per mille.		
Kuala Lumpur Ipoh		98,125 43,662	3,645 1,364	37.15 31.24	2,062 580	21.01 13.28		
Taiping		21,936 21,838	1,107 664	50.46 30.40	758 533	34.55 24.41		

43. Table showing corrected death-rates in the four principal towns during the last seven years.

Kuala L	ampur.	Ipo	oh.	Seren	ıban.	Taiping.		
Population.	Death-rate.	Population.	Death-rate.	Population.	Death-rate.	Population.	Death-rate.	
 67,980	30.00	34,357	22.64	15,006	34.05	25,484	39.90	
 81,197	27.02	38,895	21.78	13,398	27.93	21,178 21,296	50.05 35.08	
 88,009 91,381	19.19	40,399	20.12	19,210	24.78	21,462	33.45 33.91	
 94,753 98,125	15.31 21.01	42,334 43,662	13.98 13.28	20,938 21,838	17.77	21,780 21,936	29.11	
	67,930 81,197 84,476 88,009 91,381 94,753	67,930 30.00 81,197 27.02 84,476 21.36 88,009 19.19 91,381 16.74 94,753 15.31	67,930 30.00 34,357 81,197 27.02 37,194 84,476 21.36 38,895 88,009 19.19 40,399 91,381 16.74 41,047 94,753 15.31 42,384	67,930 30.00 34,357 22.64 81,197 27.02 37,194 20.38 84,476 21.36 38,895 21.78 88,009 19.19 40,399 20.12 91,381 16.74 41,047 13.89 94,753 15.31 42,334 13.98 94,753 15.31 42,334 13.98	67,930 30.00 34,357 22.64 15,006 81,197 27.02 37,194 20.38 17,479 84,476 21.36 38,895 21.78 13,398 88,009 19.19 40,399 20.12 19,210 91,381 16.74 41,047 13.89 20,074 94,753 15.31 42,334 13,98 20,938	67,930 30.00 34,357 22.64 15,006 34.05 81,197 27.02 37,194 20.38 17,479 36.16 84,476 21.36 38,895 21.78 13,398 27.93 88,009 19.19 40,399 20.12 19,210 24.78 91,381 16.74 41,047 13.89 20,074 17.34 94,753 15.31 42,334 13.98 20,938 17,77	67,930 30.00 34,357 22.64 15,006 34.05 25,434 81,197 27.02 37,194 20.38 17,479 36.16 21,178 84,476 21.36 38,895 21.78 13,398 27.93 21,296 88,009 19.19 40,399 20.12 19,210 24.78 21,462 91,381 16.74 41,047 13.89 20,074 17.34 21,616 94,753 15.31 42,334 13,98 20,938 17,77 21,780	

44. Table showing corrected deaths and death-rates from principal disease:

Town.	Malai	rin.	Dysenter diarrh		Pulmor tubercu		Beri-beri.	
men	 Deaths.	Rate.	Deaths.	Rate.	Deaths.	Rate.	Deaths.	Rate.
Kuala Lumpur	 108	1.10	179	1.82	133	1.36	28	0.29
Ipoh	 49	1.12	38	0.87	47	1.08	5	0.11
Taiping	 78	3.56	115	5.24	74	3.37	1	0.05
Seremban	 79	3.61	61	2.79	49	2.24	7	0.32

45. Infantile Mortality Table:

Towns.		Births.	un	Death der one y	ear.	Rate per 1,000 births.
Kuala Lumpur	 	3,645		469		128.67
Ipoh	 	1,364		131		96.04
Taiping	 	1,107		172		155.37
Seremban .	 	664		139		209.33

46. Table showing corrected death-rates for principal diseases in the four towns for the last seven years:

		1	Ku	ala Lump	pur.		Ipoh.			Seremban	la.	Taiping.		
Year.			Malaria,	Dysentery and distriboes.	Dysentery and durrhoea. Pulmonary tuberenlosis.		Malaria. Dysentery and diarrhoes.		Malaria.	Dysentery and diarrhoes.	Pulmonary tuberculosis,	Malaria.	Dysentery and diarrhoes.	Pulmonary tuberculosis.
1920			5,08	2.49	3.48	5,64	2.64	3.75	8.99	5,00	3,40	19.14	2.99	1.57
1921			5.25	3.63	3.22	11.24	2.39	4.54	11.76	16.71	6.23	25.06	6.06	3.94
1922			2.79	2.18	3.33	4.50	1.62	2.54	10.44	5.92	6.35	5.45	1.50	1.74
1923			2.06	1.77	2.91	1.48	1.34	3,61	2.86	3.18	2.45	6.94	2.84	3,73
1924			1.13	1.53	2.66	0.97	1.00	1.51	2.89	1.54	1.89	5.41	2.91	3.47
1925			1.46	1.14	1.75	1.77	0.66	1.51	2.58	1.62	2.72	2.85	4.40	3.81
1926	***		1.10	1.82	1.36	1.12	0.87	1.08	3.61	2.79	2.24	3,56	5.24	3.37

MOSQUITO-BORNE DISEASES.

- 47. In Malaya, these are malaria, filariasis and dengue.
- 48. Cases of filariasis are comparatively rare, dengue is frequently seen, but being non-notifiable statistics about it are not available.

MALARIA AND ANTI-MALARIAL MEASURES.

- 49. Malaria still continues to head the list of diseases as the chief cause of sickness and death in the Federation.
- 50. There was a considerable increase in the number of cases of this disease noted during the month of April to July, a slight increase is generally noted at this season but this year it was more marked than usual. Exceptionally heavy rains at the end of 1925 followed by a prolonged drought, and the increased amount of clearing which followed the rise in the price of rubber are two of the reasons suggested to account for this.
- 51. Complete incidence figures are not available and reliance has to be placed on death notifications as a basis for computation.
- 52. The number of deaths registered during the year as due to fever was 16,531, giving a death-rate per mille population of 12.57.
- 53. Nothing occurred during the year to modify the opinion that malaria prevention by mosquito reduction is the right policy where persons are grouped together in towns, in villages and on estates.
- 54. The Malaria Advisory Board of which the Senior Health Officer is Vice-Chairman met regularly during the year and its minutes were published in the press.
- 55. The Mosquito Destruction Boards continued to function and serve a useful purpose. A new Board was established at Klian Intan (Upper Perak), a small village, where malaria has been severe in the past.
- 56. The anti-malarial activities of the Health Branch included the teaching of magaintology, propaganda, investigations and research, anti-mosquito measures and quinine distribution. These activities were carried out as far as time and circumstances permitted.
- 57. Tablets of quinine to the number of 1,043,200 were issued to the various Health Officers for ultimate free distribution to the public through the Police, the Education Department, the Post Offices and the District Officers.

MEASURES TAKEN FOR PREVENTING THE INTRODUCTION AND SPREAD OF INFECTIOUS AND CONTAGIOUS DISEASES.

QUARANTINE AND PORT HEALTH WORK AT PORT SWETTENHAM.

- 58. During the year 75 ships with immigrant labourers were boarded and inspected. The labourers were landed at the Quarantine Camp. Of the 75 ships 21 were infected—eight with cholera, two with cholera and smallpox, four with smallpox, one with cerebro-spinal meningitis, one with cholera and cerebro-spinal meningitis and five with chickenpox.
- 59. The number of immigrants who entered the Quarantine Station, Port Swettenham was 99,066, the number remaining on 31st December, 1925 was 1,408 making a total of 100,474. The largest number on any one day was 7,713 on the 20th of June.

60.	The following tab					ants w			
	Discharged to Dep	ôt			***		***	98	3,184
	Transferred to Klar	ng Hosp	pital						104
	Absconded from th	ne Quan	antine	Camp					55
	Died in Hospital a								457
	Remaining on 31st							1	1,697
						Tota	1	100),497
61.	There were 457 de	aths m	ade up	as follo	ows:			1	-
	Maie adults								94
	Female adults								57
	Male minors								42
	Female minors			***					34
	Infants				***				230
									450
									457

The percentage of deaths to total arrived was 0.45.

- 62. The daily average of immigrants in the camp was 2,445.
- 63. Three thousand eight hundred and ninety-seven passengers were quarantined during the year.
- 64. Forty-four thousand four hundred and sixty immigrants received routine treatment for ankylostomiasis.
- 67. The number of infectious diseases treated were cholera 196, smallpex 5, chickenpox 11, measles 432, cerebro-spinal meningitis 4, typhoid fever 3, erysipelas 1, mumps 1. With the exception of 13 cases of cholera admitted from the Kuala Langat district, all cases came from ships or developed the disease while in the camp.
- 68. The total number of persons removed to the camp from cholera infected ships was 15,061, of these 157 either developed cholera or had the disease on admission to the camp, 39 cases remained on 31st December, 1925. Of the 196 treated 79 died giving a percentage of 40.31.

Every precaution was taken to cut short the course of the epidemics. Prophylaxis by anti-cholera inoculations, and essential oils were adopted, and I think with success, as shown by the figures, 157 cases with 15,061 contacts.

CAMP HOSPITAL.

69. Sixty-nine patients remained at the beginning of the year. One thousand nine hundred and ninety-eight were admitted during the year. Of the total treated 398 died giving a percentage of 19.92. Many of the deaths were due to pneumonia, and lung complications following influenza and measles—this was particularly noticeable in the case of children.

INFECTIOUS DISEASES OUTSIDE THE QUARANTINE CAMP.

70. The following table shows the cases of infectious diseases reported and the State in which they originated:

f no till, and g	Sma	Smallpox.		Cholera.		gue.	Diphtheria.		Cerebro-spinal meningitis.	
State.	Cases.	Deaths.	Cases.	Deaths.	Cares.	Deaths.	Cases.	Deaths.	Савея.	Deaths.
Perak Selangor Negri Sembilan Pahang	1111		13 	-6 -	1111	I FIELD	14 19 13 5	6 4 - 2	17 9 4	5 8 2
Total	-		13	6	-	-	51	12	30	. 15

71. A small outbreak of cholera occurred in March in the Kuala Langat district, there being 13 cases and six deaths. A possible cause of infection was from one of a family who had arrived from India two months previously on an infected ship, these coolies—the parents and one child—had been inmates in the non-infectious hospital at the camp. This child was suffering from measles and was much debilitated, an examination of its stool was negative for cholera. It is possible that either this child or one of its parents was an intermittent carrier. These persons absconded from the camp on March 18th; structural alterations are being undertaken which it is hoped will prevent future abscondings.

- 72. There were no cases of plague reported.
- 73. Fifty-one cases of diphtheria were reported of which 19 were in Selangor, 14 in Perak, 13 in Negri Sembilan, and 5 in Pahang.
- 74. The figures for cerebro-spinal meningitis were 30 cases and 15 deaths. All were sporadic.
- 75. The number of deaths from dysentery and diarrhoea was 2,744, giving a death-rate of 2.09 per mille population, as against 1,945 and a death-rate of 1.34 per mille in 1925.

PULMONARY TUBERCULOSIS.

- 76. There were 1,873 deaths from pulmonary tuberculosis and the death-rate was 1.42 as against 2,001 and a death-rate of 1.38 in 1925
- 77. With regard to the four States (Pahang no figures)—Perak had a death-rate of 1.52 as against 1.49 in 1925, Seiangor had 1.14 as compared with 1.20, Negri Sembilan had 1.74 as compared with 1.65.
- 78. With regard to the four large towns—Kuala Lumpur had a death-rate of 1.36 as compared with 1.75 in 1925, Ipoh 1.08 as compared with 1.51, Seremban 2.24 as compared with 2.72 and Taiping 3.37 as compared with 3.81.

HELMINTHIC DISEASES.

79. The chief one of importance from a health point of view is ankylostomiasis. At least 90 per cent. of the native population harbour the worm in small numbers but few have sufficient to give rise to symptoms, unless they are also suffering from a concurrent disease such as malaria or dysentery when the combination is a very dangerous one. Ascariasis is also very common, and though in the majority of cases is not of serious import, is sometimes responsible for convulsions in children, and acute intestinal trouble.

SCAVENGING NIGHT-SOIL DISPOSAL, DRAINAGE, ETC.

COLLECTION AND DISPOSAL OF REFUSE.

80. Scavenging in most towns is fairly well done. The best disposal is by incineration; dumping or the filling of swamps is not so satisfactory. Prolonged wet weather hinders incineration and leads to accumulation of refuse, with consequent fly breeding, this danger also occurs in dumping unless very careful supervision is exercised.

COLLECTION AND DISPOSAL OF NIGHT-SOIL.

- 81. Up to date no town in the Federated Malay States has a water carriage system. Some Government and some private installations have been built and these are being watched carefully and the effluents examined with a view to determining the most suitable types of installation for a building or small groups of buildings, such as offices, bungalows or cooly-lines.
- 82. The system in use in most towns is the two-bucket system for the shop houses and the two-bucket system or pit latrines for the bungalows standing in their own compounds. Most of these pit latrines form fly breeding grounds and they are therefore unsatisfactory. Petroleum, lime and other larvicides have been used but unless applied lavishly do not prevent the breeding. The bucket system is being recommended for use on estate wherever possible in place of pit latrines.
- 83. The ultimate disposal in the bucket system is by trenching. If not carefully watched it becomes short circuited to the Chinese vegetable gardens.

DRAINAGE.

84. Street drainage is controlled by the Sanitary Board authorities. Anti-malarial drainage is controlled by the Mosquito Destruction Boards. Many miles of sub-soil piping have been done by the anti-malaria branch of the Public Works Department.

CLEARANCE OF BUSH.

85. Clearance of bush is done on public lands by Sanitary Boards or Mosquito Destruction Boards. Clearance of bush in this country is not always a profitable proceeding for at the wet hill foots it promotes the propagation of A. maculatus, the most powerful malaria carrier in Malaya. Many a healthy area has been rendered intensely malarial by the clearance of bush, in spite of the fact that notices pointing out this danger have been published in the press and local publications.

WATER SUPPLIES.

- 86. In the majority of cases towns are supplied with drinking water brought from uninhabited areas through pipes by gravitation. In some cases filters are employed, in others they have not been considered necessary. Both slow and rapid filters are in use.
- 87. The necessity for keeping catchment areas as free from pollution as possible has long been recognised and efforts have been made to ensure these areas being kept clear of trespassers by having them gazetted as "Forest Reserves."
- 88. Arrangements have been made with the Institute for Medical Research for the monthly examination of all water supplies to the larger towns in the Federated Malay States. The two supplies for Kuala Lumpur were as in previous years analysed weekly.
- 89. Some estates have excellent supplies, others especially those in the coastal areas are not so fortunate.

PUBLIC HEALTH EDUCATION.

- 90. The Public Health Education Committee met on several occasions during the year; at Horticultural Shows exhibitions were put up which were well attended, lectures were given and pamphlets which had been prepared by the Committee were distributed in large numbers in various languages.
- Popular lectures on Malaria were given in Maiay and Chinese in various parts of the country, these were illustrated by lantern slides and were well attended.
- 92. A cinematograph film of the work done at the Infant Welfare Centre, Kuala Lumpur was also exhibited on several occasions to large and interested audiences.

SCHOOLS AND SCHOOL INSPECTION.

93. The numbers of Government and State Aided Schools in the Federated Malay States are as follows:

Government English Schools	 ****	 	 23
Aided English Schools	 	 	 24
Government Vernacular Schools	 	 ***	 482
			529

Besides the above there is a considerable number of private schools.

- 94. The duties of school inspection were shared with the Medical Branch. Altogether 213 visits of inspection were made by the Health Staff.
- 95. Each school has a stock of Government quinine which is issued free of charge to those scholars who require it.

MATERNITY AND CHILD WELFARE.

- 96. The Infant Welfare Advisory Board met several times during the year and its minutes were published.
- 97. Infant Welfare Centres are established at Kuala Lumpur, Ipoh, Taiping and Seremban, the report of the work at each Centre has been published separately.
- 98. All the Infant Welfare Centres are under the supervision of Lady Medical Officers and have specially trained European Infant Welfare Sisters. Each has a staff of Asiatic nurses who work part time at the Centre and part time on the district. These Centres are very popular and there is no doubt are doing an immense amount of good.
- 99. There are three Maternity Hospitals which have been erected and maintained by the Chinese for the benefit of the poor of that nation. One is situated in Kuala Lumpur, one in Klang and another in Ipoh. These are very popular and are doing excellent work. At each there is a training school for midwives.
 - 100. On many estates attention is paid to Maternity and Infant Welfare.

WORK UNDER THE LABOUR CODE.

- 101. It is the duty of the employer of labour to engage the staff necessary for the protection of health and the cure of disease amongst his employees. The Health Branch co-operates with the Labour Department with the object of securing an adequate standard of sanitation where labour is employed.
- 102. The total number of estates is 1,450 and the total number of estate hospitals is 156, each of these estates should be visited at least twice yearly by the Health Officer of the district but this is unfortunately not always possible.

ESTATES.

103. Details of the distribution of estate and estate hospitals and the frequency of visits by Health Officers are given below:

					Estates.		Estate hospitals.			
State-	Health dist	riet.		No.		f visits H.O.	No.	No. of visits by H.O.		
45-10 10.00	F-100 54				1926.	1925.	7.00	1926.	1925.	
(Perak North	***		267	78	48	24	15	19	
Perak	Kinta			133	64	57	6	11	10	
(Perak South	***		255	92	116	24	18	23	
Č.	Selangor East			232	122	86	23	19	19	
Selangor	Selangor Coast			207	135	130	34	33	36	
(Bernam			4	7	6	2	2	2	
Negri Sembilan	All districts			352	227	131	43	56	40	
Pahang	All districts		***			35		***	15	
Total, F.M.S	S. (Excluding Pa	hang).		1,450	725	609	156	154	164	

Supplementary visits were paid by Sanitary Inspectors

104. The distribution of labour was as follows:

			Indians.	Others.	Total.
Perak	 Perak North		23,594	 4,050	 27,644
	Kinta		8,723	 892	 9,615
	Lower Peral Batang Pad		24,818	 2,398	 27,216
Selangor	 Selangor East		20,983	 5,689	 26,672
	,, Coast		45,012	 1,575	 46,587
	Bernam .		1,783	 1	 1,784
Negri Sembilan	 All districts		23,333	 18,353	 41,686
Pahang	 			 	 _
	Total, F.M	.s	148,246	 82,958	 181,204

105. The table below sets out the mortality rates among estate labourers during the past sixteen years, that is, since the Health Branch took over the supervision of health condition on estates:

Year.		Cond		Т	otal number state labourer	8.	Deaths.		Death-rate per mille.	
1911				***	143,614		9,040		62.9	
1912				***	171,968	5	7,054	111	41.02	
1913				***	182,937		5,592		29.6	
1914					176,226	***	4,635		26.3	
1915					169,100		2,839		16.78	
1916				***	187,030		3,299		17.61	
1917		,,,			214,972		3,906		18.71	
*1918					213,425		9,081		42.55	
1919			***		216,578		3,384	***	15.16	
1920	Marie In		V		235,156	***	4,867		18.57	
1921					175,649		3,195		18.19	
1922					159,279		2,556		16.05	
1923					147,276		1,924		13.06	
1924	***	***	444		144,902		1,514	***	10.45	
1925					146,558		1,585	1000	10.81	
1926					†181,204	2	†2,632		†14.53	

^{*} Influenza year.

[†] Excluding Pahang.

^{106.} There were 2,342 deaths among the 147,646 Indian estate labourers during the year, giving a mortality rate of 15.18 per mille as against 1,429 deaths and a death-rate of 11.95 in 1925.

107. Return of malaria admissions and deaths of Indian labourers and others in estates and Government Hospitals.

Divisions.	Maharia admissions to estates and Govt., hospitals.	Malaris deaths in estates and Govt, hos- pitals,	Total admissions to estates and Govt., hos- pitals,	Total labourers employed, all nationalities.	Total deaths.	Death-rate per mille.	Indians employ- ed.	Deaths, Indian labour force,	Death-rate, Indian labour force.	Number of estates.	Number of hos- pitals,
Perak North Kinta	5,741 2,018	113 64	14,812 4,586	27,644 9,615	381 108	13.78	23,594 8,123	364 107	15.42	267 133	24 6
Lower Perak and Batang Padang	2,960	83	11,375	27,216	247	9.07	24,818	239	9.63	255	24
Selangor East	7,878	178	19,450	24,672	410	15.37	20,983	392	18.68	232	23
Selangor Coast	5,561	202	15,889	46,587	653	14.02	45,012	644	14,30	207	34
Sabak Bernam	80	4	1,194	1,784	5	2.80	1,783	5	2.80	4	2
Negri Sembilan Pahang	9,819	296	20,460	41,686	828	19.86	23,333	591	25.33	352	43
Total	34,057	940	87,766	181,204	2,632	14.53	147,646	2,342	15.18	1,450	156

· No figures.

MINES.

108. The average population engaged in mining during the year was 110,293 as against 107,257 in 1925 or an increase of 3,036 due probably to the continued high price of tin.

Mines are not required to send in sickness and death returns and the sick-rates and death-rates are not known.

GENERAL.

- 109. The Chinese Decrepit Asylum at Port Swettenham was administered by the Health Branch for the second-half of the year, the inmates being moved from the permanent camp buildings into large attap sheds on the 22nd of June so as to leave more accommodation for Indian immigrants.
- 110. Owing to this accommodation being of a temporary nature admissions were stopped from June, so only 43 new cases were admitted during the year.

111. The admissions, etc., from various States are shown in the following table:

State.		Remained.	Admitted.	Total.	Discharged.	Transforred	Absconded.	Died.	Remaining.
-	-	-	4	F	-	F	4	- 0	1 74
Perak		235	21	256	5	63 30	33	34	121
Selangor		139	11	150	7	30	23	19	71 37
Negri Sembilan .		63	7	70	3	15	6	9	37
Pahang		4	4	8	1	1	5	1	
Total .		441	43	484	16	109	67	63	229

- 112. Of the 63 deaths, 41 were due to general debility, nearly 50 per cent. of inmates remaining at the end of the year were between 60-80 years of age.
- 113. Four cases of Japanese River Fever—Tsutsugamushi—were admitted into hospital, three were Europeans from an oil palm estate—20 miles from Kuala Lumpur—of these one case proved fatal. The fourth case was a European rubber planter who had been inspecting estates in various parts of the country.
- 114. A notable event in the year was the visit of Sir Ronald Ross, K.C.B., to the Federated Maiay States in December. Staying at Klang with Sir Malcolm Watson he paid visits to Kuala Lumpur on December 13th and 18th. On the former day various anti-malarial works, completed and in progress were inspected and a visit paid to the Malaria Bureau, after which the eminent malariologist was entertained to tiffin by the medical profession in Kuala Lumpur.

Kuala Lumpur, 16th March, 1997.

A. K. COSGRAVE,
Acting Senior Health Officer, F.M.S.

REPORT OF THE REGISTRAR-GENERAL OF BIRTHS AND DEATHS, FEDERATED MALAY STATES, FOR THE YEAR 1926.

The Records of Births and Deaths for the State of Pahang for the fourth quarter, 1926 and some incomplete registers of previous years were all lost in the flood of December 27th; reports from the other States indicate that registration is up to date.

The leaves forming the Original Register of Births and Deaths were received regularly and were bound in the office.

I attach a schedule of vital statistics, unfortunately those for Pahang are not available for the reason stated above.

KUALA LUMPUR, 22nd March, 1927. A. K. COSGRAVE, Acting Registrar-General of Births and Deaths, Federated Malay States.

FEDERATED MALAY STATES.

ESTIMATED POPULATION FOR 1926 OF ALL RACES OF EACH STATE AND FOR THE WHOLE OF THE FEDERATED MALAY STATES.

State.	Europeans and Americans.	Eurasians.	Malays and other natives of the Archipelago.	Chinese.	Indians.	Others.	Total.
Perak Selangor Negri Sembilan Pahang	 2,389 3,054 1,152 352	1,040 1,775 548 132	260,177 105,875 81,797 110,211	228,461 181,071 77,943 39,258	160,136 163,246 41,748 9,785	1,976 2,149 1,069 688	654,179 457,170 204,257 160,426
Total, F.M.S.	 6,947	3,495	558,060	526,783	374,915	5,882	1,476,032

SUMMARY OF BIRTHS AND DEATHS FOR THE YEAR 1926. BIRTHS.

State.	Europeans and Americans.	Eurasians.	Malays and other natives of the Archi- pelago.	Chinese.	Indians.	Others.	Total.	Rate per mille of popula- tion.
Perak Selangor	46 78	32 64	10,175 4,004	6,118 5,332	3,676 4,418	48 18	20,095 13,914	30.72 30.44
Negri Sembilan Pahang *	7	31	3,343	1,523	916	5	5,825	28.51
Total, F.M.S. +	131	127	17,522	12,973	9,010	71	39,834	30.28
			DEAT	Hs.				
Perak	16	8	6,433	7,287	4,727	35	18,506	28.29
Selangor	16	22	2,667	5,239	5,431	15	13,390	29.29
Negri Sembilan Pahang *	4	16	2,557	2,204	1,762	6	6,549	32.06
Total, F.M.S. +	36	46	11,657	14,730	11,920	56	38,445	29.22

^{*} No figures. † Total, F.M.S. excluding Pahang.

FEDERATED MALAY STATES.

RETURN OF DEATHS OF ALL AGES FROM PRINCIPAL DISEASES FOR 1926.

Total No. of deaths.	18,506	13,390	6,549		38,445	1
1/2 1/3 1811 110			1015	MALAH	-	
Persons of other ages.	8,627	5,865	2,820		126 4,297 9,286 7,718 9,489 8,976 17,312	
Persons over 60.	2,370	1,005	109		9,976	
Persons between 20.40.	4,050	3,632	1,757	bygood.	9,439	
Children under one year.	1,924 4,562 8,459 4,050	2,888		a selatina	812,7	
Deaths from other diseases.	4,562	1,609 3,339 2,888	764 1,885 1,871	1000	9,236	
Convulsions.	1,924	1,609	192		4,297	
Promaine poison- ing.	150	91	4			
Influenza.	64	50	66		86	
Bright's discase.	88	49	118		195	
Cancer (including Sarcoma).	00	1~	6		19	108.
Tetanus.	98	22	α	MINE A	129	+ Total, F.M.S. excluding Pahang.
Ankylostomiasis.	*8	101	22		65	luding
Syphilis.	98	88	10		200	exc
Enteric (typhoid).	=	10	-1		88	M.S.
Beri-beri.	108	140	99		314	al, F
Pneumonia.	865	1,201	507		2,744 1,878 2,578	+ Tot
Phthisis (pulmon- ary tuberculosis).	966	521	926		1,873	anes.
Dysentery and distributes.	1,027	1,211	506		2,744	No figures.
Blackwater fever.	03	00	00		00	
Malaria (including fever).	8,732	5,011	2,788		16,531	
Yellow fever.	:	:	1		:	
Typhus.	:	:	1		:	
Diphtheria.	9	4	. :		10	
Cerebro-spinal meningitis.	NO.	00	01		15	
Smallpox.	:	-	:		1	
Cholera.	:	9	:	ar .	9	
Plague.	:	:	:		1 :	
40017 100	1	. :	:	:	1.1	1
i	:	:	ilan	:	L.S. +	-
State	Perak	Selangor	Negri Sembilan	Pahang *	Total, F.M.S.	

ANNUAL REPORT OF SURGICAL DEPARTMENT, IPOH HOSPITAL, 1926.

he S	urgical Wards		196 01 11	ork pe	HOIM	eu, an	d expansion of a	umissio
	or Operations.							248
-	rui major op	CT GREEN CO.						
	48.8 per cent	. over 1924;	16.9 p€	er cent.	over	1925.		
	1924; 212 for	1925.			250.000		922; 140 for 192	3; 169 f
	nese cases are	exclusive o	t Ophti	almie	Cases			
	tality.—							977
	otal deaths fol	Control of the Contro	r operat	ions .				37
	ercentage 14.9		m't for	1000 -	16 nov	cont	for 1923; 13.69	ner cer
27.5	for 1924; 15.				to per	Come	. 101 1020 , 10100	Per or-
Eme	rgency Opera							
	Total number						73.	
	deaths	following					29.	
	Percentage		***				39.8 per cent	
	rval Operation							
	Total number						175.	
	Deaths						8.	
					***		4.6 per cent	b.
	or Operations.	-s lo lavon						
	Total operation						812.	
	l Operations							
	(Minor and M		***	***	***		1,060.	
Tota	l Surgical A	dmissions			***		2,791.	
	Total deaths	to admission	s				69.	
	Percentage of	deaths to	admissi	ons	***		2.1 per cent	
	n 1925.—							
	l Surgical a	Amissions	***				2,589.	
Can	se of death for	Ilowina Maio						
Cau	Skull (decom	nression)		****		Cere	bral haemorrhag	e
9	Excision of I	Haemangiom	a (axilla	(4		Acu	te Septicaemia	
3	Ligature Br	achial arter	v			Sapi	raemia from gan	grene
	Amputation						ek and haemorr extensive injury	
5.	Ligative fem	oral artery					ek and haemorrha ultiple injuries	age due
6.	Wiring fract	ured femur	(compou	ınd'			ek and haemorrha	ige due
7	Arthrotomy	hip					te dysentery	
	" ,						iemia	
9	Laparotomy-	-general peri	itonitis	(4 case	s)	Gen	eral peritonitis	
10.		suture of	perfora	ted gu	t (2		and the same of th	
10.		cases)		***			eral peritonitis	
	Laparotomy-	-suture of gr	ut (stab	woun	d)		eral peritonitis	
	.,	stab wound	d abdon	nen	***		morrhage	
	Post gastro j	Liver abces	ss (rupt	ured)	per-	Gen	eral peritonitis	
	forated gas	stric ulcer	and ed				eral peritonitis erforated gastric	
	77	Two others	plated	novite	nitis		eral peritonitis	THE PERSON NAMED IN COLUMN TO PERSON NAMED I
	Herniotomy- Appendicecto	my-gangren	ious a	ppendi	(4		Allelin Line	
	cases)		***	***	***	Gen	eral peritonitis	
	Herniotomy- of gut	-Ing. strangu	ilated ai	nd rese	etion		eral peritonitis fr	om gan
						re	nous gut	
	Splenectomy	and auto	no of	norfo	rated			

	Intussusception—excision		***		Gangre	unous g	ut	
	,, reduction					enteritis		
	Casarian section				Cancer			
	Excision of par ovarian cy					post op	erativ	re e
	Exploratomy Laporatomy					y carsino		
					Cancer	THE PERSON NAMED IN	ALLO U	
	Opening septic haematoms				Septica			
	Calostomy—chr. dysentery						O.Mark	
						c dysent	ery	
	Hysterectomy	***	***		Septics	iemia		
	Vesical calculi	***	***		OLD !	,		
	Subphrenic and liver abces	88	***	***	Toxaer	nia		
	Imperforate ani	***	***		**			
RE	GIONAL CLASSIFICATION O	F MA	TOP SITE	OCICAI	OPET	ATTONS	FOR	1008
	Head and Neck.—	E DEPE	on sur	MOTOA	OFER	ALIUNS	FOR	1020.
*-	Wiring—cleft palate							1
	Plastic operation—hare-lip	***		****				5
	Plastic operation—palate				***	***	***	1
		····			***	***		1
	Trephining skull (decompre			***	***	***	***	
	Skull—decompression				***	***	***	1
	Schuartze's operation—chro	nie ma	astoiditis			***	***	2
	Skin grafting on neck			***	***	***		1
	Excision of cyst of neck					***	***	1
	,, tumour of nec	k				***	***	1
	,, keloids (neck)		***		***		***	1
	,, endothelioma o	f neck	with re	emovai	of cer	vical gla	nds	1
	,, suppurating se	baciou	s cyst	(head)				1
	., Epulis (Rt. up				***	***		1
	,, haemangioma							1
	"							-
						Total		19
TT I	Innas Limb -							-
11. (Jpper Limb.— Ligature—brachial artery							1
		***	***	***				4
	Amputation—fingers	***	***	***		1110	***	3
	,, arm		***	***	***	***	***	2
	,, forearm	4 1-in-				***	***	2
	Disarticulation through wris				***	***	***	1
	Excision of lipoma (shoulde	r)	***	***	***		***	-
						Total		13
						2000		_
II. I	ower Limb.—					Otto a Fi		
	Ligature—femoral artery ar	id exc	ision of	femo	ral ane	urism	***	1
	,, external iliac (an	eurisn	a)		***	***	***	1
	,, femoral artery				***		***	1
	Arthrotomy ankle-osteomy	elitis (of astrag	galus s	and tib	18	***	1
	Arthrotomy-knee (suppura	tion)	***					3
	,, hip			***		***	***	5
	Wiring (fractured femur)			***	***	***	***	1
	,, (patella)		***			***	111	1
	Platiny (fractured femur)		***		***	***	***	1
	Disarticulation toe							1
	Amputation leg (faraboeuf's			2	1112345			4
	Sequestrectomy (osteomyeli	tis of	tibia)		***		***	1
	Skin grafting on dorsum of	foot						1
	Excision of pedunculated gro	wth o	f knee io				ion	
	of patella							1
	Excision of sarcoma (thigh)						***	1
	(log)			****				1
						7		1
	,, ,, eyst (knee join	marit	(lea)			***	***	1
	,, Haemorrhagic	cyst	(reg)				10000	-
						Total	***	27
								-

IV.	. Thorax.—	
	Resection of rib necrosed	1
	Stab wound of pleura, pericardium and heart stitching of pericardium	
	Excision (fibroma breast)	1
	of carbuncle	1
	Total	4
V.	Abdomen and pelvis.—	-
	Posterior gastro jejunostomy (with one case of suture of perforated gastric ulcer)	15
	Anterior gastro jejunostomy	1
	Herniotomy (inguinal)	30
	,, (inguinal irreducible)	1
	,, (inguinal-strangulated with two cases of resection	
	of gut)	7
	,, (femoral)	1
	,, (Umbilical—Repair—Mayo's operation)	. 1
	Splenectomy (Traumatic rupture)	1
	;, (Splenomegaly)	2
	,, and suture of stomach (stab wound)	1
	Appendicectomy	11
	,, with excision of strangulated omentum	1
	Appendix abcess	3
	., ,, with general peritonitis	1
	Opening extraperitonial abscess	1
	,, Subphrenic and liver abscess	1
	,, Liver abscess	4
	,, perinephric abscess	1
	" septic haematoma over bladder region	1
	Chalecystostomy	1
	,, for biliary calculi	1
	Intestinal anastomosis for strangulated hernia	1
	,, ,. (resection)	1
	Exploratomy Laparotomy	6
	Transpleural ,, Hydatid cyst (liver)	1
	Laparotomy—general peritonitis	7
	,, removal of protruding omemtum (stab wound)	1
	,, suture of perforated gastric ulcer	1
	,, ,, small rents in small gut (stab wound)	2
	,, ,, perforated stomach (stab wound)	1
	,, ,, perforated intestine	2
	" stab wound abdomen	1
	,, haemorrhagic omental cyst	1
	Talma Drummond Morrison's operation (Cirrhosis of liver)	2
	Supra public lithotomy	9
	Reduction (intussusception)	1
	Excision ,,	1
	Caesarian section	1
	Colostomy for Chronic Dysentery	1

	Craniotomy and forceps delivery (labour)			1
	Ruptured ectopic gestation		mound	4
	Supra vaginal hysterectomy			2
	,, and excision of Rt.	tube and	left	
	amaki.			1
	,, for myoma and excis	sion of ova	rian	
	cyst			1
				1
				3
	,, bilateral cystic ovaries		v	1
				1
				1
				1
				1
	Opening and drainage bladder (traumatic rupture)			1
		Total	***	145
VI.	Perineum.—			
	Ectopic testes (lodgment and fixation)			2
	Daylones of some (Continue)			1
	Delical constitut (IT-decade tonic contratt)			12
	Colporrhaphy—laceration and haematoma of vulva .			1
	Total amputation of penis			1
	Davingamahanhy			3
	Repair of urinary fistula		elech.	6
	P-4			2
	the state of the s			1
	77 11 6 11			8
	1		***	1
				1
	Plastic operation (conginital stenosis of anus)		***	1
	Imperforate anus		***	1
		Total	200	40
		a Orali		_

Total major operations performed by Chief Surgeon including those performed in other hospitals 279.

During the year under review, twelve extra beds were added to the operation ward, and these have been fully occupied. A further extension of this ward will become necessary at an early date.

During the year I was on leave for a period of 10 months, the work being carried on by Mr. Dannatt and Dr. Wilson.

LIST OF OPERATIONS PERFORMED AT TAIPING HOSPITAL BY CHIEF SURGEON, PERAK, DURING 1926.

Appendicectomy			***		***		2
Herniotomy	***			***			4
Cholecystectomy	en luien				***		1
Removal of sabacious cyst						***	1
Perforated gastric ulcer							1
Incision of perianal fistula					***	****	1
Excision of contracted scar							1
Tonilectomy					***	***	1
Removal of ovarian cyst		***			***	***	1

LIST OF OPER	RATIONS PERFORME					ITALS	BY C	HIEF
Annondia	SURGEON, 1			NG 192	16.			0
Appendic			***	***	***	****	***	2
	ion of finger	***	- 37-	Justin.	***	100		1
	peritonitis (gangreno		ndix-	-drain:	age	111	***	1
Circumci	Particular and Addition of the land			***	***	***	***	1
Appendic			***		***	***	***	1
	of sebacious cyst		***	***	***	****		1
Circumeis								
	d gastric ulcer—gen		ritoniti	is—sat	ure an	d drai	nage	1
Splenecto			***	***	***		***	1
	of psoas abscess							1
	of intraligamentus		appen	dicecte	omy a	nd ve	ntral	
suspens			***	***	***	***	***	1
	of epithelioma (low		***	***	***		***	1
	of glands of neck				***	***	***	1
	of carbuncle back of		***	***	***	***	***	1
	of recurrent epithel	ioma of	lower	lip	***	***		1
				***	***		***	1
	of area of chronic n	nastitis	***	1100	100 C	100		1
Removal	of haemorrhoids	***						1
Appendic	ectomy				***		***	1
Drainage	of appendix abscess							1
Examinat	tion of shoulder—join	nt unde	r anae	sthesi	а			1
Drainage	of appendix abscess							1
Incision	and drainage of cell	ulitis of	neck		***			1
Radical c	eure for hydrocele							1
Caesarian	section				100	***	***	1
Inguinal	hernia-radical cure		James		1 2.2	all		1
Appendie	ectomy				***			1
	nic case							1
Extractio	n of teeth							2
	f perforated pyloric							1
	tion of haemorrhoids			Jejan		(Fees.		1
Circumcis								1
the same of the same of								
Ірон,	200					and the same	PASLI	
7th January, 15	127.				Chi	et Su	rgeon,	Perak.

ANNUAL REPORT OF THE CHIEF SURGEON, SELANGOR.

Mr. T. W. H. Burne was acting as Chief Surgeon until the end of April when he proceeded on leave. For the remainder of the year Mr. R. M. Dannatt acted for him.

The operations performed at the three hospitals were as follows:

					Major.		Minor.		Totals.
General			***		189		218		407
European				***	46		62	***	108
District	***				71		265	***	336
		*.							051
					306	***	545	***	851

SYNOPSIS OF MAJOR OPERATIONS.

	I	-AT THE	EURO	PEAN	Hos	SPITA	L.
(a)	Abdominal—						
			Total.		Deaths		Remarks.
	Appendicectomy		. 14		-		
	Appendicectomy with dra	inage	. 3		-		
	Closure of perforated ga	astro-jeju	-				
	nal ulcer and drainage	for gene	-				
	ral peritonitis				1		Gastro-jejunostomy done in Europe 12 months pre-
							viously
	Rammstedt's operation						Control to the Control of the Contro
	genital pyloric stenos	is	. 1		-		Infant 6 weeks old
	Cholecystectomy	4-5	. 1	***	-	***	For chronic cholecystitis with gall stones
	Cholecystostomy		. 1		1		
	Lumbar nephrectomy		. 2		1		
	Relief of acute intestinal	obstruct	_				
	ion due to adhesions				_		
	Ileo-colostomy		. 1		1		For excluding inoperable
							car-cinoma of splenic
							flexure of colon
	Hysterectomy		. 1		-		For multiple fibromyomater
	Sling operation for retro	version o	đ				
	uterus				-		Chronic appendix also excised
	Ovariotomy		. 1		-		
	Salpingo-oöphorectomy		-		-		For ruptured ectopic gesta- tion
	Exploratory laparotomy		1		-		
(6)	Cranial-						
	Elevation and removal						
	ments. Comminuted						
	fracture of skull	***	. 1		-		
(c)	Limbs-						
1-1	Wiring fracture of ulna		1				
	Plating tibia		1		_		
		semiluns					
	111				_		
	cartinege			***			
(d)	Perineal						
	Operations for haemorrh	oids .	5		-		
	Operations for fissure in		2		-		
			1		-		Excisions of fistulous tracts
			1				
	Radical cure of hydrocel		1				
	Radical cure of varicoce	le .	1		-		
(e)	Thorax-						
(0)	Excision fibroadenoma of	breast	. 2		_		
	Zaciolor introductionia of	Diones					

II .- AT THE GENERAL HOSPITAL

1-1	II.—AT THE	GEN	ERAI	Ho.	SPIT	AL.
(a)	Abdominal—	Total		Death		Remarks.
	Stab wounds of abdomen	1		Deats	18.	Suture of stomach
	Stab wounds of abdomen	4				Without visceral injury
	Appendicectomy	9		_		Without drainage
	Appendicectomy	3		_		With drainage (for gangre-
						nous appendicitis with
	Drainage operations for general					peritonitis)
	peritonitis of appendicular origin	6		4		
	Drainage operations for localised			1115		
	appendix abscess	4		-		· · · · · · · · · · · · · · · · · · ·
	Drainage operation for retroperi-					
	toneal iliac abscess	1		_		
	Suture of perforated gastric ulcer					
	and drainage	2		1		General peritonitis
	Posterior gastro-jejunostomy	7			***	Five for chronic gastric, 2
	Projector of concessors intus					for chronic duodenal ulcer
	Excision of gangrenous intus- susception and intestinal anasto-					
	mosis	1		_		
	Intestinal anastomosis for volvulus	1		1		
	Partial excision of jejunum and					
	lateral anastomosis	1		1	***	
	Radical cure of inguinal hernia	22		2000		jejunum
	Operations for strangulated ingui-		***	-		
	nal hernia	7				
	Intestinal anastomosis for strang-					
	ulated inguinal hernia	2		1	***	With general peritonitis
	Operations for strangulated fe-					
	moral hernia Operations for ventral hernia	1		1		
	following appendix abscess	1		1		Died of meningitis
	Cholecystectomy	2		1		Died of meanights
	Cholecystostomy	1		1		For gangrenous cholecys-
						titis
	Drainage operations for peritonitis					
	complicating liver abscess	2		1		P turn I mlan (late)
	Splenectomy	1 4		1		For ruptured spleen (late) Three for inoperable car-
	Exploratory laparotomy	4		1	***	cinoma, 1 for tubercu-
						lous peritonitis
	Ovariotomy	2		-		Banki to at talonal
	Salpingectomy	3		-		One hydrosalpinx, two
	Huntanatara	Q				ectopic gastation
	Hysterectomy Caesarean section	3 2		2		For multiple fibromyomata One in obstructed labour
	Caesarean section	-		-	***	due to atresia vaginae.
						One after prolonged in-
						trauterine manipulations
	Perineal—					
	Operations for haemorrhoids	9				
c) (Genito-Urinary—					NEWS DO DESCRIPTIONS
	Suprapuble lithotomy	5		-		
	Suprapubic cystosstomy	1		1		For gangrenous cystitis
	External urethrotomy	4	***	-	222	Three for stricture one for
	Radical cure of hydrocele	2		113		traumatic rupture
	Excision of tumour of spermatic	-	***			
	cord	1		-		
1	Plastic operation for amputation	1				
	of penis (self-inflicted)	1		-		

(d)	Gynaecological—	-				
		Total		Deatl	is.	Remarks.
	Dilation and curettage	9	***			
	Excision of fibromyoma of cervix	1	***	-		
	Excisionof vulva	1		-		For carcinoma
	Perineorraphy	1	***			
	Plastic operation atresia vaginoe	1	***	-		
13	0					
(e)	Operations on Head and Neck-					
	Elevation of fragments in com-	4		0		
	pound fracture of skull	4	***	2		
	Schwartze's operation for mastoid disease	5		1		
	Heath's operation for mastoid		***			
	disease	3		_		
	Excision malignant glands in neck	1				
	Excision of tuberculous glands of		***	-		
	neck	2				Posterior prefer pulse to
	Excision of malignant growth of	-	***	- mb		money to make the
	parotid	1	***	_		
	Tonsillectomy	9		_		Five with adenoids
	m l t	2		2	1	One late case diphtheria,
	Tracheotomy	-		hor	866	one foreign body in
						trachea
	Excision naso-pharyngeal fibroma	1		_	00.01	With preliminary laryng-
					1 1	otomy
	Nasal drainage for empyema of	-				
	maxillary antrum	2	***	-		
	Operations for hare-lip	2	***	-		
	Partial excision of lip for lymp-					
	hangioma	1		-		
	Excision of submaxillary tumour	1		-		
	Excision of sublingual	3		_		
	Kronlein's exposure of orbit	1		-		For inoperable orbital
	D					tumour
	Excision dermoid cyst	1	***			
111	Operations on Thorax—					
0,	Exploration of stab wound of chest	1				With resection of rib
	Drainage operations for empyema	3	***	-	***	"
	Resection of tuberculous rib	1		-		
	Excision of breast	1	***	1		
	Excision of lipoma of breast	1	***	-		
	Excision of bone in malunion of					
	clavicle	1		-		
(0)	Operations on Limbs—					
(9)	THE PERSON NAMED IN COLUMN TWO IS NOT THE PERSON NAMED IN COLUMN TWO IS NAMED IN COLUMN TW	1				For non-union
			***		***	For non-union
	Osteotomy for mat-united humerus	1	***	-		
	Amputations of arm	1	***	-		7
	Disarticulation of knee	1	***	-		For compound fracture
	Amputation of leg	6				,, and gang-
	Operative replacement of lower					rene
	femoral condyles (fractured)	1		-		
	Excision of inter-phalangeal joint	1				For anchylosis
	Sequestrectomies			300	1000	The state of the s
		2	***	NI TO		
	Removal of bullet from thigh	1				
	Suture of patella in compound fracture	0		1		
	fracture	2	***	-		

III .- AT THE DISTRICT HOSPITAL

(a)	Abdominal—	Dis	TRIC	T H	OSPITA	AL.	
	Drainage energtions for posterotal	Total		Death	18.	Remarks.	
	Drainage operations for perforated gastric ulcer	1		1		General peritonitis	
	Posterior gastro-jejunostomies	111				For chronic gastric ulcer	
	Exploratory laparotomy	1		1		Intestinal obstruction in tuberculous peritonitis	1
	Drainage operations for gangrenous cholecystitis	1		1	100	General peritonitis	
	Splenectomy	1		_		For rupture	
	Appendicectomy and drainage	1		1		Property of the second	
	Radical cure of inguinal hernia	12		_			
*	Operations for strangulated ing- uinal hernia	1		1			
	Operations for stab wounds of						
	abdomen	2		_			
	Drainage operations for liver						
	abscess	4	***	4			
	Inguinal colostomy	1		1	***	In operable carcinoma of rectum	E
(b)	Cranial—						
	Decompression in tracture of skull	1	***	-			
	Operations for acute mastoid disease	4		1			
(c)	Operations on Limbs-						
100	Amputation of arm	1		1			
	Excision of lipoma	2	***				
	Excision of fibroma	1		_			
(d)	Perineal—						
	Operations for haemorrhoids	11		-			
(e)	Operations on Thorax-						
	Drainage operations for empyema	6		3			
(f)	Genito-Urinary—						
47	Suprapuble lithotomy	5		_			
	Suprapubic cystostomy	1		1		For carcinoma of bladder	
	Radical cure of hydrocele	8	***	_			
	External urethrotomies	3		1			
	Amputation of penis	1		-			
	AT THE PARTY OF TH					R. M. DANNATT,	

R. M. DANNATT, Acting Chief Surgeon, Selangor.

ANNUAL REPORT OF THE CENTRAL MENTAL HOSPITAL, TANJONG RAMBUTAN.

Sir,—I have the honour to forward the sixteenth annual report of the Central Mental Hospital, that for the year 1926.

	Hospital, that for the year 19:	60.	Males.		emales.		Males.		emales
2.	There remained on December	31st.	minies.		emales.		maies.		emaies
-	1925		1,145		306				
	Admitted during 1926	***	489		155				-
	Discharged—Recovered		151		56				- 130
	Relieved		56		20				
	Not improved	***	54		22				
	Not insane		1						
	Absconded		78						
	Died	***	82		25				
	Remaining on 31st December,	1926					1,212	195.0	338
84									
1511	Remained on 31st December,	1005	115		107				
	Admitted during 1926	1020	1	***					
			-	***	2				
	Di-J	***	8	***	4				
	Remaining on 31st December,	1008	0	***	*		108		101
		1020	1 30	***	TORK	***	100	***	101
Cri	iminals.—								
	Remained on 31st December,	1925	83		-				
	Admitted during 1926	***	38	***	1				
	Discharged ,,		17	***	1 .				
	Died ,,	***	6	***	-				
	Absconded ,,		2	***	-		-		
	Remaining on 31st December,	1926	-	***	-		96	***	
Ke	dah.—								
	Remained on 31st December,	1925	104		29				
	Admitted during 1926		40	***	11				
	Discharged ,,		7		6				
	Died ,,		4		3				
	Absconded ,,		5		-				
	Remaining on 31st December,	1926	-				128		31
Ke	lantan								
	Remained on 31st December,	1925	5		1				
	Admitted during 1926		2						
	Discharged ,,		1		1				
	Remaining on 31st December,				-		6		1
	Total remaining on 31st Decem							****	1000
	1926		_	***	_		1,550		471
	Percentage of recoveries	***		32.15					-
	,, ,, deaths on total	trea	ted	. 5.	01				

- This shows an increase of 126 against 150 last year and 163 in 1924. So that although there is an increase in numbers, the increase is much less than it has been of late.
- 4. Again the increase in Federated Malay States patients is also less than last year being 99 against 131 last year and 166 in 1924. This, too, is a cause for satisfaction, as we cannot hope to look for an actual reduction in numbers.
- The Singapore patients decreased by 13, which is only to be expected as we have ceased to take patients from Singapore. Even so we still have 209 colony patients.
- 6. The Kedah patients increased by 26, against 15 last year and 22 the year before. So that Kedah has to take part of the blame for our increasing numbers.
- Kelantan patients increased by one, but there are no figures to compare this
 increase with, as we only began taking Kelantan patients in 1925.

- Criminal patients showed an increase of 13 against 6 last year and only 2 in 1924.
- Admissions.—The total of admissions during the year 1926 was 735, which shows an increase of 69 on last year, when the numbers were 666. In 1924 the admissions were 614.
- 10. In the statistical tables which follow I shall deal, as usual, only with Federated Malay States patients.
- 11. We admitted last year 644 patients, made up of 489 males and 155 females which are both records. It is a remarkable fact that the admissions increase steadily year by year. Of course our general population is increasing, but I think also the institution is becoming better known year by year, and people are more willing to bring their relations, especially now that they have not to send them through the police courts.
- 12. The month which gave us the most admissions was October, in which there were 73, while January was second with 63 and November third with 62. Last year May showed the greatest number of admissions, followed by July and August.
- 13. It is remarkable that October and November showed such a large number of admissions as they are usually two of the months with small numbers of admissions. July and August are usually the months which give the heaviest admissions. I am inclined to think that this increase in the last quarter is due to the wave of influenza we have had lately, which though rarely fatal, was at times fairly severe. The centres which sent us most cases were Kuala Lumpur, Ipoh and Taiping—all urban.
- Primary dementia again heads the list of forms of disease on admission, with recent melancholia second and recent mania third.
- 15. General paralysis of the insane has fallen to fifth place, while confusional insanity has gone into fourth. Even though general paralysis has gone to fifth place, it still shows 59 admissions.
- 16. Seeing that the classification of metal disease is not yet all that one would like to see it, we cannot place too much reliance on the primary dementia figure, as the personal equation is, as things are at present, very prone to have an undue effect, and what one man diagnoses as primary dementia might by another be classed as something else.
- 17. Still, there is no doubt there is an alarming increase in primary dementia, which should be taken as a warning that education might with advantage be taken a little more slowly. I have too many of the derelicts left in the track of the storm of modern education.
- 18. Three voluntary boarders were admitted during the year. I look as time goes on to see a large increase in this class of patient.
- Discharges.—The discharges numbered 360 of whom 207 were discharged recovered and 76 relieved.
- 20. Of the balance of 77, seventy-six were discharged "unimproved" under bond, and were in the vast majority of cases removed by their relations who were returning to their own country. One was discharged as not insane.
- The total 369 shows a great increase on last year when the numbers were 276, while in 1924 they were 232.
- 22. A large number of those discharged "relieved" would, had they been kept a little longer, have been discharged "recovered", and would have thus swelled the recovery rate, but I thought it better to allow them out to complete their convalescence at home, where their friends were anxious to have them, and obviously intended to care them.
- 23. By this means one hopes to encourage friends to bring their relations early, and also to remove the asylum idea.
- Recoveries.—Those discharged recovered numbered 207 against 177 last year and 204 the year before.
- 25. The recovery rate works out at 32.15 which is slightly higher than last year when it was only 31.38 but is lower than 1924, when it was 37.99.
- 26. The recovery rate is not as high as it should be by any means and I hope to see it greatly improved next year.

- 27. There are various reasons, some of which I have already taken steps to deal with, while another cause I mentioned in paragraph 21, that is, allowing patients out to complete their recovery at home. Yet another is that we have not enough room in our acute ward (The Conolly Norman Ward), although I have been asking to have this ward improved and enlarged for the last seven or eight years. Again, of late the number of incurable diseases, i.e., general paralysis of the insane, senile and secondary dementia has increased, and naturally reduces the recovery rate. Also the floods at the end of the year prevented the sending home of 8 or 10 patients, who had recovered.
- 28. I mentioned in paragraphs 11 and 12 the very large increase in admission in the last three months of the year, and this too tends to reduce our recovery rate, as patients admitted from October to December do not stand much chance of being discharged before the end of the year.
- 29. Of the forms from which those discharged recovered were suffering on admission, recent melancholia shows the largest number of recoveries with 57. Recent mania comes next with 50, and primary dementia third with 42, bracketed with confusional insanity. If, however, we take the proportion of recovery to incidence, we find confusional insanity the most recoverable with recent mania and recent melancholia second and third; while primary dementia shows a very small recovery rate, and even of the small proportion one wonders how many will recur.
- 30. Aetiology.—Turning to the aetiological table (No. 4) we find "gastro intestinal system" has been displaced at the head of the table by "haemopoitic system". Cardio vascular degeneration comes second and syphilis third.
- 31. Alcohol takes fifth place appearing 78 times against 71 last year. This slight increase about corresponding to the increased number of admissions, so alcohol about holds its own as a cause; but last year and this tell a very different story to 1924 when alcohol appeared only 38 times.
 - 32. Of course alcohol as a cause tends to increase during times of prosperity.
 - 33. As a contrast "privation and starvation" has quite disappeared.
- 34. A point to be noted with regard to syphilis and alcohol is that they appear more frequently as a primary cause than as a secondary, whereas the other leading causes appear more often as contributary. Alcohol appears twice as often as a primary cause, while syphilis appears almost four times as often as a primary cause than as a contributary.
- 35. A great many of the cases of cardio-vascular degeneration, which is above both syphilis and alcohol, were due to either one or other of them.
- 36. Under "deaths" I shall again touch on alcohol and will, as I have done for years, point out the danger of turning the Chinese towards alcohol. There is no need to repeat all I have previously said, but I see no reason whatever to change my opinion on the subject.
- 37. However, I would again give a warning that the day is coming, if the consumption of alcohol increases as it has done, when we shall have to deal with the crimes attributable to alcohol; and this before many more years have passed.
- 38. Of the causes which appear less frequently "critical periods" (adolescence) and mental stress make a total of 69 between them and this taken in conjunction with the amount of primary dementia dealt with leads one to fear that all is not well with the adolescent. Education beyond their capacity is responsible for a good many of the cases I fear.
- 39. Malaria appears 54 times against 43 times last year, but it only appears 9 times as a primary cause.
- 40. I am inclined to believe that a considerable number of the cases admitted during the last quarter was due to influenza.
- 41. My reasons are: first that there was a very much larger number of cases admitted in the last quarter than any of the other three, which is unusual.
 - 42. Again the large number of melancholics admitted during the last quarter.
- 43. In the first nine months we had 78 cases of melancholia admitted, whereas in the last three we had 35, so that nearly half as many were admitted in the last three months as in the first nine instead of one-third.

93

- 44. There must have been some reasons for such a large increase in the number of admission during the last three months. We know there was influenza about, and also know that influenza is a disease which tends to produce mental trouble; and we also know that the form of mental trouble usually caused by influenza in melancholia. Consequently I think we are justified in attributing the increased admissions during the last three months to influenza.
 - 45. We admitted seven cases which could definitely be attributed to influenza.
- 46. Deaths.—The total deaths were 132 giving a death-rate on total treated of 5.01 per cent. and on daily average 6.61 per cent. This is the smallest we have ever had, being less than last year when the figures were 5.31 and 7.1, respectively. In 1924 the figures were on total treated 5.96 and on daily average 6.66 per cent.
- 47. The three principal causes of death were general paralysis of the insane which encountered for 47 deaths, pulmonary tuberculosis with 16 deaths, and dysentery 15 (bacillary 13 and amoebic 2). General paralysis of the insane shows an increase of 5 deaths on last year. The extraordinary increase in G.P.I. is one of the most remarkable facts of late years as far as our patients are concerned. It must be remembered, too, that in every possible case, probably 90 per cent., the diagnosis is confirmed by post-mortem examination. The Chinese, too, have almost a monopoly of G.P.I. as we had only one Tamil G.P.I. last year and no Malay, and that is the finding year after year.
- 48. There must be some reason for this increase in G.P.I., and also for the fact that the Chinese alone, one might almost say, suffer.
- 49. All nationalities, I think I might, say, suffer equally from syphilis, and years ago I put up a possible suggestion that the fact that the Chinese participated so much more than the other nationalities in speculative business, making greater demands on their brains might account for the succeptability of the Chinese to G.P.I. Now we must find a cause for the remarkable increase in G.P.I. amongst the Chinese.
 - 50. What is the new factor?
- 51. I should say without hesitation alcohol. Syphilis has always been present or rather was present long before the Central Mental Hospital was thought of. Yet up to 1912, the existence of G.P.I. was denied. Its presence was proved in 1915 on cases collected in 1912, 1913 and 1914 though the cases were few, but they grew in number steadily and, since the war, have gone up by leaps and bounds year after year.
- 52. Sixteen years ago the Chinese had their opium and did not need alcohol. Since then opium has been, in many cases, put out of the reach of the Chinese, and discouraged in the case of those who could afford it. The result has been a steady increase in the consumption of alcohol amongst the Chinese. Along with this increase in the consumption of alcohol we had a steady increase in G.P.I. It is known that alcohol is, in the presence of syphilis, a contributary cause of G.P.I. and I am convinced that the cause of the increase of G.P.I. in the Federated Malay States is the increased consumption of alcohol by the Chinese.
- 53. Deaths from pulmonary tuberculosis have decreased by three —16 against 19. Taken in conjunction with our increased numbers I think we must say that pulmonary tuberculosis is not on the increase at Tanjong Rambutan, although it is well known that a mental hospital population is more prone to pulmonary tuberculosis than almost any other.
- 54. Deaths from dysentery too, are less though only by one, but at least we may say that we are not letting dysentery get the better of us.
- 55. Malaria only accountered for four deaths, which, seeing the extremely hard fight we had with malaria, is most satisfactory.
- 56. Suicides.—I regret to say there was one suicide. This was a case of a woman who hanged herself. This case was the result of gross negligence on the part of the night head attendant and the night attendant of the ward, as a result of which they were both dismissed.
- 57. Fatalities.—I regret to say fatalities numbered eight, which is much too long a list. Six of those were due to rupture of the spleen, four due to falls on projecting corners or the edge of bedboards, and two to blows from other patients. Of course the prevention of these—one might call them accidents—is extremely difficult. When one remembers how easily a large spleen is ruptured, one can realise the proposition one has to face.
- 58. There was a case of ruptured liver due to a kick from another patient while another patient died from shock following extensively fractured ribs due to a kick from another patient. To one who has seen some of the terrible kicks given by Chinese—a kick delivered with the heel, one might say straight from the hip—it is quite conceivable that one kick will cause death, but I am convinced that had we better attendants we would have fewer of these fatalities.

- 59. It is a remarkable fact that six out of the above list occurred in the first half of the year and five on the first three months. Attendants found that such happenings in the wards, although negligence could not be brought home to them, did not make life any easier for them.
- 60. Abscondings.—Eighty-five patients absconded during the year. This is a very large number indeed, and two more than 1925, but there were a great many more patients out at work in 1926 than in 1925.
- 61. This number is too large of course, but as I have said, many times before, I prefer abscondings to prison conditions. Of those who absconded many came back of their own accord, after having had a walk and a look round. Others were brought back by our own attendants, and others again were brought back by the police; while some came back on new certificates. Very few indeed were those of whom we heard no more.
- 62. Although I said before, I prefer abscondings to gaol conditions, I do not pretend that many of these abscondings were not due to rank carelessness, and those guilty of such were dealt with either by me or by the courts.
- 63. Criminals.—There were 38 criminal admissions in 1926 against 25 in 1925, and 21 the year before. Of these 17 were discharged against 11 last year and 13 in 1924.
- 64. Two criminals absconded. There is no excuse for a criminal absconding, so I charged the attendant responsible before the Magistrate, and I am pleased to say the court inflicted an adequate sentence on the man,
 - 65. Seven criminals were discharged as not insane.
- 66. The criminal work is a considerable strain, and also entails a considerable amount of travelling, occupying time which would be more profitably employed at the Central Mental Hospital.
- 67. The admissions from Kedah numbered 51 (40 males and 11 females) against 48 in 1925 and 63 the previous year. Seven males and six females were discharged and six males died. The Kedah patients showed no marked difference to the Federated Malay States patients, and now come in under an Enactment identical with our Mental Disorders Enactment.
- 68. New buildings.—The only new building undertaken was a double quarters for clerks, which was considerably overdue.
- 69. The only work being done, save the clerks quarters, is the conversion of the single rooms to bedrooms, by which means I hope to give some of the better patients rooms of their own.
- Farms.—The number of farms continues to increase and we have now 14, accommodating 300 patients.
- 71. The farms are, as I have before explained, places where the patients are to all intents and purposes free. There is an attendant in charge of each, who sees to the general wellbeing of his people, and keeps one informed on the condition of the various patients.
- 72. In addition to the neucleus of chronics, there is a continual stream of convalescent and recovering patients passing through on their way to the outside world.
- 73. A few of these patients wander away, but they almost invariably return next day or so, but, owing to the fact that as soon as their absence is noticed they are reported as absconded, they help to swell the number of abscondings.
- 74. The value of products sent in from the farms reached the total of \$56,013.28 so that it can be seen that, in addition to the curative value of the farms, there is a considerable saving to the various votes due to the activities of the patients.
 - 75. I attach a list of products supplied by the farms.
- 76. I also attach tables showing the work done in the workshops and sewing room, and their value.
- 77. The forest planting has progressed, and the plantations are doing well, but I have been unable to obtain more batai seeds from the Forest Department, who have now advised me to plant rain trees. Had I only known years ago that rain trees made good firewood. I could have now had a large area of timber almost ready for use, as the rain tree grows very rapidly.
- 78. Anti-malarial-work.—The work of draining, filling and oiling went on steadily, and good progress was made. In this table it will be noticed that Sungei Bulat Nos. 1, 2 and 3 are mentioned. No. 1 is the main stream and No. 2 and 3 are tributaries.

- 79. It will be noticed that only 520 feet of inverts were laid in the main stream against 533 in 1925, but then for the first two months of 1926 we were digging out inverts which had been laid and buried by sand in 1925, again we laid 436 ft. of cement inverts in No. 2 and 274 in No. 3, a total of 1,230 feet; the banks were all sloped and sodded in addition.
- 80. It was necessary in a great many places, too, to put in sub-soil drains to catch seapage which showed in the sloped banks.
 - 81. In places the cuttings were fairly deep, at times being as deep as 10 feet.
- 82. We were fortunate that all our bunds held through the great rains at the end of December, with the result that we did not have our work smothered in sand, and were able to continue work as soon as the rain finished. One large bund was on the point of giving way when one of the external attendants discovered it, and we had time to put every available man on and so prevents its going. Had it gone it would have undone months of work.
- 83. As a result of all this work the place is being steadily dried up, and the mosquito breeding places are being removed.
- 84. However, we had a great deal of malaria during 1926 which caused considerable worry. At first I thought the trouble was due to a bad tindal in charge of the Mosquito Destruction Board coolies who now do the oiling. In fact I myself found unoiled breeding places. The Chairman Mosquito Destruction Board was good enough to replace this tindal by another man who did better, but a third man had to be sent down, who again improved on the second. Still we had too many cases of malaria. Then a comprehensive mosquito survey was made both inside and outside the reserve, and it was found that the malaria was not coming from the reserve or the railway line, but from places which had never been taken into consideration in neighbouring kampongs and estates, and along the river banks.
- 85. The Chairman Mosquito Destruction Board was good enough to send down an overseer to take charge of the work and I must say the improvement was extraordinary, and we have now little or no malaria.
- 86. It will be remembered that the amount of malaria increased about the middle of 1925 and continued more or less constant to the end of the year.
- 87. In January, 1926, we had 117 cases altogether. The number went down during the next three months and was 39 in April. May and June again were very bad with 126 and 130 cases, respectively, but from July—(97 cases)—to the end of the year things improved from 38 in August to 13 in December. The great improvement coincided with the appointment of the overseer and the extensive mosquito survey; the earlier improvement with the removal of the first tindal.
- 88. The malarial incidence was 1.9 on cases and 1.6 on individuals against .73 and .66, respectively in 1925, but it must be remembered that it was an extremely bad year for malaria all over the country.
- 89. Experiments were carried out with the treatment of general paralysis of the insane by malaria but the results were anything but hopeful.
- 90. Why the malaria treatment should be apparently useless in this country it is hard to say, unless it is that most of the inhabitants of the country having had a considerable amount of the malaria an antibody has been formed, which interferes with the action on the spirochaetes which takes place in a malaria free individual.
- 91. However I shall not try to discuss this subject further, as I believe Dr. Wilson, the Assistant Medical Superintendent, intends writing a paper on it.
 - 92. Staff.—The following movements took place amongst the staff:
 - S. Subrahmanyam, Assistant Surgeon, Central Mental Hospital, transferred to District Hospital, Taiping on 1st April, 1926.
 - T. Arumugam, Assistant Surgeon, District Hospital, Taiping, transferred to Central Mental Hospital on 1st April, 1926.
 - B. Gopal Menon, Assistant Surgeon, Central Mental Hospital, transferred to Kuala Lipis on 1st July, 1926.
 - G. A. Lopez, Assistant Surgeon, Port Dickson, transferred to Central Mental Hospital on 1st July, 1926.
 - Mr. E. Mathieu, Agricultural Officer, Central Mental Hospital, transferred to Hill Gardens, Taiping on 1st January, 1926.
 - Mr. G. E. Jones, Superintendent, Hill Gardens, Taiping, transferred to Central Mental Hospital on 1st January, 1926.

Miss C. M. Ortega, Nurse, Central Mental Hospital, resigned on 11th February, 1926.

Miss S. E. Thompson, Nurse, Central Mental Hospital, appointed on 11th February, 1926.

Miss S. E. Thompson, Nurse, Central Mental Hospital, resigned on 15th July, 1926.

Miss Lee Dee Yin, Nurse, Central Mental Hospital, appointed on 1st August, 1926.

Veeramah, Female Assistant Head Attendant, dismissed on 10th February, 1926.

Coopamah, Female Assistant Night Attendant, appointed on 1st March, 1926.

Sui Raj, External Head Attendant, appointed on 1st September, 1926.

V. A. L. David, Clerk, Class III, Central Mental Hospital, transferred to Agricultural Field Officer's Office, Taiping on 1st July, 1926.

See Seng Khim, Clerk, Class III, Perak Secretariat, Taiping, transferred to Central Mental Hospital on 1st July, 1926.

Tan Seang Hong, Clerk, Class III, Central Mental Hospital, left the Federated Malay States Government to join the Straits Settlements Government service on 7th August, 1926.

Khoo Soo Chee, Clerk, Class III, Revenue Survey Office, Taiping, transferred to Central Mental Hospital on 7th August, 1926.

See Seng Khim, Clerk, Class III, Central Mental Hospital, transferred to State Engineer's Office, Taiping on 4th October, 1926.

Lam Chong Choong, Clerk, Class III, Anderson School, Ipoh, transferred to Central Mental Hospital on 4th October, 1926.

Khoo Soo Chee, Clerk, Class III, Central Mental Hospital, transferred to State Engineer's Office, Taiping on 1st December, 1926.

Lim Chooi Tee, Clerk, Class III, State Engineer's Office, Taiping, transferred to Central Mental Hospital on 1st December, 1926.

S. Saravanamuthu, Steward, went on 42 days' vacation leave from 8th January, 1926 to 18th February, 1926 and three months and 10 days' half-pay leave from 19th February, 1926 to 28th May, 1926.

Kesari Rai, Head Attendant, went on 42 days' vacation leave from 22nd February, 1926 to 4th April, 1926 and 6 months' half-pay leave from 5th April, 1926 to 4th October, 1926.

M. Kanapathipillai, Inspector, went on 42 days' vacation leave from 24th September, 1926 to 4th November, 1926 and 3 months' commuted full-pay leave from 5th November, 1926 to 4th February, 1927.

R. Kuppusamy, Probationer Dresser, District Hospital, Kuala Lumpur arrived on 20th September, 1926 as relief.

- 93. S. V. Suppiah, Probationer Dresser, Central Mental Hospital, sat for the Preliminary Examination for the Nursing Certificate of Royal Medico-Psychological Association on 20th May, 1926 and passed.
- 94. C. Mylvaganam, Probationer Dresser, and Miss Tan Guat Beng, Probationer Nurse, sat for the Preliminary Examination for the Nursing Certificate of the Royal Medico-Psychological Association on 2nd November, 1926, but the result is not yet known.
- 95. Head Attendant, Kesari Rai who sat for the Final Examination for the Nursing Certificate of the Royal Medico-Psychological Association of Great Britain and Ireland in November, 1925, was informed in February that he had passed. I would like to point out that when Kesari Rai came to Tanjong Rambutan he could not speak English, and that he is a man of fifty.
- 96. A most important appointment was that of a visiting dentist made in March, 1926, when Mr. Hitoshi Simoyama was appointed.
- 97. I look to a great improvement in the health of the patients and an improved recovery rate when the staff really realizes the importance of care of the mouth.
- 98. Attendants.—We suffered from a shortage of attendants both male and female all the year, and we have not been getting a good class, but this has been a trouble for years now. I do not know what has become of the type of attendant we used to get. The attendants dislike the discipline and the night duty, but things improved as the year advanced, and, though we are still short, we are undoubtedly better off than we were at the beginning of the year. Of 165 new attendants engaged during 1926 only 52 remained at the end of the year.

- 99. Maintenance rate.—The maintenance rate is \$176.28 per annum against \$172.96 last year which is due principally to the fact that we purchased nearly all the furniture for the 1st Class Ward out of 1926 year's vote. Had it not been for this we should have shown probably a decrease, but I do not think we can get very much lower unless contractors' prices came down. For the last few years the contractors have been steadily increasing their prices.
- 100. Singapore, Kedah and Kelantan pay the Federated Malay States Government for the treatment of their patients and a total of \$100,702.86 was collected in fees from these Governments, and Federated Malay States paying patients \$1,418.66.
- 101. Amusements.—The usual games were played in the wards—drafts, chess, dominoes, cards and in addition a few Mahjong sets were supplied to the better wards.
- 102. Sunday walks and trips to the town were taken as usual. The few cinemas that visited the town proved a great attraction to the patients, so many going that we now take the whole tent for one night.
- 103. We had to replace Sir Edward Brochman's gramaphone as it had become too decrepit for further use, and the new one now tours the wards.
- 104. The sports were held in July and as usual provided a day's outing for patients and attendants and produced some quite good sport.
- 105. The cricket XI had a much better season, and won twice as many matches as it lost.
- 106. Hockey was started this year and has so far proved very popular, some quite good talent being discovered amongst the patients as well as the staff.

I have the honour to be, Sir, Your obedient servant,

W. F. SAMUELS,

Medical Superintendent,

Central Mental Hospital, Tanjong Rambutan.

TABLE A.

GENERAL TABLE SHOWING THE MOVEMENT OF THE HOSPITAL POPULATION.

DURING THE YEAR 1926.

	Certified patients.						
	М.	F.	Total.	M.	F.	Total.	
On the hospital register, January 1st, 1926 Total cases admitted during the year	1,453 570	443 167	1,896 787	PI STORY		TO S	
Total cases under treatment during the year Cases discharged or transferred during the year—				2,023	610	2,633	
Recovered	162	63	225				
Relieved	60	21	81				
Not improved	62	23	85				
Not insane	4		4				
Died during the year	100	32	132			and or	
Absconded	85	***	85			The state of	
Total cases discharged, transferred and died			1000			1	
during the year			***	473	139	61	
On hospital register on December 31st, 1926				1,550	471	2,02	
Average daily number on the register during			less were				
the year			1,986.62				

Table B.

Showing the Form of Mental Disorder on Admissions in the Direct Admissions During the Year 1926.

		Dire	ct admis	sions.
Fo	ms of mental disorder.	М.	F.	Total.
Congenital of infantile mental deficiency idiocy or imbecility occurring as early in life as it can be observed.	/ Without onlower	6	1 6	1 12
10	1. Insanity with epilepsy	5	5	10
	2. General paralyses of the insane	. 47	3	50
	Insanity with grosser brain lesions . Acute delirium		3	5
- i		. 44	17	6
			1	
		183	33	- 21
	8. Mania—	***		_
ife	A STATE OF THE STA	58	21	7
-	(b) Chronie		2	1
		2	11	1
e e	9. Melancholia— (a) Recent	64	42	10
	(a) Recent (b) Chronic	04	42	10
90	Zal Dannant	6	1	
E	10 Aliconstinutionalis	. 6		
no.	11. Delusional insanity—			
8	(-) Suntamortical	6	****	
it,	(1) N	16	2	1
Insanity occurring later in life	12. Volutional insanity—			
ž l	(a) Impulse	377		
	(b) Obsession			
	(c) Doubt			
	13. Moral insanity			
	14. Dementia—	100		
		40	7	4
		5	***	1
	15. Not insane	1		-
	Total .	489	155	64

PATIENTS DISCHARGED AS CURED DURING THE YEAR 1926.

	Form	of me	ental disc	rder.				Males.	Females.	Total.
1. 1	ntellectual—								STORY OF A	
	(a) With epilepsy								moresimic	
	(b) Without epilep	sy							1000	
2. 1	Moral								- signite d	1000
1. 1	insanity with epileps	y							w metnik	
2. (General paralyses of	the in	isane						toosib say	
3.]	Insanity with grosser	brai	n lesion	8					1-1100	
h. 1	Acute delirium									
5. (Confusional insanity			***				34	8	42
3. 8	Stupor				***			1		1
7. 1	Primary dementia		***					38	4	42
3. 1	Mania-						19		monte	
	(a) Recent			***				33	17	50
	(b) Chronic								2	2
	(c) Recurrent							1	1	2
9. 1	Melancholia—								. bross	
	(a) Recent			***	***			35	22	57
	(b) Chronic	***		***		***			1	1
	(c) Recurrent	***	***					1	1	2
)	Alternating insanity							7		7
	Delusional insanity-						(100		page 1 may	Part Send
	(a) Systematised									10000
	(b) Non-systematis	ed								De la constante
2.	Volutional insanity—									
	(a) Impulse									
	(b) Obsession									
	(c) Doubt								Total bear	
3.]	Moral insanity									
4.	Dementia—									
	(a) Senile						2170		ancaranis.	
	(b) Secondary									
5.]	Not insane		***					1		1

An Analyse of the Discharges and Transfers During the Year 1926.

Laure delaure					M.	F.	Total.	M.	F.	Total.	M.	F.	Total.
Discharges as recov	ered fr	om-											
Direct admission	s												
First attack case	s				87	36	123						
Not first attack of	cases	***	***		30	7	37						
Cases unknown w	hether i	first at	tack o	r not	34	13	47						
Total from direct	admiss	sions			151	56	207						
From transfers—													
First attack cases	8												
Not first attack	cases												
Cases unknown w	hether i	first at	tack o	rnot									
Total from trans	fers												
Total discharges	as reco	vered						151	56				
								R	elieve	ed.	Not	impo	oved.
Discharges (not rec	overed)	as-											
Relieved	***		***	***	56	20	76	56	20	76			
Not improved			***		54	22	76	***			54	22	76
Total	***			***	110	42	152						
Reasons for such di	ischarge	·s											
To go to care of	friends		***		108	42	150						
Statutory by irreg	ularity	in rece	ption	order									
Other reasons (spec	eifying	them)	_								7724	TO YE	
Total				***	108	42	159						
Transferred as—													
Relieved		***	***	***									
Not improved	(555)	***										Draft	
Total	***		***										
Destination of such	transfe	ers-											
To leper asylum		***		***	2		2						
To single care		***											
Other destinations	(specify	ing su	ich)-									Total Control	
Total	***												
Total discharges an	d trans	fers as	_										
TO 11 T								56	20	76			
Relieved		***	****	***	***	***	***	00					

Table Showing the Form of Mental Disorder on 31st December, 1926, of those on the Register at that Date.

Forms of	nental disorder on 31st December.	M.	F.	Total.
Congenital or infantile	1. Intellectual—			
mental deficiency	(a) With epilepsy	5	2	
(idiocy or imbecility) as occurring as early	(b) Without epilepsy	33	18	5
in life as it can be				
observed.	2. Moral			
THE TO	1. Insanity with epilepsy	47	14	6
	2. General paralyses of the insane	52	4	5
Hall Fill	3. Insanity with grosser brain lesions	9	6	1
	4. Acute delirium			
	5. Confusional insanity	62	85	9
1-11-11-13	6. Stupor	5	4	
	7. Primary dementia	380	77	45
	8. Mania—	por las		
14 121 14	(a) Popont	75	35	11
I HE TO WE	(b) Chronic	52	8	6
	(c) Recurrent	15	11	9
life				
.9	9. Melancholia—	1.7000		
ater	(a) Recent	62	28	9
ng 1	(b) Chronic	59	22	8
iri	(c) Recurrent	10	7	1
Insanity occurring later in life	10. Alternating insanity	42	7	4
nit	11. Delusional insanity—			
Inse	(a) Systematised	33	6	3
	(b) Non-systematised	42	4	4
	12. Volutional insanity—	The state of		
	(a) Impulse	3		
1 - 10 - 10 HE	(b) Obsession		1	
	(c) Doubt	1		
The same of	13. Moral insanity			
	14. Dementia—	AND .		
	(a) Senile	75	24	9
THE PARTY	(b) Secondary	485	158	64
and the latest the lat	15. Not insane	3		
	Total	1,550	471	2,02

... Favourable 278
Doubtful 1,223
Unfavourable 520

Prospect of mental recovery

Table Showing the Form of Mental Disorder and Nationality on Admission in the Direct Admissions During the Year 1926.

			plan	Ma	les.					Fem	ales.			Totals.	
Form	s of mental disorder.	Chinese.	Malays.	Tamils.	Javanese.	Eurasians.	Others.	Chinese.	Malnys.	Tamils.	Javanese.	Eurasians.	Others.	Males.	Females.
Congenital or infantile mental deficiency (idiocy or imbecility) occurring as early in life as it can be observed.	1. Intellectual— (a) With epilepsy (b) Without epilepsy	3	1	2				5		1		male made		6	6
-3	Insanity with epilepsy	3		2					1	4		***		5	5
	General paralysis of the insane Insanity with grosser brain	44		1			2	3						47	3
	lesion Acute delirium	***					1	1	1				1		3
	Confusional insanity Stupor Primary dementia	24 87	33	17 59	2	 "i	 "1	8 	1 1 6	8	4			183	17 1 33
	(a) Recent (b) Recurrent (c) Chronic	26 1 	7 1	21	1		3	12 4 	2 4 	7 3 2				58 2 	21 11 2
05	(a) Recent (b) Recurrent (c) Chronic	35 2 2	8 1	19 3 3			2 1	12	2	23 	2		3 1	64 6 6	42
	Alternating insanity Delusional insanity— (a) Systematised (b) Non-systematised Volutional insanity—	1 9	1 2	3 5				2					***	6 16	2
	(a) Impulse (b) Obsession (c) Doubt Moral insanity			STARK STARK	or Town										
	Dementia— (a) Senile (b) Secondary Not insane	30 2	7	3 3 1				3		4				40 5 1	7
	Totals	269	64	142	3	1	10	62	18	64	6		- 5	489	155

Analyses of the Admissions During the Year 1926.

							Acq	uired.							
Classes of admissions.	Co	nge	enital.	Fin	st atta	ick,	Not	first a	ttack.	Unknown whether first attack or not.				Total.	
	M.	F.	Total.	M.	F.	Total.	M.	F.	Total.	M.	F.	Total,	M.	F.	Total
Direct	2	1	3	365	88	453	77	15	92	45	51	96	489	155	644
Total admissions	2	1	3	365	88	453	77	15	92	45	51	96	489	155	644

54	NO.			
	10	00	99	25
and the second second second	489155644		191	318104422
		4	4	- 55
	_			ib mbrech
24		-	1 1	1
	-			-
		-		
54				
The second secon		-		-
E .				
The second secon	9			01
14	- :	4		-
CONTRACTOR STATE				4
Married World Co.	7	2 .		
54	70	4	63	
		00	-	4
The second secon		7		
54	4	10	-	1
		60	00	10
	21	86		17
24	10	10	00	9
Control of the same of the sam	~ ~	őĭ	- 61	18 11
	**	84		18
pi _	320	00	00	1.
	86	4		53 11
	26	25	55	10 00
Pi _	61	10	21	68 44 9
	20	790	- 6	4
Total.	62	9	88	88
pi.	91,19	8	910	6102
N	8	8	2	22
LatoT	100	15,	100	6625 91
64	20	35	70	70
M.	69	8		99
Total	133		88	99 9109
2	85	50	41	91
M.	66	5	22	92
Total	92	10	00	47
9.	155	41	138	12
N.	4		10	85 12
.fatoT	56	81	57	61
54	9	4	6.	~
N.	88	18	51	15
Total	00	00	01	10
0.	*	00	-	01
M.	4	NO.	-	00
	-	01	03	-
54	-	-		
W.			01	-
	99	10	7.	23
	192	195	195	1923
	JaioT M JaioT JaioT JaioT M JaioT N JaioT A JaioT A JaioT M JaioT A A JaioT	M. F. M. M. F. M.	M. F. Touni. M.	1

AGES.

Specific	Drepre	ne wi	nica C	ATIONN 1	Drawn	Dunin	1008			
				ausau I	DEATH	DURIN	, 1020.			
General paralys		ne ins	ane	***		***	***	***	47	
Infacts of splee	n								1	
Phthisis	***				***				16	
Valvular disease	of hear	rt	***	***			***	***	1	
Leprosy					***				1	
Cirrhosis of live		***	***	***	***		***		3	
Bacillary dysen	itery		***	***	***	***			13	
Bright's disease				***					2	
Emphysema	***	***		***			***		1	
Pellagra	****	***		***		***			1	
Exhaustion from	n acute	manie	3				***		3	
Cerebral syphili	s		***	***	***	***	***		3	
Chronic interstit	tial neph	hritis					***		5	
Cardio-vascular	degener	ation				***			6	
Broncho pneum	onia	***	***	***	**	***	***	***	1	
Status epileptic	us					***			8	
Lobar pneumon	ia								1	
Cerebral malari	a			***					1	
Malignant mala	aria		***			***			8	
Emblosim in th		ary ar	tery						1	
Abscess of lung				***	***				1	
Amoebic dysent					***	***			2	
Cerebral softenii									2	
Pneumonia		***			***			***	1	
Exhaustion from				v					1	
Cachexia due to				7					1	
Toxaemia follow									1	
Accidents and a		-							10	
TAGOTACHES GIAG	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,				***					
							Total		132	
							Total		132	
							Total		132	
Description and		D. mrms	two www	nn Anu		Demons			-	
DISTRICTS FROM	which I	Patien	VTS WE	RE ADM	ITTED I	DURING			-	3.
DISTRICTS FROM Sepang	which I	Paties	2	Pu	lau Par		тне Ү		-	3.
Sepang Port Weld			2	Pu Re	lau Par mbau	rit	тне Y		1926	1
Sepang Port Weld Bruas	·		2 1 3	Pu Re Ta	lau Par mbau pah Ro	rit	тне У		1926	1 1 1
Sepang Port Weld Bruas Ipoh			2 1 3 56	Pu Re Ta Pa	lau Par mbau pah Ro pan	rit	тне У		1926	1 1 1 3
Sepang Port Weld Bruas Ipoh Kuala Pilah			2 1 3 56 14	Pu Re Ta Pa Ka	lau Par mbau pah Ro pan mpar	oad	тне У		1926	1 1 3 22
Sepang Port Weld Bruas Ipoh Kuala Pilah Kajang			2 1 3 56 14 6	Pu Re Ta Pa Ka Sit	lau Par mbau pah Ro pan mpar iawan	rit	THE Y		1926	1 1 3 22 11
Sepang Port Weld Bruas Ipoh Kuala Pilah Kajang Kuala Kubu			2 1 3 56 14	Pu Re Ta Pa Ka Sit Ta	lau Par mbau pah Ro pan mpar iawan pah	rit	THE Y		1926	1 1 3 22 11 15
Sepang Port Weld Bruas Ipoh Kuala Pilah Kajang			2 1 8 56 14 6 10	Pu Re Ta Pa Ka Sit Ta Ba	lau Par mbau pah Ro pan mpar iawan	rit	THE Y		1926	1 1 3 22 11
Sepang Port Weld Bruas Ipoh Kuala Pilah Kajang Kuala Kubu Parit Buntar Batu Kurau Kuantan			2 1 3 56 14 6 10 16 1 13	Pu Re Ta Pa Ka Sit Ta Ba Ku Se	lau Par mbau pah Ro pan mpar iawan pah tu Gaj iala Ke	oad	тне Y		1926	1 1 3 22 11 15 13
Sepang Port Weld Bruas Ipoh Kuala Pilah Kajang Kuala Kubu Parit Buntar Batu Kurau Kuantan Pulai			2 1 3 56 14 6 10 16 1 13 2	Pu Re Ta Pa Ka Sit Ta Ba Ku Se Ta	lau Par mbau pah Ro pan mpar iawan pah tu Gaj iala Ka remban iping	oad	тне Y		1926	1 1 3 22 11 15 13 20 30 48
Sepang Port Weld Bruas Ipoh Kuala Pilah Kajang Kuala Kubu Parit Buntar Batu Kurau Kuantan Pulai Klian Intan			2 1 3 56 14 6 10 16 1 13 2 5	Pu Re Ta; Pa; Ka Sit Ta Ba Ku Se; Ta Ra	lau Par mbau pah Ro pan mpar iawan pah tu Gaj ala Ko remban iping ub	rit	тне Y		1926	1 1 1 3 22 11 15 13 20 30 48 7
Sepang Port Weld Bruas Ipoh Kuala Pilah Kuala Kubu Parit Buntar Batu Kurau Kuantan Pulai Klian Intan Pusing			2 1 3 56 14 6 10 16 1 13 2 5 3	Pu Re Ta Pa Ka Sit Ta Ba Ku Se Ta Ra	lau Parmbau pah Ropan mpar iawan pah tu Gai ala Koremban iping ub entakab	rit	тне Y		1926	1 1 1 3 22 11 15 13 20 30 48 7
Sepang Port Weld Bruas Ipoh Kuala Pilah Kajang Kuala Kubu Parit Buntar Batu Kurau Kuantan Pulai Klian Intan Pusing Bentong			2 1 3 56 14 6 10 16 1 13 2 5 3 8	Pu Re Ta Pa Ka Sit Ta Ba Ku Se Ta Ra	lau Paumbau pah Ro pan mpar iawan pah tu Ga ala Kr remban iping ub entakab	rit	тне Y		1926	1 1 1 3 22 11 15 13 20 30 48 7 4 2
Sepang Port Weld Bruas Ipoh Kuala Pilah Kajang Kuala Kubu Parit Buntar Batu Kurau Kuantan Pulai Klian Intan Pusing Bentong Tanjong Rambuta			2 1 3 56 14 6 10 16 1 13 2 5 3 8 5	Pu Re Ta Pa Ka Sit Ta Ba Ku See Ta Ra Me	lau Paumbau pah Ro pan mpar iawan pah tu Ga ala Ki remban iping ub entakab oh nggong	rit pad ah angsar	THE Y		1926	1 1 1 3 22 11 15 13 20 30 48 7 4 2 3
Sepang Port Weld Bruas Ipoh Kuala Pilah Kuala Kubu Parit Buntar Batu Kurau Kuantan Pulai Klian Intan Pusing Bentong Tanjong Rambuta	n		2 1 3 56 14 6 10 16 1 13 2 5 3 8	Pu Re Ta Pa Ka Sit Ta Ba Ku Se Ta Ra Me Kr	lau Paumbau pah Ro pan mpar iawan pah ttu Ga iala Ki remban iping ub entakab oh nggong	rit	THE Y		1926	1 1 1 3 22 11 15 13 20 30 48 7 4 2 3 3
Sepang Port Weld Bruas Ipoh Kuala Pilah Kajang Kuala Kubu Parit Buntar Batu Kurau Kuantan Pulai Klian Intan Pusing Bentong Tanjong Rambuta Lahat Temerloh			2 1 3 56 14 6 10 16 1 1 13 2 5 3 8 5 7	Pu Re Ta Pa Ka Sit Ta Ba Ku Se Ta Ra Me Kr Le Bie	lau Paumbau pah Ro pan mpar iawan pah tu Ga ala Ki remban iping ub entakab oh nggong	rit	THE Y		1926	1 1 1 3 22 11 15 13 20 30 48 7 4 2 3 3 3 3 3
Sepang Port Weld Bruas Ipoh Kuala Pilah Kuala Kubu Parit Buntar Batu Kurau Kuantan Pulai Klian Intan Pusing Bentong Tanjong Rambuta	n		2 1 3 56 14 6 10 16 1 1 13 2 5 3 8 5 7 2	Pu Re Ta Pa Ra Ku Sei Ta Ra Me Kr Le Bi Sei Te Ba	lau Paumbau mbau mpah Ro pah Ro pah mpar iawan pah tu Gaj ala Ki remban iping ub entakab onggong dor lama luk An nting	rit oad ah angsar	THE Y		1926	1 1 1 3 22 11 15 13 20 30 48 7 4 2 3 3
Sepang Port Weld Bruas Ipoh Kuala Pilah Kuala Kubu Parit Buntar Batu Kurau Kuantan Pulai Klian Intan Pusing Bentong Tanjong Rambuta Lahat Temerloh Temerloh Tanjong Toalang Kuala Krau Kuala Kurau	n		2 1 3 56 14 6 10 16 1 1 13 2 5 3 8 5 7 2 4	Pu Re Ta Pa Ra Ku Sei Ta Ra Me Kr Le Bi Sei Te Ba Po	lau Paumbau pah Ro pan mpar iawan pah tu Gaj ala Ki remban iping ub entakab onggong dor lama luk An nting rt Swe	rit pad sah angsar	THE Y		1926	1 1 1 3 22 11 15 13 20 30 48 7 4 2 3 3 3 3 3 3 3
Sepang Port Weld Bruas Ipoh Kuala Pilah Kuala Kubu Parit Buntar Batu Kurau Kuantan Pulai Klian Intan Bentong Bentong Tanjong Rambuta Lahat Temerloh Tanjong Toalang Kuala Krau Kuala Kurau Ulu Selangor	n		2 1 3 56 14 6 10 16 1 13 2 5 3 8 5 7 7 2 4 2 1	Pu Re Ta Pa Ra Ku Sei Ta Ra Me Kr Le Bi Sei Te Ba Po Ch	lau Paumbau pah Ro pah	rit pad sah angsar	THE Y		1926	1 1 1 3 22 11 15 13 20 30 48 7 4 2 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3
Sepang Port Weld Port Weld Bruas Ipoh Kuala Pilah Kajang Kuala Kubu Parit Buntar Batu Kurau Kuantan Pulai Klian Intan Pusing Tanjong Rambuta Lahat Temerloh Tanjong Toalang Kuala Kurau Kuala Kurau Ulu Selangor Rasa	n		2 1 3 56 14 6 10 16 1 13 2 5 3 8 5 7 7 2 4 2 1 1	Pu Re Ta Pa Ra Ku Sei Ta Ra Me Kr Le Bie See Ch Ja	lau Paumbau pah Ro pan mpar iawan pah tu Gaj ala Ki remban iping ub entakab ongong dor luk An nting rt Swe enderis rantut	rit pad sah angsar son ttenhang	THE Y		1926	1 1 1 3 22 11 15 13 20 30 48 7 4 2 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3
Sepang Port Weld Bruas Ipoh Kuala Pilah Kajang Kuala Kubu Parit Buntar Batu Kurau Kuantan Pulai Klian Intan Pusing Bentong Tanjong Rambuta Lahat Temerloh Tanjong Toalang Kuala Kurau Kuala Kurau Ulu Selangor Rasa Batu Tiga	m		2 1 3 56 14 6 10 16 1 13 2 5 3 8 5 7 2 4 2 1 1 1 1 1	Pu Re Ta Pa Ra Ku Sei Ta Ra Me Kr Le Bie Sel Ta Ra Po Ch Jan Tro	lau Paumbau pah Ro pah	rit pad sah angsar son ttenhai	THE Y		1926	1 1 1 3 22 11 15 13 20 30 48 7 4 2 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3
Sepang Port Weld Bruas Ipoh Kuala Pilah Kuala Pilah Kuala Kubu Parit Buntar Batu Kurau Kuantan Pulai Klian Intan Pusing Bentong Tanjong Rambuta Lahat Temerloh Tanjong Toalang Kuala Krau Kuala Kurau Ulu Selangor Rasa Batu Tiga Pudu	n		2 1 3 56 14 6 10 16 1 13 2 5 3 8 5 7 2 4 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Pu Re Ta Pa Ra Sit Ta Ba Ku See Ta Ra Me Kr Le Bie See Ta Poo Ch Jan Tro Ka	lau Paumbau pah Ro pah Ro pah Ro pah Ro pah Ro mpar iawan pah Ro	rit pad sah angsar son ttenhan	THE Y		1926	1 1 1 3 22 11 15 13 20 30 48 7 4 2 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3
Sepang Port Weld Bruas Ipoh Kuala Pilah Kuala Pilah Kuala Kubu Parit Buntar Batu Kurau Kuantan Pulai Klian Intan Pusing Bentong Tanjong Rambuta Lahat Temerloh Tanjong Toalang Kuala Krau Kuala Kurau Ulu Selangor Batu Tiga Pudu Batang Berjuntai	n		2 1 3 56 14 6 10 16 1 13 2 5 3 8 5 7 2 4 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Pu Re Ta Pa Ra Sit Ta Ba Ku See Ta Ra Me Kr Le Bie See Te Ba Poo Ch Jan Tro Ka Ba Ba	lau Paumbau pah Ro pah Ro pah Ro pah Ro pah Ro mpar iawan pah Ro	ah angsar son ttenhang	THE Y		1926	1 1 1 3 22 11 15 13 20 30 48 7 4 2 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3
Sepang Port Weld Bruas Ipoh Kuala Pilah Kajang Kuala Kubu Parit Buntar Batu Kurau Kuantan Pulai Klian Intan Pusing Bentong Tanjong Rambuta Lahat Temerloh Tanjong Toalang Kuala Krau Kuala Kurau Ulu Selangor Rasa Batu Tiga Pudu Batang Berjuntai Mantin	n		2 1 3 56 14 6 10 16 1 13 2 5 3 8 5 7 2 4 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Pu Re Ta Pa Ka Sit Ta Ba Ku See Ta Ra Me Kr Le Bie See Ta Pa Tre Ka Ba Tre Tre Ta	lau Paumbau pah Ro pah Ro pan mpar iawan pah tu Gai ala Ki remban iping ub entakab oh nggong dor lama luk An nting rt Swe enderis rentut onoh ti ti thau moh	ah angsar son ttenhang	THE Y		1926	1 1 1 3 22 11 15 13 20 30 48 7 4 2 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 4 4 4 4
Sepang Port Weld Bruas Ipoh Kuala Pilah Kuala Pilah Kuala Kubu Parit Buntar Batu Kurau Kuantan Pulai Klian Intan Pusing Bentong Tanjong Rambuta Lahat Temerloh Tanjong Toalang Kuala Krau Kuala Kurau Ulu Selangor Batu Tiga Pudu Batang Berjuntai	n		2 1 3 56 14 6 10 16 1 13 2 5 3 8 5 7 2 4 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Pu Re Ta Pa Ka Sit Ta Ba Ku See Ta Ra Me Kr Le Bie See Ta Pa Tre Ka Ba Tre Tre Ta	lau Paumbau pah Ro pah Ro pah Ro pah Ro mpar iawan pah Ro pah Ro mpar iawan pah Ro pah	ah angsar son ttenhang	THE Y		1926	1 1 1 3 22 11 15 13 20 30 48 7 4 2 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3
Sepang Port Weld Bruas Ipoh Kuala Pilah Kajang Kuala Pilah Kajang Kuala Kubu Parit Buntar Batu Kurau Kuantan Pulai Kijan Intan Pusing Bentong Tanjong Rambuta Lahat Temerloh Tanjong Toalang Kuala Krau Kuala Krau Kuala Kurau Ulu Selangor Rasa Batu Tiga Pudu Batang Berjuntai Mantin Padāng Rengas	n		2 1 3 56 14 6 10 16 1 13 2 5 3 8 5 7 2 4 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Pu Re Ta Pa Ka Sit Ta Ba Ku See Ta Ra Me Kr Le Bie See Ch Jan Tru Ka Ba Tee Pe Tre	lau Paumbau pah Ro pah Ro pah Ro pah Ro mpar iawan pah tu Gai ala Ki remban iping ub entakab oh lama lauk An nting rt Swe enderis rantut onoh ti hau hau hau	ah angsar son ttenhang	THE Y		1926	1 1 1 3 22 11 15 13 20 30 48 7 4 2 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 4 4 4 4
Sepang Port Weld Bruas Ipoh Kuala Pilah Kuala Pilah Kuala Kubu Parit Buntar Batu Kurau Kuantan Pulai Klian Intan Pusing Bentong Tanjong Rambuta Lahat Temerloh Tanjong Toalang Kuala Krau Kuala Kurau Ulu Selangor Batu Tiga Pudu Batang Berjuntai Mantin Padang Rengas Sungei Besi	n		2 1 3 56 14 6 10 16 1 13 2 5 3 8 5 7 2 4 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Pu Re Ta Pa Ka Sit Ta Ba Ku See Ta Ra Me Kr Le Bie See Ta Ba Poo Ch Jan Tro Ka Ba Tee Tro Ma	lau Paumbau pah Ro pan mpar iawan pah tu Gai ala Ki remban iping ub entakab oh nggong dor lama luk An nting rt Swe enderis rantut onoh ti ti hau moh kan as	ah angsar son ttenhang	THE Y		1926	1 1 1 3 22 11 15 13 20 30 48 7 4 2 3 3 3 23 3 9 2 1 1 1 4 4 4 4 1

DISTRICTS FROM WHICH PATIENTS WERE ADMITTED DURING THE YEAR 1926—(cont.).

					(00.00.).				
Segamat	****		1	I	antau .		***		
Selim	***		1	I	Labu .			***	
Sungkai	***		3	T	g. Pian	dang	***		444
Setapak			1	G	łrik .			***	
Petaling	***		2	F	Rawang				***
Pondok Tanjo	ng		1	N	falim N	awar			+++
Chemor			4	Г	ambun			***	***
Enggor			1		Batu Ar		111		***
Sungei Siput			6	T	ampin .			***	***
Kuala Lumpu	ir		84		Bagan D			+++	
Bagan Serai			16	F	Kuala Di	pang			***
Klang			32	- 10	Canching	7	***	1000	
Salak North	***		4	-	Sabak B		1		***
Menglembu	***		6	-	epong .		***	***	***
Kuala Lipis			10		iputeh .		***	***	
Gopeng			5		Impang		***		***
Serdang			1		Kampon				***
Kota Bharu	***		4	2	Simpang	Limi	1		***
Terolak			2					70-4-1	-
Tanjong Malin			6					Total	6
Sungei Lembi	mg		2						1
T. (1.1)								2011	
Building during	ng 1926.	_						S	e.
Epsom fa	arm (er	ecting n	ewly)	***			***	1,000	28
Brick kill	n	,,		***	***			167	55
Portrane	farm re							279	84
Leigh far				***	***			366	48
Sungei B								1,286	25
Sungei B								1,081	61
			***	***	444	**	***		
Sungei B				***			***	3,499	
Dust bin,			***	***			***	110	
Dust bin,	, worksh	юр	***	***	***	***	***	45	84
Dust bin			Merry	***			***	138	84
Richmone	d farm	sinking	well			16		190	20
Ward No	. 27, cc	ncreting	bathing	place			***	85	86
Ward No	. 27, fe	nce	***					36	00
Cattleshe	d by wa	ard No.	5. re-lala	ng				199	96
Moulton			***				****	190	32
Woodside								46	
Grange-ge								202	76
Fowl hou				***		***			36
				***	***		***		
Motor ca		441-24-3			***		***	94	35
Horton i					***		***	78	
Horton f						***		111	48
Medical	Superin	tendent'	s quarte	ers ca	ttleshed	erec	eting	2000	-
newly								141	
Fowl hou				recting	newly	***		83	96
Fowl hou	se erect	ting new	y		***			69	36
Boundary	fixing	iron post	8				***	41	40
D. Ward	cement	concret	e by side	e of d	ining ha	III		163	80
Bull shed								55	
Mowing 1									18
Hatton fa								119	
	4"		CAL CREAT		***		***		
",		- 11		***	***		***	238	
Moulton	6"	,b	3	***			***	91	
Moulton					***		***	100	
11	8"						***	195	
Park 2"	sub-soil	drain	22.5		***	***	***	304	40
A DE GIRDER						m		810.040	20
						Total	***	\$10,943	18

FARM PRODUCE 1926.

				0)-0	- COL	ANT SHE		\$	c.
Milk, fresh				m.5		27,515 pints		5,910	72
Pork						12,529.04		4,761	
Fowls		-0.0				24		15	00
Eggs, hen						2,116		84	64
Curry-stuff		***				300		66	00
Onions						62.07	***	6	87
Arrow root			***			42.10		19	79
Millet	***			***		294		28	52
Dholl, Greens	eas, e	te.			***	405	3000	32	40
Paddy to fow	ls	150		·		50		3	00
Indian corn to	fowls					458		36	64
Tapioca		***		***		1,234		24	68
Sweet potatoe	18		***			2,519		75	57
Cocoanute			***		***	5,019		250	95
Yam			***		***	644	+ 1.1	70	84
Oil cocoanut						35		8	75
Oil kachang	***	***	***			25		7	50
Oil cake, coco	anut					76		5	70
Oil cake, kacl						38		2	31
Cigars						4,350		21	75
Limes					***	62	***	1	24
Monkey jack			***	***	***	49		4	90
Otahethi ches	tnut	***	***			166		1	66
Belimbing						59,441		185	88
Pomegranate						8		1	60
Carombolas						1,544		7	72
Bullock's hear	rt	***	***			267		18	35
Custardapples						3,068		153	40
Chikus	***			***		3,857		154	28
Durian beland	8.					452		36	16
Bua sala		***			***	4,835		15	76
Guava		***	***		***	10,216		51	08
Jack fruit						288		144	00
Papayas			***			7,506	***	225	18
Bananas					***	886		1,329	00
Rambutans	***		***			4,235		10	59
Jambu ayer						1,964		9	83
Pineapples		***	***		***	1,192		71	52
Buah susu			***			49		1	47
Grenadillas		***		***	***	8	***		40
Pitanji					***	3,724			73
Oranges						4,558		227	
Pomelos		***	***	***	****	62		12	40
Mulberry	***	***		***		.02 tahils			30
Buah Tanjong			***	***		530			66
Cashew		***	***	***		1,317			59
Cashew nuts		444	***	***	***	1,108	***		39
Water melon	***					54	***		80
Mangoes	***	***		***		541	***		75
Vegetables	***	***			***	289,075.06		21,633	24
									-
						Total		\$35,850	28
Firewood to l	citchen	s, disi	nfecto	rs, dh	oby-				
shed infecti					2000				
etc., etc.						1,551 carts	***	20,163	00
		-		3000	1	glerfe lion		30 200	
						Total		\$56,013	28
						20101		Annioro	-

Dambas Washs						\$ c	8 c.
Brooms, bamboo					5,889 at	30 =	1,766 70
Chinese hats					507 ,,	60 =	304 20
Baskets, carrying vegetal					49 ,,	1 00 =	49.00
Roiling fish					39 ,	90 =	35 10
,, washing rice					49 ,.	1 00 =	49 00
shovels		***			981 ,,	25 =	245 25
,, with cover	***	***			30 ,,	25 =	7 50
., carrying grass	***				61 ,,	50 =	30 50
,, earrying pigs		***	***		2 ,,	1 00 =	2 00
rations					122 ,,	20 =	24 40
medicine (rattan	1)		***	***	1 ,,	1 00 =	1.00
waste paper	***	***	***	***	2 ,,	50 = 10 =	1 00
Sieves bamboo		***	miss m	1550	6 ,,	10 =	
Tailoring.—					0.500	25 =	630.50
Bajus, female patients	***	***	***	***	2,522 2,567	25 =	641 75
Sarongs ,,	***		***	***	8,448 ,,	25 =	2,112 00
Bajus, male patients Trousers ,,					8,382 .,	25 =	2,095 50
Shirts, male attendants					261	50 =	130 00
Shorte	***				269 ,,	50 =	184 50
Kebayahs, female attend					114 ,.	75 =	85 50
Sarongs					115 ,,	75 =	86 25
Mosquito nets					115 ,,	40	
					71 ,,	50 =	81 50
Camp cot covers, canvas					18 ,,	1 00 =	13 00
Canvas cover for bath t	ub	1000	***		1 ,,	2 00 =	2 00
Pillows, coir		***			714 ,,	15 =	107 10
Pillow cases		***	***	***	1,350 ,,	15 =	202 50
Over coats for attendant					2	1 00 =	2 00
Leather slippers in pairs		***	***	***	9 ,,	20 =	1 80
Motor bus eushions cov	ers		***		6 ,,	50 =	3 00
Ovallar cover wooden					81	25 =	21 00
Qualley cover, wooden Sand boxes, wooden		***	***	***	- 00	20 =	6 60
Tyn weepens					103	20 =	20 60
Trays for feeding pigs					35 ,,	50 ==	17 50
Box for minimax					1 ,,	1 00 =	1 00
Crutches pair					1 ,,	3 00 =	3 00
Cocoanut ladles					8 .,	10 =	80
Inventry boards					14	20 =	2 80
Wooden trays	***	***			26 ,,	1 00 =	26 00
Watering cans			***		3 ,,	1 00 =	3 00
Trollies for carrying me			***	***	5 ,,	5 00 =	25 00
Palangs for buffalo's ho		***	***		3 ,	50 =	1 50
Ladders, wooden Inverts moulds	***	***	***		2 ,,	5 00 =	10 00
Yokes for bullockearts		***	***		11 ,,	4 00 = 3 00 =	
Dressing boxes, tins		***	***		0	7 00	
Racks, wooden					2	1 00 =	
Stretcher poles, wooden			***	***	6	1 00 =	
Chest of drawers, wood				***	1 ,,	5 00 =	
Mason trowels, wooden					3 ,,	50 =	1 50
Bathing tins			***		79 ,	18 =	14 22
Mending.—							
Female sarongs	***	***	***	***	3,422 ,,	10 =	342 20
bajus		***	***		2,809 ,,	5 =	140 45
Red blankets			***	***	571 .,		0.000
White blankets	***			***	688 .,	10 =	
Pillow cases Pillows, coir	***	***			884 ,,		10.000
Common lands with			***	***	749 ,,	200	
Chartoine	***		***	***	89 ,, 28	OF	
Attendants sarongs, fen	nales	***	***		140	30	
Female attendants, keb					175		
Red belts					99		
, Male trousers					4,780	-	
Male bajus			***		5,540	-	
Attendants shirts	***			***	92 ,,	10	
shorts		***	***		83 .,		

iscellaneous.—								8	0,		\$
Coffins	***					118	22	2	00	=	236 0
Tombstones		***	****			118	**		25	=	29 5
One table						1		3	00	=	3 0
Farm sign-board				***	****			1	50	-	1.5
Doors to the store						4		1	00	=	4 0
Wooden cages for			pigs			6		2	50	=	15 0
Door posts (farms					2			4	50	=	9.0
Door posts (farms						prs.		6	00	20	12 0
Rack for tea urn						15			00	_	15 0
Photo frames		***	***	***	***				00	=	3 0
Boxes for conveying		···	nn ene	aiman		12	,,	*			6 0
								4		_	4.0
Goal posts	***	***		***		pr.			1201		9 0
Lid to refuse bin	***			***	***	3			320	=	
Wooden boxes for	young	plan	its			2	12	1	00	355	2 (
uly works.—											
Laying out, weeding				nd for	plantin	g 1000	stu	IIS.			
Drain cleaning and											
Sweeping and clea	ning g	rass s	and lal	ang.							
Carting firewood a											
Gardening.											
Repairing and plan	nting.	bamb	oo and	"buns	za rave	" fen	ce.				
Clearing roads and											
Boundary clearing											
		lana									
Scything and rolli	ing pac	rang.									
Woodcutting.						-					
Woodcutting. Clearing overgrowt	ths on	sides				pipe	line				
Woodcutting. Clearing overgrowt Filling and levelling	ths on	sides holes				pipe	line				
Woodcutting. Clearing overgrowt Filling and levellin Repairing roads as	ths on ng up nd patl	sides holes hs.	and sv			pipe	line				
Woodcutting. Clearing overgrowt Filling and levellin Repairing roads at Collecting fuel for	ths on ng up nd patl	sides holes hs.	and sv	wamps.		pipe	line				
Woodcutting. Clearing overgrowt Filling and levellin Repairing roads at Collecting fuel for Making clay pipes	ths on ng up nd path kiln s	sides holes hs. and fa	and sv rms. t inver	vamps.		pipe	line				
Woodcutting. Clearing overgrowt Filling and levellin Repairing roads at Collecting fuel for	ths on ng up nd path kiln s	sides holes hs. and fa	and sv rms. t inver	vamps.		pipe	line	To the same of the			
Woodcutting. Clearing overgrowt Filling and levellin Repairing roads at Collecting fuel for Making clay pipes	ths on ng up nd path kiln s and c	sides holes hs. and fa	and sv rms. t inver	vamps.		pipe	line				
Woodcutting. Clearing overgrowt Filling and levellin Repairing roads at Collecting fuel for Making clay pipes Repairing and tha	ths on ng up nd path kiln a s and c stehing ne.	sides holes hs. and fa emen farm	and sv arms. t inver	ts.							
Woodcutting. Clearing overgrowt Filling and levellin Repairing roads an Collecting fuel for Making clay pipes Repairing and tha Mowing by machin	ths on ng up nd path kiln a s and c atching ne.	sides holes hs. and fa eemen farm	and sv arms. t inver	ts.							
Woodcutting. Clearing overgrowt Filling and levellin Repairing roads an Collecting fuel for Making clay pipes Repairing and tha Mowing by machin Working at Sunge	ths on ng up nd path kiln a s and c atching ne.	sides holes hs. and fa eemen farm	and sv arms. t inver	ts.				es.			8
Woodcutting. Clearing overgrowt Filling and levellin Repairing roads at Collecting fuel for Making clay pipes Repairing and tha Mowing by machin Working at Sunge Ploughing by mote	ths on ng up nd path kiln a s and c atching ne.	sides holes hs. and fa eemen farm	and sv arms. t inver	ts.					c.		8
Woodcutting. Clearing overgrowt Filling and levellin Repairing roads at Collecting fuel for Making clay pipes Repairing and tha Mowing by machin Working at Sunge Ploughing by mot	ths on ng up nd path kiln a s and c stching ne. ei Buls or trac	sides holes hs. and fa eemen farm at, lay ttor.	and sv rms. t inver houses	ts. s.	and su	b-soil	pip	8	c.		
Woodcutting. Clearing overgrowt Filling and levellin Repairing roads at Collecting fuel for Making clay pipes Repairing and tha Mowing by machin Working at Sunge Ploughing by moto pairing.— Bullock-carts	ths on ng up nd pati kiln as and catching ne. ei Buls or trace	sides holes hs. and fa emen farm at, lay	and sv arms, t inver houses	ts. s.	and su	b-soil	pipe	es. \$	c. 00		64 (
Woodcutting. Clearing overgrowt Filling and levellin Repairing roads at Collecting fuel for Making clay pipes Repairing and tha Mowing by machin Working at Sunge Ploughing by mot pairing.— Bullock-carts Handcarts	ths on ng up nd path kiln a s and c stching ne. ei Buls or trac	sides holes hs. and fa eemen farm at, lay ttor.	and sv rms. t inver houses	ts. s.	and su	b-soil	pipo at	es. \$ 4 3	σ. 00 00	=	64 (24 (
Woodcutting. Clearing overgrowt Filling and levellin Repairing roads at Collecting fuel for Making clay pipes Repairing and tha Mowing by machin Working at Sunge Ploughing by mot pairing.— Bullock-carts Handcarts Sledges	ths on ng up nd pati kiln as and catching ne. ei Buls or trace	sides holes hs. and fa emen farm at, lay	and sv arms, t inver houses	ts. s.	and su	16 8 8	pipe at	es. \$ 4 3 5	σ. 00 00 00	=	64 (24 (40 (
Woodcutting. Clearing overgrowt Filling and levellin Repairing roads at Collecting fuel for Making clay pipes Repairing and tha Mowing by machin Working at Sunge Ploughing by mot pairing.— Bullock-carts Handcarts Sledges Meal trolley	ths on ng up nd pati kiln as and catching ne. ei Buls or trace	sides holes hs. and fa emen farm at, lay	and sv arms, t inver houses	ts. s.	and su	16 8 8 57	pipo at	es. \$ 4 3 5 3	σ. 00 00 00 00	=	64 (24 (40 (171 (
Woodcutting. Clearing overgrowt Filling and levellin Repairing roads at Collecting fuel for Making clay pipes Repairing and tha Mowing by mach Working at Sunge Ploughing by mot pairing.— Bullock-carts Handcarts Sledges Meal trolley Meal tray	ths on ng up nd pati kiln as and catching ne. ei Buls or trace	sides holes hs. and farm farm at, laytor.	and svarms, t inver houses	ts. s. verts a	and su	16 8 8 57 137	pipe at	es. \$ 4 3 5 3	c. 00 00 00 00	=	64 (24 (40 (171 (
Woodcutting. Clearing overgrowt Filling and levellin Repairing roads at Collecting fuel for Making clay pipes Repairing and tha Mowing by machin Working at Sunge Ploughing by mot pairing.— Bullock-carts Handcarts Sledges Meal trolley	ths on ng up nd pati kiln a s and c atching ne. ei Buls or trace	sides holes hs. and farm farm at, layttor.	and svarms, t inver houses	ts. s. verts a	and su	16 8 8 57	pipo at 	es. \$ 4 3 5 3 2	σ. 00 00 00 00		64 (24 (40 (171 (274 (
Woodcutting. Clearing overgrowt Filling and levellin Repairing roads an Collecting fuel for Making clay pipes Repairing and tha Mowing by machin Working at Sunge Ploughing by moto pairing.— Bullock-carts Handcarts Sledges Meal trolley Meal tray	ths on ng up nd patl kiln a s and c stching ne. ei Buls or trac	sides holes hs. and farm farm at, layttor.	and svarms, t inver houses	ts. s. verts a	and su	16 8 8 57 137	pipe	es. \$ 4 3 5 3 2 1	σ. 00 00 00 00 00		64 (24 (40 (171 (274 (187 (
Woodcutting. Clearing overgrowt Filling and levelling Repairing roads and Collecting fuel for Making clay pipes Repairing and that Mowing by maching Working at Sunge Ploughing by mote spairing.— Bullock-carts Handcarts Handcarts Sledges Meal trolley Meal tray Benches	ths on ng up nd patl kiln a s and c stching ne. ei Buls or trac	sides holes hs. and farm farm at, layttor.	and svarms, t inver houses	ts. s. verts a	and sul	16 8 8 57 137 187	pipe	es. \$ 4 3 5 3 2 1	6. 00 00 00 00 00 00		64 (24 (40 (171 (274 (187 (163 (
Woodcutting. Clearing overgrowt Filling and levellin Repairing roads an Collecting fuel for Making clay pipes Repairing and tha Mowing by machin Working at Sunge Ploughing by mote pairing.— Bullock-carts Handcarts Sledges Meal trolley Meal tray Benches Bedboards Wooden buckets	ths on ng up nd path kiln a s and c stching ne. ei Bula or trac	sides holes hs. and farm farm at, lay tor.	and sv	ts. s. verts a	and sul	16 8 8 57 137 163 9	pipe at	es. \$ 4 3 5 3 2 1 1	c. 00 00 00 00 00 00 00 80		64 (24 (40 (171 (274 (187 (163 (7)
Woodcutting. Clearing overgrowt Filling and levellin Repairing roads at Collecting fuel for Making clay pipes Repairing and tha Mowing by machin Working at Sunge Ploughing by mote pairing.— Bullock-carts Handcarts Sledges Meal trolley Meal tray Benches Bedboards Wooden buckets Dining tables	ths on ng up nd path kiln a s and c stching ne. ei Bula or trac	sides holes hs. and farm farm at, lay tor.	and sv	ts. s. verts a	and sul	16 8 8 57 137 163 9 8	pipe at	es. \$ 4 3 5 3 2 1 1 3	c. 00 00 00 00 00 00 00 00 00 00		64 (24 (40 (171 (274 (187 (163 (7 5 24 (
Woodcutting. Clearing overgrowt Filling and levellin Repairing roads an Collecting fuel for Making clay pipes Repairing and tha Mowing by machin Working at Sunge Ploughing by moto pairing.— Bullock-carts Handcarts Sledges Meal trolley Meal tray Benches Bedboards Wooden buckets Dining tables Chairs	ths on ng up nd pati kiln a s and c stching ne. ei Buls or trace	sides holes hs. and fa semen farm at, lay tor.	and sv	ts. s. verts a	and sul	16 8 8 57 137 163 9 8	pipe at	es. \$ 4 3 5 3 2 1 1 3 2	c. 00 00 00 00 00 00 00 00 00 00 00 00		64 (24 (40 (171 (274 (187 (163 (7 ; 24 (4 (
Woodcutting. Clearing overgrowt Filling and levellin Repairing roads an Collecting fuel for Making clay pipes Repairing and tha Mowing by machin Working at Sunge Ploughing by mote pairing.— Bullock-carts Handcarts Sledges Meal trolley Meal tray Benches Bedboards Wooden buckets Dining tables Chairs Attendants bed be	ths on ng up nd pati kiln as and c atching ne. ei Bula or trace	sides holes hs. and fa semen farm at, lay tor.	and sv	ts. s. verts a	and sul	16 8 8 57 137 163 9 8 2 5	pipe at	es. \$ 4353211	c. 00 00 00 00 00 00 00 00 00 0		64 (24 (40 (171 (274 (187 (163 (7) 24 (4 (17)
Woodcutting. Clearing overgrowt Filling and levellin Repairing roads an Collecting fuel for Making clay pipes Repairing and tha Mowing by machin Working at Sunge Ploughing by moto Ppairing.— Bullock-carts Handcarts Handcarts Sledges Meal trolley Meal tray Benches Bedboards Wooden buckets Dining tables Chairs Attendants bed be Currystuff pounde	ths on ng up nd path kiln a s and c stehing ne. ei Buls or trac	sides holes hs. and fa semen farm at, lay tor.	and sv	ts. s. verts a	and sul	16 8 8 57 137 163 9 8 2 5	pipe at	es. \$ 4353211	c. 00 00 00 00 00 00 00 00 00 00 56	1111111111	64 (24 (40 (171 (274 (187 (163 (7) 24 (4 (17)
Woodcutting. Clearing overgrowt Filling and levellin Repairing roads at Collecting fuel for Making clay pipes Repairing and tha Mowing by machin Working at Sunge Ploughing by mot pairing.— Bullock-carts Handcarts Handcarts Sledges Meal trolley Meal tray Benches Bedboards Wooden buckets Dining tables Chairs Attendants bed be Currystuff pounde Farm sign-boards	ths on ng up nd pati kiln as and c atching ne. ei Bula or trace	sides holes hs. and fa semen farm at, lay tor.	and sv	ts. s. verts a	and sul	16 8 8 57 137 163 9 8 2 5 1	pipe at 	es. \$ 4 3 5 3 2 1 1 3 2 3 1	6. 00 00 00 00 00 00 00 00 00 00 00 56 00 50		64 (4 24 (4 40 (171 (187 (187 (187 (187 (187 (187 (187
Woodcutting. Clearing overgrowt Filling and levellin Repairing roads at Collecting fuel for Making clay pipes Repairing and tha Mowing by machin Working at Sunge Ploughing by mote spairing.— Bullock-carts Handcarts Handcarts Handcarts Beldges Meal trolley Meal tray Benches Bedboards Wooden buckets Dining tables Chairs Attendants bed be Currystuff pounde Farm sign-boards Screws	ths on ng up nd path kiln a s and c stehing ne. ei Buls or trac	sides holes hs. and fa semen farm at, lay tor.	and sv	ts. s. verts s	and sul	16 8 8 57 137 163 9 8 2 5	pipe at	es. \$ 4 3 5 3 2 1 1 3 2 3 1 1	6. 00 00 00 00 00 00 00 00 00 56 00 50 00		64 (24 (40 (171 (274 (187 (163 (7 2 24 (4 (17 1 1 (1 (1 (1 (1 (1 (1 (1 (1 (1
Woodcutting. Clearing overgrowt Filling and levellin Repairing roads at Collecting fuel for Making clay pipes Repairing and tha Mowing by machin Working at Sunge Ploughing by mote spairing.— Bullock-carts Handcarts Handcarts Handcarts Beldges Meal trolley Meal tray Benches Bedboards Wooden buckets Dining tables Chairs Attendants bed be Currystuff pounde Farm sign-boards	ths on ng up nd path kiln a s and c stehing ne. ei Bula or trac	sides holes hs. and fa semen farm at, lay tor.	and sv	ts. s. verts s	and sul	16 8 8 57 137 163 9 8 2 5 1	pipe at 	es. \$ 4 3 5 3 2 1 1 3 2 3 1 1	6. 00 00 00 00 00 00 00 00 00 56 00 50 00		64 0 24 0 40 0 171 0 274 0 187 0 163 0 7 2 24 0 17 5 1 0 1 0
Woodcutting. Clearing overgrowt Filling and levellin Repairing roads at Collecting fuel for Making clay pipes Repairing and tha Mowing by machin Working at Sunge Ploughing by mote Clay Real trolley Meal tray Benches Bedboards Wooden buckets Dining tables Chairs Attendants bed be Currystuff pounde Farm sign-boards Screws Screws	ths on ng up nd path kiln a s and c stehing ne. ei Bula or trac	sides holes hs. and fa semen farm at, lay tor.	and svarms, t inver houses wing in	ts. s. verts s	and sul	16 8 8 57 137 163 9 8 2 5 1	pipe at	es. \$ 4 3 5 3 2 1 1 3 2 3 1 1 1	6. 00 00 00 00 00 00 00 00 00 56 00 00 00 00 00 00 00 00 00 0		8 64 0 24 0 40 0 171 0 187 0 163 0 4 0 17 4 0 1 0 1 0 1 0 1 0 0 1 0 0 1 0 0 1 0 0 0 0 1 0
Woodcutting. Clearing overgrowt Filling and levellin Repairing roads at Collecting fuel for Making clay pipes Repairing and tha Mowing by machin Working at Sunge Ploughing by mote spairing.— Bullock-carts Handcarts Handcarts Sledges Meal trolley Meal tray Benches Bedboards Wooden buckets Dining tables Chairs Attendants bed be Currystuff pounde Farm sign-boards Screws Ladder	ths on ng up nd patl kiln a s and c stehing ne. ei Bula or trac	sides holes hs. and fa semen farm at, lay tor.	and svarms, t inver houses wing in	ts. s. verts s	and sul	16 8 8 57 137 163 9 8 2 5 1 2 2 4 11	pipe at	es. \$ 4 3 5 3 2 1 1 2 3 1 1 1 2	6. 00 00 00 00 00 00 00 00 56 00 00 00 00 00 00 00 00 00 0		64 0 24 0 40 0 171 0 274 0 187 0 163 0 7 2 24 0 17 5 1 0 1 0 2 0 4 0

REPORT ON THE WORK DONE IN COMBATING VENEREAL DISEASES IN THE FEDERATED MALAY STATES DURING THE YEAR 1926.

During the year under review considerable progress has been made with the campaign against venereal diseases. New treatment centres have been opened which were well attended, there has been an increase in the numbers attending for treatment at the old centres and re-attendance has been well maintained.

The increase made is due to the following:

- (1) The careful selection of staff for each centre. Only those who have been specially trained in this branch of medicine and who take a whole-hearted interest in their work and exhibit a sympathetic attitude towards the patients are appointed.
- (2) The combination of venereal diseases treatment with that for general diseases.
- (3) The propaganda which has been carried out during 1924, 1925 and 1926.

PROPAGANDA.

Lantern Lectures.—I delivered lectures illustrated by lantern slides in different clubs throughout the Federated Malay States. The lectures were on the lines of those given in 1925, the largest attendance was at Taiping and consisted mostly of Malays, this was mainly due to the great interest taken by the District Officer, Larut. As a result of this lecture one of the leading Malays of the State of Perak, at his own expense sent several Malays to Kuala Lumpur for treatment, and he expressed a desire to assist in propaganda as in some of the Malay kampongs venereal diseases are very prevalent, for which purpose he has been supplied with a copy of my lecture, a set of slides and a lantern.

At the Agricultural Show held in Taiping during the 14th and 15th of August a 100m was set aside for the purpose of delivering a series of lantern lectures. The number attending these lectures approximately 960. A collection of photographs, mostly taken from life showing the dangers of venereal diseases were exhibited with explanatory text in different languages. Pamphlets in different languages were distributed.

- 2. At the Agricultural Show held in Kuala Lumpur during the month of August a portion of the space allotted to the health section was partitioned off for venereal diseases propaganda and a dresser from the venereal diseases staff was in constant attendance. The photographs mentioned in the preceding paragraph were exhibited and pamphlets distributed. This portion of the health section was very well attended by all nationalities.
- 3. Posters and Pamphlets.—A new series of posters have been printed in English, Chinese, Malay, Tamil and Urdu; these are exhibited in public places. A pamphlet has been prepared explaining briefly and concisely the dangers of contracting venereal diseases; the early signs and symptoms; and the importance of early and continued treatment. In addition to the above methods opportunity is taken in the Hospitals, Town Dispensaries and Clinics to educate the public—those attending these institutions are made to understand the dangers of venereal diseases and the importance of early and continued treatment.
- 4. As in 1925 the professional education of Assistant Surgeons and dressers has been carried on. Practically the whole of the subordinate staff of Perak, Selangor and Negri Sembilan received a course of instruction in venereal diseases and their treatment. It was not found possible to give this course to the staff in Pahang but this is to a certain extent was remedied by the distribution of my lectures.

PROPAGANDA LECTURES.

SELANGOR.

KUALA LUMPUR.

Date.		To whom lectured and place.	At	tendano	e.	Time.
22-2-26	1.0	Railway Institute		70		7.00 p.m.
22-4-26				40	***	7.00
1-3-26	***	Juvenile Athletic Association		60	***	7.00 ,,
3-3-26				90		7.00 ,.
5-10-26		The second secon		55	***	6.45 ,,
		Railway Institute		60		6.30 ,,
		Confuscian School		75		6.30 ,,

SHEAS	BIG MANAGEMEN DEPARTE DIE	OW BUT NO THOMAS	
120 35	I STARTY STITT OFFICE OF TAPAH. VALUE OF	Attendance. Time.	
16-6-20	Police and Malays (in one of the hospital	1	
17-6-26	wards)		
17-0-20	Chinese (in one of the hospital wards)	. 120 7.00 ,,	
	ІРОН,		1
30-6-26	The Perak Amateur Dramatic Association	115 6.30	
	TAIPING.		
18-7-26		also labour estil (1)	
	Taiping (at the Town Hall)		
14-7-26			
	Hall)	100 0.00	
14-8-26		000	
and			
15-8-26	(10 1001000)		
	Negri Sembilan.		
01 4 00	KUALA PILAH.		
21-4-26	and the same of the same same same same same same same sam		
	school)	110 7.00 ,,	
	TAMPIN.		
27-4-26	Government servants and others	50 7.00	
28-4-26		SHEET OF THE PARTY AS THE	
	Government English school)	60 7.00 ,,	
	SEREMEAN.		
1-10-26			
	the native community (at the Miners'		
	Association, Seremban)	50 7.00	
	the to reproduce &	number altending these	
-	STAFF.	of our men neglet officers	
	re has been no increase to the staff during the year.		
and wel	trained Assistant Surgeons are necessary for each		
	Specialist, Venereal Diseases	Dr. E. A. Smith	
	Assistant Surgeon	Mr. G. H. Oorloff	
	Nurse	Mrs. V. M. de Lile	
	Dresser, Grade II	Mr. V. Nadayson	
	Dresser, Grade II	Mr. A. Sinnappu	
	Dresser, Grade II	Mr. D. Arunasalam	
	Dispensary Dresser, Grade II	Mr. Teh Tai Seng	
	Dresser, Grade III	Mr. S. Ponnusamy	
	Dresser, Probationer	Mr. T. Suppiah	
	Laboratory Dresser, Probationer	Mr. Low Pak Foon	
	Attendant	Retnam	
	Attendant	Gopal	
	Attendant	Ah Loke	
	Attendant (famale)	Toong Cai	

Attendant (female) Leong Sai

I should like to mention that the whole staff has worked extremely well, and to specially commend dresser Sinnappu's work in Klang as it was largely due to his efforts that the Klang Treatment Centre has been such a success. Dresser Arunasalam is also to be commended for the neat work he has done in printing the graphs attached to this report.

TREATMENT CENTRES.

In my annual report for 1925 six main treatment centres were suggested, four of these are now equipped and working, viz.—Kuala Lumpur, Seremban, Klang and Taiping.

2 A main centre was opened in Klang during the month of July. It will be seen from the accompanying graph No. 1 that this centre was necessary; it has quickly established itself and become popular.

The Acting Senior Medical Officer, Selangor, has shown an extremely sympathetic attitude towards the venereal diseases campaign in his State. He has rendered every help possible, especially as regards staff and equipment. Specially selected dressers are sent to the venereal diseases wards, District Hospital, Kuala Lumpur, and to the Sultan Street Clinic for training in venereal diseases; when efficient they are drafted to other main or sub-centres.

- 3. Alterations to the Town Dispensary, Seremban, were completed during the year, converting the Infant Welfare Centre into a Venereal Diseases Clinic (the Infant Welfare Centre moving into the adjoining house.) This centre was opened in October.
- 4. The Taiping Centre was opened during the early part of the year. Work here unfortunately has been of a spasmodic nature owing to shortage of staff, the Malay Assistant Surgeon having other duties to perform is not able to devote sufficient time to this centre. This is regrettable as there are many Malays in the neighbourhood who would come for treatment.

The alterations necessary to convert the old Infant Welfare Centre, Ipoh, into a Venereal Disease Clinic were completed in the middle of December.

Provided there is available staff, clinics will be shortly opened in Teluk Anson and Kampar.

The Senior Medical Officers have rendered all assistance possible over the important factors of staff and equipment.

KUALA LUMPUR.

There are two main treatment centres in Kuala Lumpur:

- (A) attached to the Government Town Dispensary, Sultan Street.
- (B) in the District Hospital for indoor and outdoor patients.

(A)—GOVERNMENT TOWN DISPENSARY AND VENEREAL DISEASE TREATMENT CENTRE, SULTAN STREET.

The two shop-houses in Sultan Street which have been rented since 1924 are still being used as the Town Dispensary and Venereal Diseases Treatment Centre.

During the year, 3,222 patients of both sexes, suffering from venereal diseases, received treatment, showing an increase as compared with 2,328 in 1925 and 35,289 received treatment for other diseases in the Town Dispensary in 1926 as compared with 29,335 in 1925.

2. Following table shows nationalities and diseases:

Nationality		Total.	Syphilis.		Per cent.	G	onorrhoe	a.	Per cent.	amb	Soft sor	e. 1	er cent.
Chinese		2,069	 1,428	***	88.77		505	****	24.40		141	***	6.91
Tamils		531	 244		45.95		246	1	46.32		41		7.72
Malays		171	 108		63.15		59		34.51		4		2.38
Sikhs		183	 . 116		63.33		58		31.69	B	9		4.91
Eurasians		48	 16		33.33		26		54.16	1	6		12.5
European	8	152	 26		17.1		113		74.84	10.	13	We	8.55
Others		68	 29		42.64								
Total	***	3,222	 1,962		60.89	de	1,036	1000	32.15		224		6.95

It will be seen from this table that the Chinese are the heaviest sufferers from syphilis and Europeans from gonorrhoea; it was the same in 1925.

3. Eighty-seven patients attended for prophylactic treatment during the year.

4. Table showing admissions during 1925 and 1926:

							1925.		1926.
Chinese		make m	777	1000	1,000	980	1,496		2,069
Tamils		olle ne			***		398	000	531
Malays		100				***	96		171
Sikhs	***	111					172	1.00	183
Eurasian	8	*******					33	100000	48
European	ns	0.0000		+441			84	- 16.	152
Others		mes.od	1		o samul		49		68
					Total		2,328		3,222
							1 (1)		

=increase of 894.

During 1926, there has been an increase of all nationalities, Malays and Europeans being specially noticeable.

5. It will be seen from graph No. 2 (showing admissions and diseases) that the largest number attended during the month of October; this was also the case in 1924 and 1925; I am unable to account for this.

Out of the 3,222 patients treated 627 were females.

RECORD OF WORK IN MALE SECTION

						 9,883
Prostatic massage	e			***		 385
Sounds			***			 484
Urethroscopic ex	caminations			***		 95
Arsenical N.A.B	. injections	***				 829
Stabilarsan injec	tions					 3,102
Bismostab inject	ions	***	***		2220	 3,541
Sulfarsenol						 29
Vaccine	M. 1.				1944	 12
Wassermanns						 353
Examination of	microscopie	slides				 3,308
	The state of the s					

FEMALE SECTION, SULTAN STREET CENTRE.

During the year 1926 six hundred and twenty-seven patients received treatment as compared with three hundred and thirty-five in 1925. The following table shows nationalities admitted during 1925 and 1926.

		Nations	lity.			1925.		1926.
Chinese		 	444	***	***	 254		475
Tamils		 				 37		64
Malays		 			***	 17		51
Sikhs	***	 	***	***		 12	***	15
Eurasia	ns	 				 3		12
Others		 	***	***		 13		10
					Total	 335	***	627
						=incre	ase of	292.

It is encouraging to note the increased attendance of Malays.

2.	RECORD OF	Work	DONE	IN FE	MALE	SECTION.		
	Douches					***		1,652
	Arsenical injections					***		689
	Bismostab injections							689
	Wassermann blood te	st					***	42
	Microscopie examinati						***	1,529

- 3. Eighteen ante-natal cases received treatment; six of these were sent by the Infant Welfare Centre, the remaining twelve came of their own accord.
- 4. During 1924, 1925 and 1926, endeavours were made to persuade the prostitutes to come for examination and treatment. A lantern lecture was given to the brothel keepers; the dangers of venereal diseases and the importance of early and continued treatment were fully explained. The keepers then visited the clinic for the purpose of discussing whether or not they would consent to their girls coming. Their wishes as regards privacy and who should examine them, etc., was met in every way. The whole of the upstairs of one of the shop-houses has entirely been allotted to them. It was possible to arrange for an entrance and exit into a back lane thus obviating their passing through the dispensary. Eventually one brothel agreed to try the experiment which was successful and when the girls saw what was being done for them and the advantages of attending were explained, no further opposition was raised. In September all the Chinese prostitutes in Selangor came for examination, and treatment when necessary; this continued to the end of the year; number of prostitutes attending the clinic since the month of September=1,406.
- Mrs. V. M. de Lile, the nurse, who speaks many Chinese dialects fluently, and the Chinese amah deserve every credit for the tact, consideration and kindness shown to the girls. It is largely due to their efforts that the prostitutes come willingly.

All the girls now realise the danger of contracting venereal diseases and the importance of early and continued treatment, in fact they now come for the most trifling ailments.

The nurse gives the girls instruction in general hygiene as well as in venereal diseases.

(B)—DISTRICT HOSPITAL, KUALA LUMPUR. VENEREAL DISEASES TREATMENT CENTRE.

During the period under review 1,414 patients of both sex indoor and outdoor received treatment.

The following table shows nationalities and diseases:

Nati	ionality.		Total.		Syphilis	8.	Per cent.	Go	norrho	ea.	Per cent.	. Se	oft sore.	Pe	r cent.
Chinese			447		305		68.2		98		22.0	***	44		9.8
Tamils			788						249				91		12.4
Malays			137	,	83		61.0		48				6		4.0
Sikhs	See.		30		10						60.0		2		7.0
Eurasia	ns		12		4		33.0		7		59.0		1		8.0
Europe	ans	***	3		1		33.0		2		67.0		-		_
Others	***		52		28		53.8		22		42.2		2		4.0
	Total		1,414		824		58.2		444	***	31.4		146		10.2

It will be seen from this table that the Chinese are the heaviest sufferers from syphilis; it was the same in 1925. Out of the total number treated 252 were females and 1,162 were males.

Indoor Outdoor	 	 	 	Males. 936 226	 120 132
				1,162	 252

Following table shows numbers receiving treatment during 1925 and 1926.

				19	25.	1926.
				1,	281	 1,414
	REC	ORD OF V	VORE			
Irrigations		200		***		 7,653
Prostatic massage				***		 346
Sounds						 15
Urethroscopic examinat	ions			****		 3
Arsenical injections (No		enobillon	Stak	oilarsan)	 3,065
Sodium thiosulphate						 69
Bismostab injections		***				 1,780
Sulfarsenol						 96
Buboes aspirated						 68
Wassermanns						 1,160
Examination of microso	opic	slides				 111
Lumbor puncture						 9
Sulphostob						 24
Contramine						 92

SEREMBAN TREATMENT CENTRE.

During the year, 782 patients of both sexes received treatment. This centre was transferred from the General Hospital to the Town Dispensary in October. It will be seen from graph No. 3 the admissions from that time onwards have greatly increased.

The following table shows the nationalities and diseases:

Nationality.	Total.	Syphilis	Per cent.	G	conorrh	oea.	Per cent.	S	oft sor	re.	Per cent.
Chinese	 421	 292	 69.35		90	***	21.37		39		9.26
Tamils	 183	 95	 51.91		69		37.7		19		10.3
Malays	 60	 31	 51.66		26		43.33	***	3		5.00
Sikhs	 85	 20	 57.14	***	10		28.57		5		14.26
Eurasians	 17	 9	 52.94		6		35.29		2		11.76
Europeans	 21	 5	 23.8	***	15		71.42	***	1		4.76
Others	 45	 28	 62.22		14		31.11		3		
Total	 782	 480	 61.38		230		29.41		72		9.2
	-	-									

	Irrigation	0.0				ORD OF						0.000	
	Prostatio					***				***		2,028	
	Urethral			***	***			***		***		379 66	
	Stabilars							***	***	***	***	1,063	
	Mercury					***			***	***	***	366	
	Bismosta							**				308	
	Sulfarser				***					***	***	195	
	Buboes					***				***	***	93	
	Wasserm		teu		***			**		***	***	299	
	Examina		f mic		mia e	lidoo			***	***		464	
	Jaamina	· ·	1 mic	SLOBOL	pro s	sinces						404	
						STAF	F.						and del
	Assistant											115 C. C. C. C. C.	etnam
	Dresser,	Grade	III							Mr. I	Kandi	ah	
	m m.					EATM							
The	Town Di	spens	ary,	Klang	g, ha	s been	alte	red	to al	low pat	ients	Sufferi	ng fron
that tim	ne to the	end e	of the	e ves	r 46	8 pati	ents	(of	both	sexes)	receiv	ed tro	eatment
Graph N	o. 1 show	s adm	ission	is and	dise	eases.	It w	ill be	e seer	n from t	he gr	aph th	at ther
	a steady												
The	following	table	show	vs na	tiona	lities a	and d	liseas	ses:				
	nality.	Total.		philis		Per cent				Per cent.	Soft	t sore.	Per cent
Chinese	95. hm	169		113		67.04		38		22.48		18	. 10.6
Tamils		286		35		36.01		110		46.61		41	. 17.3
Malays		24		11		45.83		9		37.5		4	. 15.16
Sikhs		21		7		33.33		10		47.61		4	. 19.0
Eurasian	ıs	3	***	1		33.33		1		33.33		1	. 33.33
European	ns	_		_		_		113		_			. —
Others		10		2		20.00		8		80.00			
				010		47.0			-	00.0	-	-	14.00
	Fotal	463		219		47.3		176		38.6		68	. 14.68
					RECO	RD OF	Wor	K.				0.0	
	Irrigations				***					***	***	916	
	Prostatic					•••			***	***	***	75	
	Urethral				***	***				***	***	25	
	Urethrose									***		5	
	Novarseno				8				***	***	***	202	
	Stabilarsa				***	***			***			293	
	Bismostal								***	***	***	380	
	Sulphosta		ection	8			-					26	
	Wasserms			***			1 100			***		6	
	Examinat	ion of	mic	rosco	pic s	lides	-		***	***	***	194	
						STAFF	=						
	Assistant	Surge	eon .			STAFF				M	r. V.	S. B	hattal
	Dresser,											Sinna	
								***	4100				9 Signific
A re suffering	ecord has	neen	oea	admi	tted	into	the	vene	ereal	disease	s W	ards,	District

A record has been kept during the last four months of the year of those cases suffering from gonorrhoea admitted into the venereal diseases wards, District Hospital, Kuala Lumpur, and those who were treated at the Sultan Street Clinic, showing numbers who came for treatment in the acute and chronic stages of the disease.

				Hospital.		Street.
Acute	 	 	***	 58		117
Chronie	 ***	 ***	***	 22	***	126

Table showing number of indoor and outdoor male and female patients who attended the District Hospital Clinic for arsenical injections during the year 1926.

					Number of injections.										
	Nat	ionality			1.	2.	3.	4.	5.	6.	7.	8.	9.	10.	Total.
Chinese					30	41	29	23	26	34	20	14	7	25	249
Tamils					53	46	87	58	38	39	21	15	9	20	329
Malays					6	14	4	9	11	7	7	6	1	2	67
Sikhs					2	2	2	3	3	1	1				14
Europeans	3	***							1						1
Eurasians			***			1				12.2				1	1
Others	***		***		2		3	3	3	1	1	1		2	16
			Total	***	98	103	75	91	82	82	50	34	17	50	677

It will be seen from this table that out of the total of 677 patients who received treatment 50 attended a full course.

Table showing numbers and nationalities who received arsenic injections in the Male section of the Sultan Street Venereal Diseases Treatment Centre, Kuala Lumpur, during the year 1926.

								Num	ber of	injec	ions.				
	Na	tionality.	index!	Jedges		2.	3.	4.	1. 5.	6.	7.	8.	9.	10.	Total.
Chinese					195	165	89	66	36	56	20	40	18	56	741
Tamils				411	46	27	4	8	8	4	1	3	8	7	111
Malays		***			38	16	6	3	4	2	1	1	2	1	74
Sikhs		***			16	13	111	4	3	6	4	3	2	3	65
Europeans					2	1		1	1	1		1	2		9
Eurasians					4	2	1		1					1	8
Others					2	3	2	1	2		1			1	12
			Total		303	227	112	88	58	69	27	48	27	69	1,020

It will be seen from this table that out of the 1,020 patients 69 attended a full course.

Table for the female section of the Sultan Street Venereal Diseases Treatment Centre.

							Number of injections.					25000			
	Nat	tionality	Smoot L	100	1.	2.	3.	4.	5.	6.	7.	8.	9.	10,	Total.
Chinese				***	44	39	17	16	8	9	2	4	9	14	162
Tamils	***	***			3	7		5	3	2		***	1		21
Malays				***	10	9	8	2	3	4		1	1	2	40
Sikhs					3	2	5							1	9
Eurasians		***	***		***					1			***	1	2
Others	***				1		1		***	1		1		1	5
			Total		61	57	29	23.	14	17	2	6	11	19	239

It will be seen from this table that out of the total of 239 patients 19 attended a full course.

From the above it is seen that the Chinese attendance is better than other nationalities.

FEDERAL HOME FOR CHINESE WOMEN AND CHILDREN, KUALA LUMPUR.

I paid 41 visits during the year to the above Home. A small clinic equipped with the necessary instruments and drugs was opened in March. The head matron of the Home attends to all cases who require treatment. There were 73 admissions to the clinic; of these, 32 were found to be suffering from syphilis and the remaining 41 from chronic cervicitis, gonococci were found to be present in 17 out of the 41.

Sixty-eight bloods were sent for the Wassermann test; thirty-two gave a positive reaction.

One thousand and fifty-four smears from the urethra and cervix were sent to the Sultan Street Clinic for examination.

ARSENICAL PREPARATIONS.

Stabilarsan has been used to a large extent during the year. I prefer this to other preparations being in solution it is easily administered.

- 2. Bismostab has taken the place of mercurial cream, for the excellent reason that it causes practically no pain.
- 3. A supply of sulphostab was received from Boots Pure Drugs Co., for the purpose of giving this preparation a trial. Five cases are quoted below.
 - 4. Contramine has been used with good results in treating gonorrhoeal arthritis.
- 5. Sulfarsenol.—Good results were obtained with this preparation last year in treating cases of epididymitis. I have had the same satisfactory results this year.
- Hexyl Resorcinol.—This has been found to give good results in all affections
 of the posterior urethra; both in the acute and chronic condition the urine clears
 rapidly.

SULPHOSTAB "BOOTS"-CASES TREATED WITH.

(1) Female Chinese, aged 30, suffering from cerebral syphilis: On admission her blood was positive to the Wassermann test. She presented the following symptoms: Some time before admission into hospital she suffered from headaches. On admission it was not possible to obtain many details as patient was not able to speak. She had facial paralysis right side, gait spastic, walking with difficulty. Knee jerks + +, Rhombergs sign present. Eyes: left eye absent; right eye serpiginius ulcer of cornea.

The following treatment was prescribed-

Sulphostab .-

0.3 gm. twice a week for five injections =1.5 gms. 0.45 gm. once a week for five injections =2.25 ,,

Bismostab .-

1 c.c. once a week for eight injections = .8 gms.

Potassium Iodide.—Starting with 10 grains three times a day; this was increased every third day until a maximum of 30 grains three times a day was reached, then gradual reduction of dose to 10 grains three times a day.

Her condition at the end of this course showed some improvement—being able to walk much better; gait no longer spastic, general health better, gradually increasing in weight, though she is unable to speak and the facial paralysis being still present.

Blood Examination.—Wassermann test done six weeks after last injection of sulphostab was found to be negative.

This is a difficult case to expect any marked improvement as it is most probable the affected part of the brain is damaged to such an extent that recovery of the part is impossible but her blood becoming negative six weeks after termination of one course is distinctly encouraging. She will shortly be placed on a further course.

- (2) Male Chinese, aged 40 years, admitted on 20th August, 1926, suffering from acute pain and swelling of right knee joint; he was unable to walk; a urethral discharge was present in which gonococci were found. Patient stated that this was his first attack. He received eleven injections of sulphostab .3 gm. no very marked improvement resulted but the acute condition subsided; patient is able to walk with the aid of a stick, there being a certain amount of ankylosis on the knee joints due to patient refusing to move his joint in the earlier stages of the disease but this will, I think, improve later on.
- (3) Male Malay, aged 36, admitted on 28th September, 1926, suffering from acute pain and considerable swelling of right foot; the tarsal and metatarsal joints affected; patient was unable to walk; urethral discharge was present containing genococci. He received eight injections of sulphostab 0.3 gm. The pain has completely disappeared; swelling gradually decreasing and he is able to walk. Patient was discharged on 18th December, 1926.

- (4) Male Chinese, aged 28, admitted on 16th October, 1926, suffering from pain and swelling of both knee joints, also right ankle and right metartarso phalangeal joints. Condition on admission very acute—patient unable to walk; a urethral discharge present containing gonococci. He received five injections of sulphostab of 0.3 gm., three days after the first injection a marked improvement was noticed, pain much less; patient able to walk a little; this improvement continued. Patient left hospital on 10th November, 1926, quite cured. Stay in hospital=26 days.
- (5) Male Tamil, aged 30, admitted 29th October, 1926, suffering from acute pain and swelling of both wrist joints; a urethral discharge was present which contained gonococci. Patient's mouth was in a very bad condition—pyorrhoea being present. He received five injections of sulphostab 0.3 gm. Six days after first injection a marked improvement was noticed; pain and swelling much less, this improvement continued. Patient leaving hospital quite cured on 19th November, 1926. Stay in hospital=21 days.

All these four cases of gonorrhoeal rheumatism were in the chronic stage of gonorrhoea and received in addition to the sulphostab injections, a course of prostatic and seminal vesicular massage (gonococci found to be present in smears taken from these glands) also they were made, as far as possible, to perform active movements of the inflamed joints.

In the first two cases no marked improvement occurred though progress towards recovery is being made. The last two can be considered to have done well. All the injections were given at an interval of four days.

The above cases were treated in the venereal diseases wards, District Hospital, Kuala Lumpur.

In the venereal diseases wards, District Hospital, Kuala Lumpur, the following record was kept during the second half of the year showing number of cases who suffered from a rise of temperature after intravenous injections of novarsenobillon and stabilarsan.

			Novarsenobillon.		Stabilarsan.
Number of injections		***	163		164
Temperature—numbers			102		98
Time after injection		***	11 to 2 hours		1 hour
Highest temperature recorded	***		101 F.	***	103 F.

Tamils suffered more than the other nationalities.

BURGES.

One hundred and fifty-three cases were treated during the year. Of these, 85 were treated by aspiration and injections of mercurochrome 2 per cent. 68 subsided under treatment with glycerine and ichthyol.

EPIDIDYMITIS.

Twenty-two cases were admitted during the year. Of these, 16 were treated with sulfarsenol with the usual good results. Six were treated with contramine and London Lock Hospital vaccine; the results were not so satisfactory as with sulfarsenol.

GRANULOMA INGUINALAE.

Five cases received treatment during the year. One gave a positive blood Wassermann. They consisted of the following nationalities:

Sikh	 	 	 	 -	 1
Chinese	 	 	 	 	 2
Tamils	 	 	 	 	 2

All received tartar emetic intravenously, 2 per cent. mercurochrome applied locally, and heliotherapy.

The Sikh made a perfect recovery, the remaining four very slight progress. One of the Tamils received X'Ray treatment; he showed no improvement. One of the Chinese had the growing edge of the granuloma burnt by fulguration; he received one application—this was followed by a marked improvement. I propose giving this treatment a further trial.

GONOCOCCAL ARTHRITIS.

Thirty-two cases received treatment during the year. For the first six months various drugs were tried; of these contramine gave the most satisfactory result. For the last six months a careful record of all cases has been kept. Of these, 14 received a course of contramine and 4 sulphostab. See chart. No. 4.

Gonococci were found in all cases, either in the meatal discharge or in the material from the prostate and seminal vesicles. In addition to the contramine injections patients were given a course of prostatic and seminal vesicular massage; they were also made to perform active movements of the inflamed joints.

NOMA PUDENDA (SPIROCHAETAL GANGRENE).

Two cases received treatment-

Case 1.—Occurred in a young Tamil woman who came into hospital with the whole of the left labium major and minor in a gangrenous condition extending up to the vaginal wall on the same side to some considerable extent.

Duration eight days.

On admission her condition was one of general toxaemia.

Treatment started at once.—All the gangrenous sloughs cut away until healthy tissue was reached; wound then dressed every two hours with hydrogen peroxide and bismuth formic iodide for two days; the dressing was then changed to a 2 per cent. mercurochrome three times a day for twelve days, after which it was done once daily. Patient ultimately recovered; there was a considerable loss of tissue; the whole of the left side of the vulva being completely destroyed exposing the lower portion of vagina.

Laboratory report.—Smears showed staphylococci, streptococci but the main feature was the presence of a large number of spirochaetes with loose open spirals, and fusiform bacilli.

Case 2.—Occurred in a female Chinese, aged 35 years, with the whole vulva in a gangrenous condition; also mouth, lips and left side of face affected, the latter to such an extent that left cheek had sloughed away. Her condition was one of extreme emaciation and toxaemia.

Treatment started at once.—Hourly applications of hydrogen peroxide and dusting with bismuth formic iodide; this proved of no avail; her general condition being very bad, death occurred shortly after admission.

· HAEMORRHAGIC ENCEPHALITIS.

Male Chinese, aged 29; occupation: motor salesman. Admitted into General Hospital, Kuala Lumpur, on 1st November, 1926; died at 1.30 a.m. on 10th November, 1926.

The following facts were obtained from his wife who stated he never had any illness during the time she had known him. On 30th October, 1926, at about 12 noon he is said to have had an injection of novarsenobillon; the same afternoon he felt giddy and feverish, at 5 a.m. on 31st October, 1926, he fell down unconscious.

Condition on admission.—Patient, a man of very good physique, admitted in an unconscious state; no evidence of external injury to head or elsewhere; heart, lungs, spleen, liver—nil of note, pulse: good tension and volume, knee jerks: absent, kernigs sign: not present, pupils both dilated, equal and react sluggishly to light, no paralysis, no rigidity of neck muscles, irritable, keeps rolling about the bed, resents very much being touched, twitching of muscles on left side of face, incontinence of faces and urine, temperature 99° F. Following is report on funduscopic examination.—"Both eves discs haemorrhagic to a slight extent, showing punctiform haemorrhages all over; the outlines of the discs are not lost but are slightly ill-defined; retinae also slightly pale. Blood vessels are clear and well defined but appear less full. The entire picture is suggestive of either diffuse inflammation of brain tissue or increased intra cranial pressure but more likely the latter".

Patient's condition remained unchanged, very restless, continually trying to get out of bed resenting any restraint; great difficulty in getting him to take any fluids.

On the day after admission patient's temperature became subnormal—98° F. and remained so until day prior to death when it again rose to 100.5°F.; the day he died it gradually fell; becoming subnormal just before death; which occurred at 1.30 a.m. on 10th November, 1926.

A lumbarpuncture was done the day after admission; 20 c.c. cerebro spinal fluid withdrawn under pressure; this was sent to the Government Laboratories for a report, which reads as follows: "sediment chiefly red blood cells; no organism seen". The following treatment was prescribed: venesection 20 ounces of blood; lumbarpuncture 20 c.c. cerebro spinal fluid withdrawn, and repeated injections of adrenalin.

A full post-mortem was not allowed by the relatives; consent was given to remove the brain only; which presented the following appearance: "The surface veins were intensely congested and the brain substance was softer than usual. There were extensive areas of haemorrhagic softening in both occipital lobes, left parietal and temporal lobes".

Following is the histological report received from the Government Laboratories: "section of brain shows following lesions microscopically. Numerous small haemorrhages into the brain substance. (2) Hyaline thrombosis both completely and partially obstructing various large vessels and denoting damage of the vessel endothelium. (3) Large areas of haemorrhagic softening and necrosis of brain tissue". This case is remarkable for the fact that patient lived such a length of time from 31st October, 1926, when he first became unconscious=10 days.

BALANITIS GANGRENOSA.

Two cases received treatment during the year; both are of interest on account of the intense severity and the coincidence that they come from the same village.

Case No. 1 (No. 9,381) Male Tamil, aged 24 years, admitted in the venereal disease wards, District Hospital, Kuala Lumpur, on 16th October, 1926, patient denied all history of having had sexual connection. On admission he presented the following: The glans and body of penis with exception of \(\frac{1}{2}\) an inch at root consisted of a gangrenous mass from which was oozing a very foul smelling discharge of characteristic odour; the glans, prepuce and body of penis completely disorganised; urine was passed through a small opening corresponding to the level of the sulcus; patient's general condition very poor—wasted and somewhat emaciated on account of septic absorption. Duration of condition before admission: about 25 days, according to patient's statement.

Treatment was instituted at once—as much of the gangrenous portion as possible removed; three hourly dressings with hot saline, hydrogen peroxide, and dusting with bismuth formic iodide being carried out for 'the first few days; soon a considerable change in the local condition was noticed. This treatment was carried out at more extended intervals. On 30th October, 1926, the portion of penis remaining presented a clean and healing surface; patient soon after leaving hospital.

Case No. 2 (No. 10,508) Male Siamese, aged 56 years, admitted on 19th November, 1926, in venereal diseases wards, District Hospital, Kuala Lumpur. Patient denied all history of having had sexual connection. On admission he presented the following: Penis much swollen; complete phimosis; portion behind level of the sulcus sloughing; an inflammatory circular patch on skin of prepuce corresponding to the level of the sulcus, a profuse thin yellowish discharge issuing from mouth of prepuce having a most offensive and characteristic odour; skin of scrotum which is in conjunction with the undersurface of penis consisting of a large slough. Treatment started at once—prepuce slit open; as much of slough as possible removed: dressing with hydrogen peroxide; hot saline, and dusting with bismuth formic iodide being carried out two hourly for the first few days; this soon caused a considerable change; on 23rd November, 1926, condition much improved; dressing done only 4 times a day. He steadily improved and left hospital at the beginning of 1927. According to his statement he had been ill for seven days before admission into hospital.

CEREBRAL SYPHILIS.

The following two cases are of interest—satisfactory results being obtained by energetic treatment, in case No. 2 a negative blood Wassermann did not exclude syphilis.

Case No. 1. Male Chinese, aged 30 years, admitted in venereal disease wards, District Hospital, Kuala Lumpur, on 21st October, 1926. Patient unable to walk for the last 12 days; speech stuttering and indistinct; knee jerks—right: absent; left: slightly exaggerated; pupils reacting to light and accommodation; heart, lungs, spleen and liver—nil of note; complained of intense headache and pain in the neck, head retracted, throat deeply congested.

Wassermann blood, positive. Wassermann cerebro spinal fluid, positive. Treatment: stabilarsan 0.3 gm.—five injections every third day followed by 0.45 gm. once a week=ten injections in all. Bismostab: 10 weekly injections of 1 c.c.

Potassium iodide: starting with 10 grains 3 times a day, gradually increasing dose until a maximum of 30 grains 3 times a day for one week, followed by gradual reduction to 10 grains 3 times a day. Seven days from the commencement of treatment patient was able to speak clearly; pain in neck and headaches ceased; in 17 days' time he was able to walk.

Blood examination: Wassermann test (11th January) done three weeks after last injection was found to be negative.

Cerebro spinal fluid: Wassermann test (18th January) done four weeks after last injection was found to be negative.

Case No. 2.—Male Tamil, admitted in venereal disease wards, District Hospital, Kuala Lumpur, on 13th November, 1926, suffering from hemiplegia left side, pupils react to light and accommodation; knee jerks: absent; Wassermann blood: negative.

Wassermann cerebro spinal fluid positive.

Treatment: Stabilars five injections of 0.3 gm. at 3 days' interval followed by five injections of 0.45 gm. at weekly intervals.

Potassium iodide: increasing doses as in case No. 1.

Twelve days from the commencement of treatment patient was able to walk, his condition rapidily improved; he absconded on 21st December, 1926, five and a half weeks from the date of admission.

ARSENICAL DERMATITIS.

During the year 1926 thirteen cases of arsenical demartitis received treatment in the venereal disease wards, District Hospital, Kuala Lumpur.

All recovered with the exception of case No. 6 who was admitted with exfoliative stage well developed. Patient gave a history of having received four intravenous injections (does not know the name of the drug used) during the time he was a patient in one of the Singapore hospitals. Seven days after receiving the injections, he states, he developed a skin rash. He left the hospital of his own accord and came to Kuala Lumpur seeking admission into the venereal disease wards of the District Hospital, on 28th July, 1926.

All the thirteen cases are set down concisely in tabular form. See table "A".

During the second half year reports on all cases suffering from dermatitis following the administration of one of the arsenical preparations, have been received from the varicus hospitals in the Federated Malay States—see table "B".

TABLE "A."

1	AF	ARSENICAL		DERMATITIS CAS	CASES	TREATED	N	DISTRICT HOSPITAL, KUALA LUMPUR, DURING 1926.	DURING	1926.	
Case.	Date of admission.	Nationality.	Sex.	Drugs.	No. of in- jections.	Total amount.	Appearance of first symptom after last injection.	Condition of skin.	Treatment and total amount.	Duration of skin condition.	Result.
-	15th Jan, 1926 Malay	Malay	K.	Novarsenobillon Bismostab		0.45 gras. .1 gra.	5 days	Generalised orythema	Thiestab 3.15 gms.	12 days	Discharged 27th January, 1926
01	12th March, 1926 Tamil	Tamil	X.	Stabilarsan Bismostab	00	4.5 gms. 1.0 gm.	4 days	Keratosis of palms of hands and soles of feet	Thiostab 3.6 gms.	13 days	Discharged 25th March, 1926
00	4th Feb., 1926	1	9.	Novarsenobillon Bismostab	10 10	1.5 gms. .5 gms.	2 days	Pruritis followed by a generalised crythema	Thiostab 2.5 gms.	9 days	Discharged 13th February, 1926
4	10th June, 1926	:	p.i	Stabilarsan Bismostab	00 00	2.4 gms. .8 gms.	Unknown	Admitted in exfoliative stage well developed	Thiostab 4.05 gms.	20 days	Discharged 1st July, 1926
10	17th March, 1926		M.	Novarsenobillon Bismostab	10 10	2.25 gms. .5 gms.	7 days	Generalised crythema	Thiostab 4.05 gms.	13 days	Discharged 30th March, 1926
0	28th July, 1926		M.	6.	+ 6	Unknown	7 days	uffering from a generalised exfoliative is principally affecting face, chest, lower of abdomen. Axilla and elbow joints	Thiostab 4.5 gms.	24 days	Died on 20th August, 1926
			× 12	Silventon Silven	22	10 10		nssured. Conjunctivitis present, lips alcerated, mucous membrane of mouth sore and painful. On 12th August, 1926, much improved, on 17th August, 1926, he took a turn for the worse, dying on 28th August, 1926. Post-mortem and histological reports given below.*		100	
1-	3rd August, 1926 Sikh	Sikh	M.	Stabilarsan Bismostab	01 01	0.9 gms.	15 days	Pruritis followed by a generalised crythema. Face Thiostab swollen, slight exfoliation 3.15 gm	Thiostab 3.15 guss.	13 days	Discharged 16th August, 1926

* Case No. 6: Post-mortem notes: Both kidneys—capsules strip with little difficulty exposing irregular flat projections. The surface veins are stellate and congested. Cortex—distinction between certex and medulas not epidemics—congested neutring a naturage condition. Splen—enhanged to thrive normal size. Lance—normal state of epidemics—special descriptions—specially above unablical region exuding thick pass when cut into.

Histological report received from Government Laboratories—calls of the surface layer of the epithelium give the appearance of increased activity. Stratum lendum thickened with increased production of kerstin. There is the hold vegeted from the continuence of the cortium in addition there is one large of increased activity. Stratum lendum thickened with increased production of kerstin. There is the hold vegeted of the cortium. In addition there is one large well-defined absences at the junction between the cortium and the subculatorial accordance of exceeding a neutron of surface in the contract of various lebules. Splean: Congested with presence of excessional thrombones of vessels. Kidney: Acutely congested. In addition there are magnetic actually believed to the changes occurring fellowing minute septic mixtures. In some situations there is commencing absects formation. Microscopically the appearances of the kidney lesions have a close recemblance to the changes curring fellowing minute septic mixtures.

TABLE "A"-(cont.).

	Result.		35 days Discharged 5th October, 1926	Discharged 8th Sept., 1926	Discharged 23th October, 1926	Discharged 31st Dec., 1926	To the last of the	The state of the s	Still in hospital	Still in bospital
6—(cont.)	Duration of skin condition.		35 days	4 days	10 days	48 days			Still	St.
JRING 192	Treatment and total amount.		Thiostab 2.25 gms.	Thiostab 1.8 gms.	Thiostab 3.15 gms.	Ametox 3.15 gms.	11		Ametox 2.25 gms.	Ametox 0.9 gms.
T HOSPITAL, KUALA LUMPUR, DURING 1926-(cont.).	Condition of skin,	THE RESERVE THE PERSON NAMED IN COLUMN TWO IS NOT THE PERSON NAMED IN COLUMN TWO IS NAMED IN COLUMN TWIND TWO IS NAMED IN COLUMN TWO IS NAMED IN COLUMN TWO IS NAMED IN	Generalised crythems followed by slight exfoliation. Scalp later on covered with pustules, face much swollen	Pruritis followed by a generalised erythema		35	tions of ametox dermatitis appeared. 5th November, 1926, blood examined for Van den Bergh's test. Report from Government Labo- ratory as follows: Van den Bergh's reaction is	oppassioned indicates the presence of azobilirabin and bilirabin in the blood stream. This may be taken to indicate that the jaundice is of both toxic and obstructive type	On 6th December, 1926, generalised crythema, on Ametox 11th face swellen, ordena of feet, on 12th exfolintion on chest, ulceration of lips and conjunctivits. These symptoms remain in severity. On 3rd December, 1926, patient's condition same	19th December, 1926, patient developed pruritis Ametox followed by erythema on his back, chest, arms and legs. On 3bth December, 1926, exfoliation appeared which rapidly became generalised. Patient complains of pain when swallowing
IN DISTRICT	Appearance of first symptom after last injection.		3 days	4 days	7 days	10 days			2 days	4 days
VTED	Total amount.	do lone	3.0 gms. 1.0 gm.	0.45 gms. 0.1 gm.	1.5 gms. 0.9 gms. 0.8 gms.	2.4 gms. 0.8 gms.			1.35 gms. 0.45 gms. .4 gms.	0.45 gms. 0.3 gms. .2 gms.
TRE	No. of in- jections.	10	10		10 00 00	00 00			⊗ 4	
TITIS CASES	Drugs.	September	Stabilarsan Bismostab	Stabilarsan Bismostab	Novarsenobillon Stabilarsan Bismostab	Novarsenobillon Bismostab	- September	ALIE CONTRACTOR	Stabilarsan Norarsenobillon Bismostab	Stabilarsan Novarsenobillon Bismostab
MAT	Sex.	-	ai.	M.	×	M.			K.	×
ARSENICAL DERMATITIS	Nationality.		Japanese	Eurasian	Tamil	1			:	1 Colombia
ARSEN	Date of admission.		31st Aug., 1926 Japanese	4th Sept., 1926 Eurasian	13th Oct., 1926	13th Nov., 1926	the true true	NOT CHILL AND	6th Dec., 1926	19th Dec., 1926
	Case.	-	00	6.	10	11			22	13

Ametox (May and Baker) and Thiostab (Boots) were administered according to the method advocated by Dr. Andrew Balfour, the Director of the London School of Hygeine and Tropical Medicine,

CABLE "B."

	Result.	Skin dry on 28th August, 1926 Skin clear on 19th August, 1926, deve- loped broncho- p ne um on is, died on 24th August, 1926 Skin clear on 19thAug, 1926, of dysentery No record " Condition cleared No record " Skin clear on Skin clear on Skin clear on Skin clear on	04n D0c., 1925
1926.	Duration of skin condition.	21 days 22 days No record 2 days 3 days	
LUMPUR,	Treatment and total amount.	Sodium Thiostab 3.6 gms. Thiostab 3.6 gms. Ametox no record Ametox Finosuph Contramin Noi record Contramin Nii. Nii. Ametox Anetox Outgans Contramin Nii. Anetox Outgans Contramin Nii. Thiostab	0.0 gm.
THAN DISTRICT HOSPITAL, KUALA	Condition of skin.	M & & & & & & & & & & & & & & & & & & &	chest and back of neck
OTHER TH	Appearance of first symptom after last injection.	3 days 3 days 4 days 4 days 7 days 2 days 2 days 3 days 3 days 3 days 3 days	
ATED IN	Total amount.	3.65 gms. 1.8 gms. 2.7 gms. 0.9 gms. 0.6 gms. 6 ctgs. 1.85 gms. 6 ctgs. 1.05 gms. 1.05 gms.	
EAT	No. of in- jections.	× + × × × × × × × × × × × × × × × × × ×	
ARSENICAL DERMATITIS: TRE	Drugs.	Novarsenobillon Stabilarsan Novarsenobillon Stabilarsan Novarsenobillon Stabilarsan Novarsenobillon Novarsenob	
DER	Sex.	F K K K K K K K K K K K K K K K K K K K	
SENICAL	Nationality.	Chinese Chinese Singhalese Other Indian Chinese Malay Chinese	
AR	Date of admission.	m - 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9 9	(Sultan Street) Kaala Lumpur
	Case.	1 8 8 4 10 20 11 21	

	8 9 7 1				
		fr.			
		1/2		1	
		112		21	
		1131 +		5	
		1 200		24	
		18			
		1121			
		1/3/	1100		
			# 30		
			-		
		The second			
HI IS					
1.44					
1 184 2818:					







CHART SHOWING VENEREAL DISEASES TREATED IN HOSPITALS AND DISPENSARIES OF THE FEDERATED MALAY STATES DURING THE YEAR 1926.






GRAPH SHOWING ADMISSIONS AND DISEASES.
1926.





SULTAN STREET VENEREAL DISEASES TREATMENT CENTRE KUALA LUMPUR.

GRAPH SHOWING ADMISSIONS AND DISEASES FROM JULY 1924
DATE ON WHICH THIS CLINIC WAS STARTED, TO THE END OF 1926.






SEREMBAN.

GRAPH SHOWING ADMISSIONS AND DISEASES.

1926.



THIS CLINIC WAS TRANSFERED FROM THE GENERAL HOSPITAL

TO THE GOVERNMENT TOWN DISPENSARY AT THE BEGINNING OF OCTOBER.

Drawn and Printed by F. M. S. Surveys No. 205-1927.

GRAPH SHOWING ADWISSIONS AND DISEASES.



CHART SHOWING 18 CASES TREATED FOR GONORRHOEAL RHEUMATISM. THE FIRST 4 CASES WERE TREATED WITH SULPHOSTAB 'BOOTS' AND THE REMAINING WITH CONTRAMINE

[q	AYS IN HOSPI		29/30	NATIONALITY	201111111111111111111111111111111111111
	++++	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	111	DATE ADMITTED	12. 10. 26.
1	+++	000000000000000000000000000000000000000		RESULT	IMPROVED
5	SULPHOSTAB	DOSE 3 2 3 1 3		NATIONALITY	CHINESE.
2	++++	50500000000		DATE ADMITTED RESULT	18, 10, 26,
-	++			HESULI	CURED
-	+	0 0 0 0 0 0		MATIONALITY	
-	SULPHOSTAB ++++	DOSE'3 9 9 3 3		DATE ADMITTED	TAMIL. 29, 10, 26,
3	+++			RESULT	CURED.
1	++	0000000000000	-		
5	SULPHOSTAB	DOSE, 3 3 3 3		NATIONALITY	MALAY,
4	++++	0 0 0 0 0 0 0 0 0	-	DATE ADMITTED	15. 10. 26. GURED.
	++	0 0 0 0 0 0 0 0 0			
-	CONTRAMINE	DOSE 29 95 95 95 95 95	0-0	NATIONALITY	TAMIL.
	1111			DATE ADMITTED	19.10.26.
5	+++	660000000000000000000000000000000000000		RESULT SLIGHT	IMPROVEMENT
	+				
0	CONTRAMINE ++++	DOSE 25 125 45 25		DATE ADMITTED	7AMIL. 7, 11, 26.
6	+++			RESULT	CURED.
-	‡ +	0-	H		
9	CONTRAMINE	DOSE 20 01 00 20		NATIONALITY	TAMIL
7	++++	6-9-8		DATE ADMITTED	8, 11, 26,
-	++	6-6-6-6-6-6		RESULT	CURED
-	CONTRAMINE			***************************************	
	CONTRAMINE ++++	DOSE 28 02 28 28	111	NATIONALITY DATE ADMITTED	TAMIL. 8, 11, 26,
8	+++			RESULT	CURED.
	++	******************************	0.0		
(CONTRAMINE	DOSE 25 23 25 26 25		NATIONALITY	TAMIL.
9	++++	00000	+++	DATE ADMITTED RESULT	14. 12. 26, CURED.
	++				50710.50
-	CONTRAMINE	DOSE 98 28 28 28	+++	NATIONALITY	TAMIL.
	++++			DATE ADMITTED	18, 12, 26,
10	+++	0 0 0 0	111	RESULT	CURED.
	+				
5	CONTRAMINE	DOSE 25 15 25 25	++	DATE ADMITTED	CHINESE. 24, 12, 26,
11	+++	00000			L IN HOSPITAL
-	++		++		
C	CONTRAMINE	DOSE 28 123 125 125		NATIONALITY	CHINESE.
12	++++	0 0 0 0 0	+++	RESULT STILL	24, 12, 26. L IN HOSPITAL
	++		Ш		
C	CONTRAMINE	DOSE, 48 48 48		NATIONALITY	CHINESE.
	++++	0000		DATE ADMITTED	29. 9.26.
13	+++	9 8 6 8 6 8 6 8 6 8 8 8	111	RESULT	CURED.
	+			MATIONALISM	All Indiana
10	CONTRAMINE	DOSE 24 25 25 25 25 25	+++	DATE ADMITTED	7, 10, 26,
14	+++			RESULT	CURED.
1	++	0 0 0 0 0 0 0			
C	CONTRAMINE	DOSE 25 95 95		NATIONALITY	TAMIL.
15	++++	0 0 0	-	RESULT	10. 7. 26. CURED.
-	++				
	ONTRAMINE	DOSE (03 (25) 25	H	NATIONALITY	MALAY.
0	++++	64		DATE ADMITTED	8. 7. 26.
	4 4 4	00000		RESULT	CURED.
	+++		111		
16	+ +		_		CHINESE.
16	+ +	DOSE 25 05 29		NATIONALITY DATE ADMITTED	
16	+ +			NATIONALITY DATE ADMITTED RESULT	21. 8. 26
16	+ + + + + + + + + + + + + + + + + + +	DOSE 23 85 73 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8		DATE ADMITTED	
16	CONTRAMINE ++++	DOSE 038 048 028 029		DATE ADMITTED	21. 8. 26
16	CONTRAMINE ++++ +++ ++ CONTRAMINE ++++ +++	DOSE 23 05 73 05 05 05 05 05 05 05 05 05 05 05 05 05		DATE ADMITTED RESULT NATIONALITY DATE ADMITTED	21. 8, 26 ABSCONDED TAMIL. 29. 9. 26,
17	CONTRAMINE ++++ +++ +++ CONTRAMINE	DOSE 038 048 028 029		DATE ADMITTED RESULT	21. 8, 26 ABSCONDED



ANNUAL REPORT OF THE OPHTHALMIC OF THE IPOH EYE-CLINIC, FOR THE YEAR 1926.

GENERAL REMARKS.

There has been a steady increase in the daily number of attendances at the Eye-clinic, Ipoh. The number of new cases in 1926 was 3,470, and the total number reached 10,367. The number of inpatients was 491. Only serious cases and those requiring operation were admitted.

The following figures indicate the growth of this department during the past four years:

				210	 · ·		
Year.			01	ut-patients.	 in-patients.	1	Total treated.
1923		 		1,535	 234	***	9,011
1924	***	 		1,780	 391		6,373
1925	***	 	***	2,151	 386		7,141
1926		 		3.470	491		10.867

It will be noted that the number of out-patients (new cases) has been almost trebled, and an increasing amount of work in the out-patients' department has of necessity developed.

costly developed.			IN-	DOOR.					
Diseases.—									
Conjunctivitis						1000	10.10	A.	lmissions.
Gon. ophthalm	ia								37
Pterygium		***			***	***	who I		12
Lids				***			****		9
Trachoma							***		48
Ulcer cornea						***			64
Keratitis				***		***			22
Iritis				***			***		31
Cataract					***				67
Retinitis		***							6
Glaucoma									12
Lachry. app.									10
Injuries.—									
Lids				***					3
Cornea									16
Iris						7			2
Lens									3
Eyeball									5
Refraction									
Hyperopia					***				1
General.									
Sixth nerve par	alveia								4
Facial paralysis									i
Keratomalacia					***				3
Facial erysepile									1
Herpes frontali									2
			08		-			TITLE OF	1999
Disorganised eyes	MIL I								177
One		***		***		***	***		17
Both			***	***	***			***	4
							Total	2231	491
Nationality	EVED								
Chinasa							***		314
Indiana									134
36-1									42
D									1
							02500089	-	-
							Total		491

See							
Sex.—						Ad	missions
Males							391
Females	***						100
					m-4-1		401
					Total		491
	MA	JOR.					
Operations.—							
Extraction of cataract			***	***	***	***	68
Iredectomy		***		•••		•••	29
Iridotomy		***		***		•••	11
Trephining		***		***	***		13
Enucleation of eyeball	***	***			***	***	2
Plastic operation of entropic							1
Dissection of lachrymal sa							1
Extraction of lens (dislocat	ed)	***			· · ·		2
					Total		122
					20001		
AV 311	Mi	NOR.					
Needling	***		•••	•••	***	•••	26
Dissection of pterygium	***	***		•••	***	***	12
Paracentesis			***		***		10
Dilatation of canaliculus			***	***	***		36
Incision lachrymal abscess			***				2
Tatooing					***		2
Incision abscess lid				•••			2
Expression of lid for trach	oma			•••			50
Excision of growth lid						***	1
Removal of foreign body co			***	•••		***	37
Snipping off prolapsed iris				***			3
					Total		181
						-	The same
Sub-conjunctivitis injections		***					5
Result of Cataract Operations	-						
Good result				***			55
T-111	***		***			***	3
Iritis Panophthalmitis							1
Panophthaimitis					***	***	1
					Total		68
	OTTE	DOOR.					-
Diseases.		w cases.		Repetitio	me	-	Total.
Conjunctivitis		,892		3,191			,083
Gon. ophthalmia		29		42			71
Pterygium		36		29			65
Lids		97		126			223
Trachoma		345		2,051		2	2,396
Ulcer Cornea		153		507			660
Keratitis		116		396			512
Iritis		51		172			228
Cataract		129		32			161
Retinitis		32		18			45
Glaucoma		21		14			35
Oc. muscles		12		8			20
Lachry. app		32		32			64
Carried forward		2,945		6,613		9	,558

Diseases.		1	New cases		Repetition	8.	Total.
Brough	t forward				6,613		9,558
Injury.—					18:50-71		
Lids			17		12		29
Conjunctiva			19		84		58
Cornea			64		81		145
Iris			4		3		7
Lens			4		3		7
Eyeball			9		1		10
Refraction.—							
Myop. astigmatisn	1		27		_		27
Myopia			71		19		90
Hyp. astigmatism			15				15
Hyperopia			21		1 1		21
Asthenopia			59	11	2		61
Night blindness	***		4		1000		4
Amblyopia			1		100		1
Diplopia	W COUNTY		2				2
Presbyopia			98				98
General.—			-10			100	
Sixth nerve paraly	sis		4		1		5
Third nerve paraly			1		1000		1
Facial paralysis			11		12	***	23
Sec. anaemia			3		1	***	8
Variable			9			***	9
Keratomalacia		***	2		26	***	28
Facial erysepelas		***	3		20	***	3
Herpes frontalis		***	3	***	29		32
T 32		***	- 4	***	20		
N	***	***	3			***	4
	•••		0			***	3
Disorganised Eyes			***				
One	•••	***	59		60	•••	119
Both	***	***	13	***	1		14
	Total		3,470		6,897		10,367
	Total	***	0,410		0,001		10,007
Nationality.—							
Chinese			2,470		5,704		8,174
Indians			642		858		1,500
Maiays			201		225		428
Europeans			131		7		138
Eurasians			19		59		78
Japanese			7		44		51
	Total		3,470	***	6,897		10,367
Ga							
Sex.— Males			0.051		E 010		7 070
			2,851		5,019		7,870
Females	***		619		1,878		2,497
	Total		3,470		6,897		10,367
	20001						
OLI / DI D			3 21	777 7	Si 102 10		

Chief Diseases.—From the above records it will be seen that amongst the chief diseases. Conjunctivitis holds the chief place. The main type was a koch-weeks bacillus infection. Some severe types were seen chiefly in rickshaw pullers, who live together in kongsis and pass the infection on from one to another.

There was no definite period during the year for its appearance.

The disease prevailed throughout the year.

Gonorrheal Ophthalmia.—Thirty-seven cases were admitted, ten were infants and 27 adults, eight were females and 29 males, three were Tamils and 34 Chinese.

Fifteen of the 37 had complications, i.e., corneal ulcers in one or both eyes; three had both eyes completely disorganised before admission and three had lost one eye.

Of the 22 uncomplicated cases 19 made excellent recoveries.

Of the infants one died of congenital syphilis during the treatment, but the results were uniformly better in this class of patient than in the adult.

The use of urine in the treatment of eye affections has been recorded in Ancient writings and it may be mentioned that an instance of its application was noted during the year. An old out-patient with trachoma was advised by his friends to use his own urine as an eye-wash. He did so and was admitted with Gon. ophthalmia. It was then discovered that he was suffering from gonorrhea.

Trachoma.—Three hundred and forty-five new cases were treated. All that can be said is that treatment keeps the disease in abeyance. Figures showing new arrivals from China with this disease are being compiled with a view to forwarding a report later on.

Injuries.—One hundred and seventeen cases of injury to the eye and its appendages were treated. The cornea provided 64 cases almost 50 per cent. Owing to the pain and loss of vision these cases seek advice, but unfortunately in most of them valuable time has been lost and infection had occurred.

Operations.—Three hundred and three operations were performed of which 122 were major eye operations and 181 minor. Sixty-three cataract extractions with 55 successes were the chief major operations. Three failures were due to chronic disease of the Vetina Khoroid, which could not have been forseen.

Conclusion.—As in former years, dresser Dias has been of great assistance to me. Nurses are encouraged to attend the out-patients and one is fixed for duty for a month at a time. This is a very good scheme, as practically all the nursing staff are now conversant with the methods of treatment of the more common type of eye disease. The same scheme has been extended to the dresser staff and it is to be hoped that the same satisfactory results will be attained.

P. H. HENNESSY,

Medical Officer and Ophthalmic Surgeon, F.M.S.

ANNUAL REPORT OF THE OPHTHALMIC CLINIC, GENERAL HOSPITAL, KUALA LUMPUR, FOR THE YEAR 1926.

Sir,—I have the honour to submit my annual report on the ophthalmic work in Kuala Lumpur for the year ending 31st December, 1926.

The number of patients treated were 2,962; of these 850 were in-patients and 2,112 were out-patients.

The figures according to nationalities were as follows:

Nationa	lity.					I	n-patient	8.	Out-patients.
Chinese	***		***		***	2000	351		457
Tamils				***			258	***	642
Malays		teres.			***		145		461
Sikhs		NAVE TOO	Seell	11 4 84	1		63		197
Indian Mohan	nedans			***	***		19		164
Eurasians				***			11		100
Europeans				***			Nil		65
Singhalese	1552 101	***	2000	***			3		18
Japanese	********	MORN	Beeter	(1757)	SER YO	***	1		8
					Total		850		2,112
									-

The total number of attendances were 21,941. The total number treated shows an increase by 1,000 over the number treated last year—an increase of almost 60 per cent.

All nationalities now come to realise the benefits of the work. Patients come from Tanjong Malim on the north to Johore in the south and from Pahang, Negri Sembilan and Malacca.

The disease met with are shown on anatomical basis in the attached table.

But the most common diseases met with in their order of frequency are as follows:

- 1. Conjunctivitis.
- 2. Catarrhal ophthalmia (koch-week's and morax).
- 3. Trachoma.
- 4. Errors of refraction.
- 5. Ulcer of cornea.
- 6. Iritis.
- 7. Opacities of cornea.
- 8. Meibomiah and stye abscess.
- 9. Gonorrhoeal ophthalmia.
- 10. Glaucoma.
- 11. Cataract.

Conjunctivitis was met with among all races; trachoma among the Chinese and Northern Indians; gonorrhoeal ophthalmic among Chinese; of the errors of refraction, myopia was prevalent among the North Indians and Chinese and astigmatism among Europeans.

Retinoscopy and Refraction.—Many cases of errors of refraction were accidentally discovered in examining persons joining the Government service, all of whom have their vision tested at the clinic. This has done a great deal towards greater efficiency in the public services than has been the case hitherto.

Retinoscopy also helped to discover deep-seated troubles connected with the fundus and the refractive media and the uveal tract.

A fair number of optic neuritis and atrophy and choroiditis were met with.

A table showing the details is appended herewith.

Operations.—There were done in all 295 operations; of these 83 were major and 212 minor, a distinct increase on the number for 1925 which was 217.

In the performance of operations for removal of cataract (Senile) the irrigation method of the Elliot School was abandoned in many instances with distinct advantage in saving after complications which are not infrequent in the above method.

General.—During the year under review an epidemic of catarrhal ophthalmia was dealt with. As was the case in the 1924 Epidemic the infection was of the mixed koch-week's and morax axenfeld variety and responded quickly to treatment. It affected all nationalities.

It must be said in conclusion that the usefulness of the clinic is very largely appreciated by the public. The quality of the work has greatly improved and with the appliances and conveniences that are to be provided shortly, still greater improvement and efficiency may be expected. We may then have a better equipped and well ordered and organised section of curative and preventive medicine.

I have the honour to be, Sir,

Your obedient servant,
(Sgd.) A. VISWALINGAM,
Deputy Medical Officer, Ophthalmic Clinic,
General Hospital, Kuala Lumpur.

19th January, 1927.

INCIDENCE OF DISEASES OTHER THAN THOSE SHOWN UNDER REFRACTION AND RETINOSCOPY.

	REFR	ACTIO	N A	AND RE	TINO	SCOPY.			
Disease of:									
The Lids									
Sebaceous cyst	b.			4990	***		444		7
Tarsal abscess					***			***	2
Blepharitis									38
Meibomain abs	cess			***			***	***	43
Entropion									21
Stye				***					70
Ptosis		***		***		***		***	8
Trichiasis						***	***		9
The conjunctiva									
Foreign body									34
Trachoma						-1			162
Conjunctivitis							14000		358
Pterygium				***				***	47
Catarrhal ophth									274
Haemophilia									3
Phlyctenular c									43
Follicular conju									25
Angular conju									52
Traumatic con				***		***			25
Xerosis conjunc				***	411			***	8
Pinguecula							***		7
Symblepharon					***	***			2
The Comea.									
771									127
Keratitis	***	***	***	***	***	***	***	***	31
Kerato malacia	12.224	***		***	22.	HIGHT IN	***		5
Leucoma adher		***	***					***	44
Opacities	CHS				***				15
Nebula cornea									13
Macula cornea			***						47
Staphyloma									21
Abrasions									23
				Indian			ull mo	rinoit.	-
The Lens.—									PO.
Cataract			***	***		***	***	***	72
The Iris.—	****								
Iritis—Irido cy			***	***		***	***		58
Prolapse of Ir	18	*** (0		3000	***	***			17
Uveitis		***				***		***	4

The Eyeball.—								
Panophthalmitis		***		***			***	11
Phthisis buibi	***						***	17
Contusion of eyeball	***	***			***		***	12
Glaucoma			***			***	***	26
Rupture of eyeball							***	14
Lachrymal System.—								
Dacryocystitis		***			***		***	4
The Sclera								
Episcleritis		***						2
DPT	INOSCI	ODV 4	MD D	EFRAC'	TTON			
Refraction.—	INUSCI	OP X A	IND H	EFRAC	110N.			
Danmakania								603
Anthonomia	1001	***	111	***		***		172
Astimustians	**	***	***	***	100		***	48
Myopia		***	***		***			141
Hamanania		***	***		***	***	***	18
Presbyopia	***	111	100	4 - 117	111		***	57
Aphakia			***					9
Toxic amblyopia		***	***			***		5
Squint								10
Funduscopic Examinat		No Serve	13775-1	In District	1000	I I I SEE LOOF	1110	PERM
Retinitis		Jacon						17
Retino choroiditis								9
Optic neuritis								25
Optic atrophy								17
Dislocated lens			***					2
Neuro retinitis								19
Opacity lens								7
Choroiditis								4
AND THE RESERVE TO TH								
Choroidal atrophy			***		***			3
								3
	рнтн	ALMIC	OPE	RATIO				3
on the displayer to the O		ALMIO M	OPE	RATIO	NS.			
Radical operation for	entro	ALMIC M/ opion	OPE	RATION	vs.	7)		8
Radical operation for Transplantation of p	r entro	ALMIC M/ opion um	OPE JOR. (parti	RATIO	vs. etomy	') 		8 7
Radical operation for Transplantation of p Paracentasis of anter	terygiu	ALMIC M/ opion im	OPE	RATION al tarse	etomy	') 		8 7 2
Radical operation for Transplantation of p Paracentasis of anter Removal of foreign be	r entro	ALMIC Mopion om amber	OPE	RATIO	octomy	r) 		8 7 2 5
Radical operation for Transplantation of p Paracentasis of anter Removal of foreign be Dissection of leucome	r entro	ALMIC ppion amber om eye	OPE AJOR. (parti	RATIONAL tarse	etomy	·) 		8 7 2 5 1
Radical operation for Transplantation of p Paracentasis of anter Removal of foreign be Dissection of leucoma Ablation of staphylom	r entro	ALMIC ppion amber om eye	OPE AJOR. (parti	RATIONAL tarse	octomy	r) 		8 7 2 5
Radical operation for Transplantation of p Paracentasis of anter Removal of foreign be Dissection of leucoma Ablation of staphylom Optical iridectomy	r entro	ALMIC Mopion im namber om eye ea ornea	ope Jor. (parti	RATION al tarse 	etomy	r) 		8 7 2 5 1 5
Radical operation for Transplantation of p Paracentasis of anter Removal of foreign be Dissection of leucoma Ablation of staphylom	r entro	ALMIC Mopion im namber om eye ea ornea	ope Jor. (parti	RATION al tarse 	vs.	') 		8 7 2 5 1 5 3
Radical operation for Transplantation of p Paracentasis of anter Removal of foreign be Dissection of leucoma Ablation of staphylom Optical iridectomy Perepheral iridectomy	r entro	ALMIC Mopion um namber om eye ea ornea	ope Jor. (parti	RATION al tarse 	etomy	") 		8 7 2 5 1 5 3 3
Radical operation for Transplantation of p Paracentasis of anter Removal of foreign be Dissection of leucome Ablation of staphylom Optical iridectomy Perepheral iridectomy Trephining for glauc	r entro	ALMIC Mopion um namber om eye ea ornea	ope AJOR. (parti ball and ir 	RATION al tarse 	octomy	") 		8 7 2 5 1 5 3 3 4
Radical operation for Transplantation of p Paracentasis of anter Removal of foreign be Dissection of leucome Ablation of staphylom Optical iridectomy Perepheral iridectomy Trephining for glauc Discission of catarac Evisceration of eyebs	r entro	ALMIO M/opion um camber om eye ea cornea	OPE AJOR. (parti	RATIONAL tarse	NS.	"		8 7 2 5 1 5 3 3 4 5
Radical operation for Transplantation of p Paracentasis of anter Removal of foreign be Dissection of leucome Ablation of staphylom Optical iridectomy Perepheral iridectomy Trephining for glauc Discission of catarac Evisceration of eyebs	r entro	ALMIO M/opion um camber om eye ea ornea	OPE AJOR. (parti	RATION	NS.	"		8 7 2 5 1 5 3 3 4 5 8
Radical operation for Transplantation of p Paracentasis of anter Removal of foreign be Dissection of leucome Ablation of staphylom Optical iridectomy Perepheral iridectomy Trephining for glauc Discission of catarac Evisceration of eyebs Enucleation	r entro	ALMIO M/opion um namber om eye ea ornea glaucoi	ope AJOR. (parti	al tarse	NS.	"		8 7 2 5 1 5 3 4 5 8 1
Radical operation for Transplantation of p Paracentasis of anter Removal of foreign be Dissection of leucome Ablation of staphylom Optical iridectomy Perepheral iridectomy Trephining for glauc Discission of catarac Evisceration of eyebs Enucleation Operation for symble	r entro	ALMIO M/opion um namber om eye ea ornea glaucoi	ope AJOR. (parti	al tarse	NS.)		8 7 2 5 1 5 3 4 5 8 1 2 29
Radical operation for Transplantation of p Paracentasis of anter Removal of foreign be Dissection of leucome Ablation of staphylom Optical iridectomy Perepheral iridectomy Trephining for glauc Discission of catarac Evisceration of eyebs Enucleation Operation for symble	r entro	ALMIO M/opion um namber om eye ea ornea glaucoi	ope AJOR. (parti	al tarse	NS.	"		8 7 2 5 1 5 3 3 4 5 8 1 2
Radical operation for Transplantation of p Paracentasis of anter Removal of foreign be Dissection of leucome Ablation of staphylom Optical iridectomy Perepheral iridectomy Trephining for glauc Discission of catarac Evisceration of eyebs Enucleation Operation for symble	r entro	ALMIO M/opion im amber om eye ea ornea	ope AJOR. (parti	al tarse	NS.)		8 7 2 5 1 5 3 4 5 8 1 2 29
Radical operation for Transplantation of p Paracentasis of anter Removal of foreign be Dissection of leucome Ablation of staphylom Optical iridectomy Perepheral iridectomy Trephining for glauc Discission of catarac Evisceration of eyebs Enucleation Operation for symble	r entro	ALMIO M/opion im amber om eye ea ornea glaucoi	ope Jor. (parti ball and ir ma	al tarse	NS.)		8 7 2 5 1 5 3 4 5 8 1 2 29
Radical operation for Transplantation of p Paracentasis of anter Removal of foreign be Dissection of leucome Ablation of staphylom Optical iridectomy Perepheral iridectomy Trephining for glauc Discission of cataract Evisceration of eyebs Enucleation Operation for symble Removal of cataract	r entro	ALMIO M/opion im amber om eye ea ornea glaucor	ope ajor. (parti ball and ir ma	al tarse	NS.)		8 7 2 5 1 5 3 3 4 5 8 1 2 29 83
Radical operation for Transplantation of p Paracentasis of anter Removal of foreign be Dissection of leucome Ablation of staphylom Optical iridectomy Perepheral iridectomy Trephining for glauc Discission of cataract Evisceration of eyebs Enucleation Operation for symble Removal of cataract Removal of cataract	r entro	ALMIO M/opion im amber om eye ea ornea glaucor	ope ajor. (parti ball and ir ma	al tarse	NS.	Total		8 7 2 5 1 5 3 3 4 5 8 1 2 29 83 72
Radical operation for Transplantation of p Paracentasis of anter Removal of foreign be Dissection of leucome Ablation of staphylom Optical iridectomy Perepheral iridectomy Trephining for glauce Discission of cataract Evisceration of eyebs Enucleation Operation for symble Removal of cataract Removal of foreign Removal of foreign Removal of foreign	r entro terygin rior ch ody fro a corn- na of co- tor g oma et all pharon senife	ALMIO M/ opion im namber om eye ea ornea glaucor Mi cornea onjuno	ope ope open open open open open open op	al tarse	NS.	')		8 7 2 5 1 5 3 3 4 5 8 1 2 29 83 72 54
Radical operation for Transplantation of p Paracentasis of anter Removal of foreign be Dissection of leucome Ablation of staphylom Optical iridectomy Perepheral iridectomy Trephining for glauce Discission of cataract Evisceration of eyebs Enucleation Operation for symble Removal of cataract Removal of foreign Removal of foreign Removal of foreign Stye abscess	r entro terygic rior ch ody fro a corn- na of co- y for a oma et all pharon senife	ALMIO M/ opion im namber om eye ea ornea glaucor Mi cornea onjuno	ope ope open open open open open open op	al tarse	vs.	')		8 7 2 5 1 5 3 3 4 5 8 1 2 29 83 72 54 49
Radical operation for Transplantation of p Paracentasis of anter Removal of foreign be Dissection of leucome Ablation of staphylom Optical iridectomy Perepheral iridectomy Trephining for glauce Discission of cataract Evisceration of eyebs Enucleation Operation for symble Removal of cataract Removal of foreign Removal of foreign Removal of foreign Stye abscess Meibomain abscess	r entro	ALMIO M/opion im samber om eye ea ornea glaucoi Mi cornea onjuno	ope ope open open open open open open op	al tarse	vs.	Total		8 7 2 5 1 5 3 3 4 5 8 1 2 29 83 72 54 49 34
Radical operation for Transplantation of p Paracentasis of anter Removal of foreign be Dissection of leucome Ablation of staphylom Optical iridectomy Perepheral iridectomy Trephining for glauce Discission of cataract Evisceration of eyebs Enucleation Operation for symble Removal of cataract Removal of foreign Removal of foreign Removal of foreign Stye abscess Meibomain abscess Tarsal abscess	r entro terygic rior ch ody fro a corn- na of co- y for a oma et all pharon senife	ALMIO M/opion im samber om eye ea ornea glaucoi Mi cornea onjuno	ope Jor. (parti	RATIONAL tarse	vs.	Total		8 7 2 5 1 5 3 3 4 5 8 1 2 29 83 72 54 49 34 2
Radical operation for Transplantation of p Paracentasis of anter Removal of foreign be Dissection of leucome Ablation of staphylom Optical iridectomy Perepheral iridectomy Trephining for glauc Discission of catarac Evisceration of eyebs Enucleation Operation for symbles Removal of cataract Removal of foreign Removal of foreign Removal of foreign Stye abscess Meibomain abscess Tarsal abscess Peritomy	r entro terygic rior ch ody fro a corn- na of co- y for a oma et all pharon senife	ALMIO M/opion im tamber om eye ea ornea glaucoi Mi cornea onjuno	ope Jor. (parti	RATION al tarse	vs.	Total		8 7 2 5 1 5 3 3 4 5 8 1 2 29 83 72 54 49 34 2 1 2
Radical operation for Transplantation of p Paracentasis of anter Removal of foreign be Dissection of leucome Ablation of staphylom Optical iridectomy Perepheral iridectomy Trephining for glauc Discission of catarac Evisceration of eyebs Enucleation Operation for symbles Removal of cataract Removal of foreign Removal of foreign Removal of foreign Stye abscess Meibomain abscess Tarsal abscess Peritomy	r entro terygic rior ch ody fro a corn- na of co- y for a oma et all pharon senife	ALMIO M/opion im tamber om eye ea ornea glaucoi Mi cornea onjuno	ope Jor. (parti	RATION al tarse	vs.	Total		8 7 2 5 1 5 3 3 4 5 8 1 2 29 83 72 54 49 34 2

ANNUAL REPORT OF THE RADIOLOGIST FOR THE YEAR 1926.

The new X-ray and Electro Therapy Building on the hospital site in Circular Read was completed in the latter half of the year. Although this branch, which has been urgently needed, was asked for on my arrival, in the Federated Malay States in May 1924, and the plans, both for the building and the installation of equipment, were completed at the end of June of the same year, it was not able to take patients until September 20th, 1926.

The reasons for the delay of the building and of its shortcomings have already been dealt with. In the case of the equipment installation, this, although put in as soon as the building was ready—at the end of March 1926—could not be worked until the belated arrival of a motor generator, ordered from the Crown Agents at the end of 1925, but delayed until September by industrial conditions in England. This motor generator was, in my opinion, unnecessary, but had to be bought to comply with the Electricity Enactment.

Since the end of September the new apparatus, both X-ray and Therapeutic, has been working well, although bad material in some of the treatment equipment gave trouble at first, until repaired. All the apparatus, however, is prevented from giving its best by reason of the unsatisfactory electric current supply, which cannot be remedied until the new power station is finished.

The X-ray apparatus purchased in America appears to be well adapted to the climate and the locally built accessories have proved very satisfactory.

The old, war-worn X-ray set at the General Hospital—the only source of X-rays in the Federated Malay States for some years—which had only been kept in working order by constant care, finally broke down completely in August. As the still older apparatus installed at Ipoh burnt out after a few day's use, there were, unfortunately no X-ray facilities at all in the Federated Malay States for nearly a month.

The induction coil type of X-ray machine, being most unsuitable for the tropics, should never have been brought to this country. In the future, a standard type of apparatus, especially adapted to the climate, will be installed in the Federated Malay States Hospitals, so that uniformity of results and reaching facilities may be obtained.

During 1926, 714 X-ray cases were seen, and 209 Electrical treatments given, the latter since September in the new building.

A portable (ward) X-ray set was ordered for the European Hospital, capable of easy removal to other hospitals, but this has recently arrived badly smashed in transit.

With few exceptions, consignments from England have been broken, through careless packing, faulty, badly made or incomplete; have invariably been much delayed, and have often had little or no working or assembling instructions. These faults are rendered more glaring by the fact that all consignments from America have arrived perfectly packed, invariably complete and well made, and always with the fullest possible instructions.

On the treatment side, diathermy apparatus has proved very successful in both medical and surgical cases, and work done in conjunction with other specialists has proved its great value over a very wide field, and quite justifies the purchase of further equipment, which has been ordered.

A new X-ray installation has been put in at Ipoh in the District Hospital, and is working well. There, as in Kuala Lumpur, great delay was caused waiting for a motor generator, and also much accessory equipment had to be done without as the money had been spent on these motor generators. Additional funds have been asked for and the installation will shortly be completed.

As regards staff, an Assistant Surgeon has been attached to the branch in Kuala Lumpur for X-ray and Electrical training. In Ipoh Dr. Greenwood visits the hospital twice a week for the routine X-ray work, while the Radiologist goes to Ipoh at regular intervals for special cases and for any emergency.

(Sgd.) C. F. CONSTANT, Radiologist, F.M.S.

ANNUAL REPORT OF THE MEDICAL OFFICER IN CHARGE, LEPER ASYLUM, KUALA LUMPUR.

Sir-I have the honour to submit the following report for the year 1926, concerning the Leper Asylum, Kuala Lumpur.

NUMBER OF LEPERS TREATED AND NUMBER OF DEATHS.

The total number of lepers treated was 704 with 42 deaths as compared with 635 with 31 deaths during 1925. A fair proportion of these deaths occurred among the elderly inmates some of whom had been within the asylum 15 years or more.

ADMISSIONS.

One hundred and seventy-one cases were admitted, consisting of two Eurasians, 139 Chinese, 29 Tamils and one Malay. A noticeable feature was the large proportion of early cases.

Of the 171 cases admitted, 97 had been diagnosed in some part of the Federated Malay States as suffering from leprosy and had been sent by the hospital which made the diagnosis. Seventy-four patients were self-diagnosed and voluntarily sought admission which was granted after confirmation of the diagnosis.

Of the 97 cases sent to the asylum 56 were sent on transfer from Taiping Leper Asylum. The large proportion of voluntary admissions during 1926, would appear to demonstrate the value of establishing an institution for lepers where it is known to the outside public that leprosy will not only be treated effectively, but in addition, the patient will be able to live a comparatively happy communal life.

TRANSFERRED CASES.

Five insane patients with homicidal mania were transferred to the Central Mental Hospital, Tanjong Rambutan.

ABSCONDING CASES.

Most of these were cases who would not look like lepers to the lay person. The majority returned sooner or later for various reasons. Some were brought back by the police who are notified in each case of absconding. Others returned owing to inability to find occupation. Some Chinese returned because their relatives still regarded them as legally dead, having been lepers; and others because the advantages of living within the Asylum completely outweighed those of living outside.

TREATMENT.

Tai Foong Chee Treatment.—The Tai Foong Chee treatment is being given as a routine method to patients; men, women and children. It is occasionally modified by the addition of bismuth for those who suffer from symptoms of indigestion.

This method of treatment is undoubtedly the most convenient and satisfactory form of routine treatment that could possibly be used for large numbers of lepers of mixed nationalities. It is effective, convenient to prepare and distribute, and has few disadvantages.

VARIATIONS IN RESPONSE TO TREATMENT.

The difference in response to treatment in cases with the same degree of infection and the same type of leprosy is difficult to explain and may depend upon variations in metabolizing the chaulmoogra derivatives within the body as well as other considerations, such as intercurrent infections and resistence to disease in general. Some lepers display a marked idiosyncrasy towards chaulmoogra derivatives and are in the same unfortunate position as is the patient with malaria who is unable to take quinine in any form.

For this type of leper, colloidal antimony injected intramuscularly has been of distinct benefit in arresting the disease and in some cases causing improvement.

E.C.C.O. TREATMENT.

The Ethyl Esters of the oil from Hydnocarpus Wightians combined with camphor, creosote and olive oil have been injected in a large series of more or less stationary cases. Two hundred and sixteen patients applied for a course of injections, treatment being entirely voluntary.

Nationalities were as follows:

TOTTLED 11	CTO WO	TOTTO	113 1				
Chinese				***	 		 141
Tamils			***	***	 	***	 61
Others					 		 14

Men, women and children received injections, ages varying from 5 to 55. The response to treatment depended upon the original severity of the case, the type of leprosy and the already mentioned tolerance of chaulmoogra derivatives.

In all 3,636 injections were given by the Medical Officer in charge. The sites chosen for injection were mostly the upper part of the arm and the outer side of the thigh and injections were given intramuscularly. No case of injection abscess occurred.

Results are shown as follows:

Six patients received 50 injections or more.

Of these five were much improved, one improved.

Sixteen received 40 injections or more.

Of these ten were much improved, five were improved, one remained stationary.

Twenty-one received 30 injections or more.

Of these nine were much improved, ten were improved, one stationary, one worse after 35 injections.

Thirty-six received 25 injections or more.

Of these six were much improved, 18 were improved, 11 stationary, one worse after 26 injections.

One hundred and thirty-seven received 15 injections or over.

Of these one was much improved, 121 improved, 15 were stationary.

Some of the results obtained were striking and most gratifying. The figures given above show that E.C.C.O., if persisted with, is a valuable drug in connection with treatment of the disease and, in addition, emphasises the varied response to treatment.

Some of the cases treated became bacteriologically and clinically negative and are reported as "much improved". The term "cure" is not used as it is agreed that one year at least under repeated bacteriological and clinical observation is required to form an opinion regarding cure.

DRUGS SUBMITTED FOR TRIAL.

Ethyl Esters of the oil from Gorli seed (Onchoba Echinata) were received from the Under Secretary of State, Colonial Office.

The Gorli seed had been supplied by the Commissioner of Lands and Forests, Sierra Leone.

The oil contains as its principal component, a glyceride of chaulmoogric acid and was sent for trial in a series of cases at the Leper Asylum.

After some months of use, 30 patients being injected, it was concluded that the ethyl esters of gorli seed oil were of little or no value in the treatment of leprosy. In addition, changes occurred in the oil rendering it very irritating and unsuitable for injection.

Oil from the Setumpol Tree.—The Setumpol tree is a Malayan species of taraktogenos or hydnocarpus. The oil from the seed approximates in its physical and chemical properties to those of chaulmoogra oil and was submitted for trial.

Injections of the pure oil from the seeds of the Setumpol tree gave rise to pain and local irritation and its use was discontinued. It is thought that ethyl esters of the oil might be more suitable and it is intended to conduct a future investigation as large supplies of the oil are obtainable locally.

Ethyl Esters of Chaulmoogra Oil prepared in Siam.—Both plain and iodised esters were submitted for trial. At first these were satisfactory but later, changes occurred rendering them unsuitable for injection.

Vaccine Treatment.—A large nodule was excised from one of the patients. this was dried in a dessicator, pulverized, ground with sterile sand, mixed with carbol saline and made into an emulsion containing approximately 100 million lepra bacilli per cubic centimetre. Counting of the bacilli was facilitated by mixing them with cocci from a standard emulsion, staining and then performing a differential count. The emulsion of bacilli was sterilized by heating to 60°C for 40 minutes, the sterility being tested subsequently.

This vaccine has been used in two cases of nodular leprosy who showed no improvement with other methods of treatment and who are gradually becoming worse, one being an advanced case and other less advanced.

In the less advanced case there has been a decided improvement. Treatment has been stopped for the time being in the advanced case owing to sudden disintegration of all nodules with distressing secondary infection. The use of the vaccine will be continued after healing of the skin.

Hasson's Vaccine.—During the year Dr. James Hasson of London reported the successful use of a special vaccine prepared by him from serum produced directly from leprous nodules.

The technique is briefly as follows: A leprous nodule or patch is blistered with carbon-dioxide snow and the serum collected from the blister 24 to 36 hours later. The serum and the contained lepra bacilli are incubated at 36°C for three months, the bacilli counted and then mixed in certain proportions with an emulsion containing bacillus pyocyaneus.

Dr. Hasson kindly entered into correspondence with the Medical Officer, Leper Asylum, Kuala Lumpur, and supplied further details as regards preparation of the vaccine.

The chief difficulty encountered was that of obtaining sufficient bacilli in the serum and after a number of trials the preparation of the vaccine was temporarily abandoned.

OTHER METHODS OF TREATMENT.

For the so-called leprous reaction or leprotic fever which is so distressing, hexamine given internally combined with injections of adrenalin has been found of great use in shortening the time of the reaction and in many cases arresting it completely.

Adrenalin is given as the reaction is thought to be of the allergic type depending on the patient becoming sensitized to a certain protein, whether it be a protein product due to autolysis of tissue or a sensitization depending on the presence of the lepra bacillus, it is difficult to say; but a reaction may be readily induced by potassium iodide given by mouth or large doses of E.C.C.O. administered by injection.

Various dye antiseptics are of value in rendering the leprous ulcerations comparatively free from secondary micro-organisms and promoting healing.

DISCHARGES.

At the beginning of the year it was found that there were 20 patients who had been clinically and bacteriologically negative for a year and over.

Of these, five cases who had originally entered the asylum in the early stages of the disease were selected for discharge. Three were Chinese and two Tamils. These were again carefully examined bacteriologically and found negative and after consultation with the Director of Government Laboratories and the Senior Health Officer it was decided to discharge them.

Each of the discharged patients received a small sum of money from the Lepers Aid Fund and departed with much ceremony the Tamils to India and the Chinese to China.

The moral effect on the rest of the inmates was very noticeable and proved a great stimulus both as to hope of recovery and also as regards regularity in taking treatment.

MINOR SURGERY.

A few minor operations were performed on the inmates. An extensive operation of repair was required on one patient who had severed all the structures in front of the wrist with a chopper, the bone being cut in addition. Fortunately there was good healing, an excellent functional result ensuing.

EPIDEMIC DISEASES.

During the year there has been an outbreak of chicken-pox among the children and, in addition, a generalized epidemic of influenza during which very few of the patients escaped infection. In both of these isolation and disinfection of patients and belongings were practiced as far as possible. No deaths attributable to either of these diseases occurred.

ANTI-OPIUM TREATMENT.

Sixteen cases who were opium addicts and who were unable to obtain opium were given a course of anti-opium treatment. In most of the cases, treatment was continued for a period of four to six weeks. One of the cases only reported a lessened craving for the drug.

PATHOLOGY.

Post-mortem examinations were conducted in a number of cases where the cause of death was somewhat uncertain and sections of the various lesions were examined microscopically.

A number of interesting features were noted in advanced cases of mixed leprosy among these being the enormous number of lepra bacilli contained in the spleen, and, in addition, the universal distribution of the bacilli in the nerves of the nerve-anaesthetic types. In one case with clinical involvement of the peroneal nerve only, bacilli were found high up in the sciatic nerve at the level of the gluteal fold. Sections of various specimens were sent to England for study and for teaching purposes.

BUILDINGS AND EXTENSION.

To allow for increased accommodation, the fencing has been extended at one end of the Asylum by 100 feet.

Ten new attap houses each holding 14 patients have been erected by leper labour, all are now occupied and four more are in course of erection.

A dispensary has been built by the lepers and has been stocked with most of the usual hospital drugs and mixtures. This has proved of great value in the treatment of the many minor and major ailments occurring among the 600 inmates, those who are able to attend, coming to the dispensary twice a day.

A small mosque and a special bath have been built for the Malay inmates.

Two Class VIII quarters have been built outside the asylum for the reception of two Perak Government servants who have contracted leprosy.

STAFF.

The Medical Officer in charge visits twice daily and an Assistant Surgeon once. During the year Dr. Eu Khay Hoe resigned from the service and Dr. A. Ponniah replaced him.

The remainder of the staff are lepers.

Mr. Bain, the Steward, has continued his excellent work of previous years and has been quite unsparing of himself in his devotion to the welfare and comfort of the inmates. He is especially commended for the fine work he has performed while labouring under great difficulties.

Mr. Yen Zye Yew, the schoolmaster, twice daily conducts classes in the school house for about 32 children who are receiving an excellent education. Classes in physical drill and Chinese shadow boxing are also held.

Mr. Ah Chai replaced Mr. Ah Wai as head attendant and has been of great assistance in various way including the checking of the formation of various secret societies among the Chinese inmates. In the majority of cases these secret societies are formed for no good purpose and invariably tend to upset the discipline of the Asylum.

AMUSEMENTS.

Both Chinese and Tamil theatrical performances are given at regular intervals. The new Chinese Dramatic Society is very successful and has an efficient organiser in Chan Tong Yen who is an illusionist and conjurer of extraordinary ability.

A new badminton court has been installed and the game is very popular. Ping-pong is also played and tournaments in both games are frequently held.

Athletic sports are held every six months, the programme including most of the usual events such as flat-races, hurdle races, tug-of-war, etc. Entries are received from all nationalities and men, women and children compete in the various events.

A Christmas tree and a party were given to the leper children. Each child received one or more presents. A quaint effect was lent to the ceremony by the children singing well-known Christmas carols with English melodies and Chinese words.

GENERAL.

It is estimated that there are approximately 2,500 to 3,000 lepers in Malaya and of these about 1,500 have entered Leper Colonies. This proportion of lepers segregated, compares most favourably with any tropical country in the world. In India, at a rough estimate, there are nearly one million lepers, 30,000 of these only have been confined to Leper Colonies.

The Federated Malay States Government, through the Medical Department has devoted much attention to the leper problem during the last few years and the new Sungei Buloh Leper Colony with its proposed accommodation for 2,000 lepers promises to be the best and most efficient scheme for the segregation and treatment of lepers, in the world. It should serve as a model for similar institutions in other tropical countries for many years to come.

In many places the distinction between an evil-smelling lazarette and a properly organised Leper Colony is not fully realized. The policy has generally been that of primary segregation and subsequent neglect.

In the Federated Malay States much attention is paid to treatment and conditions of living within the Asylum. The result of this is now being seen in the large proportion of early cases admitted who expect to be cured, and in the large number of patients voluntarily applying for admission.

Previously, life within a lazarette has justly been described as a living death, where neglected and untreated lepers slowly rotted and decayed. Present day conditions afford a striking contrast. The modern Leper Colony is not so much a place of confinement, but a small town which is a haven of refuge. Here are people who have been unfortunate but can still enjoy many of the amenities of their previous existence, and who no longer feel the utter misery and desolation of being both diseased and outcast.

I have the honour to be,
Sir,
Your obedient servant,
(Sgd.) RICHARD GREEN,
Medical Officer in Charge, Leper Asylum,
Kuala Lumpur.

REPORT ON INFANT WELFARE WORK IN KUALA LUMPUR FOR THE YEAR 1926.

INFANT WELFARE CENTRE, SULTAN STREET.

Four years have elapsed since this, the first Centre in the Federated Malay States was opened. The progress made by the work during that period necessitated additions to the building, increase in staff, and during the last year longer working hours, as it was found impossible to cope with the work as arranged for in the original time table.

Before describing what was attempted during the year 1926 it may be of interest to mention the general change that has taken place during these four years in the attitude of the parents towards the aim and object of the Centre. They were originally full of fears and prejudices and attended only after much persuasion and in a very hesitating manner; they vouchsafed no signs of approval or otherwise at the instructions given them and appeared completely mystified and far from happy. To-day on visiting the Centre one can see for oneself that these same people are now thoroughly at home and have lost their fears, their attendance weekly is regular and they bring not only their own babies but their neighbours; they demand instructions saying "please tell me about the feeling"; they occasionally even ask for the instructions to be written in case they forget them. Such change of attitude is due in a large measure to time, helped no doubt by the fact that they have grasped a clearer idea of the aim and object of the work and thus realise their responsibility with regard to the welfare of their children.

THE STAFF.

The staff of the Centre is as follows:

One Lady Medical Officer;

Two European nursing sisters;

Six health visitors;

One laboratory assistant;

One dispenser;

One clerk.

WORK AT THE CENTRE.

Apart from the usual activities of the Centre the following problems had to be faced at the beginning of the year. They are summarised thus:

- Keeping the actual work of the Centre definitely within control, by which
 is meant limiting it unreservedly to the preventative side of medicine;
- (ii) Extending the work among Malays, especially to those living in the distant kampongs;
- (iii) Establishing a satisfactory method of dealing with records for statistical and routine tabulations.

The difficulty mentioned in last year's report of keeping the work to the preventative side of medicine was again experienced and it was recognised that such difficulties would increase unless a definite line of action was taken. To limit such possibility it was decided:

- (i) Getting rid of all the "odd cases" such as old grandmothers, women not mothers, etc., who expected treatment for their various ailments at the clinic;
- (ii) Mothers and babies to replace these "odd cases" and the time previously given in treating them to be utilised in giving more detailed practical instructions to the young mothers.

It was at once obvious that such arrangements would not be popular and for the first-half of the year considerable difficulty was experienced. The old grandmother objected strongly to being sent elsewhere for treatment, although given a chit and provided with transport, whilst the cases requiring hospital treatment were only induced to go there after much persuasion. These people soon realised however that they had to accept the explanation given and that the change of policy was for the benefit of the baby and its mother and in keeping with the object of the Centre. In October the largest number of infants attending the clinic was recorded and there was a considerable reduction in the "odd cases" thus showing a step in the right direction. Such action will however need careful watching as it will be subject to the usual relapses which go hand in hand with any attempt at reconstruction or re-education.

The difficulty in dealing with Malays was the fact that transport had to be provided for those even living within a reasonable walking distance. A Chinese mother is prepared to walk a distance of two or three miles with a baby strapped on her back and leading a small child by the hand and such is so with other nationalities, but the Malay parents must be collected and delivered otherwise one would never see the majority of them.

To deal with work in the distant kampongs two health visitors were given the motor bus three and a half days weekly to bring mothers and babies to the Centre. The kampongs visited are within an area of twelve miles from Kuala Lumpur and the mothers are collected by the health visitors daily at 7,30 a.m. and 1,30 p.m.

This work is under the supervision of the Sister in charge of the district, who visits these places beforehand explaining to the parents the object of the work and telling them the day and hour the bus would call for them. At first difficulty was experienced in getting them to come as some had never left their kampongs before but when a few of the more heroic ones had ventured out and were returned safely the others lost their initial fears and also started attending. It was observed that the mothers were more interested when brought to the Centre to learn mothercraft than when any attempt was made to teach them in their own homes. The struggle to get their attention and the effort to entice these Malays to come to the clinic continued throughout the year but was not without success. The number of attendances for November and December being 781 and 775, respectively, against the figures for the corresponding months of 1925 of 177 and 197.

The list of Malay kampongs visited is attached.

It was found necessary to start a card index system in March as up to that time there was no very satisfactory method for keeping records required for annual reports and official returns. The Chinese surnames were the first difficulty as in order to get any accuracy and uniformity in the spelling of these it was necessary to obtain a list of Chinese surnames (commonly called "Seh") as pronounced in different dialects. and so become familiar with the commoner ones. Nearly all Chinese are able to recognise the Chinese character for their "Seh" and the ones who are not, are asked to find out from their husband or relations and to bring the information on their next visit. It was troublesome at the start and took up a considerable amount of time and patience but the consolation remained that a useful and accurate record of every mother and infant attending the clinic was then in existence.

The next obstacle to overcome was to impress on the parents the value of the reference card given to them containing the name, number, etc., of the mother and infant. In spite of the fact that its object was carefully explained to them and on the card itself was written in four languages "Please bring this card with you next visit," they invariably came without it. The chief excuses being either that the baby chewed or tore it, that they left it in a friend's house, or lost it in the bus. One mother having come twice in succession without it and not daring to appear a third time with the excuse that she couldn't find it promptly borrowed her neighbour's and triumphantly arriving at the Centre handed it in; this of course would have led to all sorts of confusion if she had not been recognised by the health visitor who knew she did not posses an infant of the name and age as indicated on the card.

In addition to the usual instruction given at the Centre an attempt was also made during the year to include some simple teaching in elementry hygiene to a few of the bigger children who came with their parents. Simple explanation of the causation of malaria, plague and ankylostomiasis were given. Posters of the mosquito, the rat, and of good children wearing shoes, to prevent hook worm infection were drawn up and proved very helpful. These simple lectures as stated were only intended for children, but they also attracted the father and mother who listened attentively and were quite interested in the childish explanation of these diseases given to the younger members of their family. To help the poor mother who could not afford to buy shoes for her children, a paper pattern was given her from which a useful and serviceable pair of shoes could be made with very little trouble from an old felt hat. Other improvised garments were also shown her such as a child's vest made from the father's old one, and child's trousers made from sleeves of a mother's baju, etc.

Owing to the large percentage of infants suffering from marasmus or malnutrition seen at the clinic it occurred to one that a great deal more illness was due to defective diets than was at first recognised and an effort was therefore made to provide such infants with a vitamin-rich diet-The result justified the contention that many cases of marasmus were due to vitamin deficiency. The food of these infants which was milk in simple dilution was enriched by the addition of a mixture of cod-liver oil, oil emulsion, marmite and orange juice in order to make sure of adequate supplies of Vitamines A, B, C and D, respectively. The dosage starting with half drachm was gradually increased to two drachms and must not exceed that amount at age two to three years. It was feared the emulsion not being a very palatable one, would be disliked by the children but such was not the case, its fishy taste making it like the Chinese sauce which is used extensively in the preparation of food by most native mothers. The vast majority of babies improved rapidly on the mixture and it became very popular. Such simple dietetic measures would not only prevent and diminish the amount of marasmus and malnutrition but should produce teeth better formed and less liable to decay-a very important factor here where at least 90 per cent. of the poorer children suffer from dental caries.

Another factor of importance that emphasized itself during the year was the discovery of a large number of infants, the majority under six months, found to be suffering from malaria. These babies generally came from distant kampongs and were brought because of their extreme palor, and the fever from which they occassionally suffered; the mothers on being questioned said that the babies took their food moderately well and did not appear at any time to be what they considered really ill. They were merely worried because of their paleness and the fact that they were not as bright as other children of the same age. On examination they had a marked degree of anaemia, a palpable spleen and their blood was full of parasites, they however improved with treatment and the mothers were carefully instructed on the cause and prevention.

The point to note about such cases is that these babies would have been left undiscovered and untreated if the kampongs were not visited by the sisters and health visitors and the mothers brought to the Centre. Many infants must have died in these distant places and the ones who survive are probably the anaemic specimens with enlarged spleens and a history of fever since babyhood so common at a school inspection. These children the school teacher will tell you are very often absent owing to fever, that they are always tired and very slow to learn. One does not know if there are any statistics to show the number of infants, say under three and under six months, found to be suffering from malaria, but the matter is worth investigation as the numbers found in the clinic during the year were alarming.

Weighing the baby has become quite popular and none of the difficulties experienced in the first years were met with, the mothers always asking the weight-gain in order that they could tell the father and grandmother as they too were very interested and were always anxious to know the progress made by the baby.

ANTE-NATAL WORK.

There is still some difficulty in getting the poorer mothers to hospital for confinement. They will explain carefully that it only means leaving a large number of young children at home with no one to look after them, the father being at work all day—and the neighbours themselves already having many of their own to supervise. It means also that the mother only worries while in hospital and is not happy until she is at home again.

They attend the clinic regularly during the ante-natal period for the routine examinations and supervisions, and if they are unable to do so the last month or two they are visited by the sister and health visitor and arrangements are made for them to have their babies at home with as much comfort and safety as is possible under the circumstances.

An obstetrical packet is given to every poor expectant mother who needs it as it helps to carry her through the first few days after her confinement, many of them living as they do in great poverty and being unable to get more than a few rags together for the event. The discovery of cases of "post-puerperal beri beri" was mentioned in last year's report and as a preventative measure treatment with Vitamin B extract prepared at the Institute Medical Research was given with good results to many ante-natal mothers during the year. The occurrence of beri beri in a woman after childbirth was commented upon by all the earlier writers on that

disease. Thus Bently (1893) in discussing the predisposing cause of beri beri says "It is rarer among women but I have seen it attack them shortly after childbirth, when it makes its appearance during the second week before the disappearance of the lochia. The most remarkable instance of this occurred in the family of a native (Malayan) prince, where three of his relations, all young women in the prime of life, were at intervals of about a year each, attacked with this disease-in one of them proved fatal after a duration of six months from the onset of acute symptoms. A very peculiar fact, and one which I am quite unable to account for, is the susceptibility of women after childbirth to be attacked with beri beri. Is it that they are more liable to any acute infectious disease as they undoubtedly are, for instance scarlet fever and other exanthemata?"

In later studies on the causation of beri beri in relation to diet and especially in the development of the order of ideas associated with the term "Vitamines" it came to be believed that the incidence of the disease is highest, other things being equal, among those upon whose metabolic processes the greatest demands are made. This is the theory now commonly held to account for the relatively high incidence of beri beri in women during the period of puerperium. With the present knowledge of the causation of beri beri success has been met with in this country in reducing nearly to the vanishing point the incidence of the disease among groups of people formerly decimated by it, such as prisoners in gaol and workers in mines, but it appears that this knowledge has not been given its due effect in the control of the disease among native women after childbirth. Owing to its frequent occurrence and very high mortality rate post-puerperal beri beri is a disease that demands renewed attention and every effort should therefore be made to obtain suggestions for its more efficient control.

VENEREAL DISEASE.

The number of cases of babies suffering from "Heredo-Syphilis" is still large, probably because the mothers themselves refused to undergo any treatment. They maintain that as they feel quite well they cannot in any way be responsible for the condition of the baby and are therefore loathe to undergo any curative treatment.

RECORD OF ATTENDANCES.

			INFANT	WELF	ARE CENTE	RE.			
	1926.		Infants.		Children.		Women.		Total.
Janua	ry		 1,623		316		287		2,226
Febru	ary		 1,317		346		275		1,938
March	1		 1,779	***	538		302		2,619
April			 1,383		329		245		1,957
May			 1,909		429		372		2,710
June			 1,875		469		399	***	2,743
July			 1,987		507		419		2,913
Augus	t		 1,382		301		276		1,959
Septer	mber		 1,594		364		371		2,329
Octob	er		 2,042		360		472		2,874
Noven	nber		 1,979		361		506		2,846
Decen	nber		 1,922		323		472		2,717
					-				
	Totals	1926	 20,792	***	4,643	***	4,396	***	29,831
	Totals	1925	 16,005		4,259	***	2,870	***	23,134
	Totals	1924	 9,106	***	3,193	***	4,039		16,238
	Totals	1928	 5,777		2,872		3,559		12,208

142

RECORD OF ATTENDENCES B	V NATIONALITIES
-------------------------	-----------------


1926.	Chinese.		Tamils.		Malays.		Others.		Total.
January	 1,469		418		206		133		2,226
February	 1,246		341	***	289		112		1,938
March	 1,880		341	***	288	15.000.00	115	5000	2,619
April	 1,363		357		162	200	75		1,957
May	 1,896	***	403		305		106		2,710
June	 1,888		417		324		114		2,748
July	 1,881		523		401	***	108		2,913
August	 1,302		354	***	203		100		1,959
September	 1,470		440		308		111		2,329
October	 1,870	***	428		477		99		2,874
November	 1,666		321		781		78	000	2,846
December	 1,633		275		775		34		2,717
Totals 1926	 19,564		4,618		4,464		1,185		29,831
Totals 1925	 15,331		4,095		2,855		1,353		23,134
Totals 1924	 11,300		3,369		582		987		16,238
					Gra	and T	otal 1926		29,831

The following graphs are attached:

- (1) Malay attendances from July, 1922—December, 1926;
- (2) Ante-natal attendances from July, 1922—December, 1926;
- Attendances of infants, children and mothers from July, 1922—December, 1926;
- (4) Attendances of infants, children and mothers for the year 1926.




INFANT WELFARE CENTRE, KUALA LUMPUR MALAY ATTENDANCES FROM JULY 1922 TO DECEMBER 1926.






KUALA LUMPUR INFANT WELFARE CENTRE ANTE-NATAL ATTENDANCES FROM JULY 1922 TO DECEMBER 1926.






KUALA LUMPUR INFANT WELFARE CENTRE ATTENDANCES FROM JULY 1922 TO DECEMBER 1926.





ATTENDANCES AT THE INFANT WELFARE CENTRE, KUALA LUMPUR 1926.





HEALTH VISITORS IN THE HOMES.

As stated this has been extended to Malays living in kampongs within a twelve-mile radius of Kuala Lumpur, and the difficulty in collecting the Malay mothers and their babies and getting them to the Centre can only be appreciated by those who have tried it. They are never ready when the bus calls for them and the health visitor very often has to dress the mother and infant and lead them through the kampongs to the car waiting on the main road. They seem to have no idea of time and never by any chance know the baby's correct age; for instance the mother will tell you her infant is two months old at its first visit to the Centre and at the second visit, which may be a fortnight later, its age has suddenly jumped to six months. They are pathetically helpless regarding themselves and hopelessly so regarding the welfare of their children but they are anxious to learn and quite appreciative of the fact that they are now getting a great deal of attention.

Having got them to the clinic for the first time they are taught one or two elementary facts which however are forgotten by the time the bus is due to take them home, however reiteration on their subsequent visits eventually produces results and after their tenth visit or so they remember most of the facts taught them. Their intelligence varies with the district they come from, thus the Gombak people about 12 miles from Kuala Lumpur are quite bright, the kampongs in that area being near the road which gives them a better chance of seeing and mixing with other people. They are a striking contrast to the Malays that come from Sungei Penchala, $7\frac{1}{2}$ miles from Kuala Lumpur, where the kampongs are 3 or 4 miles from the main road and where they have not the same opportunity of meeting people with the result that they are dull, nervous and slow. At the clinic they are provided with milk whilst waiting, special cases being given feeding bottles, patent food, vitamin emulsion, etc., and the health visitor sees them safely back to their homes. There is no difficulty with the Chinese now, they attend regularly and have grasped quite a fair knowledge of the work. They come from all parts and are very interested. The Tamils are slow, but attend regularly and definite improvement is noticeable.

District visiting is recognised as the most important part of child welfare service and the work done here by the staff has been exceptionally good throughout the year.

LIST OF MALAY KAMPONGS FOR DISTRICT VISITING WITHIN TEN MILES RADIUS FROM KUALA LUMPUR.

1. Segambut	3½ mile turning off Batu Road. Most Malays are congregated near the Malay school.
2. Batu Village	Five miles, Kepong Road. Malay centre about 100 yards from the road junction.
3. Selayang	63 mile, Batu Caves Road. Road centre for Selayang Malay Reservation in the reservation are two very thickly populated Malay kampongs about 1 mile from 63 miles and 8th mile, Rawang Road.
4. Pantai	A turning off 4½ miles, Petaling Road.
5. Petaling Station	5½ miles, a turning off New Petaling Road. This is a convenient place for Malays on the Klang River and also from Pantai.
6. Ampang Village	Five miles. The Malay centre is near the mosque 100 yards before the entrance to the village.
7. Ulu Klang	7½ miles via 4½ miles village, Ampang Road. The Malay are mostly at the northern end of the village.
8. Sungei Penchala Village	7½ miles, Old Damansara Road. This is the meeting place for Malays from the Sungei Penchala Malay Reservation to the north and small holdings to the south.
9. Gombak	4th to 10th mile.

		HEALTH	VISIT	ORS' V	VISITS	TO HO	OMES.		
	1926.								
	January						THE PARTY OF		1,763
	February				2				1,179
	March		***	***		***		***	1,607
	April	ani sal te			1.1 00		100		1,176
	May	valebby (c	***				***	***	1,435
	June	1.000							1,328
	July			***					1,294
	August		***		***	***			1,033
	September	d will and							798
	October								1,494
	November	ollprdz					111 4	***	1,060
	December		***	***			***		1,118
									-
							Total		15,285
					Grand	T-4-1	1000		15 005
									15,285
					Grand				13,221
					Grand	Total	1924		10,532
	Mothers and	babies con	veved	by tra	nsport				9,833
									124 3252 1111
					nul St				
M.	ATERNITY ANI	O CHILD Natu		ARE W	t out St		DURI	NG T	HE YEAR.
M	ATERNITY ANI	O CHILD Natu	WELF/	ARE Work.	t out St		DURI	NG T	
M	Number of cl	O CHILD Natu	WELF/	ARE Work.	VORK I		DURI	NG T	Total. 586
M	Number of cl Number of me Ante-natal	O CHILD Natu	WELF/	ARE Work.	VORK I		DURI	NG T	Total. 586
M	Number of cl Number of me Ante-natal Post-natal	Natu linics held others seen	WELFA	ARE Work.	VORK 1	 		NG T	Total. 536 398 3,998
M	Number of cl Number of me Ante-natal Post-natal Number of ba	Natu inics held others seen	welf/	ARE Work octor:	VORK I	OONE			Total. 586 398 3,998 4,419
Mal	Number of cl Number of me Ante-natal Post-natal Number of ba	Naturalinics held others seen	welf?	ARE Work octor: registe	VORK I	 		and least	Total. 536 398 3,998 4,419 13,424
M. M.	Number of cl Number of me Ante-natal Post-natal Number of ba Number of ba	Naturalinics held others seen bies entered bies weigh	welf/ re of wo by do ed on to	ARE Work octor: registe	vork i	OONE			Total. 586 398 3,998 4,419 13,424 3,248
M	Number of cl Number of me Ante-natal Post-natal Number of ba Number of ba Children norma	Naturalinics held others seen bies entere bies seen labies weigh al—general	welf?	octor: registe	vork i	OONE			Total. 536 398 3,998 4,419 13,424 3,248 2,527
M	Number of cl Number of me Ante-natal Post-natal Number of ba Number of ba Number of ba Children norms	Naturalinics held others seen bies entere bies seen bies weigh al—general incorrect	welf/ re of wo by do ed on by doo ned l advice	octor: registe	vork i	OONE			Total. 586 398 3,998 4,419 13,424 3,248 2,527 2,630
M. M	Number of cl Number of me Ante-natal Post-natal Number of ba Number of ba Number of ba Children norma Suffering from	Naturalinics held others seen the seen	welf? re of wo by do by doo hed advice feedin	octor: registe	vork i	OONE g year			Total. 536 398 3,998 4,419 13,424 3,248 2,527 2,630 6,355
M	Number of cl Number of me Ante-natal Post-natal Number of ba Number of ba Number of ba Children norm Suffering from Suffering from	bies entered bies weigh al—general incorrect worms	welf/ re of wo by do ed on ladvic feedin mothe	octor: registe	vork i	OONE g year			Total. 536 398 3,998 4,419 13,424 3,248 2,527 2,630 6,355 1,736
M. M	Number of cl Number of me Ante-natal Post-natal Number of ba Number of ba Number of ba Children norma Suffering from	bies entered bies weigh al—general incorrect worms	welf/ re of wo by do ed on ladvic feedin mothe	octor: registe	vork i	OONE g year		3223	Total. 536 398 3,998 4,419 13,424 3,248 2,527 2,630 6,355
M	Number of cl Number of me Ante-natal Post-natal Number of ba Number of ba Number of ba Children norm Suffering from Suffering from	bies entered bies weigh al—general incorrect worms	welf/ re of wo by do ed on ladvic feedin mothe	octor: registe	vork i	oone g year 			Total. 536 398 3,998 4,419 13,424 3,248 2,527 2,630 6,355 1,736
M. M	Number of cl Number of me Ante-natal Post-natal Number of ba Number of ba Number of ba Children norm Suffering from Suffering from	Naturalinics held others seen the seen	welfare of wo	octor: registe etor e giver	vork i	oone g year en			Total. 536 398 3,998 4,419 13,424 3,248 2,527 2,630 6,355 1,736
M. M	Number of cl Number of me Ante-natal Post-natal Number of ba Number of ba Number of ba Children norm Suffering from Suffering from	Naturalinics held others seen the seen	welfare of wo	ARE Work octor: registe extor e given	vork I	oone g year en			Total. 536 398 3,998 4,419 13,424 3,248 2,527 2,630 6,355 1,736 261
M. M	Number of cl Number of me Ante-natal Post-natal Number of ba Number of ba Children norms Suffering from Suffering from Suffering from Referred to h	bies entered bies seen bies weigh al—general moorrect worms malaria, cospitals	welf Are of wo	octor: registe etor e giver	or during	oone g year en			Total. 536 398 3,998 4,419 13,424 3,248 2,527 2,630 6,355 1,736 261
M. M	Number of cl Number of me Ante-natal Post-natal Number of ba Number of ba Number of ba Children norm: Suffering from Suffering from Suffering from Referred to h	bies entered bies seen bies weigh al—general worms malaria, cospitals VISIT mothers	welf Are of wo	ork octor: registe etor e given	vork I	oone g year en			Total. 536 398 3,998 4,419 13,424 3,248 2,527 2,630 6,355 1,736 261
M. M	Number of cl Number of me Ante-natal Post-natal Number of ba Number of ba Number of ba Children norms Suffering from Suffering from Suffering from Referred to h	bies entered bies seen bies weigh al—general incorrect worms malaria, cospitals VISI mothers ler one yeared one-five	welf Are of words a by document of the distribution of the distrib	octor: registe ctor e giver	or during	oone g year en			Total. 536 398 3,998 4,419 13,424 3,248 2,527 2,630 6,355 1,736 261
M. M	Number of cl Number of me Ante-natal Post-natal Number of ba Number of ba Number of ba Children norm: Suffering from Suffering from Suffering from Referred to h	bies entered bies seen bies weigh al—general incorrect worms malaria, cospitals VISIT mothers ler one year aid by sist	welf Are of wo	octor: registe etor e given	vork I	oone g year en		and a second and a second a se	Total. 536 398 3,998 4,419 13,424 3,248 2,527 2,630 6,355 1,736 261 686 8,542 6,057
M. M	Number of cl Number of me Ante-natal Post-natal Number of ba Number of ba Number of ba Children norm Suffering from Suffering from Suffering from Referred to h	bies entered bies seen bies weigh al—general incorrect worms malaria, cospitals VISI mothers ler one yeard one-five aid by sist aid by nur	welf Are of wo a by do ed on ed ladvice feedin mothe ar years ters	are work octor: registe ctor e giver	or during	oone g year en			Total. 536 398 3,998 4,419 13,424 3,248 2,527 2,630 6,355 1,736 261 686 8,542 6,057 4,115 15,285

EDUCATION AND PROPAGANDA.

The fourth infant welfare exhibit and the third baby competition took place at the Malayan Agri-Horticultural Exhibition held in Kuala Lumpur in August.

The Exhibit was prepared by the staff of the Infant Welfare Centre on the lines of previous years with the following additions:

- (1) A vitamin section;
- (2) Elementary hygiene section;
- (3) Practical demonstrations in
 - (A) Feeding of infants including the following:
 - (i) Preparation of cow's milk;
 - (ii) Condensed milk;
 - (iii) Patent food;
 - (iv) Thin cereal food for an infant of about seven months;
 - (B) Bone and vegetable soup;
 - (C) Albumen water;
 - (D) Raw meat juice;
 - (E) Fruit juice.

The demonstrations were held every morning from 12-1 and every afternoon from 4-6 and the lectures accompanying them being given in Malay, Chinese and Tamil by the health visitors. The preparation of food, etc., came in for a lot of attention, the cooking being done on charcoal and oil stoves in a fashion possible to all the lookers-on.

INFANT WELFARE FILM.

This proved by far the most useful form of propaganda. It was arranged by the Secretary of the Infant Welfare Advisory Board and shown at the local Cinemas. It depicts the work as daily carried out at the Centre and in the district.

BABY COMPETITION.

The total entries for this numbered 275; the figures for Malays being nearly double that of other years.

A thousand dollars was handed in at the Centre for prize money for the baby show which facts fully demonstrates not only its popularity but the appreciation of all nationalities to the Government's generous effort made on behalf of the welfare of their children.

Miss Tan Kim Eng, health visitor, joined the staff in May, 1926.

M. JOSEPHINE WERE,

Lady Medical Officer, Infant Welfare Centre,

Kuala Lumpur.

PREMISES.

On 1st May, the Welfare Centre was removed from the shop-houses in Treacher Street to the Yau Tet Shin building in Club Road. This building, though by no means ideal for a Welfare Centre is a great improvement on the previous premises. It consists of two wings—one being used for the Town Dispensary and dresser's quarters—the other having the Health Visitors quarters on the first floor with waiting, consultation, examination and dressing rooms on the ground floor. These latter rooms are well ventilated, well lighted and spacious but in spite of all efforts have not that appearance of scrupulous cleanliness possible in a building specially built for a Welfare Centre.

STAFF.

One part time Lady Medical Officer.
One European Nursing Sister.
Two Health Visitors.

One Amah.

The dispensing is done by the dresser-in-charge of the Town Dispensary.

On December 4th, Miss E. M. Goulding, European Nursing Sister, commenced her long leave. Her place has been filled by Miss M. Grice. On May 1st, Mrs. M. G. Paul began duty as a Health Visitor. Neither Health Visitor has the necessary qualifications for the post. Nurses with the necessary qualifications are very hard to get as not only must they have completed their hospital training and be certified midwives but also they must be able to speak English, Malay, Chinese or Tamil.

WORK AT THE CENTRE.

The centre still continues to be more an out-patient department for the treatment of disease than an institution for its prevention. The work is by no means confined to maternal and child welfare. Any woman wishing to consult a Lady Medical Officer must either be admitted to hospital or wait for one special afternoon set aside each week for examination at the hospital of gynaecological patients, or come to the centre. Many women refuse admission to hospital and cannot be turned away to wait for perhaps another week for medical attention; so the only alternative is to treat them at the centre. The teaching at the centre is mostly individual. So far there have been no demonstrations on a large scale. Each mother as she presents herself or child for examination is instructed about infant feeding and general hygiene. Not many women bring their babies for advice, if the latter are perfectly well, unless they have been previously so advised by the Health Visitors. The same may be said of ante-natal cases. There is usually some condition other than pregnancy for which they come to seek treatment. All cases of suspected venereal disease are treated at the District Hospital. Simple chemical examination of urine is done at the centre but any specimens of faeces blood or urine must be sent to the District Hospital for microscopical examination.

RECORD OF ATTENDANCES-INFANT WELFARE CENTRE.

ALLE CO.	er or	 ******	C. Auto	****				
1926.		Infants.		Children.		Women.		Total.
January	***	 604		286	***	411		1,301
February		 443		204		287		934
March		 617		307	***	385		1,309
April		 458		207		318		983
May		 662		275		366	***	1,303
June		 446		197		361		1,004
July		 627		299		391	***	1,317
August		 568		276		403		1,247
September		 582		264		443		1,289
October		 544		244		359		1,147
November		 597		205		336	***	1,138
December		 135		617		356		1,108
Tota	1 1926	 6,283		3,381		4,416		14,080
Tota	1 1925	 7,215		3,585		4,773		15,523
						-		

HEALTH VISITING IN THE HOMES.

During the year, this the most important part of maternal and child welfare work, has progressed considerably. The appointment of a second Health Visitor has enabled the visiting of homes to be greatly extended. Each week a list of birth notifications is sent by the Police Department and births within a radius of approximately 3½ miles are entered in the welfare birth register. It is impossible to extend the work beyond this area whilst the staff consists of only one European Nursing Sister and two Health Visitors whose only means of conveyance is by rickshaw. Chinese are visited by the Chinese Health Visitor, Malays and Tamils by the Tamil Health Visitor. Each visitor spends the whole of alternate days visiting homes. The intermediate days she assists at the centre except for 1½ hours in the morning when in company with the European Nursing Sister she visits particularly newly born babies. One and half hours each morning is the most the Sister-in-charge can be spared from the centre. It is by visiting the homes that most of the preventive work is done. Babies are seen a few days after birth. Mothers are instructed regarding infant feeding and general hygiene. Children under 12 are inspected and sent for medical attention if necessary. Puerperal women if ill are advised to enter hospital. If this advice is not taken the Lady Medical Officer is informed and visits the patient in question. The instruction in the Malay language received by the previous Sister-in-charge has proved invaluable to her in her work in the homes.

				HEALT	TH VIS	SITS.				
January	7	***	***		***			***	***	143
Februar	у									126
March										175
April			***		***					117
May	***	***								419
June										487
July										532
August		***								534
Septem	ber									760
October										520
Novemb	er				***					742
Decemb	er			***		***		,		1,342
						Grand	Total	1926		5,897
						Grand	Total	1925		1,309

MOTHERS AND BABIES BROUGHT BY TRANSPORT.

Nil.

INFANTILE MORTALITY RATE-IPOH

		Tren	THITTE	DIL.C.	TA TATAL T T	****			
Year					Births.		Deaths.	R	late per millie.
1915					641	***	122		190.00
1916	***	***			725		126		173.79
1917					881		168		190.00
1918					843		173		205.22
1919					792		137	***	173.00
1920			***		866		187		158.19
1921					898		155		172.61
1922					974		152		156.05
1923				***	904		119		131.64
1924			***		1,072		126		117.54
1925					844		120		142.18
1926					1,364		181		96.04

MATERNAL AND CHILD WELFARE WORK	DONE DUE	RING ?	THE	YEAR.
Nature of work.				Total.
Number of clinics held				543
Number of mothers seen by doctor ante-r	natal .			154
Number of mothers seen by doctor post-r	natal			200
Number of babies on birth register durin	g year .			928
Number of babies under one year seen b	y doctor .		2	,201
Number of babies weighed				613
Children normal general advice given				493
Suffering from incorrect feeding				175
Suffering from worms			4	,421
Suffering from malaria				262
Referred to hospital				108
s of leading startly because of heregonite was				
VISITS BY NURSE	IS.			
To expectant mothers				146
To infants under one year			3	,999
To children 1-5 years	***		1	,752
Total visits by Sister			1	,324
Total visits by Nurses			5	,897
Number of mothers and babies brought by	transport .			Nil
Visits are paid once a week to the Gopeng				
Matal and took 1	and the same			789
Total cases treated	***			100

Iрон, 25th January, 1927. M. HEWITSON,

Lady Medical Officer,

Infant Welfare Centre, Ipoh.



ANNUAL REPORT ON INFANT WELFARE CENTRE AND CLINIC FOR MOTHERS, TAIPING, FOR THE YEAR 1926.

STAFF.

One Lady Medical Officer (part time).

One European Nursing Sister.

Two Health Visitors (during nine months, one only has been available).

One Amah.

One punkah-puller.

One peon.

The dresser-in-charge of the Town Dispensary, assisted by an attendant, does the dispensing. Microscopical and chemical examinations are made at the General Hospital Laboratory.

PREMISES.

These consist of a shop-house. Besides a small entrance space and passage there are three rooms.

 A small cubicle, downstairs, where minor dressings are done.
 A large room, up steep and narrow stairs, used for waiting, general and exhibition room as well as my office.

III. A small room, used for examination of patients when privacy is indicated.

THE WORK.

The welfare work is beginning to show definite signs of progress.

The number of infant attendances at the weight-clinic is over 800 more than in 1925.

Babies weighed ... 4,768

1925 -

Babies weighed ... 3,946

The parents are bringing their children much more willingly and without such frequent remindings by the Health Visitor-they show quite an eager interest in the progress of their babies' weight and seek our advice on various matters of infant upbringing.

ANTE-NATAL CLINIC.

This most important branch of our work is proving almost the most difficult to establish. There is the usual trouble of trying to induce the prospective mother who thinks herself perfectly well, to make sure not only that she really is so, but that she remains so.

Prospective mothers frequently come up with some small complaint connected with their pregnancy, but unless the women give some promise of continued attendance, they are not included in the ante-natal returns. The number of cases which have been kept under observation for several months before confinement and then visited afterwards by the Health Visitor is small.

MIDWIVES.

There are six certified midwives practising in Taiping but they complain that much of the work that they should get is done by untrained bedans.

HEALTH VISITING.

A great deal of time is given to the instruction of the parents in our methods and routine, this in their homes as well as at the centre-but results come very slowly. However in a good number of cases we have induced mothers to breast feed their babies where their previous children had been brought up on the bottle.

It is difficult to instil our ideals of cleanliness, scientific feeding, clothing and general upbringing of children into a poor coolie whose home is a small inadequate airless room, in which the furnishing and equipment is almost negligible. The most constantly occurring obstacles to progress are:

I. The long rubber tube and the "dummy".

II. Sweetmeats.

III. As far as their structure permits, living rooms in which all possible ventilation is prevented—especially is this condition met with in post-natal cases.

Now that a second Health Visitor has been appointed to Taiping it is becoming possible to cope more satisfactorily with the visiting in the homes and thus, besides keeping the children under more strict observation, maintain the interest of the parents in the Infant Welfare Work.

Consultation at Port Weld are being held twice a month and continue to be well attended. It will be more satisfactory when these cases can be brought in to the centre, as will be possible when a motor-bus is obtained.

EXHIBITION.

An elementary but comprehensive welfare exhibition was held at the Agri-Horticultural Show in Taiping on August 14th and 15th.

The show-room was constantly crowded, chiefly with Malays, all very interested and anxious to be instructed. Many of the Malays would have liked their families to attend the Welfare Centre regularly but the distance to their homes and the expense of transport is too great for them to do so.

Of the exhibits shown, some were kindly lent by the Ipoh Welfare Centre and others were obtained locally. Several of the Nurses from the General Hospital gave up their spare time to help at the exhibition and it was thus possible to give a continuous series of little lectures, demonstrations and explanations in all the necessary languages.

CHANGES IN STAFF.

Mrs. Bentinck left on transfer on June 16th.

I assumed duty on June 16th.

An untrained Health Visitor Chan Cheng Swah left on March 23rd.

Nurse Pung Seh Moy commenced duty as Health Visitor on November 1st.

EVELYN B. JACQUES, Lady Medical Officer, Infant Welfare Centre, Taiping.

RECORD OF ATTENDANCES. INFANT WELFARE CENTRE, TAIPING.

1926.	Infants of 5 year	Childre 5-14 yea	Women		Total.	Health visiting total.		Weight.	881	ases at to pital.
January	 845	 162	 178		1,185	 461	***	67		18
February	 610	 194	 180		984	 442		44		5
March	 919	 216	 206	***	1,341	 584		47		7
April	 581	 275	 164		1,020	 217		28		2
May	 684	 330	 164		1,178	 250		32		7
June	 810	 287	 202		1,299	 231		32		3
July	 804	 122	 223		1,149	 299		_		-
August	 700	 148	 170		1,018	 272		-		_
September	 751	 103	 240		1,094	 310		-		-
October	 681	 113	 180		974	 306				_
November	 785	 106	 182		1,073	 509				-
December	 876	 145	 223		1,244	 510		-		-
Total	 9,046	 2,201	 2,812		13,559	 4,391		250		42
						Total		2	92	0.10

Grand Total of work ... 18,242

			C		NATIO	NALITIES.			
1926.		Chinese.		Tamils.		Malays.		Others.	Total.
January	***	984		136		437		111	 1,668
February		897	***	381		99		27	 1,404
March		1,263		471		133		75	 1,942
April		917		403		99		27	 1,446
May		1,044		355		137		37	 1,573
June		949		327		134		41	 1,451
July		886		436		99		27	 1,448
August		863		314		87		26	 1,290
Septembe	r	885	****	339		144		36	 1,404
October		839		280		124		87	 1,280
November		1,051		401		105		25	 1,582
December		1,045		437		213	2000	59	 1,754
Total		11,623	0000	4,280		1,811		528	 18,242

(1) Ante-natal cl	inic					 	 . 135
(2) Babies on reg	gister					 	 520
(3) Babies weigh	ed-						
January			***			 ***	 504
February						 	 290
March	***		***			 	 507
April				***		 	 338
May				***		 	 266
June						 	 367
July						 	 465
August						 	 351
September						 	 405
October			***			 	 414
November		***				 	 468
December		***				 	 393
						Total	 4,768
(4) Health Visiti	ng—						
Primary vi	isits	***				 	 754
Subsequen	t visi	ts to	hildren	under	1 year	 	 2,718
Visits to c	hildre	en age	1 1-5			 	 791
Visits to a	inte-n	natal cases			***	 	 128
						Total	 4,391

August Marine San Register San

PRINTED AT THE FEDERATED MALAY STATES GOVERNMENT PRESS, EUALA LUMPUR.

