Obstetric department clinical report : 1947 / Camborne-Redruth Miners' & General Hospital.

Contributors

Camborne-Redruth Miners' & General Hospital. West Cornwall Hospital Management Committee.

Publication/Creation

[Place of publication not identified]: [publisher not identified] 1947

Persistent URL

https://wellcomecollection.org/works/spkepjwa

License and attribution

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



CLINICAL REPORT

OF THE

OBSTETRIC DEPARTMENT


0

Camborne - Redruth

Miners' & General Hospital

Redruth

FOR THE YEAR 1947





CONTENTS

				Page
Abortions			 	13
Accidental Antepartum Haemorrhage			 	20
Antenatal Admissions	(cinoxa.re)		 	4
Antepartum Haemorrhage		***	 	18
Antepartum Haemorrhage of Unknown	n Origin		 	21
Blood Transfusion			 	52
Breech Deliveries—complicated			 	34
Breech Deliveries—uncomplicated			 	33
Caesarean Section			 	36
Eclampsia			 	48
External Version			 	23
Face Deliveries			 	45
Fibroids and Pregnancy			 	15
Forceps Deliveries	din't		 	40
Heart Disease			 	16
Hydramnios			 	46
Induction of Labour			 	25
Induction of Labour for Other Causes			 	30
Induction of Labour for Pre-Eclampt	tic Toxae	emia	 	26
Internal Version			 	46
Manual Removal of Placenta			 	51
Maternal Deaths			 	60
Methods of Delivery			 	31
Morbidity of Patients			 	56
Neonatal Deaths			 ***	69
Ophthalmia Neonatorum			 	71
Ovarian Cyst and Pregnancy			 	15
Perineal Tears			 	55
Placenta Praevia			 	18
Postnatal Admissions			 	13
Postpartum Haemorrhage	17.00 NO		 	50
Premature Babies			 	61
Previous Caesarean Sections			 	38
Prolapse of Cord			 	47
Pulmonary Tuberculosis			 	17
Stillbirths			 	66
Syphilis in Pregnancy	4	***	 	15
Transverse Lie	***		 	45
Twins			 	. 44
-			-	-

ABBREVIATIONS

Alb	 Albumin.	L	 Living.
A.R.M.	 Artificial rupture of	N.L.	 Normal Labour.
	the membranes:	P.P.H.	 Postpartum
B.P	 Blood pressure.		Haemorrhage.
	Caesarean Section.		Breech Presentation.
	Dead or Died.		Stillborn.
	Haemoglobin.		Vertex Presentation.
			Wasserman Reaction,

MEDICAL STAFF

Honorary Physicians

R. H. BLAIR, B.Sc., M.B., Ch. B. (Livp.).
L. W. HALE, M.D. (Lond.), F.R.C.P. (Lond.).

Dermatologist

F. A. WHITLOCK, M.B., B. Chir. (Cantab.), M.R.C.P. (Lond.).

Neurologist

N. S. ALCOCK, M.D., M.R.C.P.

Honorary Surgeons

G. P. O'DONNELL, M.B., B.Ch., B.A.O. (Dub.).
 T. M. REID, M.B., Ch. B., F.R.C.S. (Edin.).
 J. H. DONOVAN, M.B., Ch. B. (Leeds), F.R.C.S. (Edin.).

Honorary Ear, Nose and Throat Surgeon

M. R. SHERIDAN, M.B., Ch. B., F.R.C.S. (Edin.), D.L.O. (Eng.).

Honorary Ophthalmic Surgeon

C. HEATH, M.A., M.D. (Cantab.), D.O.M.S.

Consulting Pathologist

C. RIVERS, M.A., B.Sc. (Lond.), M.D., B.Ch. (Cantab.), M.R.C.P. (Lond.).

Consulting Thoracic Surgeon

N. R. BARRETT, M.A., M.Ch. (Cantab.), F.R.C.S. (Eng.).

Honorary Anæsthetists

A. A. MACDONALD, M.B., Ch. B. (Glas.). E. O'REILLY, M.B., B.Ch., B.A.O. (Dub.).

Honorary Medical Officer

D. P. CLAHANE, M.B., B.Ch., B.A.O. (Dub.).

Pathologist (Acting)

R. SALM, M.D. (Amsterdam), L.R.C.P. & S. (Edin.).

Radiotherapist

J. A. G. HAIR, M.B., B.S. (Lond.), D.M.R.E. (Cantab.).

Radiologist

A. C. P. D. THOMSON, M.C., M.B., Ch. B., D.M.R.

Honorary Dental Surgeon

W. H. C. LEAN, L.D.S., R.C.S. (Eng).

Obstetricians and Gynæcologists

J. G. HASTINGS INCE, M.B., B.S. (Lond.), M.R.C.O.G.

R. POLLOK, L.R.C.P., L.R.C.S. (Edin.), L.R.F.P. & S. (Glas.), L.D.S. (Glas.), D.Obst. R.C.O.G.

Resident Obstetric Officer

DR. N. NOBLE, L.R.C.P., M.R.C.S.

OBSTETRIC REPORT

1947

THIS REPORT deals with the cases discharged from the County Maternity Unit at Redruth Hospital during the year 1947. "False alarms" of labour are excluded.

Booked cases are cases accepted for institutional delivery, on account of unsuitable home conditions or on medical grounds, and have attended the antenatal clinics. Emergency cases are all cases admitted direct who have not been under antenatal supervision in the clinics.

During the year 896 women were discharged from the ward. In addition 20 women who had been discharged undelivered were readmitted, of these 19 were booked cases which after investigation in the wards or on completion of treatment were returned to the antenatal clinic and subsequently readmitted for their delivery; I was an emergency case who was treated for pyelitis of pregnancy and readmitted at the request of her doctor for an external version, she was subsequently delivered at home. Thus there were 916 discharges from the ward.

		Boo	ked	Emer	gency	To	tal
		P	M	P	M	P	M
Delivered in hospital		246	334	142	97	388	431
Antenatal only	 	I	8	9	22	10	30
Postnatal admissions	 	2	16	7	- 8	9	24
Abortions	 	I	I	0	2	I	3
		250	359	158	129	408	488

ANTENATAL ADMISSIONS.

The following table shows all cases admitted during the year for treatment of complications of pregnancy.

Booked cases 93.

Emergency cases 141.

ANTENATAL ADMISSIONS

Remarks	*	Readmitted for delivery.	Readmitted postnatally.	Morbid.	T. C.	I WILL GEHVELY.								Readmitted for delivery.					Readmitted for delivery.		Readmitted for delivery. Forcens delivery. Morbid.	
Days in Hospi- tal befere deliv- ery or discharge M B B C B L S C	· · · · · · · · · · · · · · · · · · ·			28: L : L	2: L: L 33: L: SB	10: 1: 1 T	1.	17: L: L		. L .	9: L: L			33 : L : L		: T:			4: L: L		3: L : SB	. L
Freatment					Rest, etc. Surgical induction	Surgical induction	Rest, etc.	Kest, etc.	Rest, etc.	Rest, etc.	Rest, etc.	Medical induction	Medical induction	Kest, etc.	Rest, etc.	Surgical induction	Surgical induction	Surgical induction	External version	External version	External version Rest. etc.	Penicillin injections
Reason for admission			Pre-eclamptic toxaemia Abdominal pain	Mitral stenosis Abdominal pain	Pre-eclamptic toxaemia Intrauterine death	Pre-eclamptic toxaemia	Pre-eclamptic	Pre-eclamptic toxaemia	Pre-eclamptic toxaemia	Mitral stenosis	Antepartum haemorrhage	Pre-eclamptic toxaemia		Antepartum haemorrhage Deformed pelvis	Antepartum haemorrhage	Old vesico-vaginal fistula	Diabetes mellitus	Twin pregnancy	Breech presentation		Breech presentation Pre-eclamotic toxaemia	Syphilis
Maturity		388 : 388 : 39	. 36	. 38	: 39	: 37	: 38	: 36	30	: 34	: 27	. 30	: 39	. 32	. 35	: 36	. 33	: 36	: 3	: 34	. 34	38
Para	D	6 7 6	5	3 0	о н	н	0	2			١	0 0	2	н н		н	0	0 6	I :	0	00	н
- 9gA	KE	35	41 22	25	30	35	22	29	35	21	26	19	58	32	25	25	23	31	23	20	20	27
No.	BOOKED	8051 18251 7238	18886	18753	17179	18956	18444	18880	10320	18018	18232	3940	9785	19265	8624	12480	19088	19301	4771	18249	19979	4462

	SB Prolapsed cord. SB Readmitted for delivery.	L Readmitted for delivery. L Readmitted for delivery. L L	E Readmitted for delivery. L Morbid. L Readmitted for delivery. L Delivered in Looe.	E Forceps delivery. L Readmitted for delivery. L L L L L L L L	E Forceps delivery. SB Caesarean section. Morbid. L Readmitted for delivery. (L X-Ray showed twin pregnancy. (L Readmitted for delivery. L Readmitted for delivery. L Readmitted for delivery.
	29 : L 50 : L 38 : L	36 2 4 5 24 2 5 8 5 24 2 5 5 5 5	10 1 7 60 E 0 17 17 17 17 1		21.88 4 9 6 6 11.0 LT
	section rersion	Rest, etc. Drugs, etc. Hysterotomy & sterilisation Rest, etc.	Surgical induction Rest and stabilisation Surgical induction Rest Rest	Surgical induction Surgical induction External version Surgical induction Medical induction Rest, etc.	Rest, etc. Surgical induction Rest, etc. Rest, etc. Rest, etc. Nil Investigation Rest, etc.
		H4HH-	Pre-eclamptic toxaemia Pulmonary tuberculosis Diabetes mellitus Diabetes mellitus Renal investigation Threatened abortion	Hydramnios Pre-eclamptic Breech present Hydramnios Pre-eclamptic Pre-eclamptic	Pre-eclamptic toxaemia Pre-eclamptic toxaemia Pre-eclamptic toxaemia Marked oedema Pre-eclamptic toxaemia Hyperemesis gravidarum Breech presentation Haematuria Hyperemesis gravidarum Hyperemesis gravidarum
	1 : 34 3 : 34 0 : 37 2 : 34				0 37 0 38 0 37 0 37 1 28 1 32 1 32
KED	37 : 26 : 40 : 41 : 41 : 41 : 41 : 41 : 41 : 41	36 : 24 :	31 24 25 35 25 25 25 25 25 25 25 25 25 25 25 25 25	32 24 26 27 28 28 29 29 31 26 29 31 26 31 31 31 31 31 31 31 31 31 31 31 31 31	20 19 23 40 23 19 20 20 34
BOOKED	19624 4626 19299 18442 19368	13361 19534 20279 7117 18522	19910 19962 20326 19472 19872 20738	20058 20739 20473 19138 18883 18523 20252	18594 21086 21317 19878 20254 13646 21788 21518

	Readmitted for delivery. Readmitted. Missed abortion at home.	the state of the s	Readmitted for delivery 1948.
	7 : L : L 30 : L : – 21 : L : –	# 4 ~ & \$ 4 4 8% 2 5 5 1 5 5 5 5 5 5 4 4 4 4 4 6 8 4 8 4 4 4 4 4 6 8 4 8 4 4 4 4	1 : L : L 2 : L : SB
	Investigation Investigation Rest	Removed Caesarean section Investigation Rest, etc. Caesarean section Rest, etc. Rest etc. Rest Rest Rest Drugs, etc. Surgical induction Drugs Rest Rest Rest Rest Rest Rest Rest Res	External version Rest
	Abdominal pain Essential hypertension Essential hypertension		Breech presentation Antepartum haemorrhage
KED	17: 0: 36 38: 2: (24 31	28	19: 0: 39 36: 0: 32
BOOKED	22002		

19 1 1 1 1 1 1 1 1 1
Rest, etc. Drugs, etc. Drugs, etc. Drugs, etc. Orugs, etc. Drugs, etc. Nargical induction Nil Rest, etc. Medical induction Nest, etc. Rest, etc.
Pre-eclamptic toxaemia Antepartum haemorrhage Acute hydramnios Pre-eclamptic toxaemia Antepartum haemorrhage Antepartum haemorrhage Pyelitis Pre-eclamptic toxaemia Pyelitis Pre-eclamptic toxaemia Antepartum haemorrhage Antepartum haemorrhage Antepartum haemorrhage Fre-eclamptic toxaemia Antepartum haemorrhage
Secondary Seco

Disseminated sclerosis.	Twins. Self discharged.	For delivery at home.	0	Blood transfusion. Delivered at home. Blood transfusion. Twins.	Face delivery. X-Ray showed vertex presentation. For delivery at home	Ovariotomy on 4th day of puerperium. Blood transfusion.	For delivery at home. Discharged at own request. For delivery at home.	
SB	الميم	L I SB	SB	J.S.J.J.	۱ ابر	11 H	188 1.	1
14: L 33: L	2 . L	3 21 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	13:1 13:1 2:1	1 2 I L		1,0 4,0 T.	18 6 4 4 6 10 1 1 1 1 1 1	
Rest, etc. Surgical induction	Rest, etc. Rest, etc. Rest, etc.	Rest, etc. External version Rest, etc. Rest, etc.	Surgical induction Surgical induction Internal version Rest, etc.	Rest, etc. External version Rest, etc. Rest, etc.	Rest, etc.	Nil Rest, etc.	Blood transfusion Rest, etc. Rest, etc. External version Drugs, etc.	1000
Antepartum haemorrhage Diabetes mellitus	Pre-eclamptic toxaemia Antepartum haemorrhage Hyperemesis gravidarum	Antepartum haemorrhage Breech presentation Anaemia Pre-eclamptic toxaemia	Pre-eclamptic toxaemia Pre-eclamptic toxaemia Antepartum haemorrhage Pre-eclamptic toxaemia	The second second			Anaemia Hb. 20% Hysteria Debility Breech presentation	1 years
3ENCY 22: 0: 35 42: 1: 32	1 0 0 1	41: 10: 40 24: 0: 38 38: 1: 36 25: 4: 28	31: 0: 37 32: 0: 35 37: 2: 40 24: 0: 40	0000	00	o m · o	23 2 37 35 2 37 23 1 (34 (37	
EMERGENCY 19423 22 : C	19538	19767 19873 19411 19422	19703 19738 19908 19871	19916 20037 19911 19946	19455	19870 19959 20113	19041 20275 20320 20379	1991/

Forceps delivery For delivery at home. For delivery at home. Morbid. Blood transfusion	Eclampsia. Forceps delivery. Morbid. Forceps delivery. Morbid. Discharged herself.	For delivery at home. For delivery at home. Undelivered.
7777777777 77777 77777	777777 774674 777770 716177	
H 2 2 2 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	98 48 H & & H & & O O O O	8 48 8 6 L H C
Surgical induction Rest, etc. Drugs, etc. Rest, etc. Rest, etc. Rest and diet External version Rest Rest Rest Rest Rest	Rest, etc. Rest, etc. Rest, etc. Fest, etc. Rest, etc. Surgical induction Rest, etc. Nil Rest, etc. Nil Rest, etc. Nil Rest, etc. Medical induction Medical induction	Resc. etc. External version Resc. etc. External version Rest, etc. Surgical induction External version Rest, etc.
Pre-eclamptic toxaemia Antepartum haemorrhage Pyelitis Pre-eclamptic toxaemia Antepartum haemorrhage Hyperemesis Breech presentation Pulmonary tubercu.os.s Femoral thrombosis		Anaemia Anaemia Breech presentation Degenerating fibroids Breech presentation Pre-eclamptic toxaemia Pre-eclamptic toxaemia Breech presentation Antepartum haemorrhage
34 37 37 37 38 38 38	36 4 4 6 9 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	330 11388
NCX	0 H O O H O O H O O H O O H O O H O	0 H O O O D D O D
EMERGENCY 20147 22 : 0 20321 30 : 1 20219 26 : 0 20427 34 : 0 20567 28 : 0 20567 28 : 0 20568 33 : 0 20733 21 : 1 20748 20 : 0 20481 33 : 0 14653 35 : 3	28 26 26 36 36 36 36 36 37 37 37 37 37 37 37 37 37 37 37 37 37	220 27 267 257 256 37 256 37 39 37 39 37 39 37 30 37 30 37 30 37 37 37 37 37 37 37 37 37 37 37 37 37
20147 20321 20321 20427 20427 20567 20686 20733 20748 20748	20428 20791 20869 20543 21018 20944 20732 12432 210455	21020 21267 21413 21556 21412 21112 21424 21755 21768

	For delivery at home. Discharged herself. Forceps delivery For delivery at home. Died undelivered 5 hours	Plac For Bre Disc	Forceps delivery. Morbid. Blood transfusion.	Disc	Forceps delivery. Breech delivery.
	24 17 17 17 17 17 17 17 17 17 17 17 17 17		28 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1		35 4 4 5 1 1 1 2 3 5 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
	Drugs Rest, etc. Surgical induction Rest, etc. Rest, etc. Surgical induction Rest, etc. Rest, etc.	Caesarean section Rest, etc. Rest, etc. Rest, etc.	Kest, etc. External version Rest, etc. Rest, etc. Rest, etc. Rest, etc.	Rest, etc. Rest, etc. Rest Rest	Rest, etc. Rest, etc. Rest, etc. Rest, etc. Rest, etc.
	5 Pyelitis 2 Habitual abortion 8 Pre-eclamptic toxaemia 6 Pre-eclamptic toxaemia 5 Pre-eclamptic toxaemia 6 Pre-eclamptic toxaemia 6 Antepartum haemorrhage 7 Hyperemesis gravidarum 8 Eclampsia		6 Antepartum haemorrhage 9 Breech presentation 9 Pre-eclamptic toxaemia 8 Pre-eclamptic toxaemia 4 Pre-eclamptic toxaemia 6 Eclampsia	Pre-eclamptic toxaemia 8 Hyperemesis 9 Abdominal pain 2 Pre-eclamptic toxaemia	3 Pre-eclamptic toxaemia 9 Pre-eclamptic toxaemia 9 Pre-eclamptic toxaemia 5 Antepartum haemorrhage 6 Essential hypertension
EMERGENCY	21482 32 5 35 21418 22 3 12 21474 25 0 38 21478 28 1 36 21425 27 0 35 21675 29 0 26 21675 27 0 30 21675 27 0 20 21675 27 0 20 21675 27 0 20 21675 27 0 20	15016 25: 2: 38 21908 41: 1: 26 21944 18: 0: 28 22255 23: 0: 36	22047 27: 0: 36 22092 31: 1: 39 22256 39: 1: 39 222282 43: 0: 38 21961 23: 0: 34 22091 37: 0: 36	222064 27: 0: 37 222404 27: 1: 38 22576 30: 2: 39 22039 29: 1: 32	40: 0: 17: 0: 40: 0: 25: 1: 39: 1:

	Forceps delivery. Died undelivered. Discharged herself. For delivery at home	Breech delivery.	No living child. Morbid. Eclampsia. Vertex presentation by X-rays. For delivery at home.	Morbid. Morbid. Discharged herself after 6 hrs. Twins.	For delivery at home.
	797771	プロリプロプロト	ושרח	77 75	SB
	חשחחחשח	コロコロココロ	HHHH I	7111	777
	8 25 39 5 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	W W W V H 4 W U	0 + 0 · 0 · 0	12 12	1 6 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
	Rest, etc. Surgical induction Rest, etc. Rest, etc. Rest, etc. Rest, etc. Deep X-Ray therapy Rest, etc.	Rest, etc. Rest, etc. Medical induction Surgical induction Rest Rest Medical induction	Caesarean section Medical induction Surgical induction Nil	Surgical induction Surgical induction Nil Rest	External version Rest Rest
	Pre-eclamptic toxaemia Infantile paralysis Antepartum haemorrhage Antepartum haemorrhage Pre-eclamptic toxaemia Antepartum haemorrhage Secondary carcinoma of heart Threatened abortion		Essential hype Pre-eclamptic Pre-eclamptic Breech present	Pre-eclamptic toxaemia Pre-eclamptic toxaemia Pre-eclamptic toxaemia Pre-eclamptic toxaemia	Breech presentation Pre-eclamptic toxaemia Hyperemesis gravidarum
1388	300 30 48 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6	68 0 4 8 0 4 8 0 0 0 0 0 0 0 0 0 0 0 0 0	383333	37 37	39
CY	онюонон	0400400	9 40 0 4	0 0 0 1 10	ном
SEN	22 22 23 33 33 34 25 25 25 25 25 27 27 27 27 27 27 27 27 27 27 27 27 27	37 : 19 : 19 : 28 : 28 : 28 : 28 : 28 : 28 : 28 : 2	40 330	32 : 36 : 19 : 21 :	32 33
EMERGENCY	22549 21629 22190 22575 22626 22017 21571 23015	222934 222935 23000 22747 23277 23113 23086	23269 23189 23310 23604	23110 23182 23710 23710	15995 23433 23241

POSTNATAL ADMISSIONS.

During the year there were 35 women admitted after delivery.

The reasons for admission were :—

BOOKED CASES.

Primiparae 2. Multiparae 16.

Delivered at home and admitted for nursing.

EMERGENCY CASES.

	Primiparae	Multiparae	Total
Unsuitable home condition	ns I	3	4
Retained placenta	3	2	5
Acute liver atrophy	2	0	2
Perineal tear	I	. 0	I
Premature baby	0	2	2
Jaundiced baby	0	ı	I

ABORTIONS.

Threatened abortions admitted for treatment are included in the table of antenatal admissions. Most abortions are admitted to the Gynaecological wards and are treated there and omitted from this report. One booked primigravida was treated with an inevitable abortion. One booked multiparae had a therapeutic abortion performed for essential hypertension and two emergency multiparae were treated with inevitable abortions.

Details of therapeutic abortions are appended.

THERAPEUTIC ABORTION

Remarks	Sterilization.
Result M	Ĥ
Method	Abdominal hysterotomy
Indication	20 Essential hypertension B.P. 250/150
Maturity in weeks	
Para	O :
Age	BOOKED 7117 32: 4
No.	ВО 7117

FIBROIDS AND PREGNANCY.

There were three cases admitted with fibroids in pregnancy.

BOOKED.

20661 Aged 42 para 2 Normal labour.

21085 Aged 40 para o Normal labour.

EMERGENCY.

21556 Aged 37 para 2 Antenatal only for red degeneration.

OVARIAN CYST AND PREGNANCY.

EMERGENCY.

19959 Aged 44 para 3 Normal labour, pseudomucinous.

ovarian cyst removed on the 3rd day of puerperium.

22282 Aged 43 para o Normal labour, ovarian dermoid found at post-mortem examination.

SYPHILIS IN PREGNANCY.

A routine Wasserman reaction is done on all booked cases in the antenatal clinic and on all emergency cases after admission. There were eight positive results.

> Booked primigravidae 2 Booked multiparae 6

HEART DISEASE IN PREGNANCY

There were five booked cases treated in hospital.

Remarks							
C		T	D	T	SB	T	
Result		L	ı	1	1	h	
Delivery		Forceps	Normal	Normal	Normal	Normal	
Days in Hospital before delivery		91	40	0	0	15	
Exercise		Good	Pood	Good	Fair	Cood	
Auricular		Nc	No	No	No	No	
Congestive heart failure		INI	Nil	Nil	Nil	INI	
Enlarge- ment		Slight	Slight	Slight	Slight	Slight	
Lesion		Mitral	stenosis	Stenosis	Patent	Septum Mitral stenosis	
Maturity at delivery		: 40	. 40	: 40	. 32	40	1
Para	0	0 :	0		0	. н	
93A	BOOKED	25	21	7434 27	3 23	26	100
No.	BOG	18753 25 :	18018	7434	19038	22480	1

PULMONARY TUBERCULOSIS IN PREGNANCY.

Five booked and one emergency case were treated in the wards. Two of the booked cases were admitted after delivery at home, four cases were in an active stage of the disease.

Remarks	Morbid. Blood transfusion. Postnatal admission. Morbid. Induction. Postnatal admission.	Morbid.
Result M C		L : L Morbid.
Labour	Normal Normal Normal Normal	Normal
Days in hospital before delivery	о н о г	e
State of disease	Quiescent Active Active Active Quiescent	Active
Maturity at delivery	35 39 40	. 40
Para	C 1 4 1 2 1	CY C
- Age	BOOKED 595 37 : 317 30 : 326 24 : 754 31 : 209 37 :	GEN 20:
N _O	BOOKE 9595 37 5317 30 20326 24 21754 31 22209 37	EMERGENCY 20748 20 : 0

ANTEPARTUM HÆMORRHAGE.

A case is only classified as Placenta Prævia when there is definite evidence by palpation of the edge of the placenta that it was attached in the lower uterine segment.

Cases of concealed hæmorrhage or those in which on examination of the lower uterine segment under anæsthesia no placenta was felt are classified as accidental hæmorrhage.

All cases in which no complete examination of the lower uterine segment was made are grouped under the heading antepartum hæmorrhage of unknown origin.

PLACENTA PRÆVIA.

Cases are classified as:

ist degree When the edge is felt in the lower uterine segment but does not reach the internal os.

and degree When the edge of the placenta reaches the internal os but does not cover it.

3rd degree When the placenta completely covers the internal os when closed but not when fully dilated.

4th degree When the placenta completely covers the fully dilated os.

PLACENTA PRAEVIA

		Morbid. Morbid.
Remarks	Blood transfusion.	Blood transfusion. Mc Blood transfusion. Prolapsed cord. Blood transfusion. Blood transfusion. Mc Undelivered. Toxic myocarditis.
Result M C	T : T	1 11 11 10 EU118 11 11
Treatment	Caesarean section	Rest, etc. Caesarean section Caesarean section Internal version Rest, etc. Internal version for transverse lie Caesarean section Rest, etc.
Degree	2nd 2nd	1st 3rd 3rd 1st 1st 1st 2nd 3rd
Dilatation of os on admission	Closed	Closed
Condition on admission	Fair	Fair Good Good Good Fair Good
Maturity	37	33 34 40 40 38 39 39 39 39 39 39 39 39 39 39 39 39 39
Para	н о	2 2 2 2 2 2 2 2 2 2 3 3 3 3 3 3 3 3 3 3
9gA	BOOKED (938 27 :	GEN 31: 32: 37: 37: 20: 20: 20: 25: 35:
No.	BOOKED 21938 27 : 15879 35 :	EMERGENCY 18655 38 : 5 18994 31 : 2 18995 32 : 5 19908 37 : 2 20113 20 : 0 21666 23 : 0 15016 25 : 2 12432 35 : 2

ACCIDENTAL ANTEPARTUM HAEMORRHAGE

Remarks			SE SE Blood transfusion. SB Blood transfusion.
Days in Hospital Defore delivery or discharge		I:L:SB	72: L: SB 1: L: SE 0: L: SB 0: L: SB
Treatment		Blood transfusion	Rest. Artificial rupture of membranes
B.P. on admission		100/80	178/120 220/140 120/80 120/80
Albumin on admission		8 0/00	1 0/00 2 0/00 Nil +
Variety		Mixed	External Mixed Mixed External
In		Yes	No No Yes No
Condition on admission		Poor	Fair Fair Fair
Maturity		33	36 36 37 37 32
Para		24	80000
əgA	KED	33 :	28: 28: 24: 32: 39: 39: 39: 39: 39: 39: 39: 39: 39: 39
No.	ВООКЕВ	18585 33:	EMERGENCY 18873 28 : 0 : 14912 32 : 0 : 23014 24 : 0 : 23602 39 : 6 : :

ANTEPARTUM HAEMORRHAGE OF UNKNOWN ORIGIN

Remarks		Recurrent bleedings.	Readmitted for delivery.		Postpartum haemorr-	Morbid.	
Days in Hospital Defore delivery or discharge		J: T: 06	33: L: L	40 : L : L	I:L:L		2: L: SB
Treatment		Rest	Rest	Rest	1		1
B.P.		146/80	130/90	120/60	180/120		148/90
Albumin		No	No	No	No		No
Variety		External	External	External	External		External
In labour		No	No	No	Yes		Yes
Condition on admission		Good	Good	Good	Good		Cood
Maturity		: 27	: 32	: 35	: 40		32
Para	-	4	I	3	0		0
- 9gA	BOOKED	26:	32 :	25:	32 :		36:
No.	BOO	18232	19265	8624	19462 32:		22475 36: 0: 32

ANTEPARTUM HAEMORRHAGE OF UNKNOWN ORIGIN (continued)

				Blood transfusion.			Twins.				Twins.			Blood transfusion.	Blood transfusion.	Face delivery.																		Forceps delivery.			
	10. T. O.	. r.	20: L: L				1: T: I	٦): ':	- 7		8: L:(L		2: L: L	I:L:L			1: I: D			. T :						 	· ·			 	: 7:	. न		: T:	65 : L : L		
1	Rest etc		Rest, etc.			Kest, etc.	Rest, etc.				Rest, etc.			Rest, etc.	Rest, etc.	Rest, etc.	1						Rest, etc.	1		Rest, etc.		Kest, etc.		Rest, etc.		Rest, etc.	Rest	Rest	Rest	Rest	
	144/84	134/96	120/90	178/94	140/94	150/90	110/74		152/112	148/100	150/90		134/86	106/64	06/011	120/80	148/88		124/90	110/74	120/90	08/811	140/100	130/70	130/84	156/100	02/011	132/100	150/98	140/84	110/85	120/84	08/811	140/88	102/64	120/90	
	I.N.	NE	Nil	Nil	Nil	TZ.	Nil		Nil	NIL	Nil		Nil	Nil	Nil	NII	Nil	INI	IN	Nil	Nil	Nil	NI	Z	N	IZ.	IN.	TZ.	TN	NII	Nil	N	Nil	Nil	Nil	Nil	
	Evternal	External	External	External	External	External	External		External	External	External	The same of	External	External	External	External	External	External	External	External	External	External	External	External	External	External	External	External	External	External	External	External	External	External	External	External	
	No	No.	No	Yes	No	No	No		No	No	No		No	No	No	No	Yes	No	No	No	No	No	No	Yes	Yes	oN;	Yes	No	Yes	oN:	Yes	No	No	No	No	No	
	Cond	Poop	Good	Fair	Good	Good	Good		Good	Good	Good		Good	Good	Good	Good	Good	Good	Good	Good	Good	Good	Good	Good	Good	Good	Good	Cood	D005	Good	Good	Good	Good	Good	Good	Cood	
, ADIAL.		36: 0: 36	. 4 :	: 9 :	: I :	. 4 :	. 2 :		: 0 :	. 0 :	25: 0: 34		: OI :	22: 0: 40	24: 0: 38	: 0 :	: 0 :	: 0 :	: I :	: 0 :	: I :	: 0 :		. 0 .		: 0 :	. 3 .	. 4 .	. 0 .		: 0 :	: I :		30:0:38	20: I: 30	30: 2: 28	
- Committee	EMERGENCI	18570	18754	18161	18590	19415	19464		19465	19423	15019		19767	91661	11001	19455	20073	19870	20321	20567	20428	20791	20869	21081	10080	21408	21585	21156	21475	22047	22183	22062	22190	22575	22017	23277	

EXTERNAL VERSION.

Only cases admitted for external version are included in the table. All cases found to have a breech presentation have an attempt to correct this in the antenatal clinic whenever the presentation is recognised and the pregnancy has reached the 32nd week.

EXTERNAL CEPHALIC VERSION

Remarks	Readmitted for delivery. Face presentation. Readmitted for delivery. Readmitted for delivery. Readmitted for delivery. Readmitted for delivery. Readmitted for delivery.	For delivery at home. For delivery at home. Delivered at home. For delivery at home.
Result M C	1 11111 1 188	
Labour	Forceps delivery Normal Normal Normal Normal	Normal Normal Normal Normal
Attitude of foetus	Extended legs Extended legs Extended legs Extended legs Extended legs Extended legs	Extended legs Extended legs Extended legs Extended legs Extended legs Flexed Extended legs Extended legs
Version	Yes Yes No Yes Yes Yes Yes	Yes Yes Yes Yes Yes Yes Yes
Complica- tions during version		
Maturity at delivery Anaesthesia given	40 : Yes 40 : No 35 : Yes 38 : Yes 40 : Yes 40 : Yes	No No No Nes No Nes No Nes No No No No No No No N
Maturity at version	34 37 37 37 37 37	39 39 37 37 37 37 39 39
Para	н 00000	CY 33 11 11 11 11 11 11 11 11 11 11 11 11
- 93A	20 : 20 : 20 : 20 : 20 : 20 : 20 : 20 :	3EN 339: 330: 225: 227: 227: 227: 227: 227: 227: 227
No.	BOOKED 4771 23 : 18249 20 : 19979 26 : 18442 20 : 19138 22 : 23707 19 :	EMERGENCY 6771 39 : 2 19611 30 : 3 19873 24 : 0 20037 25 : 0 20733 21 : 1 21412 22 : 0 21412 22 : 0 21413 39 : 5 21755 33 : 1 15995 27 : 2

INDUCTION OF LABOUR.

The following tables give details of cases induced and the reason for the induction.

INDUCTION OF LABOUR FOR

No.	Age	Para	Maturity on admission	Condition on admission	Progress	Method of induction
вос	KEI)				
18956	35 :	I	: 37	B.P. 142/92. Albumin nil. Oedema nil.	Albuminuria appeared	Drugs and A.R.M.
14825	42 :	4	: 39	B.P. 134/92. Albumin 2 0/00. Oedema nil.	Hypertension increased	Drugs
3946	19:	0	: 38	B.P. 122/92. Albumin 2 0/00. Oedema slight.		Drugs
19045	36:	0	: 36	B.P. 180/120. Albumin 5 0/00. Oedema slight.	At term	Drugs
9785	28 :	2	: 40	B.P. 136/84. Albumin 2 0/00. Oedema slight.	At term	Drugs
19962	21 :	0	: 34	B.P. 160/104. Albumin 5 0/00. Oedema of legs.	Hypertension increased	A.R.M.
20473	32 :	1	: 34	B.P. 176/130. Albumin 2 0/00. Oedema nil.	Hypertension persisted	A.R.M.
18523	26 :	0	: 37	B.P. 160/100. Albumin nil.	Albuminuria appeared	Drugs
21317	23:	0	: 37	B.P. 130/90. Albumin trace. Oedema of legs & abdomen.	At term	A.R.M.
21319	35 :	0	34	B.P. 150/94. Albumin nil. Oedema ankles.	At term	Drugs
23040	27 :	I	: 37	B.P. 156/90. Albumin nil. Oedema ankles.	No improvement	Drugs
23185	38 :	3	: 40	B.P. 164/92. Albumin nil. Oedema ankles.	No improvement	Drugs
22774	35 :	0	: 36	B.P. 160/94. Albumin + +. Oedema generalised.	At term	Drugs
20758	28 :	0	: 37	B.P. 160/110. Albumin nil. Oedema feet.	Albuminuria appeared	Drugs

PRE-ECLAMPTIC TOXAEMIA.

Highest B.P. (excluding labour)	Albumin on discharge	B.P. on discharge	Length of induction	Length of labour	Days in Hospital	Re	esult C	Remarks
		miles in land						
148/96	Nil	118/82	3½ hrs.	9½ hrs.	18	: L	: (L	Twins.
148/96	-	-	Stat.	. 3 hr.	7	D		- Land
166/122	Nil	120/80	Stat.	12½ hrs.	14	L	L	100
190/110	Nil	182/98	Stat.	2½ hrs.	29	L	: D	Control of the contro
136/84	Nil	124/80	Stat.	2 hrs.	3	L	L	
180/118	Nil	116/86	2 hrs.	16 hrs.	10	L	L	
182/132	Nil	120/82	16½ hrs.	3½ hrs.	11	L	L	
176/110	Nil	154/104	Stat.	10 hrs.	17	L	L	
140/90	Nil	144/96	Stat.	16 hrs.	17	L	L	Forceps delivery.
170/108	Nil	112/68	Stat.	15 hrs.	43	L	L	Forceps delivery.
162/112	Nil	140/110	Stat.	5½ hrs.	14	L	L	
210/140	Nil	160/110	Stat.	5 hrs.	2 :	L	L	
170/124	Nil	102/74	Stat.	39 hrs.	14	L	L	
180/118	Nil	130/90	Stat.	11½ hrs.	20	L	L	
	8 8 9 1							

INDUCTION OF LABOUR FOR

		1-1-1	-			
No.	Age	Para	Maturity on admission	Condition on admission	Progress	Method of induction
EMER	GEN	CY		Alth		COLLEGE STATE
18771			: 39	B.P. 170/120. Albumin	At term	Drugs
18605	25	0	: 36	3 o/oo. Oedema to knees. B.P. 168/7o. Albumin nil. Oedema to knees.	No improvement	Drugs and A.R.M.
18006	20	0	: 30	B.P. 190/140. Albumin nil. Oedema nil.	Hypertension persisted	Drugs and A.R.M.
18913	31	I	: 36	B.P. 190/120. Albumin 2 0/00. Oedema to knees.	Albuminuria increased	Drugs
19080	23	0	: 40	B.P. 192/120. Albumin 5 0/00. Oedema generalised.	At term	Drugs
19212	35	3	: 38	B.P. 184/124. Albumin 7 o/oo. Oedema nil.	Eclampsia	Drugs and A.R.M.
18970	33	1	: 32	B.P. 180/110. Albumin 5 0/00. Oedema slight.	Albuminuria increased	Drugs
19703	31	0	: 37	B.P. 194/110. Albumin trace. Oedema generalised.	No improvement	A.R.M.
19738	32	0	: 35	B.P. 220/132. Albumin 7 0/00. Oedema slight.	No improvement	A.R.M.
20147	22	0	: 38	B.P. 146/100. Albumin 9 0/00. Oedema nil.	Albuminuria increased	A.R.M.
20944	27	. 0	: 35	B.P. 164/102. Albumin 6.5 0/00. Oedema generalised.	Oedema increasing	A.R.M.
21055	37	1	: 28	B.P. 146/90. Albumin 2 0/00. Oedema of legs.	Hypertension increased	A.R.M.
21020	37	2	: 37	B.P. 144/92. Albumin trace. Oedema of legs.	At term	Drugs
21424	20	0	: 39	B.P. 170/110. Albumin 6 o/oo. Oedema of legs.	No improvement	Drugs and A.R.M.
21474	25	0	: 38	B.P. 160/70. Albumin 0.5 0/00. Oedema of legs.	Hypertension increased	A.R.M.
21475	29	0	: 36	B.P. 172/84. Albumin 1 0/00. Oedema slight.		A.R.M.
23000	19	0	: 40	B.P. 160/116. Albumin +. Oedema generalised.	At term	Drugs
22747	21	0	: 33	B.P. 208/134. Albumin 6 o/oo. Oedema generalised.	Albuminuria increased	A.R.M.
23086	28	0	:	B.P. 226/130. Albumin 4.5 0/00. Oedema of legs.	Albuminuria increased	Drugs
23189	26	: 0	. 1 2	B.P. 190/100. Albumin trace. Oedema nil.		Drugs
23310	23			B.P. 200/100. Albumin 10 0/00. Oedema generalised.	No improvement	A.R.M.
23110	33	: 2	: 36	B.P. 160/1co. Albumin trace. Oedema of feet.	No improvement	A.R.M.
23182	36	I	: 38	B.P. 180/122. Albumin 1.5 0/00. Oedema slight.	Albuminuria increased	A.R.M.
Water William	STALL S	-			The second second second	

PRE-ECLAMPTIC TOXAEMIA (continued).

Highest B.P. (excluding labour)	Albumin on discharge	B.P. on discharge	Length of induction	Length of labour	Days in Hospital before delivery	Ro	sult C	Remarks
	Da For Ge	SALES OF						
170/120	Nil	146/92	Stat.	18 hrs.	2	L	L	
170/90	Nil	138/92	96 hrs.	18 hrs.	II	L	L	
190/140	Nil	160/100	46 hrs.	37 hrs.	32	L	L	
190/120	Nil	140/90	Stat.	II hrs.	8	L	L	
190/130	Nil	130/90	Stat.	5 hrs.	4	L	L	
190/144	Nil	140/102	1½ hrs.	10 hrs.	4	L	SB	
180/110	Nil	140/120	Stat.	7 hrs.	28	L	L	
194/110	Nil	154/100	3 hrs.	14 hrs.	11	L	L	
220/132	2 0/00	120/84	22 hrs.	78 hrs.	13	L	L	
146/106	Nil	110/74	4 hrs.	17 hrs.	11	L	L	
194/130	Nil	128/100	7 hrs.	7 hrs.	3	L	D	Intrapartum eclampsia. Forceps.
152/100	Nil	180/110	6 hrs.	3½ hrs.	10	L	L	cetampsta. 2 orecps.
142/92	Nil	120/80	Stat.	24 hrs.	19	L	L	No living child.
170/120	Nil	114/70	1½ hrs.	56 hrs.	7	L	L	
170/110	Nil	130/90	Stat.	3 hrs.	12	L	L	
174/84	0.5 0/00	130/80	ı hr.	113 hrs.	25	L	L	Forceps delivery.
160/116	Nil	110/70	Stat.	7½ hrs.	3	L	L	
208/134	Nil	124/80	14 hrs.	20 hrs.	7	L	L	
226/130	trace	178/120	Stat.	8 hrs.	8	L	D	
190/100	Nil	124/84	Stat.	8 hrs.	9	L	L	Morbid.
200/100	Nil	112/78	2 hrs.	17 hrs.	5	L	D	Eclampsia.
160/112	Nil	120/78	4½ hrs.	7½ hrs.	24	L	L	Morbid.
180/122	Nil	140/92	2½ hrs.	7½ hrs.	16	L	L	Morbid.

INDUCTION OF LABOUR FOR OTHER CAUSES

	Remarks			Anencephalic, Previous	caesarean section.	Droloneed over	From Deed Cold.			Forceps delivery.	Morbid.	No living child.	Forceps delivery.	B.P. 200/124. Forceps delivery.				Breech delivery.
	sult	Result M C		: SB	1	1-		. L	O :		. T	r	. L	r	-	: SB	. L	. D
	Re			J	1	H	4	Г	1	L	T	r	I	1		T	L	J
	.	Length		1	21"	20"	22"	20″	1	22"	101"	194"	19,"	1		151,"	18,,	"41
	Child	ht		TO OZ.	7 oz.	2 oz.	5 oz.	2 oz.	I oz.	6 oz.	12 OZ.	4 oz.	5 oz.	TO OZ.		2 oz.	13 oz.	lb. ro oz.
		Weight		I lb. I	8 lb.	7 lb.		8 lb.	5 lb.	11 lb. 6 oz.	lb.		5 lb.	8 lb. 1		4 lb.	4 lb. 1	2 lb. 1
-	Duration of	labour		roo hrs.	15 hrs.	14½ hrs.			2½ hrs.	17 hrs.	58 hrs.	94 hrs.		36 hrs.		I hr.	19 hrs.	2 hrs.
	Duration of Duration of	induction		*24 hrs.	24 hrs.	ra hrs.			5 hrs.	3 hr.	3 hrs.	36 hrs.		19 hrs.		72 hrs.	Statim	48 hrs.
-		Method		A.R.M.	A.R.M.	A.R.M.	A.N.M.	A.R.M.	A.R.M.	A.R.M.	A.R.M.	A.R.M.	A.R.M.	A.R.M.		A.R.M.	Drugs	A.R.M.
		Indication		35 Intrauterine	0	00	t ast mouth	39 Oedema of legs	35 Pulmonary	36 Hydramnios	Hydramnios	37 Diabetes mellitus	Essential	nypertension Diabetes mellitus		34 Diabetes mellitus	田	nypertension 34 Infantile paralysis
-	Para			: 35	: 37	38	· · ·	: 39	: 35	36	35	: 37	: 36	: 37	1	: 34	: 37	. 34
-			D	I	н	0 0		2	I :				0 :	0	ZCX	I :	39: 'I	н
-	-ge	¥	BOOKED	. 30	25	3 23	_	3 4I	24	24	38	_	_	31	RGEI	1 42		28
	No.		BOC	18834 30 :	12480	19988	19301	19368	20326	20739	18883	19472	22048	20602	EMERGENCY	: 42 :	22474	21629

METHODS OF DELIVERY.

	Booked		Emerg	gency	то То	Total	
Spontaneous delivery	P	M	P	M	P	M	
occipito-anterior	218	302		71	313	373	
Spontaneous delivery persistent							
occipito-posterior	3	3	0	4	3.	7	
Breech delivery	4	4	9	10	13	14	
Face delivery	0	2	I	3	I	5	
Forceps delivery	18	5	30	I	48	6	
Cæsarean section	2	6	2	4	4	10	
Twins	I	12	5	4	6	16	
	246	334	142	97	388	431	

BREECH DELIVERIES.

The following tables give details of all cases delivered as a breech. The case is only regarded as complicated when there is some maternal or foetal complication. Primiparity and extension of the limbs are not considered by themselves as complications.

BREECH DELIVERIES - UNCOMPLICATED.

	n	'n.
S	External version unsuccessful 1946. External version twice in clinic.	Awaiting external version.
Remarks	rsion t	ternal
н	nal vel 46. nal ve nic.	ng ex
	External version unsucces 1946. External version twice in clinic.	Awaiti
Lit C	S SB TITI	חחקחח
Result M 0	ח ח חחחח	
70	5 oz. 6 oz. 14 oz. 14 oz. 9 oz.	02. 02. 02.
Weight of	. 5 oz. 6 oz. 14 oz. 14 oz. 9 oz.	. 2 0z. . 10 0z. . 15 0z. . 14 0z. . 9 0z.
We	7 lb. 7 lb. 6 lb. 8 lb.	7 Pb.
	Extended arms brought down 7 lb. 5 oz. Arms brought down 7 lb. 6 oz. Nil 7 lb. 6 oz. Nil 7 lb. 0 oz. Nil 8 lb. 14 oz. Nil 8 lb. 9 oz.	
	ught n	
Treatment	down	
Trea	Extended arms brou Arms brought down Nil Nil Nil	
	tende	
	NI NI NI Ext	BEEEE
	So	
Attitude of Breech	l arm	arms
e of]	rs and ms	and sis
ttitud	ed leg ed arr	ed legs ed legs ed legs ed legs ed legs
A	40 Extended legs and arms 40 Extended arms 40 Flexed 40 Extended legs 37 Flexed 38 Flexed	Extended leg and arms Extended legs Extended legs Extended legs Extended legs
Maturity	36 H 40 H 4	04 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4
Para	0 2 0 H	CY 2 :: 0 :: 0 :: 0 :: 0
- 9gA	BOOKED 5804 32: 7583 41: 9378 20: 9621 29: 1630 30: 1184 37:	GEN 27 : 19 : 24 : 22 : 25 :
No.	BOOKED 16804 32: 17583 41: 20378 20: 19621 29: 21630 30: 22184 37:	EMERGENCY 12367 27 : 2 8964 19 : 1 20108 24 : 0 20882 22 : 0 20729 25 : 0

BREECH DELIVERIES - COMPLICATED.

Remarks			Morbid.		Declared cond	Floiapsed cold.				*		
rit C		SB	11		1-	11	T	SB	11	T :	r.	: SB
Result M (11	1		1-	4	1,	1-	11	L		Т
Weight of		5 lb. 9 oz. 4 lb. 4 oz.			2 4	9 lb. 5 oz.	6	IO	11	6	3	1
Attitude		Flexed Extended arms	Extended arms Extended legs	and arms	Extended legs	Flexed	Flexed	Extended legs	Flexed	r extended leg	I extended leg	Extended legs
Treatment during labour		E	Nil Extended arms	ought	IN.	NII	IN	Z.	NI	NI	E.	N:
Complication		Twins. 2nd twin Twins. (1st twin	Twins. 2nd twin		Ist	I wins. 1st twin	puz .	terine death	Twins. Ist twin	[ıst	[sud	Prematurity
Maturity		39	36		38	39	: 39	: 35	39	: 39	-	: 32
Para	1	24 20	-		3	01	I	0	1	0		0
93A	KEI	34:			30 :	31	34:	26 :	34	25		38 :
No.	BOOKED	14322	6823	Con	17647	19301	19625	62661	21108	21788		23222

BREECH DELIVERIES - COMPLICATED (continued).

Remarks	1	Morbid. Blood	transfusion.	Blood transfusion											Prolapsed cord.				*				Induction.				
Result M C		T: T		L: D	L : SB			I : D		Q :			I D			I SB									L . D	T: T	
Weight of		5 lb. o oz.	5 lb. ro oz.	4 lb. 8 oz.	1	4 lb. 3 oz.	3 lb. 6 oz.	2 lb. 4 oz.		2 lb. 6 oz.		1	3 lb. 15 oz.	3 lb. I oz.	7 lb. 4 oz.	lb.	5 lb. 1320z.		2 lb. 11 oz.	to Ib. 7 oz.	1	8 lb 12 oz.	2 lb. ro oz.	13		IC	
Attitude		Flexed	Flexed	Flexed	Extended legs	Extended legs	Flexed	Extended arms		Extended arms		Extended legs	Flexed	Flexed	Flexed	Fiexed	Extended leg	and arms	Flexed	Flexed	Flexed	Extended legs	Flexed	Flexed	Flexed	Fle red	
Treatment during labour		III	Nil	EN.	Nil	Nil	Nil	Extended arms	brought down	Extended arms	brought down	Spinal theca tapped	Nil	Nil	Nil	Nil	Extended arms	brought down	Internal version	Internal version	Nil	Nil	Nil	Nil	Nil	Nil	
Complication		Twins.	(2nd twin	-		-	Toxaemia	Twins. (1st twin		(2nd twin		-	-				Twins. 2nd twin		-	Transverse lie			Infantile paralysis		Twins.	Twins. 1st twin	
Maturity	1	: 38		: 34	: 32	: 32	: 32	: 30				: 37	: 34	: 34	: 40	: 40	: 36		: 30	: 40	: 30	: 40	: 34	. 39	36	: 38	
Para	CY	0		5	0 :	3	H	5				0 :	0 :	4		0 :	: 3		0 :	7	0 :	0 :	I	4	H	0	4
∍g√	GEN	31:		38	25 :	30	31	33				27	56	25	37	25	35		23	29	18	40	28	38	_	-	
No.	EMERGENCY	18574		18655	19014	19189	19458	19464				66961	15019	19422	19908	20544	14653		21666	22008	21944	22337	21629	22935	23375	23316	

CAESAREAN SECTION.

Caesarean section was performed 14 times, 8 being booked and 6 emergency cases.

Incidence of Caesarean Section in booked cases 1.4%. Incidence of Caesarean Section in emergency cases 2.5%.

CAESAREAN SECTION.

Remarks		Scoliosis. Sterilisation. Morbid. Blood	Blood transfusion.			Morbid. Blood	Blood transfusion.	Blood transfusion.	No living child.
c		SB	пп	SB		Г	11	11	ı
Result		. ההחה	니니	HH		л 	חח	חח	L
it of		11 oz. 9 oz. 13 oz.	7 oz. 15 oz.	2 oz. 15 oz.		2 oz	8 oz.	zo 6	5 oz.
Weight of		7 lb. 1 6 lb. 1	6 lb. 1	8 lb. 14 lb. 1		5 lb.		. P.	6 lb.
Type of operation		Upper segment Lower segment Upper segment Lower segment	Upper segment Upper segment	Lower segment Upper segment		Upper segment	segment	segment	2 hrs. Upper segment
No. of hrs. membranes ruptured		5 hrs.	11	48 hrs.		1	11	: 40 hrs. : 72 hrs.	2 hrs.
lo noitstud tuodsl		Nil 48 hrs Nil 72 hrs.	EE	EE		Nil	IN IN	48 hrs. 63 hrs.	Niil
Indication		Deformed pelvis Contracted pelvis Contracted pelvis Uterine inertia	Placenta praevia Suspected abdom-	C.S. Contracted pelvis Placenta praevia		Placenta praevia	Placenta praevia Placenta praevia	0	Essential hypertension
Previous		I C.S. I C.S. 2 C.S., I Ectopic o	I N.L. 3 abortions	I Forceps, I C.S.		2 N.L.	1 ab., 4 N.L. 2 N.L.	00	r forceps, r ab., Essential
Maturity		39	04	40 34	1.00	36	38	40	37
9gA	KED	40 35 40	35	35	BINCA	31 :	32	33 :	39
No.	BOOKED	19421 4626 19910 19878	21938	22366 15879	EMERGENCY	18994	18995	22410	15977

The following table gives details of the treatment of cases which had had a previous Caesarean section.

PREVIOUS CAESAREAN SECTIONS.

Remarks	Anencephaly. Postpartum haemorrhage.	
Result C	77 77 7 7 777 77 7 8 7 77 7 7 777 77 7	L SB L SB
Birth	1 lb. 10 oz. 7 lb. 11 oz. 6 lb. 4 oz. 9 lb. 9 oz. 6 lb. 13 oz. 8 lb. 0 oz. 7 lb. 10 oz. 8 lb. 4 oz. 6 lb. 2 oz. 6 lb. 2 oz. 6 lb. 8 oz.	8 lb. 2 oz. 6 lb. 4 oz. 1 lb. 8 oz.
Method of delivery	12 oz. Gaesarean section 8 oz. Normal 15 oz. Caesarean section 0 oz. Caesarean section 0 oz. Normal 2 oz. Normal 2 oz. Normal 0 oz.	oz. Caesarean section oz. Normal oz.
Weight of Caesarean baby	5 lb. 12 oz. Caesa 5 lb. 12 oz. Caesa 5 lb. 8 oz. Norm 7 lb. 15 oz. Caesa 7 lb. 9 oz. Caesa 7 lb. 0 oz. Norm 6 lb. 12 oz. Norm 9 lb. 0 oz. Norm 9 lb. 2 oz. Norm 9 lb. 8 oz. Norm 6 lb. 12 oz. Norm 7 lb. 2 oz. Norm 7 lb. 2 oz. Norm 6 lb. 8 oz. Norm 7 lb. 2 oz. Norm 7 lb. 0 oz. Norm 7 lb. 0 oz. Norm 7 lb. 0 oz. Norm 8 lb. 0 oz. Norm 9 lb. 0 oz. Norm	7 lb. o oz. 6 lb. 3 oz. 8 lb. 8 oz.
Indication for Caesarean section	Disproportion Deformed pelvis Placenta praevia Contracted pelvis Disproportion Disproportion Disproportion Disproportion Disproportion Disproportion Disproportion Hyperemesis	Contracted pelvis Disproportion Contracted pelvis
Previous	1 C.S. 1 C.S. 1 C.S. 1 C.S. 1 C.S. 1 C.S. 2 C.S., 1 ab. 1 forceps, 1 C.S., Disproportion 2 N.L. 1 C.S. 1 C.S. 1 forceps Disproportion 1 C.S. 1 C.S. 1 forceps Disproportion 1 C.S. 1 C.S. 1 Kuptured vein in broad ligamen 2 N.L. 1 C.S. 1 Hyperemesis	r forceps, r C.S. Contracted per r C.S., r forceps Disproportion r C.S.
Age	30 40 40 29 26 35 30 31 31 31	30 29
No.	BOOKED 18834 19421 18015 4626 19910 19531 20034 20034 20034 20256 21021 21158 19470 4460	22366

PREVIOUS CAESAREAN SECTIONS (continued).

		SB Toxaemia, Rhesus aggultinins present	No living child			
	1	SB	1	L	L	D :
		L	ר	.:	. J	. J
	7 lb. 2 oz.	1	9 lb. 10 oz.	7 lb. 4 oz.	7 lb. 7 oz.	2 lb. 6 oz.
	Breech	Normal	Normal	Normal	Normal	
	to lb, 13 oz. Breech	1	7 lb. 12 oz. Normal	7 lb. 8 oz. Normal	7 lb. o oz. Normal	8 lb. 4 oz. Normal
	Disproportion	Disproportion	Fibroids	Failed forceps	Disproportion	Disproportion
	I C.S., I twins	I C.S., I forceps Disproportion	4 N.L., I C.S.,	I C.S., 2 N.L. Failed fo	I C.S., I N.L.	I C.S., I N.L.
BNCY	27	. 32	34	28	25	30
EMERGENCY	12367	19412	78101	15445	9333	23277

FORCEPS DELIVERIES.

There were 56 deliveries by forceps including one set of twins, both being extracted with forceps for persistent occipito-posterior position. 23 casese were booked and 32 were emergency cases.

Incidence of forceps deliveries in booked cases	4.0%.
Incidence of forceps deliveries in emergency cases	13.4%.
Incidence of forceps deliveries in all cases	6.7%.

FORCEPS DELIVERIES.

	Remarks				Mannal dilatation of cervix	***************************************		Morbid.	-6	of placenta. Morbid	Dec.	Face presentation,	Manual dilatation of cervix	Morbid	Morbid	Blood transfusion.		Induction for toxaemia.	Essential hypertension.	Induction.	Diabetes mellitus			Morbid.		
. #	O		L	L	1	11	L	1	L	-	SB.	7.	7,	1,	1-	1	L	r	L	-	L	SB.	٦,	1	1	
Result		1																								
	M		1		7-	-	-	H	-					1-		-	_	I	-					7-		
	Circ. of head		1	1	1	13"	1	1	1		1:	15,	13	15,	142	11/1	- 1	13"	13"		1:	13,	14	15,		
P	Length		1	1	1	21"	1	1	1		13	214 ::	21.	22	. "00	. "00		: "61	: "61		1:	21" :	21"	22"		
Child			6 oz.	oz.	.zo	OZ.	OZ.	oz.	oz.		OZ.	oz.	.20	OZ.	020			OZ.	oz.		OZ.	OZ.	OZ.	OZ.		
	Weight		lb. 6	6 .					5 .). IO		P. o	4 0			-					-	0.0		
			8 11		7 lb.		6 lb.		9 lb.		8 lb.	7 ib.	7 Ib.	H .		S Ib		7 lb.	5 lb.					9 P		
of Labour	2nd stage		6 hrs.	2 hrs.	7 hrs.	hrs.		24 hrs.	4 hrs.		1;	I hrs.	1	3 hrs.	Zg mrs.			5 hrs.				ry hrs.	4 hrs.	4 hrs.		
Duration of Labour	ıst s.age		51 hrs.	27½ hrs.		57 hrs.		-	27 hrs.			35 hrs.		I4 hrs.		TE hee					35 hrs.	130 hrs.		72 hrs.	30 ms:	
	Indication		Me	distress Maternal distress	Deep transverse arrest	Foetal distress	Foetal distress	Foetal distress	Persistent	occipito-posterior	Maternal distress	Delay in 2nd stage	Foetal		Door	Foetal distress	Foetal distress	Deep transverse arrest	Deep transverse arrest		Foetal distress	Foetal distress	Delay in 2nd stage	Delay in 2nd stage	occipito-posterior	
irity	Matu		: 40	. 40	: 40	40	40	04 :	: 40		: 40	: 40	: 40	36	40	40	04 .	40	36		: 37	: 40	: 40	. 40		
LIS	Pa,	0	0	0	0	0 0	0	0 :	0 :		4	I	H	I	0	0 0		0	0		0 :		0 :	0 0		
- Se	A	BOOKED	25 :	-	-	22	36	25 :	30 :		42 :	23 :	34:	-	3 100	30.	-		-	-	31	-	-	31	_	
	No.	BOC	17907	18753	18885	19087	19457	17909	19124		3963	4771	15691	20739	16750	10953	199/2	21317	22048		20602	19705	21319	21899	20/22	-

FORCEPS DELIVERIES (continued).

	Failed forceps before admission Manual dilatation of cervix. Blood transfusion.	Morbid.	Failed forceps before admis- sion. Blood transfusion.	Morbid.	Failed forceps before admission. Morbid.	Manual dilatation of cervix. Morbid.	Eclampsia. Eclampsia. Failed forceps before admission	Failed forceps before admission	
	11	SB	Г	D	r	חח	רוט	11	The second
	11	71	J	J	ı	H	777	77	Manage Park
	12½" 13½"	134"	15"	1	1	13"	13"	1444	
	18" : 20" :	: "IZ	81	1	1	"6I	20"	20"	
	0 0Z. 18" 2 0Z. 20"	8 oz 7	6 oz. 18"	2 oz.	II oz.	9 oz. 19"	12 02. 18" 6 02. 20" 14 02. 20"	3 oz. 20" 3 oz. 22	
	6 lb. c	lb.	7 lb. (8 lb.	8 lb, I	. in		. lb.	
	3½ hrs. 6	1 hr. 7	14 hrs. 7	4 hrs. 8	0	½ hr. 5	14 hrs. 7 15 hrs. 7 1 hr. 7	3 hrs. 7	
	51½ hrs. 103 hrs.	120 hrs. 25½ hrs.	36 hrs.	46 hrs.	15½ hrs.	114 hrs. 75 hrs.	5 hrs. 52 hrs. 24 hrs.	17 hrs. 56 hrs.	
	36 Foetal distress 40 Foetal distress	40 Foetal distress 40 Persistent	40 Persistent	40 Persistent	40 Delay in 2nd stage	40 Foetal distress 40 Foetal distress	Ma Ma Pe	Fo	occipito-posterior
	36	. 40	40	: 40	: 40	04 04	335	04 04	
ACX	00	00	0	0	0	00	0 0 н		
GEN	1 35	5 35	2 24	19700 24	20429 26	7 34 9 20	7 28 7 27 27 27 27	1 34	_
EMERGENCY	18731 35 19013 26	19015	19452	19700	2042	20427	20944 21017 21477	21483	

FORCEPS DELIVERIES (continued).

Manual dilatation of cervix.	Manual dilatation of cervix, Morbid.	Manual dilatation of cervix. Morbid. Morbid. Eclamosia Morbid Blood	transfusion. Manual dilatation of o	Failed forceps before admission Antepartum haemorrhage. Eclampsia. Manual dilatation of cervix. Morbid.	Eclampsia. Morbid. Blood transfusion. Morbid.
חו	1 1	L D L	L SB	STITE	TO T
11	ם ם			חחחחח	77 7
132"	14 ½" 14 ½"	14½" 	1 1 1 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4	"4" 13# 14" 41 14" 41 14" 41 14" 41 14 14 14 14 14 14 14 14 14 14 14 14	14"
5 oz. 20" : 12 oz. 18½" :	13 0z. 22" 0 0z. 21"	oz. 22½" : oz. 10" :	02. 20" 02. 20" 02. 21"	0z. 23" 0z. 20" 0z. 19½" 0z. 23" 0z. 23"	to oz. 22" 0 oz. 19"
5 0 21		9 oz. 12 oz. 14 oz.		4 2 0 1 0 2 2 1	
7 lb. 5 lb.		8 lb.	6 lb. 7 lb. 7 lb.	8 lb. 8 lb. 7 lb. 11 lb. 7 lb.	3 lb. 7 lb.
2 hrs.	4½ hrs.	2 hrs.	th. 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	3 hrs	2 hrs. 2 hrs. 2 hrs.
36 hrs.	96 hrs.	96 hrs. 48 hrs.	120 hrs. 96 hrs.	38 hrs. 144 hrs. 29 hrs. 17 hrs. 100 hrs. 53 hrs.	14 hrs. 100 hrs. 57 hrs.
Deep transverse arrest Foetal distress	Maternal and foetal distress Foetal distress	Maternal and foetal distress Delay in 2nd stage Persistent			36 Delay in 2nd stage 40 Persistent occipito-posterior 40 Deep transverse arrest
6 4	40	37	- 6 6	040404	40 40
»°°°	0 0	0 00		000000	00 0
EMERGENCY 21715 27 : 0 21475 29 : 0	0 33	AND REAL PROPERTY.		35 32 32 3 3 4 4 4 4 3 3 4 4 4 4 4 4 4 4 4	2 27 27
EMER 21715 21475	21830	22259	22438 21673	22550 22172 22575 22873 22745 23243	23013 23085 23402

TWINS.

	Remarks	Toxaemia. Blood transfusion. Morbid. Prolapsed cord. Surgical induction. Toxaemia. Morbid. Blood transfusion. Postpartum haemorrhage. Blood transfusion. Morbid. Blood transfusion. Morbid. Blood transfusion. Postnatal admission.
It	2nd	Rundan and a been ben
Result	M rst	PAPAPAPA PAPAPA PAPAPAPA PAPAPAPA PAPAPAPA PAPAPAPAPA PAPAPAPAPAPAPAPAPAPAPAPAPAPAPAPAPAPAPA
	Type	Binovular
		9 02. 8 02. 5 02. 5 02. 7 02. 9 02. 9 02. 9 02. 10 02. 13 02. 10 02. 11 02.
	znd	10. 10 10 10 10 10 10 10 10 10 10 10 10 10
Weight	-	3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3
1	ıst	EE 424210 0 0 2 414 8801
		55 66 66 66 66 66 66 66 66 66 66 66 66 6
	2nd	THEMPHEM TEMME TO MENTAL THE
Sex	-	
	ıst	METERNAM PAFFF MERNTHM
tation	znd	breech vertex vertex vertex vertex breech breech vertex breech vertex breech vertex breech breech breech breech vertex breech vertex breech vertex breech vertex breech vertex
Presentation	ıst	vertex
rity	Matu	38 38 38 38 38 38 38 38 38 38 38 38 38 3
rs	ья	OHH 000000 0 00 00 00 00 00 00 00 00 00 0
e di	3A	KED 34 34 33 34 34 34 34 34 34 34 34 34 34 3
	No.	BOOKED 14322 34 : 2 18179 21 : 1 18887 27 : 3 4011 30 : 1 18956 35 : 1 17647 30 : 3 6823 37 : 1 19301 31 : 2 19301 31 : 2 21198 34 : 7 21788 25 : 0 21788 25 : 0 21788 25 : 0 18574 31 : 0 18655 38 : 5 19464 33 : 2 15019 25 : 0 19946 24 : 0 19946 24 : 0 19946 24 : 0 23375 41 : 1 23398 24 : 1 23398 24 : 1 23316 21 : 0

FACE PRESENTATIONS.

Result Weight of Remarks M C	L : SB 1 lb. 10 oz. Anencephaly. L : L 7 lb. 10 oz. External version. L : L 7 lb. 10 oz. Morbid.	L : L 7 lb. 6 oz. L : SB 7 lb. 7 oz. L : L 3 lb. 0 oz. Antepartum haemorrhage. L : L 8 lb. 10 oz. L : L 7 lb. 7 oz. Blood transfusion.
Treatment	Nil Forceps delivery Nil	Nil Nil Nil Caesarean section
Position	L.M.A. L.M.A. R.M.A.	L.M.A. L.M.A. L.M.A. L.M.A.
Maturity	40 40	40 40 40 40
Para	ннн	0 H D C O
Age	30 23 33	32 32 35 26 30 30
No.	BOOKED 18834 4771 22205	EMERGENCY 19086 32 19185 35 19455 26 20319 30 22410 33

TRANSVERSE LIE.

Remarks	For delivery at home.
Result C	1 : 1 - 1 - 1 - 1 - 1 - 1
Treatment	External version Internal version Internal version
Complications	Nil Placenta praevia Prolapsed arm
Maturity	39 40
Para	000
Age	36 23 29
No.	EMERGENCY 6771 36 21666 23 22008 29

INTERNAL VERSION.

-	Age	Para	Maturity	Indication	Weight of child	Result M C	Remarks
SG	BMERGENCY		1				
-	37	2	40	Prolapsed cord	7 lb. 4 oz.	L : SB	7 lb. 4 oz. L : SB Placenta praevia.
21666	. 23	0	30	Transverse lie	2 lb. 11 oz.	L: L	Placenta praevia.
22008	29	24	40	Transverse lie	Ito lb. 7 oz. L : SB	L : SB	

HYDRAMNIOS.

Remarks	Forceps delivery.	L : - Discharged undelivered.
Result M C	L : L	L: -
Treatment	Artificial rupture of membranes L : L Forceps delivery.	39" Rest, etc.
Girth of abdomen	11	39"
Maturity	36	. 28
Para	н 2	5
Age	2D 24 38	38
No.	BOOKED 20739 18883	EMERGENCY 7513 38

PROLAPSE OF CORD.

Remarks							SB Cord not pulsating on admission.
Result M C		L : SB	L : L	L : SB		L : SB	L : SB
Treatmen		IIN	IIN	Nii		Internal	Version
Size of os when first diagnosed		Full	Full	4 dilated		2 fingers	4 dilated
Complication		NI	Twins—breech	Nil		Placenta praevia	Nil
Maturity		40	38	40		40	34
Para		0	7	.3		64	0
Age	KED	24	.31	90	BENCY	37	26
No.	BOOKED	17912	19301	19299	EMERGENCY	19908	21882

ECLAMPSIA

			20				F	its			Urine	
No.	Age	Para	Maturity	Condition on admission	Before	-	Total	Onset	Albumin on Adm. Dis.		Quantity in first 24 hrs.	
EMER	GEN	CY										
19212	(35 :	3	: 36	Conscious	0	:	I	Antepartum	7 0/00	:	Nil	32 ozs.
15705	21:	I	: 38	Comatose	8	:	31	Antepartum	Solid	:	-	_
20944	27 :	0	: 35	Conscious	0	:	12	Intrapartum	6.5 0/00	:	Nil	40 ozs.
21017	28 :	0	: 38	Drowsy	3	:	3	Antepartum	5 0/00	:	Nil	_
21155	23:	0	: 38	Drowsy	2	:	2	Intrapartum	7.5 0/00	:	Nil	44 ozs.
. 22155	39 :	0	: 40	Comatose	7	:	16	Intrapartum	Solid		-	-
22091	37 :	0	: 37	Comatose	5	:	5	Antepartum	10 0/00	:	Nil	26 ozs.
	:		:			:				1		
22873	24:	0	: 40	Drowsy	2	:	2	Intrapartum	12 0/00	:	Nil	20 OZS.
23013	36 :	0	: 36	Comatose	I	:	2	Intrapartum	-	:	Nil	26 ozs.
23310	23:	0	: 34	Conscious	0	:	1	Postpartum	10 0/00	:	Nil	68 ozs.
	:		:		18 19	:				-		Hame '

Oedema	Highest B.P.	Headache	Eye-signs	Days in hospital before delivery	Type of delivery	Res	sult C	Remarks
Nil	190/144	Slight	No	: 4	Normal	L	: SB	Surgical induction
Nil	170/130	Slight	No	:-	Normal	D:	SB	
Generalised	194/130	No .	No	: 3	Forceps	L	D	Surgical induction
Feet	180/118	Yes	No	: 3	Forceps	L:	L	Morbid.
Slight	150/120	No	Yes	: I	Normal	L:	L	
Moderate	160/110	_	-	:-		D:	_	Died undelivered.
Marked	180/120	No	No	: 4	Forceps	L :	L	Twins. Morbid.
Generalised	134/80	No	No	: 0	Forceps	L:	L	
Generalised	188/132	No	No	: 1	Forceps	L:	L	
Generalised	200/100	No	No	: 5	Normal	L	D	Retinal detachment.

POSTPARTUM HAEMORRHAGE.

Remarks	Morbid.	Morbid.	Morbid. Forceps delivery. Morbid.
Result M C	77 77 777 777 777 777 777 777 777 777		11 11 1
Treatment	Blood transfusion Blood transfusion Blood transfusion Manual removal of placenta. Blood transfusion Blood transfusion Blood transfusion Manual removal of placenta. Blood transfusion	Manual removal of placenta. Blood transfusion Blood transfusion Blood transfusion	Blood transfusion Drugs and massage Drugs, etc. Blood transfusion Drugs, etc. Blood transfusion Drugs, etc. Blood transfusion
Predisposing cause	Nil Twins Antepartum haemorrhage Nil Nil Nil Nil	Nil Nil Nil	Antepartum haemorrhage Nil Placenta praevia Nil
Character of labour	Normal Normal Normal Normal Normal Normal	Normal Normal Normal	40 Normal twins 36 Normal 40 Prolonged : 40 Normal
Waturity.	0 0 0 0 0 0 0 0 0 0	04 04	38 38 0 0 4 0 4 0 0 0 0
Para	ен оо оно н	4 H	٥٠ ٥٠ ٥
98v	KED 23: 35: 35: 30: 30: 25: 25: 28: 28: 31: 23: 23: 23: 23: 23: 23: 23: 23: 23: 23	36:	22 24 24 20 33 33 17 17 17
	BOOKED 17451 23: 18956 35: 19462 32: 19124 30: 20874 25: 21021 28: 18882 31: 22063 23:	22277 23039	EMERGENCY 19916 22 : 0 19946 24 : 0 20013 20 : 0 21830 33 : 0 22465 17 : 0

MANUAL REMOVAL OF PLACENTA.

Remarks	Blood transfusion. Blood transfusion. Blood transfusion. Blood transfusion.	Postnatal admission. Blood transfusion. Postnatal admission. Blood transfusion. Postnatal admission. Blood transfusion. Postnatal admission.
Result M C	11111 11111 8	L SB L L L L L L
Morbidity	Nil Nil Nil Nil	Nil Nil Nil 3 days
Indication	Haemorrhage Haemorrhage Haemorrhage Delay Delay	Delay Delay Delay Delay
3rd stage	4 hrs. 5 hrs. 2 hrs. 2½ hrs. 2½ hrs.	4½ hrs. 8 hrs. 12 hrs. 15 hrs.
Labour	40 Forceps delivery 40 Normal 40 Normal 40 Normal	40 Normal 40 Normal 40 Forceps delivery 40 Forceps delivery
Maturity	044444	0 0 0 0 0
Para	00000	CY CY
ogA	BOOKED 8802 19: 3124 30: 8882 31: 777 36: 7238 42:	GEN 42 42 22 22 22 25 25 25
°Z	B00 18802 19124 18882 11737 23238	EMERGENCY 19996 42: 11 21111 22: 0 3639 32: 1 21832 25: 0

BLOOD TRANSFUSION.

Remarks		Manual removal of placenta. Twins.	Ruptured uterus. Caesarean section. Antepartum haemorrhage. Forceps delivery. Manual removal of placenta.	Pulmonary tuberculosis.	At delivery and 4 days later.	Caesarean section.	Caesarean section.	Caesarean section.	Erythoblastosis foetalis. Erythoblastosis foetalis.
C		L SB	94444	144	111	SB 7	111	HHA	77
Result		חחחח	ВРРРР	חחחר	ייייי	יייי	77-	חחח	11 }
Morbidity		BBBB	Nil Nil 9 days 12 days 63 days	EEE		Nul 12 days Nil	222	Nil 2 days Nil	
Volume of citrated blood given		2 pints 3 pints 2 pints 2 pints	I pint 3 pints 5 pints I pint 14 pints	4 pints 2 pints 2 pints		4 pints 3 pints 3 pints	3 pints 6 pints r pint		150 ccs. 90 ccs. 100 ccs.
Indication		Postpartum shock Postpartum haemorrhage Antepartum haemorrhage Postpartum haemorrhage	Shock Hb. 43% Postpartum haemorrhage Hb. 60% Postpartum haemorrhage			Postpartum Hb. 52% Hb. 36%	Postpartum haemorrhage Placenta praevia Postnartum haemorrhage		Hb. 49% Hb. 76% Hb. 62%
Maturity		40 33 40	044444	26 40		04 04 04	. 37		
Para		3	4+000			000	н н с		
Age	Q	19 33 35	320 320 300 300	37	10 00 CH	19 40 23	23	30,2	ro days 2 days 13 days
No	BOOKED	18802 17451 18595 18956	14825 19421 19462 17909 19124	19983 9595 20071	21021	19878	22063	15879 22277 23039	12535

BLOOD TRANSFUSION (continued).

Remarks		Morbid. Twins, 24th day.	Postnatal admission. 28th day	Acute liver atrophy. Twins.		Caesarean section.	Failed forceps before	admission.		Postnatal admission. Manual	removal of placenta.
Result C		1):	 1	0	11				: L		J
M M		1	1	1	1-	111	H	1	11	ר	1
Morbidity		Nil	Nil	Nil	9 days	2 days	EE	Nil	ZZ	Nil	Nil
Volume of citrated blood given		2 pints	3 pints	6 pints	3 pints	4 pints	2 pints r pint	2 pints	3 pints 2 pints	4 pints	.5 pints
Indication		Hb. 47%	Hb. 28%	Placenta praevia	Hb. 45%		Accidental haemorrhage Shock			Shock	38 Antepartum haemorrhage Hb. 38%
Maturity		38	40	34	40	36	36	40	40	40	38
Para	CZ	0	0	5	0 4	0 4 m	000	н	0 7	: II	c
98A	BEN	31 :	34 :	38	26:	31:	32 :	33 :	37:	42 :	24:
No.	EMERGENCY	18574 31	18600	18655	19013	18994	14912	7204	80661	9666:	He

BLOOD TRANSFUSION (continued).

	L : L D : SB L : [T	L : L Postnatal admission. 2 pints	L : L Postnatal admission. 2 pints	L : L Forceps delivery. 3 pints on	L : L Forceps delivery. 2 pints on		L : L Caesarean section.	L : L Caesarean section.	SB For	L : SB Forceps delivery.
	3 days I Nii 14 days I	NII	NIII I	8 days I	15 days I	NII			ys	2 days I
100	8 pints 2 pints 6 pints	4 pints	4 pints	5 pints	4 pints	2 pints 2 pints	2 pints		2 pints 4 pints	3 pints 2 pints
	Placenta praevia Anaemia, Hb. 20% Anaemia, Hb. 28%	40 Retained placenta	Retained placenta	40 Postpartum haemorrhage	Hb. 54%	Hb. 38% Hb. 56%		Fostpartum naemonnage Hb. 54%		40 Prolonged labour 32 Antepartum haemorrhage
	368	. 40	. 40	. 40	. 40	: 40		. 40	: 37	. 32
SNCY	0 : 0 2 2 5 : 3	22 : 0	2 : I	33: 0	29: 0	37: 12	3:0	0 : 6	0 · H	27: 0
EMERGENCY	20113 20275 14653 3	211112	3639 32	21830 3.	21782 2	3553 3 22091 3	22410 3	22405 1	23014 2	23085 2 23502 3

PERINEAL TEARS.

Perineal tears following delivery occurred as under :-

Booked primigravidae

Emergency multiparae

100 in 244 vaginal deliveries 41%.

Booked multiparae 62 in 328 vaginal deliveries 19%.

Emergency primigravidae 68 in 140 vaginal deliveries 48%.

17 in 93 vaginal deliveries 17%.

247 in 805 vaginal deliveries

MORBIDITY OF PATIENTS.

The following table gives details of all cases which were notified as morbid in the wards. A patient is considered morbid when the temperature of 100.4 is maintained for a full twenty-four hours or recurs within twenty-four hours. A catheter specimen of urine and a high vaginal swab are taken on all cases which have a temperature over 100, or in which a temperature of 99.4 occurs on two successive days.

Morbidity of booked cases

Deaths not included below Pyrexia	1 21 — 22 —	5.5%
Morbidity of emergency cases		
Deaths not included below Pyrexia 'Morbidity of all deliveries	6 26 — 32 —	13.4%
Deaths not included below Pyrexia	7 57 64 —	7.8%

In addition one booked case and one emergency case were admitted after delivery and were morbid in the puerperium.

MORBIDITY OF PATIENTS DELIVERED IN HOSPITAL.

Remarks			Ante & postpartum haemorrhage. Blood transfusion. Manual removal of placenta. Blood transfusion.
C		<u>нананар</u>	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
Result		חחחחחח	חחחחח
Cause of temperature		Mastitis Unknown Unknown Mastitis Pyelitis Breast abscess Pneumonia	Mastitis Anaemia Sepsis Unknown Pyelitis Parametritis
Duration of temperature		3 days 2 days I day 2 days 9 days I day 6 days	2 days 2 days 2 days 3 days 23 days 63 days
Day when temperature reached morbidity		######### ############################	8th 4th 2nd 2nd 1st
Method of delivery		Normal Normal Normal Normal Normal Normal	Normal Normal Normal Forceps Forceps
Maturity		04444448 0400000	444444
Para	0	0.6000 мн	000000
93V	воокер	24 26 20 23 20 37 37	24 24 20 20 30
, N	BOC	17659 7654 18681 17149 17736 18589 6823	18174 19462 17277 19533 17909 19124

MORBIDITY OF PATIENTS DELIVERED IN HOSPITAL (continued).

	L Postnatal admission.	ـ د د د	111-	SB Blood transfusion. L Diabetes mellitus.		Blood transfusion.
	니니	ННН	HHHH	1777	1111	
	Cystitis Pulmonary tuberculosis	Pneumonia Mastitis Sepsis	Sepsis Sepsis Sepsis Macritic	Sepsis & pyelitis Pyelitis Pyelitis	Unknown Mastitis Mastitis	Pyentis Sepsis Bronchitis
	5 days 4 days			days days days	4 days 2 days 5 days 6 days	2 days 2 days 19 days
	5th 4th	rst r3th 2nd	3rd 6th	7th	344 844 844 844 844 844 844 844 844 844	4th 2nd 3rd
	Normal Normal	Normal Normal Forceps	Forceps Forceps Normal	Caesarean section Normal Normal		Forceps Normal Normal
	. 40	40	0 0 0 0 c	37	5 4 4 4	04 04
0	7 : T	H H B	0000	0 7 1 1	 9	0 4 0
BOOKED	9595 37	20109 3 20462 2, 20739 2.	18750 23 18953 38 18594 20	-	19789 28 17830 28 20879 19	22766 20 22277 30 7375 17

MORBIDITY OF PATIENTS DELIVERED IN HOSPITAL (continued).

EMERGENCY	GEN	CY	-						
18574	31:	0	38	Normal breech	7th	17 days	Pyelitis and	L: (L	Blood transfusion.
18771	28:	00	40	ZZ	4th 15th	2 days 2 days	Unknown	רר הרה	Postnatal admission. Blood
19013	26:	0	40	Forceps delivery	2nd		Sepsis	1. 1.	Blood transfusion.
18994	31 :	о н	36	Caesarean section Face delivery	3rd 6th	2 days	Sepsis	L : SB	Blood transitusion.
19211	24:	. 0	40	Normal delivery	4th		Sepsis		
19015	35 :	0 0	40	Forceps delivery	54	z days	Pyelitis	1	
20113	20 :			-	3rd		Sepsis		Placenta praevia.
20429	: 92	: 0			2nd		Pyelitis		Failed forceps before admission.
20424	20:	0.0	40	Normal delivery	rith	2 days	Mastrus	ייי	Pulmonary tuberculosis.
14653	35 :		36	_	ıst	20 days	Pyelitis	•••	Blood transfusions.
18699	20:		40		ıst		Pyelitis		
21017	28:	. 0	38		znd	4 days	Pyelitis		Eclampsia.
21830	33 :		40	Forceps delivery	3rd	s days	Pyentis		Blood transfusions
22250	. 67) H	40	Forceps delivery	2nd		Sepsis		Diod dansiusions.
22091	37:		37		3rd		Pyelitis		Eclampsia. Blood transfusion.
22567	38:		26	Inevitable	6th	3 days	Pyelitis	T : T	
22745	41	0	40	abortion Forceps delivery	7th	4 days	Pyelitis & septic	L : SB	Blood transfusion.
23189	26:	0	40	Normal delivery	6th	2 days	Perineum Sepsis		
23085	-	. 0	40		ıst	2 days	Sepsis		
23402	27:	. 0	40	Forceps	5th	3 days	Pyelitis		
23182	33.	и н	40	Normal delivery	r4tn 2nd	2 days	Pyelitis	ייי	
,	-		-2						
					-			-	

MATERNAL DEATHS.

Booked.

14825. Aged 42, para 4. Admitted at the 39th week for mild toxaemia of pregnancy. Medical induction at term and rapid delivery, the labour lasting about three-quarters of an hour. Some slight bleeding immediately after delivery but the patient collapsed, became restless, pale, with evidence of severe haemorrhage. Blood transfusion immediately commenced and there was some slight improvement in the general condition a half-hour later but she had a further collapse and died 40 minutes after delivery. Post-mortem showed a tear in the right side of the uterus extending to the cervix with a large sub-peritoneal haemorrhage in the right broad ligament.

Emergency.

12432. Aged 35, para 1. Admitted at the 32nd week with a history of slight antepartum haemorrhage. Her general condition which was poor remained about the same for three weeks when she had a further slight loss, this was repeated the following day when she collapsed and died an hour later. Post-mortem showed a bilateral hydronephrosis and a large chronic tubo-ovarian abscess. There was also a complete placenta praevia.

15705. Aged 21, para 1. Admitted early in labour after an eclamptic fit. Blood pressure was, 170/140 and the urine almost solid with albumen. She delivered herself normally 8 hours after admission of a premature stillborn child and died 5 hours later.

Post-mortem showed acute liver atrophy.

20275. Aged 30, para 2. Admitted at the 37th week with severe anaemia, R.B.C. 1,300,000, haemoglobin under 20%. She was given a 2 pint blood transfusion and had some generalised respiratory disturbance after 2 pints had been infused. The condition improved somewhat over the next 12 hours and she expelled a premature macerated foetus the following day. Her respirations became shallow and laboured some two hours later and she died 10 hours after delivery. Post-mortem showed oedema of all organs with some generalised oedema and oedema of the lungs.

21571. Aged 38, para o. Admitted at the 20th week with sickness and vomiting after all meals and pain in the back radiating around to the chest. She had had a radical mastectomy performed 2 months previously and had had three weeks deep X-Ray therapy. Her condition deteriorated, she became drowsy and uraemic and died in coma 24 days after admission. Post-

mortem refused.

22155. Aged 39, para o. Admitted at approximately term with a history of six eclamptic fits. She had eight further fits after admission and died undelivered five hours later. Post-mortem

showed liver atrophy.

22282. Aged 43, para o. Admitted at the 38th week for mild toxaemia which was adequately controlled by rest. She delivered herself normally and died the following day. Post-mortem showed pulmonary embolus.

PUERPERAL INSANITY

7560. Booked case, aged 27, para 1. Some albuminuria and hypertension with oedema for the last four weeks of her pregnancy. Refused admission for rest during this period. Normal delivery at term of a 6lb. baby. Became restless and confused on the sixth day with typical confusional insanity and after attempts to leave the hospital was discharged home to her husband on his responsibility. Subsequently transferred to a mental hospital for observation.

INFANT REPORT.

During the year there were 841 babies born.

Details of all premature babies born alive $(5\frac{1}{2})$ lbs. weight or less) and those admitted after delivery outside are appended in the following table. There were 25 deaths of premature babies out of a total of 84 premature babies.

PREMATURE BABIES.

Remarks	Died 4th day. Died 2nd day. Morbid. Postnatal admission.
Method of feeding Breast : Bottle	rly: 4 hourly: 4 hourly: 4 hourly: 14 hourly: 15 hourly: 17 hourly: 18 hourly: 19 hourly
Method Breast	4 hourly 4 hourly 4 hourly 4 hourly 5 hourly 6 hourly 7 hourly 8 hourly 8 hourly 9 hourly 1 hourly 1 hourly 1 hourly 1 hourly 2 hourly 4 hourly
Days in hospital	26 26 44 43 44 15 17 17 30 30 15
Weight on discharge	4 lb. 12 oz. 4 lb. 14 oz. 4 lb. 14 oz. 3 lb. 15 oz. 4 lb. 2 oz. 5 lb. 1 oz. 5 lb. 4 oz. 5 lb. 5 oz. 5 lb. 64 oz. 5 lb. 64 oz. 5 lb. 13 oz. 5 lb. 2 oz.
Birth	5 lb. 3 oz. 5 lb. 3 oz. 5 lb. 3 oz. 3 lb. 14 oz. 4 lb. 15 oz. 5 lb. 18 oz. 5 lb. 19 oz. 6 lb. 19 oz. 6 lb. 19 oz.
Cause of premature birth	(1st twin 1st twin 2nd twin 2nd twin 2nd twin
Cause	Twins. Unknown Unknown Unknown Unknown Unknown Unknown Unknown Toxaemia Toxaemia Toxaemia
Maturity	4 % 6 4 4 6 6 6 6 8 6 6 8 6 8 6 8 6 8 6 8 6
Mother's pregnancies	н оно во оно н м м м м м м м м м м м м м м м м
Mother's	2I
No.	BOOKED 18179 18283 18879 18887 17519 19241 17023 18880 18524 6823

PREMATURE BABIES (continued).

4 hourly 4 hourly — Died 3rd day. — Died 3rd by parents	Postnatal admission. Forceps delivery. Died 12th day. W.R.+ Died 2nd day. Discharged herself.
4 hourly: 4 hourly: 4 hourly: 3 hourly: ————————————————————————————————————	4 hourly: 3 hourly 4 hourly: — 4 hourly: — 3 hourly: 3 hourly 6 — — — — — — — — — — — — — — — — — — —
38 16 16 14 14 13 13 13 13 13 13 13 13 13 13 13 13 13	17 13 14 17 17 17 17 17 17 17 17 17 17 17 17 17
5 lb. 6 oz. 5 lb. 7 oz. 5 lb. 7 oz. 3 lb. 12 oz. 5 lb. 3 oz. 5 lb. 3 oz. 5 lb. 4 oz.	5 lb. 15 lb. 16 lb. 15 lb. 16 lb. 15 lb. 16 lb. 16 lb. 17 lb. 18
13 02. 13 02. 15 02. 15 02. 16 02. 17 02. 18 02.	15. 12. 12. 12. 13. 14. 15. 15. 15. 15. 15. 15. 15. 15. 15. 15
Unknown Toxaemia Unknown Induction for pulmonary tuberculosis Induction for toxaemia Twins Unknown Inknown	for essential rtension and twin praevia pric toxaemia
24 0 40 27 0 36 27 0 40 24 1 35 32 1 36 32 1 36 32 3 40	- 1000000000000000000000000000000000000
18159 19962 19235 20326 20473 8008 21201	21198 19138 22009 22048 20152 20152 22573 15879 16367 11528

PREMATURE BABIES (continued).

	Died 6th day. Died 25th day. Died 6 hours. Died 8th day. Died 3 hours. Died 3 hours. Died 4 hours.	Died 20 hours.
	4 hourly	4 hourly: — 4 hourly: — 4 hourly: —
	25 6 hours 39 11 21 28 8 8 8 26 3 hours 3 hours 17 17	17 17 63 1
	6.55 1 1 1 1 1 1 1 1 1	4 lb. 6 oz. 4 lb. 10½oz. 4 lb. 13 oz.
	16. 15 02. 16. 15 02. 16. 16. 16. 16. 16. 16. 16. 16. 16. 16	120
	haemorrhage svia. 1st t znd t toxaemia toxaemia toxaemia toxaemia toge znd toxaemia toge znd toxaemia toxaemia toxaemia toxaemia toxaemia	Pre-eclamptic 1st twin toxaemia 2nd twin Antepartum haemorrhage Antepartum haemorrhage
	38 4 58 88 8 8 8 8 8 8 8 8 8 8 8 8 8 8 8	33 38
CY	33 33 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	26 0
EMERGENCY	18576 18627 18627 18006 19085 18913 18939 18970 19464 15019	19946 19455 20073

PREMATURE BABIES (continued).

	Died zoth day. Died znd day. Died 5 minutes. Died 8 hours. Died 6th day. Postnatal admission. Died 20 hours. Died 2nd day. Died 2nd day. Died 3th day. Died 1st day] Postnatal Died 1st day. Died 2nd day.	
	3 hourly	
	2 E E E I 2 0 0 0 E E E E I 2 T T T E E E E E E E E E E E E E E E E	
	4 Hb. 15 02. 4 Hb. 15 02. 5 Hb. 16 02. 5 Hb. 17 02. 6 Hb. 17 02. 7 Hb. 1 0 02. 8 Hb. 1 0 02. 9 Hb. 1 0 02. 1 Hb. 2 02. 1 Hb. 2 02. 1 Hb. 8 02.	
	5 b. 2 oz. 5 b. 1 b. 12 oz. 5 b. 14 b. 14 oz. 5 b. 16 b. 17 oz. 5 b. 16 b. 17 oz. 5 b. 16 b. 17 oz. 6 b. 17 oz. 6 b. 18 oz. 6 oz. 6 b. 18 oz. 6 oz. 6 b. 18 oz. 7 c. 7	
	Antepartum haemorrhage Antepartum haemorrhage Eclampsia Eclampsia Unknown Unknown Pre-eclamptic toxaemia Unknown Pre-eclamptic toxaemia Essential hypertension Unknown Unknown Pre-eclamptic toxaemia Essential hypertension Unknown Tre-eclamptic toxaemia Essential hypertension Unknown Induction for infantile paralysis Acute liver atrophy Pre-eclamptic toxaemia Antepartum haemorrhage Pre-eclamptic toxaemia Eclampsia Twins Ist twin Twins Ist twin Twins Ist twin Twins Ist twin Tre-eclamptic toxaemia Ist twin Twins Ist twin Tre-eclamptic toxaemia Ist twin Tre-eclamptic toxaemia Ist twin Induction Ist twin Induction Ist twin Ist twin Ist twin Ist twin Induction Ist twin Ist twi	
	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	
VCY	23 24 23 23 24 24 24 24 24 24 24 24 24 24 24 24 24	
EMERGENCY	20791 20869 21155 20944 21392 21408 21588 21679 21679 21679 21679 21747 222474 21629 222777 22747 22747 23310 23310	

STILLBIRTHS.

There were 58 stillbirths during the year.

per 1,000 total births

	A 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
Booked primiparae	12 per 246 total births = 48.8	
Booked multiparae	12 per 334 total births = 35.9	
Emergency primiparae	17 per 102 total births = 166.7	
Emergency multiparae	17 per 97 total births = 175.2	

STILLBIRTHS.

Remarks	Prolapsed cord. External version one week previously.
Weight	5 lb. 9 oz. 4 lb. 10 oz. 5 lb. 10 oz. 5 lb. 10 oz. 6 lb. 11 oz. 6 lb. 11 oz. 6 lb. 11 oz. 7 lb. 4 oz. 2 lb. 4 oz. 2 lb. 7 oz. 7 lb. 13 oz. 7 lb. 14 oz. 2 lb. 4 oz.
Cause of stillbirth	Tentorial tear Placental infarcation Anencephaly Placental separation Tentorial tear Asphyxia Placental separation Prolonged labour Prolapsed cord Unknown
Maternal complications	Twins Syphilis Caesarean section scar Accidental haemorrhage Nil
Method of delivery	Normal breech Normal face Normal face Normal breech Normal breech Forceps Normal
Mother's age Mother's previous previous pregnancies	BOOKED 322 34 2 324 34 6 334 30 1 5595 33 2 5979 24 6 9779 26 0 9979 26 0 9979 26 0 878 42 4 878 40 3 878 40 0 653 34 0 876 47 0 878 40 0 878 40 0 878 40 0 878 40 0 878 44 0 705 38 2 229 30 5 238 2 0 222 38 0 222 38 0
No.	BOC 14322 19016 18834 18595 17912 19979 19038 19299 19038 19795 22623 22623 22825 22825 22825 22478 19705 22478 17739

STILLBIRTHS (continued).

	Prolapsed cord. Failed forceps before admission. Failed forceps before delivery.	
	140 18W 00 124 H 40 11 18 10 41 0 41 4 1	10 0Z. 10 0Z. 0 0Z.
	444 778 889 247 87 88 89 87 87 88 89 87 87 88 89 89 89 89 89 89 89 89 89 89 89 89	4 6 E E
	Hydrops foetalis Placental infarcation Placental separation Placental infarcation Placental infarcation Placental separation Placental infarcation Placental infarcation Placental infarcation Hydrocephaly Hydrocephalos Placental infarcation Placental infarcation Placental separation Multiple abnormalities Tentorial tear Placental infarcation Unknown Asphyxia Tentorial tears Placental infarcation Unknown Placental infarcation Unknown Placental infarcation	
	Nil Nil Accidental haemorrhage Nil Nil Nil Nil Nil Nil Pre-eclamptic toxaemia Nil	Nil Toxaemia Antepartum haemorrhage
	breech delivery breech breech breech breech breech	delivery
	Normal	Forceps Normal Normal
	4400H00WH012H044400H 0HH04000H0	0000
ENCY	3,20,03,30,00,00,00,00,00,00,00,00,00,00,00	3 4 4 2 8
EMERGENCY	18949 12010 18873 19014 19185 19422 19422 19422 19422 19422 19422 19422 19422 19422 19422 19524 20378 20378 20378 21961 21081 21081 21081 21081 21081 21081 21081 21081 21081 21081 21081 21081 21081 21081 21081 21081 21081 21081	23014 22745 23433 23602

NEONATAL DEATHS.

There were 45 neonatal deaths including 3 babies admitted after delivery. Neonatal death rate 42 in 783 live births = 53.7 per 1,000 live births.

Remarks		Haemangioma of liver, Tracheo-oesophageal	
Cause of death		Prematurity Prematurity Pneumonia	Pneumonia Malformation of heart Congenital abnormalities Atelectasis Prematurity Spina bifida Inhaled vomit Spina bifida Syphilis Prematurity
Age at death		4 days I day 3 days	14 days 7 days 13 hours 22 hours 3 days 2 days 19 days 10 days 12 days 2 days 2 days
Birth		3 lb. 14 oz. 4 lb. 8 oz. 5 lb. 14 oz.	5 lb. 11 02. 5 lb. 11 02. 7 lb. 5 02. 6 lb. 2 02. 5 lb. 1 02. 7 lb. 1 02. 7 lb. 1 02. 7 lb. 6 02. 4 lb. 15 02.
Method of delivery		Normal Normal Normal	Normal Normal Normal Normal Normal Normal Caesarean section
Maternal		Nil Nil Mitral stenosis	Toxaemia Nil Nil Nil Pulmonary tuberculosis Nil Nil W.R.+ Placenta praevia
Mother's parity		1 OI	омооноомоо
Mother's age	KED	42 42 211	36 25 26 27 27 28 32 33 33 33
No.	BOOKED	18879 18997 18018	19045 19704 17277 18170 20326 18525 18951 18674 22573

NEONATAL DEATHS (continued).

		Postnatal admission.	Postnatal admission. Mother rhesus positive	
	- 0 TT	Tentorial tear Tentorial tear Duodenal atresia Cerebral haemorrhage Atelectasis Prematurity Prematurity Atelectasis Broncho-pneumonia	Icterus gravis Tentorial tear Prematurity Prematurity Pneumonia Meningitis.	Prematurity Tentorial tear Intraventricular haemorrhage Tentorial tear
	6 days 25 days 6 hours 9 days 10 days 3 hours 3 days 4 hours 3 days	2 days I day 6 days II day 7 days 7 days 7 days 8 minutes 8 hours 12 hours 6 days	5 days 4 days 7 day 6 hours 2 days 8 days	
	40000000404010	12 02. 12 02. 12 02. 13 02. 14 02. 11 02. 10 02.		5 5
		7 + 0 + 7 + 1 × 2 × 2 × 4 × 4 × 4 × 4 × 4 × 4 × 4 × 4		3 8 8 7 7 B.
	breech breech breech	breech	breech	breech
	Normal Normal Normal Normal Normal Normal Normal Normal Normal	Normal Normal Forceps Normal Normal Normal Forceps	Normal Normal Normal Normal Normal	Normal Forceps Normal
	Antepartum haemorrhage Placenta praevia Nil Nil Toxaemia Antepartum haemorrhage Toxaemia Antepartum haemorrhage Nil	Antepartum haemorrhage Femoral thrombosis Eclampsia Toxaemia Nil Nil Nil Nil Nil Nil Infantile paralysis	Nil Toxaemia Antepartum haemorrhage Toxaemia Twins	Nil Eclampsia Nil
	000 0000 000	000000moH	W 4400 H	000
BENCY	36 21 30 24 24 24 33 33 34 35	33 27 33 39 39 39 39 39 39 39 39 39 39 39 39	22 41 230 32	27 23 39
EMERGENCY	18627 18655 19047 19189 19304 19464 19538 15019	20073 20073 20073 20074 20732 21382 21588 21588 21679 21679	22935 23277 23086 23375	23085 23310 23410

OPHTHALMIA NEONATORUM.

No case of ophthalmia neonatorum occurred during the year.

