Notes on preventive medicine for medical officers, United States Navy : instructions to medical officers / Department of the Navy, Bureau of Medicine and Surgery.

## Contributors

United States. Navy Department. Bureau of Medicine and Surgery.

## **Publication/Creation**

Washington : Government Printing Office, 1918.

### **Persistent URL**

https://wellcomecollection.org/works/rytfprq5

## License and attribution

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org CONFIDENTIAL.

BULLETIN NO. 50. (Division of Sanitation.)

INSTITUTION.

LIBRARY

WelMOmec

[M. & S. No. 129655-

WELLCOME

LIBRARY

**General** Collections

015

## Notes on Preventive Medicine for Medical Officers, United States Navy.

#### INSTRUCTIONS TO MEDICAL OFFICERS.

DEPARTMENT OF THE NAVY, BUREAU OF MEDICINE AND SURGERY, Washington, D. C., November 9, 1918.

#### NOTES ON THE PRESENT EPIDEMIC OF INFLUENZA.

Up to and including November 9, 1918, there have been reported from all shore stations in the United States 54,585 cases of influenza with 2,795 deaths. For the week ended November 9, 1918, there were 1,295 cases reported from shore stations as against 1,793 cases reported during the week ended November 2, 1918. Except for four new cases which developed in the Pacific Fleet no recent reports have been received from other ships. The total number of cases so far reported from the force afloat is 8,564 with 332 deaths, making a grand total for ships and shore stations of 63,149 cases of influenza with 3,127 deaths.

The following table furnishes the latest information of a statistical nature:

Automatica	inf	Total luenza to date.	New cases week ended Nov. 9.	New cases week ended Nov. 2.	Total deaths.	Case fatality rate per cent.
Nind mand district	37	5,235	136	267	270	5, 10
Third naval district	131		100	128	108	4. 13
faring harmole Quantice Variation Variation Variation (Variation Variation)	198	2,615 2,802	284	195	100	3.7
darine barracks, Quantico, Va	111	2, 002	201	190	101	0.11
sixth naval district, exclusive of training camp, Charleston,	10	905	48	19		- DIMENSION
S. C., and marine barracks, Paris Island, S. C	441		40	12 74	64	3.19
Naval training camp, Charleston, S. C	137	1,100.	146	283	19	2.43
Iarine barracks, Paris Island, S. C.	12	781	140	200	10	4. 9
Eighth naval district, exclusive of naval station, New Orleans,	18			5 1 200 m		
naval station, Pensacola, and naval training camp, Gulfport,	182	200	0 0,0	33	1	2. 3:
Miss.	1	300	6	11	34	3. 3
Vaval station, New Orleans, La		1,012	H	20	24	
aval station, Pensacola, Fla		1,474	5	20	24 20	1.6
aval training camp, Gulfport, Miss	Pro-	875	and a state	3.	20	2. 2
welfth naval district, exclusive of naval training camp, San	12			10109 10000		IN SALES
Diego, Cal., naval station, Mare Island, Cal., and naval	12.		101	004	00	0
training station, San Francisco, Cal		1, 553	121	284	39	2.5
Naval training camp, San Diego, Cal	120	1,087		74	6	. 5
Vaval station, Mare Island, Cal		1,415	123	144	71	5.0
Vaval training station, San Francisco, Cal		0				
Naval training station, San Francisco, Cal Other shore stations		33, 431	338	- 365-		6. 00
the second s	-		1 005	1 700	0 705	- 17
Total, shore stations		54, 585	1, 295	1,793	2,795	5.1
ships	-	8,564			332	3. 88
Grand total	-	63, 149			3,127	4. 9

The following figures relative to the results obtained after the use of influenza vaccine at the marine barracks, Paris Island, S. C., are furnished by the senior medical officer at the marine barracks, Quantico, Va.: A draft of 756 men was received from Paris Island, in the early part of November. Of these, 304 had not been vaccinated against influenza, and 39 or 12.8 per cent of them have

since contracted the disease. Four hundred and fifty-two men of this draft had received from one to four inoculations of influenza vaccine and 72 or 15.9 per cent afterwards contracted influenza as follows:

After one inoculation, 11 out of 75 men (14.6 per cent) developed influenza. After two inoculations, 30 out of 226 men (13.2 per cent) developed influenza.

After three inoculations, 8 out of 57 men (14 per cent) developed influenza.

After four inoculations, 23 out of 94 men (24.4 per cent) developed influenza.

Deaths (stillbirths excluded) reported during the week ended Nov. 2, 1918, with death rates, in large cities of the United States.

		ed Nov. 2, 18.	191		nding week, average of 3-1917.		Week end	ed Nov. 2, 18.	19	orrespon 17. λ= 9ek, 1913	ding week average o 3-1917.
Cities.	Total deaths.	Annual rate per 1,000.		otal aths.	Annual rate per 1,000.	Cities.	Total deaths.	Annual rate per 1,000.		otal aths.	Annual rate per 1,000.
Albany	184	90.0	C	36	17.6	Newark	402	50.1	A	98	12.8
Atlanta	79	21.0	č	38	23.4	New Haven	223	76.4	Ö	34	11.0
Baltimore	613	53.8	Ă	176	15.7	New Orleans	455	62.9	A	149	21.2
Birmingham	123	33.8	A	54	16.3	New York	5, 819	52.9		1,209	11.
Boston	319	21.7	A	215	15.0	Oakland	279	70.5	A	37	10.0
Buffalo	613	67.2	A	107	12.1	Omaha	128	37.5	C	27	8.
lambridge	42	19.2	A	28	12.9	Philadelphia	1,771	53.2		466	14.
Chicago	2, 191	44.9	A	576	12.3	Pittsburgh	1, 132	100.7	C	205	18.
Sincinnati	382	48.1	C	119	15.0	Portland, Oreg	213	36.0	C	43	7.
Neveland		65.6	C	163	12.3	Providence	201	40.3	C	45	9.
Columbus		32.9	C	58	13.7	Richmond	100	32.9	C	51	16.
Dayton	148	59.9	A	31	12.7	Rochester	254	50.0	C	76	15.
Denver	155	30.1	A	60	12.4	St. Louis	426	28.9	C	227	15.
Fall River	69	27.7	C	39	15.7	St. Paul	149	30.8	C	39	8.
Frand Rapids	49	19.2	C	29	11.4	San Francisco	804	89.0	C	178	19.
ndianapolis	169	31.1	C	96	17.7	Seattle	148	21.1	A	46	7.
Kansas City, Mo		49.6	C	91	15.5	Spokane	42	13.9	C	26	8.
los Angeles	533	51.9	A	99	11.0	Syracuse	95	31.3	C	36	11.
Louisville	159	34.4	C	38	8.2	Toledo	175	45.2	A	58	16.
femphis		28.7	AC	36	16.6	Washington, D. C	302	42.6	A	116	16.
filwaukee	105 208	36.0		49	16.8	Worcester	102	32.0	C	52	16.
finneapolis	208	24.4	AC	86 69	10.5	Tatal	01 000	10 0	1		
Nashville	203	28.3 41.0	lö	69 42	9.6 18.5	Total	21, 020	48.0			

In the cities listed above there have been 74,232 deaths from influenza and pneumonia during the eight weeks ended November 2, 1918. The figures by weeks are as follows:

Disease.	Sept. 14.	Sept. 21.	Sept. 28.	Oct. 5.	Oct. 12.	Oct. 19.	Oct. 26.	Nov. 2.
Influenza	19	193	904	2, 716	6, 809	12, 289	13, 110	9, 743
Pneumonia	299	405	855	2, 044	4, 606	7, 648	7, 734	4, 858



#### A Report on the Findings in 116 Necropsies Performed in the U.S.S. Mercy on Bodies DEAD OF INFLUENZA.

#### By Lieut. THOMAS L. RAMSEY (Medical Corps), U. S. N. R. F.

The epidemic of influenza started in the Atlantic Fleet early in September. The cases first coming to autopsy were those of simple broncho-pneumonias with a mixed bacteriology. These were characterized by nodular consolidated areas in one or more lobes around central bronchioles which were filled with pus. These areas were surrounded by congested and oedematous lung tissue involving sometimes an entire lobe. At times two or three lobes were involved.

Material taken from affected areas was first streaked on blood agar plates and then smeared on glass slides and stained. Gram stains of the direct smears showed a mixed bacteriology, the pneumococcus predominating. Other bacteria found were staphyolococci, Micrococcus catarrhalis, streptococci, Bacillus mucosus, and influenza bacilli.

The blood agar plates were incubated and colonies studied from time to time as they made their appearance. In this manner the various microorganisms were identified as the colonies became visible.

Colonies identified as pneumococci were fished and transplanted into Avery's rabbitblood broth which after eight hours' incubation was subjected to agglutination tests with specific type sera. In this manner organisms were grouped according to type. Intraperitoneal inoculation of white mice was resorted to when agglutination reactions in the Avery media were obscure. Other organisms were also identified from day to day as they made their appearance on the plates. It was noticed that the fine colonies of influenza bacilli could not be detected until after three or four days' incubation.

This routine procedure of making direct smears from the affected parts and of growing cultures for study was continued on all cases which came to autopsy during the epidemic.

After the epidemic had lasted for a few days it was noticed that broncho-pneumonic areas were increased in extent as compared with earlier cases, that is more lung tissue was involved and hemorrhages into the lung around affected areas were so extensive that whole lobes were filled with a combination of consolidated areas and blood clot. Pleurisy with effusion, and the formation of purulent plastic exudates with adhesions were noted more frequently and pericardial involvement, the result of direct extension, was found in many cases. Cultures now showed the infection to be more truly pneumococcic as pure cultures of this organism were obtained from the infected parts in nearly every case. Cultures from the heart's blood showed that septicemia was present in every case where purulent fluid was found in the chest. This proved to be due to a Type II pneumococcus in every instance. When septicemia cases began to appear autopsy findings indicated that the patients had

succumbed to infection with a much more virulent microorganism and the lung findings were not nearly so pronounced. Some of the cases proved to be hemorrhagic speticemia with no lung involvement other than extensive hemorrhages which sometimes occurred into the lung tissues. The important lesions found in this group of cases were hemorrhages into the lungs, ventricles of the brain or suprarenal glands. Later, cases of meningeal involvement with or without broncho-pneumonia, began to be seen. Many of these cases before death had no signs of meningeal involvement other than those of the accompanying septicemia, but at autopsy they showed a purulent spinal fluid containing pneumococci. From these findings one can readily see that the infection which started as a rather mild complicating broncho-pneumonia rapidly increased in virulence until it reached its climax in the involvement of the blood stream and meninges.

the involvement of the blood stream and meninges.

The cases autopsied were divided into the following groups:

Group I. Simple broncho-pneumonia.

Group II. Lobar pneumonia.

Group III. Massive broncho-pneumonia.

Group IV. Hemorrhagic broncho-pneumonia.

Group V. Broncho-pneumonia with septicemia.

Group VI. Broncho-pneumonia with meningitis.

Group VII. Hemorrhagic septicemia.

Group VIII. Pneumococcus meningitis.

In 9 of the 116 cases no organisms were recovered, because the bodies had been kept in the mortuary ice box before coming to autopsy.

The pathology of these groupings is submitted separately in order to bring out more clearly the differential findings.

#### GROUP I. SIMPLE BRONCHO-PNEUMONIA (8 CASES).

Pathology.—The findings are those of broncho-pneumonia; namely, one or more lobes showing areas of consolida-tion around the smaller bronchioles which were filled with pus, the air spaces themselves being filled with purulent material. In the early stages of the disease the consolidated area was reddish to reddish brown, turning gray as resolution set in. The surrounding lung tissue was congested and oedematous sometimes extending to other consolidated areas in the same lobe, thus simulating consolidation of the entire lobe. The heart was rarely involved in this type of cases except when acutely dilated as a terminal process. Cultures from the heart's blood were negative. There were eight cases in this group, and pneumococci were found in lung cultures from seven. In one case the bacteriology was not determined.

The types of pneumococci found were: Type I, 3 cases; Type II, 2 cases; type not determined, 2 cases. Associated with pneumococci there were found stapholococci, streptococci, B. influenze, B. mucosus, and M. catarrhalis, indicating a very mixed bacteriology in this group.

#### GROUP II. LOBAR PNEUMONIA (7 CASES).

Pathology.—The consolidation found in this group was typically that of true lobar pneumonia, the lung involve-ment being usually unilateral, with no hemorrhages into the lung tissue, no complicating pleurisy except of a dry fibrous character which was found in one case. There was no myocardial involvement, although acute dilatation of the heart was found in two cases with extensive lung involvement. The heart's blood was negative for bacteria.

Direct smears from lungs were positive for pneumococci in five cases. Type II pneumococcus was found in one case. Pneumococci in the other four cases were not typed. Staphylo-cocci, B. influenzæ and B. mucosus were found associated in several of the cases. The bacteriology in two instances was not determined.

#### GROUP III. MASSIVE BRONCHO-PNEUMONIA (33 CASES),

Pathology.-That of a very extensive broncho-pneumonia where the areas of consolidation around the pus filled bronchioles were so numerous and adjacent as to cause consolidation of nearly, if not all, of the affected lobe. Several lobes were involved, with congestion and edema of the other parts, and the breathing capacity was much impaired. The body for the most part was very much discolored, and the right heart was acutely dilated in almost every instance. In 14 cases the pleura was involved by direct extension of the inflammatory process with adhesions fibrous in character. Myocarditis and nephritis occurred in a small percentage of these cases, probably due to febrile disturbances. Heart blood was negative for bacteria in this series. No case with effusion or purulent exudate was classified in this group.

The bacteriology was determined in 30 cases. Pneumococci found in 26 cases: Type I, 5 cases; Type II, 2 cases; Type III, 5 cases; Type IV, 4 cases; type not determined, 10 cases.

Staphylococcus occurred alone in 2 cases, associated in 12. B. influenze occurred alone in 2 cases, associated in 9. Bacteriology not determined in 3 cases."

#### GROUP IV, HEMORRHAGIC BRONCHO-PNEUMONIA (18 CASES).

Pathology.-That of massive broncho-pneumonia type, associated with extensive hemorrhages into the lung tissues so that affected lobes were completely filled with blood. Several lobes were involved in many cases; those not involved in the hemorrhagic process were very much congested and edematous, thereby impairing the breathing space. The pleura was sometimes involved by direct extension of the inflammatory process. Adhesions, if any, were fibrous in character. There was no effusion or purulent exudates in this series, and the heart's blood was negative for bacteria. Associated myocarditis and nephritis occurred in many of the cases.

Bacteriology determined in 18 cases.

Pneumococci found in 18 cases: Type I, 1 case; Type II, 5 cases; Type III, 4 cases; Type IV, 1 case; type not determined, 7 cases.

Staphylococcus associated in 9 cases; streptococcus associated in 3 cases; B. influenza associated in 3 cases; B. mucosus associated in 1 case.

#### GROUP V. BRONCHO-PNEUMONIA WITH SEPTICEMIA (26 CASES).

Pathology.—The broncho-pneumonia found had not developed to any great extent before the more septic process terminated fatally. Empyema occurred in a large percentage of this group along with plastic purulent exudates into the thoracic cavity, pericardium, and even into the peritoneum. Adhesions of lung to thoracic wall occurred in most of these cases. Cultures from exudates and heart's blood contained pneumococci in every case. Myocarditis and endocarditis with vegetations on the heart valves occurred in a large percentage of the cases. Nephritis was found in 20 of the 26 cases.

Bacteriology determined in 25 cases.

Pneumococci were isolated in 23 cases: Type I, 2 cases; Type II, 7 cases; Type III, 6 cases; type not determined, 8 cases

Staphylococcus associated in 11 cases; streptococcus associated in 4 cases; M. catarrhalis associated in 2 cases; B. influenza associated in 6 cases; B. mucosus associated in 2 cases; streptococcus viridens pure culture, in 2 cases. Bacteriology not determined in 1 case.

10.00

#### GROUP VI. BRONCHO-PNEUMONIA AND MENINGITIS (12 CASES).

Pathology.—The broncho-pneumonic process had not progressed extensively before associated meningitis occurred through invasion of the blood stream by the pneumococcus. In many cases death occurred early, even before marked meningeal symptoms had developed, the meningeal involvement being found post-mortem by spinal puncture, purulent fluid containing pneumococci being obtained. On opening the cranium, the brain was found covered with thick purulent material just beneath the pia, cheifly involving the vertex of the brain, the base being involved to a lesser extent. This purulent meningitis extended down the cord also, just beneath the pia mater. Sometimes this exudate was so thick in the spinal canal that nothing could be obtained by spinal puncture. No involvement of the myocardium was found in this group. Heart's blood cultures were positive in a large percentage of the cases. The bacteriology was determined in 12 cases. Pneumococcus was found in 12 cases: Type II, 8 cases; type not determined, 4 cases.

NOTE.—All pneumococci found in this condition proved to be Type II. Staphylococcus was found associated in 3 cases. Streptococcus viridens was found associated in 3 cases. B. influenza was found associated in 1 case. B. mucosus was found associated in 1 case.

#### GROUP VII. HEMORRHAGIC SEPTICEMIA (5 CASES).

Pathology.—In this group no broncho-pneumonia was found, the lungs not being involved in any pneumococcic process although extensive hemorrhages had occurred into the lung tissue in two cases. Blood was obtained from the spinal canal by puncture in two cases. On opening the cranium in one case the ventricles of the brain were found filled with blood, due to hemorrhage from the choroidal plexus into the lateral ventricles. In four cases extensive hemorrhage was found in the suprarenal glands, entirely destroying the gland substance; in one of these cases no other pathology was found except this suprarenal apoplexy. Intraventricular apoplexy was found in one case, no involve-ment of the suprarenals being found. In no case was there involvement of the myocardium. Kidneys showed inflam-matory involvement in a majority of these cases. Cultures from the heart's blood were generally negative, only one positive blood culture being found which proved to be a Type I pneumococcus. Pneumococcus found in lung smears in 2 cases, type not determined. No bacteria found in 1 case. Staphylo-coccus was found associated in 1 case. B. mucosus was found associated in 1 case.

#### GROUP VIII. PNEUMODODCIC MENINGIRIS<sup>1</sup> (7 CASES).

Pathology .- In this group meningitis was the predominating post-mortem finding. The brain and spinal cord were extensively involved. Spinal fluid obtained by puncture contained large numbers of pus cells and pneumococci. There was no lung involvement. Cultures from the heart's blood were positive, pneumococci being found in five cases. In two cases small hemorrhages into the lung tissue were found. In one case hemorrhage into one of the suprarenal glands was found.

The bacteriology was determined in 7 cases. Pneumococcus was found in 7 cases: Type I, 1 case; Type II, 1 case; Type IV, 1 case; type not determined, 4 cases.

Staphylococcus found associated in 1 case. Streptococcus associated in 1 case. B. mucosus associated in 1 case.

<sup>1</sup> It has since been reported from the U.S. Naval Hospital, Norfolk, Va., that nine patients were admitted to hospital from the U.S. S. Mercy with diagnosts of meninglits, cerebrospinal. The health records of four contained notes stating that pneumococci were found in the spinal fluid. There were no notes regarding laboratory findings in the other cases. All had received polyvalent antipneumococci serum intraspinally, and most of them intravenously also. All except one had been treated on the Mercy for periods of from tweive to twenty-one days. Che had been on the Mercy only five days. Chadmission to this hospital two cases were convalescent. The remainder ranged from moderately to seriously ill. Che died the night after admission. Meningorocci were found in four of the cases, including the one terminating fatally. Spinal puncture was not done in the two con-valescent cases. To of the negative cases showed clear fluid, and the others, slightly cloudy fluid. No pneumococci were found in any case.

91483-18-2

				6					
Note	Bacteriology very mixed.	Probably Type II infections.	Bacteriology very mixed.	Pneumococcus Types II and III predominating.	Two cases streptococcus viridens. Types II and III pneumococcus predominate.	All pneumococci identified were Type II.	Positive blood culture in only one case. Con- dition one of toxemia.	Three heart's blood cultures positive preumo- cocci, various types.	Predominating Types II and III.
Associated.	63	0	63	H	63	0	0	0	1
M. catarrhalis alone.	:	:		-		:	:		0
Associated.	1	-	53	1	63	Ч	Ч	1	10
B. mucosus alone.	:	3010		by and	1.1.1		-244	:	0
Associated.	4	63	6	00	9	1	0	0	25
B. influenzæ alone.	:	1	63		1		17.2		63
Associated.	-	0	00	00	4	00	0	-	15
Streptococcus alone.	6	: .		1	63		:21		63
Associated.	63	63	12	6	H	00	63	H	42
Staphylococcus alone.	:	:	63			- :	.:		63
Bacteriology not determined.	-	61	60	0	-	0	61	0	6
No type identified.	64	4	10	-	00	4	61	4	41
- Contraction of the	1000	0100		-110	1000	0000	-000-		15 113
Types of pneumococcus.	THHY								
Number cases pneumococcus	2	5	26	18	23	12	00	4	101
Total number of cases.	œ	2	33	18	26	12	10	7	116
Group.	I. Simple broncho-pneumonia	II. Lobar pneumonia	III. Massive broncho-pneumonia	IV. Hemorrhagic broncho-pneu- monia.	V. Septic broncho-pneumonia	VI. Broncho-pneumonia w i t h meningitis.	VII. Hemorrhagic septicemia	VIII. Pneumococcus meningitis	Total

#### REPORT OF INFLUENZA OUTBREAK IN THE THIRTEENTH NAVAL DISTRICT.

(A report submitted by the medical aid to the commandant of the thirteenth naval district.)

Influenza first appeared in serious form in this naval district September 17, 1918. September 16 a draft of 334 men was received from Philadelphia. This draft reached Seattle, Wash., at about 9 p. m., and was ferried across the sound on barges, disembarking at about 11 p. m. Many of the men were inadequately clothed. Few, if any, had overcoats, nor were they properly supplied with blankets. A number complained of cold on the trains. Eleven were sick upon arrival, and, while waiting around the detention camp for their clothing, blankets, and disinfection of their effects, some became so ill that it was necessary to remove them at once to the sick bay and later to hospital. The men were up almost all night in the process of reception.

On September 18 and September 20 approximately the same number of men were received at the same time and under practically the same circumstances. These three drafts from Philadelphia were kept in isolation for a period of one month in the detention camp, navy vard, Puget Sound.

Of the first draft, 11 were sick upon arrival; of the second, it was necessary to remove 8 to sick quarters by ambulance, and 23 had to fall out during muster; of the third draft, 8 arrived by ambulance and 5 fell out during muster. Most of those who fell out during muster, or "dropped in their tracks," to use the words of the medical officer, were suffering from exhaustion, for they recovered the following day. Our experience with the disease has been similar to that in other districts throughout the

Our experience with the disease has been similar to that in other districts throughout the United States. The disease has shown special predilection for the enlisted personnel, especially those between the ages of 18 and 26, large numbers of whom have come down within a very short space of time.

On September 24, 1918, at a meeting of the board of health, thirteenth naval district, the question of quarantine against influenza was brought up and discussed. The following recommendations were made:

(a) That cases of influenza be isolated as early as possible.

(b) That all men remain away from public gatherings, such as theaters, dance halls, etc., for a period of 30 days.

(c) That attention be called to the dangers of coughing, sneezing, and expectorating.

(d) That men be instructed through their chief petty officers, or in such other way as seems best to their commanding officers, relative to sneezing, coughing, and expectorating, and that men found to be sneezing or coughing be sent to sick bay at once for treatment.

(e) That men be warned against using drinking fountains in public places outside of the navy yard.

(f) That the roads and parade grounds be sprinkled at least twice daily.

These recommendations were officially approved.

In the meantime, Surg. B. J. Lloyd, United States Public Health Service, sanitary adviser and inspector, thirteenth naval district, addressed the various units of the training camp on the subject of prevention in relation to influenza, at which time it was pointed out that the epidemic then raging in the East would undoubtedly reach the Puget Sound station very soon.

Early in the epidemic it became apparent that the naval hospital, navy yard, Puget Sound, would be able to care for only the more serious cases, and that of necessity it could receive patients only from the 7,000 men attached to the navy yard, Puget Sound. September 28 the medical aid canvassed the various hospitals in Seattle with a view toward securing accommodations for the sick of the command at the Seattle training camp who might require hospital treatment. This survey was most unsatisfactory, as practically no outside hospital accommodations were available.

Tuesday, October 1, the president of the University of Washington was interviewed, and he very kindly allotted Lewis Hall, an unoccupied dormitory, for use as a temporary hospital. By this time the sick bay, hostess house, and armory were crowded with the sick, and as immediate results were imperative assistance was asked from the Red Cross through the local director. By October 2 Lewis Hall was completely equipped for service as a temporary hospital. Women nurses and a sufficient number of medical officers were secured with the approval of the bureau and our prospects became brighter. This epidemic has been characteristic in so far as its tendency to attack large numbers in a short space of time is concerned. Doctors and nurses were not immune. At the naval hospital 5 doctors and 6 nurses sickened in a few days. One doctor and 1 nurse succurabed. At the naval training camp, navy yard, Puget Sound, 3 doctors were attacked. At the naval training camp, Seattle, 4 doctors and 5 nurses became ill. About 20 Hospital Corpsmen also contracted the disease. All but those infected at the naval hospital recovered. The rapidity with which some have been seized with this disease and have succumbed to the infection has been appalling. Many who suffered with severe attacks, after they had been ill for three days, bore a striking resemblance to patients in the third week of typhoid fever.

The effect of the epidemic upon the city of Bremerton and its influence upon the civilian personnel attached to this navy yard was appreciated. On September 27 the medical aid personally consulted with the physicians of Bremerton, at which time they were warned to take measures to control the epidemic. It was suggested that churches, moving pictures, theaters, lodges, dance halls, skating rinks, and schools be closed; that the streets be sprinkled frequently, as at this time there had been quite a dry spell.

The effects of the disease have, at this time, been so protean that it seems best to discuss them in only a general way, reserving detailed observations for a later report. The lungs, pleuræ, kidneys, brain, gall bladder, larynx, stomach, and intestines have all been attacked. The outstanding feature in serious cases is toxemia. In 75 per cent of the cases examined the hemolytic streptococcus has been obtained from the blood and from the organs mentioned above. Organisms have been recovered from the blood as early as 18 hours after the attack. Pneumococci have been recovered from the blood in only one case. The bacillus of Pfeiffer has not been found in any case.

At the suggestion of the sanitation officer, a vaccine was prepared from a hemolytic streptococcus isolated on September 25, and tried out on a few volunteers. There were no ill effects. It was then decided to vaccinate as many men as the facilities of our bacteriological laboratory would permit. Accordingly, from September 30 to October 25, 3,200 enlisted men, including marines who had not had influenza were given three injections at 48-hour intervals, starting with 60,000,000 for the first dose, 120,000,000 for the second dose, and 240,000,000 for the third dose. In our experience there have been practically no untoward effects from the use of this vaccine.

The following record has been compiled from data procured in the training camps in connection with the use of the vaccine. Of 2,800 men who were vaccinated, 45 suffered mild attacks of influenza following the first injection; of the remainder, 8 were attacked after the second injection and 2 after the third. A draft of 111 Filipinos reported on October 4, during the height of the epidemic, and were isolated at once. They received three prophylactic injections, and only 2 of them were attacked. Eighty-three machinist's mates for aviation, who arrived on October 3, were isolated and vaccinated. Thirty-one contracted influenza after the first injection; 1 only after three injections. Three hundred and sixty-one Marines were vaccinated; 2 came down with influenza after the first injection; none were affected after the third. Sixty-two marines at the ammunition depot were vaccinated; 2 were taken with influenza after the second injection, and 1 after the third, no others being affected. Six hundred and sixty-two blue jackets, members of the naval unit attached to the University of Washington, were vaccinated; 3 came down after the first injection; 1 after the second, and 7 after the third. No deaths have occurred among those who have been vaccinated, and pneumonia has thus far not developed in those inoculated. Hemolytic streptococcus vaccine was first administered on September 30, 1918. It was impossible for us to prepare the vaccine in requisite amounts in time to inoculate the 11,000 men attached to the two training camps mentioned above. However, an effort was made to inoculate those who had not had the disease and who could be controlled for observation. Later on vaccine was furnished to the health departments of the cities of Seattle, Portland, and Bremerton. Copies of letters received from the Washington State commissioner of health and the city health officer of Seattle, Wash., are inclosed. It is believed that the administration of this vaccine has been justified by the results obtained. Our experience has forced us to the conclusion that while the influenza bacillus may or may not have been the original factor in connection with the epidemic in the East, the hemolytic streptococcus has been responsible for the deaths here and our efforts have been directed toward immunization against this organism.

Naval training camp, Seattle, Wash.:       4, 159         Complement.       4, 159         Number of admissions for influenza from Sept.       813         25, 1918, to Oct. 21, 1918.       813         Deaths.       33         Percentage of complement attacked       19, 54         Case fatality rate       4. 05         Naval training camp, navy yard, Puget Sound,       Wash.:         Complement       6, 466         Number of admissions for influenza from Sept.       25, 1918, to Oct. 21, 1918.         25, 1918, to Oct. 21, 1918.       633         Deaths.       634	Philadelphia draft (included in complement for navy yard):
---	---

In a general way it may be stated that we are still in doubt as to the etiology of this outbreak. The influenza bacillus has not been isolated in any case. If we are dealing with two distinct types of disease it is our opinion that if the influenza bacillus was responsible for the beginning of this pandemic, as far as our experience in this district is concerned, this organism is without importance. We are dealing here, especially in the serious cases, with a hemolytic streptococcus, which may be a primary or a secondary invader and which is responsible for our mortality. The pneumococcus has been isolated in only one instance. In practically all other cases examined the hemolytic streptococcus has been found in the blood as early as 18 hours after invasion, and in the blood of some who have recovered. Our results with the streptococcus as the culprit. We feel certain that overcrowding, with its increased opportunity for cross infection, is of the greatest importance as a predisposing factor.

It is further submitted that greater care in transporting troops under certain circumstances should be observed. Contracts with railroads should provide for adequate heating, sufficient bed clothing, and one berth per man. Troops to the number of 100 or over should be accompanied by a medical officer. Troops should not be detrained at night. Trains arriving late at night should be sidetracked and remain intact until morning, if in the judgment of the medical officer and officer in charge of the detachment such procedure is warranted. It is believed that the high mortality in the Philadelphia draft was due in part to the conditions under which they were transported to this yard.

Too much can not be said by way of appreciation with regard to the whole-hearted assistance which has been rendered at a time of great emergency by commanding officers, doctors, nurses, Hospital Corpsmen, the president of the University of Washington, and the director of the Seattle division of the American Red Cross.

	Diseases 1	reported i	n 1 ship oj	f the fleet	, week end	led Sept.	21, 1918.			50
Complement Fotal admissions, disease Fotal admissions, venere Number of communi	al diseases		sive of ve	nereal di	iseases, 0.	······				59
	Diseases 1	eported is	n 3 ships o	of the fleet	, week en	ded Sept.	28, 1918.			. 2.
Complement Fotal admissions, disease Fotal admissions, venere Number of communi	al diseases		sive of ve	nereal di	iseases, 0.					54 
	Diseases :	reported i	in 14 ships	of the fle	et, week e	nded Oct.	5, 1918.			
omplement otal admissions, disease	s and injuri	08								10, 48
otal admissions, veneres	al diseases									2
Number of communi	cable diseas	es, exclu	isive of ve	nereal di	iseases, 6	1, as 10110	ws:	5	Surger St.	223
	Cerebro spinal fever.	Diph- theria.	Influenza.	Malaria.	Measlas.	Mumps.	Pneu- monia, broncho.	Pneu- monia, lobar.	Scarlet fever.	Typhoi fever.
J. S. S. De Kalb			56							
I S S Denver	Con the second states of		4							

91483-18-3

U. S. S. Quinnebaug .....

Diseases reported i	in 14 ships o	f the fleet, week	k ended Oct. 12, 1918.
---------------------	---------------	-------------------	------------------------

Complement Total admissions, diseases an Total admissions, venereal d Number of communicab	nd injuries	8			 			169
P.L. marker and and and a state	Cerebro	Diph	al local	-	 Pneu-	Pneu-	Scarlet	Typhoid

and the second s	spinal fever.	Diph- theria.	Influenza.	Malaria.	Measles.	Mumps.	monia, broncho.	monia, lobar.	fever.	fever.
U. S. S. Denver			11					1		
		and the second se						1		
U. S. S. De Kalb			12							
U. S. S. Susquehanna			41				2			
U. S. S. Florida						1				
U. S. S. Texas						1 28.1				

446 36

Diseases reported in 15 ships of the fleet, week ended Oct. 19, 1918. Complement. Total admissions, diseases and injuries. Total admissions, venereal diseases. Number of communicable diseases, exclusive of venereal diseases, 115, as follows: 12,020

Chip American States - 1	Cerebro spinal fever.	Diph- theria.	Influenza.	Malaria.	Measles.	Mumps.	Pneu- monia, broncho.	Pneu- monia, lobar.	Scarlet fever.	Typhoid fever.
U. S. S. Minneapolis			2	L PAULTA		2.1.52				
U. S. S. Marblehead			67							
U. S. S. Chattanooga		10104.000	1	40.005.0	1000000	2230000				
U. S. S. Denver		D. C L	4	THE PERMIT	19 19 19	1 10 10 10 10 10 10 10 10 10 10 10 10 10				A ST LAN
U. S. S. De Kalb	1.18 10. 16	1	2	20. 1.20	· · · · · · · ·	1	Hered.			12000100
U. S. S. Arizona			· Storeday				2			
U. S. S. Finland			3							
U. S. S. Princess Matoika			-5		2					
U. S. S. Frederick	1		5	habitet) et		Ten Inch				
U. S. S. Mount Vernon	Sume	1.19	20	63064 . 34	2	2	. Ariana			
					and the second second second	1	1			

Diseases reported in 57 ships of the fleet, week ended Oct. 26, 1918.

Complement. Total admissions, diseases and injuries. Total admissions, venereal diseases. Number of communicable diseases, exclusive of venereal diseases, 113 as follows: ... 38, 685 .....

	Cerebro- spinal fever.	Diph- theria.	Influenza.	Malaria.	Measles.	Mumps	Pneu- monia, broncho.	Pneu- monia, lobar.	Scarlet fever	Typhoid fever.
U. S. S. North Dakota			6							
U. S. S. Kentucky								1		
U. S. S. Mississippi			5			2				
U. S. S. Ohio						2	1			
U. S. S. Wisconsin		2				1	1			
U. S. S. Illinois							1			
U. S. S. Maine				1						
U. S. S. South Carolina					1					
U. S. S. Minneapolis			3							
U. S. S. De Kalb			5			3				
U. S. S. North Carolina		1.000	9			172				
U. S. S. Marblehead			9	Caller Number	1.4.4. 1. 1. 1. 1. 1.	Sector Sector				
U. S. S. Louisville			2						Anna (	
U. S. S. Cleveland			8.							
U. S. S. Pastores			1							
U. S. S. Calamares			1	The second						
U. S. S. Hancock			ī							
U. S. S. Mount Vernon			7							
U. S. S. Lenape			2							
			15							
U. S. S. Powhatan			1							
U. S. S. America			î				1			
U. S. S. Manchuria			2							
U.S.S. George Washington.			7						A STATE OF THE STATE	
U. S. S. Princess Matoika	State States		4					1		
U. S. S. Mongolia			2					-		
U. S. S. President Grant			ĩ							
U. S. S. Cheyenne			3				3			
or or oneycane			0				9			

### INSTRUCTIONS TO MEDICAL OFFICERS.

Hereafter triple typhoid lipo-vaccine will be furnished in place of the triple typhoid saline vaccine which has been used in the past.<sup>1</sup> The Army is prepared to furnish the lipo-vaccine and is now using it to vaccinate Army personnel against typhoid and para typhoid fevers. The following instructions covering its use will be sent with each consignment:

Each c. c. of this vaccine contains 2,500 million bacillus typhosus, 2,500 million bacillus paratyphosus A and 2,500 million bacillus paratyphosus B. On standing in the cold some of the fats may separate and cause a precipitate. This will disappear on standing a short time at room temperature. A single dose of 1 c. c. of this vaccine is sufficient. It is especially important that this vaccine be given subcutaneously, and not intravenously, intramuscularly, or under the fascia. In order to insure this, you will pick up a fold of skin and inject into the subcutaneous tissue of that fold. Practically all the severe reactions that have been reported have been due to neglect of this precaution. The deep injection of this vaccine may lead to fat embolism and diffeats the object of the inoculation.

The precautions to be taken regarding absence of temperature or disease are the same as are given for typhoid vaccine in Circular No. 16, War Department, Office of the Surgeon General, March 20, 1916. It is advisable to give the vaccine about 4 o'clock in the afternoon, and the man should be required to remain in quarters for 24 hours after the injection.

Extract from Circular No. 16, War Department, Office of the Surgeon General, Washington, March 20, 1916:

The site of the inoculation is the arm at the insertion of the deltoid muscle. If for any reason this site can not be used, the needle may be introduced in the back, over the lower portion of the scapula, or in the chest below the clavicle. The dose is to be given *subcutaneously* and not into the muscles nor into the skin. The arm should be cleansed as for any other operation. Tincture of iodine painted over the dry skin, before and after the injection, has proved satisfactory.

The ampul should be washed off in an antiseptic solution and opened after making one or more cuts near the top The ampul should be washed off in an antiseptic solution and opened after making one or more cuts near the top with a file. The vaccine can be drawn out of the container with a syringe. The syringe and needle should be ster-ilized by boiling. To insure perfect sterilization draw the piston out to its full length, or remove it entirely, so that the barrel is full of water during the boiling. A fresh needle should be used for each man, or, if one needle must be used on two or more men, it should be resterilized before each injection. No person should be vaccinated who is not perfectly healthy and *free from fever*. The temperature will be taken before vaccination is begun, and, in doubtful cases the urine should be examined; if fever or any other symptoms of illness are present the procedure should be postponed. This precaution is necessary to avoid vaccinating men who may be in the incubation stage of typhoid or other fever. Neither beer nor alcohol in any other form should be drunk on the day of treatment.

on the day of treatment.

Under certain nomenclature titles the following terms are frequently used in entering the diagnosis on Form F cards: Condyle, cuneiform, phalanx, scaphoid, etc. To enable the Sta-tistical Section to make proper classification, the foregoing terms, when used on Form F cards, should be amplified by definitely stating the location, viz: Condyle: State whether internal or external and the bone involved.

Cuneiform: Carpal or tarsal.

Phalanx: Proximal, middle, or distal, and first, second, third, etc., finger or toe, and whether right or left hand or foot, as indicated.

Scaphoid: Carpal or tarsal.

Medical officers are invited to submit to the bureau any suggestions that they believe may be useful in extending the scope of this bulletin to render it of greater value. Questions in the field of preventive medicine will be welcomed and will be answered in accordance with the best available authority.

Criticism of anything contained in the body of the bulletin or in any of the statistical tables will be appreciated. In referring to any article appearing in a bulletin, reference should be made to the letters and numbers in the upper right-hand corner, as well as to the serial number.

The following statistics are furnished for the information of medical officers:

The annual rates shown in the tables are obtained in the following way: The figure representing the total original admissions to the sick list or the number of deaths reported during the week is multiplied by 1,000 and divided by the complement. The quotient is then multiplied by 52. As weekly figures always fluctuate widely, caution must be used in interpreting annual rates calculated on a weekly basis. In the following tables it may be taken for granted where no figures appear that the disease did not occur, or, if in reference to hospitals, that no case was admitted. The figures in column "Total admissions for communicable diseases" do not include venereal diseases.

W. C. BRAISTED.

<sup>1</sup> Medical officers who require additional saling vaccine to complete a series of inoculations already begun with the triple typhoid saline vaccine should so state in letters or telegrams requesting vaccine.

TABLE 1.—ADMISSIONS TO SICK LIST AND ADMISSION RATES, TRAINING STATIONS, CAMPS, AND OTHER LARGER SHORE STATIONS, WEEK ENDED NOV. 2, 1918.

	81	Diseases	Diseases and injuries.		100			Venet	Venereal diseases.				
(has they here not Granding and the state of a colection with a colection with a colection of a colection a to the ortical may Fig for here in the second of ant is show it is necession of a colection o	Complement.	Total admissions, all causes.	Annual rate per	Average rate since July 1, 1918.	Total admissions, venereal.	190 Jair Der Der 1,000,1	Average rate since 1117 1, 1915.	Chancroid.	1900, 1000,	Gonococcus infec- tion.	Annual rate per 1,000.1	.silinqy8	Annual rate per
FIRST NAVAL DISTRICT. FIRST NAVAL DISTRICT. Air station, Chatham, Mass. Ammunition depot, Hingham, Mass. Aviation school, Cambridge, Mass.	428 350 1,700 1,091	125-1-2	1, 040.00 1, 376.44 571.38	484.19 3,068.36 1,707.02	5 H A	121.16 60.84	43. 42 21. 32 25. 58 40. 32			H 61	121.16		
District headquarters, Boston, Mass. District section headquarters, Boston, Mass. (Lockwood's Basin). District section headquarters, Bar Harbor, Me District section headquarters, Portland, Me District section headquarters. Portland, Me	511 1,815 358 364	20 20 4	1, 220. 97 572. 52 1, 016. 60 570. 96	264. 549. 240. 437.	8-18	57.20 145.08 285.48			145.08	F3	28.60 285.48		28.60
	350 1,596 1,596	1222525	1, 209, 00 57. 72 390, 00 814, 32	1, 366, 75 3, 061, 82 1, 065, 07 2, 045, 50 2, 042, 08								1111	
Navy yard, Portsmouth, N. H. Officers' material school, Cambridge, Mass. Radio school, Harvard University. Receiving ship Boston, Commonwealth Pier. Rifle range, Wakefield, Mass. Training camp, Bumkin Island, Boston, Mass Training camp, Hingham, Mass.	1, 211 4, 780 5, 303 506 1, 090	2 4 8 7 7 8 8 7 R	85.80 429.52 739.44 1, 233.92 1, 049.36 1, 049.36			$\begin{array}{c} 42.64\\107.12\\21.32\\48.88\\102.44\\158.08\end{array}$	$ \begin{array}{c} 11.88\\ 26.52\\ 43.26\\ 71.29\\ 70.60\\ 21.72\\ 21.72 \end{array} $		9.36		$\begin{array}{c} 42.04\\ 107.12\\ 21.32\\ 29.12\\ 102.44\\ 158.08\end{array}$		9.36
Other stations sECOND NAVAL DISTRICT. Bistrict section headquarters, New London, Comn Receiving barracks, Newport, R. I. Submarine base, New London, Conn Training station, Newport, R. I.	132 . 4,000 1,916 6,599	39 29 39 39	377.00 216.84 306.80	196. 75 875. 80 836. 22 1, 123. 16 851. 12	H SH	13.00 7.80	36. 27 5. 45 75. 24 33. 01			H 101 H	13.00 38.48 7.80		
THIRD NAVAL DISTRICT. Air station, Bayshore, Long Island, N. Y. Air station, Montauk, N. Y. Air station, Rockaway, Long Island, N. Y. Ammunition depot, Iona Island, N. Y.	880 387 1, 230 380	8 10 1	472. 68 422. 76 273. 52	822.00 366.25 708.46 1, 296.08	T	58.76	23.85 58.96 19.53			T	58.76		

40. 04 4 31. 72 50. 96	73.32 58.24 30.16	65.00 29.12 74.36 20.80	34.34 1 34.34 80.60	173.16 45.24 146.64 	208.00 13.00 48.88 2 9.36
	1 30.16	9.36 3 3 1		7.28 6 4 4	4.68
101. 87 37. 43 43. 08 1	. 16.86 72.35 59.42 28.65 38.39	43. 14. 16. 32. 32. 32.	49. 64 29. 64 37. 70 453. 21	5.87           74.21           122.82           43.64           7.80	37. 14 37. 14 31. 99 66. 25
9 71.76 1 50.96 1 39.52	1         73.32           2         58.24           2         60.84	65. 39. 74. 20.	2 69.16 1 80.60	1 173.16 9 67.60 4 146.64	1 208.00 2 13.00 3 63.44
601.00 1,004.95 900.08	1, 877.06 1, 877.06 1, 800.09 1, 961.14 1, 550.86 973.18	$\begin{array}{c} 1,029,29\\ 1,011,04\\ 3,337,23\\ 876,11\\ 876,11\\ 656,92\\ 2,074,83\\ 1,566,76\\ 1,566,76\\ 596,23\\ 596,23\\ \ldots\end{array}$	1, 931, 24 1, 931, 28 582, 34 2, 365, 97 1, 655, 97 4, 106, 05	1, 235, 43 1, 312, 18 733, 63 733, 63 733, 63	$\begin{array}{c} 1,129,12\\ 1,962,87\\ 1,550,51\\ 1,265,37\\ 1,434,70\\ \end{array}$
232. 44 975. 00 474. 76	355. 68 355. 68 606. 84 879. 84 416. 00 645. 32 523. 64 337. 28	585.00 586.32 86.32 707.72 707.72 707.72 521.04 728.00 674.96 1,485.64	1, 040.00 242.32 242.32 1, 446.64	346. 510. 440.	$\begin{array}{c} 374.40\\ 728.00\\ 416.00\\ 1,677.00\\ 1,007.24\\ \end{array}$
3 19	88 9 12 33 3 22 22 22 23 3 13 99 12 3	1	524 0004 -	1	4 0 0 258 258 258 258
1, 1, 6,	н н	$\begin{array}{c} 1,600\\ 600\\ 5,600\\ 5,330\\ 608\\ 608\\ 12,476\\ 12,476\\ 35\\ 858\\ 698\\ 800\\ 12,476\\ 858\\ 858\\ 858\\ 858\\ 858\\ 858\\ 858\\ 85$	1,500 1,500 1,500 1,500 1,152	6, 1,	694 500 350 8,000 10,581
Armed draft detail, New York: In barracks	Island, N. Y. District section headquarters, New Haven, Conn. District section headquarters, Sayville, Long Island, N. Y. District section headquarters, Tompkinsville, N.Y. District section headquarters, Whitestone, Long Island, N. Y. Federal rendezvous, Brooklyn, N. Y. Fleet Supply base, Brooklyn, N. Y. Navy yard, including marine barracks, New York.	Office of supervisor, auxiliary reserve, New York. Officers' material school, Princeton, N. J. Recruiting station, New York. Reserve force headquarters, New York. Receiving ship New York, including Ellis Island. Rifle range, Peekkill, N. Y. Steam engineering school, Stevens Institute, Ho- boken, N. J. Training camp, Pelham Bay, N. Y.	FOURTH NAVAL DISTRICT. Air station, Cape May, N. J	Air station, Hampton Roads, Va.	Ammunition depot, Portsmouth, Va

TABLE 1.—ADMISSIONS TO SICK LIST AND ADMISSION RATES, TRAINING STATIONS, CAMPS, AND OTHER LARGER SHORE STATIONS, WEEK ENDED NOV. 2, 1918—Continued.

Complement.		control from the constant of		-	aller on		Ven	Venereal diseases.	ises.	States		
	T'otal admissions, all causes.	Annual rate per 1,000,1	Average rate since July 1, 1918,	Total admissions, venereal.	Annual rate per 1,000.	A verage rate since July 1, 1918.	Chancroid.	Annual rate per 1,000.	Gonococcus infec- tion.	Annual rate per 1,000.1	Syphilis.	Annual rate per 1,000.
FIFTH NAVAL DISTRICT—continued.           Navy yard, including marine barracks, Norfolk, Va.         2, 215           Receiving ship, including St. Helena, Norfolk, Va.         7, 572           Rifle range, Glenburnie, Md.         7, 572           Rifle range, Virginia Beach, Va.         790           Other stations.	38.5	117, 00 260, 52 554, 32 263, 12	1, 177. 86 702. 02 1, 249. 78 688. 65	9	41.08	15.01 89.35 89.35 34.77			4	27.04	5	13.52
SIXTH NAVAL DISTRICT.		10 10 1 1										
Marine barracks, Paris Island, S. C.       8, 549         Navy yard, Charleston, S. C.       1, 000         Receiving ship, Charleston, S. C.       1, 000         Rifle range, Charleston, S. C.       1, 000         Rifle range, Charleston, S. C.       1, 000         Other stations.       1, 273         SEVENTH NAVAL DISTRICT.       1, 273	389 5 18 18 110 110 22	2, 366, 00 1, 398, 80 1, 312, 00 1, 315, 08 1, 315, 08	$\begin{array}{c} 1,153.82\\ 353.60\\ 1,739.61\\ 541.94\\ 235.32\\ 3,306.78\end{array}$	15 16 1 1	91.00 52.00 11.96	62.15 10.40 324.48 10.40 121.54 41.18		232.96	11 13 1	66. 56 1, 010. 36 11. 96	4	23. 92
Air station, Miami, Fla. (naval). Air station, Miami, Fla. (marine). Training camp, Key West, Fla.	45	$\begin{matrix} 1, 124. 24 \\ 839. 80 \\ 1, 046. 24 \end{matrix}$	$1, 416. 53 \\592. 21 \\1, 485. 84$	0 60	74.88 69.68	43. 95 30. 56 85. 38	- : : : :	37.44	3	37.44		
EIGHTH NAVAL DISTRICT.		18-11-1	A PARTY									
Training camp, Gulfport, Miss. 1, 901 Training camp, New Orleans, La. 2, 177 Training camp, Pensacola, Fla. 5, 508 Other stations.	17 48 91	464.88 1,146.60 859.04	$\begin{array}{c} 3,119.73\\ 2,081.40\\ 1,361.33\end{array}$	10	54.60 238.68	115.50 156.31 23.07	[ ]	27.04	5	119.08	4	27.04 95.16
NINTH, TENTH, AND ELEVENTH NAVAL DISTRICTS.		1. W. W.	A Store									
1,	20	669.24	1, 366. 21	00	100.36	33.36	:		60	100.36	:	
Buffalo, N. Y. 700 District section headquarters, Cleveland, Ohio 1, 200 District section headquarters, Detroit, Mich 650	13-7	520.00 563.16 639.60	1, 271. 39 819. 59 769. 40		43.16 79.56	47.59 69.30 73.19	111	79.56	1	43.16		

1 53.56 3 3.90		1 10.40 1 10.52				28 5.72
			-	:	:::	63
128.44		39.52 27.56 10.40 16.12		47.32	17.16	37.44
52		4444		1	F3	178
107.64 2.60						3.64
10101			1. 21		111	18
40. 17 43. 16 81. 83 78. 14		50.06 51.67 51.67 51.67 54.63 54.63 54.63 57.27 92.09 67.43		40.50	53.72	66.91
128.44 161.72 73.84		39.52 27.56 20.80 16.12 10.92		47.32	17.16	46.80
01 00 10			1 · · ·	1	67	224
$\begin{array}{c} 1, 376. 33 \\ 608. 21 \\ 1, 762. 92 \\ 959. 81 \end{array}$		$\begin{array}{c} 1,038.08\\ 1,991.00\\ 1,769.77\\ 1,769.77\\ 2,485.89\\ 2,485.89\\ 2,488.89\\ 1,385.45\\ 1,385.45\end{array}$	1 11		637.00 850.56	1, 043. 48
2, 046. 20 515. 32 323. 96 426. 40		$\begin{array}{c} 6,000.80\\ 2,742.48\\ 1,551.16\\ 2,011.36\\ 2,011.36\\ 1,402.96\\ 1,402.96 \end{array}$			274.56 796.64	776.36
61 8 328 328		176 51 47 128 128 139 130 128		1.	32 62	3, 690
1, 550 807 962 39, 975		$\begin{array}{c} 1,525\\ 967\\ 3,000\\ 1,307\\ 7,504\\ 4,912\\ 3,157\\ 4,706\end{array}$		1,	6, 044 4, 046	247, 020
Dunwoody Industrial Institute, Minneapolis, Minn Rifle mage, Camp Logan, Waukegan, III. Training camp, Detroit, Mich. Training station, Great Lakes, III. Other stations.	TWELFTH NAVAL DISTRICT.	Air station, San Diego, Cal. District section headquarters, San Diego, Cal. District headquarters, San Francisco, Cal. Submarine base, San Pedro, Cal. Training camp, Mare Island, Cal. Training camp, San Diego, Cal. Training camp, San Pedro, Cal. (municipal pier). Training station, San Francisco, Cal.	THIRTEENTH NAVAL DISTRICT.	Navy yard, including marine barracks, Puget Sound, Wash	Sound, Wash	Total

TABLE 1.—ADMISSIONS TO SICK LIST AND ADMISSION RATES, TRAINING STATIONS, CAMPS, AND OTHER LARGER SHORE STATIONS, WEEK ENDED NOV. 2, 1918—Continued.

Annual rate per 1,000.	43.1	15.60
'sduny	4	60
Annual rate per 1,000.		7. 80
Measles.		
Annual rate per 1,000.		19.24
German measles.		
Annual rate per 1,000.	32, 24	2.80
Malaria.		
Annual rate per 1,000.		
Diphtheria.		
Annual rate per 1,000.		
Cerebrospinal lever.		
Average rate since July 1, 1915.	131.04 114.92 86.93 38.02 38.93 38.02 38.93 38.93 30.55 10.99 117.03 10.99 10.99 113.07 30.16 30.16 30.16 30.16 30.16 31.67 31.77 31.77 31.77 31.77 31.77 31.77 37.77 37.77 37.77 37.77 37.77 37.777 37.777 37.7777 37.77777777	5, 62 64, 42 20, 62 20, 62 20, 62 24, 65 28, 35 28, 35 29, 57 20, 67 20,
Annual rate per 1,000.	60. 84 47. 32 28. 60 32. 24 43. 16	13.00 19.24 39.00 58.76 42.12
Total admissions com- municable diseases, e.c. ciusive of influenza.		1 1 1
	FIRST NAVAL DISTRICT. Air station, Chatham, Mass Ammunition depot, Hingham, Mass Aviation school, Cambridge, Mass Detention camp, Deer Island, Mass District headquarters, Boston, Mass. (Lockwoods Basin) District section headquarters, Boston, Mass. (Lockwoods District section headquarters, Bortland, Me District section headquarters, Portland, Me District section headquarters, Portland, Me District section headquarters, Rockland, Me District section headquarters, Rockland, Me Radio School, Quincy, Mass Navy yard, Boston, Mass Officer material school, Cambridge, Mass Radio School, Harvard University Receiving ship Boston, Oambridge, Mass Training camp, Hingham, Mass. Other stations.	SECOND NAVAL DISTRICT. District section headquarters, New London, Conn- Receiving barracks, Newport, R. I. Submarine base, New London, Conn. Training station, Newport, R. I. Other stations. Newport, R. I. Other stations. Newport, R. I. Air station, Newport, R. I. Air station, Montauk, Long Island, N. Y. Air station, Montauk, Long Island, N. Y. Ammunition depot, Iona Island, N. Y.

1

16

(

	9.36	22.36	6, 24 6, 24 24, 44 6, 76 654, 32
			L 10 L 4
			4,44
			52 52
			6.24
	· · · · · · · · · · · · · · · · · · ·		
	9.3		4,6
	4.10		
34. 57 58. 34 58. 34 58. 34 11. 41 11. 41 216. 32 39. 62 39. 62 123. 68	600, 14 600, 14 227, 100 227, 100 227, 74 2202, 19 2202, 19 2202, 19 220, 101, 10 104, 81 104, 81 206, 09 26, 09 26, 09 26, 00 26, 00 20, 14 10, 10 20, 14 10, 10 20, 14 10, 10 20, 14 10, 10 20, 10	18.08 89.12 89.12 64.48 64.48 17.43 17.43 17.43	$\begin{array}{c} 31.46\\ 90.93\\ 26.00\\ 114.67\\ 832.76\\ 332.06\\ 331.00\\ 208.02\\ 2522.59\end{array}$
7.80	174, 20 30, 16 32, 50 71, 76 29, 12 33, 28	22.36	$\begin{array}{c} 104.00\\ 32.24\\ 102.96\\ 13.52\\ 554.32\\ 65.52\\ \end{array}$
-	m ⊷ ⊷ 10 m 00		22 23 11
<ul> <li>Armed draft detail, New York.</li> <li>Auxiliary reserve school, Pelham Bay, N. Y</li> <li>Auxiliary reserve school, Pelham Bay, N. Y</li> <li>District section headquarters, Bath Beach, Long Island,</li> <li>N. Y</li> <li>District section headquarters, Montauk, Long Island, N. Y</li> <li>District section headquarters, New Haven, Conn.</li> <li>District section headquarters, Wentschne, Long Island, N. Y.</li> </ul>	Federal rendezvous, Brooklyn, N. Y Fleet supply base. Brooklyn, N. Y Fleet supply base. Brooklyn, N. Y Navy yard, including marine barracks, New York. Office of supervisor, auxiliary reserve, New York. Office of supervisor, auxiliary reserve, New York. Officer material school, Princeton, N. J Recruiting station, New York, including Ellis Island. Reserve force headquarters, New York. Reserve force headquarters, New York. Reserve force headquarters, New York. Receiving ship, New York, including Ellis Island. Rifle range, Peekskill, N. Y Steam-engineering school, Steven's Institute, Hoboken, N. J Training camp, Pelham Bay, N. Y Other stations	FOURTH NAVAL DISTRICT. Air station, Cape May, N. J. Ammunition depot, Dover, N. J. District section headquarters, Cape May, N. J. District section headquarters, Lewes, Del. Distributing barracks, Piter No. 19, Philadelphia, Pa. Navy yard, Philadelphia, Pa. Navy yard, Philadelphia, Pa. Radio school, Philadelphia, Pa. Radio school, Philadelphia, Pa. Training camp, Cape May, N. J.	FIFTH NAVAL DISTRICT. Air station, Hampton Roads, Va. Ammunition depot, Portsmouth, Va. District section headquarters, Baltimore, Md. District section headquarters, Cherrystone, Va. Marine barracks, Quantico, Va. Training station, Hampton Roads, Va. Navy yard, including marine barracks, Norfolk, Va. Receiving ship, including marine barracks, Norfolk, Va. Riffe range, Glenburnie, Md. Riffe range, Virginia Beach, Va.

TABLE 1.—ADMISSIONS TO SICK LIST AND ADMISSION RATES, TRAINING STATIONS, CAMPS, AND OTHER LARGER SHORE STATIONS, WEEK ENDED NOV. 2, 1918—Continued.

Annual rate per 1,000.		5.72		37.44 162.76		23.40		66. 56	502.84	33.80
.sqmuM		1		1			-	63	15	26
Annual rate per 1,000.										
Measles.										
Annual rate per 1,000.										
German measles.										
Annual rate per 1,000.				22.88		47.32				
Malaria.						10 10		:		
Annual rate per 1,000.										
Diptheria.							-	:		
Annual rate per 1,000.										
Cerebrospinal lever.										
Average rate since July 1, 1918.		$\begin{array}{c} 35.96\\ 35.96\\ 45.50\\ 10.40\\ 10.40\\ 263.77\end{array}$		56. 81 21. 03 38. 99		42.57 111.05 15.38		127.70	44. 24 76. 75 29. 90 189. 58	380. 09 191. 68 182. 27
Annual rate per 1,000.		188. 24 52. 00 47. 32		74. 88 69. 68 184. 60		119.08	121 AR	66. 56	159.64 603.72	36.40
Total admissions com- municable diseases, ex- clusive of influenza.		31 		00 H FO		10 00	-	63	18	28
	SIXTH NAVAL DISTRICT.	Marine barracks, Paris Island, S. C. Navy yard, Charleston, S. C. Receiving ship, Charleston, S. C. Rifle range, Charleston, S. C. Training camp, Charleston, S. G.	SEVENTH NAVAL DISTRICT.	Air station, Miami, Fla. (Naval). Air station, Miami, Fla. (Marine). Training camp, Key West, Fla.	EIGHTH NAVAL DISTRICT.	Training camp, Gulfport, Miss. Training camp, New Orleans, La. Training camp, Pensacola, Fla.	NINTH, TENTH, AND ELEVENTH NAVAL DISTRICTS.	Auxiliary reserve school, Chicago, Ill	N. Y District section headquarters, Cleveland, Ohio District section headquarters, Detroit, Mich. Dunwoody Industrial Institute, Minneapolis, Minn.	Training camp. Logan, waukegan, III. Training camp, Detroit, Mich. Training station, Great Lakes, III.

	214.76	152.36	165.36		94.64	91.96	
-	4 21	22 15 47 49	15 16	:	2 9		
-			:::	:	:		
	33.80					68.64	
	1			:		80	:
		31.72			-		
-	::	· · · · · · · · · · · · · · · · · · ·		:			
1				:			
-			::	:			
-	::		::	:		::	:
-			11	-			
			-				
			::	:	:		
	23.58 74.70 06.89	48.47	39. 11 68. 17		29.46	80.42	
	76	52 36 84 52 84	28 3	:		64 32 1	
	33.8	39. 5 152. 3 528. 8	176. 2	-	94.64	68. ( 372. 3	
	14	22 22			53	8 29	
TWELFTH NAVAL DISTRICT.	Air station, San Diego, Cal. District section headquarters, San Diego, Cal. District section headquarters San Francisco, Cal	Submarine base, San Pedro, Cal. Training camp, Mare Island, Cal. Training camp, San Diego, Cal.	Training camp, San Pedro, Cal. (municipal pier) Training station, San Francisco, Cal	Under Stauons THIRTEENTH NAVAL DISTRICT.	Navy yard, including marine barracks, Puget Sound, Wash Provide shin including training comm Puget Sound	Wash Training camp, Seattle, Wash	

3.12 171 35.88

15

. 52

4

2.08

10

.41

63

.20

1

66.04 114.67

316

Total. .....

TABLE 1.-ADMISSIONS TO SICK LIST AND ADMISSION RATES, TRAINING STATIONS, CAMPS, AND OTHER LARGER SHORE STATIONS, WEEK ENDED NOV. 2, 1918-Continued.

Annual rate per 1,000.	485.68	580.84	406.64	143. 00 435. 24	241.80 57.72 52.00 65.00	42. 64 228. 28 48. 88	210.60 715.32	March Inc.	117.00 54.08	39.00	10 011	212.68		277.16	355.68
Influ- enza.	4	19	4	0.00	01-0101	1 21 5	4 15	2 12	6.61	5		N 10	0 01 05 X	1	00
Annual rate per 1,000.								12 12							
Ty- phoid.															
Annual rate per 1,000.								2.08							
Small- pox.								an in							
Annual rate per 1,000.		00 LY						2	13.00						
Scarlet fever.								10	1						
Annual rate per 1,000.								1 N							
Tuber- culosis.								H III							
Annual rate per 1,000.		60.84		28.60	26.00			10		7.80	1	08.76 42.12	7.80		
Pneu- monia.		2			1			100		1		1			
	Air station, Chatham, Mass.	Aviation school, Cambridge, Mass.	District headquarters, Boston, Mass	woods Basin) District section headquarters, Bar Harbor, Me. District section headquarters, Portland, Me.	District section headquarters, FrovIncetown, Mass. District section headquarters, Rockland, Me Fuel-oil school, Quincy, Mass. Naval Prison, Portsmouth, N. H.	Navy yard, Portsmouth, N. H. Officer material school, Cambridge, Mass. Radio school, Harvard University Receiving ship, Boston, Commonwealth Pier.	Training camp, Bunkin Island, Boston, Mass. Training camp, Hungham, Mass.	. SECOND NAVAL DISTRICT.	District section headquarters, New London, Conn Receiving barracks, Newport, R. L.	Training station, Newport, R. I.	THIRD NAVAL DISTRICT.	Air station, Daysnore, Long Island, N. 1 Air station, Montauk, N. Y. Air station, Rockaway, Long Island, N. Y.	Ammunition depot, fona Island, N.Y. Armed draft detail, New York. Auxiliary reserve school Polham Bay N.Y.	District section headquarters, Bath Beach, Long Is- land, N. Y.	N.Y. District section headquarters, New Haven, Conn

514.80	208.00 58.24 290.68 130.00		azu, 00 83, 20 1, 485, 64	8 WE H	903.76	68.12	1, 2416.00 1, 241.24 117.00 20.28	1, 538. 68	154.96	860.60 244.92
.4	- 1 61 10 9 <del>4</del>	155	581		20	6	191 60 55 33	253	5	12
			4, 16				4. 68			the second
			1							
	32.50						6.76		52.00	
*	1						I			
10.20	174. 20 30. 16	71.76 9.36	24.96		as en		104.00 19.24 39.00	182.00		47.32
	0 H	1	9				ο φ τη	30		4
District section headquarters, Sayville, Long Island, N.Y. District section headquarters, Tompkinsville, N.Y.	Pland, N. Y. Landquarters, mitteeboure, 1008 12- land, N. Y. Federal rendezvous, Brooklyn, N. Y. Fleet supply base, Brooklyn, N. Y. Navy yard, including Marine barracks, New York. Office of supervisor, auxiliary reserve, New York.	Recruiting station, New York. Reserve force headquarters, New York. Receiving ship, New York, including Ellis Island Rifle range, Peekskill, N. Y Steam engineering school, Stevens Institute, Hobo-	Training camp, Pelham Bay, N. Y. Other stations. FOURTH NAVAL DISTRICT.	Air station, Cape May, N. J. Ammunition depot, Dover, N. J. District section headquarters, Cape May, N. J. District section headquarters, Lewes, Del	Distributing barracks, Pier No. 19, Philadelphia, Pa Navy yard, Philadelphia, Pa Officer material school, University of Pennsylvania, Philadelphia, Pa.	Radio school, Philadelphia, Pa Receiving ship, Philadelphia, Pa Training camp, Cape May, N. J Other stations.	FIFTH NAVAL DISTRICT. Air station, Hampton Roads, Va Ammunition depot, Portsmouth, Va District section, headquarters, Baltimore, Md District section headquarters, Cherrystone, Va. Marine barracks, Quantico, Va. Training station, Hampton Roads, Va. Navy yard, including marine barracks, Norfolk, Va. Receiring ship, including St, Helena, Norfolk, Va. Rifle range, Glenburnie, Md. Rifle range, Virginia Beach, Va.	SIXTH NAVAL DISTRICT. Marine barracks, Paris Island, S. C. Navy yard, Charleston, S. C.	Receiving ship, Charleston, S. C. Riffe range, Charleston, S. C.	Training camp, Unarleston, S. U.

TABLE 1.—ADMISSIONS TO SICK LIST AND ADMISSION RATES, TRAINING STATIONS, CAMPS, AND OTHER LARGER SHORE STATIONS, WEEK ENDED NOV. 2, 1918—Continued.

	Annual rate per 1,000.	149.76 279.76 604.24	109.20 310.44 235.56	234.00 520.00 159.64 67.40 52.00	5, 796, 44 2, 366, 00 1, 232, 92 1, 232, 92 304, 72 296, 40	42. 64 230. 88	332.80
	Influ- enza.	4 4 26	25 25	- 100 4 0 <del>4</del>	170 447 31 31 66 66 68 18	185	1, 581
	Annual rate per 1,000.						
	Ty- phoid.						
	Annual rate per 1,000.						
	Small- pox.						
· manim	Annual rate per 1,000.		10 L				.52
nonminion.	Scarlet fever.						4
DTAT (#	Annual rate per 1,000.		23.40				.52
	Tuber- culosis.		1				4
ALLAND	Annual rate per 1,000.	37. 44 69. 68	23.40 9.36	159.64 100.36	39.52 10.92	282.36	21.84
	Pneu- monia.			69 00 69	1	22	105
Labor with the property gas and a second		SEVENTH NAVAL DISTRICT. Air station, Miami, Fla. (naval) Air station, Miami, Fla. (marine) Training camp, Key West, Fla Other stations.	H NAV port, M Orlean acola, ]	NINTH, TENTH, AND ELEVENTH NAVAL DISTRICTS. Auxiliary reserve school, Chicago, III. Districts. Aviation detachment, Curtiss Aeroplane Plant, Buf- falo, N. Y. District section headquarters, Cleveland, Ohio. District section headquarters, Detroit, Minch. District section headquarters, Detroit, Minch. District section headquarters, Ill. Dinwoody Idustrial Institute, Minneapolis, Minn. Rifle range, Camp Logan, Waukegan, III. Training camp, Great Lakes, III. Other stations.	TWELFTH NAVAL DISTRICT. Air station, San Diego, Cal. District section headquarters, San Diego, Cal. District section headquarters, San Francisco, Cal. Submarine base, San Pedro, Cal. Training camp, Mare Island, Cal. Training camp, San Diego, Cal. (municipal pier). Training tauton, San Pedro, Cal. (municipal pier).	THIRTEENTH NAVAL DISTRICT. Navy yard, including marine barracks, Puget Sound, Wash Receiving ship, including training camp, Puget Sound, Wash. Training camp, Seattle, Wash. Other stations.	Total

1

E

# TABLE 2.—SUMMARY OF REPORTS FROM NAVAL HOSPITALS AND SICK QUARTERS, WEEK ENDED OCT. 26, 1918.

<sup>1</sup> Report not received.

## TABLE 3.—NUMBER OF ADMISSIONS AND ANNUAL RATES, ENTIRE NAVY, WEEK ENDED<br/>OCT. 19, 1918.

#### [Rates based on an estimated complement of 550,600.]

Class.	Number of cases.	Annual rate per 1,000.	Class.	Number of cases.	Annual rate per 1,000.
Diseases of blood Diseases of circulatory system Diseases of digestive system Diseases of ductless glands and spleen Diseases of ear Diseases of ear Diseases of genito-urinary system (nonvenereal) Diseases of infective type (non- venereal) Diseases of infective type (ve-	36 687 61 46 165 8, 945		Diseases of motor system Diseases of nervous system Diseases of respiratory system Diseases of skin, hair, and nails Hernia. Miscellaneous diseases and condi- tions. Parasites (fungi and certain animal parasites). Tumors. Injuries. Poisons.	$78 \\ 68 \\ 1, 337 \\ 70 \\ 37 \\ 126 \\ 68 \\ 6 \\ 456 \\ 6 \\ 6$	126.36
nereal) Diseases of lymphatic system Diseases of mind	729 32 35	68.64		12, 988	1, 227. 72

Chancroid infection. 155 Gonococcus infection. 450 Syphilis. 124



TABLE 4.—TOTAL ADMISSIONS REPORTED BY FORM F CARDS FOR CERTAIN COMMUNICABLE DISEASES, OTHER THAN VENEREAL DISEASES, AND ANNUAL RATES, ENTIRE NAVY, WEEK ENDED OCT. 19, 1918.

[Rates based on an estimated complement of 550,000.]

Disease.	Number of admissions.	Annualrate per 1,000.	Disease.	Number of admissions.	Annual rate per 1,000.
Cerebrospinal fever	15	1.41	Chicken pox	6	0.56
Diphtheria	19	1.79	Dengue	19	1.79
Malaria	3 45	4.23	Influenza	8, 315	785.72
Measles	9 32	3.01	Paratyphoid fever	1	. 09-
Mumps	209	19.76	Whooping cough	2	. 19
Pneumonia, broncho	J 784 584	55.12	Tuberculosis	31	2,92
Pneumonia, lobar		18.72		+	
Scarlet fever	8	. 75	Total	9,491	906.36
Typhoid fever	E 10 2	.19			marken

#### TABLE 5 .- DEATHS REPORTED, ENTIRE NAVY, WEEK ENDED NOV. 2, 1918.

[Rates based on an estimated complement of 600,000.]

Influenza-pneumonia	1 mile and the line	T. Jackson and J	and Salaran material	638
Influenza-meningitis, cerebrospinal 1		In all I loud	and and the set of the local of	2
Cerebrospinal fever		hand and a starter		- 5
Typhoid fever				1
Other infections				2
Drowning Other accidents and injuries			A state of the sta	4
orner accidento and injuries		I levelene and	1	and the
Total		mul		658
Annual death rate per thousand, all causes		has been produced of		56.99
Annual death rate per thousand, diseases only.		hand the proved		56.16
and the second second second free preserves	1 Pneumococcus, 2	2	a character and a second state	TONOS

And a located state to a state of the

14 star broad on any antimated court leakent of 550,80

		WASHINGTON	GOVERNMENT	PRINTING OFFICE : 1918
--	--	------------	------------	------------------------