

**The anatomy of the human gravid uterus exhibited in figures. = Anatomia
uteri humani gravidi tabulis illustrata. / By William Hunter, Physician
Extraordinary to the Queen, ...**

Contributors

Hunter, William, 1718-1783.

Publication/Creation

London : Printed for E. Cox and Son, by T. Bensley, 1815.

Persistent URL

<https://wellcomecollection.org/works/wc7gxkcu>

License and attribution

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>

HUNTER'S

GRAVID UTERUS,







36

A N A T O M I A

UTERI HUMANI GRAVIDI

TABULIS ILLUSTRATA.

AUCTORE

GULIELMO HUNTER,


SERENISSIMAE REGINAE CHARLOTTAE MEDICO EXTRAORDINARIO,

IN ACADEMIA REGALI ANATOMIAE PROFESSORE, ET SOCIETATUM,
REGIAE ET ANTIQUARIAE, SOCIO.

LONDINI:

IMPENSIS EDITÆ ELIZABETHÆ COX FILIIQUE, IN VICO VULGO VOCATO, ST. THOMAS'S STREET, BOROUGH.
EXCUDEBAT T. BENSLEY, BOLT COURT, FLEET STREET.

1815.



THE ANATOMY
OF THE
HUMAN GRAVID UTERUS
EXHIBITED IN FIGURES.


BY

WILLIAM HUNTER,
PHYSICIAN EXTRAORDINARY TO THE QUEEN,
PROFESSOR OF ANATOMY IN THE ROYAL ACADEMY, AND FELLOW OF THE ROYAL AND
ANTIQUARIAN SOCIETIES.

LONDON:

PRINTED FOR E. COX AND SON, ST. THOMAS'S STREET, BOROUGH;
BY T. BENSLEY, BOLT COURT, FLEET STREET.

1815



TO
THE KING.

SIR,

THIS Work had no other claim to the honour with which it is distinguished by Your Majesty, than as it illustrates one part of Science hitherto imperfectly understood, and as it contains the foundation of another part of Science, on which the lives and happiness of millions must depend. It gave me an heart-felt satisfaction to observe how much this consideration recommended it to Royal Patronage.

When respect, reverence, and gratitude rise to a certain degree, and the occasion is public and solemn, the freedom of language is restrained: and I should be guilty of an abuse of that gracious condescension with which I am honoured, were I to attempt to express by words the sentiments of my heart. But, for the sake of those who, placed at a distance from the Royal Presence, may be more able to serve the public than I have been, let me be permitted to say, that, although to acquire knowledge, and to communicate it to others has been the pleasure, the business, and the ambition of my life; yet, since I have had the honour to be a servant in Your Royal House, I have felt an additional incitement to assiduity, being convinced that an unremitting endeavour to be useful to society is the readiest and the most certain way to the favour of our Sovereign.

That the peculiar blessing of this Country, the influence of a King who sincerely wishes to encourage every improvement, a King who takes a part with all his subjects in their happiness, and in their distress, may be long preserved to us, is the fervent prayer of

SIR,

YOUR MAJESTY'S

Most faithful Subject,

And most dutiful Servant,

WILLIAM HUNTER.

To the Publishers of this Impression of *Dr. Hunter's Plates on the Gravid Uterus, with the annexed Notes*, much credit is due, for the great care and pains they have taken to preserve or restore them to their primitive state of exactness and beauty; and for the neatness with which the work in general is completed. They have, with more earnestness than the affair seemed in my opinion to deserve, solicited me to make some addition to the Preface and Notes, in the way of explanation and recommendation. With their solicitation I am concerned I cannot comply; not thinking myself competent to interfere with, or justified in making any alteration or addition to what DR. HUNTER himself had thought sufficient and proper to do on the occasion; and as to any recommendation of the Work, none can be required, nor any advantage gained, by attempting to exalt a character so well known and established as DR. HUNTER's is, not only in this country, but in every part of Europe.

BUT as this request of the Publishers affords me an opportunity of bearing my testimony to the general character of DR. HUNTER, founded upon a long attendance on his Lectures, I hope to escape censure if I presume to indulge my inclination, by giving a short view of the talents and habits of a man, who excelled all his competitors in those branches of the profession which he cultivated.

OF DR. HUNTER's education, it is enough to say that it was fully adequate to all the purposes of his profession. His assiduity and perseverance in the acquisition of knowledge were scarcely ever exceeded; his faculties were acute, and his judgment penetrating and accurate; and these co-operated to qualify him for the important pursuits in which he was engaged, especially in his anatomical disquisitions, all of which I consider as excellent, though none seem preferable to his account of the Cellular Membrane in the second volume of Medical Observations and Inquiries. His style and manner of writing, and his method in delivering his Lectures, were explicit and comprehensive; his language was correct; his pronunciation distinct and harmonious; and his illustrations appropriate. His knowledge of the principles of Midwifery, to the practice of which he dedicated himself, was enlightened and just, of which the present Work may be considered as a striking example. With these talents and habits, it cannot surprize that he should have contributed in a more than ordinary degree to the edification of students, and the improvement of general practice. Nor was this confined to his own time, for there has scarcely been a book since published on that subject, which has not some reference to the doctrines or works of DR. HUNTER. His diet and manner of living were particularly abstemious; and in all that respected his person, there was a natural and an attentive regard to elegance.

I do not here take notice of his other pursuits; of his collections of Natural History; of Paintings; of Books, which were select and considerable; nor of his collection of Coins and Medals, which being large and valuable, was highly esteemed, and purchased at a very great expence. These merely prove his unceasing attention to science in general; they to many would have been of chief consequence, but the reputation of them has been absorbed in his character as a most skilful, dexterous, and scientific Anatomist.

THOMAS DENMAN.

MOUNT STREET,
September 1, 1815.

PROÆMIUM.

QUOD omnes fere homines, in studiis scientiarum versati, diu multumque desiderarunt, linguam scilicet per universum terrarum orbem cognitam atque perceptam, id nobis ex parte aliqua suppeditat tabularum aenearum sculptura, ad veras rerum formas exhibendas adprime accommodata. Imo hujusce artificii ea est utilitas, ut in naturæ explicacione sepenumero plus valeat quam quævis verborum copia, aut vis dicendi. Quodcumque enim oculo fidelí inde subjicitur (spectet modo aliquis argumenti haud ignorans) animo confessim arripitur, memoriaque alte impressum haud porsus excidit. Lentior sermonis mora, et vis hebetior.

Prinde ex quo tempore hoc artificium usibus hominum communiter inservit, via brevior faciliorque patet; est ad inventa quilibet tum divulganda, tum conservanda; et rerum simul naturalium cognitio, ab exiguis prospecta initii, paulatim eo usque incrivit, ut seculi hujusce, omnium ornatissimi, summum fere ornamentum jam tandem extiterit.

Ceteraum artium incrementa, aequis saltē passibus, comitata est anatoma. Etenim non pauci recentiorum, viri excellentis ingenii atque judicii, labore plurimo, nec minori solertia, corporis humani tum fabricam tum economiam pervestigarunt. Ex his plurimi artem hanc graphicam ancillantem sibi adhibuerunt; præcipisque inde repertis suis feliciter illustratis, effecerunt, ut ars anatomica, hominum generi, si qua alia, utilis et fructuosa, intellectu simul facilior fieret, simul cultu jucundior.

Restabat tamen pars una humani corporis, eaque sive structure sue perelegantis, sive muneri officiique ratione, primas tenens, a discipline hujus professoribus non satis explorata. Restabat scilicet uterus humanus. Absit vero, ut viris hisce egregiis vitio detur, quod rem quasi neglectam habuerint, cuius penitus perspicienda copia rara erat et infrequens. Paucissim enim contigerit necesse est, ut tot cadavera secuerint, quot ad conditionem uteri gravidi, variis temporibus variam, pervestigandam, nedum demonstrandam, sufficerent. De aliis itaque silendum, ubi laudare nefas. Quod ad auctorem attinet, anno ille primum MDCCLII, felici usus est occasione id in humano corpore explorandi, quod in brutis jam olim quæsiverat. Mulier quedam, graviditatis sue mensibus prope exactis, repentinam obierat mortem. Cadaver, adhuc integrum, cultro subjectum est. Operi favit tempestas. Cera, vasis rite injecta, voto respondit. Repertus est pictor, vir ille in suo genere adprime solers. Partes singule, multis coram adstantibus, exploratae sunt atque excusse, ut omnis rei veritas firma satis auctoritate comprobari posset atque stabiliri.

Sub his auspiciis, aliquot menseum spatio decursu, decem prime tabulae perfecte delineatae fuerant; partes ipsae identidem publice exposite; et ea que de singulis animadversa fuerant ac notata, palam dicta. Auctoris inde animum subiit, se rem discipline hujusce fautoribus haud ingratis facturum, si historiam uteri gravidi anatomicam ederet, his decem tabulis illuminatam. Consilium placuit. Opus in manus artificium, maxima laude floruentum, traditum est; et pecunia collatitia a plurimis pro more reprezentata.

Alterum interea cadaver in manus incidit, quod quidem, coelicit tempestas minus jam anatomicæ amica esset, haud nullo in pretio habendum fuit; quandoquidem huic acceptæ referuntur figuræ aliquot, ab hoc instituto haud alienæ, dignæque adeo vise, que operi, supplementi loco, adjicerentur. Dum autem super his operam navabat sculptor, en tertium cadaver quam opportune! Etenim illud non modo quedam, prius obscura, luci dedit, sed novis etiam figuris adornandis materiam aptissimam subministravit.

P R E F A C E .

THE art of engraving supplies us, upon many occasions, with what has been the great desideratum of the lovers of science, an universal language. Nay, it conveys clearer ideas of most natural objects, than words can express; makes stronger impressions upon the mind; and to every person conversant with the subject, gives an immediate comprehension of what it represents.

From the time when this art came more generally into use, it has been much more easy both to communicate and to preserve discoveries and improvements; and natural knowledge has been gradually rising, till it is at length become the distinguishing characteristic of the most enlightened age of the world.

Anatomy has, at least, kept pace in improvement with the other branches of natural knowledge. Many of the moderns, through much labour and patience, as well as ingenuity and judgment, have thrown considerable lights upon the structure and operations of the human body; and they have particularly by engravings made the study of that art, in which humanity is so much interested, both more easy and pleasant. Most of the principal parts of anatomy have, in this manner, been successfully illustrated.

One part however, and that the most curious, and certainly not the least important of all, the pregnant womb, had not been treated by anatomists with proportionable success. Let it not, however be objected to them, that they neglected what in fact it was rarely in their power to cultivate. Few, or none of the anatomists, had met with a sufficient number of subjects, either for investigating, or for demonstrating the principal circumstances of utero-gestation in the human species. But let what cannot be praised, in others, be passed over in silence. With respect to the present undertaking, in the year 1751 the author met with the first favourable opportunity of examining, in the human species, what before he had been studying in brutes. A woman died suddenly, when very near the end of her pregnancy; the body was procured before any sensible putrefaction had begun; the season of the year was favourable to dissection; the injection of the blood-vessels proved successful; a very able painter in this way was found; every part was examined in the most public manner, and the truth was thereby well authenticated.

In the course of some months, the drawings of the first ten plates were finished, and from time to time the subject was publicly exhibited, with such remarks as had occurred in the examination of the several parts. Many lovers of this study approved of the author's proposal to publish the anatomy of the gravid uterus, illustrated by those ten plates; the work was immediately put into the hands of our best artists; and subscriptions were received.

In the mean time a second subject was procured; which, though the weather happened to be very unfavourable, afforded a few supplemental figures, of importance enough to be taken into the work. And before the engravings were finished, a third subject occurred very opportunely, which cleared up some difficulties, and furnished some useful additional figures.

PROCEMIUM.

Cum auctor huic se accingebat negotio, animo destinavit decem tabulas, modo dictas, in lucem proferre, jactum quasi aliquod fundamen, cui alia plura, de hac parte anatomie, postea adjicere ac superstruere sperabat, si quando per otium ipsi licet hanc rem altius investigare. Jam vero majora quadam animo completi, et etiam spem fovere ausus est, fore ut opus aliquod ex adificare posset, quod ad perfectionem absolutionemque proprius perveniret. Prievidit enim, in hac urbe, tanta hominum multitudine referta, annorum aliquot spatio, occasionem omnia ea que ad gravidum uterum pertinent, observandi viro, hoc agenti, haud fortasse defuturam; unde sibi facultas dari posset aliquod quasi sistema condere, variasque illas mutationes, singulis graviditatis mensibus proprias, figuris exprimere. Ejusmodi fere est hoc opus; quod, longo tempore promissum, tandem aliquando in medium prodit. Ut id sane ad exitum perduci posset, moras trahere necesse fuit, præter opinionem diurnas. Quoties autem auctor secum reputat has moras operi minime infructuosas fuisse, abunde sentit, quo se consoletur.

Quod autem hoc argumentum illustrandi uni alicui mortali tam pulchra occasio toties obtigerit, in causa fuerunt tum vari casus, tum amicorum multorum gratissima subsidia. Tabularum harum ultima et manu artificis vix excesserat, cum datum fuit delineare embryonem mole minorem, quam quem antea vidisse auctori unquam contigerat; et haud ita pridem conceptus in tuba fallopiana, res haec animadversione egregie digna, figurarum numerum adauxit. Haec figure, et si qua alia, ejusdem argumenti, occurrent, vel in tabula his adjicienda, vel cum uteri gravidi descriptione anatomica, quam auctor secum meditatur, data occasione, in publicum aliquando proferentur. Hactenus quidem ipsi id otii non concessum est, quod ad hanc operis partem absolvendam esset satis. Quod si casu aliquo improviso impeditus fuerit, quo minus id, quod cupiat, possit exsequi, persuasum sibi habet, haud defuturum aliquem, huic muneri obeyendo parem. Multi enim sunt, qui auctorem de his rebus publice disserentem sepe sepius audierint; multi quos ille opinionis ac sententiae suee participes fecerit.

FIGURARUM fere anatomicarum duplex est et diversa ratio. Ab altera simplex rei imago, qualis oculo visa est, talis delineatur. Altera, una cum visis, etiam excogitata exhibet; nec tam ad oculum artificis, quam ad ingenium memoriamque accommodatur. Prioris generis exempla apud Bidloo extant; posterioris apud Eustachium.

Fieri non potest, quin figura ista, que fit ex fideli naturæ imitatione, et ex rei unius contemplatione enascitur, in quibusdam sepe partibus indistincta aut minus perfecta esse videatur. Altera autem, ex ingenio nata, que autem non fit, nisi per laborem, et observationem et experientiam multiplicem, id uno intuitu videndum exhibet, quod fortasse non nisi ex rerum varietate colligendum fuerat. Porro haec est ea, que et meliorem ordinem, et compendium facilius, et majorem perspicacitatem admittit. Alteram quidem contemplans oculus, elegantia ac concinnitate nature oblectatur: in altera, geometrici cujusdam diagrammatis, accurati quidem exactique, sed nudi, sed inornati similitudinem agnoscit. Illa rem quamlibet spectatori, tanquam presentem, subjicit, et dat percipere: haec tantummodo describentis vice fungitur, remque delineatam in animi conspectu ponit. Prior autem posteriori in hoc prestat maxime, quod ea, que revera visa sunt, exhibens, speciem quandam veritatis præ se fert: et fit plementum æque nescia fallere, ac illud ipsum quod assimilatur.

De hoc opere, quod naturæ ipsius speciem proponit, alii fortasse aliud judicium ferent. Nisi enim auctorem fallat vana spes sua atque opinio, multi sunt apud quos ille nec censebitur operam inutilem posuisse, nec inepte erogasse pecuniam; apud quos scilicet figurarum magnitudo, elegantia, ac varietas aliqua in laude ponentur. Contra, non deerunt, quibus magna pars sumptus supervacanca omnino atque infructuosa fuisse videbitur. Contentant scilicet operis hujus utilitatem ad plures potuisse pertingere,

P R E F A C E.

The original plan having been only to publish the first ten plates, as a ground-work for farther improvements in this branch of anatomy, to be added whenever good opportunities should be offered, the author now began to entertain hopes of being able to give a much more complete work. He foresaw that, in the course of some years, by diligence he might procure in this great city, so many opportunities of studying the gravid uterus, as to be enabled to make up a tolerable system; and to exhibit, by figures, all the principal changes that happen in the nine months of utero-gestation. Such is the work which, at length, is offered to the public. The execution of it has indeed taken up more time than what was at first expected: but it gives the author no small satisfaction to reflect, that the delay of publication has contributed not a little to the value of the work.

That one person should have had so many opportunities of illustrating this subject, has been owing to fortunate circumstances, as well as to the assistance of many friends. Even since the last plate was finished, he had an opportunity of making a drawing of a younger embryo than he had before seen; and he has likewise made some figures from a very curious case, which he lately met with, namely, a conception in the fallopian tube. These, and whatever may hereafter occur, he will offer to the public upon some future occasion; probably in the way of a supplemental plate, or with the anatomical description of the gravid uterus, which he proposes to publish at full length. He has not hitherto had leisure to finish that part of the work. But if he should be prevented from doing it, by any unforeseen accident, it will be in the power of many gentlemen of the profession to do it for him, as he has constantly explained his observations on this subject in his public lectures.

ANATOMICAL figures are made in two very different ways; one is the simple portrait, in which the object is represented exactly as it was seen; the other is a representation of the object under such circumstances as were not actually seen, but conceived in the imagination. Bidloo has given us specimens of the first kind; Eustachius of the latter.

That figure which is a close representation of nature, and which is finished from a view of one subject, will often be, unavoidably, somewhat indistinct or defective in some parts: the other, being a figure of fancy, made up perhaps from a variety of studies after Nature, may exhibit in one view, what could only be seen in several objects; and it admits of a better arrangement, of abridgement, and of greater precision. The one may have the elegance and harmony of the natural object; the other has commonly the hardness of a geometrical diagram: the one shews the object, or gives perception; the other only describes, or gives an idea of it. A very essential advantage of the first is, that as it represents what was actually seen, it carries the mark of truth, and becomes almost as infallible as the object itself.

With regard to this work, which is a faithful representation of what was actually seen, the judgment of the public will probably be divided. Many will approve of the labour and expense which have been bestowed upon it, and commend the largeness, elegance, and variety of the figures. Others will think that a great part of the expense might have been spared, and the work thereby rendered of more general use, if the figures had been made to a smaller scale, if the engravings had been less finished,

P R O C E M I U M.

si figure omnes in formam arctiorem contractae fuissent; si sculputra manu minus delicata fuisset elaborata; si aliquot etiam figure, utpote alias non absimiles, omnino omissae.

Cum autem figurae, de quibus agitur, ea mente atque consilio fingi soleant, ut ab iis, quantum fieri potest, explicetur tam vera natura partium (nempe earum habitudo et structura) quam forma externa, positura, ac nexus; magnitudo illis aliqua concedatur necesse est; aliter enim partes, ut aiunt, componentes haud satis distincte delineari possunt. Quod si magnitudo rei propriis sculputræ quodammodo apta sit ac conveniens, ea ipsa, utpote quæ observatu dignissima est, haud dubio eligenda est precipue. His de causis figure omnes, quotquot hic proferuntur, ad justum naturæ modum perfectæ sunt, paucis tantum exceptis, quæ in formam naturali angustiorem coguntur, nec non una atque altera ex rebus minutissimis, quas auctor idcirco ampliandas curavit.

Iisdem etiam de causis constat hujuscemodi figuris deberi accusationem quandam atque elegantiam. Profecto in pictura saepè fit, ut oculi levissima adumbratione oblectentur; maxime ubi vel ipsum tabule argumentum adeo per vulgatum est, ut vis imaginandi facile excitetur, id, quocunque defuerit, statim adimpleatur (sicuti iis in tabulis, ubi hominum effigies, aut aliae res, pariter note, exhibentur) vel ubi partes rerum minutiæ aciem pene oculorum fugiant, ut ea in tabula chorographica, quæ, propter distantiam suam, parum cernuntur. Sed in anatomia, ut et in rei naturalis historia, argumentum tabule spectatori vel omnino esse incognitum ponitur, vel non satis exploratum; partesque minimiæ, pariter cum majoribus, studium diligentiamque requirunt. Operam quidem dare, ut summa arte elaborentur et figurarum partes, per quas nihil, aut parum exprimitur, nisi quod prius sit expressum, hominis esset vehementer et opera sua et pecunia abundantis. Harum igitur figurarum pars maxima ultra mediocritatem perfecta est; aliquot summa expeditio atque artificio ornata; leviora, vel ea quæ prius fuerant aliqua ex parte illustrata, leviter tacta; ea vero quæ secunda vice proferuntur, lineis fere exterioribus adumbrata.

Cilibet homini, cui judicium sit paulo severius, in proclivi erit id auctori vitio dare, quod opus hoc haud satis justo ordine processerit, nec formam pressam satis atque concinnam induat. Quod si ille difficultates, quotquot auctorem circumvenient, contemplatione fuerit complexus, conamina hæcce, qualiaquæ sint, animo magis aquo excipiet. In plurimis aliis operibus, ad rem anatomicam que spectant, veniam sibi haud facile impetrat vel defectus aliqua ex parte, vel materies confuse permisteque disposita. Quivis enim, si modo laborem pati decreverit, primum quidem exercitatione et experientia intimam argumenti sui cognitionem adipisci poterit; deinde autem operis ejusdam, perfecti undique omnibusque numeris absoluti, formam atque imaginem animo effingere; et denique ex variorum cadaverum contemplatione rem institutam pro arbitrio suo ad exitum perducere.

Humani autem corporis uterum gravidum otiose secare paucis datum est. Pauci sunt, quibus, in omni vita spatio, nisi semel, aut etiam fortassis iterum hoc fuerit indulsum. Ponamus quidem talen occasionem, faustum omnino atque felicem, cultori cuiquam anatomiae esse oblatam; necesse est ut continuo et sine mora de ratione ille instituti fixum aliquod certumque destinet; et uno eodem tempore duo consilia, inter se non bene convenientia, prosequatur. Primum enim, ut ipse rem penitus perspectam habeat, cadaver sua manu socet necesse est; ita tamen instituenda est investigatio, ut artifici simul adstanti, præcipuas partes comode delineandi plena sit copia. Proinde vix fieri potest, quin consilium, optime susceptum, immutari oporteat, variisque rebus accomodari quæ de improviso et præter expectationem possint occurrere. Ut autem id rite perfici possit, opus est longi temporis mora; partes interea aeri externo, inter pingendum, expositaæ plurimum detrimeni capient; præcipue si vir talis negotio obeundo presit cui cune est, ne facti quidpiam, aut ex memoria deprompti, ipsi rei veritati inducatur, et ut nihil ibi locum habeat, nisi quod ex natura ipsius observatione, re alia nulla intercedente, arreptum fuerit.

P R E F A C E.

and if some of the figures, which are very similar to others, had been omitted.

Anatomical figures being intended to shew, as much as possible, the true nature, that is, the peculiar habit and composition of parts, as well as the outward form, situation and connection of them, should certainly be large; otherwise the smaller component parts cannot be distinctly represented; and if the natural size of the object be tolerably fit for an engraving, that must be of all others the very best, as it has the advantage of shewing such an important circumstance. Upon these considerations, all the figures in this work were made of the natural size, except a few which were reduced in size, and one or two of minute objects, which, on that account, were magnified.

For the reason already given it should seem obvious that anatomical figures ought to be likewise well finished. In many subjects of painting, indeed, the slight manner of producing an effect, without labour, is very agreeable; particularly when, either the subject is so well known, that a mere hint is enough to the imagination, which easily supplies all that is wanting; as, in designs where human figures and other well known objects are introduced; or, when the smaller parts of objects are not to be seen distinctly, as the objects, especially the more distant ones, in a landscape. But in anatomy, as in natural history, the subject is supposed to be new, or only imperfectly known; and the smaller parts are to be studied with care, as well as the larger masses. In those parts of anatomical figures, indeed, which are mere repetitions, or nearly so, the labour and expence of finishing may well be saved. Accordingly in this work the greater part is tolerably well finished, some very highly and delicately; matters of less moment, or approaching to repetitions, are executed in a slight manner; and what is merely a repetition, is commonly put down in bare outlines.

Any judicious person, who examines this work with care, will naturally think that the plan might have been more regular and more compact: but it is presumed that he will be less dissatisfied with the author's best endeavours, when he shall consider the difficulties which lie in the way, and render the subject almost unmanageable. In many other works of anatomy, imperfections, and particularly a want of method, are not readily excused. For, if the author will but take the pains, he may first of all make himself master of his subject by repeated dissections and experiments; he may, at his leisure, lay down a studied plan for a complete and regular work; and with his own convenience he may execute his plan from a variety of dead bodies.

But opportunities of dissecting the human pregnant uterus at leisure very rarely occur. Indeed, to most anatomists, if they have happened at all, it has been but once or twice in their whole lives. Upon such occasions, therefore, even when attended with the most favourable circumstances, the anatomist must fix upon a plan, without loss of time, and at once carry on two schemes which are hardly compatible; that is to say, he must dissect for his own information, in the first place, and yet conduct the enquiry so as to have good drawings made of the principal appearances: and it is more than probable that he must alter any plan that he might have proposed, and adapt it to a variety of circumstances in the subject that could not be foreseen; and much time must be lost, and the parts must be considerably injured by long exposure to the air before the painter; especially if the work be conducted by an anatomist who will not allow the artist to paint from memory or imagination, but only from immediate observation.

PROCEMIUM.

Si auctor tot opportunitates, hanc ornandi provinciam, sibi prospexisset futuras, quot ipsi bona sua fortuna indulserit, figuras ille quidem aliquas emendatores reddere potuisset; alias alius melius accommodare, et ita carum seriem in minorem numerum cogere. Figuras quidem nonnullas, jam delineatas, duasque tabulas, a sculptore prorsus perfectas, apud se prudens detinuit; ne operis scilicet moles atque impendium plus aequo increaserent. Et profecto plures, eadem ratione inductus, detinuisse, nisi ipsi mentem subiisset, longum fortasse fore tempus, priusquam aliquod figurarum systema, hoc suo minus imperfectum, in lucem sit exiturum. Potius igitur duxit culpae in periculum incurere, eo quod rem nimis copiose, quam quod parcus ac negligenter tractaverit.

Operi, nimis jam magnitudinis, nimique sumptus, internam fetus anatomiam adjicere, supervacaneum fore visum est; praesertim cum alii isto munere sunt functi; et cuiilibet, id qui de integrō meditatur, haec omnia cognoscendi atque tractandi hand facile defutura est occasio.

In arte typographicam sumptus, major solito factus est, non quidem solam ob elegantiam, qua operaceleberrimi artificis JOANNIS BASKERVILLE ornanturomnia; sed praeципue charte, qua ille utitur, atramentique ratione habita. Talis est enim charte hujuscē levor, atramentumque tam cito arescit, ut folia, literis licet plenisima, tabule cujusque interpositae nitorem integrum atque illāsum conservatura sint.

Ad summam, si auctori ea laus fuerit concessa, quod, in hac primaria parte humani corporis illustranda, neque opere sue, neque impensarum, neque temporis rationem habuerit, voti sane fit compos, nec aliud gratiae sibi apponi postulat. Harum autem lundum in partem plurimum merito suo evocandus est frater auctoris solertissimus JOANNES HUNTER; nam in plurimis caderibus secundis partem ille laborum sustinuit: cuius quidem viri acumen et diligentia in hujuscemodi p̄vestigationibus adeo omnium sermone celebrantur, ut de illo si sileretur adjutore, operis ipsius pretio atque existimationi parum esset consultum. Artificibus etiam singulis, quorum ingenii atque industrie monumenta hic extant, gratias agit quas habet maximas; p̄e ceteris autem viro optimo ac peritissimo artifici ROBERTO STRANGE, non modo quod is manu sua tabularum harum duas tanquam aternitati consecrari, sed etiam quod, ab incepto operis usque ad exitum, auctorem amicitia stabili atque incorrupta dignatus, vel consilium vel auxilium expetenti haud unquam defuerit.

P R E F A C E.

If the author could have foreseen the numerous opportunities which have fallen to his share, he might have easily improved some of the figures; and by adapting them better to one another, he might have completed the series in a smaller number. He has actually kept back several drawings which had been made, and two plates which had been engraved, that the work might not be overcharged; and would have withheld more, for that reason, if he had not thought that it would probably be long before a more perfect system of figures would be offered to the public. This consideration induced him to risk the being censured rather for having done too much than too little.

In a work which was already become too large and expensive, it was thought proper to omit the internal anatomy of the child, especially as that part has been executed by others, and opportunities of studying it may be easily procured.

The additional expense of Mr. BASKERVILLE's art was not incurred for the sake of elegance alone; but principally for the advantage of his paper and ink, which render a leaf of his press-work an excellent preservative of the plates between which it is placed.

If it be allowed that the author has spared neither labour, nor time, nor expense, in improving an important part of anatomy, this is all the merit which he can claim. In most of the dissections he was assisted by his brother, Mr. JOHN HUNTER, whose accuracy in anatomical researches is so well known, that to omit this opportunity of thanking him for that assistance, would be in some measure to disregard the future reputation of the work itself. He owes likewise much to the ingenious artists who made the drawings and engravings; and particularly to Mr. STRANGE, not only for having by his hand secured a sort of immortality to two of the plates, but for having given his advice and assistance in every part with a steady and disinterested friendship.



J. V. Rymsdyk delin.
F. S. Barron sculp.
TAB. I. Deem priores Saluile Malorum exhibent, non feri gravitatis. Inveniuntur subiecta morte extindam, alterius a venis circa diversi coloris
injecta distincte erant. In huius prima contenta. Abdomen, integumentis decapsulatum incisis et reflexis, ab inspectum veniant. Raspiniatum est ita, ac
sive situ, cadaver, ut superior Irani pars haud paulo est clausa, et viscera inde abdominis ducius contemplanda obicerentur.

T A B U L A I.

D E C E M priores tabulae mulierem exhibent, nona fere graviditatis mense exeunte, ann. 1750, subita morte extictam. Arteriae a venis, cera diversi coloris injecta, distincte erant. In hac prima, coptenta abdominis, integumentis decussatum incisis et reflexis, sub conspectum veniunt. Resupinatum est acclivi situ cadaver, ut superior trunci pars hand paulo eset elatior, et viscera inde abdominis clarius contemplanda objicerentur.

Femora et pudenda, utpote partes externae, eam, quam cæteræ, notationem non desiderant. Pudenda vero, in hoc cadavere, ut ex figura concipi potest, aqua inter cutim intumuerunt.

A A Pars anterior thoracis infra mammas, qua trunca erat transverse discussus, figuram a parte superiori terminans.

B B Integumentis, musculis, et peritoneo decussatum incisis, eorum anguli duo superiores in marginem thoracis cartilaginosum reflexi.

C C Eorundem anguli duo inferiores, in inguen, femur, et ossis ilium spinam anteriorem, utrinque reclinati. In angulis illis, ad marginem inferiorem medio femori oppositum, observare licet partem rugosam et teusam, qua filum adhucbitum fuit, ea mente, ut, angulo detracto, uterus et partes vicinae manifestius apparent.

D Incisionis longitudinalis, ad scrobiculum cordis a mucrone cartilagineo ensiformis ductæ, extremitas superior.

E Ejusdem incisionis, ad symphysin ossium pubis, extremitas inferior.

F Hepatis ligamentum falciforme, super faciem internam musculorum abdominis, ad illius visceris fissuram, magnum inter lobum et parvum, protensum.

G G Arteria et vena epigastrica, in utrisque muscularum angulis inferioribus trans peritoneum sese projiciens.

H Hepatis lobus sinister, in parte summa regionis epigastricae situs. Illius in superficie, arterie quedam parvæ, ex substantia illius glandulosa exeunt, et in peritoneum quo vestitur diffusæ, delineantur.

I Omentum in regione epigastrica situm, intestinis tenuibus incumbens. Vasa quamvis cera vividi coloris impleta, ob partis adipem cerni vix potuerunt.

K Omenti pars inferior et media, ab utero antea elevata, quæ in plicaturas multas quam minimas contracta jacuit et compressa.

L Omentum a latere dextro, post partem eam uteri, quæ incipit tuba dextra, descendens.

M Omentum quod in latere sinistro ante tubam fallopiam, &c. descenderat, scorsum positum, quo partes ex in conspectum prodirent.

NN Duo intestinorum tenuium gyri, quorum utriusque, maxime autem inferior, omenti parte praedicta tecti fuerant.

O Uterus, regionem totam umbilicalem et hypogastricam possidens. Superficies ejus vasorum rami tenuibus intertexta: horum pleraque vene. Multa sanguine sollemmodo impleta, cadavere diutius asservato, paulatim evanuerunt. Situs uteri paulo obliquus, et ad latus dextrum. Umbilicus parte qua stat haec litera aliquanto altior fuit. Forma uteri quodammodo irregularis; aliae ejus partes aliis eminentiores. Id ex eo oritur, partum quod visceribus viciniis incumbens, præsentim ad fundum, se ad corum formam fixerat; partum etiam quod contenta ejus formam irregularem pre se tolerant.

P Tumor ad latus sinistrum uteri, in ea scilicet parte cui media placenta interne adhaerescebat.

Q Uteri tumor in latere dextro, ubi fortis nates jacebant, ubi etiam fetus per substantiam ejus satis distincte tactu sentire licuit. Undique circum hanc partem uterus aliquanto depresso, et sacculi instar aqua repleti, tactum afficiebat.

RR Ligamenta rotunda. Sinistrum, propter uteri situm obliquum, dextro longius. Exitus amborum, angulis inferioribus integumentorum &c. filo deductis, in conspectum veniebat. Quæ strictræ ibi videtur, nihil aliud est, quam peritonei in forma theœ duplicitatio et protrusio, ea ligamenta per paululum spati sequens, quæque ab eo peritonei processu, funem spermaticum canis vel tauri comitate, gradu tantum differt.

SS Tube fallopiæ. In latere dextro, solum fere tube principium cernere licet, parte reliqua post uterum descendente. In latere sinistro, tube media pars sola in conspectum venit; tegitur enim ejus principium intestino ibi posito; finemque ejus, vel fimbrias, vasa spermatica occupant. Ea vasa, in latere sinistro, satis aperte conspicuntur; spatium illud totum inter ligamentum rotundum et tubam occupant: vas parvum arteria; magnum ei adhaerens vena. Utraque ad uteri latus, in peritonei duplicatione illa, quæ ligamentum latum appellatur, procedunt.

P L A T E I.

The first ten plates were made from the dissection of a woman, who died suddenly, in the end of her ninth month of pregnancy, in the year 1750. The arteries and the veins were injected with wax of different colours.

This plate represents the object, as it appeared when the abdomen was opened by a crucial incision, and the four angles of the containing parts turned outwards; the subject lying on its back, but so that the upper part of the trunk was considerably higher than the rest.

The thighs and pudenda require no explanation: the latter of these were edematous in the subject, as they appear in this figure.

AA The forepart of the chest, just below the breasts, where the trunk was cut through, and which therefore terminates the figure above.

BB The two upper angles of the integuments, muscles, and peritoneum, turned back over the cartilaginous margin of the chest.

CC The two inferior angles, turned down over the groin, thigh, and anterior spine of the os ilium, on each side. In each of these, at its lower edge and opposite to the middle of the thigh, a part may be observed to be stretched and puckered, where a thread was fixed, which pulled the flap down, in order more fully to expose the womb and adjacent parts.

D The upper end of the longitudinal incision, at the pit of the stomach, beginning just at the point of the ensiform cartilage.

E The lower end of the same incision carried down to the symphysis of the ossa pubis.

F The umbilical or falciform ligament of the liver, which runs upwards, on the inside of the abdominal muscles, to the fissure between the great and the small lobe of that viscus.

GG The epigastric artery and vein, projecting through the peritoneum, upon each of the lower angles of the muscles.

H The small lobe of the liver, situated in the highest part of the epigastric region. Upon its surface are represented some small arteries, which emerge from the glandular substance, and are diffused in the peritoneum that covers it.

I The omentum, spread over the small intestines in the epigastric region. The vessels of this part, though injected, were hardly visible, being obscured by fat.

K The lower and middle part of the omentum, which had been pushed up by the womb, and lay in numerous small folds pressed together.

L The omentum, in the right side, passing a little way down behind that part of the womb from which the right tube begins.

M The omentum, in the left side, which came down before the fallopian tube, &c. and which was thrown a little aside, that those parts might be seen.

NN Two turns of the small intestines; both of which, but especially the lower, were covered by the above-mentioned part of the omentum.

O The womb, possessing all the umbilical and hypogastric regions. Its surface is interspersed with a number of small vessels, the greatest part of which are veins. They were filled with blood only, and therefore gradually disappeared after the subject had been preserved some time. Its situation is a little oblique, and towards the right side. The navel was a little higher than the part upon which this letter is placed. The form of the womb is a little irregular, at some places projecting more than at others. This seemed partly owing to its being moulded upon the neighbouring bowels, particularly at its fundus; and partly to its contents; whence,

P A swelling towards the left side of the womb, where the middle of the placenta adhered; and

Q A swelling on the right side of the womb, where the buttocks of the child lay; and where the child could be felt, pretty distinctly, through its substance. All around this part the womb was somewhat depressed, and felt like a bag of water.

RR The round ligaments. The left is longer than the right, because of the oblique situation of the womb. The exit of each was brought to view by drawing down the lower angles of the integuments, &c. with threads: and the appearance of a stricture there, was occasioned by a sheath-like doubling and protrusion of the peritoneum attending those ligaments some little way; which differed only in degree from that process of the peritoneum, which accompanies the spermatic rope in a dog, or bull.

SS The fallopian tubes. In the right side, little more than the beginning of the tube is seen, the rest running down behind the womb. Of the left tube the middle part only is exposed; its beginning at the womb being concealed by the gut which lies upon that part, and the end or fimbriae being covered by the spermatic vessels. Those vessels are pretty distinctly seen in the left side: they fill up all the space between the round ligament and the tube: the small vessel is the artery, and the great, to which that clings, is the vein: both of them are passing up to the side of the womb, in that doubling of the peritoneum called, the broad ligament.

T A B U L A II.

C O N S P E C T U S viscerum abdominis a latere dextro, partibus continentibus thoracis et hypochondrii, ac maxima omenti parte sublatiss.

- A A* Femora, ubi transverse scissa, linteo cooperata.
- B* Spina, supra diaphragma, transverse discissa.
- C* Costa octava pars.
- DD* Integumenta et musculi, ad partem thoracis posteriorem, eo loco quo negligenter sejunctus erat truncus, rescessi.
- E* Musculorum abdominis angulus inferior, in femur, ut in praecedente figura, devolutus.
- F* Montis, ut dicitur, veneris pars, una cum musculis abdominis, devoluta.
- G* Ad sinistrum latus pars huic respondens.
- H* Thoracis margo, ad latus scrobiculi cordis sinistrum, peritoneo et musculis inversis, tectus.
- I* Cartilago mucronata.
- KK* Costae inferiores, partesque cæteræ continebentes sectione longitudinali sejunctæ, adeo ut, iis que sectionem et scrobiculum cordis interjacerent remotis, viscera hypochondrii dextri sese vindenda offerrent.
- L* Pars ima atque posterior cavi dextri thoracis.
- M* Aorta descendens, eo loco, quo cadaver transverse discissum fuerat, deligata; nec non, colorata cera impleta.
- N* Oesophagus ibidem deligatus.
- O* Superficies superior convexa diaphragmatis; quod, pleura tectum, in situ suo naturali manet.
- P* Diaphragmatis centri tendinei pars qua in dextro latere sita est.
- Q* Vene cave inferioris trunca, juxta diaphragmatis superficiem superiorem deligatur.
- RR* Pleuræ sectæ margo, loco quo reflexa de superficie superiori diaphragmatis, faciem internam costarum contegebatur.
- SS* Peritonei secti margo, loco quo scilicet a superficie diaphragmatis inferiori, membrana investiens more, ad faciem internam musculorum abdominis cursum tenebat.
- Superficies inter margines pleuræ (RR) et peritonei (SS) connexio est diaphragmatis ex facie interna costarum disrupta.
- T* Facies interna musculi transversi peritoneo tecta.
- U* Hepatis lobus magnus, vel dexter; in cuius crepidine, ex adverso litera U, exiguae videre est fossulan, quam vesicula fellis figura sua aptaverat, facie interne hepatis hic loci apposita.
- V* Hepatis lobus parvus, in hypochondrium sinistrum sese extendens.
- W* Ligamentum hepatis falciforme, seu vene umbilicalis reliquæ, loco quo fissuram hepatis intrat, recisum.
- X* Colon ex caeco ad hepatis sursum transiens.
- YY* Colon, cum appendiculis suis adiposis, sinistrorum currens, hepatis inter et intestina tenuia.
- zz* Omenti principium vel radix, quod ex intervallo hepatis et intestinorum tenuis nascitur.
- aaaa* Illorum intestinalium gyrum aliqui, omento paulo infra marginem hepatis excisso, in conspectum veniunt.
- b* Uterus. In facie ejus externâ, venas, aliquantulum obscurias, notare licet. Hæc, cum cadaver primum apertum posui, cera etiæ implete, colore tantum distinguebantur. Tempore autem quo tabula hæc erat delineata, venæ, parte jam arefacta, sese magis projiciebant, ut in figura quoddammodo se videndas offerent.
- c* Ligamentum rotundum in inguen descendens.
- d* Tuba fallopiana, juxta faciem internam osis ilium retrorsum currens.
- e* Tubæ extremitas fimbriata, pone latus uteri jacens, hic paulum educta, nempe ut conspiciantur. Ligamentum rotundum inter et tubam, vasa spermatica, praesertim vena, conspiciantur, ad partem uteri anteriorem, cui placenta adhaerebat, præcipue tendentia.
- f* Utteri angulus prominens, loco quo fetus nates, utero nondum adaptato, tactu sentire licet.
- N. B. Costis notis lateris dextri remotis, illæ quo ad latus erant sinistrum, ex vi resiliendi, marginem thoracis (*H*) et exinde cartilagineum ensiformem elevarunt; unde facies ejus anterior in dextrum latus oblique tendebat.
- Figura hac nondum delineata uterum observavi sensim subse disse; quapropter ope quam minimæ cannulae, uterum, spiritu vinoso injiciendo, in pristinam figuram restituere conatus sum.


P L A T E II.

A V I E W of the same object from the right side, after the upper abdominal flap, and the containing parts of the right hypochondrium had been removed, that the whole mass of the abdominal viscera might appear in its natural situation, the subject lying on its back.

- AA* The thighs, covered with a cloth where they had been cut through.
- B* The back-bone cut through above the diaphragm.
- C* Part of the eighth rib.
- DD* The integuments and muscles at the back part of the chest, cut rudely through in severing the trunk at that place.
- E* The inferior flap of the abdominal muscles, &c. turned down over the thigh, as in the preceding figure.
- F* Part of the mons veneris turned aside with the abdominal muscles.
- G* The respective part of the left side.
- H* The margin of the chest at the left side of the scrobiculus cordis, covered by the peritoneum and muscles, which are turned over it.
- I* The ensiform cartilage.
- KK* The lower ribs and other containing parts, cut down longitudinally, by which means all the parts, between this section and the scrobiculus cordis were removed, that the viscera of the right hypochondrium might be exposed to view.
- L* The posterior and lower part of the right cavity of the chest.
- M* The trunk of the descending aorta, tied up where it was cut, and filled with injection.
- N* The oesophagus, tied up at the same place.
- O* The upper convex surface of the diaphragm, which remains, covered by the pleura, in its natural situation.
- P* That part of the centrum tendinosum of the diaphragm which is in the right side.
- Q* The trunk of the lower cava tied, close to the upper surface of the diaphragm.
- RR* The cut edge of the pleura, where it was reflected, from the upper surface of the diaphragm, to line the inside of the ribs.
- SS* The cut edge of the peritoneum, where it was continued as an investing membrane, from the inferior surface of the diaphragm, to the inside of the abdominal muscles.
- The surface which is seen between the edge of the pleura (RR) and that of the peritoneum (SS) is the attachment of the diaphragm cut off from the inside of the ribs.
- T* The inside of the transversus muscle covered with the peritoneum.
- U* The great, or right lobe of the liver; at the lower edge of which, over against the letter U, there is a small notch formed by the concave bed of the gall bladder, which lies upon the inside of the liver at this place.
- V* The small lobe of the liver, which stretches over into the left hypochondrium, and is therefore seen in perspective.
- W* The falciform ligament, or remains of the umbilical vein, cut off as it enters the fissure of the liver.
- X* The colon passing up from the caecum to the liver.
- YY* The colon with its appendiculus adiposus running towards the left side, between the liver and small guts.
- zz* The beginning or root of the omentum, coming downwards from the interspace of the liver and small guts.
- aaaa* Some turns of those small intestines brought to view by cutting off the omentum a little below the edge of the liver.
- b* The womb. Upon its surface we can trace, though obscurely, its veins. These, though injected, were distinguishable only by their colour when the subject was first opened: But when this drawing was made, the part, having been some time in the air, had become a little dry, and the veins projected, as they appear in the figure.
- c* The round ligament running down to its passage into the groin.
- d* The fallopian tube running backwards on the inside of the os ilium.
- e* The fimbriated end of the tube, lying behind the side of the womb; but drawn a little out, that it might be seen.
- Between the round ligament and the tube, are seen the spermatic vessels, especially the vein, which go chiefly, or by their largest branches, to the forepart of the womb where the placenta adhered.
- f* The prominent corner of the womb, where the buttocks or loins of the child were distinctly felt before the part was opened.
- N. B. The false ribs of the right side having been removed, those of the left side, by their elasticity raised up the margin of the chest (*H*) and consequently the ensiform cartilage; the anterior surface of which was thereby turned obliquely to the right side.
- Just before this figure was made, I had observed that the womb had insensibly subsided, and therefore by the cannula of a very small trocar I injected some spirits to raise it up, as nearly as I could guess, to the figure it had when the abdomen was first opened.

TAB. II. *Conquedas micrum-dileminatum a latere dextra, partibus continentibus Thoracis et Hypochondri.*

ad matrem omni parte sublatum.









J. C. G. 1800

TAB. III. Sectionem ventri abdominalis a sinistro latere, nisi cum Diaphragmate in superficie facile apparet in delineanda hoc figurari.
Injectum oculi fixiss. a parte lateris superiori obliquè dorsum, utrusque quippe punctorum senum doctrinam viuum intercludat.

T A B U L A III.

- PARTIS TRUNCI INFERIORIS PROSPECTUS;** oculo scilicet a parte sinistri lateris superiori, oblique deorsum ita collinato, ut universa viscerum compages, una cum diaphragmate, in situ suo naturali apparet.
- A* Femur sinistrum. Uteri prominentia femur dextrum oculo subtrahit.
B Clavis sinistra.
C Integumentorum et muscularum abdominis, lateris sinistri, angulus inferior in inguen et femur inversus.
D Integumentorum et muscularum abdominis sectio transversa.
EE Partium abdominis continentium sectio longitudinalis, ab ilium osse sursum ducta. Hac sectione, angulus superior partium continentium omnino auferitur, ut viscera hypochondrii sinistri in conspectu ponantur.
- FF* Partium thoraci continentium posteriorum sectio transversa.
G Vertebra transverse scissa.
H Medulla spinalis.
II Cavorum thoraci partes inferiores postice, quae, pulmonibus sublatiss., vacue conspici possunt.
KK Diaphragma superficiem convexam hepatis complectens.
L Cartilago mucronata, diaphragmati adhuc adhaerens.
M Diaphragmatis ad latus dextrum convexitas magna; quae in hoc cadavere, parte huic consentanea in latere sinistro elatior comparuit. Ibi etiam vasa nonnulla conspicuntur; quorum majora, vena scilicet phrenica, cera impleta sunt a vena cava; hinc altius extantia.
N Diaphragmatis superficies, qua pericardio annectitur, quæque eorū sustinet.
O Vena cava, ubi per diaphragma transit, deligata.
P Oesophagus supra diaphragma deligatus. Hunc cera injecta, dum per arterias fluebat, materia colorante amissa, in ventriculum transiens, implevit.
Q Aorta, eodem modo, ubi corporis truncus transverse fuerat scissus, supra diaphragma, deligata.
 Oesophagus et aorta, una cum spine prominentia, cavum satis magnum in diaphragmatis parte postica et superiori efficiunt.
- RSTU* Diaphragma in hypochondrio sinistro recisum. Ex parte posteriori, labia (*TU*) ejusdem fissure in diaphragmate mutuo reedunt; et ex parte anteriori, anguli (*RS*) inversi sunt, nempe ut viscera hypochondrii sinistri in conspectum veniant.
- V* Hepatis extremitas sinistra.
W Ventriculi extremitas sinistra.
XX Lienis margo anterior denticulatus.
 a Hepatis lobus dexter.
 b Ligamentum falciforme, fissuram hepatis intrans.
 c Hepatis lobi sinistri ea pars quæ in regione epigastrica sedem habet.
dddd Omenti radix dextrorum de ventriculo et colo, hepar inter et intestina tenuia, descendens.
- eff* Omenti radix in hypochondrio sinistro ita disposita, ut cavum suum exhibeat. Ea pars, quæ marsupii omenti anteriorem partem facit (*ee*), quæque de ventriculo descendit, ventriculo jam et hepatis incumbit replicata: omentum autem pars posterior, quæ colo connectitur (*ff*) ante intestina tenuia in proprio manet sita.
- gg* Omenti radix, ventriculum inter et lienum.
hh Omenti pars ad latus sinistrum, in situ suo naturali, tenuia inter intestina et peritoneum faciem internam transversalis abdominis integens.
- iii* Intestinorum tenuium gyri multo inferiores et frequentiores hac in parte, quam eorum, quæ in latere dextro jacent; partim propter obliquitatem uteri, præcipue autem quod in latere dextro hepatis moles minus spatii intestini permittit.
- k* Uterus. Tempore quo figura haec delineabatur, pars uteri superior arescere coepit, unde vasa infarcta, magis conspicua se exhibebant: minorum pleraque arteriae sunt; majora omnia vene.
- l* Ligamentum rotundum.
m Tuba.
n Arteria spermatica ramus magnus, trans ligamentum latum ad uterum repens.
oo Ramorum, vena scilicet spermaticæ, majorum fasciculus, per ligamentum latum ad uterus ascendentium.

P L A T E III.

- A VIEW**, from the left side and downwards, of the lower part of the trunk; so prepared, as to shew the whole mass of the bowels, with the diaphragm, in their natural situation.
- A* The left thigh. The right thigh is covered from our view by the projection of the womb.
B The left buttock.
C The lower flap of the integuments and abdominal muscles on the left side, turned down over the groin and thigh.
DD The section of the integuments and muscles, made by the transverse incision of the belly.
EE A longitudinal section of the containing parts of the belly, carried upwards from the os ilium, or posterior extremity of the transverse incision, parallel to the spine; whereby the upper flap of the containing parts is entirely removed, to shew the viscera of the left hypochondrium.
- FF* A transverse section of the posterior containing parts of the chest.
G The body of the vertebra cut through.
H The spinal marrow.
II The lower posterior parts of the cavities of the chest, which are seen empty, the lungs being removed.
KK The diaphragm, embracing the convex surface of the liver.
L The ensiform cartilage left adhering to the diaphragm.
M The great convexity of the diaphragm in the right side, which in this subject was much higher than the analogous part in the left. Upon this part, some vessels are seen: the larger are the phrenic veins, which were filled with wax from the cava, and are therefore prominent.
N The surface of the diaphragm, where the pericardium is united with it, and upon which the heart rests.
O The vena cava tied, as it passes through the diaphragm.
P The oesophagus, tied above the diaphragm. It was filled by the injection, which passed without the red colour, from the arteries into the stomach.
Q The aorta, in the same manner tied above the diaphragm, where the trunk of the body was cut through.
 The oesophagus and aorta, as well as the projection of the spine, make a considerable notch upon the upper posterior part of the diaphragm.
- RSTU* The diaphragm, slit up in the left hypochondrium. At the posterior part, the lips of the fissure in the diaphragm (*TU*) recede from each other; and forwards, the corners (*RS*) are inverted, to shew the viscera in the left hypochondrium, viz.
- V* The left extremity of the liver;
WW The left, or great extremity of the stomach; and
XX The spleen: its anterior edge indented.
 a The right, or great lobe of the liver.
 b The falciform ligament, entering the fissure of the liver.
 c That part of the small lobe of the liver which lies in the epigastric region.
- dddd* The root of the omentum towards the right side, coming down from the stomach and colon, between the liver and the small intestines.
- eff* The root of the omentum in the left hypochondrium, so disposed as to shew its cavity. What makes the anterior part of the pouch of the omentum (*ee*) which comes down from the bottom of the stomach, is turned up over the stomach and liver: and the posterior part of the omentum (*ff*) which is fixed to the colon is left, in its natural situation, before the small intestines.
- gg* The root of the omentum, between the stomach and spleen.
hh Part of the omentum in the left side, in its natural situation, between the small intestines and the peritoneum which lines the transversalis muscle.
- iii* The convolutions of the small intestines, which are considerably lower and in greater quantity in this side than in the right; partly because of the obliquity of the womb, but chiefly because in that side the liver comes low down, and leaves little space for intestines.
- k* The womb: which had begun to dry at its upper part when this figure was taken; so that the injected vessels began to appear more remarkably. The smaller are chiefly arteries; the larger are all veins.
- l* The round ligament.
m The tube.
n A large branch of the spermatic artery going across the broad ligament of the womb.
oo A group of large veins from the spermatic, mounting in the broad ligament to the womb.

T A B U L A IV.

UTERUS, et contenta pelvis a parte anteriori: ossa pubis, cum integumentis, et superiori pudendorum parte, omnino fuerant absissa, quo cervix uteri, et vesica simul urinaria ei praetensa, oculo incidenter.

- A A** Femora transverse abscissa.
- B B** Musculorum abdominis, et peritonei, anguli inferiores, in latera reflexi.
- C** Uterus; cuius in tota superficie, perque substantiam, vasa cera impleta, et exinde prominentia, conspicuntur.
- D D** Os pubis utrinque supra foramen magnum ischii resecatum.
- E E** Ischii processus parvus, eo in loco ubi ossi pubis committitur resectus.
- F** Perineum.
- G G** Labia pudendi: horum pars superior, nymphæ etiam, et clitoridis extremitas, absinduntur.
- H H** Caruncula myrtiformis insignis.
- I** Vagina. In parte hujus introitus inferiore (nigriore scilicet in hac tabula), vaginae cavum, tam retrorsum quam sursum ducens, representatur. Ad literam (*L*) conspicitur vagina pars superior et anterior, que cum vesica urinaria et urethra connectitur, quaque a visceribus incumbentibus deprimitur. Ad punctum etiam orificii hujuscem, fusi, ovati, superius, urethrae extremitas prominens, meatusque urinarii orificio conspicuntur.
- K** Clitoridis corpus transverse discissum: in medio, duas arterias fere contiguas, scilicet in utroque corpore cavernoso unam, cernere licet.
- L L** Clitoridis crura et erectores. Haec, ossibus pubis quibus annectuntur sublati, magis horizonti ad libellam, quam in situ suo naturali, respondentia jacent. Sejunctione autem ossium pelvis, nempe ut contenta ejus melius possint conspicui, extenduntur.
- M** Vesicae urinariae pars superior, simul utero, pone sito, et caput fetus continent, simul ossibus pubis unitis, eam exadversum ambientibus, compressa.
- NN** Peritoneum reflexum, telaque cellulosa, latera integens pelvis, et ei uteri parti quam ossa pubis amplectuntur, circumjacent.
- O** Ex peritoneo et tela cellulosa resectis, margo, eo loco ubi de vesica urinaria ad faciem internam muscularum abdominis tegendam, supra ossa pubis, ascendunt.
- P P** Ligamenta rotunda, abdominis intra cavum.
- QQ** Ligamenta rotunda, eo in loco ubi ex abdomen recedentia sese in montis veneris lateribus sensim recondunt.
- RR** Musculi femoris, ossibus pubis exorti, rescissi.

P L A T E IV.

A FORE-VIEW of the womb, and of the contents of the pelvis; the ossa pubis, with the muscles and integuments which cover them, being removed.

- A A** The thighs cut through.
- B B** The lower angles, or flaps of the abdominal muscles and peritoneum turned aside.
- C** The womb. All over its surface the injected vessels are seen projecting through its substance.
- D D** The os pubis of each side cut through, above the foramen magnum ischii.
- E E** The small branch of the ischium which joins the os pubis, cut through.
- F** The perineum.
- G G** The labia pudendi. The upper part of those, the nymphæ and the extremity of the clitoris are cut off.
- H** A large caruncula myrtiformis.
- I** The vagina. At the lower part of this passage, the deeper black part of the surface, in this figure, represents the cavity of the vagina leading backwards as well as upwards: and where this letter (*J*) stands, is seen the upper or anterior part of the vagina which is united to the bladder and urethra, and which is pressed downwards by the viscera that lie over it. At the upper point of this dark oval orifice the tuberous extremity of the urethra is seen, with the orifice of the meatus urinarius.
- K** The body of the clitoris cut through; in the middle of which two small arteries are seen almost contiguous, viz. one in each corpus cavernosum.
- L L** The crura and erectores clitoridis. These lie more horizontally than in the natural state, the ossa pubis, &c. which suspend them, being removed: and they are stretched or lengthened, from the bones of the pelvis being drawn a little asunder, that the contents might be better seen.
- M** The upper part of the bladder, which was compressed between the womb containing the child's head, which lay behind it, and the united ossa pubis, which had embraced it forwards.
- NN** The reflected peritoneum and cellular membrane, which lines the sides of the pelvis, surrounding that part of the womb which was embraced by the ossa pubis.
- O** The cut edge of the peritoneum and cellular membrane, where these ascend from the bladder to line the abdominal muscles above the ossa pubis.
- P P** The round ligaments within the cavity of the abdomen.
- QQ** The round ligaments after they have passed out of the abdomen, where they are insensibly lost in the sides of the mons veneris.
- RR** The muscles of the thigh, which spring from the ossa pubis, cut through.



J. C. Rymsdyk delin.
R. Strange sculp.
TAB. IV. Uterus, et contenta Pilvis a parte anteriori, ossa pubis, cum integumentis, et superiori pudendorum parte,
omnino fuerant abscisæ, quæ cervix uteri, et vesica, simul urinaria ei prætensa in visum incurrent.







S. S. Agnew delin.
J. D. Collyer sculp.
TAB. V. Uteri pars anterior et dextra, autem manu ab adhaerentibus secundinis separata, et in sinistrum latus reflexa. Id eo factum
est consilio, ut pars membranæ decidua ac Placenta, cum vasis carunculis corâ repletis, et superficies uteri, cui adhaeserant,
interna eadis expulerentur. Partes lencis designatas proximi sequens tabula sicut integras.

T A B U L A V.

- U**TERI pars anterior et dextra, cauta manu ab adhaerentibus secundinis separata, et in sinistrum latus reflexa. Id eo factum est consilio, ut pars membrana decidua ac placente, cum vasis excurrente cera repletis, et superficies uteri cui adhaerenter interna, oculis exponerentur. Partes lineis designatas proxime sequens tabula exhibit integras. Priorem autem hic habet locum linearum de sequenti tabula explanatio, ne literas insculptae nimis illam obscurarent.
- A A* Femorum superiora.
B B De angulis inferioribus partium continentium abdominis, quod superest, in ossa ilium reflexum. Anguli utrinque, peritoneo vasis distincto integrantur; quod vero, ex parte, praesertim in latere sinistro, ad cursum vasorum epigastricorum retegendum, excidebatur.
C C Ossa pubis supra foramina ischium transverse scissa.
D D Ischium processus ascendentis transverse scissi.
Ossa hic pelvis, symphysi pubis sublata, laxa evaserunt; et in latere sinistro, ad descensum uteri in cavum pelvis magis illustrandum, extrosrum fuerunt evusa.
E E Arteriae inguinales.
F F Vena inguinalis.
G G Arteriae epigastricae.
H H Vene epigastricae.
I I Arteria obturatoria sinistri lateris, epigastrice scilicet ramus.
K K K Ligamenta rotunda ex abdomen ad exteriora vasorum epigastricorum descendenta.
L L Sectio integumentorum et muscularum, in utroque inguine, facia minima, dum partes pelvis anteriores continentis amovebantur.
M M Perineum.
N N Labiorum pudendi partes inae.
O O Hymenis reliquiae.
P P Vagina.
Q Q Clitoridis crura.
R R Urethra ex parte superiori in longum diffissa et expassa.

- SSSS* Vesica urinaria, anteriorem inferioris extremitatis uteri portionem ambiens.
T T Vesica urinariae cavum, partem ejus anteriorem ab urethra sursum incidendo, oculis subjectum.
U U Uteri extremitas parva vel inferior, in cavo pelvis posita. Ea pars quo circundata fuerat margine pelvis, satis evidenter constricta fuit: quam super uterus statim eminentior evasit, quippe partes que facilius cedunt enim ibi amplectebantur.

- VVV* Uteri facies exterior.
X X Vasorum spermaticorum in latere sinistro chorus.
Y Y Uteri substantia, ad dextrum latus, ab fundo penitus dissecata. In hac sectione vasa resecta occurunt; quod autem ad figuram attinet inordinatum, quippe cera friabili adimpluta. Venae coloris sunt lucidioris ac grandes; exiliores arteriae, atque fuscae; fundum versus uteri utraque minores quam in parte inferiori, ubi a lateribus uteri ad placentam transibant. Uteri crassitudo cum magnitudine vasorum his in locis proportionis legem quodammodo servata.
Z Z Decidua, sive membrana exterior, ramulis venarum ab facie interna uteri ad eam transeuntibus distincta.
a a Hic decidua vasis referta et opaca, cauta manu sublata fuit. Nulla hic loci, vel in chorio, vel in amnio, venarum, ex injectione cerae, species fuit. Altera enim pars per alteram adeo perlucet, ut fetus cutem distincte cernere licet.
b b Externae superficie placenta portio, hic cera extra venas uteri effusa, illuc venis ipsis cera impletis, a facie interna uteri ad eam transeuntibus interstincta.
c c Arteria cera rubri coloris impleta, quo ab utero abrupta fuit, et semel circumvoluta, in placentam se immisit.
d d Ejus partis uteri, quo deciduum (*Z Z*) anteza obtexerat, facies interna. In hac superficie vene quo per deciduum ramos suos diffuderunt, abruptae conspicuntur.
e e Ejus partis uteri, quo placentam (*b b*) obtexerat facies interna. Haec quidem superficies cera extra vasa effusa specimen patefuit. Quae autem aspectu primo, extra vasa effusio videbatur, recta, venarum complanatarum, cum anastomosibus frequentioribus, ex utero ad placentam peroblique transeuntium, cera adimplatarum, species fuit. Cujus quod sequitur sit specimen.
f f In substantia uteri vena.
g g Ejusdem vena in placentam continuatio.
h h Uteri facies interna hic cavata, et vena dimidium, quo cursu peroblique transibat, formans.

P L A T E V.

THE first view of the opened womb. Its substance is cut through on the right side and turned to the left; by which means part of the decidua and of the placenta, as well as the inside of that portion of the womb, are exposed in the injected state. All the parts represented in outlines are seen finished in the next plate. The explanation is given here, that the other might not be obscured with engraved letters.

- A A* The upper part of the thighs.
B B Part of the lower corners of the containing parts of the abdomen turned down over the ossa ilium. They are covered by the vascular peritoneum; which (in the left side especially) was partly cut away, to shew the course of the epigastric vessels.
C C The ossa pubis cut through above the foramina ischium.
D D The ascending processes of the ischia cut through. These bones of the pelvis became loose, from the loss of the symphysis at the pubes; and on the left side they were drawn outwards, to shew more fully the descent of the womb into the cavity of the pelvis.
E E The inguinal arteries.
F F The inguinal veins.
G G The epigastric arteries.
H H The epigastric veins.
I I The obturator artery of the left side, a branch of the epigastric.
K K K The round ligaments descending from the abdomen upon the outside of the epigastric vessels.
L L The section of the integuments and muscles in each groin, which was made in removing the anterior containing parts of the pelvis.
M M The perineum.
N N The lower extremities of the labia pudendi.
O O The remains of the hymen.
P P The vagina.
Q Q The crura clitoridis.
R R The urethra slit on its upper part, through its whole length, and spread out.
SSSS The bladder spread round the forepart of the lower end of the womb.
T T The cavity of the bladder exposed by an incision through its forepart from the urethra upwards.
U U The small or lower extremity of the womb, lodged in the cavity of the pelvis. There was a manifest constriction upon that part of it which was surrounded by the brim of the pelvis; above which the womb swelled out immediately, being there embraced by more yielding parts.
VVV The outside of the body and fundus of the womb.
X X The cluster of spermatic vessels of the left side.
Y Y The substance of the womb cut quite through, from the fundus downwards on the right side. In this section the cut vessels bear an irregular figure, which was occasioned by the crumbling of the wax with which they were filled. The veins are of a light colour, and large; the arteries dark, and small in proportion. At the upper part of the womb, both of them are small in comparison with what they are lower down, where they were passing from the sides of the womb to the placenta: and the thickness of the womb bears some proportion to the size of its vessels in these different places.
Z Z The external membrane, or decidua, full of small branching veins, which passed into it from the internal surface of the womb.
a a At this place the vascular and opake decidua was carefully removed: there was no appearance of injected veins upon the chorion, or in the amnion; both of which were so transparent, that the child's skin could be seen distinctly through them.
b b A portion of the external surface of the placenta, interspersed partly with extravasated wax, from the venal system of the womb, and partly with injected veins, passing into it from the interior surface of the womb.
c c An artery filled with red wax, which was broken off from the womb. It made one circumvolution, and plunged into the placenta.
d d The inner surface of that part of the womb, which was in contact with the decidua (*Z Z*). Upon this surface the veins are seen broken off, which dispersed their branches through the decidua.
e e The inside of that part of the womb which covered the placenta (*b b*). Upon this surface there was indeed some extravasated wax: but the greatest part of what seemed, at first sight, to be extravasation, proved to be injected veins, of a flattened figure with numerous anastomoses, passing from the womb to the placenta in a very slanting direction. The following instance may serve as a specimen.
f f A vein in the substance of the uterus.
g g The same vein continued into the placenta.
h h The inner surface of the womb hollowed, and forming half of the vein, which passed here in a very slanting direction.

T A B U L A VI.

Fœtus in utero prout a natura positus.

Omnis hujus figuræ partes, utero et iis quæ in eo continentur exceptis, eadem fere sunt cum iis, quæ, in tabula precedentia lineis exterioribus, literisque jamjam explanatae sunt. In hoc tantum discrepant, nimurum hic vesice urinaria pars superior omnino sublata est, ad fœtus caput in parte uteri inferiori exhibendum. Uteri et secundarum pars anterior, una cum placenta, omnino sublata est. Funis umbilicalis est dissensus, deligatus, et in marginem uteri resecti sinistrorum reflexus. Ad fundum uteri membranæ investientes etiam in marginem uteri quo melius possint conspici inversæ sunt. Caput fœtus in uteri parte inferiori, in pelvis scilicet cavo, ponitur; corpus ejus præcipue in latere dextro jacet. Positio obliqua, vel ad lineam diagonalem appropinquans; unde partes posteriores antrorum sunt et ad matris latus dextrum; partes anteriores retrorsum et ad latus sinistrum versæ. Pes dexter inter femur sinistrum et tibiam occurrit. Corpus, uti infantibus denuo utero exclusis plerumque fit, muco albo pinguinculo fædatum: scilicet hic mucus, ad dorsi partem superiorem, ex rugis in corpore fœtus, motuque suo lineis intersecta, oculis subjicitur. Repræsentatur quæque pars eodem modo quo primum occurrebat, ne quidem vel digitii articulo turbato, partem aliquam magis illustrandi, aut tabulam elegantiorem reddendi gratia.

P L A T E VI.

This represents the child in the womb, in its natural situation.

All the parts of this figure, except the womb, and its contents, are nearly the same as those represented by outlines in the fifth plate, where they are lettered and explained. The only difference is, that here all the upper part of the bladder is cut away, in order to shew the situation of the child's head in the lower part of the womb. All the forepart, both of the womb and of the secundines (which included the placenta), is removed. The navel-string is cut, tied, and turned to the left side, over the edge of the womb. At the fundus the investing membranes are likewise turned over the edge of the womb, that they might be more apparent. The head of the child is lodged in the lower part of the womb, or in the cavity of the pelvis; and its body lies principally in the right side. Its position is diagonal or oblique; so that its posterior parts are turned forwards, and to the right side of the mother, and its foreparts are directed backwards, and to the left side. Its right foot appears between its left thigh and leg. Its body was covered with a white, greasy mucus, which is commonly seen on children at their birth. This is represented at the upper part of its back, where it was intersected with lines, from the wrinkles and motion of the child's body. Every part is represented just as it was found; not so much as one joint of a finger having been moved to shew any part more distinctly, or to give a more picturesque effect.




J.A. Rymsdyk delin'

TAB. VI. *Fetus in utero, prout a natura positus, rescissis omnino parte uteri anteriori,
ac Placentâ, cù adharente.*

R. Wenzel sculp.







F. S. Rennaldi delineavit
TAB. VII. Sublato Foetu parus uteri conspectus intimus et posterior membranis suis adhuc vestitus.
Partes levias adumbratae praeudenti tabula se ad plenum dedere.

London: Published July 1745 by E. CHI and Son. M. Thomas Street. Fetter Lane.

T A B U L A VII.

SUBLATO foetu, paries uteri conspicitur intimus et posticus, membranis suis adhuc vestitus. Partes lineis adumbrate in praecedenti tabula se ad plenum dedere.

A Partis uteri projectio versus interna. Hæc nata est a spina, vasis magnis, et præcipue ex aorta, quæ hic loci in arterias duas iliacas se dividit. Pars projecta, cadaveris situ supinato adaucta, superiorem et posticam uteri partem in cava duo lateralia retrosum dividit. Cavum dextrum, propter obliquitatem hujus uteri, sinistro majus fuit, et maximam fœtus partem continuit.

B B Partes duæ eodem modo projectæ, scilicet ob projectionem musculorum psoe, vasorumque iliacorum. Obliqua hæc juga, uteri cava superiora lateralia ab inferiore azygo, in pelve positæ, secernunt; in parte hujus ima posteriore oculis subjicitur.

C Os tincæ, paulo ad latus dextrum versum. Uterus in statu laxo, rugoso, uti fœtu vacuus apparebat, repræsentatur. Paries totus internus membranis suis vestitus. Per eas venæ cera impletæ, tales quales in hac figura, distinete se videndas exhibuerunt.

P L A T E VII.

A FORE-view of the cavity of the womb, as it appears when the child was taken out, and the investing membranes left adhering. The parts which are expressed by outlines are nearly the same as in Plate V. and VI. the first of which may be consulted for the explanation.

A An internal projection of the womb, occasioned by the spine and great vessels, especially the aorta, which divides at this place into the two iliac arteries. This projection, which is increased by the supine position of the subject, divides the upper part of the womb backwards into two lateral cavities. The right cavity, from the obliquity of the womb in this case, was larger than the left, and contained the greatest part of the child.

B B Two similar projections occasioned, in the same manner, by the projection of the psoæ muscles, and of the iliac vessels. These oblique ridges divide the upper lateral cavities of the womb from the azygos lower cavity which is situated in the pelvis; at the lower posterior part of which is seen

C The os tincæ, in this case a little on the right side. The womb is represented in the loose rugous state, as it appeared when empty; and its whole internal surface is covered by the membranes. Through these the injected veins were distinctly seen, as they appear in this figure.

T A B U L A . V I I I .

PARTES proxime post gravidum uterum sitae; utero ipso sede sua immoto, et in pudenda devoluto. Ossa pubis, &c. prius furent abscissa, ut in tabula quarta.

Iuniora, partesque ceteras lineis adumbratas, in tabula quinta juxtam explanata, cernere licet.

AAA Uterus.

BBBB Intestina tenuia que fundo uteri arte circumjacebant.

CCCC Intestinorum tenuium (cum mesenterio *DDD*) gyri, qui, post uterum, in latere sinistro, spinam inter et os ilium, supraque partem obliquam prominentem, e musculo scilicet psoa et vasis iliaca formatam, jacabant.

E Intestini ilei pars inferior (cujus mesenterium *F*), uterum inter et furcam aortae decurrens ad

G Terminationem suam in caecu.

HHH Caecum in cellas corrugatum propter ligamenta carnosa per longitudinem ejus tendentia.

I Cæci ligamentum anterius.

K Appendix cæci vermiformis.

L Cæcum in colon continuatum.

M Colon cum omento, loco in quo sursum et sinistrorum sub hepar ascendit, lineis adumbratum.

N Sinistri lateris colon, post intestina tenuia in faciem internam ossis ilium descendens.

OO Colon vasa iliaca transiens, et in pelvis cavum decurrens rectum continuo appellatum. Id suspensus cernitur de parte extrema mesocoli, supra vasa iliaca sinistra expansi. Hoc in loco intestinum coactatum fuit, et adipem involutum.

PP In cavo sacri, intestini recti priores gyri.

Q Trunci vene cavae extremitas, que hic in iliaca se dividit.

R Vertebrae lumborum infimæ facies anterior.

S Substantia, ligamenti natura prædicta, vertebraem inter proxime dictam et sacri partem summam.

T Ossis sacri summi facies anterior.

U Vena sacra.

Partes que literis *Q*, *R*, *S*, *T*, *U*, notantur, per peritoneum illis incumbens conspicuntur.

V Ureter dexter vasa iliaca transiens, et in pelvis latus decurrens.

W Arteria dextræ iliaca truncus.

X Venea dextræ iliaca truncus.

Y Psoas dexter.

Z Arteria iliaca externa.

a Vena iliaca externa.
Ut partes, qua literis *V*, *W*, *Y*, *Z*, et *a* notantur, in conspectum venirent, peritoneum illas obtegens fuerat percissum, membranæ cellulosa que circumjacebat cultro anatomico sublata.

b Vena iliaca interna.

c Ureter sinister per membranam cellulosa post peritoneum, et ante arteriam iliacan, ad pelvem transiens. Quo hic in conspectum prodiret, peritoneum et mesocoli pars inferior fuerant perissata, paululum etiam membranæ cellulose sublatum.

dd Vasa spermatica dextri lateris post cæcum, peritoneo tecta, descendenta.

ee Eadem vasa ad terminum superiore ligamenti lati, in uterum, tubam, et ovarium se diffundentia.

ff Tuba dextra pars.

g Ovarii dextri pars.

h Sinistri lateris vasa spermatica.

i Tuba sinistra.

k Ovarium sinistrum.

P L A T E . V I I I .

A view of the parts which lay immediately behind the womb, and which were in contact with it. They were brought to view by turning down the empty womb over the pudendum; the ossa pubis, &c. having been removed, as in the preceding figures.

The thighs and other parts expressed by outlines, are nearly in the same state as in Plate V. where the explanation is given.

AAA The womb.

BBBB The small intestines which lay around, and in contact with the upper part of the womb.

CCCC Those turns of the small intestines (and *DDD* the mesentery) which lay behind the womb in the left side, between the spine and os ilium, and above the oblique ridge formed by the psoas and iliac vessels.

E The lower part of the intestinum ileon, and (*F*) its mesentery. It passed between the womb and the bifurcation of the aorta to

G Its termination in the cæcum, where it was covered externally with an appendicula epiploica.

HHH The cæcum, corrugated into cells by its longitudinal fleshy bands, or ligaments.

I The anterior band or ligament of the cæcum.

K The appendix cæci vermiformis.

L The continuation of the cæcum into the colon.

M The colon and omentum represented in outlines, where this intestine mounts upwards, and to the left, under the liver.

N The colon in the left side, coming down, from behind the small intestines, upon the inside of the os ilium.

OO The colon passing over the iliac vessels into the cavity of the pelvis, where its continuation is called rectum. It is here suspended by the extremity of the mesocolon, which is expanded over the left iliac vessels. At this part the gut was contracted, and involved in fat.

PP The first turns of the rectum in the hollow of the sacrum.

Q The extremity of the trunk of the vena cava, which divides at this place into the iliac veins.

R The forepart of the lowest vertebra of the loins.

S The ligamentous substance between the last mentioned vertebra and the upper part of the sacrum.

T The forepart of the uppermost bone of the sacrum.

U The vena sacra.

The parts which are marked *Q*, *R*, *S*, *T*, *U*, are seen through the peritoneum that covers them.

V The right ureter, passing over the iliac vessels into the side of the pelvis.

W The common trunk of the right iliac artery.

X The common trunk of the right iliac vein.

Y The right psoas muscle.

Z The external iliac artery.

a The external iliac vein.

The parts marked *V*, *W*, *X*, *Z*, and *a*, were brought to view by slitting the peritoneum which covered them, and dissecting away the cellular membrane at that place.

b The internal iliac vein.

c The left ureter passing in the cellular membrane behind the peritoneum, and before the iliac artery, to the pelvis. This was brought to view, by making a long slit in the peritoneum, and lower part of the mesocolon, and taking away a small quantity of the cellular membrane.

dd The spermatic vessels of the right side, coming down from behind the cæcum and covered by the peritoneum.

ee The same vessels at the upper end of the broad ligament, diffusing themselves into the womb, tube, and ovary.

ff Part of the right tube.

g Part of the right ovary.

h The spermatic vessels of the left side.

i The left tube.

k The left ovary.



TAB. VIII. *Partes immediatae post impregnationem uterum sive utero ipso sede sua amoto, et in pudenda devoluta. Opia pubis, &c.
prius fuerant absissa. Confer TAB. IV.*

London Printed July 1774 by R. and V. Dodsley's Sons in New-Bond-Street.







J. J. Keenagh delin.

TAB. IX. *Sinuosa vertice per rectum spinam ducere, hoc figura representat pedem quaque ventre, sive seu naturaliter, cum aliisque effervore ducto et leviori parte.
Ostia ad fundum rectum, ad diaphragma procoelium latibus, manutinet medium, aliunde suppeditat. Tabula 22.*

London Printed by E. and T. Thos. for Thomas Bowes.

J. J. Keenagh delin.

T A B U L A I X.

PELVIS a latere ad plenum conspecta, una cum contentis ejus, partibusque configuis; per medium scilicet divisa, ut ossium flexure, et situs partium internarum oculo faciliter exhiberi possent. Imperfecta est hæc figura ad pubem, extreosum pariter aique intromissum. Partes que hic deficient, cultro separate sunt, dum cadaver ad aliquas tabularum precedentium pingendas parabatur. Defectum autem abunde supplebit Tab. XXII. in hunc foemina delineata.

- A* Femur dextrum a parte interiori.
- BBB* Lumborum vertebrae quatuor inferiores. Has inter cernit substantia ligamentosa, extans, inaequales, circa medium praesertim ubi molliissima est, et facile tumescit, si quando secta fuerit, et in aqua macerata.
- CCC* Vertebrarum earundem processus spinæ.
- D* Vertebrarum lumborum canalis, in quo reliqua quadam cauda equina, obsecure licet, conspicuntur.
- E* Musculorum lumbos extensissimam partes inferiores.
- F* Lumborum membrana adiposa et cutis.
- GGG* Sacri ossa tria superiora.
- HH* Sacri ossa duo inferiora, primumque coccygis, firmiter coalita.
- I* Coccygis os secundum, quod, ope medie cartilagine, ad primum facile movetur.
- K* Coccygis ossa duo inferiora in unum coalita, parique ad secundum mobilitate praedita.
- L* Sacri spina.
- M* Sacri canalis.
- N* Musculorum abdominis et integumentorum anguli inferioris pars, in ossis ilium spinam reflexa.
- O* Vena inguinalis magna epigastricam emittens, quam etiam arteria ejusdem nominis constitutus.
- P* Femoris musculi, ex osse pubis orti, quale in osse eo sejungendo facunt percussi.
- Q* Os pubis supera foramen magnum dissiccum.
- R* Idem os, ubi ramo paro ischiis committitur, dissiccum.
In loco quo litera *P* occurrit, lineis punctis adumbrata representatur pars ejus ossis pubis, que abscissa fuerat, figura. Nota * iisdem lineis adumbratum symphysis ostendit.
- S* Pars carnosæ anus inter et os coccygis.
- T* Anus omnino apertus.
- U* Recti pars inferior lac sectione exposita. In hac figura videoas, intestinum idem ab ano sursum progrederis oculo dum persequeris, ad se retrorsum deflectisse, nempe ut ad faciem internam ossis coccygis veniret.
- V* Hic rectum ad latus sinistrum se deflexit, solamque membranam suam cellulam in hac sectione cernere licet.
- WW* Rectum sectione apertum, ubi ante coccygis os primum, sacrifice ossa tria inferiora, decurrebat.
- X* Hic rectum omnino in latere dextro jacebat, nec id aperuit secessu.
- Y* Rectum dissiccum, ubi de latere dextro ad os secundum sacri decurrebat, et in flexuram coli signoideam cursum suum tenebat.
- Z* Vena cava.
- a* Arteria iliaca dextra.
- b* In hoc loco vena iliaca sinistra dissicca; cera autem, quæ injecta fuerat, utpote fricta, jam delapsa, vessa haud satis clare se in conspectum dedit.
- c* Perinei sectio.
- d* Labii dextri, ad partem inferiorem, facies interna.
- e* Nymphæ dextræ, ad partem inferiorem, facies interna.
- f* Coeratatio, hymenische reliquia, ubi incipit vagina.
- g* Meatus urinarius extremitas.
- hh* Vagina carum.
- i* In hoc loco vagina et rectum sunt coalita. Rectum vagina densius.
- k* Vagina pars anterior, cum urethra et vesica urinaria conjuncta.
- ll* Os uteri in vagina, ad os coccygis vergens.
- Quavis in hoc cadavere fundus uteri dextrosum inclinatus fuerat, os uteri tamen adeo ad dextrum se tenebat, ut cultrum anatomicum has partes secantem fugeat.
- m* Ad os tincæ, uteri et vaginae substantia coalita.
- n* Uteri in duas partes reuelas secti margo posterior.
- o* Uteri secti pars anterior. Sectio hæc nequaquam per medium uteri se habebat, sed ad latus dextrum haud paululum; etenim ut figure precedentes commode posse delineari, cadaver hoc modo necesse habui disponere.
- p* Os uteri intus.
- q* Uteri paries internum membranis suis vestitus.
- Fundus uteri in hac figura non representatur; tunc quoniam id a proposita ratione alienum duxi, tunc quoniam uterus adeo flaccidus et tener hunc temporis erat, ut in situ suo naturali, ad arbitrium et usum pictoris, minime potuerit retineri.
- Membranæ, tempore quo tabula hæc delineabatur, ipsæ se ab utero free perforauerant. In parte superiori, decidua, ubi se ab utero receperat, venulis abundavit; justa os uteri, vix ullus horum vasorum vestigium apparuit.
- r* Meatus urinæ.
- Vesica urinaria pars inferior, uteri cervicem inter et vaginae partem superiorem sita. Omnis vesicae portio post symphysis pubis sita jamdudum abscissa fuerat. Melius autem nihil visum est, figuram vel curtam dare, quam aliquid ingenuo excogitatum pingere.

P L A T E I X.

A FULL side-view of the pelvis with its contents, and adjacent parts, cut down through the middle, to shew the turns of the bones, and the situation of the internal parts. This figure is imperfect at the pubes, both externally and internally; the defective parts having been cut away, in preparing the subject for some of the preceding figures. But what is wanting in these, may be seen in Plate XXII. which was intended as a supplement to this.

- A* The inside of the right thigh.
- BBBB* The bodies of the four lowermost lumbar vertebrae. Between these the ligamentous substances are represented prominent, and unequal, especially near the centre, where they are softest and most disposed to swell when soaked in water, and cut through.
- CCCC* The spinal processes of the same vertebrae.
- D* The canal of the lumbar vertebra, in which some remains of the cauda equina are indistinctly seen.
- EE* The lower part of the extending muscles of the loins.
- F* The adipose membrane and skin of the loins.
- GGG* The three superior component bones of the os sacrum.
- HH* The two lowermost bones of the sacrum, and the first of the coccyx, firmly ankylosed.
- I* The second bone of the coccyx, moveable on the first by means of an intermediate cartilage.
- K* The two last bones of the coccyx grown into one, and moveable on the second bone by the intervention of a cartilage.
- L* The spine of the sacrum.
- M* The spinal canal in the sacrum.
- N* Part of the lower flap of the abdominal muscles and integuments, turned over the spine of the os ilium.
- O* The great inguinal vein, sending off the epigastric, which is accompanied with the artery of the same name.
- P* The muscles of the thigh which arose from the os pubis, and which were cut through when that bone was removed.
- Q* The os pubis cut through above the foramen magnum.
- R* The same bone cut through at its conjunction with the small branch of the ischium.
- Where the letter *P* stands, is represented, in dotted outlines, the figure of that part of the os pubis which was cut off, and the mark * is upon the symphysis in the same outlines.
- S* The fleshy part between the os coccygis and the anus.
- T* The anus considerably opened.
- U* The lower part of the rectum laid open by the section. In tracing the gut from the anus upwards, we see from this figure that it takes a bend backwards, to get at the inside of the os coccygis.
- V* Here the rectum made a turn to the left side, and its surrounding cellular membrane only was seen in the section.
- WW* The rectum laid open by the section, where it runs before the first bone of the coccyx, and the three lowermost pieces of the sacrum.
- X* Here the rectum lay entirely in the right side, and was not opened by the section.
- Y* The rectum cut through, by the section, where it passed from the right side, across the second bone of the sacrum, and was continued into the sigmoid flexure of the colon.
- Z* The vena cava.
- a* The right iliaca artery.
- b* Here the left iliaca vein was cut through: but the brittle wax with which it was filled had fallen out, and the vein was seen indistinctly.
- c* The section of the perineum.
- d* The inside of the lower part of the right labium.
- e* The inside of the lower part of the right nymphæ.
- f* The stricture at the beginning of the vagina, or remains of the hymen.
- g* The extremity of the meatus urinarius.
- hh* The cavity of the vagina.
- i* The compound substance of the vagina and rectum, the latter of which is considerably the thickest.
- k* The forepart of the vagina united with the urethra and bladder.
- ll* The mouth of the womb in the vagina, directed towards the os coccygis. Though the bottom of the womb, in this case, was directed towards the right side, its mouth lay so much on the right side, that it was not touched in making the middle section of all the parts.
- mm* The substance of the womb and vagina blended, at the os tincæ.
- nn* The edge of the bisected womb backwards.
- oo* The edge of the womb forwards. This section was not in the middle, but considerably on the right side. It had been made to prepare the subject for some of the preceding figures.
- p* The mouth of the womb internally.
- q* The inside of the womb lined with the membranes.
- The bottom of the womb is not represented in this figure; both because it was not very material here, and because it was become so flaccid and tender, that it could not be well kept out, in its natural situation, before the painter.
- The membranes were almost entirely separated from the womb, of themselves, when this figure was made. In the upper part the decidua was full of small veins, even where it parted off from the womb; but near the mouth of the womb, hardly any such vessels appeared.
- r* The meatus urinæ.
- The lower part of the bladder, placed between the neck of the womb and the upper part of the vagina. The upper part of the bladder, which was situated behind the symphysis of the pubes, had been cut away before this section was made; and we chose to leave the figure imperfect, rather than venture to delineate from fancy.

T A B U L A X.

F I G . I.

UTERI pars anterior et extima, prout se videndam praebuit, omnino siccata, exhibens faciem vasorum uterinorum, quem illa pre se ferunt, eo loci ubi placenta utero adhaeret.

Vene cera flava impletæ, ab arteriis, magnitudine, colore lucidiori, anastomosibus pluribus, facile dignoscuntur. Arteriae enim, minores, pauciores, flexuose, coloris magis luridi hic loci sunt. Arteriae omnes convolute; pauciores autem videbantur, quia multi earum ramuli, cera flava, venis prius injecta, impletæ fuerint. Hujus exemplum unus est ad literam *A*.

Venas profundius sitas, per uterum siccatum cernere non licuit; indistinctæ igitur, nulloque ordine, ita uti visæ sunt delineantur.

F I G . II.

Facies interna placenta, cuius vasa per funiculum umbilicalem cera sunt repleta.

A Cannula ænea in arteriarum alteram umbilicalium inserta, per quam ultrae arteria cera viridi impletatur.

B In vena cannula, per quam cera cœrulea illi injecta fuit.

CC Amnion, in parte quadam placenta, laxum relictum.

DD Placenta, qua amnion sublatum est, superficies intima chorio producto coeperta. Hoc in loco rami primi vasorum umbilicalium conspicuntur. Arteriae colore lucidiori facile possunt distinguiri.

Hie arteriae quidem lucidiores, ut in figura, apparebant; tunicis scilicet earum pellucidis colorem cerae injectæ viridem lucidorem transmittentibus: In funiculo autem umbilicali aliter se res habet quippe vena, in medio posita, undique glutinosæ funiculi substantia fuit intacta; unde cera cœrulea ad livorem propius accedens visa est.

E Decidua pars in margine placenta relictæ, inque faciem ejus internam reflexa; nempe ut superficies ejus exterior, utero que adhaerent, vasis parvulus (cera flava per venas uteri impletis) referat, possit exhiberi.

Placenta cera flava ubique turgida, que substantiam ejus celulosam adeo implevit, ut in facie interna ubique per chorion transluceret. Cera ubique quasi in grana abiit, quod inde cava naturalia impleverat, indicium certum exhibens: si enim placentam, vasis effusa impletisset, in massas majores, irregulares formatam, se ad prospectum disset. Cera in hanc spongiosam, sive celulosam partem, injecta, placenta crassitudo naturali duplo major evaserat.

F I G . III.

Mulieris, die secundo post partum extinctæ, uteri superficie internæ pars.

Decem his tabulis jam delineatis, eas statim in lucem edere proposui; et simul hancce tertiam figuram, quasi supplementum, addere, utpote solum quod tunc temporis potui. Discrimen exhibet inter superficiem uteri (*A, B, C*) cui membranæ adhaerent, et superficiem (*D, E, F*) cui placenta se immiserat; illam quidem levem et unius formæ; haec autem orificiis magnis venarum referat. Ubi primum haec orificia sub examen venerant, omnia grunis coagulati sanguinis referata comperti, in cavum uteri nonnihil extantibus. Utero vero aliquandiu in aqua macerato, coagula haec magis albida evaserunt; nonnulla exciderunt; et aliqua etiam extracta sunt, ut venarum orificia clarius monstrari possent. Ita se res habuit, cum figura haec delineabatur.

P L A T E X.

F I G . I.

A View of the outside of the forepart of the womb, as it appeared when quite dry; exhibiting a specimen of the uterine vessels, at the part where the placenta adhered.

The veins were filled with yellow wax, and are readily distinguished by the largeness, light colour, and frequent anastomoses, from the arteries, which are here smaller, less numerous, serpentine, and of a darker colour. The arteries were all convoluted; but they did not appear to be so numerous, because many of the branches were filled with the yellow wax, which had been thrown into the veins. This is seen, in one instance, at the letter *A*.

The deeper-seated veins could not be distinctly seen through the dried substance of the womb, and are therefore represented with the same obscurity and confusion which appeared in the object itself.

F I G . II.

The inside of the placenta, which was injected by the umbilical vessels after it was taken out of the womb.

A A brass pipe tied into one of the umbilical arteries, by which the two arteries were filled with green wax.

B A pipe in the vein, by which it was filled with blue wax.

CC The amnion left loose upon one part of the placenta.

DD The internal surface of the placenta, where the amnion was removed, covered by a continuation of the chorion: upon this are seen the first branches of the umbilical vessels. The arteries are sufficiently distinguishable by their lighter colour.

They appeared so from being filled with a lighter coloured green wax, and from the transparency of their coats: but on the navel-string it was the contrary; because the vein, being the most central, was covered with a greater thickness of the gelatinous substance of the string, and therefore did not allow the darker-coloured blue wax to be so well seen.

E Part of the decidua left at the edge, and turned over the inside of the placenta, so as to shew its outer surface (which had adhered to the womb) full of small vessels, injected with yellow wax from the uterine veins.

The body of the placentæ was uniformly filled with the yellow wax. This had so completely filled all the cellular substance of the placenta, that upon the inside it appeared every where through the chorion. And the wax was every where manifestly granulated; so that it had plainly filled natural cavities: for if it had filled the substance of the placenta by common extravasation, it would have formed itself into irregular and large masses. From this injection of the spangy or cellular part, the placenta had acquired at least double its natural thickness.

F I G . III.

A portion of the internal surface of the womb, from a woman who died two days after delivery.

When these ten plates were finished, I intended to have published them immediately, and this third figure was added, as the only supplement which I could give at that time. It shews the difference between the surface of the womb (*A, B, C*) where the membranes had adhered, and that surface (*D, E, F*), to which the placenta had been fixed. The first was much more smooth and uniform; the latter was full of large venous orifices. At the first examination, all these orifices were found filled up with plugs of coagulated blood, which projected some way into the cavity of the womb. The part being steeped some time in water, they became whiter, and many of them fell out, or were pulled out, to shew the orifices of the veins more clearly. In this state the drawing was made.


Fig. III.



Fig. II.





Fig. I.



J. T. Rymer delin.^s T. C. Cooper sculp.
TAB. X. Fig. I. Ultori pars anterior et extimata, prout se probuit omnino siccata, exhibens faciem vasorum uterinorum, qualem prie se
ferunt eo loci, ubi Placenta uteri adhaeret. Fig. II. Facies interna Placentae, cuius vasa per funiculum umbilicalem ceri sunt repleta.
Fig. III. Ab aliâ muliere biduc post partum extintâ, exhibet partem superficiem intime uteri.

London: Published July 1st 1815 by E. CHURCH and Son, 57, Fleet Street, Breamore.







J. J. Rymer delin.
TAB. XI. ab aliis muliere, nono gravitatis mense hemorrhagiā uterina e medio sublatā. Conspicitur
uterus et osa pubis juxta naturam sita. Vasa injecta cœri turgida, quoque ad partem inferiorem et
cervicem uteri digressantur ope amplissima, arguant Placentam parti illi adhaerere.

London: Published by J. D. and T. G. Newell, 1805.

T A B U L A XI.

As alia muliere, nono graviditatis mense haemorrhagia uterina e medio sublata. Conspicitur uterus, et ossa pubis juxta naturam sita. Vasa, injecta cera, turgida; queque ad partem inferiorem et cervicem uteri dignoscuntur esse amplissima, arguant placentam parti illi adhaesisse.

- A** Fundi uteri pars media et summa.
- B** Uteri pars qua scilicet in cadavere fuit ex adverso umbilici.
- C** Symphysis ossium pubis.
- D** Membrana adiposa ad fundum vesicæ urinariæ.
- E E** Os pubis utrinque, ad locum fere quo ossi ilium committitur, percissum.
- F F** Utriusque ossis ischii ramus parvus, ad locum quo ossi pubis adjungitur, percissus.
- G G G** Contenta pelvis omnia, antrorum ossibus pubis obtecta.
- H** Infra hanc literam meatus urinarii orificium.
- I** Ab hac litera sursum ad meatum urinarium vaginae principium; infraque hanc literam aditus ad rectum.
- K** Strato intimo ani sphincteris circumtectus.
Labia pudendi, perinæum et anus hic non habent locum; in situ enim suo relicta erant in cadavere, quippe illud externe matutum et deforme reddere non licet.
- L L** Tube.
- M M** Fimbriae.
- N N** Ligamenta rotunda.
- O** Ovarii pars.
- P P** Vasorum spermaticorum utrinque ad uterum ascendentium fasciculus.
- Q Q** Vasa hypogastrica, in latere uteri, ad conjunctionem cum spermaticis ascendentia.
- R** Uteri venæ magnæ in loco cui placenta interne adhaerebat.

P L A T E XI.

FR OM a second subject, viz. a woman who died of a flooding in the ninth month of pregnancy. It represents a fore-view of the womb, with the ossa pubis in the natural situation. The situation of the large vessels, which were injected, shews that the placenta was attached forwards, and to the lower part of the womb.

- A** The middle and highest part of the bottom of the womb.
- B** That part of the womb which, as it lay in the dead body, was directly opposite to the navel.
- C** The symphysis of the ossa pubis.
- D** The adipose membrane upon the fundus of the bladder.
- E E** The os pubis of each side, cut through near its union with the os ilium.
- F F** The small branch of each os ischii, cut through near its conjunction with the os pubis.
- G G G** The whole contents of the pelvis covered forwards by the ossa pubis.
- H** Immediately below this letter is seen the orifice of the meatus urinarius.
- I** From this letter upwards to the meatus urinarius is seen the beginning of the vagina; and immediately below this letter *I* is seen the passage into the rectum, surrounded by
- K** The inner stratum of the sphincter ani.
The labia, perinæum, and anus were left in the body, that it might not be disfigured outwardly.
- L L** The tubes.
- M M** The fimbriae.
- N N** The round ligaments.
- O** Part of the ovary.
- P P** The group of spermatic vessels, on each side, approaching the womb.
- Q Q** The hypogastric vessels mounting upon the sides of the womb to join the spermatic.
- R** The great veins of the womb in that place to which, internally, the placenta was attached.

T A B U L A XII.

Az eadem, qui præcedens, muliere, exhibet apertum plane a parte postica uterum cum vagina, quo situs fœtus, parsque inferior placente, sub fœtus capite, indicarentur. Placenta scilicet orificio uteri interno acreverat, eoque sub finem graviditatis dilatato, separata est: unde haemorrhagia, et mors.

- AA* Tuba fallopiana.
B Ovarium sinistrum; cuius ad extremitatem inferiorem videre est
C Corpus luteum extans.
D Ovarium dextrum.
E Vasorum spermaticorum ad uterum ascendentium fasciculus.

- FFF* Substantia uteri sectio.
G Vaginae per incisionem aperta et expassæ facies interna.

HH Os uteri.

I Placentæ partis inferioris facies externa, in lobos partita, quæ quondam cervici et ori uteri interne adhaeserat; tempore autem parturiendi instante, partim illarum dilatatione sejuncta fuerat, unde necessario haemorrhagia.

KK Membranae de margine placentæ exentes, capitisque fœtus partem inferiorem amplectentes, percissæ. Spatium, quod inter cervicis uteri partem interiorem, et secundas, vacuum occurrit, in cadavere sanguine coagulato refertum compéri.

Situs partesque fœtus explanatione non egent.

P L A T E XII.

FROM the same subject, a view of the womb and vagina fully opened on the back part, to shew the situation of the child, and of the lower part of the placenta at the inside of the mouth of the womb, under the child's head, and detached from the womb; the occasion of the fatal hemorrhage.

- AA* The fallopian tubes.
B The left ovary, at the lower end of which is seen
C The projecting corpus luteum.
D The right ovary.
E The group of spermatic vessels approaching the sides of the womb.

- FFF* The section of the substance of the womb.
G The inside of the vagina, which is laid open by a longitudinal incision, and spread out.

HH The mouth of the womb.

I The external lobulated surface of the lower part of the placenta; which had originally stuck to the inside of the neck and mouth of the womb; but as parturition approached, the dilatation of these parts occasioned a separation, which was necessarily followed by an hemorrhage.

KK The membranes cut through where they were coming out from the border of the placenta, and enclosing the most depending part of the child's head. The vacant space which is seen between the inside of the neck of the womb, and the secundines, was full of clotted blood in the dead body.


The situation, and the several parts of the child, require no explanation.



J. J. Rengacharthy delin.
TAB. XII. ab eadem, qui precedens, muliere, exhibet apertum planum a parte posteriori uterum cum vagina,
quo situs Fœtus, parque inferior Placenta sub Fœtus capite indicarentur. Placenta scilicet orifice
uteri interno acreverat, quoque sub finem graviditatis dilatata, fatali modo divisione separata est.

London Printed July 17th 1805 by E. CHAPMAN and Son, 57, Fleet-street, W.C.







Engraved after
 TAB. XIII. A tertio cadavere, nono gravitatis mense. Aperto Utero (vagina et vesica urinaria integris) incastratus, scilicet, una cum anteriori Uteri paride, secundinis, et adhaerentibus, et in fundum Uteri reclinatis, situs Fetus (qui hic erectus est) conspicitur. Vasa Uteri eorum erant repleta.

T A B U L A XIII.

A TERTIO cadavere, nono graviditatis mense. Uteri conspectus ab anteriori, cum vagina et vesica urinaria. Pariete hujus anteriori, cum secundis ei adhaerentibus, inciso, et in fundum ejus reclinato, situs fœtus erecti conspicitur. Vasa uteri, cera prius injecta, turgida fuerunt.

- A Vesica urinaria in situ suo naturali, respectu habito ad uterum.
- B Vaginæ pars superior et extima, quæ sub symphysi ossium pubis jacet, quaeque ibi cum urethra connectitur.
- C Vaginæ cavum in conspectu positum, cum labia partesque aliae externe fuerant abscissæ, quo specimen hoc e cadavere eximiretur.
- DD Vasa spermatica, ad uteri latera ascendentia.
- EE Tubæ; quarum extremitates, seu fimbriæ, sub racemo vasorum spermaticorum, occultantur.
- F In uteri latere dextro, vena magna, ab anastomosi venarum, hypogastricæ scilicet et spermaticæ, ortum suum dicens.
- GG Ligamenta rotunda.
- HH Uteri pars anterior, cum portione placentæ quæ ei adhaerebat, incisa, et in fundum uteri reclinata, ut fœtus in conspectum prodiret.
- III Uteri substantiæ, et membranarum investientium, sectio.
- KK Eadem sectio per placentæ substantiam ducta, quæ uteri parti anteriori in latere dextro adhaerebat. Ex figura inversa latus dextrum sinistrum fit.
- L Funis umbilici, a fœtu ad placentam transiens.

P L A T E XIII.

FROM a third subject, in the ninth month of pregnancy. A fore-view of the womb (with the vagina and vesica urinaria) in which all the inclosing parts were cut through, and turned up, to shew the situation of the child, with its head upwards. The vessels of the womb had been previously injected.

- A The bladder, in its natural situation with respect to the womb.
- B The upper and external part of the vagina, which lies under the symphysis of the ossa pubis, and where the urethra is united with it.
- C The cavity of the vagina exposed, where the labia and other external parts had been cut off, in taking this part from the dead body.
- DD The spermatic vessels, passing up towards the sides of the womb.
- EE The tubes, of which the extremities, or fimbriae, are concealed behind the group of spermatic vessels.
- F The great vein on the right side of the womb, formed by the anastomosing hypogastric and spermatic veins.
- GG The round ligaments.
- HH The forepart of the womb, with that part of the placenta which adhered to it, cut up, and turned back over the fundus of the womb, to bring the child to view.
- III The section of the substance of the womb, and of the investing membranes.
- KK The same section carried through the substance of the placenta, which, in this case, was fixed to the forepart and right side (which is the left in this inverted figure) of the womb.
- L The navel-string passing from the child to the placenta.

T A B U L A / XIV.

A QUARTO cadavere, nono itidem graviditatis mense. Fasciculi fibrarum muscularium, in parte interna uteri, triplici figura exhibentur.

Hac parte uteri per aliquot dies in aqua macerata, decidua jam tenera abstergebatur.

F I G . I.

Uteri inversi, et in latus paululum reclinati, facies posterior interna.

A Uteri os relaxatum; supra quod, paulo obscurius licet, cævæ ruge, ramorum ad instar dispositæ, cernuntur.

BB Uteri pars angusta, margine pelvis ante circumscripta.

CC Seta in orificio utriusque tubæ; quam circum fasciculi fibrarum ad musculos pertinentium, in orbæ, eodem centro terminatos, se disponunt.

D Hic fasciculi, in orbem dispositi, uterum cingunt.

E Uteri fundi pars media, qua ordines hi duo fibrarum, tubas cingentium, inter se coèunt, sive se invicem contingunt.

F Interstitium triangulare, tres inter ordines fibrarum prædictarum, ubi mutuo inter se coèunt, diversumque ordinem et cursum tenent.

F I G . II.

Ejusdem uteri, forma minuta, conspectus a latere.

A Uteri os.

B Seta in orificio tubæ.

C Fibrarum, sine ordine dispositarum, stratum, faciem internam fasciculorum, qui in circulum fleetuntur, obtengentes.

F I G . III.

Ejusdem uteri partis anterioris facies interna.

A Os uteri; supra quod rugæ anteriores, in ramos abeunt, se in conspectum dant.

B Superficie cui placenta adhaeserat pars media, ubi fibrae fascias sine ullo ordine formabant; quas inter, venarum orificia, in placente partem spongiosam cursum tenentum, passim occurserunt.

Ad cervicem uteri, fibrae musculosæ, in fasciculos collectæ, nullæ conspicuntur.

P L A T E / XIV.

FR OM a fourth subject at nine months. This shews the disposition of the muscular fasciculi on the inside of the womb, in three different views.

The part was steeped in water some days, whereby the decidua was made tender, and then brushed off.

F I G . I.

The inner surface of the posterior part of the womb, turned a little to one side.

A The relaxed mouth of the womb, over which are faintly seen the branching rugæ of the neck of the womb.

BB The narrow part of the womb, which was embraced by the brim of the pelvis.

CC A bristle put into the orifice of each tube, round which the fasciculi of muscular fibres are ranged into concentric circles.

D The fasciculi going circularly round the body of the womb.

E The middle of the fundus of the womb, where the two orders of circular fibres, which surround the tubes, meet, or become reciprocally tangents.

F The triangular interstice between the three orders of circular fibres, where they are all blended, and take a variety of directions.

F I G . II.

A side-view of the same womb, in miniature.

A The mouth of the womb.

B A bristle in the orifice of the tube.

C An irregular stratum of fibres, upon the inside of the circular fasciculi.

F I G . III.

The inner surface of the anterior part of the same womb.

A The mouth of the womb: over which are seen the anterior branching rugæ.

B The centre of the surface to which the placenta adhered, where the fibres formed irregular, interwoven bands; in the interstices of which were the orifices of the veins which went into the spungy part of the placenta.

At the cervix no distinct muscular fibres appear on the internal surface of the womb.


Fig. I.



Fig. III.

*opposite side.*

TAB. XIV. *Al quarto Cadavere, nono idem graviditatis mense. Fasciculi fibrarum muscularium, in parte internâ uteri, triplici figura exhibentur.*







Ex opere ab aliis.
TAB. XV. d' quanto Cadavere, non diam gestationis mense, systema arteriosum Uteri quintuplici figura exprimum. Fig. I. Fasces Uteri inverso, quo praevertim loco Placenta adhaeserat. Fig. II. Arteria partem Uteri anteriorem perreplantata. Fig. III. Eadem per partem posterioram distributrix. Fig. IV. Eadem Fundum pernuntius. Fig. V. Corpus sicutum in Ovario per medium diviso, & Febris Tuba Fallopiana.

London: Published July 1780 by J. & W. and 1803 by Thomas & Newgate.

TABULA XV.

A QUINTO cadavere, nono etiam gestationis mense, sistema arteriosum uteri quintuplici figura expressum.

Mulier haec, post partum neutiquam difficultem, in animi deliquium inciderat, causa quidem adstante latente; horis autem duabus a partu nondum lapsa, illam mors abstulit. Arterie cera rubri coloris implete.

FIG. I.

Facies uteri inversi, quo praescitum loco placentas adhaeserant.

A Pars media superficie, cui placentas adhaeserant, in fundi parte postica, et paulo ad latus sinistrum sita.

B Arteriae convolutae quanplurimae, quarum extremitates, quo loco per deciduam ad placentam decurrent, ruptae sunt.

C Orificia venarum uteri (vulgo sinus appellata) que per deciduam ad placentam decurrent; haec multo plura quam quae in figura representantur; minora praesertim innumerata. Ex venarum illarum cursu, uteri scilicet superficie parallelo, aere non prius inflato, orificia rix possunt conspicui. Earum igitur statu distensum maxima pars delineatur, nimirum ut umbra insertis, magis conspicuer evadent, et cursus etiam earum obliquus, nec non orificia ramorum inter se coextintum, in conspectum venient.

D Uteri superficie pars cui membranae adhaeserant. Vasa hujus partis, omissa quam minima. Pars etiam ipsa laevior, quam ea cui placentas se immiscerant. Facies deciduae externe perfectam quandam similitudinem praeseferunt.

E Uteri internae lamellae pars, cultro anatomico sublata, et reclinata, ea mente, ut conspicerentur.

F Uteri fibrarum fasciculi, quorum maxima pars circulos conficiunt.

G Oris uteri dilatati margo, vesiculis glutinosis distinctus.

H Cervix uteri, ad partem posticam, rugae pennatae, quae vulgo valvulae appellantur.

I Vaginae inversae pars.

FIG. II.

Arteriae partem uteri ejusdem anterioem perpenitantes. Uteri facies externa jam siccata fuit, et inde pellucida adeo facta, ut arteriae in conspectum distincte venirent, pariter ac si in superficie extima dispersa fuissent. Uterus hic duodecim fuit pollices longus: unde facile patet, quantum mox contrahatur in hac figura: quantum etiam in figura tercia et quarta, que ad eandem normam delineantur. Pars haec dum crescet, in longitudinem quam latitudinem magis distensum fuit: quapropter uterus, in hac, et in figura proxime sequenti, longior est quam ex natura.

A Peritonei resecti margo, eo loci, quo ab utero ad vesicam urinariam reflectitur.

B Tuba.

C Ovarium sinistrum.

D Ovarium dextrum. Vid. Fig. V.

E Ligamenta rotunda: quorum in sinistro est arteria magna flexuosa, ab ramo spermatico, per anastomosis conjunctio cum hypogastrica, transmissa: in dextra aliquot sunt minores.

F Truncus arterie sinistre hypogastricae, ad cervicem uteri latus ascendens.

G Truncus arterie sinistre spermatica, in ligamenti lati duplicationem transeuntis.

Arteriarum distributionem in utero explanare haud opus est. Cursus enim earum flexuosi, anastomosique plurimae, per se satis patent.

FIG. III.

Eundem exhibet uterum a parte postica visum.

B B. **C. D. E E.** Vid. Fig. II.

F Arteria hypogastrica dextra.

G Arteria spermatica ad ligamentum lata excurrentes.

H In ea parte uteri cui placentas adhaeserant arteriae flexuose confertae.

Si arterias in hac figura, eas in figura precedenti conferamus, eas multo maiores, magisque convolutas esse, in latere eo uteri, cui placentas adhaeserant, certe habebit. In utero gravido, cursus sanguinis maxime in placentam diriguntur.

FIG. IV.

Fundum exhibet uteri, parte postica sursum versus.

B B. **T**uba.

D Ovarium sinistrum.

C Ovarium dextrum.

Vid. Fig. III.

FIG. V.

Ovarium dextrum et tuba ejusdem uteri.

A Tuba.

B Tubae ligamentum, mesenterio quodammodo respondens.

C Tubae orificium extreum, fimbriae folia referentibus circumdatum.

D Fimbriae a latere utroque orificio tubae, per ligamentum, ad ovarii extremitatem externam, continuatae.

E E. Ovario jam aperio, arterie flexuose, per substantiam eius sparsae, conspicuntur.

F Hac in parte corpus luteum, per medium divisum, in ovarii substantia cernitur.

Nulla conspicuntur vasa in centro, quod colore est albido. Circa centrum illud undique, substantia eius rasculis abundat.

PLATE XV.

FIVE figures, from a fifth subject at the full time, to explain the arterial system of the pregnant womb.

This woman, immediately after a natural labour, grew faint, as was said, without an apparent cause; and died within the space of two hours. The arteries were injected with red wax.

FIG. I.

The womb, inverted.
A The centre of the surface to which the placenta had been fixed; which was at the back-part of the fundus, and a little on the left side.

B Innumerable convoluted arteries, with broken extremities, where they had passed through the decidua into the placenta.

C The orifices of those sericeous veins, commonly called sinuses, which had passed through the decidua to the placenta. These orifices were much more numerous than they are represented in this figure. The smaller especially were innumerable. The passage, or direction of these veins, was almost parallel to the surface of the womb; so that their orifices were scarce perceptible, without being inflated. Most of them are represented in that state; both that they may be more distinct, by having a little shade, and that their oblique direction, and the orifices of their communications, or anastomosing branches, may be seen.

D Part of the surface of the womb where the membranes had adhered. This was furnished with very small vessels only; and was much smoother than that part to which the placenta had adhered. In every respect it was like the outer surface of the decidua.

E Part of the inner lamella of the womb, raised by dissection, and turned to one side, to shew

F The fasciculated fibres of the womb, most of which, at this place, were nearly circular.

G The border, formed by the dilated mouth of the womb, and studded with gelatinous vesicles.

H The branching rugae, commonly called valves, at the posterior part of the neck of the womb.

I Part of the inverted vagina.

FIG. II.

A fore-view of the outside of the same womb, after it was dried, and become so transparent, that the arteries could be seen distinctly, as if they had been dispersed on its outer surface. The object was twelve inches long; from which it will be understood, how much the scale is reduced, in this, as well as in the third and fourth figures, which are delineated to the same proportion. In drying, the preparation was unavoidably more stretched in its length, than in its width; whence in this, and in the following figure, the womb is more oblong, than it is naturally.

A The edge of the cut peritoneum, where it was reflected from the womb to the bladder.

B B. The tubes.

C The left ovary.

D The right ovary. See Fig. V.

E E. The round ligaments. In the left there is one large serpentine artery, sent down from an anastomosing branch of the spermatic and of the hypogastric: in the right there are several small arteries.

F The trunk of the left hypogastric artery, mounting upon the side of the neck of the womb.

G The trunk of the left spermatic artery, passing into the duplicature of the broad ligament.

The distribution of the arteries upon the womb requires no explanation. Their serpentine course, and frequent anastomoses are apparent.

FIG. III.

A back-view of the same subject.
B B. **C. D. E E.** See Fig. II.

F The right hypogastric artery.

G The spermatic arteries passing into the broad ligaments.

H H. The crowded serpentine arteries in that part of the womb where the placenta adhered.

By comparing the arteries in this figure, with those in the preceding, we see that they are much larger, and more convoluted, in that side of the womb to which the placenta is fixed; the great current of blood being towards the placenta.

FIG. IV.

A full view of the fundus of the same womb, the back part being upwards.

B B. The tubes.

D The left ovary.

C The right ovary.

H H. See Fig. III.

FIG. V.

The right ovary and tube of the same womb.

A The tube.

B The ligament of the tube, analogous to the mesentery.

C The external orifice of the tube, surrounded by foliaceous fimbriae.

D The fimbriae continued from each side of the orifice of the tube, all along the ligament, between that and the outer end of the ovary.

E E. The ovary cut open, with serpentine arteries interspersed through its substance.

F F. Here, in the substance of the ovary, the corpus luteum is seen, split through the middle. No vessels appear at its centre, which is of a white complexion; but all around that centre, its substance is very vascular.

T A B U L A XVI.

A SEXTO cadavere, octavo mense, uterus a latere dextro visus, atque ita injecta cera præparatus, et scalpello anatomico denudatus, ut vasorum ad eum accessus, eorumque primæ ramifications quam clarissime appareant. A parte postica, adhuc peritoneo lævi copertus est uterus. In parte autem anteriori, qua placenta interne adhærebat, stratum externum uteri substantiae cultro anatomico sublatum fuit.

AA Cervicis uteri, nullo modo dilatati, facies exterior. Os uteri literie inferiori A oppositum; ab qua parte deorsum, pars vaginæ, adipi tecta, relinquitur.

BB In uteri superficie strictura, ubi margine pelvis antea fuit circumdatus.

CC Tuba; post quam ovarium oblitescerat.

DD Ligamentum rotundum, cultro anatomico dissecatum; nimirum ut vena, ex parte inferiore tortuosa, binseque arteriæ convolutæ, ex vasis spermaticis omnes ortum ducentes, in conspectum venient.

E Arteria hypogastrica.

F Vena hypogastrica.

Vasa haec ramos quam plurimos ad vaginam dimittunt, et per latus uteri ascendunt, nimirum ut per anastomosin cum spermaticis suis propriis coeant. Partem utriusque trunci, administratione infusoria necessario destructam, ingenium pictoris supplevit.

G Arteria spermatica.

H Vena spermatica.

Vasa haec dum ad uterum accedunt, ramos quam plurimos, anterius ad tubam, et retrosum ad ovarium mittunt; deinde, ad uteri latus adventantia, per anastomosin cum hypogastricis coeunt. Rami precipui utriusque, hic, ad fundi uteri partem anteriorem, cui placenta adhærebat, cursum tenent.

II Peritonei, posticam uteri superficiem tegentis, margo.

P L A T E XVI.

FR OM a sixth subject, at eight months. A full view, from the right side, of the womb, so injected and dissected, as to shew the approach, and first general branchings of the uterine vessels. The back-part of the womb is still covered by the smooth peritoneum; but at the fore-part, to which the placenta adhered internally, the outer stratum of the substance of the womb was removed by dissection.

AA The outside of the neck of the womb, which was not in the least dilated. The os uteri was opposite to the lower letter A; from which downwards, there is a part of the vagina left, covered with fat.

BB A stricture on the womb, where it was surrounded by the brim of the pelvis.

CC The tube; behind which the ovary lay concealed.

DD The round ligament dissected, to shew a vein in its middle (twisted at the inferior part) and two convoluted arteries: all from the spermatic vessels.

E The hypogastric artery.

F The hypogastric vein.


These vessels send down numerous branches to the vagina, and mount upon the side of the womb, to anastomose with the respective spermatics. By fixing pipes, for the injection, and tying up these vessels, a part of each trunk was lost, and then supplied by fancy.

G The spermatic artery.

H The spermatic vein.

These vessels, in their approach to the womb, send up numerous branches, forwards to the tube, and backwards to the ovary; then, passing to the side of the fundus uteri, they anastomose with the hypogastric. The principal branches of both, in this case, go to the fore-part of the fundus uteri, where the placenta was attached.

II The edge of the peritoneum which covers the posterior surface of the womb.



*TA B. XVI. A sexto Cadavere, octavo mense. Uterus a latere dextra visus, atque ita
injecta ceni propagatus, et scalpulo anatomico denudatus, ut vasorum ad eum accessus,
et cum primis ramifications quam clarissime apparant.*

London Published July 1782 by E. CHURCH and Son, 37 Thomas-street, Finsbury.





*J.S. Agarwala delin.**London Published July 1st 1816 by T. Cope and Son, 37 Thoresby Street, Liverpool.**Engraved by J. S. Agarwala.*

TAB. XVII. Idem Uterus a fronte. Primo sive extimo substantia strato sublato, vasorum amplicum distributio, etiamque versus. Placentam (quae hic parti anteriori et fundo uteri adhaerbat) progressus melius conspicuntur.

T A B U L A XVII.

IDE M uterus a fronte. Primo, sive extimo substantiae strato sublato, vasorum distributio majorum, eorumque versus placentam (quae hic parti anteriori et fundo uteri adhaerebat) progressus, melius conspicuntur.

- A* Vaginæ extremitas summa, incisione aperta.
- B* Uteri orificium contractum.
- C* Uteri cervix nondum dilatata.
- DD* Ligamenta rotunda.
- EE* Tube: ovarii pone absconditis.
- FF* Fasciculus vasorum spermaticorum, ad latus uteri ascendentium.
- GG* Fasciculus vasorum hypogastricorum, ad latus cervicis uteri accedentium.
- HH* Trunci vasorum spermaticorum et hypogastricorum, ex anastomosi communes, ad uteri latera decurrentium. In fundi parte anteriori, arteria et vena, cursum versus placentam tenentes, conspicuntur: vasa minora, rotundiora, magisque convoluta, arteriæ sunt; majora, et planiora, venæ.

P L A T E XVII.

FROM the same subject. A direct fore-view of the womb, after the outer stratum of its substance had been dissected off, to shew the distribution of the larger uterine vessels in their way to the placenta; which, in this case, adhered to the fore-part and fundus of the womb.

- A* The upper extremity of the vagina laid open.
- B* The orifice of the womb, contracted.
- C* The neck of the womb, not stretched.
- DD* The round ligaments.
- EE* The tubes: the ovaries being concealed behind them.
- FF* The group of spermatic vessels, mounting up towards the sides of the fundus of the womb.
- GG* The group of hypogastric vessels, coming to the side of the neck of the womb.
- HH* The common, or anastomosing trunks of the spermatic and hypogastric vessels, running along the sides of the womb. On all the fore-part of the fundus are seen the arteries and veins, passing towards the placenta: the smaller, rounder, and convoluted vessels are the arteries; the larger and flattened vessels are the veins.

T A B U L A X V I I I .

Ex eisdem uteri fundus, eodem modo preparatus, ut amplissimum vasorum, venarum presertim, placenta ex adverso objectarum, crebrae inter se commissuræ contemplandæ prostarent.

- AA* Sectio per fundum uteri transverse ducta; inter
AAB Uteri parietem posteriorem, in statu suo naturali (scilicet, peritoneo levi cooperatum), et
AAC Uteri parietem anteriorem, ubi stratum substantiae ejus extimum sublatum est, ut vasa, ex adverso placentae sita, conspici possent. Vasa majora et planiora, ut in figura praecedenti, venæ sunt.
DD Tube. Post, sive infra eas, ovaria istiusmodi situm habebant, qui illa conspici non sineret.


- EE* Ligamenta rotunda, deorsum, paulo ante tubas, cursum tenentia.

In hoc cadavere, uteri paries anterior posteriori tumidior fuit; ita ut si uterus a vertice deorsum per tubas transcessus fuisset pars anterior, altera major extitisset. Re vera tamen, inter eas discrimen haud tantum fuit, quantum in hac tabula videtur; quippe uterus, paulo oblique oculo se offerens, plus superficie anterioris, quam posterioris, exhibebat.

P L A T E X V I I I .

From the same subject. The very fundus, or upper part of the womb, which was previously dissected to shew in the preceding figure as well as in this, the plexus of large vessels, especially the veins, opposite to the placenta.

- AA* The transverse section across the fundus of the womb, between *AAB* The posterior side of the womb left in its natural state; that is, covered with the smooth peritoneum, and
AAC The anterior side of the womb at its fundus, the outer stratum of its substance being removed, to shew the vessels opposite to the placenta. The larger and more flattened vessels, as in the preceding figure, are veins.
DD The tubes. The ovaries lay behind; that is, below them in this attitude, and farther from the eye of the spectator, so as to be hidden.
EE The round ligaments, going downwards, a little before the tubes.
- In this case, the anterior side of the womb made a greater swelling, or was more stretched than the posterior; so that if the womb had been vertically cut down through the two tubes, the anterior portion would have been larger than the other. Yet the difference was not so considerable, as may be imagined from a view of this figure, because the object was placed a little obliquely to the eye, which thence took in more of the anterior, than of the posterior surface.




J.C. Rymsdyk delineavit

M. Smith sculpsit.

TAB. XVIII. *Ex usdem Uteri Fundas, adem modo preparatus, ut amplissimorum vasorum.*
Venarum praecepsim Placenta ex adverso objectarum frequentes inter se commisurae
contemplanda posstantur.

London Published July 2^d 1765 by E. and N. S. Thomas & Son, 87 Fleet Street, Borough.







J. van Rossum dicit.

J. Forster Sculp.
TAB. XIX. Item Uterus, ejus facie anteriori incisa et in Fundum revoluta illa sis secundinis, per
quas caput Partus, obscurias licet, videtur. Hoc igitur visui sese offert iam. Superficies externa Placentae
ad dimidiatum item parum superficie interna uteri, ei respondens, una cum vasis que inter Uterum et
Placentam intercedunt. Partes Uteri extiores solis linearum duobus exprefas sunt.

London Printed July 1780 by S. Dods and T. & J. Cadell for Thomas Bowles.

T A B U L A X I X.

IDEM uterus; ejus facie anteriori incisa et in fundum revoluta, illæsis secundis, per quas caput fetus, obscurius licet, cernitur. Hic igitur visui sese offert cum superficies externa placenta, diuidatae quidem, tum pars superficiæ internæ uteri, ei respondens, una cum vasis quæ inter uterum et placentam intercedunt. Partes uteri exteriores lineis solummodo expressæ sunt.

- A** Vaginæ extremitas superior incisione aperta.
- B** Uteri orificium contractum.
- C** Uteri cervix nondum dilatata.
- D D** Ligamenta rotunda.
- E E** Tubæ; ovaria, pone iis sita, oculum fugiant.
- F F** Vasa spermatica.
- G G** Vasa hypogastrica.
- H H H** Uteri substantia incisa.
- III** Uteri paries anterior a secundis separatus, et supra fundum uteri reflexus.
- K** Chorion, amnion tegens. Per membranas illas pellucidas fœtus caput conspicitur.
- L L L** Decidua, qua uteri parieti intimo adhaerebat. Vasa quædam parva, in eam ex utero transmissa, conspicuntur.
- M M M** Uteri superficies interna, parti huic deciduae respondens.
- N N** Superficies externa partis illius placenta, quæ ab utero avulsa fuit.
- O O** Superficies uteri interior huic respondens.
- P P** In uteri superficie interna arteriæ convolutæ, quæ ad placentam antea transierant, jam vero disruptæ, dum uterū a placenta separabatur.
- Q Q** Arteriæ in placenta superficie illis respondentes.
- R R** Venæ ex placenta substantia excurrentes; in ea scilicet parte superficie ejus disruptæ, qua ad uterum transibant.
- S S** Vena in facie uteri interna, venis proxime dictis respondentes.
- T** In angulo, superficiem externam placenta, inter et superficiem internam inversam uteri, arteria convoluta, ab utero in placentam continuata, conspicitur.
- V** Vena, arteriam proxime dictam juxta posita, et illi respondens, ex placenta in uterum continuata.

P L A T E X I X.

From the same subject. A fore-view of the womb. Its substance is cut through and turned up over the fundus, to shew a part of the membranes, through which the child's head is obscurely seen, and about half of the placenta; together with the corresponding internal surface of the womb, and the vessels passing between the womb and the placenta. The external parts of the womb are represented in outlines only.

- A** The upper extremity of the vagina laid open.
- B** The orifice of the womb, contracted.
- C** The neck of the womb, not stretched.
- D D** The round ligaments.
- E E** The tubes; the ovaries concealed behind them.
- F F** The spermatic vessels.
- G G** The hypogastric vessels.
- H H H** The substance of the womb, cut through.
- III** The fore-part of the womb raised up from the secundines, and turned over the fundus.
- K** The chorion covering the amnion; through which transparent membranes the child's head is seen.
- L L L** The decidua where it adhered to the inner surface of the womb, in which are seen some small vessels sent into it from the womb.
- M M M** The corresponding inner surface of the womb.
- N N** The outer surface of that part of the placenta from which the womb was separated.
- O O** The corresponding internal surface of the womb.
- P P** Convoluted arteries upon the inner surface of the womb, which had passed to the placenta, and were broken through in separating the womb.
- Q Q** The corresponding arteries on the surface of the placenta.
- R R** Veins emerging from the substance of the placenta, and broken through at its surface, where they were passing into the womb.
- S S** The corresponding veins on the inside of the womb.
- T** In the angle between the outer surface of the placenta, and the inverted inner surface of the womb, is here seen a convoluted artery, continued from the womb into the placenta.
- V** A corresponding vein, near that artery, continued from the placenta to the womb.

T A B U L A X X.

IDEM denuo uterus, sed omnino apertus, ut fœtum in situ naturali ostendat. Circa fundum quaqueversus substantia tam placentæ quam uteri incisa conspicitur.

Uteri partes exteriores, lineis leviter adumbratæ, eadem sunt, quæ in praecedenti figura, et in tabula decima septima ad plenum sunt expresse.

A A A Utero per medium diviso, dimidii inferioris substantiæ sectio, ad eam partem scilicet qua membranæ adhærebant.

B B B Uteri dimidii superioris substantiæ sectio, ad partem qua placenta adhærebat.

C C C Placentæ sectio; illius cava spongiosa cera flava, ex uteri arteriis, et cera rubra, ex venis impleta sunt.

D D D Gyri nonnulli funis umbilicalis, brachia inter et crura fœtus positi.

E E Sectioni uteri superimposita funis umbilicalis extremitas, ad partem illam placentæ, quæ fœtum præjacebat, excisa

Fœtus iconem explanare supervacaneum foret.

P L A T E X X.

From the same subject. A fore-view of the womb, fully opened, to shew the child in its natural situation. All around, at the fundus, the substance of the placenta, as well as that of the womb itself, is seen cut through.

The external parts of the womb, slightly sketched in outlines, are the same as in the preceding plate, and were fully expressed in Plate XVII.

A A A The section of the substance of the lower half of the womb, where the membranous part of the secundines adhered.

B B B The section of the substance of the upper part of the womb, where the placenta adhered.

C C C The section of the placenta: the spongy cavities of which contained both some of the yellow wax, from the arteries of the womb, and of the red wax, from the veins.

D D D Some turns of the navel-string, lying between the arms and the legs of the child.

E E The end of the navel-string laid over the section of the womb. It was cut off from its attachment at that part of the placenta which lay before the child.

The parts of the child require no explanation.

*J.H. Rymsdyk delin.**H. Rogers sculpsit.*

TAB. XX. *Idem denauo Uterus, sed omnino apertus, ut Tum in situ
naturali ostendat. Circa Fundum quaquaversus substantia, tam
Plurante quam Uteri incisa conspicitur.*

London: Published July 1788, by H. Cox and Son, 2, Thomas's Street, Borough.







Alio. Gegeen delin. TAB. XXI. A. Septimo cadavere, Septimo graviditatis mense. Aperto cruciformi incisione Utero, & redinatis rite cornubus,
per membrinas Fartum continentis Tinctus ipso aqua innutans conspicitur. Reliquae partes nudo tinas distincte sunt.

Pro. discent. scip.

T A B U L A / X X I .

A SEPTIMO cadavere, septimo graviditatis mense. Aperto cruciformi incisione utero, et reclinatis rite cornibus, per membranas fætum continentis, fœtus ipse aquis innatans conspicitur. Relique partes nudis lineis distinctæ sunt.

AA Femora.

B Mons veneris, ad partem qua abdominis incisio longitudinalis desinit.

C Scrobiculus cordis, qua abdominis incisio incipit.

DDDD Partium continentium abdominis anguli quatuor reflexi, ut viscera plene paterent.

EE Omentum, in latere dextro abdominis,

FF Colon, ad latus sinistrum abdominis transiens.

GGG Intestina tenuia, uteri latus sinistrum versus, quorum gyri nonnulli, tempore quo cadaver primo apertum fuit, ante uterum jacebant.

HHHH Uteri, cruciformi incisione aperti, anguli quatuor, a secundis subtus positis liberati et reflexi, ut contenta ejus in conspectum prodirent.

I Uteri superficie intima pars exigua, cui placenta adhaeserat, minus quidem complanata, magis autem sanguine imbuta quam pars illa que membranis vestita fuerat.

K Lobi parvi placente superficies parti proxime dictæ respondens.

L Membrane liquor amnii continentis, que ad partem uteri inferiorem, propter cadaveris declivem situm, fœtus partes superiores e conspectu fere tollunt; ad fundum autem uteri subsidentes, fœtus partes satis distincte in conspectu locant.

M In hoc angulo, uterum inter et secundas, pictor id exprimere voluit quod in cadavere satis manifestum fuit, scilicet, uteri et secundarum substantiam inter se mutuo continuatam: dum autem hanc partem separabamus, tenera illa internectens membrana, decidua, in duas lamellas abiit; quarum haec utero, illa chorio, adhaerebat.

P L A T E / X X I .

FR OM a seventh subject, at seven months. The womb opened by a crucial incision, and the four corners carefully separated, and turned aside from the secundines, so as to shew the child, and waters, through the enclosing membranes.

The adjacent parts of the subject are represented in outlines.

AA The thighs.

B The mons veneris, where the longitudinal incision of the abdomen terminates.

C The pit of the stomach, where the longitudinal incision of the abdomen begins.

DDDD The four angles of the containing parts of the abdomen, inverted, to expose the bowels.

EE The omentum, in the right side of the abdomen.

FF The colon passing towards the left side of the abdomen.

GGG The small intestines on the left side of the womb; some turns of which, when the body was first opened, lay before the womb.

HHHH The four angles of the womb (which was opened by a crucial incision), separated from the subjacent secundines, and inverted, to expose the contents.

I A small part of the inner surface of the womb to which the placenta had adhered, more unequal and more bloody than what had been lined by the membranes.

K The corresponding surface of a small lobe, or extremity of the placenta.

L The membranes containing the liquor amnii, which, at the lower part of the womb, from the declivity of the object, cover the upper parts of the child from our view; and towards the fundus uteri collapse so as to shew the parts of the fœtus not indistinctly.

M In this angle between the womb and secundines, the artist endeavoured to express what was very apparent in the object, viz. the continuity of the substance of the womb and of the secundines; in parting which, the tender connecting medium, the decidua, separated into two layers, one of which clung to the womb, and the other to the chorion.

T A B U L A XXII.

IDEM cadaver. Contenta pelvis, quo situ disposita sunt, ex duplice patet figura: in altera, lineis nudis expressa, vaginam integrum vides; in altera, magis elaborata, dextra vaginæ latere dissecto et devoluto, os uteri in conspectum prodit.

- A* Femoris sinistri facies interna.
- B* Nates sinistra.
- C* Abdominis pars anterior.
- D* Mons veneris.
- E* Labium sinistrum.
- F* Labii dextri pars.
- GGG* Cutis et membrane adipose: sectio, eo consilio facta, ut in latere dextro, partes continentis pelvis removerentur.
- H* Ossis sacri superficies cartilaginea, qua ossi ilium dextro commissa fuerat.
- II* Os pubis dextrum, prope symphysin, transverse scissum.
- KK* In pelvis parte postica, musculi externi secti.
- LL* Musculi in facie ossis pubis externa etiam secti.
- M* Clitoridis crus dextrum, ubi ischio, quod hic sublatum est, coniectebatur.
- N* Vaginæ sphincter.
- O* Musculus transversalis perinei.
- P* Ani sphincter.
- Q* Tumor ex utero in cavi pelvis parte superiori.
- R* Vesica urinaria, fere vacua, parte anteriori uteri et latere superiori vaginæ compressa.

Litteræ sequentes ad lineas adumbratas spectant.

- SS* Vaginæ facies externa situ suo naturali vesicam urinariam inter et
- TT* Rectum.
- VV* Musculi, levator ani scilicet et coccygeus lateris dextri, cultro anatomico a pelvis contentis separati, et in superficiem recisam devoluti.
- XX* Totius vaginæ substantia ex uteri cervice et vesica urinaria excisa et in rectum devoluta, ut os uteri in conspectum prodiret.
- Y* Oris uteri labium anterius.
- Z* Ejusdem labium posterius.
- Vagina ad hunc modum aperta, os uteri, nimirum adminiculò suo jam a latere sublato, dextrorsum propendebat.

P L A T E XXII.

FRom the same subject, shewing the situation of the contents of the pelvis. In the figure which is expressed by outlines, the vagina is whole; but in the more finished figure, its right side is cut out, and turned down to shew the mouth of the womb.


- A* The inside of the left thigh.
- B* The left nates.
- C* The anterior part of the abdomen.
- D* The mons veneris.
- E* The left labium,
- F* Part of the right labium.
- GGG* The section of the skin and adipose substance, which was made to remove the containing parts, on the right side of the pelvis.
- H* The cartilaginous surface of the sacrum, by which it was joined to the right os ilium.
- II* The right os pubis, cut through near the symphysis.
- KK* The external muscles, on the back part of the pelvis, cut through.
- LL* The muscles which lie on the outside of the os pubis, cut through.
- M* The right crus clitoridis, where it was fixed to the ischium, which is removed.
- N* The sphincter vaginae.
- O* The transverse muscle of the perineum.
- P* The sphincter ani.
- Q* The swelling formed by the womb, in the upper part of the cavity of the pelvis.
- R* The bladder, almost empty, compressed between the anterior part of the womb and the upper side of the vagina.

The following letters refer only to the figure of outlines, viz.

- SS* The outside of the vagina in its natural situation between the bladder and rectum.
- TT* The rectum.
- VV* The levator ani and coccygeus muscles of the right side, by dissection turned down from the contents of the pelvis over the cut surface.

The following letters relate only to the more finished figure, viz.

- XX* The whole substance of the vagina, in the right side, cut from the neck of the womb and bladder, and turned down over the rectum, to shew the orifice of the womb.
- Y* The anterior lip of the orifice.
- Z* The posterior lip of the same.
- When the vagina was thus opened, the lateral support being removed, the os uteri pushed out towards the right side.



Ridley delin. *Prof. Charnier sculp.*
TAB. XXII. *Idem Cadaver Contenta Pelvi, quo situ disposita sunt, ex duplice patet figura.*
In illa elaboratori, dextra Vagina latere dissecto & devoluto, Os Uteri in conspectum prodit.





*J. C. Leydig sive didit.**M. H. K. Schaeffer sculps.*

TAB. XXIII. Octavum Cadaver, Sesto Gestationis mense. Uteri ac membrinarum parte anteriori sublata, exhibetur Fetus cum parte Placentae et Tuniculae umbilicalis..

Uteri Vasa aeria impletar erant.

London Published by AND PHILIPps, 57 Cheapside, & Thomas Bowles, Strand.

TABULA XXIII.

OCTAVUM cadaver, sexto gestationis mense. Uteri ac membranarum parte anteriori sublata, exhibetur fetus cum parte placentae et funiculi umbilicalis. Uteri vasa cera impleta fuerant.

- A Vesica urinaria, respectu habitu ad uterum, in situ, suo naturali; modice distenditur, et ramis quibusdam majoribus venarum hypogastricarum circumtegitur.
 B Vagine partis postice facies interna.
 C C Vasa hypogastrica ad cervicem uteri decurrentia et ramos ad vesicam urinariam et vaginam transmittentia.
 D D Vasa spermatica, duplicationem ligamenti lati intrantia.
 E E Tuba.
 F F Fimbriae.
 G G Ligamenti lati lamella posterior: anterior cultro anatomico sublata fuerat, ut vasa spermatica, ad fundum uteri ascendentia, oculo plenius occurrerent.
 H H Ligamenta rotunda. In sinistro, arteria praecipua, convoluta, a spermatica descendens conspicitur.
 III Uteri substantiae totius et membranarum sectio, qua uteri et secundarum paries anterior, ad eorum contenta exhibenda, sublatus fuerat.
 K Funis umbilicalis, prope locum ubi in placenta desinit.
 Placenta uteri parti postice, versus fundum, adhaerebat. Funis umbilicalis, primo deorsum, super humerum sinistrum, deinde sursum, pone corpus infantis, ad finem suum in placenta transibat.

PLATE XXIII.

FROM the eighth subject, at six months. A fore-view of the womb, which was injected; the anterior part, both of the womb and of the membranes, having been cut away, and the liquor amnii taken out to shew the fetus, with a part of the placenta and of the navel-string.

- A* The bladder, in its situation with respect to the womb. It is moderately distended, and is covered with some large branches of the hypogastric veins.

B The inside of the posterior part of the vagina.

C C The hypogastric vessels, going into the neck of the womb, and sending branches to the bladder and vagina.

D D The spermatic vessels, going into the duplicature of the broad ligament.

E E The tubes.

F F The fimbriae.

G G The posterior lamella of the broad ligament: the anterior had been removed by dissection, to give a clearer view of the spermatic vessels, in their ascent to the fundus of the womb.

H H The round ligaments. In the left is seen a large convoluted artery, coming down from the spermatic.

III The section of the whole substance of the womb, and of the membranes, by which the fore-part of the womb and of the secundines was removed, to expose their contents.

K The navel-string, near its termination in the placenta.
The placenta adhered to the posterior part of the womb, towards the fundus. The navel-string passed first downwards, over the left shoulder, and then upwards, behind the body of the child, to its termination at the placenta.

T A B U L A XXIV.

Ae eodem cadavere.

FIG. I.

Placenta uteri fundo et parti posticæ adhaerescens; substantia ejus spongiosa, injecta per vasa uterina cera, turgescit.

Pleraque partes lineis adumbratae, in tabula praecedenti ad plenum sunt expositæ.

- A* Vesica urinaria.
- B* Vagina.
- CC* Vasa hypogastrica.
- DD* Vasa spermatica.
- EE* Tube.
- FF* Fimbriae.
- GG* Ligamenta lata.
- HH* Ligamenta rotunda.
- III* Uteri et membranarum sectio.
- K* Funiculus umbilicalis.
- LL* Placenta utero adhaerescens.

Nulla pars cœriæ, in vasa uterina injecta, ad ramos vasorum illorum que funem umbilicalem conficiunt, transierat; vasa illa sanguinem solummodo continentia, obscure designata erant, ubi ex fune umbilicali in faciem internam placente se immitebant. Cellulæ autem in placente parte spongiosa, omnes eodem modo cera, vel corulea in venas uteri injecta, vel rubra in arterias infusa, turgescabant. Haec rem indicat figura secunda.

M Membranæ ex margine placente excuntes, et uterum undique investientes.

Placentæ margo hic paulo elatior est, et magis conspicuus, substantia ejus spongiosa cera distensa.

FIG. II.

Portio aliqua placente transverse secta, ut substantia ejus spongiosa appareret, ejusdemque cera adimplete crassitudo.

- A* Placentæ superficies interna.
 - B* Placentæ superficies externa.
 - C* Membranarum portiuncula, ex margine placente, cera injecta turgescantis, et in figuram rotundiorem adactæ, exiens.
- Cerae coruleæ, primum per venas uteri injecta, maxima pars ad superficiem ejus internam propulsa fuerat; ceraque rubra, postea per arterias injecta, in partibus exterioribus restabat; per totam autem ejus substantiam, colores hi duo plus minusve commixti sunt.

FIG. III.

Arterias exhibet illius partis membranæ deciduae, quæ parietem uteri anteriorem investiverat.

FIG. IV.

Vene ejusdem partis deciduae.

Figuræ hæc due separatis, aliqua ne fieret confusio, delineatae sunt.

PLATE XXIV.

From the same subject.

FIG. I.

The placenta, adhering to the fundus and back-part of the womb; its spongy substance is filled by the injection of the uterine vessels.

Most of the parts in outlines were more fully represented and explained in the preceding plate.

- A* The urinary bladder.
- B* The vagina.
- CC* The hypogastric vessels.
- DD* The spermatic vessels.
- EE* The tubes.
- FF* The fimbriae.
- GG* The broad ligaments.
- HH* The round ligaments.
- III* The section of the womb and membranes.
- K* The navel-string.
- LL* The placenta, adhering to the womb.

None of the wax, injected into the vessels of the womb, had passed into the branches of those vessels which compose the navel-string; and as they contained only some blood, they were not distinctly marked, where they spread, from the navel-string, over the internal surface of the placenta. But the cells, or interstices in the spungy part of the placenta, were universally loaded with wax; either the blue, which was injected into the veins of the womb, or the red, which was thrown into the arteries. This is illustrated by Fig. II.

M The membranes, coming out from the edge of the placenta, and investing the womb all around.

The edge of the placenta, in this case, was much more elevated and distinct, its spongy substance being distended.

FIG. II.

A section of half of the placenta, principally to shew what thickness it had acquired, by its spongy cavities being filled with wax.

- A* Its internal surface.
 - B* Its external surface.
 - C* A small portion of the membranes, going off from the edge of the placenta, which was thickened, and rounded, by the injected wax.
- Most of the blue wax, which was first injected by the veins of the womb, was driven on towards the internal surface; and the red wax, which was afterwards injected by the arteries, was lodged principally in the outer parts; but the two colours were, more or less, blended through the whole.

FIG. III.

The decidua, from the fore-part of the same womb, with the convoluted uterine arteries, which are dispersed through it.

FIG. IV.

The same part of the decidua, with the uterine veins, which ramify through it.


These two figures were drawn separately, to avoid confusion.

Fig. I.




Fig. IV.

Fig. III.



Ab eodem Cadavere. Fig. I. Placenta Fundo et postice Uteri parti adhaesens. Substantia ejus spongiosa injecta per vasa uterina cori turpissim. Fig. II. Portio aliqua Placentae transversim secta, ut substantia ejus spongiosa apparet, ejusdemque cori adimplita crufitio. Fig. III. Sistit Arterias. Fig. IV. Venas illius parti membranae Dividitur, qua parietem Uteri anteriorem investient.







TAB. XXV. . Nonum cadaver quinte mense. Uterus apertus omnino. ac Tinctus inde exemptus, quo subtilius
judicium feruntur de Feto ipso; de partibus ejus, cum invicem comparatis et dimensis; nec non de conditione
Cervicis Uterina quo sub hoc tempore obtinet.

J. C. Stompholus did.

Hand. sculp.

London Published July 1st 1788 by E. Cole and Son. 37 Threadneedle Street, Bishopsgate.

T A B U L A XXV.

N O X U M cadaver, quinto mense. Apertus omnino uterus, ac fœtus inde exemptus; quo subtilius judicium feratur de fœtū ipso, de partibus ejus secum invicem comparatis et dimensis, nec non de conditione cervicis uterinæ quæ sub hoc tempore obtinet.

Fœtus ipse, funisque umbilicalis illustratione non eagent.

A A A Vulnus per membranas factum, quo fœtus eximeretur; unde internæ faciei placentæ pars, circa finem funis umbilicalis, cernitur.

B B B Amnion et chorion collapsa, et decidua reflexa (quæ nondum caeperat deciduae ipsi adhaerescere) externe coopta.

C C Tubæ.

D D Ligamenta lata.

Literas sequentes alteri figuræ separatim insculptas invenies.

E Decidua, osnem illam partem uteri, quam placenta non occupaverat, investiens.

F F F F Substantia cervicis uteri, ab altero latere ad alterum secta.

G Pars superior, angustior, et laevior introitus cervicis uteri, ubi decidua manifeste continuata fuit in membranam ejus internam.

H Pars inferior, latior, et rugosa, ejusdem.

I Facies interna posterioris labii oris uterini, vesiculis, glutine repletis, bullata.

K Facies interna vicinæ partis vaginæ.

L L Portiunculae haec duæ membranarum fluitantium, exhibent, imperfecte quidem, partes duas cuticulae vaginam investientis, quæ in cadavere, de quo agitur, distinete admodum, separabatur usque ad ipsam oris uteri prominentiam.

P L A T E XXV.

FROM the ninth subject, in the fifth month. The womb fully opened, and the fœtus taken out; to shew the exact dimensions and proportions of the child, and the state of the cervix uteri at this period of utero-gestation.

The child and its navel-string require no explanation.

A A A The opening made in the membranes for the extraction of the child, through which is seen the inside of the placenta around the termination of the navel-string.

B B B The collapsed amnion and chorion, covered externally with the decidua reflexa, which had not as yet contracted an adhesion to the decidua.

C C The tubes.

D D The broad ligaments.

The following letters are engraved on a separate figure in outlines.

E The decidua, lining that part of the womb where the placenta did not adhere to it.

F F F F The section, from side to side, of the substance of the cervix uteri.

G The upper, narrower, and smaller part of the passage in the cervix uteri, where the decidua was evidently continued down into the inner membrane of that passage.

H The lower, wider, and rugous part of that passage.

I The inside of the posterior lip of the os uteri, studded with small bags of gelly.

K The inside of the adjacent part of the vagina.

L L The two small pieces of floating membranes represent, though very imperfectly, two portions of the cuticular lining of the vagina; which, in this subject, was separated distinctly and evidently as far up as the projection of the os uteri.

TABULA XXVI.

DECIMUM cadaver, quinto graviditatis mense. Exhibetur status uteri retroversi.

FIG. I.

Apertum cruciali incisione abdomen, contractiori autem forma exhibitum, cum vesica urinaria præter morem distensa.
A Incisio longitudinalis a scrobiculo cordis incipiens.

- B Qua ad montem veneris desinentem.
- CC Partium continentium anguli superiores, super hypochondria reflexi.
- DD Anguli inferiores utriusque, supra spinam ossis ilium reflexi.
- E Vesica urinaria, ita urina sanguinolenta distensa, ut fundus ejus ad dimidium spatii, umbilicum inter et scrobiculum cordis, porrigeretur.

FIG. II.

- Vesica urinaria, forma pariter minutiori, per medium divisa, et ad imam partem aperta, ut situs oris uterini, hic præter naturam sursum spectantis, appearat.
- AA Pars dimidia vesicae super spinam ossis ilium utrinque reflexa.
 - B Incisionis, a vertice ductæ, qua vesica urinæ aperta fuit, finis anterior.
 - C Unus recti intestini circuitus, qui in conspectum prodit ad ejusdem finem posteriorem.
 - Two inter has literas B, C, vesicie pars ima tumore, qui universum pelvum occupavit, elevata fuit: hic quoque, facta cruciali incisione parva, inter et paulo ante fines ureterum, os uteri in conspectum venit, pone situum, ut in hac figura, et paulo elatius quam pars superior symphysis ossium pubis.

FIG. III.

- Contenta pelvis omnia a parte postica; inter quæ, uterus retroversus maximum locum habet.
- A Rectum, ubi in cavum pelvis transibat, deligatum.
 - BBB Margines posteriores sectionis vesicæ a vertice ductæ.
 - CC Rectum, uterum inter et ossa, sacrum et coccygis, compressum.
 - DD Peritoneum, quod, ad sinistrum recti intestini latus, pelvum investiverat, ope catheteris, hanc partem inter et uterum immissi, elevatum.
 - EEE Pars peritonei, parti modo descriptæ respondens, ad dextrum recti intestini latus, percissa, et margo ejus F sepositus.
 - G Uteri facies anterior, retrorsum versa.
 - H Tuba ascendens ab uteri fundo, qui scilicet ad partem pelvis infimam devoluta fuerat.
 - I Ovarium ante tubam positum.

FIG. IV.

- Uteri portione, ac deciduae parte sublatis, fetus per pellucidas membranas, chorion nempe et amnion, conspicitur.
- A Os uteri sursus versum. Circum undique conspicuntur partium vaginae contiguarum interiora.
 - BB Uteri substantia divisa.
 - C Facies externa partis alieojus deciduae. Membrana haec densa, opaca, et arteriis parvis, injecta ab utero cera impletis, abundavit.
 - D Facies interna alijs partis membranæ ejusdem. Qua parte ablata fuerat decidua, per chorion pellucidum et amnion conspicitur fetus, cuius caput dependet, et collum fuso umbilicali circum-cingitur.
 - Lineæ albæ super chorio leviter adumbratae, vasorum tenuum, qua inde in deciduum transeunt, reliquias representant.
 - Cum hæc tabula delineabatur, liquor amnii sanguine exudante infectus fuerat. Membranarum autem pellucidarum facies convexa figuram satis distinctam, forma quidem contractam, fenes-træ, lucem admittentis, dedit.
 - Tubarum et ovariorum figuram huicce tabulæ inserere super-vacuum duxi.

PLATE XXVI.

From the tenth subject, in the fifth month, shewing the circumstances of a retroverted womb.

FIG. I.

- Shews, in miniature, the abdomen fully opened by a crucial incision, and the bladder enormously distended.
- A The beginning of the longitudinal incision at the pit of the stomach.
 - B Its termination in the mons veneris.
 - CC The upper angles of the containing parts, turned over the hypochondria.
 - DD The lower angles; each turned over the spine of the os ilium.
 - E The bladder, so distended with bloody urine, that its fundus reached half way between the navel and the pit of the stomach.

FIG. II.

- Drawn in miniature to the same scale, shews the bladder cut down through its middle, and opened, at its lower part, to shew the situation of the os uteri.
- AA Half of the bladder, on each side, turned over the spine of the os ilium.
 - B The anterior extremity of the vertical incision by which the bladder was opened.
 - C One turn of the rectum which was seen at the posterior end of the same incision.
 - Between these two letters, B, C, the depending part of the bladder was raised up by a tumor which possessed the whole cavity of the pelvis: and here, a small crucial incision having been made, between, and a little before the ends of the ureters, the os uteri was seen, as in this figure, situated behind, and a little higher than the upper part of the symphysis of the ossa pubis.

FIG. III.

- A back-view of the whole contents of the pelvis, consisting principally of the retroverted womb.
- A The rectum tied, where it passed down into the cavity of the pelvis.
 - BBB The posterior edges of the vertical section of the bladder.
 - CC The rectum, where it was pressed and flattened, between the womb and the sacrum and the os coccygis.
 - DD The peritoneum, which lined the pelvis on the left side of the rectum, raised up by a large catheter, introduced between it and the womb.
 - E EF The corresponding part of the peritoneum on the right side of the rectum, slit down, and the edge F turned off.
 - G The anterior surface of the womb, turned backwards.
 - H The tube, coming up from the fundus uteri, which was turned down, into the lowest part of the cavity of the pelvis.
 - I The ovary placed before the tube.

FIG. IV.

- The womb opened, to shew the secundines and their contents.
- A The os uteri, turned upwards. All around it is seen the inside of the adjacent parts of the vagina.
 - BB The substance of the womb, cut through.
 - C The external surface of part of the decidua. This membrane was thick and opaque, and full of small arteries, which were injected from the womb.
 - D The internal surface of another part of the same membrane. Where the decidua was removed, through the transparent chorion, and amnion, is seen the fetus, with its head depending, and the funis round its neck.
 - The white loose lines on the chorion are the remains of the filamentous vessels, which pass from it into the decidua.
 - The liquor amnii was become bloody, by transudation, when this figure was drawn; and the convex surface of the transparent membranes reflected a distinct miniature picture of the window which gave light.
 - It was thought unnecessary to introduce the tubes and ovaria.



London. Published July 1718 by E. and J. Thomas, near St. Paul's Church, Strand.

TAB. XXVI. Decimum cadaver, quinto graviditatis mense. Exhibetur status Uteri retrorsum. Fig. I. Uteri decupatio incisione. Abdomen, sed contractiori forma exhibatum, cum Utricula Utrinaria propter murem distentia. Fig. II. Utricula Utrinaria, forma pariter minutiori, per medium divisus et ad inam partem aperta, ut sedis Ovis Uterini hic propter naturam suorum spectantis apparet. Fig. III. Contenta Pelvis omnia a parte postica. Fig. IV. Uteri portione ac Dextor parte subtulata, fixata per pollucidas membranas, Cherion nempe et Amnion, non spicitur.

J. S. Sympath. del.

Engraved by





Fig. I.



Fig. II.



J. R. Greenhalgh delin.
TAB. XXVII Undecimum Cadavere, mense quinto aevante. Fig. I. Uterus cum appendicibus suis a tergo visus. Vagina secundum longitudinem incisa, Cervix et Uterus interne
statim apparet. Fig. II. Uterus omnino aperio, membrana Dordue Reflexa, Chorion invaginatum, cornutum, et Cervix Ovaque utrinque facies internae.

London Printed and Printed by G. C. and Son STATIONERS to the Royal

T A B U L A XXVII.

UNDICIMUM cadaver, mense quinto ineunte. Due figure. Arteriae et venae cera diversi coloris impletæ sunt.

FIG. I.

- Uterus a tergo visus, una cum vagina eo consilio incisa, ut status cervicis orisque uteri oculo objiceretur.
- A* Clitoris.
 - BB* Nymphæ.
 - C* Proxime hanc supra literam, orificium urethrae conspicitur.
 - DD* Vaginæ pars extrema inferior, rugis aspera.
 - EE* Ejusdem pars extrema superior, altera lievior, præcipue a parte postica.
 - F* Uteri orificium, in terminum vaginæ superiorem se projiciens.
 - GG* Tube.
 - HH* Fimbriæ.
 - II* Ovaria.
 - KK* Vasorum spermaticorum fasciculus, per ligamenta lata, ad ovaria, tubas, et fundum uteri ascendentium.

FIG. II.

- Utero omnino aperto, membrana decidua reflexa, chorio super extensa (per quas membranas fœtum videoas) et cervicis orisque uterini facies interna apparent.
- A, BB, C, D, D, EE, F, GG, H, II, KK*, vide fig. I.
 - L* Cervicis uteri facies interna rugosa, per gluten pellucidum cernitur.
 - MM* Substantia tam uteri quam decidua percissa.
 - NN* Decidua reflexa, membranas pellucidas investiens, alba, opaca, striata, distenta adeo tenuata, ut in locis plurimis fere pellucida videretur: deciduae, quæ hanc obtegebant, adhaerescere nondum ceperat.
 - O* Arteria convoluta, per deciduam reflexam, a margine placenta excurrens.
 - P* Vena huic similis.
 - Vasa fœtus sub cute repentina, sanguine adeo distincte adimpta sunt, ut profecto vasorum cadaveris, cera injecta turgentium, imaginem præ se ferrent.
 - Funis umbilicalis pars altera fœtus collum ambibat, talum sinistrum altera.

PLATE XXVII.

Two figures from the eleventh subject, in the beginning of the fifth month. The arteries and veins were injected with wax of different colours.

FIG. I.

A back-view of the womb, with the vagina slit up to shew the state of the cervix and os uteri.

- A* The clitoris.
- BB* The nymphæ.
- C* Immediately over this letter is the orifice of the urethra.
- DD* The lower end of the vagina, which is rugous.
- EE* The upper end, which is more smooth, especially behind.
- F* The orifice of the womb, projecting into the upper end of the vagina.
- GG* The tubes.
- HH* The fimbriae.
- II* The ovaries.
- KK* The bundle of spermatic vessels, passing up, in the broad ligaments, to the ovaries, tubes, and fundus of the womb.

FIG. II.

The same womb fully opened, shewing the decidua reflexa upon the chorion, (through which the child appears), and the inside of the neck and orifice of the womb.

- A, BB, C, D, D, EE, F, GG, HH, II, KK*, see fig. I.
- L* The rugous inside of the neck of the womb, seen through the transparent cement.
- MM* The substance, both of the womb and of the decidua, cut through.
- NN* The decidua reflexa, covering the transparent membranes, in white and opake striæ. It was become so thin, by extension, as to be rendered almost transparent in many places. It had not as yet contracted an adhesion with the decidua which covered it.
- O* A convoluted artery, branching through the decidua reflexa, from the edge of the placenta.
- P* A vein of the same kind.
- The cutaneous vessels of the child were as distinctly filled with blood as if they had been injected.

One turn of the navel-string was round the child's neck, and another was twisted round the left ankle.

T A B U L A XXVIII.

FIG. I.

IDEM uterus quem exhibet figura secunda tabulae praecedentis, sed inversus (i. e. fundo decliviori); ut, contentis suo pondere a cervice fundum versus devolutis, margo placenta clarus distinguatur; ac decidua, que ibidem ab interno uteri pariete in externam chorii superficiem reflexa, nomen decidua reflexa assumit, melius conspiciatur.

Partes lineis adumbratae in tabula praecedenti modo expositae sunt atque explanatae: scilicet *A, BB, C, DD, EE, F, GG, HH, II, et KK*, in figura prima; atque *L et MM* in figura secunda. *NNN* Decidua crassa, faciem internam uteri, ubi placenta non adhaerescat, investiens.

OOO Placenta margo, ubi lamina interna decidua a facie interna uteri, ad faciem membranarum externam, inflexa fuit atque continuata; quo in loco efficit.

PP Deciduam reflexam, eo magis attenuatam, quo longius a margine placentae distat.

Q Hic patuit, deciduam non eodem modo, quo ceterae membrane, trans aditum in cervice uteri se porrexisse; sed per paullulum spatiis intra hunc aditum, more investientis membrane, esse continuatam; tandemque sensim evanuisse, aut cum glutine illo, quod vice cementi fungitur, esse confusam.

FIG. II.

Ejusdem uteri facies interna, qualis apparuit, placenta jam separata.

A, BB, C, DD, EE, F, GG, HH, II, KK, Vide tabulae praecedentis figuram primam.

L, MM, Vide ejusdem tabule figuram secundam.

NNN Pars decidua, undique circa placenta marginem, etiamnum utero intus adhaerescens.

O Centrum superficie orbiculariae, in uteri parte anteriori, cui placenta hoc in cadavere adhaerescebat. Haec arteriis venisque abundabat, que scilicet utero ac placenta communies fuerant, quasque necessario perrupi, duas has partes dum separabam. Arteriae quidem parvæ erant, convolutæ, colorisque magis lucidi, ut ad literam *P*; vena autem perruptæ speciem macularum fuscarum, vel foraminum, magnitudinis haud exiguae, præ se ferebant, ut ad literam *Q*.

Linea inaequalis quoæ superficiem hanc searam, cui placenta adhaerescebat, ambit, extremum marginem placentæ a membranis abscisse designat; eadem etiam indicat locum quo decidua lamina interna reflexa fuerat, circum undique, in faciem chorii externam.

PLATE XXVIII.

FIG. I.

THE womb, from the same subject, turned upside down, that the weight of its contents might carry them towards the fundus; to shew the border of the placenta, where the decidua is reflected from the inside of the womb, to the outside of the chorion, at which last place it makes the decidua reflexa.

The parts represented in out-lines were seen and explained in the preceding plate, viz. *A, BB, C, DD, EE, F, GG, HH, II, and KK*, in the first figure, and *L, MM*, in the second.

NNN The thick decidua, lining the inside of the womb, where the placenta was not attached.

OOO The edge of the placenta, where the inner layer of the decidua was continued, by inflexion, from the inside of the womb to the outside of the membranes; at which last place it constitutes

PP The decidua reflexa, which becomes thinner, in proportion as it is farther distant from the edge of the placenta.

Q Here it was evident that the decidua was not, like the other membranes, extended across the passage in the neck of the womb, but was continued a little way down that passage, and there insensibly lost, or blended with the glutinous cement.

FIG. II.

The inside of the same womb, after the placenta had been separated.

A, BB, C, DD, EE, F, GG, HH, II, KK, See the first figure of the preceding plate.

L, MM, See the second figure of the preceding plate.

NNN Part of the decidua all around the edge of the placenta, left adhering to the inside of the womb.

O The centre of the circular surface at the forepart of the womb, to which the placenta, in this case, adhered. It was full of arteries and veins, which had passed between the womb and placenta, and which we broke through in separating these two parts: the arteries were small, and convoluted, and of a lighter colour, as at the letter *P*; the broken veins had the appearance of dark spots, or holes, of some considerable size, as at the letter *Q*.

The irregular line, which surrounds this rough surface to which the placenta adhered, points out the extreme border of the placenta, which was cut off from the membranes, and where the inner layer of the decidua, all around, was reflected upon the outer surface of the chorion.




FIG. I.


FIG. II.

Exemplaria
TAB. XXVIII. Fig. I. *Uterus quem adhuc sanguinem secundum Tabulae praeudenti, sed inverso (i.e. fundo doliori) ad ventritis suo pandere a Cervice.*

*Tundam vescas dentata, marginis Placenta clavis distinguatur, ac Dividit, quo dividit ab interno Uteri parite in extream dorsi superficem reflexa.
nonum. Dividit. Placenta operculum, medias con�ciatur. Fig. II. *Uteri quodcum uteri interni propterea avulsa Placenta se cohicit. Sicut illa irregulariter
asperam et irregulariter superficem circumferens, extremis inducit Placenta laniata, ab Dividit ab Utero etiamque in thoracem suarum efficit.**







*J. C. Gmelin delineauit
TAB. XXIX. Quinque figuræ exhibet, quarum tri primæ ex eodem sanguine Cadavere. Fig. I. Ovum
ittere exceptum. Fig. II. Pars Ductus vitro adhuc amplificata. Fig. III. Ovarium cum Corpus lateo.
Fig. IV. Portio membranarum nona mense. Fig. V. Pars Uteri et Placenta cum membranis.*

London Printed July 27th 1765 by E. CHAPMAN and SONS at Thomas's Street Stationery.

T A B U L A XXIX.

TRES figuræ ad idem exemplar depictæ; quibus adjiciuntur duæ aliae, tanquam appendices.

F I G . I.

Ovum utero exemptum, faciem placenteæ representans, laciniasque marginem ejus undique ambientes, membrane scilicet decidua reflexa dilacerata reliquias. Hæc figura cum figura secunda præcedentis tabulae congruit, faciem ovi exhibens, quod isti utero intus adhæserat.

AAAA Decidua reflexa, undique a margine placenteæ, ad chorion integendum, exiens.

BB Limbus, marginem placenteæ circum-ambiens, ortus scilicet ex sectione decidua reflexæ, ubi illa separata fuit a decidua, aut, si ita loqui mavis, a facie uteri interna.

Circulus, limbo isto circundatus, faciem externam placenteæ exhibet, que utero adcreverat. Has partes dum separabamus, arteriae venaeque plurimæ dilaceratae sunt, quarum pars altera utero adnexa est, altera placenteæ. Arteriae quidem, ut in figura præcedenti, parvæ, convolutæ, et colore sunt lucidiori, ut ad literas *CCC*; venas autem maculae latæ ac fuscae representant, ut ad literam *D*.

F I G . II.

Partis membranæ deciduae facies interna, ope vitri optici amplior visa, ut textura ejus propria, cibrum haud male referens, explicari posset.

F I G . III.

Ovarium, corpusque luteum percissa. Corpus luteum, etiam ad hoc temporis, cavatum esse manifeste apparuit.

F I G . IV.

Figura addita.

Pars aliqua membranæ deciduae, pertinentis ad partum legitimum, mensibus scilicet novem rite exactis.

F I G . V.

Alia figura addita.

Portio aliqua uteri et secundarum, nono mense, ita disposita, ut earum partium strata diversa distincte conspiciantur. Arteriæ cera rubra, venis flava est injecta.

AA Utter substantia discissa.

BB Placenta ei adhaerens, prope marginem percissa.

CC Decidua ea in parte utero adhaerens.

DD Decidua in illa parte ab uteri facie interna elevata, chorioque adhaerens.

E Chorion, supra faciem internam placenteæ continuatum, ubi separari non potest.

F Amnion elevatum a chorio, cui non, nisi per gluten aliquod mollissimum, adhæsit.

P L A T E XXIX.

THREE figures from the same subject, and two supplemental.

F I G . I.

The ovum taken out of the womb, shewing the external surface of the placenta, and the ragged edge all round, where the decidua reflexa was torn through. This figure corresponds with the second figure in the preceding plate, shewing the surface of the ovum which was attached to the inside of that womb.

AAAA The decidua reflexa going off, all round, from the edge of the placenta, to cover the chorion.

BBB The edge, formed all round the brim or border of the placenta, by cutting the decidua reflexa where it came off from the decidua, or, in other words, from the inside of the womb.

The round surface, enclosed by that edge, is the outer surface of the placenta, which had adhered to the womb. In separating those two parts, many arteries and veins were torn through, one part of each remaining with the womb, and the other with the placenta. The arteries, as in the preceding figure, are small, convoluted, and of a lighter colour, as at the letters *CCC*; the veins make broad dark spots, as at the letter *D*.

F I G . II.

The internal surface of a portion of the decidua, considerably magnified, to shew its peculiar cibriform or lace-like appearance.

F I G . III.

The ovarium and corpus luteum cut through; the latter of these, even at this time, had an apparent cavity.

F I G . IV.

A supplemental figure.

A portion of the decidua, from a common delivery at nine months; its vessels filled with red blood.

F I G . V.

Another supplemental figure.

Part of the womb and secundines, at nine months, so disposed as to exhibit distinctly the different strata of those parts. The arteries had been injected with red wax, the veins with yellow.

AA The substance of the womb cut through.

BB The adhering placenta cut through, near its edge.

CC The decidua, at that part left adhering to the inside of the womb.

DD The decidua, at this part raised from the inside of the womb, and adhering to the chorion.

E The chorion, which is continued upwards over the inside of the placenta, where it cannot be separated.

F The amnion, raised up from the chorion, to which it adhered by a tender gelly only.

T A B U L A XXX.

A DUODECIMO cadavere, quarto mense decurrente. Uterus a parte antica apertus, vasis cera repletus. Praecipue hic notanda occurunt, superficies placentaæ externa; vasa ex utero se immersentia in placentam; cervicis uteri externa, qualis facies, sit, et respectu ad vesicam urinariam urethramque habito, quem situm obtineat.

Placentæ superficies, sectione uteri substantia circumdata, adeo refert precedentem, scilicet Fig. I. ut explanatione minime indiget.

- AA** Vasa spermatica.
- BB** Fimbriæ, sive tubarum partes extremae.
- CC** Ovaria. Horum sinistrum corpore luteo nonnihil auctum fuit.
- DD** Ligamenta rotunda.
- E** Cervix uteri contracta.
- F** Facies interna portionis istius vesicæ, quæ vaginæ subjecta adhaeret. Utroque in latere, et hac litera paulo superius, cernitur orificium urethrae.
- G** Urethra universa omnino aperta.

P L A T E XXX.

FR OM the twelfth subject, in the fourth month, shews the injected womb, opened on its fore-part, to give a full view of the external surface of the placenta, with the vessels passing into it from the womb. It shews likewise the state of the cervix uteri externally, and its relation to the bladder and urethra.

The surface of the placenta, surrounded by the section of the substance of the womb, is so much like that of the first figure in the preceding plate, that it requires no explanation.

- AA** The spermatic vessels.
- BB** The fimbriæ, or ends of the tubes.
- CC** The ovaria. The left was enlarged by the corpus luteum.
- DD** The round ligaments.
- E** The cervix uteri, in a contracted state.
- F** The inside of that portion of the bladder which adheres to the subjacent vagina. On each side, and a little higher than this letter is seen the orifice of the ureter.
- G** The urethra opened in its whole length.



London Published July 1st 1760 by S. Cox and Son 87 Thosmore Street Borough.
TAB. XXX. A duodecimo Cadavere, quarti mense decurrente. Uterus a parte antica apertus, varis cori repletus.
Principae hic notandæ occurant, superficies Placenta externa; Vasa in Placentam ex Utero se immersentia;
Cervix uteri externa fœtus, quibus sit et respectu ad Uterum Uninatum leviter insque habito, quem sicut obtinat;







Fig. I.

Fig. II.

Fig. III.

L.F. Bourguignon del.

TAB. XXXI. Tres figure ab uestim Calavre. Fig. I. Uterus a parte postea utræ apertæ at. Fœtus Membrane Æntricæ uterine vix exponatur.

Fig. II. Parte postea Uteri. membrane Æntricæ Æntricæ. Totæ spæ per posticadas membranas apparet. Fig. III. Ovarium apertum corporis Lutei cævum ostendit.

T A B U L A XXXI.

TRES Figureæ ab eodem cadavere.

F I G. I.

Uterus a parte postica ita apertus, ut facies membranæ deciduae externæ, ac status cervicis uterinae visui exponantur.

- Literas tabulae huic explanandæ accommodatas omnes præter *II, K et L*, in figura secunda videas.
- A A* Vasa spermatica.
 - B B* Fimbriae.
 - C C* Ovaria.
 - D* Vagina universa, quaqua in longum extenditur, omnino aperta.
 - E* Os uteri in vaginam se protendens.
 - F* Via rugosa, per cervicem uteri ducens, unde gluten illud consolidans jam exciderat.
 - G G* Sectio substantiae cervicis uteri.
 - H H* Sectio substantiae uteri, unde
 - II* Pars posterior totius substantiae uteri de decidua sublata et in fundum uteri revoluta cernitur.
 - K* Decidua, adhuc membranis adhaerens, superficiem suam scabram externam exhibens, unde uterus fuerat separatus. Membrana hæc vasis, cera impletis, venis maxima de parte, abundavit. Plurimis autem in locis, cera, vasis effusa, hæc visu indistincta reddiderunt.
 - L* Membrane pellucideæ, amnion scilicet et chorion, decidua reflexa, tenuissima et pellucida obtectæ, in aditum cervicis uteri, ubi decidua ipsa deficit, viam sibi facientes.

F I G. II.

Idem uterus; postquam scilicet inversa pars uteri fuerat absissa, et decidua ablata, ea mente ut fœtus, in liquore amnii natans, per pellucidas membranas apparet.

Decidua reflexa hic adeo tenuis fuit, ut, nisi ad marginem placenta, vix percipi posset.

Fœtus, cuius crura fune umbilicali circum-vincta sunt, per se satis est manifestus.

F I G. III.

Corpus luteum ovario in sinistro percissum, ut appareat qualiter ejus cavum, sub hoc tempus gestationis. Qualem se habeat cicatrica illa, in externa facie corporis lutei ovarii sinistri, ibi cernitur, ubi stat litera *C*, in figuris prima et secunda.

P L A T E XXXI.

THREE Figures, from the same subject.

F I G. I.

A back-view of the womb, so opened as to expose the outer surface of the decidua, and to shew the state of the neck of the womb.

The letters for explanation, except *II, K*, and *L*, are seen in Fig. II.

- A A* The spermatic vessels.
- B B* The fimbriae.
- C C* The ovaries.
- D* The vagina laid open in its whole length.
- E* The os uteri protuberating into the vagina.
- F* The rugous passage through the neck of the womb, from which the viscous cement had, by this time, fallen out.
- G G* The section of the substance of the cervix uteri.
- H H* The section of the substance of the womb; whence
- II* The posterior part of the whole substance of the womb, raised up from the decidua, and turned over the fundus of the womb.
- K* The decidua left upon the membranes, shewing its scabrous outer surface from which the womb was separated. It was crowded with injected vessels, especially veins; and in many places the wax was extravasated, which occasioned indistinctness.
- L* The transparent membranes, viz. amnion and chorion, covered with a very thin and transparent decidua reflexa, pushing down into the passage of the cervix uteri, where the decidua is wanting.

F I G. II.

The same object, after the inverted portion of the womb had been cut off, and the decidua removed, to shew the fœtus, in the liquor amnii, through the transparent membranes.

The decidua reflexa in this subject was so thin, that it was scarcely perceptible, except near the edge of the placenta.

The fœtus, with the navel-string round its legs, requires no explanation.

F I G. III.

The corpus luteum, in the left ovary, cut through, to shew its cavity at this period. The appearance of the cicatricula, upon the outside of the corpus luteum in the left ovary, is seen where the letter *C* stands in the first and second figures.

T A B U L A XXXII.

FIGURE duæ, a decimo tertio cadavere, tertio mense prorsus exacto.

FIG. I.

Uterus a parte antica apertus, unde fœtus cervicisque uteri status in conspectum prodeunt.

- AA* Ligamenta lata.
- BB* Ligamenta rotunda.
- CC* Tubæ.
- DD* Fimbrie.
- EE* Ovaria.
- F* Vagina omnino aperta.
- G* Oris uterini labium posterius.
- HH* Via per cervicem uterinam.
- III* Sectio substantiæ uteri, portione quadam ovali excisa, ut contenta ejus sub examen venirent.
- K* Membrane percissa.
- L* Facie interne uteri pars.
- M* Pars ista deciduæ, qua inde separata fuit. Prope hanc literam cernitur foramen quoddam (locus nimurum ubi decidua deest), via que per cervicem uteri ducit, exadvorsum situm.
- N* Decidua reflexa, nec non chorion et amnion, percissa.
- O* Placenta parti posteriori uteri adhærescens.
- Fœtus cum collo fune umbilicali his circundato, abunde patet.

FIG. II.

Uteri, placente et membranarum secundum longitudinem sectio, cum foro juxta sito, funiculo autem umbilicali nondum diducto.

- AA* Vaginæ facies interna.
- BB* Os uteri percissum.
- CC* Cervix uteri percissa.
- DE* Via que per cervicem dicit, ejus, in parte posteriori *E*, rugæ penniformes conspicuntur.
- FFFF* Sectio fundi uteri, ubi conceptus sedem habebat.
- GG* Sectio placentæ, que retrorsum adhaeserat, ubi uterus, de quo agitur, preter morem crassus visus est.
- HH* Placentæ superficies interna, amnio et chorio obtecta.
- I* Uteri superficie interioris pars.
- K* Pars parva deciduæ, inde separata.
- LL* Decidue facies interna, sive cavum ipsam inter et deciduam reflexam.
- M* Angulus prope marginem placentæ, ubi decidue lamella interior super chorion reflectitur, nempe ad deciduam reflexam faciendam.
- N* Decidue reflexæ facies externa; sive cavum, membranam illam inter et deciduam.
- O* Margines secti deciduæ reflexæ et chorii, que inter se intime connectuntur.
- P* Margo amni sectus.

P L A T E XXXIII.

Two Figures from the thirteenth subject, at full three months.

FIG. I.

A fore-view of the womb opened, to shew the child, and the state of the cervix uteri.

- AA* The broad ligaments.
- BB* The round ligaments.
- CC* The tubes.
- DD* The fimbriae.
- EE* The ovaries.
- F* The vagina laid open.
- G* The posterior lip of the os uteri.
- HH* The passage through the cervix uteri.
- III* The section of the substance of the womb, where an oval piece was taken out to shew its contents.
- K* The membranes cut through.
- L* Part of the internal surface of the womb.
- M* That part of the decidua which was separated from it. Near this letter is the hole, or deficiency of the decidua opposite to the passage through the cervix uteri.
- N* The decidua reflexa, chorion, and amnion, cut through.
- O* The placenta, adhering to the posterior part of the womb.
- The fœtus, with two turns of the navel-string round its neck, requires no explanation.

FIG. II.


A longitudinal section of the womb, placenta, and membranes; with the child near it, but still attached by the navel-string.

- AA* The inside of the vagina.
- BB* The os uteri cut through.
- CC* The cervix uteri cut through.
- DE* The passage through the cervix; in the posterior part of which *E*, the penniform rugæ are conspicuous.
- FFFF* The section of the fundus uteri, in which the conception was lodged.
- GG* The section of the placenta, which adhered backwards, where this womb was remarkably thick.
- HH* The internal surface of the placenta, covered with the amnion and chorion.
- I* Part of the inner surface of the womb.
- K* A small part of the decidua separated from that inside of the womb.
- LL* The internal surface of the decidua; or the cavity between the decidua and decidua reflexa.
- M* The angle, near the edge of the placenta, where the inner layer of the decidua is turned over the chorion to form the decidua reflexa.
- N* The outside of the decidua reflexa; or the cavity between that membrane and the decidua.
- O* The cut edges of the decidua reflexa and chorion, which are intimately connected.
- P* The cut edge of the amnion.



TAB. XXXII. *Quae adhuc totum Cadaver. Figure, recto mense. Adhuc. Fig. I. Membrana a parte anteriori palpata, totam cum Cervice Membrana contumplandam dicitur. Fig. II. Idem Membrana remota longitudinem cum Placenta et membranis樵ivis, adhaerente etiam Totu. ac per fundum amniosum affinitate.*

London Printed by JAMES COOK and Son, STATIONERS, near Temple.







TAB. XXXIII. *Ova trax abortu perdita, sex figuris expedita.* Fig. I. *Conceptus novem circiter hebdomadarum, si parte que tota membranarum est oculus obiecta.* Fig. II. *Eiusdem sectio secundum longitudinem.* Fig. III. *Conceptus alter post octo fore hebdomadas.* Fig. IV. *Idem Conceptus ad extremum paucis curiosis revolutus.* Fig. V. *Chorion nudum, i.e. ab ambiente Decidua expeditum.* Fig. VI. *Idem Conceptus aperitus.*

T A B U L A XXXIII.

O V A I T A A B O R T I V A , S E X F I G U R I S E X P R E S S A .

F I G . I .

- Ovum non circiter hebdomadæ abortivum, ab ista scilicet parte que membranaceæ est adspicuum. Decidua lacerata est, & aliquantulum revoluta, ut decidua reflexa levius et opaca in conspectum prodiret.
- A* Decidua ab utero separata facies externa scabra.
 - B* Facies externa portianculæ cuiusdam placenteæ, cujus reliqua pars sedem suam exadvorū habebat.
 - CCCC* Decidua facies interna cribriformis, quæ primis mensibus graviditatis membranis inclusa non adhaerescit.
 - DDDD* Decidua margo laceratus, qui in marginem *EEE* membranæ ejusdem custum suum tenuerat.
 - F* Decidua reflexa per faciem externam chorii diffusa.
 - GG* Angulus partis reflexæ, ad marginem scilicet placenteæ, ubi lamella interior decidua in chorion reflectitur, simili fere modo quo lamella interior pericardii reflectitur ad superficiem externam cordis obtegundinatur.
 - H* Extrema pars decidua ad cervicem uteri.

F I G . II .

- Ejusdem sectio a vertice ad imum usque.
- A.A* Sectio placenteæ, quam superficie superiori et posteriori uteri adhaesisse supponendum est.
 - BB* Decidua pars anterior sectio.
 - CC* Decidua pars posterior sectio.
 - D* Pars extrema decidua ad cervicem uteri.
 - EE* Amni cava, in quo fetus, ope funiculi tenuissimi umbilicalis, a facie interna placenteæ suspenderatur.
 - F* Sectio trismembranarum, scilicet amni, chorii et decidua reflexæ, que non solum sunt contiguae, sed et sibi invicem adhaerescunt.
 - GG* Angulus ad marginem placenteæ, ubi lamella interior decidua in faciem externam chorii reflectitur.
 - H* Hic tres istæ membranae paululum sejunguntur, ut situs earum, respectu habitat ad placentam, manifestius apparet.

F I G . III .

- Ovum octo circiter hebdomadæ abortivum. Pars quedam exigua decidua excisa est, et sursum reflexa, ut cava inter eam et ceteras membranas emeretur.
- A* Decidua pars exigua excisa.
 - B* Conceptus pars ubi deest decidua, exadversum scilicet aditus per uteri cervicem ducenti.
 - CCC* Decidua superficies externa.
 - D* Decidua reflexa obtegens chorion et amnion, quæ membrane se protendentes cavum decidua adimplent.
 - E* Partis superioris placentæ facies extensa.

F I G . IV .

- Idem ovum, decidua jam incisione dessudatum aperta, angulis quatuor revolutis exindeque reflexæ portione rotunda cultro sublata et devoluta, nempe ut vascula in superficie externa chorii in conspectum prodirent.
- A, B* Vide Fig. III.
 - CCCC* Facies interna quatuor angulorum in quos decidua cruciali incisione fuerat divisa.
 - DDD* Decidua reflexa cæstæ membranæ investiens.
 - EE* Angulus ad placenteæ marginem, ubi lamella interior decidua protenditur supra faciem externam chorii, nonnihilque decidua reflexæ obtinet.
 - F* Decidua reflexæ portio rotunda, ex facie externa chorii cultro anatomico sublata et devoluta.
 - G* Chorion cum vasis suis hirsutis expositum. Vasa ista decidua reflexæ arte adhaerent, et eorum nouissima una cum membrana illa abscissa sunt.

F I G . V .

- Ovum eodem fere tempore abortu separatum, constans ex chorio solo, cum vasis et contentis ejus; quippe decidua, vel pars secundarum uterina, in utero remanebat.
- AAA* Vasorum hirsutorum rami majores et frequentes, ex quadam parte superficie externe chorii emergentes, et cum decidua, vel parte uterina, ad placentam conficiendam se commiscentes.
 - BB* Chorii ea portio quæ postea membrana uniformis et pellucida evadit; vasis paucioribus, subtilioribus, fluitantibus, quæ in decidua reflexa evanescent, absunt: per eam conspicitur embryo.
 - C* Vesicula umbilicalis in facie externa amni per chorion conspecta: filum exalbidum (reliquie arterie et venæ) ab ea ad umbilicum embryonis transit.

F I G . VI .

- Idem ovum apertum. Membranæ antice parti insidentibus a placenta abscissa et reflexis, distincte conspicitur embryo.
- A* Vesicula umbilicalis, ex qua.
 - BB* Reliquie arterie et venæ, filo albo similes, ad umbilicum embryonis transiunt. Intestinorum gyri quidam in principio funis umbilicalis includuntur.

P L A T E XXXIII.

SIX FIGURES OF THREE DIFFERENT ABORTIONS.

F I G . I .

- An abortion, of about nine weeks, seen on that side which is membranous. The decidua is torn, and turned somewhat aside, to shew the smooth and opaque decidua reflexa.
- A.A* The rough, external surface of the decidua, which exfoliated from the womb.
 - B* The outside of a small portion of the placenta, the rest of which was situated on the back-part of this object.
 - CCCC* The internal cribriform surface of the decidua, which, in the first months of pregnancy, does not adhere to the membranes which it encloses.
 - DDDD* The lacerated edge of the decidua, which had been continued into the edge *EEE* of the same membrane.
 - F* The decidua reflexa spread over the outside of the chorion.
 - GG* The angle of reflection at the edge of the placenta, where the inner layer of the decidua is turned over the chorion; much in the same manner as the inner lamella of the pericardium is reflected, to cover the outer surface of the heart.
 - H* The termination of the decidua at the cervix uteri.

F I G . II .

- A vertical section of the same.
- A.AA* The section of the placenta; which, we must suppose, had adhered to the upper and back-part of the womb.
 - BB* The section of the anterior portion of the decidua.
 - CC* The section of the posterior portion of the decidua.
 - D* The termination of the decidua at the cervix uteri.
 - EE* The cavity of the amnion, in which the embryo hangs by a slender navel-string, from the inside of the placenta.
 - F* The section of the three membranes, which are not only contiguous, but adhere to one another, viz. the amnion, the chorion, and the decidua reflexa.
 - GG* The angle, at the edge of the placenta, where the inner layer of the decidua is reflected over the outside of the chorion.
 - H* Here those three membranes are a little separated, to shew their course at the placenta.

F I G . III .

- An abortion of about eight weeks. A small strap of the decidua is cut out, and turned up, to shew the cavity between it and the other membranes.
- A* The cut slip of the decidua.
 - B* The part of the conception where there is no decidua, viz. opposite to the passage through the cervix uteri.
 - CCC* The external surface of the decidua.
 - D* The decidua reflexa covering the chorion and amnion, which project and fill up the cavity of the decidua.
 - F* The outside of the upper part of the placenta.

F I G . IV .

- The same, when the decidua had been opened by a crucial incision, and the four angles had been turned off, and then a round piece of the decidua reflexa dissected off, and turned to one side, to shew the loose vessels on the outside of the chorion.
- A, B* See Fig. III.
 - CCCC* The inside of the four angles or flaps into which the decidua was reduced by a crucial incision.
 - DDD* The decidua reflexa, covering the other membranes.
 - EE* The angle, at the edge of the placenta where the interior lamella of the decidua is continued over the outside of the chorion, forming the decidua reflexa.
 - F* A round portion of the decidua reflexa dissected from the outside of the chorion and turned aside.
 - G* The chorion, with its shaggy vessels, laid bare. These vessels adhered firmly to the decidua reflexa, and parts of them were cut off with that membrane.

F I G . V .

- An abortion of the same age, consisting of the chorion only, with its vessels and contents; that is, without the decidua, or uterine part of the secundines.
- AAA* The larger and more crowded branches of the shaggy vessels which shoot from the external surface of one part of the chorion, to mix with the decidua, or uterine part, to form the placenta.
 - BB* That portion of the chorion which afterwards becomes the uniform transparent membrane. It is covered with fewer and more delicate floating vessels, which lose themselves in the decidua reflexa. The embryo is seen through it.
 - C* The vesicula umbilicalis on the outside of the amnion, seen through the chorion; with a whitish thread (the remains of an artery and vein) leading from it, towards the navel of the embryo.

F I G . VI .

- The same opened. The membranes, which were at the fore-part, being cut from the placenta, and turned up, the embryo is distinctly seen.
- A* The vesicula umbilicalis, from which *B*, the remains of an artery and vein, in the form of a white thread, pass to *C*, the navel of the embryo, with some turns of the intestines lodged in the beginning of the navel-string.

TABULA XXXIV.

FIG. I.

Ovum, scilicet chorion, cum omnibus ejus contentis, quintam, ut licet conjicere, circiter hebdomadam. Litera, anteau uitat, ad dignoscendas partes, minime desideratur. Macule quae maxime fusce sunt, sanguinis coagulati reliquias parvas representant. In medio, globi instar, conspicitor amnios, per quod fokus indistincte appetit. Ovum ipsum hanc paululum mole sua complanatum exparsum fuit, coram pictore in patella dum jacebat.

FIG. II.

Idem ovum preparatum, magna satis parte chorii absissa, amnioque cum fuctu inclusa, ut partes circa umbilicum ejus conspicerentur, in latus reflexo, vitri optici ope, ut partes minores magis distincae possent conspici, amplificatum fuit. Vasa hirsuta fluitantia, ex facie externa chorii excutientia, per se satis patent.

A A Spatium inter chorion et amnion. Hoc refertur erat gelatina tenera, adeo pellicula ut vix cerni posset; unde arteria venaeque ramosa, sanguine rubro impletata, in facie interna placentae distincae per eam conspicibantur.

B Amnion liquore pellucido colorisque experie, purissima veluti aqua fontana, distendebatur; per quem fucus partes minores apparetur.

C Vesicula umbilicalis humore distensa: sequo annio illa, neque chorio connexa fuit; gelatina autem tessera circumdata, umbilico fucus per arteriam et venam, quasi per pediculum, adnexa; quibus quidem vasa tantum inter se vicinitas fuit, ut vas unum sanguine rubro impletum, et ramos in vesiculam umbilicalem solam emittens, videbantur.

Caput fortis trunco ejus longius fuit; brachia et crura non nisi per paululum emiserantur; viscera abdominis nondum fuerunt vestita; pars eorum magis fusca habet fuit; fons umbilici nondum orto, fortis faciei interne chorio et amnii, que eo loci contigua erant, medio abdonim, adnectebatur.

FIG. III.

Ovum integrum, quartae hebdomadae, ut licet conjicere. Ovi pars superior, latior et plana, sacculus fuit, textura tenera, per deciduum solam formatas: hinc ad angulos superiores foramen dentatum utrinque fuit, ubi tubas intrasse conjiceretur; et ad ovi partem inferiorem portio crassa angustaque ex membranis omnibus contentisque eorum constituta.

FIG. IV.

Ejusdem ovi, sectione a vertice ad imum ducta aperti, facies interna. Ovum totum adhuc fundi uteri formam triangulare retinebat.

Pars lata superior, decidue faciem internam levioriem, cribro similem, ostendit; et pars angusta inferior, omnium membranarum sectionem caravagum in conspectu ponit. Fetus primordia omnisimo soluta fiscerant. Amnion a chorio junctum fuerat, et in cavo ejus fluctabat. Ex duobus ad eodem centro circulus albus, interior est chorii sectio; cuius in facie externa stratum spongiosum, cui vasa ex chorio se immittunt, cernitur: circulus albus exterior, ad partem inferiorem, deciduum, placentam obtegenciam, et ad partem superioris deciduum reflexum representat.

FIG. V.

Ovum integrum tertiam circiter hebdomadam.

A A Seta in cavum ovi trajecta per foramen ad utrumque angulum superiore, quod scilicet tubae fallopianae extremitatem esse conjicere licet.

B Eadem sete per foramen majus exentes ad angulum inferiorem, quod cervici uteri exadversum esse ponimus.

C Hydatis, molis admodum exiguae, per deciduum sese projiciens, et filamenta gracilia ramosa ex superficie sua emittens: chorion esse, vix dubitandum.

FIG. VI.

Idem ovum, maxima parte parietis anterior excisa.

A A, B B, C. Vide Fig. V.
Manifestum est deciduum, hocce in ovo, membranam fuisse densam, texture glutinosa, quae toti cavo triangulare fundi uteri, tanquam vestimentum, adhaeserat; constabat etiam tubas in facie ejus interna fuisse terminatas; chorion inter duplicitum ejus membranam situm fuisse, vel substantia ejus circumdatum; parique passu cum incremento chorii, procedente uteri gestatione, carum decidue coartasse, lamellam ejus interioris (sive deciduum reflexum) distendens, donec tandem, delecto cavo, lamella illa interna facie interior decidue fuisse contigua. Progressus hic, de quo agitur, mutationis planius intelligetur, si proxime his figuris inter se conferantur, ordine earum inverso, primo scilicet nonam, deinde octavam, et postremo septimam.

FIG. VII.

Status ovi (qualem scilicet animo licet concipere) in utero sestate provectionis, ubi chorion, decidua reflexa obiectum, per pluminum spatii in cavum decidua porrigitur.

A A Decidue sectio, ubi certo temporis progressu, lamellam extenam duplicitis membranis, spongiosum vel notum chorion vulgo dicit, confecisset.

B In hoc loco, decidua partem placenter, uterinam dictam confecisset.

C Amnii cavum.

D Cavum, vel spatium, amnion inter et chorion.

E Cavum decidue, vel spatium membranam illam inter et deciduum reflexum.

F F Tuba in faciem internam decidue sese aperientes.

G Cervix uteri nullius partem ovi continens.

FIG. VIII. et IX.

Due haec figure, in hoc fere solo ab ultima differant; nimisimum chorii in decidua cavum projectio eo minor est, quo ovum sestate minus.

PLATE XXXIV.

FIG. I.

A CONCEPTION, viz. the chorion, with all its contents, supposed to be about five weeks. The parts will be understood without letters of reference. Some of the darkest spots represent the little remnants of clotted blood. In the middle the amnion is seen like a round ball, through which the fetus appears indistinctly. The whole object was considerably flattened and spread out, by its own weight, as it lay in a little dish before the painter.

FIG. II.

The same conception, prepared by cutting away a considerable part of the chorion, and turning aside the amnion with the enclosed fetus, that the parts about the navel might be seen. It is considerably magnified, to shew the small parts more distinctly.

The sluggish floating vessels, which shoot from the outer surface of the chorion, require no explanation.

A A The space between the chorion and amnion. This was filled with a tender jelly, so transparent as to be almost invisible; whence, the branching arteries and veins, filled with red blood, upon the inside of the placentae, were distinctly seen through it.

B The amnion distended with a liquor as transparent and void of colour as the clearest water; through which the minute parts of the fetus were distinctly seen.

C The vesicula umbilicalis, distended with a fluid. It was neither attached to the amnion nor the chorion; surrounded with the tender jelly; connected, as by a pedicle, to the navel of the fetus by an artery and a vein; which lay so close together, as to appear like one vessel filled with red blood, and dispersing its branches on the vesicula umbilicalis alone.

The head of the fetus was longer than the trunk: the arms and legs had shot out but a little way: the abdominal viscera were not covered: the darker part of this was the red liver: there being no navel-string, the fetus was attached at its abdomen to the inside of the amnion and of the chorion, which were contiguous at that place.

FIG. III.

An entire conception, supposed to be in the fourth week. The broad and flat upper part of the object was a bag of a tender texture, formed of the decidua only, with a ragged perforation at each of the two upper angles, where the tubes were supposed to have entered; and the thick and narrow portion, at the lower part of the object, consisted of all the membranes and their contents.

FIG. IV.

The inside of the same object, laid open by a vertical section. The whole conception retained still the triangular shape of the fundus uteri. The broad upper part shows the smoother and cribriform internal surface of the decidua; and the narrower part below shews the section and cavity of all the membranes. The rudiments of the fetus were dissolved. The amnion had separated from, and lay loose in the cavity of the chorion. Of the two, concentric, white circles, the interior is the section of the chorion; upon the outside of which is seen the stratum of spongy substance into which the vessels shoot from the chorion. The outer white circle represents, at its lower part, the decidua covering the placenta; and, at its upper part, the decidua reflexa.

FIG. V.

A complete conception, of about three weeks.

A A A bristle passed into the cavity of the conception, through a hole at each of the upper angles, which was supposed to be the termination of the fallopian tube.

B B The same bristles coming out through a larger hole at the lower angle, supposed to be opposite to the cervix uteri.

C A small hydatid, projecting through the substance of the decidua, which had slender branching filaments shooting from the surface, supposed to be the chorion.

FIG. VI.

The same conception, after a considerable portion of its fore-part had been cut out. A A, B B, C. See Fig. V.

It plainly appears that the decidua, in this case, was a thick membrane (of a gelatinous texture) which had lined and adhered to the whole triangular cavity of the fundus uteri; that the tubes terminated on its internal surface; that the chorion was lodged in its duplicitate, or was surrounded with its substance; and that in proportion as the chorion would have been extended, in the progress of gestation, it would have encroached upon the cavity of the decidua, stretching its interior lamella (or decidua reflexa) till at length the cavity being obliterated, that interior lamella would have come into contact with the inside of the decidua. This progressive change will be more clearly understood by comparing the three following figures in the inverted order, viz. IX. VIII. and VII.

FIG. VII.

The more advanced state of a supposed conception in the womb; when the chorion, covered by the decidua reflexa, is stretched a great way out into the cavity of the decidua.

A A The section of the decidua, where, in process of time, it would constitute the external lamella of the double membrane, which has been commonly named the spongy or false chorion.

B Here the decidua constitutes the uterine part of the placenta.

C The cavity of the amnion.

D The cavity or space between the amnion and chorion.

E The cavity of the decidua, or the space between that membrane and the decidua reflexa.

F F The tubes, opening upon the inside of the decidua.

G The cervix uteri, containing no part of the conception.

FIG. VIII. and IX.

These two figures differ from the preceding almost in this respect alone, that the projection of the chorion into the cavity of the decidua is less in proportion as the conception is younger.



J. C. Rengacharya

London Published July 1st 1803 by E. CHAT and Son, 50, St. Thomas Street, Bowsgate.

A. Cook engraver

TAB. XXXIV. Fig. I. *Conceptus, quinta circiter hebdomadae, nempe Chorion cum omnibus suis contentis.* Fig. II. *Idem conceptus con-*
spicillo amplificatus, ab orbe parte anteriori exasperia, et Utriusque Embryonem contineat, ut latius reflexo. Fig. III. *Conceptus integer, quartie-*
circiter Hebdomadis. Fig. IV. *Partes ejusdem interiores, sectione a vertice ad insim ducta, patetfactae.* Fig. V. *Conceptus integer, circa*
tertiam Hebdomadom. Fig. VI. *Idem, pars anteriori sublati.* Fig. VII VIII. IX. *Figures quadram fetus Uteri, in gravidis, secundam longi-*
tudinem defecte, ut ejus diversitas, in primis gravitatis Hebdomadis, status, qualis mihi videbatur, innotesceret.

