# The morbid anatomy of some of the most important parts of the human body / [Matthew Baillie].

#### **Contributors**

Baillie, Matthew, 1761-1823

#### **Publication/Creation**

Albany: Printed by Barber & Southwick, for Thomas Spencer, bookseller, 1795.

#### **Persistent URL**

https://wellcomecollection.org/works/x95gwhuk


#### License and attribution


This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org





## ZEITLIN & VER BRUGGE

JACOB ZEITLIN
JOSEPHINE VER BRUGGE ZEITLIN
Cable Address: Jabberwock

815 No. LA CIENEGA BLVD. • OLympia 2-0784

LOS ANGELES, CALIFORNIA 90069

BOOKSELLERS

Importers and Dealers in
Rare Books and Manuscripts,
Old Master Drawings and Prints.
Wants searched for and reported. Books and
Libraries bought. Catalogues issued.





We offer:

### BAILLIE, MATTHEW.

The Morbid Anatomy of Some of the Most Important Parts of the Human Body ... The First American Edition. Albany: Printed by Barber & Southwick, for Thomas Spencer, 1795.

8vo. [2]ff., viii, 248pp., [6]ff. Contemporary calf, red morocco label (worn). Hinges weak, lightly browned, ex-lib., but very good copy.

\$75.00

First American Edition of "the first systematic textbook of morbid anatomy, treating the subject for the first time as an independent science." - Garrison-Morton: 2281 (London, 1793 ed.).

Baillie (1761-1823), the nephew of William and John Hunter, was educated at the University of Glasgow and later at Oxford, where he studied under William Hunter. Baillie practised in London with great success and was the last and most eminent member owner of the Gold Headed Cane. He is best remembered for The Morbid Anatomy, in which he described many pathological conditions including emphysema of the lung and cirrhosis of the liver. See: Major, Classic Descriptions of Diseases, p.637.

The present edition is also of interest for the printers, John Bariber and Solomon Southwick, Jr. John Barber and his brother Robert, with the help of Governor De Witt Clinton and his political party, became the publishers of the newspaper, The Albany Register, in opposition to the older newspaper of Albany, The Gazette of the Federalists. Solomon Southwick, Jr., son of a patriot printer of Newport and Providence, went into partnership with John Barber (1792-1800), as publishers and editors. - Douglas C. McMurtrie, A History of Printing in the United States. II, p. 176.

Austin: 95. Evans: 28226.

Subject to prior sale; carriage extra

# BAILLIE, MARTHEW.

The Morbid Anatomy of Son Body ... The First American week, for Thomas Spageer, 17


Swo. [2]ff., vill, Zafipp., (wors), Mingos weak, Highnly

First American Edition of the

Baillie (1761-1823), the university of Ginsmow and the University of Ginsmow and Humber. Saillie processed in a for The Marinant ersenberg course of the including emphysees of the in the including emphysees of the including the including emphysees of the including the includin


The present edition is else of Solomon Southwick, Er. 1000 of Governor De Witt Citation in the newspaper, The Albany, The Carette of the Fill Patrick printer of Newsport and (1792-1800), as publishers and Printing in the United States.

Austin 95. Evens: 25226.










A. Thornson April 1822 "75 Nº 51



# MORBID ANATOMY

OF

SOME OF THE MOST IMPORTANT PARTS

OF THE

# HUMAN BODY.

BY

MATTHEW BAILLIE, M. D. F. R. S. FELLOW OF THE ROYAL COLLEGE OF PHYSICIANS, AND PHYSICIAN OF ST. GEORGE'S HOSPITAL.

THE FIRST AMERICAN EDITION.

### ALBANY:

FOR THOMAS SPENCER, BOOKSELLER, MARKET-STREET.

M,DCC,XCV.

SUMOTAMA CHAROM

THATIZOTIAL TRONG THE TOTAL

OF THE W

HUMAN BODY.

Tb 16 795 b

SHIDE OF HERE THE GREEK

Change To Real Property lies

Tanking in

237

# SIR GEORGE BAKER, BART.

PRESIDENT,

AND

THE FELLOWS OF THE ROYAL COLLEGE OF PHYSICIANS,

THIS VOLUME

IS,

WITH THE GREATEST RESPECTS

INSCRIBED

BY THEIR MOST OBEDIENT

HUMBLE SERVANT,

Matthew Baillie.

# PREFACE.

THERE are some diseases which consist only in morbid actions, but which do not produce any change in the structure of parts: these do not admit of anatomical inquiry after death. There are other diseases, however, where alterations in the structure take place, and these become the proper subjects of anatomical examination.

The object of this work is to explain more minutely than has hitherto been done, the changes of structure arising from morbid actions in some of the most important parts of the human body.

This, I hope, will be attended with some advantages to the general science of medicine, and ultimately to its practice. It is very much to be regretted that the knowledge of morbid structure does not certainly lead to the knowledge of morbid actions, although the one is the effect of the other; yet surely it lays the most solid foundation for prosecuting such inquiries with success. In proportion, therefore, as we shall become acquainted with the changes produced in the struc-

ture of parts from diseased actions, we shall be more likely to make some progress towards a knowledge of the actions themselves, although it must be very slowly. The subject in itself is extremely difficult, because morbid actions are going on in the minute parts of an animal body excluded from observation; but still the examination of morbid structure seems to be one of the most probable means of throwing light upon it.

Another advantage, arising from the more attentive examination of morbid structure, is, that we shall be able to distinguish between changes, which may have some considerable resemblance to each other, and which have been generally confounded. This will ultimately lead to a more attentive observation of symptoms while diseased actions are taking place, and be the means of distinguishing more accurately diseases. When this has been done, it will be more likely to produce a successful inquiry after a proper method of treatment.

Another advantage, arising from a more attentive observation of morbid structure, is, that we shall be fitted to detect diseased alterations in the organization of parts, which are but little or not at all known. This will lay the foundation of our inquiry into the diseases themselves, so that



with the disease, when he comes to examine the body after death, and finds some of the appearances that are described in this Treatise, will acquire a knowledge of the whole disease. He will be able to guide himself on such knowledge in similar cases, and also to inform others. It may, perhaps, too, lead him to a proper method of treatment.

When a person has become well acquainted with diseased appearances, he will be better able to make his remarks, in examining dead bodies, so as to judge more accurately how far the symptoms and appearances agree with each other; he will be able also to give a more distinct account of what he has observed, so that his data shall become a more accurate ground of reasoning for others.

The natural structure of the different parts of the human body has been very minutely examined, so that anatomy may be said to have arrived at a high pitch of perfection; but our knowledge of the changes of structure produced by disease, which may be called the Morbid Anatomy, is still very imperfect. Such changes have commonly been observed only in their more obvious appearances, and very seldom with much minuteness or accuracy of discrimination.

Any works explaining morbid structure, which I have seen, are very different in their plan from the present: they either consist of cases, containing an account of diseases and dissections, collected together in periodical publications, without any natural connection among each other; or consist of very large collections of cases, arranged according to some order. In some of these periodical works, the diseased structure has been frequently explained with a sufficient degree of accuracy, but in all the larger works it has been often described too generally. The descriptions, too, of the principal diseased appearances, have been sometimes obscured by taking notice of smaller collateral circumstances, which had no connection with them, or the disease from whence they arose. Both of these faults even too frequently occur in the stupendous work of Morgagni de Causis et Sedibus Morborum, upon which, when considered in all its parts, it would be difficult to bestow too high praise; besides, the bulk of these very large collections prevents them from being generally in the possession of practitioners, and also renders them more difficult to consult.

In the present work we propose to give no cases; but simply an account of the morbid changes of structure, which take place in the thoracic



very little of now, will afterwards be known perfectly.

The principal motive, which has induced me to undertake this work, is to render the morbid structure of parts more accurately and generally known, as one of the best means of advancing our knowledge of diseases.

Although I have ventured to lay this work before the Public, yet I am very sensible of its imperfections. There are some appearances described, which I have only had an opportunity of seeing once, and which, therefore, may be supposed to be described less fully and exactly than if I had been able to make repeated examinations. There are others, which I have seen long before I had any idea of undertaking this work, and which I may be supposed to have observed less accurately than if there had been a particular object in view. There are others still, which I have only had an opportunity of examining in preparations. In some of these, certain appearances may be supposed to be lost, which might have been observed had they been examined recently after death. All of these are sources of inaccuracy, which may be said in some degree to be unavoidable. I have endeavoured, however, to be accurate; and if the Public

should approve of the plan of this work, I shall be very careful, by the addition of new materials, and by repeated observations, to render it more perfect.

Show that the state of the stat

THE RESERVE OF THE PERSON NAMED TO SERVE OF T

and the same of the party of the same of t

### CHAPTER I.

Diseased Appearances of the Pericardium.

The pericardium, or the membrane, which furrounds the heart like a bag, and is reflected upon its furface, giving it a fmooth external covering, is liable to inflammation. This is not a very common difease, although it happens sufficiently often to afford frequent opportunities of examining its effects after death. The disease, from its nature, cannot be confined to any particular periods of life; yet, from what I have seen, I should believe that it takes place more commonly when the body has for some time arrived at the adult state, than either in childhood or in advanced age.

In inflammation of the pericardium, the membrane is frequently thicker than in its natural state, and I think is also a little more pulpy. This change depends upon additional matter being thrown into the membrane by the increased action



lace work, and junctions are often formed between that portion of it lining the pericardium, which is reflected like a bag, and that other portion lying upon the pericardium, which is the immediate covering of the heart. This matter has a very close resemblance, both in colour and structure, to the coagulable lymph of the blood, and is probably nothing else than this substance separated from the blood by a peculiar action of the small vessels of the pericardium.

At the fame time that this layer of pulpy matter is thrown out upon the inner furface of the pericardium, there is accumulated in its cavity, more or less of a brownish or yellowish fluid. There is fometimes only a few ounces of it; at other times more than a pint. In it there are floating loofe fhreds of the pulpy matter formerly explained, and there is also occasionally some mixture of pus. This fluid refembles in its properties the ferum of the blood, and has commonly been confidered as the ferum. It is probably feparated in part from the coagulable lymph while it forms the folid layer, on the infide of the pericardium, fimilar to the spontaneous separation of the different parts of the blood after bleeding; but I should believe that it was not wholly accumulated in this manner, because it is often in very large proportion to the quantity of the coagulable lymph.

Inflammation of the pericardium fometimes advances to form pus, although rarely. Of this I have feen one inftance. The pericardium was very much thickened, was inflamed, and lined with coagulable lymph, but there was no fign of ulceration in any part of it. This last circumstance will be more particularly noticed, when we come to speak of the diseased appearances of the pleura. The pericardium in this case, contained more than a quart of common pus. When the pericardium is inslamed, which forms the immediate covering of the heart, the muscular substance of the latter is ocasionally inslamed to some depth.

## Adhesions of the Pericardium to the Heart.

In opening dead bodies, adhesions of the pericardium to the heart, are not uncommonly found. The adhesion is sometimes at different spots; at other times is extended over the whole surface. It either consists of a thin membrane, or of a more solid matter. When it is a thin membrane, it resembles very much, the common cellular membrane of the body, and when the matter is solid, it differs little from the coagulable lymph of blood. Whether the adhesion be in the one way or the other, the matter of the adhesion is, in both

cases, capable of being rendered vascular from injection. The adhesion too, is, in both cases, formed from the pulpy matter formerly explained, for I have oftener than once had an opportunity of tracing its gradual changes into each. Such adhesions are to be confidered as the consequence of previous inflammation, and shew that an inflammation of the pericardium may be furvived. They connect the pericardium in different cases, more closely or loosely to the furface of the heart; and where the connection is close, the inflammation has probably been more recent: where it is loose, the inflammation has probably been of older date, so that time has been given for the adhesions to be elongated by the motion of the heart. It is worthy of remark, that where there is an adhesion of the pericardium to the heart, the latter fometimes pulfates fo violently, that it is impossible to diftinguish it from the pulsation of an aneurism.

## Dropfy of the Pericardium.

This difease is not uncommon, and I believe is most frequent at an advanced period of life. I have seen it however in persons considerably under the age of thirty; and it probably also happens occasionally in childhood. I have seen oftener than once both anasarca and ascites in children

under twelve years old, which is as improbable as the accumulation of water in the pericardium. Water is fometimes found accumulated in the pericardium, while there is none in any other cavity; but generally it is accompanied with the accumulation of water in the other cavities of the thorax.

This water varies a good deal in quantity, a-mounting in some cases hardly to two ounces, and in others to more than a pint. Although the quantity be large, which may happen to be accumulated, yet I do not recollect to have seen the pericardium very much stretched, but it has always appeared as if it could contain a greater quantity. It is probable therefore that the pericardium may really grow so as to keep pace with the accumulation.

The fluid, which is accumulated, is of a brown colour, having a darker or lighter shade in different cases, and resembles in its properties the serum of the blood. If the person should happen at the same time to have jaundice, then the sluid has a pretty deep yellow tinge from bile. It has however frequently a yellowish colour, like the serum, without there being any reason to suppose that bile has been circulating with the blood.

The accumulation of water in the cavity of the pericardium, may proceed from two causes.

The one is, that the small exhalant vessels, or pening upon the inner surface of the pericardium, may throw out an unusually large quantity of sluid into its cavity, which is not absorbed in the same proportion by the absorbent vessels of that part: the other is, that the sluid may be thrown into the cavity in the natural quantity, but may not be absorbed in the natural proportion, from a defect in the action of the absorbent vessels.

I once had an opportunity of feeing two or three fcrofulous tumours growing within the cavity of the pericardium, one of which was nearly as large as a walnut. They confifted of a white foft matter, fomewhat refembling curd, or new cheefe. The pericardium is a very unufual part of the body to be attacked by fcrofula, and therefore this must be a very rare appearance of disease. The tumours had probably been slow in their progress, as in scrofula generally, and this disease could not have been guessed at in the living body.

I have twice found (and it has been feen much oftener by an anatomist\* of the best authority) the pericardium so changed as to resemble a common ox's bladder in some degree dried, or like a common pericardium, which had been for some time exposed to the air. As the thorax and abdomen were entire in both cases, no opening whatever

<sup>\*</sup> Mr. Hunter.



## CHAP. II.

# Diseased Appearances of the Heart.

INFLAMMATION of the substance of the heart is a rare difease, and is most commonly connected with an inflammation of the pericardium. When the pericardium covering its furface is inflamed, the inflammation fometimes passes a little way into the fubstance of the heart. That part of it becomes much more crowded with fmall veffels than in its natural state, and there are fometimes to be feen a few spots of extravafated blood. The substance of the heart may however be inflamed without inflammation of the pericardium. I recollect an instance of this fort where no marks of instammation could be observed in that membrane, but where there was a little more water than usual accumulated in its cavity. In this case the pulsation had been fo strong, that it was impossible to distinguish it from the pulsation of an aneurism. Authors have mentioned cases of abscesses and ulcers\* of the heart, but these I am persuaded are ex-

<sup>\*</sup> Vid. Morgagni, Epist. XXV. Artic. 17. Vid. Bonet, Tom. 1, p. 849; and also Lieutaud, Tom. 2, p. 27.

tremely rare. It happens still more rarely that the heart becomes mortified, although this diseased state of it has also been observed.

In opening dead bodies there is very often to be feen upon the furface of the heart, a white opaque spot, like a thickening of the pericardium. This is sometimes not broader than a fix-pence; at other times as broad as a crown. It is most commonly on the surface of the right ventricle, and is very rarely to be seen either on the surface of the left ventricle, or of the auricles, although it is occasionally on both.

It confifts of an adventitious membrane, formed on a portion of the pericardium, which covers the heart, and may eafily be diffected off fo as to leave the pericardium entire. It is an appearance, I believe, of no confequence whatever, and is fo very common that it can hardly be confidered as a difeafe.

### Polypus.

This has been confidered by the older anatomists, as a very common and a very fatal disease. By many of the moderns it has been rejected as a disease altogether. It consists in a mass of the coagulable lymph silling up some of the large cavities

§ Vid. Lieutaud, Tom. 2, p. 33.

of the heart, particularly the ventricles, and extending into the neighbouring large vessels.

The coagulable lymph is of a yellowish white colour, fometimes of a very yellow colour, and has confiderable firmness. It fills up the cavity completely, or nearly fo, in which it is found; and in the ventricles it shoots out processes between the fasciculi of the muscular fibres. From this circumstance, probably, it has derived its name. It also extends into the large arteries, which arife from the ventricles, and is often moulded to the shape of the semi-lunar valves at their origin. Any examples of this appearance, which it has occurred to me to observe, have been chiefly in preparations, and had undoubtedly taken place after death. In order that the circulation may be carried on it is necessary that the cavities of the heart be free for the transmission of blood; and if any one of its cavities should be plugged up, the circulation would necessarily be stopped altogether. A polypus, however, plugs up the cavity of the heart in which it is formed, fo entirely as to prevent the circulation. It may be faid, perhaps, that polypi are formed gradually, and the circulation is carried on for fome time, although very imperfectly. We have no general evidence however of coagula of blood being formed in the ordinary circulation where there is

a healthy structure of the parts concerned in this function. When polypi are examined, there is the same fort of appearance through their whole substance, which shews that the whole coagulum had been formed at the same time. Both of those circumstances seem to contradict very strongly the opinion that polypi are formed during life. When polypi are formed, I believe that the coagulation of the blood does not take place very quickly after death. They are without any admixture of the red globules of blood, and therefore the blood has been sufficiently long in coagulating to allow these globules to separate from the other parts in consequence of their greater specific gravity.

The ordinary coagulations of the blood, which commonly do not fill up very fully the cavities of the heart (although inflances occasionally occur of this fort) take place pretty soon after death, because the red particles of the blood are generally arrested in the coagulum. It may be worth while to remark, that there is sometimes found a portion of a coagulum in one of the ventricles of a yellow colour, and with an oily appearance so as to resemble exactly fat. There is however no admixture of oil in it, and it possesses all the ordinary properties of the coagulable lymph. The colour of a coagulum sometimes depends on a portion of

bile having circulated with the blood during life, as in cases of jaundice; but it takes place also when there is no reason to suppose that bile is mixed with the blood. These appearances depend probably on certain circumstances of the coagulation; but what those circumstances are, it is very difficult to determine.

It fometimes happens, although I believe very rarely, that the heart becomes aneurifinal. This difeafe confifts in a part of it being dilated into a pouch, which is commonly more or lefs filled with coagulated blood. Of this difeafe I have only feen one inftance. The apex of the left ventricle was dilated into a pouch large enough to contain a finall orange, was much thinner than in the healthy structure, and was lined with a thick white opaque membrane. There was hardly contained in it any coagulated blood; but the quantity of the coagulated blood depends commonly on the fize of the bag.

This disease most probably arose from the muscular structure at the apex of the ventricle becoming weaker than in any other part, so that when the ventricle contracted upon the blood it was pushed against the weakened part, which was not fully able to resist its impetus, and therefore was gradually dilated. Had the strength of the apex of the left ventricle been in due proportion

to that of the other parts, it is impossible that the aneurismal swelling should ever have taken place.

The most frequent situation of aneurism within the cavity of the thorax, is at the arch of the aorta. In this difease the arch of the aorta is much enlarged beyond its usual fize, fometimes forming an uniform tumour, and at other times there are fmaller aneurifmal fwellings rifing out of the larger one. This enlargement of the artery, if very confiderable, is more or lefs filled with coagulated blood, which is disposed in concentric laminæ. The coats of the dilated artery are nearly of the same thickness with those in its natural state, and therefore in proportion as the fwelling increases, new matter must be deposited in the coats of the artery. This new matter is undoubtedly deposited with a view to prevent the artery from being fo foon ruptured as it would be otherwise, and is formed by the action of the vafa vaforum. A portion of the new matter may perhaps also be formed by the action of the parts immediately in contact with the artery.

The coats of the artery, both at the place where the aneurism is formed, and near it, are considerably altered from their natural structure. They are more readily divisible into different layers, and have often formed in them spots of bony matter. These spots are frequently of a yellowish



as large as a child's head at birth; the fkin then becomes in some measure dead, and cracks from distention at the highest point of the tumour, a portion of the coagulated blood is forced out by the impetus of the circulation, and the person is cut off instantaneously. The blood sometimes oozes out slowly, and the person sinks gradually under its loss.

Aneurisms at the arch of the aorta, as well as in every other part of the arterial system, arise from the coats of the artery being previously discased, which are thereby unable to resist sufficiently the impetus of blood that strikes against them. This is obvious, both from the discased structure of the coats of an aneurism itself, and of the artery in its neighbourhood.

I have also found very frequently diseased appearances in the arch of the aorta, which were not advanced far enough to produce aneurism. These consist in little white opaque spots being formed in the inner membrane of the artery, and its coats are more easily separable from each other than in the healthy state.

The reason why aneurisms take place more frequently in the arch of the aorta, than in any other part of the arterial system, is its curvature, which exposes it to the full impetus of the blood, propelled by the strength of the left ventricle.

Aneurisms hardly ever happen in the pulmonary artery, because there is no arch formed by the pulmonary artery, and the blood readily passes by two large branches into the substance of the lungs. It may not be improbable too, that the pulmonary artery may not be so liable as the aorta to those diseased alterations of structure, upon which aneurism ultimately depends.

Aneurisms in the arch of the aorta, as well as in every other part of the arterial system, happen much more rarely in women than in men.—
This arises from two causes. The one is, that women, from their sedentary life, are less liable to an encreased impetus of the blood, occasioned by excited circulation: the other is, that the arteries in this sex appear to be less liable to diseased alterations of structure. This is not at all peculiar to aneurism. Some other diseases prevail in the one sex from which the other is in a great measure exempt.

The time of life at which aneurisms are most frequent, seems to be about the middle age.— When very strong pulsation is to be felt of the heart, we are not always to consider the disease as aneurism, especially if the person be young and of the semale sex. I have known an instance where an adhesion of the pericardium to the heart, was attended with that very strong pulsa-

tion, which is usual in aneurism; and I have known another instance where there was an unufually strong pulsation at the time the heart was a little inslamed upon its surface, and there was a small quantity of water in the pericardium.

The three femi-lunar valves at the origin of the aorta, or of the pulmonary artery, are often found difeafed. This confifts in the deposition of a bony or earthy matter. When a small portion of this bony or earthy matter is deposited, the valves are only somewhat impaired in their function, but when the quantity of matter is considerable, they must loose their valvular function altogether. The communication between the ventricles and the arteries becomes very narrow, the circulation is extremely interrupted, and the person is at length destroyed. This disease most commonly takes place towards an advanced period of life, but I have seen an instance of it in a boy of ten years old.

It fometimes happens that the femi-lunar valves are confiderably thickened, and of an opaque white colour: in this case the coats of the artery in the neighbourhood I believe are commonly thickened and diseased.

There is a preparation in Dr. Hunter's collection, where one of the femi-lunar valves is thickened, having at the fame time little tenacity, and being of a brown colour, in which a confiderable rupture had taken place. It is very rare that fuch an occurrence happens, and in the prefent instance the rupture was so large that I believe it must have proved very soon fatal.

The valvular apparatus between the auricles and ventricles, is liable to the formation of bony and earthy matter as well as the valves, which are fituated at the origin of the two large arteries, but by no means fo frequently. What this depends upon it is very difficult to determine.—

These valves may perhaps be considered as belonging more to the veinal than the arterial system, and it is very certain that offisication takes place very seldom in veins, although very often in arteries.

The valvular apparatus between the auricles and ventricles, is also occasionally thickened, having lost all its transparency, and having an opaque white colour. The chordæ tendineæ are also thicker than natural; and the internal membrane lining the ventricles, is frequently at the same time a good deal thickened, appearing like a firm white membrane. Under such circumstances the heart is often found to be considerably enlarged beyond its usual size. I have also seen the valvular apparatus between the auricle and the ventricle in a state of inflammation, and co-

vered with a layer of coagulable lymph. This I-believe to be very uncommon.

It fometimes happens, and I believe chiefly in those who are advanced in life, that the heart at some part becomes thinner, and upon any great exertion bursts: the blood escapes into the cavity of the pericardium, and the person is instantly destroyed.

Of fuch cases I have seen one instance only, but have heard from the best authority of another. They both happened to men; and I mention these circumstances because men appear to be more subject to diseases of the heart and blood vessels than women. It is probable that persons dying from this cause have, on account of the suddenness of their death, been supposed to die of apoplexy.

Cases have occurred, although very rarely, in which a large quantity of blood has been accumulated in the cavity of the pericardium, but where no rupture could be discovered, after the most diligent search, either in the heart itself, or in any of its vessels. This appears very wonderful, and not at all what any person would expect a priori. Upon the supposition of there being no rupture, two conjectures only have occurred to me with regard to the possibility of such

an effect taking place, and they are both attended with confiderable difficulty.

The one is, that the veffels upon the furface of the heart, may have loft a part of the compactness of their texture, so that the blood may have escaped through their coats by transudation. The other is, that blood may have been thrown out by the extremities of the small vessels, opening upon the surface of that portion chiefly of the pericardium which forms the immediate covering of the heart, whose orifices may have been to a very uncommon degree relaxed.\*

It also happens, although I believe very rarely, that a heart is so imperfectly formed as to allow of life being continued for some length of time in a very uncomfortable state, but to be ultimately the cause of death. There are two cases of this fort described by the late Dr. Hunter,† and there is one specimen of this malformation preserved in his collection. The malformation preserved in the collection, consists in the right ventricle of the heart being extremely small, and the pulmonary artery being very small also which arises from it. At its origin from the right ventricle it is completely impervious. The ductus arteriosus is

<sup>\*</sup> See Med. Observations, Vol. 4, p. 330. Memoirs of Med. Society, Vol. 1, p. 238.

<sup>†</sup> Vid. Medical Observations, Vol. 6, p. 291.

open, but forms likewise a small canal, and terminates in the lest branch of the pulmonary artery. The right auricle is larger than its natural size, probably from the frequent accumulation of blood in it; and the communication between the two auricles, by means of the foramen ovale, is much larger than usual. The child in whom this malformation was found had its skin of a very dark colour, had very laborious respiration, and violent action of the heart. It lived only thirteen days.

In another case related by Dr. Hunter, the pulmonary artery was very small, especially at its origin, and there was a desiciency in the septum cordis, at the basis of the heart, large enough to allow a small thumb to pass through it. The person in whom this malformation of the heart was found lived about thirteen years. He never had a fresh complexion, but it was always dark, or tending to black. He was often seized with fits, especially when there was any hurry upon his spirits, or there had been any brisk motion of his body.

It is obvious that in these deviations from the natural structure, too small a quantity of blood must pass through the lungs to receive the benefit of respiration, and this will be more or less according to the degree of the deviation. The blood will from this cause be of a dark colour, as it is well known that it receives the slorid hue

from the influence of the air upon it in the lungs. Hence the colour of the skin must be necessarily fallow or dark, and this will be increased when the blood is more than usual accumulated in the veins. It is natural to think that in such structures of the heart, the circulation will be carried on with much more difficulty when it is excited beyond its usual standard. This may even be increased to such a degree that the circulation must for a short time be suspended altogether. It was from this cause probably that sits occasionally were produced, as related in one of the cases.

There is an example also in Dr. Hunter's collection, of a heart from a child, which had a hole in the septum ventriculorum at the basis of the heart, large enough to allow a goose quill readily to pass through it. The child was still-born at fix months, and the hole in the septum evidently arose from original malformation. This too is described by Dr. Hunter in the fixth volume of the Medical Observations. An instance somewhat similar to this has likewise been published by Dr. Pulteney in the third volume of the Medical Transactions, where the person, to whom this monstrosity belonged, lived to near sourceen years of age.

I do not know how far I ought to mention, confistently with the plan of this publication, that





ever may act upon these external vessels, so as to excite contraction in them, may be supposed capable of producing an accumulation of blood, as well as an increased action in the inner branches of the intercostals, many of which are distributed upon the pleura. Perhaps too, upon another principle, there may be a greater confent between the action of the external and internal vessels of the cheft, than in the body generally. If these observations be just, they would account for the very frequent inflammation of the pleura; but they are only to be confidered in the light of a conjecture. Whatever be the cause of it, the fact is undoubted that the pleura is more liable to inflammation than any other membrane investing cavities which have no external opening. This is fo much the cafe that one can hardly examine the cheft of any person, who has arrived at the adult state, without perceiving more or less the traces of the prefent or former inflammation.

When the pleura is inflamed, it becomes thicker than it is naturally, and is in some degree pulpy. There is also interspersed through it a great number of very small vessels containing slorid blood, and a layer of coagulable lymph is at the same time thrown out upon its surface. This layer is sometimes very thin, and at other times

of confiderable thickness. It is either smooth upon its surface, or it throws out many fine small slocculi, which exhibit the appearance of a rich lace work. There is also poured into the cavity of the thorax a serous sluid, in which are sloating many small broken laminæ of the coagulable lymph; and there is occasionally some mixture of pus.

The coagulable lymph, covering the pleura which forms the external membrane of the lungs, frequently adheres to that which covers the pleura that is reflected on the infide of the ribs and the intercoftal fpaces, either in small portions, or by extended surfaces. Upon such occasions I have sometimes been able to trace the gradual change of the adhesion from the nature of a coagulable lymph to that of a cellular membrane. This coagulable lymph is capable of being rendered vascular from injection, as we have already mentioned in the inflammation of the pericardium. When the pleura is inflamed which covers thelungs, the substance of the latter is frequently inflamed to some depth.

Adhesions in the Cavity of the Thorax.

Adhesions are often found between that portion of the pleura which covers the lungs, and that other portion of it which lines the ribs and the intercostal spaces, while there is no sign whatever of present inslammation. These adhesions are often partial, and then they are most commonly to be found at the upper and posterior part of the chest; but they are sometimes extended over the whole cavity. They either connect the parts together closely, and then they often consist of a firm thick membrane, or they connect them loosely, when they consist of a soft spungy membrane, which exactly resembles the common cellular membrane of the body. Such adhesions are the consequence of inflammation, and are perhaps the most common morbid appearance to be found in dead bodies.

## Empyema.

Pus is not unfrequently found accumulated in the cavity of the cheft, forming the disease called empyema. This may either arise from the blood vessels of the pleura being in such a state of inslammation as to form pus, or from the bursting of some abscess in the lungs, so as to evacuate its pus into the cavity of the thorax. When pus is formed by an inslamed state of the pleura, there is no occasion for ulceration to take place. The pleura is found entire, but is covered with a layer

of the coagulable lymph. This fact has been long ago afcertained by the late Dr. Hunter. The formation of the pus depends on a certain state of action in the veffels of the pleura. The pus may either be accumulated in the whole cavity of the cheft, or may be confined to a part of it by adhefions taking place between the lungs and the pleura, which invests the ribs and the intercostal fpaces. When pus is evacuated into the cavity of the cheft by the burfting of an abfcefs in the lungs, it is almost always confined within certain limits by adhesions. In cases of empyema, there is frequently no particular appearance of the cheft observeable on the outside: there is sometimes, however, a fullness to be perceived externally on the fide where the matter is accumulated, and even occasionally an evident swelling between two of the ribs, as of matter pointing. Ulceration has also been known to take place in one or more of the intercostal spaces, so that the matter has been evacuated externally. There is an example in Dr. Hunter's collection, where the matter had been evacuated from the cheft by a great many openings in the intercostal spaces.

# Hydrothorax.

A watery fluid is not uncommonly found in one or both cavities of the cheft, forming the disease called hydrothorax. It is often attended with the accumulation of water in other parts of the body, especially in the pericardium, and the cellular membrane of the lower extremities .-The fluid in hydrothorax is commonly of a brown or yellowish colour, but occasionally has a red colour arifing from the mixture of the red globules of blood. It refembles in its properties the ferum. It is found to vary a good deal in quantity in different cases, sometimes amounting only to a few ounces, and at other times to feveral quarts. When it is accumulated in very large quantity in either fide of the cheft, that fide appears to be fuller to the eye externally, and when the cavity is laid open after death, the lungs on that fide are found more or less compressed. I have feen a lung fo compressed, as not to be larger than the closed fift.

Water is likewise found in the cavity of the chest where there are considerable adhesions.—
This shews that a good deal of inflammation had formerly taken place, which had probably, by

throwing out a confiderable quantity of ferum, laid the foundation of the hydrothorax.\*

## Offification of the Pleura.

It fometimes happens, although I believe rarely, that a portion of the pleura is converted into bone. This confifts of a thin plate, and fometimes extends over a pretty broad furface of the pleura. In all cases which I have seen, the bony matter feemed to me to be exactly like ordinary bone. I have never feen it form a thick irregular knob, but always a thin plate. The caufe which first excites this diseased process it is very difficult to determine; but there can be no doubt that the bone is formed by the finall veffels of the pleura taking on the fame mode of action which vessels do in the formation of ordinary bone.-This process is not peculiar to the pleura, but takes place in almost every part of the body: I believe, however, that it is more common in the pleura than in any other fimilar membrane. In the cases which I have observed, this process feemed not to have been attended with much in-

<sup>\*</sup> This circumstance is illustrated in several instances by Mr. Cruikshank, in his Treatise upon the Absorbent System. See 2d edition, p. 116.

convenience. There was no inflammation found in the pleura furrounding the bone, nor in the fub-flance of the lungs under it. One could imagine, however, if the bone were to grow irregularly, fo as to form pretty sharp processes, that it might excite inflammation, and lay the foundation of a fatal disease.

AND THE WAY TO PERSON AS THE P

the first of the section is the state of the section in

the real religions of the same sealest

and the case of the second designation

#### CHAP. IV.

# Diseased Appearances of the Lungs.

## Inflammation.

INFLAMMATION of the substance of the lungs, I believe, feldom takes place without some similar affection of the pleura; at least in the instances which I have feen, this has been most frequently the case. When a portion of the lungs is inflamed, its spungy structure appears much redder than usual, the colour being partly florid and partly of a darker hue. This arises from a much greater number of small vessels than usual, being distributed upon the cells of the lungs which are capable of admitting the red globules of the blood. There is also an extravalation of the coagulable lymph into the fubstance of the lungs, and sometimes of the blood. The extravalated blood has been faid upon fome occasions to be in very large quantity; but this has never fallen under my own observation.

In confequence of the greater quantity of blood being accumulated in the inflamed portion of the lungs, they become considerably heavier, and will frequently sink in water. The pleura covering the inslamed portion of the lungs has generally a similar affection; it is crowded with sine red vessels, and has generally lying upon it

a layer of coagulable lymph.

This inflamed state of the lungs is to be distinguished from blood accumulated in some part of them after death, in consequence of gravitation. From the body lying in the horizontal posture after death, blood is often accumulated at the posterior part of the lungs, giving them there a deeper colour, and rendering them In this case there will be found no heavier. crowd of fine veffels filled with blood, nor any other mark of inflammation of the pleura. Where blood too is accumulated in any part of a lung after death from gravitation, it is always of a dark colour; but where blood is accumulated from inflammation, portions of the inflamed part will appear florid.

It is very common to find abscesses formed in the lungs. These sometimes consist of small cavities containing pus, and at other times the cavities are very large, so that the greater part of the substance of the lungs has been destroyed. These cavities sometimes communicate only with branches of the trachea, which are destroyed in the pro-

gress of the ulceration; at other times they open into the cavity of the cheft, emptying their contents there, and forming the disease which is called empyema. Where abscesses are deeply seated in the substance of the lungs, the pleura is commonly not affected; but where abscesses are formed near the furface, it is almost constantly inflamed. The lungs round the boundaries of an abfcefs, when it has arisen from common inflammation, are more solid in their texture, in consequence of coagulable lymph being thrown out during the progress of the inflammation. When the abscesses are scrofulous, the texture of the lungs in the neighbourhood is fometimes not firmer than usual, but prefents the common natural appearance.-This I believe to be principally the case when the abfeeffes are small, and placed at a considerable diftance from each other. When a portion of the lungs is crowded with tubercles, and fome of these are converted into abscesses, the intermediate fubstance of the lungs is often of a very folid texture. When blood vessels are traced into an abscess of the lungs, I have found them, upon examination, very much contracted, just before they reach the abscess, so that the opening of their extremities has been closed up entirely. fuch occasions it will require a probe to be pushed with a good deal of force, in order to open

again their extremities. The late ingenious Dr. Stark\* has found in some of these vessels the blood coagulated. This change in the blood vessels is no doubt with a view to prevent large hæmorrhages from taking place, which would certainly be almost immediately fatal.

When abscesses of the lungs are the consequence of common inflammation, they are comparatively under the most favourable circumstances for recovery; but they are much more frequently the consequence of an inflammation depending on a particular constitution, viz. what is called scrosulous; and in this case they are almost always fatal.

## Tubercles.

There is no morbid appearance so common in the lungs as that of tubercles. These consist of rounded sirm white bodies, interspersed through their substance. They are I believe formed in the cellular structure, which connects the air cells of the lungs together, and are not a morbid affection of glands, as has been frequently imagined. There is no glandular structure in the cellular connecting membrane of the lungs; and on the

<sup>\*</sup> See Dr. Stark's works, p. 28.



cells in a great measure obliterated. The texture of the lungs on many occasions, however, round the boundaries of an abscess, is perfectly natural.

I have fometimes feen a number of small abscesses interspersed through the lungs, each of
which was not larger than a pea. The pus in
these is rather thicker than what arises from common inflammation, and resembles scrosulous pus.
It is probable that these abscesses have been produced by a number of small scattered tubercles
taking on the process of suppuration. The lungs
immediately surrounding these abscesses are often
of a persectly healthy structure, none of the cells
being closed up by adhesions.

When tubercles are converted into abfceffes, it forms one of the most destructive diseases in this island, viz. pthis pulmonalis. Tubercles are sometimes found in the lungs of children at a very early age, viz. two or three years old, but they most frequently occur before the completion of the growth. They are apt likewise to be formed at rather an advanced age.

In cutting into the lungs, a confiderable portion of their structure sometimes appears to be changed into a whitish soft matter, somewhat intermediate between a solid and a sluid, like a scrosulous gland just beginning to suppurate. This appearance I believe is produced by scrosulous

matter being deposited in the cellular substance of a certain portion of the lungs, and advancing towards suppuration. It seems to be the same matter with that of the tubercle, but only diffused uniformly over a considerable portion of the lungs, while the tubercle is circumscribed.

I have feen another fort of tubercle in the lungs, which I believe to be very rare. It confifts of a fost tumour, formed of a light-brown smooth substance. This is not contained in any proper capsule, but adheres immediately to the common structure of the lungs. In cutting through several of these tumours I did not find any of them in a state of suppuration. They were commonly as large as a gooseberry, and were chiefly placed upon the surface of the lungs; some, however, were feattered through their substance, of a smaller size. These are very different in their appearance from the common tubercle last described, and are the effect of a diseased process, which probably is very impersectly known.

In opening into the cheft, it is not unufual to find that the lungs do not collapse, but that they fill up the cavity completely on each side of the heart. When examined, their cells appear full of air, so that there is seen upon the surface a prodigious number of small white vesicles. The branches of the trachea are at the same time much

filled with a mucous fluid. This is not uncommonly the case in persons who have laboured for some considerable time with difficulty of breathing, but without any symptoms of inslammation; such persons would appear to die for want of supply of atmospheric air sufficient to produce the proper change in the blood which is necessary for its useful circulation through the body.

The lungs are fometimes, although I believe very rarely, formed into pretty large cells, fo as to refemble fomewhat the lungs of an amphibious animal. These cells in the only instance which I have seen of this disease, were most of them of the size of a common garden pea, and some sew were so large as to be able to contain a small gooseberry. They were surrounded by a sine transparent capsule, and were so numerous as to occupy more than one half of the portion of the lung which I saw. The only specimen of this fort of disease which I am acquainted with, is in the collection of Mr. Cruik-shank; and the person in whom it was found, had been very long subject to difficulty of breathing.

The lungs are fometimes converted into a folid fubstance very much resembling the liver. It has nearly the same folidity, and the same general appearance. When examined more minutely, the air cells appear to be filled with a brownish solid matter. It is very obvious, that as this process

foreads over the lungs, respiration must become more difficult; there will be less room for the admission of air, in order to produce that change upon the blood which is necessary for the preservation of life, and this circumstance will produce a more frequent respiration, in order to endeavour to remedy the desiciency.

Part of the lungs is occasionally converted into a bony fubstance, but this is a very rare disease. The fmall veffels disposed through the substance of the lungs under fuch circumstances separate bony matter from the blood. In the only instance which I have known of this complaint, the process would appear to have been rapid. There was great difficulty of breathing before the person died, but this difficulty had only begun for a very few weeks. Each lung was undergoing the fame procefs, which had made cousiderable advancement. In the particular case to which I allude, there had been a very strong disposition to form bone in the constitution. A very large bony tumour had been formed round one of the knees of this person; and very foon after the knee and leg were removed by amputation, the difficulty of breathing began, which was occasioned by a part of the lungs being converted into bone. Here was a transference of this peculiar disease from an ex-

It is very obvious, that as this process

ternal to an internal part, fimilar to the translation of gout or rheumatism.

I have also seen a tumour as large as an orange, attached to the lungs on one side by a loose membranous connection, and in some degree compressing them; this tumour consisted of a yellowish, porous substance, which neither resembled the structure of what is commonly understood to be a schirrous or scrosulous tumour, but had an appearance somewhat peculiar to itself. It was probably the effect of a morbid action with which we are very little acquainted.

Earthy concretions have occasionally been found in the lungs, although it is a rare appearance of disease. These are often small, but sometimes form pretty large masses.\* Even a considerable portion of the lungs has been known to be changed into an earthy substance.†

Hydatids are also formed in the lungs, and are many of them brought up by coughing. Of their nature I propose to treat particularly afterwards.

<sup>\*</sup> Vid. Morgagni, Epist. XVII. Art. 19. Epist. XV. Art. 25.

<sup>†</sup> Vid. Morgagni, Epist. XXII. Art. 15.

### CHAP. V.

Diseased Appearances in the Posterior Mediastinum.

By the posterior mediastinum, is meant that space which lies between the laminæ of the pleura, that pass from the root of the lungs to each side of the spine. The space is of considerable size, and contains a portion of the trachea arteria, of the æsophagus, of the thoracic dust, of the descending aorta, and the vena azygos, besides some absorbent glands.

Discased Appearances of the Trachea.

The inner membrane of the trachea is not uncommonly inflamed to a greater or lefs degree. In this state it is crouded with minute slorid vessels, which give it a general appearance of vascularity. When there is no inflammation, it appears a white pulpy membrane, but there are rarely to be seen any red vessels ramifying through it. While the inner membrane is inflamed, the secretion from

its glands is very much increased, and therefore its cavity is found a good deal filled with a mucous fluid; even pus is sometimes formed, and both fluids are mixed with globules of air. This is probably the situation of the trachea in a very violent catarrh, and also in some cases where there are scrofulous abscesses of the lungs, attended with hoarseness, and a sense of soreness along the trachea.

When the inner membrane of the trachea is inflamed, it is fometimes lined with a layer of a yellowish pulpy matter. This does not adhere very firmly to the inner membrane, but may be easily separated. It extends from the upper part of the cavity of the larynx, into the small branches of the trachea, which are distributed through the substance of the lungs. There is at the same time a good deal of mucus in the trachea and its branches, together with a mixture of pus. This is the appearance of the inside of the trachea, in patients who have died from the croup.

## Polypus.

The trachea and its branches are fometimes lined with a layer of a yellowish matter, forming

a fort of tube, which is applied to the inner furface loofely. It has not occurred to me to fee any instance of it in the dead bodies which I have examined, but I have feen feveral examples of it in preparations. The inner membrane of the trachea feems to be perfectly natural, and the layer of adventitious membrane refembles exactly the coagulable lymph, that is thrown out in other parts of the body: I have therefore no doubt of its being that fubstance. Since this disease (which is called polypus) lasts for a long time, and is not attended with fymptoms of inflammation, it appears probable, that the vessels of the inner membrane of the trachea poffess a power of separating the coagulable lymph from the blood, and that this disease confists in a peculiar action of these vessels.

The trachea has been faid to be fometimes filled with a folid fubstance of the same kind with what we have described; but in the instances which I have seen, it has been tubular; and I believe this to be by much the most common appearance.

The trachea I have seen narrowed in diameter for two or three inches, thickened in its substance, and the inner membrane has been raised into a number of little hard tubercles. This state of the trachea was accompanied with a schirrous affection of some absorbent glands, which closely adhered to it; and it appeared to me, that the difease in the glands had spread so as to affect the trachea.

The cartilaginous rings of the trachea occasionally become offified, although this is not a very frequent appearance. The change is so natural from cartilage into bone, that we should be led to expect it more commonly. When the offification is inconfiderable, the function of the trachea will hardly be affected, but where the rings are entirely offified, the flexibility of the trachea must be much lessened, and its cavity will not admit of being fo much contracted as in the healthy state, by the action of the muscular fibres, which form a part of its structure. The consequence of this must be, that the mucus, which is occasionally accumulated, will not be fo readily expelled by coughing, and the air will not be thrown out in fo fmall a column, nor with fo much momentum.

The trachea is no doubt liable to be destroyed in part by ulceration, from causes acting immediately upon itself, but in the instances which I have feen, the ulceration has been connected with ulceration of the cefophagus. As the cefophagus is more liable to this difeafe, it is probable that in fuch cases the ulceration has begun in the œfophagus, and fpread to the trachea.

## Diseased Appearances of the Oesophagus

The cefophagus is frequently lined with a layer of the coagulable lymph, which is continued from the cavity of the mouth.

This, it is faid, fometimes extends over the whole intestinal canal, but I should believe this appearance to be extremely rare, and it commonly terminates at the lower end of the The inner membrane of the œsophagus. mouth is much more vascular than in its natural state, shewing an intense dark red colour, but in examinations after death the appearance of greater vascularity is sometimes little observable in the cefophagus. This difease is known under the name of apthæ, and is much more often to be observed in the living than the dead body. Portions of the coagulable lymph are thrown off, and other portions formed, feveral times in the progress of the disease.

The cesophagus is liable to stricture, produced by the contraction of its muscular fibres at some particular part. This disease is I think most common in women whose constitutions are delicate, and much subject to nervous influence. When such a disease is examined in the dead

body, the æsophagus is sound to be more or less contracted in some part of it, and it seels harder than usual, as all muscles do in a contracted state. There is no appearance of diseased structure usually combined with it. I can suppose, however, that this contraction might lay the soundation of a permanent, and even a statal disease. The muscular sibres of the æsophagus might so press on the inner membrane, as to excite inflammation in it, which might advance to suppuration, and would most probably terminate statally.

I once faw a very unufual stricture of the copphagus. It consisted in its inner membrane being puckered together, so as to form a narrowness of the canal at a particular part. The canal at that part, was so narrow, as hardly to allow a common garden pea to pass. There was no appearance, however, of diseased structure in the inner membrane which was so contracted, and the muscular part of the cesophagus surrounding it was perfectly sound. I know that this disease was very slow in its progress, for the person in whom it took place had been for many years affected with a difficulty of swallowing, and could only swallow substances of extremely small size.

The most common appearance of disease in the cesophagus, is that of an ulcer in its cavity.

Ulcers of the cefophagus are sometimes of a common nature, but most frequently they are attended with a schirrous affection. When ulcers of the cefophagus arife from common inflammation, the structure of the cosophagus, immediately furrounding the ulcer, is little thickened, and there is the appearance of the usual erosion in ulcers. When the ulcer is of a fchirrous nature, the œsophagus in the neighbourhood is very much thickened, and is very hard in its texture. When this texture is examined, it either confifts of an hard, uniform, fleshy substance, or is a little interfected by a membranous appearance, or it is griftly. Under fuch circumstances, the canal of the æsophagus is always more or less narrowed, and in some cases is almost wholly obliterated. It is worthy of remark, that these ulcers happen most frequently, either immediately under the pharynx, or near the cardia.

Any fubstance capable of irritating the inner membrane of the œsophagus, by having sharp hard projections, will no doubt be more likely to essect the œsophagus, where it sirst enters into it. In an œsophagus, therefore, predisposed to schirrous, such an accident may prove an exciting cause, and the disease will more frequently take place at its upper end. At the cardia too,

there is a peculiar arrangement of the mufcular fibres, which are capable of acting in some degree like a fphincter, and which probably produce on many occasions a narrowness of canal there. This will render the cefophagus at the cardia more liable to be injured by the passage of any hard substance, and may ultimately lay the foundation of a schirrous ulcer. This is the account which the late Dr. Hunter used to give of the frequent situation of ulcers at the upper and lower extremities of the æfophagus, and feems to have great weight. It happens, however, most commonly that ulcers of the cofophagus arife spontaneously, or in other words, from causes within itself which we cannot ascertain. When an ulcer takes place at the upper end of the œfophagus, it is apt to spread into the fubstance of the thyroid gland. In this case the gland becomes hard, enlarged, and ulcerated.

A portion of the cefophagus has been observed by some anatomists to be converted into cartilage, and to have its diameter at that part very much diminished in size.\* This was, perhaps, only a strong example of the gristly texture which we have above described, or it may have been a change into a substance like commmon cartilage.

<sup>\*</sup> Vid. Bonet, Tom. 2. p. 32.

In this case it is a very uncommon appearance of disease.

I have feen an instance of a fungus arising on the inside of the œsophagus, which is to be considered as a rare disease. When cut into, it appeared to have a fibrous structure, disposed in some measure at right angles to the inner membrane upon which it was formed, and was ulcerated on its surface.

The pharynx, at its lower extremity, has been known to be dilated into a pouch of confiderable fize, which paffed behind the œfophagus. This may be supposed to be very rare, but there is an instance of it preserved in Dr. Hunter's collection. The pouch in this cafe began to be formed in confequence of a cherrystone having rested there for some time, which had made a kind of bed for itself. It remained in that fituation for three days, and then was brought up by a violent fit of coughing. A part of the food always rested afterwards in the cavity made by the cherry-stone, by which it was gradually enlarged. At length, in the course of about five years, the cavity was enlarged into a bag of a confiderable fize, fufficient to contain feveral ounces of fluid. This bag paffed down a good way behind the cefophagus, and the œfophagus necessarily acquired a val-



more readily divisible into layers. There are also frequently deposited behind the inner membrane little thin laminæ of bony matter. This appearance of disease has been formerly explained more particularly. It is rare that this part of the aorta becomes aneurismal, unless there be a general aneurismal affection over the arterial system.

# The Vena Azygos.

The vena azygos is very feldom diseased. I have seen it however varicose, and very much enlarged. This change in it took place from particular circumstances. A considerable portion of the vena cava inserior had become obliterated; in consequence of this, the usual vena azygos, together with an uncommon one on the lest side, were the only channels through which the blood could return by a circuitous route to the heart; they were therefore necessarily, from the impetus of the blood, much enlarged in size, and for the same reason likewise varicose. This case I have more particularly described in the Medical and Chirurgical Transactions.\*

The vena azygos has been known to be ruptured, when very much diffended with blood.†—Such a cafe has not come under my own observation, and I should believe it to be very uncommon.

### The Thoracic Duct.

The thoracic duct is also subject to very few diseases. I have never seen any other, except that of its being very much enlarged beyond its natural size, and varicose.

In the instance to which I allude, it was very nearly as large as the natural side of the subclavian vein, but nothing could be detected in the neighbouring parts, capable of accounting for this appearance. There was no obstruction at the entrance of the thoracic duct into the venal system, which might naturally have been expected. This diseased appearance of the thoracic duct has already been taken notice of in Mr. Cruikshank's Treatise on the Absorbent System.\*

The thoracic duct has been known to be obfiructed by an earthy matter deposited in its cav-

<sup>+</sup> Vid. Morgagni, Epist. XXVI, Art. 29.

<sup>\*</sup> See fecond edition, p. 207; and is represented in an engraving, plate V.

ity.† It does not necessarily happen, when the thoracic duct is at some part obstructed, that chyle is prevented from entering into the system of blood vessels. The thoracic duct not unsrequently sends off one or more considerable branches, which unite again with the principal trunk. If under such circumstances an obstruction should take place in a part of the principal trunk, between the origin and the termination of those branches, no bad essel would follow; one or more of those branches would become enlarged and convey the chyle in its full quantity to the blood.

The thoracic duct has also been known to be ruptured, although this is exceedingly rare.

## Absorbent Glands.

The absorbent glands in the posterior mediastinum, as well as in everyother part of the body, are liable to several diseases. The most common morbid affection is scrosula. In this case they are frequently a good deal enlarged, and feel somewhat softer to the touch than in their healthy structure. When cut into, however, they frequently exhibit very much the natural

+ Vid. Lieutaud. Tom. 2, p. 93.

appearances; but it is more common to find that fome of them contain a white, foft, cheefy matter, mixed with a thick pus; this is the most decided mark of scrofulous affection. When the absorbent glands in this situation are very much enlarged, they necessarily produce some difficulty of breathing, both by the pressing on the lungs and the trachea. They might produce also some difficulty of swallowing.

I have seen the absorbent glands in the neighbourhood of the trachea affected with schirrus, although it is a rare disease in them. They were much enlarged, and very hard to the touch.—When cut into, they exhibited a hard brownish structure, somewhat intersected by white sirm membrane, so as to resemble exactly what is called schirrus in other parts of the body. The trachea in contact with these glands was also affected. In this case the thyroid gland was schirrous, and it is probable that the disease spread from the thyroid to the absorbent glands, and so to the trachea.

The absorbent glands near the trachea are sometimes converted into a bony or earthy matter, from a peculiar secretory action in their blood vessels; and I think that this disease is more common in the absorbent glands at the root of the trachea, than in any other part of the body.

These glands when so diseased, by pressing against the trachea or œsophagus, occasionally produce ulcers in them.

## The Anterior Mediastinum.

By the anterior mediastinum is meant the space inclosed between the laminæ of the pleura, which pass from the sternum to the pericardium; it contains little else than cellular membrane, with perhaps a small portion of sat; and in the younger subject, the thymus gland.

It is feldom found with any difeafed appearances in it. Abfcesses are occasionally formed there, but rarely. Water too is sometimes found in the cells of its cellular membrane. I have also seen air accumulated in these cells

Fat is occasionally deposited in the mediastinum in considerable quantity. When the quantity is very large, it has been known to disturb a good deal the functions both of the heart and lungs. The thymus gland I have known to be much enlarged beyond its natural size, and to have a schirrous hardness.

#### CHAP. VI.

Diseased Appearances within the Cavity of the Abdomen.

### Ascites.

Ascites, or dropfy of the cavity of the abdomen, is a very frequent disease, and is not confined to any fex or age. I have feen feveral inflances of it in children under ten years old; but it is much more common at the middle, and the more advanced periods of life. It is also more common in the male than the female fex. When water is accumulated in a very confiderable quantity in the cavity of the abdomen, the fuperficial veins of the belly are generally a good deal distended with blood; this most probably arifes from the pressure of the water upon the deeper feated veins; it is however fometimes hardly observable, even when the accumulation of the water is very considerable. The skin at the navel is also often protruded, yielding easily

to pressure, but this is not universally the case. On many occasions the protrusion can hardly be feen, when the water is accumulated in large quantity. In opening into the cavity of the abdomen, there is to be feen a larger or less quantity of an aqueous fluid, generally of a brownish colour, but its colour varies according to circumstances. When there is a schirrous liver accompanying the dropfy, the water is commonly of a yellowish or greenish colour. This arises from a mixture of the bile with the water, and under fuch circumstances there is almost always a jaundiced colour of fkin. I have feen the water in afcites of a chocolate or coffee colour; but this appearance is rare. In the case to which I allude, the water was thicker than that of afcites usually is, but it had the common properties, as far as could be known from the application of heat or of acids. When there are none of the vifcera of the abdomen difeafed, the water in afcites refembles the ferum of the blood in its colour, as well as in its other properties.

While water is accumulated in the cavity of the abdomen, the intestinal canal is frequently found to be somewhat in a contracted state; but often too this is not observable. In many cases of ascites the liver is diseased, being hard and tuberculated, as we shall explain particularly when



florid blood. When a portion of the inflamed peritonæum is feparated from the abdominal mufcles, there is commonly no appearance whatever of the inflammation having fpread into the mufcles; but where the peritonæum covers the intestinal canal, the inflammation is sometimes found to have penetrated not only into the muscular coat of the intestines, but even into the villous membrane. The reason of this difference probably is, that the peritonæum is less connected with the abdominal muscles than with the intestinal canal, so that the inflammation is more easily confined in the one case than in the other.

The inflammation of the peritonæum is fometimes slight and partial; at other times is great and universal. When it is slight and affects that part of the peritonæum which is connected with the intestinal canal, it often forms broad bands of inflammation which run along the course of the intestines, and are bounded by the contact of the different portions of the intestines among themfelves. In this case, the coats of the intestine are not thicker than usual, the inflammation being slight and confined to the peritonæum itself. Where the inflammation is great, the intestines are much thicker and more massy. This evidently arises from the greater accumulation of blood travalation of fluids into the fubstance of the intestines, in consequence of the strong inslammatory action of the vessels. The mesentery and mesocolon are much thicker than in their natural state, and there is also a remarkable change in the omentum. It is frequently as thick as a person's hand, and lies as a circumscribed mass along the great curviture of the stomach. The principal cause of this change in these parts, is the extravalation of coagulable lymph into the cellular membrane, between the laminæ of the peritonæum which form them.

In many places there is thrown out a layer of a yellowish pulpy matter, gluing different portions of the viscera together. This is sometimes a thin layer, at other times is of considerable thickness; and it appears to be the coagulable lymph of the blood. The layer of the coagulable lymph is often found to extend beyond the immediate surface which is inflamed. This arises from the coagulable lymph being thrown out from the vessels in a sluid state, and therefore need not be limitted by the boundaries of the inflamed surface when it becomes solid. There is also a considerable quantity of brownish sluid in the cavity of the abdomen, resembling the serum, which is mixed with small shreds of the coagulable lymph, and sometimes

with pus, giving it a turbid appearance. The quantity of the coagulable lymph, and of the fluid, is fometimes large, in proportion to the degree of the inflammation. Air is frequently too found accumulated in the stomach and the intestinal canal, which had been formed in the progress of the disease. At other times this air is wanting.

Adhesions in the Cavity of the Abdomen.

When there has been inflammations of the peritonæum, either generally or partially, fufficient to have thrown out coagulable lymph, and the patient has furvived the disease, the coagulable lymph is changed into a fine transparent membrane, which is the membrane of adhesions. The time, which is occupied in the change of the coagulable lymph into the membrane of adhesions, is not very long, for I have had feveral opportunities of tracing the gradual progress of the change from the one into the other, while the inflammation appeared to have been recent. This membrane confifts of a cellular fubstance, similar to the general cellular membrane of the body, and has a moderate share of vafcularity. It does not naturally shew many veffels large enough to admit the red globules of the blood, but it shews its vascularity upon slight

degrees of inflammation, or by using the fine injection. This membrane is capable of elongating gradually by the motion of the viscera upon themselves, so as ultimately to be attended with very little inconvenience. I have very often had an opportunity of observing these adhesions, either joining all the viscera of the abdomen more or less together, or joining some particular viscera to each other.

I have observed also pus formed in an inslamed state of the peritonæum; but I have never seen this membrane mortissed independent of the mortiscation of some of the viscera.

I have feveral times had an opportunity of obferving a white, foft, granulated matter, adhering
universally behind the peritonæum. In some
places it formed a mass of considerable thickness;
in others it was scattered in single small masses.
In one case I recollect it formed a substance as
thick as my hand, between the peritonæum and
the abdominal muscles, while it was scattered in
small separate portions in the mesentery and the
peritonæum, covering the intestinal canal. The
omentum I have sometimes seen changed into a
cake of this substance. The matter itself appeared
to me to be scrosulous, for it resembled exactly
the structure of a scrosulous absorbent gland before pus is actually formed. I am not at all cer-

tain how far this appearance of disease should have been classed along with those of the peritonæum: it does not take place (at least in the cases which I have seen) in the peritonæum itself, but behind it, yet at the same time adhering to it. It appears, however, upon the whole, to pe placed here with more propriety than it could have been any where else.

I have also seen some small cancerous tumours growing from the peritonæum. These were extremely hard, of a white colour, and resembled exactly in their structure the cancerous masses which are formed in the stomach. What puts the appearance I allude to beyond doubt, is, that in the same body I found a cancerous tumour of the stomach. The cancerous tumours of the peritonæum were not at all connected with this other, but were in that part of the membrane which lines the recti abdominis muscles, nearly opposite to the region of the stomach.

Hydatids have occasionally been found to occupy a portion, or even the whole, of the cavity of the abdomen. In such cases many of them are connected with the viscera, which were probably the basis of their formation, of which the chief are the liver, and the spleen. This appearance of disease is very uncommon.

Air has been occasionally said to be accumulated in the cavity of the abdomen, while little or none is contained in the intestines.\* This I believe to be a very rare occurrence. Air is not unfrequently sound accumulated in considerable quantity in the intestinal canal, while there is none at all in the cavity of the abdomen.

\* Vid. Lieutaud, Tom. 1, p. 432.

### CHAP. VII.

Diseased Appearances of the Stomach.

## Inflammation.

It fometimes happens, although not very frequently (unless poisons have been swallowed) that inflammation takes place in the stomach, so as to fpread over a very confiderable portion of its inner membrane, or perhaps the whole of it. It is much more common, however, for inflammation to occupy a fmaller portion of the stomack. In fuch cases too, the inflammation is generally not very violent. The stomach upon the outside, at the inflamed part, shews a greater number of finall vessels than usual, but frequently not much crowded. In opening into the stomach, it is found to be a little thicker at the inflamed part, the inner membrane is very red from the number of finall florid veffels, and there are frequently spots of extravafated blood. It does not often occur that

a common inflammation of the stomach proceeds to form pus, or to terminate in gangrene.

When arfenic has been fwallowed (which is the poison most frequently taken) the stomach is affected with a most intense degree of inflammation. Its fubstance becomes thicker, and in opening into its cavity there is a very great degree of redness in the inner membrane, arising partly from the very great number of minute vessels, and partly from extravafated blood. Portions of the inner membrane are fometimes destroyed, from the violent action that has taken place in confequence of the immediate application of the poifon. I have also seen a thin layer of the coagulable lymph thrown out upon a portion of the inner furface of the stomach. Most commonly too some part of the arfenic is to be seen in the form of a white powder, lying upon different portions of the inner membrane.

In opening the bodies of perfons who have died from hydrophobia, the inner membrane of the stomach is frequently found inslamed at the cardia, and its great end. This inslammation, however, is generally inconsiderable, is, I believe, never very great, and in some instances has been faid to be wanting.

## Ulcers of the Stomach.

Opportunities occasionally offer themselves of observing ulcers of the stomach. These fometimes refemble common ulcers in any other part of the body, but frequently they have a peculiar appearance. Many of them are hardly furrounded with any inflammation, nor have they irregular eroded edges as ulcers have generally, nor is there any particular diseased alteration in the structure of the stomach in the neighbourhood. They appear very much as if some little time before a part had been cut out from the stomach with a knife, and the edges had healed, so as to prefent an uniform fmooth boundary round the excavation which had been made. ulcers fometimes destroy only a portion of the coats of the stomach at some one part, and at other times destroy them entirely. When a portion of the coats is destroyed entirely, there is fometimes a thin appearance of the stomach surrounding the hole, which has a smooth surface, and depends on the progress of the ulceration. At other times the stomach is a little thickened furrounding the

hole; and at other times still, it seems to have the common natural structure.

Schirrus and Cancer of the Stomach.

This affection of the stomach is not very uncommon towards an advanced period of life, and, I think, is more frequently met with in men than in women. This, perhaps, arises from the greater intemperance in the one fex, than in the other. It cannot, however, be produceed entirely by intemperance; there must be added a considerable predisposition of the parts towards this disease. Hence, when there is no previous disposition, the stomach does not become affected with this disease, whatever be the intemperance. When, however, there is this previous disposition, there is reason to think that it is encouraged and brought forwards by this mode of living.

Schirrus fometimes extends over almost every part of the stomach, but most commonly it attacks one part. The part which is affected with schirrus has sometimes no very distinct limit between it and the sound structure of the stomach, but most commonly the limit is very well marked. When schirrus attacks a portion of the stomach only,

it is generally towards the pylorus. The principal reason of this probably is, that there is more of glandular structure in that part of the stomach than any other; and it would appear that glandular parts of the body are more liable to be affected with schirrus, than parts of the body generally.

When the whole stomach, or a portion of it, is schirrous, it is much thicker than usual, as well as much harder. When the difeafed part is cut into, the original structure of the stomach is frequently marked with fufficient distinctness, but very much altered from the natural appearance. The peritonæal covering of the stomach is many times thicker than it ought to be, and has almost a griftly hardness. The muscular part is also very much thickened, and is interfected by frequent pretty strong membranous septa. These membranous fepta are, probably, nothing elfe than the cellular membrane intervening between the fasciculi of the muscular fibres, thickened from disease. The inner membrane is also extremely thick and hard, and not unfrequently fomewhat tuberculated towards the cavity of the stomach.

It very frequently happens that this thickened mass is ulcerated upon its surface, and then a stomach is said to be cancerous. Sometimes the inner membrane of the stomach throws out a process which terminates in a great many smaller processes, and produces what has been commonly called a fungous appearance.

It also happens that the stomach at some part loses entirely all vestige of the natural structure, and is changed into a very hard mass, of a whit-ish or brownish colour, with some appearance of membrane intersecting it; or it is converted into a gristly substance, like cartilage somewhat softened. The absorbent glands in the neighbourhood are at the same time commonly enlarged, and have a very hard white structure.

I have feen feveral inflances of a schirrous tumour being formed in the stomach about the
fize of a walnut, while every other part of it was
healthy. This tumour has most frequently a
small depression near the middle of its surface,
and appears a little radiated in its structure.—
While this tumour remains free from irritation,
the functions of the stomach are probably very
little affected by it; when, however, it is irritated, it must occasion very considerable disorder
in the functions of the stomach, and perhaps lay
the foundation of a fatal disease.

A part of the stomach is occasionally formed into a pouch by mechanical means, although very rarely. I have seen one instance of a pouch being so formed, in which sive halfpence

had been lodged. The coats of the stomach were thinner at that part, but were not inslamed or ulcerated. The halfpence had remained there for some considerable time, forming a pouch by their pressure, but had not irritated the stomach in such a manner as to produce inslammation or ulceration.

The orifice of the stomach may be almost, or perhaps entirely, shut up by a permanent contraction of its muscular fibres, either at the cardia or the pylorus. It is much more likely, however, to occur most frequently at the pylorus, from natural circumstances: there is both less opposition to prevent the contraction at this orifice of the stomach than at the cardia, and there is also belonging to it a stronger and more direct circular muscular power. A less contraction too, at the pylorus, will produce an obstruction in the canal, than at the cardia. I have feen one instance of this contraction at the pylorus, which, even there, is a very rare disease. The contraction was fo great as hardly to admit a common goose quill to pass from the stomach into the duodenum, and it had prevented a number of plum stones from passing, which were therefore detained in the stomach.

The stomach is sometimes found so contracted through the whole of its extent as not to be

larger than a portion of the small intestine; at other times, it is enlarged to much more than its ordinary size. Neither of these appearances are to be considered as arising from disease. They depend entirely on the muscular sibres of the stomach being in a state of contraction or relaxation, at the time of death. It happens, I think, more frequently that the stomach is dilated than contracted.

The stomach is very commonly found, in a dead body, flaccid and almost empty, but not unfrequently it is found more or less distended with air: this air may have been formed after death, but it is often formed during life. When this is the case, we may suppose it to be produced by a new chemical arrangement in the contents of the stomach; but, I believe, it more frequently happens that air is separated from the blood in the blood vessels of the stomach, and poured by the fmall exhalants into its cavity. This has been more particularly taken notice of by Mr. Hunter, in his Essay upon Digestion,\* and by myfelf, in a paper which is published in the Medical and Chirurgical Transactions.+

<sup>\*</sup> See Mr. Hunter's Observations on certain parts of the animal economy, p. 164.

<sup>+</sup> See case of emphysema, p. 202.

In looking upon the coats of the stomach at its great end, a finall portion of them there appears frequently to be thinner, more transparent, and feels fomewhat more pulpy than is ufual, but those appearances are feldom very strongly marked. They arise from the action of the gastric juice resting on that part of the stomach in greater quantity than any where elfe, and diffolving a fmall portion of its coats. This is therefore not to be confidered as the confequence of a difeafe, but as a natural effect, arifing from the action of the gastric juice, and the state of the stomach after death. When the gastric juice has been in confiderable quantity, and of an active nature, the stomach has been dissolved quite through its subflance at the great end, and its contents have been effused into the general cavity of the abdomen. In fuch cases the neighbouring viscera are also partially diffolved. The inflances, however, of fo powerful a folution are rare, and have almost only occurred in perfons who, while in good health, had died fuddenly from accident. The true explanation of these appearances was first given by Mr. Hunter, and published, at the request of Sir John Pringle, in the Philosophical Transactions.\*

<sup>\*</sup> See Philosoph. Transact. Vol. 62, p. 447.

Tumours, confisting of a suetty substance, have been sometimes sound in the stomach, but they are to be considered as a very rare appearance of disease. Ruysch relates, that he has seen a tumour from the stomach of a man which contained hair, together with some dentes molares; and this he has preserved in his collection.†

Calculi with different appearances have been described as being occasionally sound in the stomach. They have never come under my own observation, and are to be reckoned very uncommon.‡

Papillæ and puftules, fomewhat refembling the fmall-pox, have also been described as being formed on the inner membrane of the stomach, but these are exceedingly rare.

Even true fmall-pox pustules have been said to be formed in the stomachs of persons who have died from this disease. In later dissections, however, this appearance has not been observed, so that, at least, it must be considered as very uncommon.

<sup>+</sup> Vid. Ruysch, Tom. 2. Adversar. Anatomicor. Decad. Tert.

<sup>‡</sup> Vid. Lieutaud, Tom. 1, p. 17.

<sup>§</sup> Vid. Lientaud, Tom. 1, p. 23.

<sup>||</sup> Vid. Lieutaud, Tom. 1, p. 371.

### CHAP. VIII.

Difeased Appearances in the Intestines.

### Inflammation.

I HE intestinal canal is subject to inflammation from a variety of causes, and therefore it is very usual to observe its effects after death. When a portion of the intestine is inflamed, there is spread upon its outer furface a number of fmall vessels, many of which are carrying a florid blood. When the intestine is cut into, so as to exhibit its inner membrane, it appears highly vascular, from the fmall veffels of the villi being loaded with blood, and there are frequently to be feen a few spots of blood extravafated. In inflammation of the inteftines, the peritonæum is often very little, or not at all affected. When the inflammation, however, is very great, the peritonæum is also inflamed, and covered with a layer of coagulable lymph. I have likewise seen, in violent inflammation, scatupon the furface of the villous membrane. The intestine is at the same time much more thick and massy than in a healthy state, and its colour is sometimes very dark, from a large quantity of black extravasated blood. This state of an intestine has often been mistaken for mortification.

It very commonly happens that inflammation of the intestines advances to suppuration and ulceration. Ulceration, however, does not appear to be fo common in the small as in the great intestines. When it takes place either in the one or the other intestine, it is attended with considerable variety in its appearance: the edges of the ulcer have fometimes confiderable thickness; at other times, they are not thicker than the healthy structure of the intestine; the edges and general cavity of an ulcer are fometimes ragged, and at other times they are fmooth, as if a portion had been cut out from the intestine with a knife. Sometimes, through a confiderable length of the intestine (especially the great one) the inner membrane hangs in tattered shreds, occasioned by the great ravage of the ulceration. I have also seen, a confiderable portion of the intestine completely stripped of its inner membrane, from the extent, of this process, and its muscular coat appeared as distinct as if it had been very carefully dissected.

In the follicular glands, which are gathered together in little oval groups, I think I have feen ulceration more frequently than in the other parts of the intestine.

When ulceration advances very actively, it fometimes eats through the coats of an intestine entirely. When this is the case, a portion of the contents of the intestine occasionally passes into the general cavity of the abdomen, producing there inflammation. This, however, does not very often happen; most commonly that portion of the gut where the ulcer is fituated, adheres by inflammation to some other portion, or to a neighbouring vifcus, and a communication is formed between the one and the other. I have feen communications formed in this manner between the rectum and the bladder in a male, and between the rectum and the vagina in a female. I have even feen a communication formed between the kidney and a portion of the intestine from this cause, by which the pus produced in the kidney was evacuated through the intestine. Those communications are the means of preferving life (although in a very uncomfortable state) for a much longer time than it could be, were the mattar to pass into the general cavity of the belly. It would there produce peritonæal inflammation, which would very foon deftroy.

Inflammation of the intestines sometimes, although rarely, advances to mortification. When this is the case, the mortified part is of a dark livid colour, and has lost its tenacity; it is in this state very easily torn through, or the fingers will pass through it as through a rotten pear. The want of the natural tenacity, when attended with the other circumstances which we have mentioned, is the only fure criterion of a part being mortified, in examinations after death. A portion of inteftine may be of a very dark colour, and yet may not be mortified. This darkness of colour may be occasioned by a large quantity of extravalated blood thrown out during a high degree of inflammation, where the principle of life is maintained in full vigour. Thus we fee blood effused into the cellular membrane under the fkin producing a very dark appearance, yet the parts are all alive. It has often happened too, that a very dark portion of intestine has been returned in the operation for the bubonocele, and yet the parts have recovered their natural functions. This could never have happened if the black portion of the intestine had really been mortified. Under such circumstances, the mortified part would have separated from the living, and the function of the gut must have been destroyed. When a portion of gut has been for fome time mortified, there is formed a considerable quantity of air, which is accumulated in its cavity. This is a part of the matural process which takes place in all dead animal substances.

### Intus Susceptio.

This is a difease which is not very uncommon, and which is frequently fatal. It confifts in a portion of gut paffing for some length within another portion, and there also enters along with it a part of the mesentery. The portion of gut which is received into the other, is found generally in a contracted state, and is fometimes of confiderable length. It usually happens that an upper portion of intestine falls into a lower; but the contrary likewife occurs, although rarely. Intus fusceptio may take place in any part of the intestinal canal, but it happens most frequently in the fmall intestine, and also where the ilium terminates in the colon. In this last situation, it appears to me to happen more frequently than any where elfe. This, perhaps, depends on the great disproportion in bulk between these two portions of intestine. In opening into the bodies, particularly of infants, an intus susception

is not unfrequently found, which had been attended with no mischies: the parts appear perfectly free from inflammation, and they would probably have been easily disentangled from each other by their natural peristaltic motion. At other times, however, violent inflammation takes place, the parts are thickened and glued together by adhesions, and the passage of the intestines is obstructed. This is the satal state of the disease.

#### Ruptures.

A portion of the viscera of the abdomen frequently passes out of that cavity, being lodged in a bag of elongated peritonæum, and this disease is called a rupture. It happens most commonly from some sudden and violent concussion of the body, where the weaker parts of the parietes of the abdomen give way; it is a more frequent disease too, I think, among fat than lean people: this depends on the increased size of the viscera, which therefore press with more force against the sides of the abdomen. There is hardly any viscus which has not, at some time or other, been found in the sack of a rupture, but most frequently it is either a portion of the omentum, or of the

intestine. The bag formed by the elongated peritonæum may be thrust out almost at any part of the belly, but this happens more frequently at the ring of the external oblique muscle, under Paupart's ligament, and at the naval: it also, sometimes, takes place at other parts of the abdomen. The most usual fituation of a rupture in the male, is at the ring of the external oblique muscle; and this arifes probably from the larger fize of that opening in the male, than in the female. The most usual situation of a rupture in the semale is either under Paupart's ligament, or at the navel. The reason of the frequency of the first situation, is the particular shape of the pelvis in the female, by which there is a larger empty space under Paupart's ligament, than in the male, fo that the vifcera here are less firmly supported. The reason why the fecond fituation of a rupture occurs often in the female, is, probably, frequent child-bearing. During pregnancy, at its advanced period, the navel opens, or gives way, and where pregnancies have been frequent, it probably never recovers its original strength.

The vifcus which is most commonly found in the fack of a rupture, is the omentum. This, perhaps, arises from its being a loose mass, not tied down to any particular situation, and therefore it readily passes into any cavity which communicates with the abdomen. When it has once fallen down, it has no means of pulling itself out, like a portion of intestine, which is another reason it is so often found in a rupture. When it has remained long in any sack, it forms a pretty compact mass, sometimes having no connection with, but at other times adhering to the inner surface of the sack. There is frequently no inflammation produced in the omentum while in this situation, but occasionally violent inflammation takes place, which may even advance to mortification.

A portion of gut is very often lodged in the fack of a rupture, either by itself, or along with a portion of the omentum. The portion of gut is fometimes very fmall, but at other times it is very confiderable. Very often the functions of the intestines go on properly in this situation, but occasionally violent inflammation is produced, interrupting their function, and often terminating fatally. This inflammation is produced by the pressure of the narrowest part of the sack against the gut, viz. that part of the fack which immedidiately passes out of the cavity of the abdomen. This inflammation exhibits the different appearances, upon diffection, which we have so often explained. The gut too, is frequently found mortified: this is fhewn by its dark colour, by its want of proper tenacity, and by the air which is

formed. When the inflammation of the gut in a fack has not been very violent, and has receded, it frequently leaves adhesions behind it, connecting the gut with the inner surface of the sack. It is perhaps possible too, that adhesions may be formed by long close contact, without inflammation.

When the fack of a rupture has not been of long standing, it confists of a thin, firm, white, opaque, membrane; this is an elongation of the peritonæum, fomewhat thickened by pressure.-When the fack has been of long standing, it is often very thick, and evidently confifts of a number of layers. The fack upon the infide has a very fmooth furface, and the membrane which forms this furface can be readily traced into the peritonæum, lining the cavity of the abdomen; the outer furface of the fack is more rough and coarfe in its texture: the fack, where it presses out of the cavity of the abdomen, has frequently a narrow neck, or aperture, and is distended below into a bag of confiderable fize. At other times, the communication between the fack and the cavity of the abdomen, is by a larger opening. Under these circumstances there is less danger of inflammation being produced by pressure against the gut.

In bubonocele, or that species of rupture which takes place at the ring of the external oblique muscle, the fack is usually quite distinct from the fack of the tunica vaginalis testis. At other times, there is no feparation between them, but the contents of the rupture are immediately in contact with the body of the tefficle: this kind of rupture is called the hernia congenita. It was formerly supposed, when it happened, to arise from a portion of the fack of the rupture and of the tunica vaginalis giving way, fo that the contents of the rupture fell into the cavity of the tunica vaginalis testis, and came in contact with the testicle. Upon a little reflection, it might have been feen that this could hardly take place; but the account of this appearance was not known till it was explained by Dr. Hunter. Baron Haller discovered, that till about the eighth month the testicles do not descend into the scrotum, but are fituated in the cavity of the abdomen under the kidneys. When they defcend into the fcrotum, the peritonæum which covers them is necessarily drawn down along with them through the ring of the external oblique muscle; it then forms a bag, whose upper extremity communicates with the cavity of the abdomen. Baron Haller had also observed, that in infants a portion of intestine fometimes falls down into this

bag after the testicle, or along with it, producing what he called the hernia congenita. The communication between the bag and the abdomen is commonly foon closed, for it is not open at birth. It appears, however, if it is prevented from closing at the usual time, that it does not close afterwards, but remains open through life .--Hence, if any portion of an intestine, or of the omentum, falls into the elongated fack of the peritonæum, it must be in contact with the testicle. When Dr. Hunter became acquainted with the observations of Baron Haller upon the descent of the testicles, he saw at once that the species of rupture fometimes to be met with in adults, where a portion of intestine or omentum is in contact with the testicle, might be easily explained. His explanation corresponded with what we have just given, and has been univerfally adopted by anatomifts and furgeons.

Schirrus and Cancer of the Intestines.

Schirrus is a disease which takes place much more commonly in the great than in the small intestines, but the latter are occasionally affected by it. I have seen both a schirrous tumour, and a cancerous ulcer in the duodenum. In the great

intestine at an advanced period of life, schirrus is not uncommon; it is not equally liable to affect every portion of this intestine, but is to be found much more frequently at the figmoid flexure of the colon, or in the rectum, than any where else; the reason of this is, perhaps, difficult to determine. There is certainly more of glandular structure in the inner membrane of the great intestine, towards its lower extremity, than in any other part of it, and I think that this fort of structure has a greater tendency to be affected with schirrus, than the ordinary structures of the body: the gut, too, is narrower at the figmoid flexure than at any other part, and therefore must be more liable to be injured by the paffage of hard bodies; these, by their irritation, may excite the disease of schirrus in a part which was pre-disposed to it. What we have now faid, however, is merely conjectural.

The schirrus sometimes extends over a considerable length of the gut, viz. several inches; but generally it is more circumscribed. When it is affected with schirrus, it exhibits the same appearances of structure which were described when speaking of schirrus of the stomach. The peritonæal, muscular, and internal coats are much thicker and harder than in a natural state. The muscular too is subdivided by membranous septa,

and the internal coat is fometimes formed into hard irregular folds. It often happens that the furface of the inner membrane is ulcerated, producing cancer. When schirrus affects the gut, the paffage at that part is always narrowed, and fometimes fo much fo as to be almost entirely obliterated. The obliteration, or stricture, would fometimes appear to be greater than in proportion to the thickness of the fide of the diseased gut : this, most probably, depends upon the contraction of the mufcular fibres of the gut, which, although difeafed, have not altogether lost their natural action. Where the passage is very much obstructed, the gut is much enlarged immediately above the obstruction, from the accumulation of contents in that part of the intestine. While this difease is going on in a portion of the intestine, adhesions are formed between it and the neighbouring vifcera, and the ulceration fometimes fpreads from one to the other.

The inner membrane of the great intestine I have seen a good deal thickened, and formed into small irregular tubercles, some of which were of a white, and others of a yellowish colour; the peritonæal and muscular coats were also thicker and harder than in a natural state. This is not a frequent appearance of disease, but it has sometimes been found to take place in very severe

dysenteries, such as in that which has been defcribed by the late Sir John Pringle.\*

I have also seen the internal membrane of the great intestine formed into broad thick folds, in which a considerable quantity of blood was accumulated; these folds were perfectly independent of the state of the contraction in the muscular coat, and were very different in their appearance from the irregular puckering which is often seen in the inner membrane of the great intestine.—

When these folds were examined: they were found to consist of an accumulation only of cellular membrane, lying behind the inner coat of the gut. What was the effect of this diseased structure in the living body, I had no opportunity of learning.

Upon the inner furface of the great intestine, about two inches above the anus, little processes fometimes grow from the internal membrane: they generally surround the gut at short distances from each other, so as to form a fort of circle.

Piles, and fiftulæ in ano, are diseases which are extremely common, but which hardly ever become an object of examination after death; they have therefore not been so commonly introduced into accounts of morbid appearances

<sup>\*</sup> See Pringle's Difeases of the Army, p. 246.





should be an external opening of the canal only; or, that there should both be an external opening, and another into the gut.

It is a species of monstrous formation, not very uncommon, that the rectum does not terminate in the anus, but in a cul-desac, without reaching the external surface. Sometimes the extremity of the gut lies immediately behind the skin, which is there thinner, and shews where the anus ought to have been placed. Under these circumstances, it is easy to make an artificial anus. It happens often, however, that the rectum terminates more than an inch behind the skin, and then a remedy by an operation is much more difficult and uncertain.

I have also seen the rectum terminate in the bladder, from an original error in the formation, so that there was no other external opening to the rectum than by the urethra: this was in a child at birth; the mal-formation was of such a kind, as neither to admit of a remedy by art, nor to allow of life being continued. The rectum has also been known to terminate in the vagina, from a defect in the original formation; but this is very uncommon.

#### Worms.

Worms are formed in the intestines of man, as well as of many other classes of animals, but not so frequently in the former as in the latter. In most quadrupeds and fishes it is extremely common to find a number of worms upon opening their intestines.

The worms which are found in the human fubject, may be reduced to three general classes, viz. the lumbricus teres; tænia, and ascaris.

### Lumbricus Teres.

The lumbricus teres, or round worm, is much more commonly found in the intestines of children, than in those of persons full grown, or advanced in life: it is very usually met with in the first, but rarely in the two last. The lumbricus teres has been often confounded with the common earth-worm, to which it bears a general resemblance, although it is really very different. By most practitioners, indeed, at present, there is known to be a difference, yet it is not very well understood in what this difference confists. The two species of worms, if attentively

examined, will be found to differ a good deal from each other in their external appearance. The lumbricus teres is more pointed at both extremities, than the common earth-worm; the mouth of the lumbricus teres confifts of three rounded projections, with an intermediate cavity. The mouth of the earth-worm confifts of a finall longitudinal fissure, fituated on the under surface of a finall rounded head. Upon the under furface too of the worm there is a large semilunar fold of skin, into which the head retreats, or out of which it is elongated, which is entirely wanting in the lumbricus teres. The anus of the lumbricus teres opens upon the under furface of the worm, a little way from its posterior extremity, by a transverse curved fissure; the anus of the earth-worm opens by an oval aperture at the very extremity of the worm. The outer covering or skin in the lumbricus teres is less fleshy, and less strongly marked by transverse rugæ, than in the earth-worm. In the latter there is often to be feen a broad white band, furrounding the body of the worm; but in the lumbricus teres, this is entirely wanting. On each fide of the lumbricus teres there is a longitudinal line very well marked; in the earth-worm there are three longitudinal lines upon the upper half of its surface, but these are faintly marked, fo as to be hardly observable.

The lumbricus teres has nothing corresponding to feet; whereas the earth-worm has on its under furface, and towards its posterior extremity, a double row of processes on each side, very sensible both to the eye and the singer, which manifestly serve the purposes of feet in the locomotion of the animal.

The internal structure of both animals is also extremely different. In the lumbricus teres, there is an intestinal canal, nearly uniform and smooth in its appearance, which paffes from one extremity of the worm to the other;\* in the earth-worm, there is a large and complex stomach, confisting of three cavities; and the intestinal canal in the latter is likewise larger, and more formed into facculi, than the former. The parts fubfervient to generation are very different in both: in the lumbricus teres there is a distinction of sex, the parts of generation being different in the male and in the female; in the common earth-worm the organs of generation are the same of each individual, as this animal is hermaphrodite. The appearance too of the organs of generation, is extremely different at first fight in the one species of animal, and the other: There is an oval mass

<sup>\*</sup> Near the head of the lumbricus teres, the canal is narrower than it is any where else, and also somewhat distinct in its limits, which may be considered as the œsophagus.

fituated at the anterior extremity of the earth-worm, refembling a good deal the medullary matter of the brain; in the lumbricus teres this fubstance is wanting.\* These are the greater differences between the one species of animal and the other, which are obvious upon a very moderate attention to each. Many other differences would, no doubt, be found by a person who might choose to prosecute their anatomy minutely.

### Tænia.

The tænia, or tapeworm, is rarely found in the intestines of the inhabitants of this country, but is very common in those of some countries on the continent, particularly Switzerland. It consists of a great many distinct portions, which are connected together so as to put on a jointed appearance; these joints are commonly of a very white colour, but are occasionally brownish, which depends on a sluid of this colour that is found in their vessels. The worm is usually very long, extending often many yards, and seldom passes entire from the bowels. This circum-

<sup>\*</sup> What this substance is I do not know, and I have only mentioned its resemblance to the medallary matter of the brain, in order to give a clearer description of it.

stance has prevented the extremities of the tænia from being often seen.

The head of a tænia is somewhat of a square form, with a narrowed projection forwards; in the middle of this projecting part there is a distinct circular aperture, around which is a single or a double row of curved sharp processes. At the square edge of the head, there are situated four round projecting apertures, at equal distances from each other: this head is placed upon a narrow jointed portion of the worm, of considerable length, and which gradually spreads itself into the broader joints, of which the body of the worm is composed.

The body of the tænia, which is most commonly found in the human subject, consists of thin, slat, pretty long joints, on one edge of which there is a projection, with a very obvious aperture. In the same worm some of these joints appear considerably longer than others; this probably depends on one joint being contracted, while another is relaxed. The apertures which we have just mentioned are generally placed on the alternate edges of the contiguous joints; but this is not uniformly the case; they are sometimes placed on the same edges of two, or even several contiguous joints. When these joints are examined attentively, there are frequently seen, in each of them, veffels filled with a brownish fluid, and disposed in an arborescent form. Around the edges of each joint, there is also a distinct serpentine canal.\* The last joint of a tænia resembles very much a common joint rounded off at its extremity, and without any aperture. The description which I have given, is chiesly taken from what I have seen myself, and corresponds, I believe, with what is most common. There are differences, however, in tæniæ described by authors, chiesly with regard to the number and situation of the oscula of the joints, so as to distinguish them into different species; but they are all formed upon one general plan.

### Ascaris.

The afcaris is a very small worm, which is often found at the lower end of the rectum in children, and even more frequently in adults than is commonly imagined. It is white in its colour, and about half an inch in length; it is a little narrowed at the extremity where its head is placed,

\* This, as well as the vessels disposed in an arborescent form, is very distinctly seen injected in some preparations which have been made, and given to me by an ingenious young surgeon, Mr. Carlisse. and at the other extremity it terminates in a long, very fine, transparent process. These worms are more or less surrounded with mucus, and this is probably secreted in greater quantity by the glands in the inner membrane of the rectum, from the irritation occasioned by the worms.

There is nothing in the economy of animals more obscure than the origin of intestinal worms; were they found to live in fituations out of the bodies of living animals, one might readily suppose that their ovula were taken into the body along with the food and drink, and there gradually evolved into animals. This, however, is not the case; they do not feem capable of living for any length of time in any fituation, except within a living animal body, which appears to be the proper place for their growth and residence. We might therefore be led to another supposition, viz. that intestinal worms are really formed from the matter contained in the intestines, which previously had no regular organization; but this idea is widely differ-. ent from all analogy in the production of animals, where there has been any fatisfactory opportunity of examining this production. The origin therefore of fuch animals is a subject of much obscurity, nor do I pretend at all to throw any light upon it. When the whole evidence, however, in support of the one and the other opinion, is compared together, I own, that the grounds for believing that in some orders of animals equivocal generation takes place, appear stronger than those for a contrary opinion.

It is not unufual to find air accumulated in the intestinal canal, in greater or less quantity; this air is sometimes accompanied with a slight inslammation of the peritonæum, and at other times it is not. In such cases the blood vessels, creeping upon the intestines, are frequently silled with air, but not uncommonly they are without it. Air is often let loose into the intestines after death by putresaction; but that which we wish particularly to consider here, has been formed during life.

There are only two ways in which we can well conceive air to be formed in the intestines: the one is, some new arrangement in the contents of the intestines, by which air is extricated: the other is, the formation of air in the blood vessels of the intestines by a process similar to secretion, and which air is afterwards poured out by the extremities of the exhalant arteries into the cavity of the intestines. That the blood vessels of an animal body have this power there can be no doubt; and I own I am inclined to think that this is a very frequent mode by which air is accumulated in the intestines. This air probably differs somewhat at different times: in several

of containing any proportion of inflammable air, but always a very fensible proportion of fixed air. It requires, however, to be examined by some person well acquainted with chemical experiments, in order that its ingredients may be

exactly ascertained.

These are the more common appearances of diseased, or preternatural structure in the intestines; but I have likewife had an opportunity of observing others, which are of rarer occurrence. In one or two instances, I have seen a fort of bony matter thrown out upon the furface of the inner membrane of the gut: I have even feen an adhesion between two portions of intestine, converted into bone. It would appear, that almost every part of the body is endued with a power of taking on this process. It may not improperly be confidered, as a natural process misplaced. An adhesion being once formed, has the same power (as far as we know) of running into different processes, as the cellular membrane, which makes a part of the original structure. It may therefore form bone, as readily as cellular membrane, or fome other membranes of the body, which have a refemblance to the membrane of adhesions, as the pleura, and the peritonæum.

I have feen one of the valvulæ conniventes much larger than ufual, and paffing round on the infide of the jejunum, like a broad ring. The canal of the gut was necessarily much narrowed at this ring, but no mischief had arisen from it.—

This mal-formation, however, might have laid the foundation of fatal mischief. Some substance too large to pass, might have rested on the ring, and produced there inflammation, ulceration, and ultimately death.

Calculous matter has fometimes been known to be accumulated in some part of the cavity of the intestinal canal, especially the great intestine; but this has not come under my own observation, and, at least in the human subject, it is a very rare occurrence.\*

Small-pox puffules have been faid to be fometimes formed in the intestines of persons who have died from this disease.† How far this may have occasionally taken place, I will not pretend to say, but late dissections, upon the best authority, have not confirmed this sact.

<sup>\*</sup> Vid. Lieutaud, Tom. 1, p. 77, 78.

<sup>+</sup> Vid. Lieutaud, Tom. 1, p. 3. 71,



in mortification are the fame when it affects the mesentery, as any other part, and they have been already explained.

The absorbent glands of the mesentery are frequently found to be scrofulous, and this is more apt to take place in children, than at a more advanced age. When affected with this difease, the glands exhibit different appearances, according to its progrefs: they are enlarged in their fize, and are fofter to the touch, than in a natural state. When cut into, they fometimes flew very much the natural structure; but more frequently they are changed, in part, into a white, foft, curdly matter, and this is not uncommonly mixed with pus. When the absorbent glands of the mesentery are univerfally affected with fcrofula, and very much enlarged in their fize, the abdomen is very tumid, and the face and extremities appear a good deal emaciated.

When a portion of the intestinal canal becomes cancerous, some of the absorbent glands in the mesentery generally become affected with the same disease: this is in consequence of the matter of cancer being conveyed to those glands by absorbent vessels. The glands become enlarged in size, and are changed into hard masses of a schirrous, or cancerous texture.

The absorbent glands of the mesentery are sometimes found filled with an earthy, or bony matter; but this is to be considered as a rare occurrence.\* The absorbent glands at the lungs, are more liable to be affected with this disease.

Hydatids have also been occasionally found adhering to the mesentery.

Tumours, likewise, consisting of a fatty matter, have been seen attached to the mesentery; but these I believe to be very uncommon.

AND THE PERSON OF THE PERSON O

\* Vid. Med. Transactions, Vol. 1, p. 361.



ances which have been described when speaking of the inflammation of the peritonæum, of which it is a part. It is crowded with a great number of very minute vessels, which carry a florid blood, and is thicker than natural. There is also thrown out upon its surface, a layer of coagulable lymph; this layer is thicker on some occasions than others, and often glues the liver more or less completely to the neighbouring parts. There is at the same time thrown out, some quantity of serous sluid.

## Adhesions.

It is more common to fee adhesions formed, which are the consequence of a previous inslammation in the membrane of the liver, than to see the membrane in an actual state of inslammation. These adhesions are formed from the coagulable lymph of the blood, which undergoes a gradual progress of change, as we have formerly described. They consist very commonly of a thin, transparent membrane, which joins the surface of the liver to the neighbouring parts. This junction may either be general, over one extended surface of the liver, or it may consist of a number of processes of adhesion: the adhesion is sometimes by a membrane of considerable length; at other times, the adhesion

is very close, the surface of the liver being immediately applied to the neighbouring parts. The surface of the liver, where these adhesions are most commonly found, is the anterior, by which it is joined to the peritonæum lining the muscles at the upper part of the cavity of the abdomen.

When an abscess is formed in the substance of the liver, and points externally, these adhesions are of great use in preventing the pus from escaping into the general cavity of the abdomen. Adhesions are also frequently found connecting the posterior surface of the liver to the stomach, and the duodenum; and these may also be useful in abscesses of the liver, near its posterior surface, by preventing the matter from passing into the general cavity of the abdomen, and conducting it either into the stomach, or the upper part of the intestinal canal.

Inflammation of the Substance of the Liver.

It does not often happen, at least in this country, that the substance of the liver is actually found in a state of inflammation. Where its membrane is inflamed, the substance is sometimes inflamed which lies immediately under it; but it rarely happens that the general mass of the liver is in-

flamed. In warmer countries, the fubstance of the liver is much more liable to inflammation than in Great-Britain. When the liver is generally inflamed through its substance, it is a good deal enlarged in fize, and of a deep purple colour.\* It is also harder to the touch, than in its healthy state. Its outer membrane is fometimes affected by the inflammation, and at other times is not. It is frequently accompanied with a jaundiced colour of fkin, arifing from the bile not getting readily into the ductus communis choledochus, on account of the pressure of the inflamed liver on the pori biliarii. When this inflammation has continued for fome time, abfceffes are formed, and then the active state of the inflammation very much fubfides. These abscesses are sometimes of large fize, fo as even to contain fome pints of pus. Sometimes the whole of the liver is almost converted into a bag containing pus. When inflammations of the liver have been of confiderable standing, they are not uncommonly attended

\* May not the purple colour arise from the accumulation of blood in the branches of the vena portarum?

As this vein most probably performs the office of an artery in the liver, is it not probable, that its small branches take on the same actions as the small branches of an artery during inflammation?



The tubercles which are formed in this disease occupy generally the whole mass of the liver, are placed very near each other, and are of a rounded fhape. They give an appearance every where of irregularity to its furface. When cut into, they are found to confift of a brownish or yellowish white folid matter. They are fometimes of a very fmall fize, fo as not to be larger than the heads of large pins; but most frequently they are as large as a small hazel nut, and many of them are sometimes larger. When the liver is thus tuberculated, it feels much harder to the touch than natural, and not uncommonly its lower edge is bent a little forwards. Its fize, however, is generally not larger than in a healthy state, and I think it is sometimes fmaller. If a fection of the liver be made in this state, its vessels seem to have a smaller diameter than they have naturally. It very frequently happens that in this state the liver is of a yellow colour, arifing from the bile accumulated in its substance; and there is also water in the cavity of the abdomen, which is yellow, from the mixture of bile. The gall-bladder is generally much contracted, and of a white colour, from its being empty. The bile, from the pressure of the hard liver upon the pori biliarii, does not reach the ductus hepaticus, and therefore cannot país into the gall-bladder. The colour of the skin in such cases is jaundiced, and it remains permanently so, as it depends on a state of liver not liable to change. This is the common appearance of what is generally called a schirrous liver: but there are other hard tubercles formed in it, which resemble more the structure of schirrus in other parts of the body.

# Large white Tubercle of the Liver.

Hard white masses are sometimes formed in the liver, of a considerable fize. They are often found as large as a chesnut, but I have seen them both a good deal larger and smaller than this size. They are to be found near the surface of the liver in greater number, than near the middle of its substance: two or three frequently lie contiguous to each other, with a considerable portion of the liver, in a healthy state, interposed between them and a cluster of similar tubercles. They consist of a very sirm, uniform, opaque, white substance, and generally somewhat depressed, or hollow, upon their outer surface. The liver in this disease is frequently a good deal enlarged beyond its natural size.

These tubercles appear to be first formed round the blood vessels of the liver, as is seen in making



Such tumours might perhaps be called fcrofulous, but there is no strong evidence in support of this opinion; and there is certainly no resemblance between this fort of tumour and either a scrofulous tubercle of the lungs, or a scrofulous absorbent gland. It would be worth while, as such appearances occur, to examine them particularly, so that at length the nature of the disease may be ascertained. This inquiry will be much affisted by an accurate knowledge of the symptoms, and of the general tendency of the constitution in the living body.

## Scrofulous Tubercles of the Liver.

Tubercles are occasionally found in the liver which resemble exactly the tubercles of the lungs; but this is a very rare appearance of disease. They have the same size, the same structure, and the same seeling to the touch. In the only instance which I have seen of this disease, the tubercles were generally dispersed through the substance of the liver at pretty regular distances, and did not render the surface of the liver irregular, as in the common fort of tubercle.

I have likewise seen the liver much more slaccid in its substance than is natural, with reddish tumours, of confiderable fize, interspersed through it, which contained a thick fort of pus. I am inclined to confider this liver as scrofulous, because it was found in a person whose general constitution had strong indications of scrofula, and where there were found many scrofulous absorbent glands on examining the body.

# Liver very foft in its Substance.

The liver is not unufually found much more flaccid in its fubstance than natural, without any other appearance of disease. It feels upon such occasions nearly as soft as the spleen, and is commonly of a leaden colour. This change must arise from a process which takes place through its whole substance, and seems to be what Mr. Hunter has called the interstitial absorption. By this process is meant, where the absorbents remove insensibly small ingredient parts out of the general mass of any structure in an animal body without ulceration. This state of liver is very rarely, if ever, found in a very young person, and is most common in persons who are advanced in life.

## Liver very hard in its Substance.

There is a very contrary state of the liver, not at all unufual, viz. where it is much harder than natural, and when cut into, exhibits no peculiar mode of structure. Upon the surface of these livers, there is not uncommonly a thready appearance of membrane, disposed somewhat in a radiated form, and the lower edge is bent a little forwards. This I believe to be the first step in the progress towards the formation of the common tuberculated liver. I have fometimes feen fmall tubercles formed upon a part of the furface of fuch a liver, which were exactly of the common fort. From this appearance, it is probable, that additional matter is deposited in the interstices, through the general mass of the liver, rendering it much harder, and that this matter, together with part of the ordinary structure of the liver, is converted into tubercles. What we have now faid, however, is to be confidered as entirely conjectural. This hardened state of liver is sometimes accompanied with a beginning afcites, and at other times is without it.

### Hydatids.

There is no gland in the human body in which hydatids are so frequently found as the liver, except the kidneys, where they are still more common.\* Hydatids of the liver are usually found in a cyst, which is frequently of considerable size, and is formed of very firm materials, so as to give to the touch almost the feeling of cartilage. This cyst, when cut into, is obviously laminated, and is much thicker in one liver than another. In fome livers it is not thicker than a fhilling, and in others it is near a quarter of an inch in thickness. The laminæ which compose it are formed of a white matter, and on the infide there is a lining of a pulpy fubstance, like the coagulable lymph. The cavity of the cyft I have feen fubdivided by partitions of this pulpy fubstance. In a cyst may be found one hydatid, or a greate: number of them. They lie loofe in the cavity, fwimming in a fluid, or fome of them are attached to the fide of the cyft. They confift of a round bag, which is composed of a white, femi-opaque,

<sup>\*</sup> Although the hydatids of the liver, and the kidney, have got the same name, yet there is reason so believe that they are different from each other.

pulpy matter, and contain a fluid capable of coagulation. Although the common colour of hydatids be white, I have occasionally seen some of a light amber colour. The bag of the hydatid confifts of two laminæ, and possesses a good deal of contractile power. On the infide of an hydatid, there are fometimes found fmaller ones, which are commonly not larger than the heads of pins: these are attached to the larger hydatid, either at fcattered irregular distances, or form small clusters: they are also found floating in the liquor of the larger hydatids. Hydatids of the liver are often found loofe and unconnected with each other; but at other times, they inclose each other in a feries, like pill-boxes. The most common situation of hydatids of the liver, is in its substance, and inclosed in a cyst; but they are occasionally attached to the outer furface of the liver, hanging from it, and occupying more or less of the general cavity of the abdomen.

The origin and real nature of these hydatids are not fully ascertained; it is extremely probable, however, that they are a fort of impersect animals. There is no doubt at all, that the hydatids in the livers of sheep are animals: they have been often seen to move when taken out of the liver, and put into warm water; and they retain this power of motion for a good many hours after a sheep has

been killed. There is a great analogy, however, between hydatids in the liver of a sheep, and in that of the human subject. They are both contained in strong cysts, and they both consist of the fame white pulpy matter. There is undoubtedly fome difference between them in simplicity of organization; the hydatid in the human liver being a fimple uniform bag, and the hydatid in that of the sheep having a neck and mouth appended to the bag. This difference need be no confiderable objection to the opinion above stated. Life may be conceived to be attached to the most simple form of organization. In proof of this, hydatids have been found in the brains of sheep, which refemble exactly those in the human liver, which have been feen to move, and therefore are certainly known to be animals. The hydatids of the human liver, indeed, have not, as far as I know, been found to move when taken out of the body and put into warm water; were this to have happened, no uncertainty would remain. It is not difficult to fee a good reason why there will hardly occur any proper opportunity of making this experiment. Hydatids are not very often found in the liver, because it is not a very frequent difeafe there; and the body is allowed to remain for so long a time after death before it is examined, that the hydatids must have lost their living prin-

ciple, even if they were animals. The probability of their being animals, however, is very strong; and it appears even more difficult to account for their production according to the common theory of generations, than for that of intestinal worms. We do not get rid of the difficulty by afferting, that hydatids in the human liver are not living animals, because in sheep they are certainly such, where the difficulty of accounting for their production is precifely the same. I have mentioned, however, already every thing I have to fay upon this fubject, when speaking of intestinal worms. If any person should wish to consider hydatids more minutely, he will find an excellent account of them published by Dr. John Hunter in the Medical and Chirurgical Transactions.\*

Upon the infide of a cyft, exactly refembling that which contains hydatids, I have feen adhering a white, friable, earthy matter; what was its exact nature, I cannot determine, but it was in fome degree foluble in the marine acid.

Worms\* have been faid to be found in cysts of the liver, as well as in the biliary ducts. Instances of this fort are extremely rare, and have not come under my own observation.

<sup>\*</sup> See Medical and Chirurgical Transactions, p. 34.

Vid. Lieutaud, Tom. 1, p. 194.

#### CHAP. X.

## Diseased Appearances in the Gall-bladder.

## Inflammation of its Coats.

I HE coats of the gall-bladder are very rarely inflamed, independently of the inflammation of the membrane which covers the posterior furface of the liver. When inflammation attacks this membrane, it naturally fpreads over the outer coat of the gall-bladder, which is a continuation of it, and may affect the other coats of the gall-bladder, if it should have arisen to a violent degree. Inflammation, however, of the outer coat of the gall-bladder will not commonly be attended with inflammation of the others, because it is not closely applied to them, there being interposed a considerable quantity of cel-The appearances of inflam lular membrane. mation in the coats of the gall-bladder, are ex actly the same with what take place in the in

flammation of fimilar structures. These we have already fully described, and shall not here repeat them.

#### Adhesions.

It is a very common appearance, upon diffection, to find the gall-bladder connected by adhesions, either to the small end of the stomach, or the beginning of the duodenum. These are the consequence of a previous inflammation in the outer coat of the gall-bladder, and resemble exactly the adhesions which we have already described.

It is rare that inflammation of the gall-bladder advances to ulceration: the accumulation of gall-stones in it, very rarely produces any effect of this kind. There is but one instance of ulceration of the gall-bladder which has fallen under my own observation, and this is preserved in Dr. Hunter's collection.

## Coats of the Gall-bladder fchirrous.

I believe it to be very uncommon that the coats of the gall-bladder become fchirrous. It

has occurred to me, however, to observe one instance of it. The coats of the gall-bladder were in this case above a quarter of an inch in thickness, and studded with hard white tubercles, exactly similar to what we have described in the liver: indeed the liver to which this gall-bladder belonged was affected with the same disease. It is probable that the gall-bladder is scarcely ever affected with schirrus, unless it has previously taken place in the liver.

# Coats of the Gall-bladder bony.

I have likewise seen the coats of the gall-bladder very much thickened, and converted in many parts into a sort of bony substance; but this is to be considered as a very rare appearance of disease.

# Diseased State of Ducts.

There are two diseases in the excretory ducts of the liver and the gall-bladder which have come under my observation: the one is their obliteration, and the other their dilation. The first is extremely rare, and the only instance which I have seen of it was in the ductus cysticus; the other is not uncommon. The ductus hepaticus, ductus cylticus, and ductus communis choledochus, are fometimes dilated to an almost incredible size. I have seen the ductus hepaticus and choledochus so much dilated as to be nearly an inch in transverse diameter. These dilatations of the biliary ducts hardly ever take place but from one cause, viz. the passage of gall-stones; and it is associated have been known to pass into the duodenum.— This ought to afford a good ground of comfort to persons who are labouring under this complaint.

It may not be improper to take notice here, that I have once feen an immediate communication, by a short canal, between the gall-bladder and the small end of the stomach. This lusus naturæ is very rare, and is probably attended with little inconvenience to the animal functions.

#### Gall-Stones.

It is not an uncommon appearance of difease in examining dead bodies, to find gall-stones, either in the gall-bladder, or in some of the biliary ducts. The gall-bladder is sometimes much enlarged in its size, and full of them. In this

case its coats are often a good deal thickened, and this probably arises chiefly from the efforts of the contractile power of the gall-bladder to expel them. The number of stones accumulated in the gall-bladder is fometimes very great; above a thousand have been taken out of one gall-bladder, which are preserved in Dr. Hunter's collection. When there is a folitary gall-stone in the gall-bladder, it is occasionally very large; I have known an inftance of one which was fully the fize of a hen's egg. When there is but one gall-stone, either in the gall-bladder, or in the biliary ducts, it is generally of an oval shape; when there is a confiderable number, they acquire, by rubbing upon each other, a great many fides and angles.

There is a great variety in the external appearance of gall-stones with respect to colour: some are whitish, others are black; they are also of a yellowish, a greenish, a light brown, a dark brown, and a redish brown colour. These are the principal varieties in colour, but there are many other small differences which it would be very difficult to express in words. Gall-stones differ also very much in the smoothness of their surface, some being very smooth, and others a good deal suberculated.

When cut or broken, gall-stones are commonly found to confift of concentric laminæ upon the out fide, and in the centre of a radiated texture. The laminated part bears fometimes a large proportion to the other, and at other times the contrary happens. The laminated and radiated structures are sometimes compact, and at other times confift of loofe matter. It likewife occasionally happens that both the laminated and the radiated structures are very obscure, and the gall-stone appears a good deal like an unformed folid mass. The laminated part on the out fide very frequently confifts of a different substance in appearance, from the radiated structure in the centre; and it is not unufual to find the structure in the centre to consist of shining white crystallizations, which have a good deal the look of spermaceti.

Gall-stones being very different both in their outward appearance and their internal structure from each other, naturally lead us to suppose that they may also differ in their chemical properties. Upon this subject I can only speak very generally; but any trials which I have made incline me to this opinion. Very sew gall-stones yield a bitter taste, which shews that they do not consist of inspissated bile; but in some I have sound the taste intensely bitter. Almost all of them melt in the

flame of a candle; but I have met with one fort, of a very black colour, which did not melt, but burnt exactly like a cinder.

All the gall-stones which I have examined disfolve in the nitrous acid. They are separated into a fine black powder when put into the vitriolic acid, especially if exposed to a fand heat. By the marine acid they are not acted upon at all in the common heat of the atmosphere, and are even but little affected by it when exposed to a fand heat for a considerable time.

Most of the gall-stones which I have examined are somewhat soluble in oil of turpentine, in the common heat of the atmosphere, but still not in a considerable degree; and one fort I have sound to be in this heat altogether insoluble. When put into this oil, and exposed to a sand heat, they are much more readily acted upon. Some are converted into a kind of oil, which sinks to the bottom of the oil of turpentine; others are partly soluble, tinging the oil of turpentine of a brownish colour, and are partly separated into a powder.

Most gall-stones appear to be very little affected by spirit of wine in the common heat of the atmosphere, but are either partly or entirely soluble in it in a boiling heat.\* Such are the gen-

<sup>\*</sup> When some biliary calculi are exposed to spirit of wine in a boiling state, white slaky crystals are soon

eral refults from a good many trials of my own; but I speak upon this subject with very little confidence. It ought to be taken up by an able chemist, who is much in the habit of making experiments, and his experiments should extend to a great variety of gall-stones, which differ from each other in their appearance.

#### Bile.

The bile in the gall-bladder is found to differ in different bodies; but this is too common to arife from difease, and must depend on natural circumstances. It is sometimes of a green, at other times of a brownish yellow or a purer yellow colour. The brownish yellow colour is the most common. It is always more or less visited, and the variety in this respect is considerable; in man it generally appears a good deal more visited than in most other classes of animals. Upon one occasion, I have seen it as ropy as the mucus

formed upon its cooling. When they are exposed to spirit of wine in the common heat of the atmosphere, it is some weeks before crystals begin to be formed, and they appear to be more pointed in their shape than the former. These crystals were, I believe, first observed by M. Poulletier de la Salle. See Elemens d'Histoire Naturelle et de Chimie, par M. de Fourcroy, Tom. p. 354.

which is commonly coughed up from the trachea. In opening dead bodies, the bile is almost always found to have transuded in small quantity through the coats of the gall-bladder, so as to tinge the neighbouring parts, especially the small end of the stomach, and the beginning of the duodenum. This is to be considered as a natural esset, which has taken place after death, and not as a diseased appearance. The coats of the gall-bladder, in consequence of death, have lost that compactness by which they were formerly able to confine the bile; it therefore transudes in small quantity, and tinges the neighbouring parts.

The gall-bladder is fometimes distended with bile so as to be of nearly twice its usual fize; at other times, there is no bile at all in its cavity, and under such circumstances it is white in its colour, and contracted into a very small fize.

It has been known to be diffended to an immense size, and to contain hydatids;\* but this state of it must be looked upon as extremely uncommon.

orpers' to be more pointed in their fluipe than the for-

Poulterier de la Salle. See Flemens d'Hitlaire Noturelle.

crede Chimie, par M. de Fourcroy, Tom poster, and

<sup>\*</sup> See Medical Communications, Vol. 1, p. 101.

#### CHAP. XI.

formed between the sploy a god sittle neighbon

parts, than to find its coats in an afficial flans

It is more conunou, however, to

Diseased Appearances of the Spleen.

Inflammation of the Coats of the Spleen.

The coats of the spleen are liable to inslammation; but this rarely takes place unless the peritonæum in the neighbourhood be also assected. The proper capsule of the spleen is so intimately connected with the peritonæum which is reflected over it, that it must necessarily partake of any inslammation affecting that portion of the membrane. When inslammation attacks the coats of the spleen, it exhibits exactly the same appearances which have been so often explained. They become much more crowded with vessels than in a natural state, are somewhat thicker, and throw out a layer of coagulable lymph.

### Adhesions.

It is more common, however, to find adhesions formed between the spleen and the neighbouring parts, than to find its coats in an actual state of inflammation. These adhesions consist of a white transparent membrane of more or less sirmness, and generally connect more or less closely the broad surface of the spleen to the diaphragm. They often connect too the spleen to the great end of the stomach, and a part of the transverse arch of the colon.

Inflammation of the Substance of the Spleen.

It is very rare to find the substance of the spleen either in a state of inflammation or suppuration; but such cases have occasionally been observed and related by authors.\* Instances also have been related where the spleen has been observed to be mortified, but this I should believe to be even much more rare than the former.

<sup>\*</sup> Vid. Lieutaud, Tom. 1, p. 222.

Vid. Lientaud, Tom. 1, p. 223.

### The Spleen extremely foft.

There is an appearance of the fpleen which is very common, and which perhaps is hardly confidered as a difease, yet surely it is a very obvious deviation from its healthy structure. The appearance to which I allude, is an extreme foftness of the spleen, so that when its capsule is broken (which under fuch circumstances is very tender) the substance of the spleen seems to be little else than a very fost, brownish-red mucus, intermixed with a spongy fibrous texture. This appearance of the spleen is hardly ever to be observed at a very early period of life, but is very common in middle and more advanced age. I should believe that such a state of the spleen is not marked by any peculiar feelings, so as to make the persons conscious of any disease taking place, and is probably of very little confequence in the general economy of the animal. Still, however, it is a very obvious deviation from the healthy structure of the spleen, and is not a state into which the spleen naturally degenerates in the gradual decay of the body.

T

### The Spleen very hard.

The spleen is sometimes much harder than natural, and at the same time is generally a good deal enlarged. When cut into, the natural structure seems to be preserved, except that it is much more compact, the parts being much more closely applied together.

This state of the spleen is generally considered as schirrous; and it very much resembles the hardened state of the liver, before the more common tubercles are formed in it. I do not recollect, however, to have seen in this state of the spleen, either schirrous tubercles formed, or any derangement of structure similar to schirrous in other parts of the body: it may therefore be a question, whether this is to be considered as the kind of derangement which a schirrous affection is capable of producing in this viscus, or whether it is to be considered as a disease really different. In this state of the spleen, water is sometimes accumulated in the cavity of the abdomen.

### Tubercles in the Spleen.

The spleen I have seen, once or twice, studded in its substance with small tubercles, exactly similar to the scrosulous tubercles of the lungs. These were placed at pretty regular distances from each other, not running into masses; and I do not recollect to have seen any of them in a state of suppuration.

#### Coats of the Spleen cartilaginous.

The coats of the spleen are sometimes converted into cartilage; and this disease may be considered in a great measure as peculiar to the spleen. It is, at least, much more common in this viscus than in any other.

The cartilage is generally to be found on the convex furface of the spleen, and extends over more or less of it, in different cases. It is likewise much thicker in one case than another; in some being not thicker than a shilling, and in others being four times as thick. It is generally formed in a smooth layer, but occasionally it is somewhat irregular. It is probable, that offisica-

tilage, but in the cases which have come under my own examination, bony matter was not to be observed. The cartilage into which the coats of the spleen are changed, does not resemble the cartilage at the extremities of the bones, but much more that of the nose and ears, although it is generally of a whiter colour. This diseased process, it is natural to think, is slow in its progress, can hardly produce any impediment to the functions of the spleen, and is probably not marked by any peculiar feelings to the patient.

### Spleen very large.

The spleen is sometimes sound to be three or sour times larger than its natural size, but with a structure perfectly healthy; and this more commonly happens to the spleen than to any other viscus. Although it may be looked upon as a monstrous growth of the spleen, rather than a disease, yet it may produce inconvenience by its pressure and by altering in some degree the situation of the neighbouring viscera.

\* Morgagni has seen offisication of a part of the capsule of the spleen. Vid. Epist. X. Art. 19. Epist. XIV. Art. 23.

Hydatids are occasionally formed in the spleen, which are the same kind with those of the liver; but they are much more common in the latter viscus than in the sormer.

Stony concretions have been feen occasionally in the spleen; but such cases are very rare, and have not fallen under my own observation.

The spleen has been said to be occasionally wanting, as a defect in the natural formation; but this too is very uncommon.

§ Vid. Morgagni, Epist. XXXVIII. Art. 34.

† Vid. Lieutaud, Tom. 1, p 231.

‡ Vid. Lieutaud, Tom. 1, p. 234.

#### CHAP. XII.

### Difeased Appearances of the Pancreas.

THE pancreas is subject to very few diseases. It seldom happens, upon examining dead bodies, that it exhibits any other than the healthy structure.

#### Pancreas hard.

It is not very uncommon to find the pancreas much harder than in its natural state, but without any appearance to the eye of the structure being altered. This I believe to be the beginning of a process, by which the pancreas becomes truly schirrous. It very seldom in this state shews, in any part, the real schirrous structure. I have seen this, however, to be the case, which renders it very probable, that the one is the beginning of a change into the other. When the pancreas in

any part assumes the schirrous structure, that part loses entirely its natural appearance, and is converted into a hard, uniform, white mass, somewhat intersected by membrane, like schirrus in other parts of the body.

## Calculi of the Pancreas.

Calculi are occasionally formed in the ducts of the pancreas. Of this I have only known one instance. The calculi were about the fize of the kernel of a hazel nut, with a very irregular surface, and of a white colour. When one of the stones was put into the marine acid, it dissolved very quickly, with the extrication of a large quantity of air. The stones, therefore, in this case differed very much from the nature of urinary calculi. It is probable, that calculi formed in the pancreas may differ somewhat from each other, as we find to happen in other calculi which are formed in the body; but as this is a very rare disease in the pancreas, it must be a long time before this point can be fully ascertained.

Abscesses have also been occasionally found in the pancreas, but very seldom.\*

<sup>\*</sup> Vid. Lieutaud, Tom. 1, p. 224.

## [ 140 ]

Worms have been faid to have been found in its excretory duct.

The pancreas has been faid to be entirely wanting, as a defect in the original formation.+

> § Vid. Lieutaud, Tom. 1, p. 248. + Vid. Lieutaud, Tom. 1, p. 247.

hadi of the Bereight and

the pancious. Of this Libert only longer one

the flower was put into the marine acid, in difficived

as we find to happen in other cateur which are

diledle in the paner are, it much be a long time be-



#### Abscesses of the Kidneys.

The fubstance of the kidneys is not often inflamed without advancing to suppuration, and perhaps there is no considerable gland in the body so liable to form abscesses as the kidneys. In some cases which I have seen, the abscesses have appeared to be of a common nature; but in the greater number of cases they have been scrosulous.

When a kidney is attacked with fcrofula, and the difeafe has advanced to suppuration, it exhibits different appearances, according to the degree of the advancement. Sometimes there are only one or two circumscribed abscesses containing a curdly pus, without any thing being particularly observable in the inner surface of the abscesses. Very frequently, however, the inner surface of the abscesses is lined with a pulpy granulated matter. These abscesses generally first destroy the mamillary portion of the kidney; and when they advance very far, they destroy almost the whole structure of the kidney, converting it into capsules, which surround a number of imperfect cavities that are lined with this pulpy substance.

The capfule, into which the kidney is changed by the progress of this disease, is on some occafions thicker than on others, is frequently of a confiderable hardness, and seems sometimes to be laminated. At the time when a kidney is so affected, it is not uncommon for the pelvis and ureter to partake of the disease, and a calculus is often found either in the abscess, or in the pelvis of the ureter.\*

#### Kidney Schirrous.

The kidney I have once feen converted into a firm, brown, uniform substance, somewhat interfected by membrane, in which the natural structure of this gland was entirely lost. The kidney was at the same time very much enlarged in size. This alteration of structure I should call schirrus, because it exactly resembles schirrus in other parts of the body, especially in the testicle. What esfect it had upon the secretion of the urine, I have had no opportunity of being informed. This disease I am persuaded is very rare in the kidneys.

\* In such cases, it is very probable that the calculus is the immediate cause of the other disease, the constitution being at the same time disposed to it. By the irritation of the calculus, inslammation and suppuration are produced in the kidneys, and these partake of the nature of the constitution.

## Kidneys very foft.

I have also seen the substance of the kidney converted into a foft loofe mass, resembling almost exactly the appearance of common fponge. On the furface there were many round interstitial cavities scattered at irregular distances; and when the fubstance of the kidney was cut into, it exhibited the fame fpongy structure. The blood veffels of the kidney were feen ramifying very diffinctly through the fpongy mass. There was no appearance of pus in the kidney, nor was there the most distant resemblance between this process, and the effects produced by suppuration. It was a process of a peculiar kind, by which a confiderable portion of the kidney was removed by the action, probably of absorbent vessels, and it feemed to act much more on the cortical than the tubular part of it. I am not at all exaggerating the effect of this diseased process, when I say that the kidney was rendered fully as foft as a common fponge. When shaken in water, the parts all feparated from each other, fomewhat like the unravelling of the fhaggy veffels of the placenta. Such an appearance of kidney has fallen under my observation two or three times, but I have only had once an oportunity of know-

ing any thing of the fymptoms which it produced during life. I was once fent for to a woman about two days before she died, who, besides other ailments, complained very much of a fevere pain in the region of the kidneys, and the urine which fhe paffed had a very large proportion of a fediment fomewhat refembling cream. How long fhe had complained of these symptoms I do not now recollect, but the time was confiderable. This very imperfect account is only thrown out to be some guide to future inquiry, that this diseased process may be more particularly afcertained. One thing furprifed me at the time, and perhaps without reason, viz. that this process had been attended with fevere pain. We are naturally led to expect in processes which go on slowly, and as it were imperceptibly, that the pain fhould be inconfiderable.

#### Hydatids of the Kidneys.

The formation of hydatids is not an uncommon difease in the kidneys. There are sometimes one or two considerable hydatids on the surface of the kidney, lying between its substance and capsule; and at other times, the natural structure of a kidney is almost entirely lost,

and is changed into a mass of small hydatids .-When this is the case, the mass is commonly much larger than in the natural fize of a healthy kidney. These hydatids do not appear to be of the fame nature with the hydatids of the liver: they are not inclosed in firm cyfts, like those of the liver: their coats are also thinner, and less pulpy, and not uncommonly they are almost as thin as any membrane of the body. I do not recollect likewife to have feen any instance of small hydatids attached to the coats of larger hydatids in the kidney, as may be frequently observed in the liver. It is therefore probable that the hydatids of the kidney depend on a difeafed alteration of the structure of this organ, and are not diftinct organized fimple animals.

## Calculi of the Kidneys.

The formation of calculi is not peculiar to the kidneys, but it is a more frequent disease in them than in any other part of the body. Small granules of stone are sometimes sound in the tubular portion of the kidneys, which may either be attended with inflammation, or be without it. Most commonly, however, there is sound a calculus of considerable size lodged either in some part of the

fubstance of the kidney, or in the pelvis of the ureter.

The last situation is by much the most frequent. When a stone in this situation is so large as not to be capable of paffing through the ureter, it is afterwards gradually increased in fize, from the contact of the urine. In its growth, it necessarily follows the branches of the pelvis, which are called infundibula, and is therefore of an arborescent form. Such calculi vary in their colour and furface; they are fometimes of a light brown, fometimes of a dark brown, and fometimes of a white colour. They are also fometimes smooth, and at other times a little roughened on their furface. Of the nature of urinary calculi we shall speak afterwards, when we come to take notice of the difeafed appearances of the bladder. When a stone in the pelvis of the ureter has increased to a very confiderable fize, it prevents the urine almost altogether from paffing into the ureter. The urine is therefore accumulated in the pelvis above the stone, and hence enlarges it very much in size, as well as the cavity of the kidney. From the preffure too of the urine behind the stone, the pelvis of the ureter, befides being enlarged, is thrust out from the fubflance of the kidney. If the interruption to the paffage of the urine from the kidney arises from some obstruction in the lower extremity of the ureter, or at the neck of the bladder, or in any part of the urethra, not only the pelvis of the ureter is then enlarged, but the ureter itself. I have seen the ureters of both kidneys enlarged from this cause to twice or thrice their natural size.

Whatever be the nature of the obstruction, if the pelvis of the ureter be very much enlarged from the accumulation of the urine, the cavity of the kidney is at the fame time enlarged. As this process advances, the substance of the kidney becomes more and more compressed, and its cavity becomes enlarged in proportion. The fubstance of the kidney is, at length, in a great measure loft, and is converted into a capfule, containing a great many cells, which communicate with each other. The capfule is fometimes very thin, and the whole mass is a great deal larger than the natural size of a healthy kidney. It is worthy of remark, that the urine is fecreted even when the natural structure of the kidney is almost entirely lost. This is both feen in the derangement of the kidneys now under confideration, and when they are converted into a mass of hydatids. It would appear from this fact, that either a very fmall portion of the natural structure of the kidneys is capable of fecreting very nearly the ordinary quantity of the urine, or that the urine





fluid, and this is probably what is meant by authors when they fay that they have found in the cavity of a renal capfule a fluid like ink. In fuch cases we may either suppose the dark coloured substance to be converted into the black-ish fluid; or we may suppose that the former is removed by absorption, and the latter is deposited.

The renal capfules have been observed to contain pus.\*

Little granules of stone have been found in the substance of the renal capsules.§

Both of these appearances are very rare, and have not fallen under my own observation.

\* Vid. Lieutaud, Tom. 1, p. 285.

CARRY OF BUILDING A SHOULD TO BE

#### CHAP. XIV.

Difeafed Appearances of the Bladder.

Inflammation of the Peritonwal Covering.

The bladder is covered on the outfide partially by the peritonæum, which is very liable to be inflamed. That portion of it which belongs to the bladder, is not very often inflamed by itself, but it partakes of the inflammation which extends over the membrane generally. The appearances accompanying its inflammation have been sufficiently often described. When the inflammation subsides, adhesions are frequently lest behind, connecting the bladder more or less to the neighbouring parts; in a semale, to the uterus, and in a male, to the rectum.

Inflammation of the inner Membrane.

The inner membrane too is fometimes inflamed. When this happens, the inflammation is either extended over the whole cavity, or is confined to a particular portion of it. The portion which is most frequently inflamed is near the neck of the bladder. This may arise from two causes; the one is, that in this fituation, or near it, fome obstruction is frequently found to the passage of the urine, which may produce irritation, and bring on more or less of inflammation; the other is, that inflammations of the urethra occasionally extend fome way within the cavity of the bladder, and even fometimes over the whole of it. It is well known that the inner membrane of the bladder hardly shews vessels which carry red blood in its natural state, but when it is inflamed it is crowded with a prodigious number of extremely fine blood veffels, and accompanied fometimes with fmall fpots of extravafated blood. When the inflammation is in a high degree, the muscular coat of the bladder may be affected; but as this is only loofely attached to the inner membrane, the inflammation will not very readily pass from the one to the other.

#### Ulcers.

Inflammation of the inner membrane of the bladder advances fometimes to the formation of

pus, and abfeeffes and ulcers are occasionally produced. Thefe, when the inflammation has been of the common fort, exhibit the ordinary appearances which have been often mentioned. They fometimes advance fo far as to destroy a portion of the bladder entirely, and to form communications between it and the neighbouring parts: as with the general cavity of the abdomen, with the reclum of the male, and the vagina in the female. When the communication is formed with the general cavity of the abdomen, the urine escapes into it, producing there general peritonæal inflammation, of which I recollect a very striking example. When the communication is formed with the vagina or the rectum, the urine will escape by these passages, producing in them more or less of irritation and inflammation.

When abscesses take place in the bladder, they are produced more frequently from local violence, than from a previous spontaneous inslammation. One of the most common causes of violence is the incision of the bladder in the operation of lithotomy. When the part has been very much irritated in the operation, or the constitution is such that it is excited into violent action from the common degree of irritation, an ulcer is formed at the lips of the wound, and

fpreads more or less into the cavity of the bladder.

It fometimes happens, although I believe very rarely, that the whole of the inner membrane of the bladder is destroyed by ulceration, and its muscular fibres appear as bare as if they had been nicely diffected. In the case where I recollect this process to have taken place most completely, the bladder was almost filled with a scrofulous pus. This had exactly the same appearance as when a scrofulous absorbent gland suppurates, viz. there was a curdly white matter mixed with pus.

#### Schirrus and Cancer.

I do not think that schirrus or cancer often attack the bladder by itself, but it sometimes partakes of this disease from its contiguity to parts which are very liable to it. The disease on some occasions spreads to the bladder from the rectum, and on others from the uterus: under such circumstances the bladder becomes very thick and hard, and exhibits the ordinary cancerous structure. Communications too are generally formed either with the rectum, the uterus, or the vagina.

#### Fungous Excrescences.

Sometimes fungous excrescences arise from the inner furface of the bladder, either in one mass, or in separate portions. When examined, they are found to confift of a loofe fibrous structure. When they are fituated a little behind the neck of the bladder, which is not uncommonly the case, they must produce a considerable obstruction to the passage of the urine. A stronger . action will, therefore, be required in the bladder to expel the urine, and its muscular coat will be confequently thickened: accordingly it is frequently found thickened; and it is not improbable that even where the fituation of the fungus may not obstruct the passage of the urine into the urethra, its presence may still irritate the bladder so as to excite it to more frequent and stronger actions than in a natural state, and the mufcular coat become thereby thickened.

# Elongations of the inner Membrane.

I have also known the inner membrane of the bladder elongated in some parts, so as to form irregular processes. These when cut into, were found to confift of a confiderable quantity of cellular membrane, intermixed with a little fat.—
The process producing such an appearance was probably a slow one, and was probably also attended with no pain. If these elongations were to be situated at a distance from the neck of the bladder, one can hardly perceive how they could produce any inconvenience, but if situated near the neck of the bladder, they might occasion extreme difficulty in making water, and even lay the foundation of a fatal disease.

### Muscular Coat Thickened.

One of the most ordinary changes in the bladder, from its natural structure, is the great thickening of its muscular coat. In a natural state, the muscular coat of the bladder (when it is moderately distended) consists of thin layers of muscular fibres, running in different directions.—

These are probably altogether, not more than the eighth of an inch in thickness. The muscular coat of the bladder, however, is occasionally found at least half an inch thick. This arises from an additional quantity of muscle being formed in consequence of extraordinary efforts being necessary in the bladder. The efforts take place

when there is any confiderable difficulty in making water, as happens when the proflate gland is a good deal enlarged, when there is a stone in the bladder, or when there are strictures of the urethra. It is usual, therefore, to find this thickening of the muscular coat of the bladder when there is any of these diseases. When the bladder is thickened, the fasciculi of which its muscular coat is composed become much larger, but never, or at least very feldom, acquire the full red colour which muscles have in other parts of the body. This is a deviation from the general plan of nature with regard to the increase of muscles from exercife. When muscles are enlarged in fize from exercise, they also become of a There is no other instance deep red colour. too in the body, as far as I recollect, of a muscle being fo much enlarged beyond its natural fize in confequence of increased exertion, as the muscular coat of the bladder. Between the fasciculi of the muscular fibres, little pouches are formed by the inner membrane. This arifes from the pressure of the urine against the inner membrane of the bladder which is impelled by the ftrong powers of the muscular coat. These pouches are often large enough to admit the end of the finger, and contain occasionally small calculi.-The bladder in this state admits of very little diftension, so that it is capable of containing little water; hence the inclination to make water is frequent, and frequent efforts of the muscular coat are required, which increase more and more its thickness. It is much more common to find this appearance of the bladder in the male than in the female, because in the latter there are fewer causes to produce it. There is in that sex a want of the prostate gland altogether, and the urethra being short and wide, obstructions seldom take place in it. When the muscular coat of the bladder is thickened, I believe that it has been sometimes mistaken for schirrus.

#### Calculi.

Calculi are not uncommonly found in the bladder, and are confined in their formation to no particular period of life. They are formed in very young children, and also in persons of middle and advanced age. This disease is not so frequently met with in the semale as in the male, which may depend on two causes; the one is, that there is not so strong a tendency to their formation in that sex; and the other cause is, that stones escape through the urethra in women, which would be detained in the bladder of men, and lay the soundation there of larger calculi.

The stones which are found in the bladder, are either originally formed in the kidneys, and pass through the ureters into the bladder, or they are first formed in the bladder itself. When the last circumstance takes place, the earthy matter is fometimes first deposited round some extraneous body, which becomes the nucleus of the calculus, although most frequently no nucleus whatever is to be observed. The nuclei which I have seen have been fmall portions of lead (probably broken off from a leaden bougie) fmall nails, and little maffes of hair. In fhort, any extraneous body which may happen to be introduced into the bladder, may become a nucleus. It is natural to think that fuch nuclei are more common in the calculi found in the bladder of women, than of men; their urethra is wider and shorter, so that an extraneous body can be much more eafily introduced into the bladder; and women too, from the particular frame of their minds, are more apt to make experiments of this fort than men.

The calculi of the bladder either lie loofe in it, or are confined to some fixed situation, from particular circumstances. When they are of a small size, they are sometimes lodged in pouches, or sacculi, formed by the protrusion of the inner membrane of the bladder between the fasciculi of its muscular sibres. A calculus also

is occasionally attached to an excrescence of the bladder, so as to be kept in a fixed situation.

There is frequently one calculus only in the bladder at a time, and then it is usually of an oval form; but there are often more, and the calculi by rubbing upon each other acquire flat fides and angles. Calculi have fometimes a fmooth uniform furface, but most frequently the furface is granulated. These granules are commonly placed very near each other over the whole furface of the calculus, giving it a certain degree of roughness. I have also seen them gathered into clusters on particular parts of the furface of a calculus. These granules are also fometimes of a fmaller, at other times of a larger fize, and in different calculi are more or less elevated. Some calculi have an irregular porous structure upon the surface, instead of being granulated.

Calculi when divided by the faw, or broken, exhibit most commonly a laminated structure. These laminæ are disposed in concentric curves, and are applied together with more or less compactness. They also differ in their thickness in different calculi: The laminated structure sometimes pervades uniformly the whole mass of the calculus; at other times different portions of

it are interrupted by a coarse porous texture. In some calculi no laminated structure whatever is observable, but it is entirely porous.

They are most frequently of a brown colour, which is sometimes of a lighter, and at other times of a darker shade. They are also sometimes of a white, and often of a yellowish colour. It is remarkable, that different portions of the same calculus are frequently of a different colour. Some laminæ, for instance, are perfectly white, while the other laminæ are brown. In this fort of mixture, I have most commonly found the white laminæ on the outside, and the brown laminæ in the middle; and I do not recollect seeing one instance of laminæ of different colours, disposed in alternate strata.

The specific gravity of urinary calculi differs wery considerably, as they differ a good deal in their compactness; but they are generally a little more than twice the specific gravity of water.

The chemical properties of different urinary calculi, I am apt to believe, differ more from each other than has generally been imagined. Any trials which I have made have led me to this opinion; but I am so little accustomed to chemical experiments that I do not rest upon it with much confidence.

The celebrated Scheele made a number of experiments to afcertain the nature of urinary calculi, of which the following are the chief.

- 1. Diluted vitriolic acid does not act on urinary calculi at all, but the concentrated dissolves them in heat.
- 2. Marine acid, whether diluted or concentrated, does not act on urinary calculi, not even when boiled with them.\*
- 3. Nitrous acid diluted, or aqua fortis, act a little on urinary calculi in the cold, but on the application of heat act with effervescence and with vapours.
- 4. This folution tinges the skin with deep red spots within half an hour after being applied, is not precipitated by alkalis, but on the application of lime-water yields a white precipitate.
- 5. Ponderous earth dissolved in marine acid, does not occasion any precipitate from this solution.
- 6. Acid of fugar, or falt of forrel, does not produce any precipitate from this folution.
- 7. A calculus when pounded and boiled in a folution of alkali of tartar, remains unchanged;
- \* I have met with an urinary calculus capable of being distolved in marine acid, even in the common temperature of the atmosphere.

but perfectly pure, or caustic alkali, such as does not shew the least mark of the ærial acid, dissolves the calculus even in the cold.

- 8. Lime-water dissolves the urinary calculus, by digestion, in the proportion of four ounces of lime-water to twelve grains of the calculus.
- 9. Pure water dissolves an urinary calculus, in the proportion of five ounces of the water to eight grains of calculus; and this solution changes tincture of lacmus to a red colour.
- 10. On distilling, in a small glass retort, one drachm of calculus in the open fire, there was obtained a volatile alkaline liquor like that of hartshorn, but no oil. In the neck of the retort there was a brown fublimate. Upon heating the retort thoroughly red hot, and then leaving it to cool, there was obtained a black coal, weighing twelve grains, which when put upon red hot iron in the open air, retained its black colour. The fublimate, which feemed to have been fomewhat fused, weighed twenty-eight grains; and upon being purified by a new fublimation, it grew white. It had no fmell, but a fomewhat fourish taste, and was eafily foluble in boiling water: it also dissolved in spirit of wine, but a larger quantity than of Lime-water water was requifite for this purpofe. was not precipitated. The fublimate feemed to agree in fome respects with the sal succini.

From these and some other experiments Scheele concludes, that urinary calculi are neither gypseous nor calcareous, but consist of an oily, dry, volatile acid concrete, mixed with some gelatinous matter.

Bergman, by burning the charcoal of urinary calculi to a white cinder, obtained some calcareous earth, and calculates that the proportion of it in these stones is about 1-200 part.

Although the world is much obliged to these two celebrated chemists, Scheele and Bergman, for their labours upon this as well as many other subjects, yet I am inclined to think that a sufficient number of urinary calculi has not yet been examined, to ascertain all the variety of their constituent parts.

Although the matter of calculus in the bladder is generally formed into one or more circumfcribed maffes, yet I have feen the whole bladder filled with a fubstance like mortar. This could not be removed entirely from it; but a great many small irregular portions still adhered to the sides of its cavity. I believe this matter to have been of the same nature with a common calculus, because it exhibited the same affinities with the common acids.

In opening dead bodies, the bladder is occasionally found to be very much distended, and domen. This might arise from some accidental circumstance of the water being accumulated, while the muscular coat of the bladder still possessed its proper powers; or the muscular coat of the bladder may have been paralytic, and therefore not capable of expelling the water. I do not think it is possible to discriminate between these two disferent situations by any examination after death, but they can always be ascertained by inquiry into the history of the case.

The bladder is also found contracted to such a degree as hardly to have any cavity. This is generally not to be considered as a disease, but simply as having arisen from a very strong action of the muscular coat of the bladder previously to death.

The anterior part of the bladder is occasionally found wanting, and instead of it there is a very soft vascular sless, situated externally at the lower part of the abdomen. This soft vascular sless is usually formed into irregular projecting masses, and in the living body is covered with a thick ropy mucus. The two ureters open somewhere upon this vascular sless, distilling gradually the urine as it is secreted upon its surface, which the mucus is intended to protect against the stimulus of that sluid. When there is such a formation of the bladder, I

believe that there is always found a deficiency of the bone at the fymphysis pubis, and also a monstrous formation of some of the organs of generation. This species of monstrosity I have described at large in the Medical and Chirurgical Transactions.\*

Another kind of monstrous formation in the bladder occasionally happens, viz. that at its depending part there is a communication between it and the rectum, the latter being continued into the former. Of this I have seen one instance, and it has been already taken notice of, when treating of the diseased and preternatural appearances of the intestines.

A portion of the bladder at its fundus, has been known to be lodged in a hernial fack, both at the abdominal ring and Paupart's ligament; but this is very rare, and has not fallen under my own obfervation.

<sup>\*</sup> See Medical and Chirurgical Transactions, p. 189.

See Pott on Ruptures, p. 226.

#### CHAP. XV.

Diseased Appearances of the Vesicula Seminales.

The diseased appearances of the vesiculæ seminales are not very commonly known, because from their situation these bodies cannot be seen without a good deal of dissection; whereas many of the viscera come immediately into view when the cavity is simply laid open in which they are lodged: diseased appearances, however, have been occasionally observed in them.

It has never occurred to me, to observe the vesiculæ seminales inslamed by themselves, although they are, no doubt, liable to this disease like every other part of the body. I have seen them, however, involved in the natural consequences of inslammation with the surrounding parts. Thus I have seen the posterior surface of the bladder, the vesiculæ seminales, and a portion of the rectum adhering with unusual sirmness toge-

ther, in the same manner as other parts of the body frequently do after inflammation.

The veficulæ feminales are also affected with scrofula. Upon one occasion I recollect to have feen one of the vesiculæ feminales filled with true scrofulous matter, the distinguishing characteristic of which has been often mentioned.

The ducts of the vesiculæ seminales open naturally by two distinct orifices into the cavity of the prostate gland, but they are occasionally wanting, and the vesiculæ seminales terminate in a cul-defac. The vasa deserentia are at the same time without their natural termination, for they end in the cul-de-sac of the vesiculæ seminales. This is a species of monstrosity which is very rare, but it is of great consequence, because it prevents the semen from passing into the urethra, and frustrates one of the most important functions in the animal economy.

The veficulæ feminales differ a good deal in their fize in different adult bodies, and indeed it is very common for the one to be confiderably fmaller than the other; but I have oftener-than once feen both of them fo small that they must have been very little able to fulfil the intentions for which they were formed.

One of the vesiculæ seminales is occasionally wanting altogether. Under such circumstances I should believe that the extremity of the vas deferens upon that side is most frequently enlarged and tortuous, becoming a sort of substitute for it. This was at least the case in the instance which I have seen of this mode of formation. The extremity of the vas deferens has at all times a structure similar to that of the vesiculæ seminales, and renders therefore this conjecture very probable.

The veficulæ feminales have also been observed to be schirrous; but this is not at all common.\*

<sup>\*</sup> See Morgagni, Epist. XLVI. Art. 5.

#### CHAP. XVI.

Diseased Appearances of the Prostate Gland.

Abscess in the Prostate Gland.

The prostate gland is not often found in a state of common inflammation. I have seen, however, an abscess in it, without any uncommon thickening and enlargement of the gland, and where the pus appeared to be of the common fort. This must be considered as being a common abscess, and must have been preceded by the ordinary fort of inflammation.

Scrofula of the Prostate Gland.

The proftate gland is fometimes fcrofulous. I have feen, in cutting into it, precifely the fame white curdly matter, which takes place in a fcrofulous abforbent gland. In fqueezing it also,

I have forced out from its ducts a ferofulous pus.

# Schirrus of the Prostate Gland.

The most common disease of the prostate gland is a schirrous enlargement of it. The prostate gland, it is well known, is naturally about the fize of a large chefnut, but when it is attacked by schirrus, it is often enlarged to the fize of the fift. In this enlarged state, the external appearance of its structure is not different from what is natural, but when cut into, it exhibits a very firm, whitish, or brown fubstance, with membranous feptarunning through it in various directions, which are often very ftrongly marked. This is the appearance of schirrus in other parts of the body. When the proftate gland is a good deal enlarged, its cavity becomes deeper from the growth of its fides, and the posterior extremity forms a confiderable projection into the cavity of the bladder, which interrupts the paffage of the urine into the urethra. According to the degree of this projection, the urine is paffed with greater or less difficulty, as well as an inftrument for drawing it off. When the projection is very great, it has fometimes been found impoffible to pass an instrument over the projection, and an artificial passage has occasionally been made through it accidentally, by which the urine has been evacuated. Under such circumstances the gland has been known not to be irritated by the violence used in making this new passage, and life has been prolonged for a greater length of time than it would have been otherwise. Still, however, the instrument ought to be made to pass over the projection, if possible; and we should never run the risk, by injuring the gland, of bringing on immediate satal consequences.

Sometimes in the progress of the enlargement the proftate gland grows irregularly, and a winding paffage is formed through it, by an alteration in the shape of its cavity. This increases the difficulty to the patient of making water, and to the furgeon of introducing an inftrument. When the proftate gland is enlarged, its internal furface is fometimes ulcerated, but commonly it is not. Fiftulous communications are fometimes formed between an enlarged prostate gland and the rectum. It is obvious too, from what has been mentioned, that in an enlarged state of the prostate gland, the difficulty of making water must be very great. This difficulty excites extraordinary and very frequent

efforts in the bladder to overcome it. Its mufcular coat becomes confequently much stronger and thickened. A prostate gland, therefore, is never found enlarged to any considerable degree, without the bladder having undergone this change in its muscular coat. This disease is hardly ever to be found in a young person, but is not at all uncommon at an advanced period of life.

# Calculi in the DuEs of the Prostate Gland.

There is another disease of the prostate gland, which occasionally takes place, although it is by no means so frequent as the former, viz. a formation of small calculi, which are lodged in its ducts. They are usually of the size of a small pea, and those which I have seen have been of a brown or black colour. What is their exact nature I cannot positively ascertain, because to do this requires a much nicer management of chemical experiments than I pretend to posses, but from some very impersect trials which I have made, they seemed to differ in their properties from urinary calculi.\* What I say, however, re-

<sup>\*</sup> They were almost entirely dissolved in the vitriolic acid, but were only separated into a fine powder by the nitrous.

lates only to the brown calculi; with regard to the other, I have had no opportunity of examining them.

# Ducts of the Prostate Gland enlarged.

The proftate gland is fometimes feen with its cavity very much widened, and its ducts enlarg-In the natural state of the gland, the orifices of its ducts can hardly be feen, but they fometimes are fo much enlarged as to be capable of admitting a crow quill. When the ducts are fo enlarged, there is always a great obstruction to the passage of the urine through the urethra, arising most commonly from stricture there. The urine either paffing in very fmall quantity, or being entirely obstructed, is accumulated in the cavity of the proftate gland and the bladder. The effect of this accumulation is, that the cavity of the proftate gland is widened, and the ducts very much enlarged. The bladder too, from making extraordinary efforts to overcome the obstruction, has its muscular coat gradually thickened, and often to a very confiderable degree. Attending, therefore, this state of the prostate gland there is a thickened bladder, and an obstructed urethra.

## The Prostate Gland preternaturally small.

I have also seen the prostate gland of an extremely small size, so that one could hardly suppose it from this circumstance to be fit for its office. It was attended with a monstrous formation of the urinary bladder and some of the organs of generation, and has been particularly described by me in the Medical and Chirurgical Transactions.

§ See p. 194

#### CHAP. XVII.

Diseased Appearances of the Urethra.

## Abscesses.

ABCESSES are occasionally formed in the membranous part of the urethra. These may arise from an inflammation, produced by fome latent cause, as abscesses are formed in any part of the body; but they happen most frequently from an obstruction to the passage of the urine through the urethra. This obstruction is produced generally by a stricture in some part of this canal, but most frequently it is a little anterior to the membranous portion of it. The urine being forced by the efforts of the bladder behind the structure, irritates that part, producing inflammation and suppuration; the abfcefs breaks externally, and the urine is evacuated by this opening. While it remains in the state of an abscess, it is not at all different from any common abfcesses which may communicate with the bladder.

#### Fistulæ.

While the obstruction in the urethra continues, the opening made by the breaking of the abscess is not allowed to close, but a fistulous orifice is gradually formed. This is surrounded with parts somewhat thickened and hard, as sistulæ are generally. The most common situation for these sistulous openings is behind the scrotum, near the membranous part of the urethra, because the most common situation of the stricture is a little anterior to this part of the urethra. Not uncommonly there are more than one of these openings leading to short canals which run in different directions.

The membranous part of the urethra I have also seen formed into a bag large enough to contain a hen's egg, and consisting of a pretty firm substance. This bag was occasioned by a large stone which had lodged there.

I do not recollect to have seen Cowper's glands diseased, which are situated near this part of the urethra. They must be liable to changes from disease, like other parts of the body; but they are small and difficult of access, so that they have very seldom become an object of examination.

Morgagni mentions one of them being converted into a ligamentous fubstance;\* and one of the excretory ducts in another instance being obliterated.

The inner membrane of the urethra is very liable to be inflamed, particularly at its anterior extremity, and the inflammation occasionally fpreads over the whole extent of the canal. This exhibits no appearance different from the inflammation of membranes lining fecretory canals which open externally. The inner membrane of the urethra is much more crowded with fmall blood vessels than in a natural state, and there is an increased secretion of the glands which open it. The inflammation is often not confined to the inner membrane of the urethra, but fpreads into the substance of the corpus spongiosum, affecting both its cellular structure and its glands. Under these circumstances, the corpus spongiosum is enlarged and harder from the extravafation of the coagulable lymph into its cells, and is more vascular than in a natural flate. The glands being increased in fize from the inflammation, become fenfible to the touch, like very fmall rounded tubercles.

<sup>\*</sup> See Morgagni, Epist. XLIV. Art. 3.

<sup>§</sup> See Morgagni, Epist. XLIV. Art. 12.

Ulcers are also seen occasionally in laying open the urethra, but these are not frequent.

#### Stricture.

The most ordinary diseased appearance of the urethra is stricture. This confists in a part of the canal being narrowed, or perhaps altogether obliterated. It may take place in any part of the canal, but it is most frequent a little anterior to the membranous part of the urethra. This stricture confifts fometimes of an approximation fimply of the opposite sides of the canal, so as to form a line of obstruction, and at other times the canal is narrowed for fome length. The inner membrane, at the stricture, fometimes exhibits the natural appearance of furface; at other times the furface is abraded or ulcerated. These effects are generally produced by bougies, and fometimes false paffages have been made into the corpus spongiosum urethræ in consequence of employing too much violence in the use of this instrument. There is very commonly more than one stricture in the fame urethra. It fometimes happens too, that the stricture is more on one fide of the canal than the other, fo that the passage there is crooked.

teneral determined. This devestion in the firme-

# busing don ei bas street en diem ein ben einen

A fmall fleshy excrescence sometimes grows in the urethra. This is called caruncle, and used formerly to be considered as the most common cause of obstruction in this canal, but since diffections of dead bodies have become more frequent, it has been found in reality to be very rare.

I have known one instance of a thin layer of earthy matter extending from the bladder through the whole length of the urethra.

### Preternatural Orifice of the Urethra.

The urethra fometimes does not open at the projecting extremity of the glans penis, but under it, where the frenum is naturally fituated; and in fuch cases there is no frenum. It consists of a small rounded opening, much less than when there is the natural termination in the glans. I have known an instance in this structure of parts, of a canal being formed besides the urethra, about two inches in length, which terminated at one extremity in a cul-de-sac, and at the other opened on the glans where the urethra commonly does. How far this variety may be frequent, I cannot pre-

tend to determine. This deviation in the structure is not to be considered as a disease, but simply as a mal-formation of parts, and is not attended with any material inconvenience, as far as I know.

There are some other diseased appearances of the penis, such as ulcers, phymoses, paraphymosis, &c. These are external, are very well known, and do not properly fall within my plan; I shall therefore omit them entirely.

#### CHAP. XVIII.

Diseased Appearances of the Testicles, and the Spermatic Chord.

#### Hydrocele.

Hydrocele, or a collection of water in the tunica vaginalis testis, is a very common complaint, and is confined to no particular period of life. It is not unfrequent in very young children, and in them most commonly disappears without any chirurgical treatment. In persons who are grown up, the disease scarcely ever goes away spontaneously, but requires the assistance of art. The bag in which the water is accumulated is of a pyramidal shape, and approaches more or less towards the ring of the abdominal muscle, according to the degree of the accumulation. It sometimes extends almost to the ring itself. The bag is also more or less thick in different cases: it is often scarcely thicker than the tunica vaginalis in its na-

tural state; at other times, when the accumulation is large, it is three or four times thicker, and is obviously laminated. In such cases too the testicle is a good deal compressed, and has sometimes been known to waste in consequence of this compression. The sluid which is accumulated is of a yellowish, a greenish, or brown colour, and resembles in its properties the serum of the blood. This disease, in persons who are advanced in life, is frequently combined with a schirrous state of the testicle, which will be afterwards particularly explained.

#### Hydatids.

Hydatids have fometimes been found in the cavity of the tunica vaginalis testis, either loofe or adhering; but I have had no opportunity of examining them, and they are not at all common.\*

#### Adhesions.

The tunica vaginalis is frequently found adhering to the furface of the testicle. The adherion is sometimes extended over the whole surface,

<sup>\*</sup> See Morgagni, Epist. IV. Art. 30.

but frequently confifts only of scattered processes of membrane. The adhesions are sometimes fine, but at other times have considerable thickness, and connect the tunica vaginalis to the body of the testicle more or less closely in different cases. They are produced by some previous inflammation, as in the cavity of the chest and belly; but it does not often occur that a tunica vaginalis is examined after death, which immediately before had been in an active state of inflammation.

## Testicle inflamed.

The substance of the testicle itself is very frequently inslamed, but this is commonly removed by art, and therefore hardly ever becomes an object of examination after death. It exhibits, however, precisely the same appearances as the inslammation of the substance of other parts, and therefore does not require to be particularly described. When the testicle is inslamed, the vas descrens sometimes partakes of the inslammation, its coats becoming considerably thickened, and in some instances the veins of the spermatic chord have been known to become varicose.\* After the inslammation of the testicose.

<sup>\*</sup> See Mr. Hunter on the Venereal Disease, p. 54.

and fullness of the epidydimis to remain for a confiderable length of time, or even through life. This depends on the matter, which had been extravalated during the inflammation, not being afterwards entirely absorbed.

Abfceffes too are not unfrequently formed in the testicles, from the progress of common instammation, and are attended with the same circumstances as abfceffes in other parts.

## Testicle scrofulous.

The testicle is sometimes completely changed from its natural structure, and converted into a mass of truly scrosulous substance. Upon such occasions it is generally enlarged in its size, and when cut into, shews a white, or yellowish white, curdly substance, which is sometimes more or less mixed with pus.

#### Testicle enlarged and pulpy.

The testicle is sometimes much enlarged in size, and converted into a brown, uniform, pulpy matter, in which its natural structure is entirely lost. This sort of change has been some-

times mistaken for schirrus, although it is very different from what is called schirrus in other parts of the body, and what is also found in the testicle itself.

# Schirrus and Cancer of the Testicle.

The testicle is often found much enlarged, where its natural structure is lost, and where it is changed into a hard mass of a brownish colour, which is generally more or less intersected by membrane. There is sometimes mixed in this structure cartilage; and sometimes there are cells formed in it containing a sanious sluid. This state of the testicle I consider as the true schirrus, and according to the progress of the disease, the epidydimis and the spermatic cord are more or less, or not at all, affected. This disease not unfrequently advances to form a foul deep ulcer, or throws out a sungus, and then it is called the true cancer of the testicle.

#### Testicle cartilaginous.

The testicle I have seen much enlarged in size, and changed into a mass of cartilage. This does not seem different in any essential property

from common cartilage, but is only a little fofter. This I should consider as depending upon the same general diseased process with the schirrus just described, for sometimes both structures are blended together in the same testicle.

#### Testicle bony.

The testicle is sometimes converted into bone. This is not a very unfrequent disease, and is more commonly confined to some part of the testicle, than extended over the whole of it.

I have seen a testicle with a small sirm cyst adhering to it, which contained a worm of that fort called vena medinensis. This is a worm of considerable length, with a smooth surface, and an uniform appearance; at the posterior extremity it terminates in a slender hook-like process, and at the anterior, there is a rounded opening or mouth. It very commonly burrows under the skin of the inhabitants of some warm countries, particularly Guinea, and is very troublesome to them. This testicle had probably belonged to a man who had visited some of those climates in which the vena medinensis is sound, and who had brought it over with him to this country.

# The Epidydimis ending in a Cul-de-Sac.

The testicles have sometimes this fort of malformation, that the epidydimis does not terminate in a vas deserens, but in a cul-de-sac. In these cases it is evident that the semen can never be evacuated by the urethra, and the person must therefore be incapable of procreation.

# Stricture of the Vas Deferens.

I have also seen a portion of the canal of the vas deferens obliterated by stricture. This had not been an original fault, but was the effect of a diseased process, and must have prevented the semen of that testicle from reaching the urethra.

## Testicles very fault and wasted.

The testicles are sometimes exceedingly small in their size. I have known one case, in a person of middle age, where each of them was not larger than the extremity of the singer of an adult. This, as appeared from its history, arose from a sault in the original formation, and was attended with a total want of the natural propensities. It is

much more common for a testicle to waste either spontaneously, or in consequence of a former inflammation, or compression, so as gradually to disappear entirely.\* When this takes place in one testicle only, the natural powers are preserved, but when it takes place in both, they must be altogether lost.

Sometimes one testicle, and sometimes both remain in the cavity of the abdomen through life, so that a person appears to have only one testicle, or to be without them altogether. The testicle or testicles, I believe, in these cases are of a small size, and Mr. Hunter suspects that they are by no means so persect as when they descend into the scrotum.

# Diseased Appearances of the Spermatic Chord.

The spermatic chord is also liable to diseased alterations of structure: one of the most common is that of its becoming schirrous. This I believe to be very rarely, if at all an original disease of the chord, but always, or almost always, spreads to it from the testicle. In the early state of a schir-

<sup>\*</sup> See Hunter on the Venereal Disease, p. 209.

<sup>§</sup> See Mr. Hunter's Observations on certain Parts of the Animal Economy, p. 18.

and this is the proper season for the extirpation of the testicle; but when the disease has taken place for a considerable time, and does not remain stationary, the chord becomes at length affected. Under such circumstances it is changed into a large hard mass, exhibiting the same appearance of structure with the testicle itself. During the last stage the disease advances to the loins, so as to affect the absorbent glands there.

When the testicle is scrofulous, the spermatic chord sometimes partakes of the same disease, and exhibits also the same appearance of change with the testicle itself.

A difease of the spermatic chord which is not uncommon, is an enlargement of its veins. The veins of the spermatic chord are numerous, and support a very long column of blood. This last circumstance, added to some impediments which occasionally take place to obstruct the return of the blood, renders the veins frequently enlarged. This enlargement varies very much in different cases, arising from the degree and the continuance of the impediment. When the enlargement of the veins is very considerable, they also become varicose, and the spermatic chord is changed into a bulky mass, soft to the feeling, and capable of being readily diminished upon pressure. In this

state of the spermatic chord, the testicle is sometimes wasted.

Water has fometimes been known to be accumulated in the cells of the cellular membrane, which envelopes the vessels of the spermatic chord. The cellular membrane of this part of the body is in confiderable quantity, and when water is accumulated in its cells, there is a large fwelling formed in the fituation of the spermatic chord, which is readily diminished upon pressure. When pressure is used, the swelling is diminished, not only by a part of the water being forced into the cells of the chord within the abdominal ring, but also by its being forced into the cellular membrane under the skin of the lower part of the belly. Many pints have been known to be accumulated in these cells. It has never occurred to myself to see this disease, and therefore I have had no opportunity of examining the nature of the fluid, but I presume it is of the same fort with what is usually found in anasarca.

A fack has also been known to be formed in the spermatic chord, consisting of a firm white membrane, and containing a sluid which most probably is of a serous nature. Both of these cases have been particularly described by Mr. Pott in his Treatise upon Hydrocele.\*

\* For the first case, see Pott on Hydrocele, p. 39. For the second see Pott on Hydrocele, p. 57.

#### CHAP. XIX.

Diseased Appearances in the Female Organs.

Inflammation of the Uterus.

When the uterus is inflamed, it takes place almost always under the same circumstances, viz. very soon after parturition. The inflammation is sometimes consided to the uterus itself, or its appendages, but the peritonæum in the neighbourhood is most commonly affected, and frequently over the whole extent of the cavity of the abdomen. The uterus when inflamed, exhibits the same appearances as the inflammation of the substance of other parts, and these are principally observable in its body or fundus. The inflammation is frequently sound to creep along the appendages of the uterus, especially the Fallopian tubes and ovaria. It often advances to suppuration, and the pus is generally sound in the large veins of

the womb.\* When the peritonæum is also affected by inflammation, it exhibits the same appearances which we formerly described particularly, when treating of the inflammation of this membrane; but the extravasated sluid, and the coagulable lymph, are not uncommonly in very large proportion to the degree of the inflammation.

## Schirrus of the Uterus.

One of the most frequent diseases of the uterus is schirrus. In this state it is increased more or less in bulk, and often to a very large size. Its substance when cut into is thick and hard, and when its structure is examined, it shews a whitish sirm substance, intersected generally by strong membranous divisions. This is the common appearance of the structure of schirrus in other parts, and it differs less from the natural appearance of the structure of the uterus than that of any other part of the body. In this state of the uterus, its internal surface is commonly found

<sup>\*</sup> See Dr. Clark's Effays, p. 69 and 70.

<sup>§</sup> Dr. Clarke, who has examined a great many women that have died after parturition with inflammation of the peritonæum, has observed this particularly. See Dr. Clarke's Eslays, p. 136.

ulcerated, forming cancer, and sometimes so much so as to throw out long ragged processes. The ulceration also sometimes extends to the neighbouring parts, as the vagina, the bladder, and the rectum, making communications between them, and producing dreadful havock. The uterus is occasionally almost entirely destroyed by the progress of the disease. I have seen several instances where the fundus of the uterus only was remaining, and where the rest was changed into a tattered ulcerated mass.

It is worthy of remark, that schirrus or cancer attacks generally the cervix uteri before the sundus. This may perhaps depend upon a general principle, which we have often mentioned: the cervix uteri is more glandular than any other part of the uterus, and schirrus or cancer appears to be peculiarly a disease of glandular parts.

# Tubercles of the Uterus.

Hard tubercles often grow from the uterus, which are either imbedded in its substance, or arise from its outer surface. They vary a good deal in their size, viz. from that of a hazel nut to more than the size of the sist. They are irregular in their shape, but are commonly rounded, and are

in some degree tuberculated. These when cut into, shew a whitish very firm substance, intersected by membranous septa, which are commonly very thick and strong. It is extremely rare that those circumscribed masses are sound ulcerated. The uterus in this state of disease is generally of the natural size.

A mass of the same kind is sometimes sound in the cavity of the uterus, and often grows to a very large size. I have seen it a good deal larger than a child's head at birth. This mass when cut into, exhibits precisely the same appearances which we have so lately described. It is remarkable, that those masses within the cavity of the uterus commonly do not adhere in any part closely to it, but are connected with it loosely, by the intervention of the cellular membrane, so that they can be very easily peeled off, without injuring the structure of the uterus. The uterus itself is more or less enlarged according to the bulk of the mass it contains, but appears to be persectly healthy in its structure.

#### Polypus.

Polypus forms a very common disease of the uterus, and may take place almost at any period

of life; it is more frequent, however, at the middle or advanced age, and very rarely happens in persons who are young. By a polypus is meant a diseased mass, which adheres to some part of the cavity of the uterus, by a fort of neck or narrower portion. It is of different kinds: the most common kind is hard, and confifts evidently of a white fubstance, divided by very thick membranous fepta. When cut into, it shews precisely the fame structure with the tubercle of the uterus lately described; so that a person looking upon a fection of the one and the other, out of the body, could not at all diftinguish between them. This fort of polypus varies very much in its fize, some being not larger than a walnut, and others being larger than a child's head. It adheres by a narrower portion or neck, which varies a great deal in its fize, and in its proportion to the body of the polypus. The largest polypus I ever saw was fuspended by a neck hardly thicker than the thumb; and I have feen a polypus, much less than the fift, adhering by a neck fully as thick as the wrift.

The place of adhesion also differs considerably. It is most commonly at the fundus uteri, but it may take place in any other part; and I have seen a small polypus adhering just on the inner part of the lip of the os uteri. When a polypus

is of any confiderable fize, there is generally but one; but I have occasionally seen on the infide of the uterus, two or three fmall polypi. Another fort of polypus takes place in the uterus, which confifts of a bulky, irregular, bloody mass, with a number of tattered processes hanging from it. This when cut into exhibits two different appearances of structure: the one appearance is that of a spongy mass, consisting of laminæ, with fmall interstitial cavities between them; the other is that of a very loofe texture, confifting of large irregular cavities. These are the different varieties in polypi, which I have observed; but perhaps there may be others, which have not come under my notice. It is very obvious, that in proportion as a polypus grows, the cavity of the uterus must be enlarged, and the same change must take place in the vagina, when a polypus leaves the uterus, and paffes into this canal.

# The Inversion of the Uterus.

The invertion of the uterus occasionally takes place, and principally from two causes, viz. from the weight of a polypus, or from violent pulling in attempts to remove the placenta. When the invertion is incomplete, the fundus uteri forms

a tumour within its cavity; there is at the same time an appearance of fissure upon the outside of the uterus, where the fundus usually is; and the Fallopian tubes, round ligaments, and ligaments of the ovaria, are drawn inwards at both edges of the fissure. The uterus, particularly after labour, is sometimes inverted entirely, the inner surface being exposed, and the sundus uteri forming a large tumour on the outside of the vulva.

# Prolapsus Uteri.

The uterus fometimes leaves its natural fituation and falls downwards, so as either to get to the external parts, or out of the body entirely. This is most apt to happen when women have a large pelvis, and where the soft parts have been very much relaxed by repeated and severe labours. This disease is called prolapsus uteri, and will be explained more particularly when we come to treat of the diseases of the vagina. It is much more frequent than the other disease called the inversio uteri.

# Stricture in the Cavity of the Uterus.

A stricture is fometimes formed within the cavity of the uterus, fo that its cavity at one part is obliterated entirely. This I believe almost always to take place at one part, viz. where the cavity of the fundus uteri terminates, and that of the cervix begins, for in this place the cavity of the uterus is narrowest. As the fides of the cavity round this place lie very near each other, and form naturally a small aperture, it is probable that fome flight inflammation may unite the parts together, and shut up the aperture; or the parts may gradually approach together without this cause, as in strictures of the urethra. uteri has been found to be fo contracted, as to have its passage in a great measure obliterated;† and it has even been known to be closed up, by the growth of an adventitious membrane. I

#### Uterus bony.

The uterus has fometimes its substance more or less converted into bone. This arises from a par-

<sup>+</sup> Vid. Morgagni, Epist. LXVII. Art. 11.

<sup>‡</sup> Vid. Morgagni, Epist. XLVI. Art. 17.

ticular morbid action of its blood vessels, by which they secrete from the blood bony matter, and is a very rare disease.

The Uterus changed into an earthy Substance.

The uterus has also been known to be converted into an earthy substance. It is probably of the same kind with the earth of bones, and this disease probably differs only from the former, in there being a less proportion of animal gluten, to combine the earthy particles together.

A bony Mass in the Cavity of the Uterus.

In the cavity of the uterus a bony mass is sometimes found. When this is the case, I suspect that the hard sleshy tubercle within the cavity of the uterus, such as we lately described, had been converted into bone. This at least had taken place in the only instance which I have known of it (for a great part of the tubercle had still remained unchanged) and I think it very probable, that it most frequently happens where such bony tumours are found.

|| Vid. Lieutaud, Tom. 1. p. 323-

#### Stones in the Cavity of the Uterus.

Stones\* have fometimes been found in the cavity of the uterus. These are described by authors as varying in their appearance, some being of a dark, and others of a light colour. About their nature they are silent, and I can say nothing of it from my own knowledge, as it has never occurred to me to see an instance of this disease. I should believe, that these concretions are formed from matter thrown out by the small arteries which open upon the internal surface of the uterus, and are in some degree analogous to the concretions which are formed in some glands of the body.

It has also been known to happen, that a dead fectus has remained for a long time in the cavity of the uterus, and has there been gradually changed into an earthy mass preserving the shape of the child. §

Water in the Cavity of the Uterus.

Water has fometimes been known to be accumulated in the cavity of the uterus in very large

<sup>\*</sup> Vid. Lientaud, p. 339.

See Chefelden's Anatomy of the Bones, plate LVI.

quantity.+ In some cases fifty, fixty, or even a hundred pints, have been faid to be accumulated. This water is fometimes bloody in its appearance, and at other times is of a yellowish colour. Of its nature I cannot speak particularly, as I have never feen an instance of this disease. I should believe, however, from analogy, that the water accumulated in the cavity of the uterus, refembles in its properties the ferum; and I should believe alfo, from the same ground of conjecture, that it is poured out by the finall curling arteries of the uterus. In cases where water is really accumulated in the cavity of the uterus, one must suppose a stricture of the cervix, otherwise the water would escape gradually into the vagina as it is formed. I am apt to believe, however, that where water has been faid to be accumulated in the cavity of the uterus, it has frequently been really in large hydatids formed in that cavity.

#### Hydatids in the Uterus.

Large maffes of hydatids have also been found in the cavity of the uterus. Whether these be of the same kind with what occasionally grow in the

<sup>+</sup> Vid. Lieutaud, Tom. 1, p. 319, p. 333.

<sup>‡</sup> Vid. Lieutaud, Tom. 1, p. 335.

placenta, or like those in the other parts of the body, I cannot determine, as it has not occurred to me to fee an example of this difease. The hydatids of the placenta are a good deal different from those of the liver, kidneys, and some other parts of the body. They confift of vehicles of a round or oval shape, with a narrow stalk to each, by which they adhere on the outfide of one another. Some of those hydatids are as large as a walnut, and others as fmall as a pin's head. A large hydatid has generally a number of fmall hydatids adhering to it by narrow processes. Of their real nature nothing is known, but they are not improbably animals. There is in quadrupeds a difference in hydatids, and this is even the case in the same species of quadruped; and yet these have been determined to be animals. I should believe that the hydatids said to be found in the uterus, have not uncommonly been only hydatids of the placenta, which had been retained there.

#### Rupture of the Uterus.

These are the various diseased appearances which are well ascertained to take place in the uterus. I have to add, that the womb is not un-

frequently ruptured, which is rather to be confidered as an accident than a difeafe. This, perhaps, never takes place but in the pregnant uterus, and at the time of delivery. It may arife either from too violent an action of the muscular fibres of the uterus upon the child, or upon the hand of an accoucheur, who may for some reason or other have introduced it into its cavity, and pressed upon some part of the uterus with a good deal of force. The ruptures which I have feen have been commonly in the fide of the womb, and of confiderable extent. The - peritonæum covering the womb is often not ruptured, and there is a large mass of black coagulated blood lying between it and the uterus, where the rupture has taken place. This black appearance is occasionally mistaken for mortification.

#### Two Uteri.

It has fometimes happened, although very rarely, that two uteri have been formed in the fame perfon instead of one. In this case there is but one ovarium and one Fallopian tube to each. The vagina is at the same time divided by a septum into two canals, each of which conducts to its pro-

per uterus. In the case which is described in the Philosophical Transactions, a communication was formed at one part through the septum; but how far this generally takes place in such a kind of monstrosity I cannot determine.

The uterus varies a good deal in its fize in different perfons, in fome being fully twice as large as it is in others. It differs also somewhat in the thickness of its substance. There is some difference too in its situation, being often placed much nearer one side of the pelvis than the other. All of these are to be considered as varieties in the natural formation, and not as disease.

|| See Philosophical Transactions, Vol. 64, p. 474.

It has fomerance happened, although very rare,

perfou inflend of one, in this case there is but one to aring and one I allowed take to pech. The

erquire of the dealer distributed to the relation of

#### CHAP. XX.

Diseased Appearances of the Ovaria.

Inflammation of the Peritonæal Covering of the Ovaria.

That portion of the peritonæum which covers the ovaria I believe is feldom inflamed, unless where the inflammation has spread to it from the uterus, or where it has attacked this membrane generally. It is not unusual, however, for it to be inflamed under either of these circumstances, and it shews the same appearances as the inflammation of the peritonæum covering any other part. Adhesions too are frequently found, joining the ovaria to the neighbouring parts, which had been the consequence of such an inflammation.

Inflammation of the Substance of the Ovaria.

Where the uterus has been inflamed to a confiderable degree, as after parturition, the fubstance of the ovaria has also been occasionally attacked by the inflammation spreading to it. The ovaria are then enlarged in size, are harder than in a natural state, and are highly vascular; and very commonly pus is found to have been formed.

### Schirrus of the Ovaria.

Schirrus is a disease which sometimes attacks the ovaria, although very seldom in comparison of its attacking the uterus. Under such circumstances the ovaria become enlarged, and are converted into a whitish hard mass, which is more or less intersected with membranous septa. These schirrous masses have sometimes a disposition to be converted into bone; and in this way most frequently, I believe, the ovaria become bony. The bony substance into which they are converted has sometimes a greater admixture of earth than the natural bones of the body.

The ovaria are fometimes very much enlarged in fize, and converted into a uniform, brown, pulpy matter. Cells are at the fame time formed in fome part of it which contain a fluid.

I have also seen the ovaria partly changed into a scrosulous matter, intermixed with cells.

# Dropfy of the Ovaria

The most common disease of the ovaria is dropsy. The whole substance of an ovarium is sometimes destroyed, and it is converted into a capsule containing a sluid. These capsules are not unfrequently of a very large size. They consist of a white sirm membrane, and contain an aqueous sluid, capable of being partly coagulated.

When the substance of the ovaria is destroyed, and they become dropfical, it is very common for them to be converted into a number of cells which communicate with each other by confiderable openings, and to be prodigiously enlarged in their fize. An ovarium in this case may be fo enlarged as to occupy almost the whole cavity of the abdomen. The ovaria are also sometimes converted into a congeries of complete cyfts. These vary a good deal in their fize, some being not larger than a hazel nut, and others as large as an orange. Their coats are fometimes thin, at other times are of considerable thickness, and confift of a compact, white, laminated membrane. They contain either a serous fluid, with which I have feen fome flimy matter mixed, or a thick ropy fluid, or a kind of jelly; and, what

one would not expect to find a priori, different cyfts in the same ovarium will sometimes contain a different fort of sluid.

Thefe cyfts, I believe, have been occasionally confounded with hydatids, to which they bear fome refemblance. They are however really very different. They have much firmer and less pulpy coats than hydatids; they contain a different kind of fluid, and they are differently connected among themselves. Hydatids either lie loose with regard to any connection among each other, or they inclose each other in a series; or small hydatids adhere to the coats of larger ones. Cysts of the ovarium adhere to each other laterally by pretty broad furfaces, do not envelope each other in a feries, and appear to have no power analogous to generation like hydatids, by which fmaller cyfts are formed, that are attached to those of a larger fize. It appears not improbable, that these cysts are formed by a gradual enlargement of the small veficles which make a part of the natural structure of the ovaria.

The Ovaria changed into a fatty Substance with Hair and Teeth.

The ovaria are sometimes converted into a fatty substance, intermixed with long hair and teeth,

which is furrounded by a capfule confifting of a white strong membrane. The hairs are most of them loofe in the fatty fubstance, but many of them also adhere on the inside of the capsule. Teeth too are formed, but are generally incomplete, the fangs being wanting. These sometimes arise immediately from the inner membrane of the capfule, and are fometimes connected with an irregular mass of bone. Such productions have been commonly confidered as very imperfect impregnations, but there is good reason to believe that they can take place without any intercourse between the fexes. I have described a case, which has been published in the Philosophical Transactions, where it was hardly possible that impregnation could have happened. The girl in whom this change of the ovarium was found, could not from all appearances be more than twelve or thirteen years old; the hymen was perfect; and the uterus had not received that increase of bulk which is usual at puberty. The other marks of puberty were also wanting. these circumstances I should judge the womb to be incapable of the stimulus of impregnation. A tumour, confifting of teeth and hair, was preferved by the celebrated Ruysch\* in his collection, which he fays was found in a man's stomach. If this be

<sup>\*</sup> Vid. Ruysch, Tom. 2, Adversar. Anatomicor. Decad. Tert.

true (which there feems to be no reason to doubt) it puts my conjecture beyond dispute. This production could not possibly, under such circumstances, have any connection with impregnation; and if it is produced without it in one part of the body, there can be no good reason why it may not also take place without impregnation in another part. These productions are much more frequent in the ovaria than any where else, probably because the process which forms them bears some analogy to generation, in which the ovaria are materially concerned. I must still therefore, whatever objections have been made to it, retain my former opinion. These masses in the ovaria are commonly about the fize of a large orange.\*

#### A Fatus in the Ovarium.

A fœtus is fometimes found in the ovarium.— This feldom arrives at the full fize, but its formation as far as it goes is commonly perfect. When

\*I have very lately met with the same kind of fatty substance intermixed with hair, and the body of one tooth covered with enamel, in the ovarium of a young woman about eighteen years of age. In this case the uterus was rather less than its usual size in the adult when unimpregnated, and there was no membrana decidua whatever formed in its cavity. It appeared, therefore, to be under-



minished to half their natural fize, are somewhat tuberculated on their surface, and are very hard. When cut into, the vesicles which make a part of their natural structure, are sound to be filled with a white solid matter.

An Ovarium wanting.

An ovarium on one fide has been known to be wanting, but this is extremely rare.

on the lame changes, althoughting manifeld, which it does in the needings on continuations of program cyange Tru factoristic well-alse are also enterpress or der to supply a fulficient quantity of blood to the owns which is growing in the ownium.

Surinking of the Oxonia.

ed are obanged in finite thurstone in They are dig

coing no change health in which happens to hen there is a same action when the health plant in the continue of the succeiment.

ing remarkably finall. Their circumfinaces do not a-

of the state of the state of the truth of the truth of the

#### CHAP. XXI.

Diseased Appearances of the Fallopian Tubes.

Inflammation of the Fallopian Tubes.

When the uterus is inflamed to a confiderable degree, the inflammation not uncommonly fpreads along the Fallopian tubes: they become highly vascular, and when cut open, sometimes contain blood in their cavities. The inflammation may even advance to suppuration, and their cavities will be found loaded with pus.

# Adhesions.

When the peritonæum generally, or some part of it, in the neighbourhood of the Fallopian tubes, is inslamed, the external covering of these tubes, which is a continuation of the peritonæum, also partakes of the inslammation. This, when it subsides, generally terminates in adhesions of the Fallopian tubes to the contiguous parts. It is not

unufual to find the fimbriated extremity of the Fallopian tubes adhering to the ovaria, or when the previous inflammation has been confiderable, to find the fimbriated appearance entirely loft, and the body of the Fallopian tube feems to terminate on the furface of the ovarium. Under fuch circumftances there is no aperture towards this end of the Fallopian tube, and it has loft its power of conveying the ovum from the ovarium to the uterus.

The Fallopian tube communicates by a small aperture with the cavity of the uterus. This aperture is sometimes obliterated, but not so often as the aperture of that extremity towards the ovarium.

## Dropfy of the Fallopian Tubes.

When the Fallopian tube has its aperture closed at both extremities, it is sometimes dilated into a confiderable tortuous cavity. This when laid open appears occasionally subdivided by small partial septa, and contains an aqueous sluid, which is capable of being partly coagulated. This sluid is undoubtedly supplied by the secretion of the small arteries belonging to the inner membrane of the Fallopian tube, which is naturally very vascular. It may be called dropsy of the Fallopian tube.

The Fallopian Tubes terminating in a Cul-de-fac.

The Fallopian tubes I have feen without any aperture or fimbriated extremity, from a defect in the original formation, and terminating in a cul-defac. Under fuch circumstances they were incapable of performing their office as subservient to generation.

#### An Ovum in the Fallopian Tube.

The Fallopian tube is fometimes dilated into a bag containing an ovum. This arifes from the ovum being stopped in its progress from the ovarium to the uterus. When it is so stopped it does not die, but is gradually evolved as if it had been lodged in the cavity of the uterus. This, among many others, is a proof that the uterus is not the only organ which is sitted to evolve an ovum, but that other parts of the body can perform this office. While the ovum is enlarging, the Fallopian tube is more and more dilated, forming a thin bag round the ovum. The blood vessels passing to the ovarium and the Fallopian tube where the ovum is contained, are gradually enlarged, in pro-

portion to the increase of the ovum, in order to fupply it with a fufficient quantity of blood. While this process is going on in the Fallopian tube, the uterus increases in bulk so as to be even twice its natural fize, and becomes more vafcular. The cavity of its fundus is also lined by a membrana decidua, and the cervix uteri is plugged up with jelly. The uterus therefore undergoes a variety of changes, exactly fimilar to what take place in natural pregnancy, being thrown into this progrefs of action from the original stimulus of impregnation. The ovum fometimes makes confiderable progrefs in the Fallopian tube, and even has been known to advance to the full period of gestation, but more commonly it dies at an early period. In the course of the evolution of the ovum, the Fallopian tube has been known to rupture, and the person to die from internal hæmorrhage. A very clear and accurate account of fuch a cafe has been published by Dr. Clarke in the Medical and Chirurgical Transactions-page 216.

Hard Tumour growing from a Fallopian Tube.

I have seen a hard round tumour growing from the outer surface of one of the Fallopian tubes. This when cut into exhibited precisely the same appearance of structure with the tubercle which grows from the surface of the uterus, viz. it confisted of a hard white substance, which was interfected with strong membranous septa. This, however, I believe to be a very rare appearance of disease.

The round ligaments partake of the inflammation of the uterus, when it is confiderable, and has fpread to its appendages. They are also, doubtless, subject to other diseases, but these are very rare, and have not fallen under my own observation, nor do I know of their having been particularly noticed by authors.

beth known to make place in the vaging which

· not beneft in beginster a granuous discomo promi

#### CHAP. XXII.

Diseased Appearances of the Vagina.

Inflammation of the Vagina.

THE internal furface of the vagina, near the outward opening, is frequently inflamed, especially from the application of the venereal poison, but this hardly ever becomes the subject of examination after death.

## Adhesions of the Sides of the Vagina.

A very violent inflammation has fometimes been known to take place in the vagina, which has terminated in the adhesion of the sides of the cavity. This adhesion is sometimes extended over a great part of the cavity, but most frequently is more confined, producing a stricture in some one part. The vagina under such circumstances loses its office as a canal subservient to the uterus; and according to the degree of the extent of the adhe-

## [ 221 ]

fion, the disease becomes more or less easy to be remedied by art.

# Ulcers of the Vagina.

Ulcerations are not unufual in the vagina.—
They fometimes appear like spots of the internal furface, removed as it were by a knife, and at other times there is a foul ragged ulcer. When this last is the case in any considerable degree, the ulcer has commonly not originated in the vagina, but has spread from the womb, which is in a cancerous state. When the ulcer spreads very much, communications are often made with the neighbouring parts, producing a most miserable state of existence. Thus communications are sometimes formed between the vagina and the rectum, or the vagina and the bladder.

# Schirrous Tumours in the Vagina.

Schirrous tumours occasionally arise in the vagina itself (although, I believe, rarely) when the uterus is unaffected. When cut into they exhibit the true schirrous structure which has been often described. Such tumours may ulcerate, and produce the fame dreadful havock which we have fo lately mentioned.

## Inversion of the Vagina.

One of the most common diseases of the vagina is its invertion, or prolapfus: this is more apt to happen where the natural formation of the pelvis is large, where the external opening at the vulva is wide, and where the parts are generally relaxed. The prolapfus is more or less in different cases; in some the uterus does not pass out at the external parts, and in others the inversion of the vagina is complete, at the extremity of which is fituated the os uteri. The protrufion has then different shapes; it sometimes forms a large rounded mass, and at other times it is narrowed and more elongated, extending, perhaps, five inches from the body. When this last has been the case, it has been fometimes mistaken for that species of monstrous formation called hermaphrodite. We may here take an opportunity of mentioning, that although in fome of the common quadrupeds a real hermaphrodite structure has been found, yet it has never been discovered in the human subject. When the vagina has been long in the habit of being inverted, its inner furface becomes in many



in it: these are chiefly polypi; and when they have been removed by art, the vagina, if it has not been for a long time stretched, recovers nearly its natural size.

#### The Vagina very narrow.

The vagina has occasionally been found to be very much contracted with regard to transverse diameter, from a defect in the original formation. This, however, occurs very rarely.

#### CHAP. XXIII.

Diseased or preternatural Appearances of the external Parts.

## The Hymen imperforated.

The hymen is sometimes sound without a perforation in it, so that the vagina is completely shut up at its external extremity. This is an original mal-formation, which is frequently not discovered till the age of puberty, when the menstrual blood is accumulated behind it. It is of little consequence, as it can be easily remedied by art.

## The Clitoris enlarged.

An enlarged clitoris is also a natural defect, less common than the other, but a more unfortunate one. At birth, the clitoris in such a case is often larger than the penis of a male child of the same age. It has a well formed prepuce and glans,

together with a fiffure at its extremity, fo as to refemble almost exactly the external appearance of the male organs. These cases have given rise to a mistake, with regard to the fex, and females have been often baptized for males. On most occafions, however, where there is an enlarged clitoris, the fex may be determined by the following circumstances. The labia are well formed, and when handled, no round bodies are felt in them, like the tefficles. The fiffure at the extremity of the glans does not lead to any canal of the urethra, but under the glans, and at the posterior extremity of the fiffure, there is an opening which leads immediately to the bladder. I should believe, that by putting a fmall straight probe into this orifice, and passing it into the bladder, it could be at once determined on most occasions, whether the child was male or female. If the child should live to grow up, the clitoris enlarges, but, I believe, not in the same proportion as the penis would do. It is a most unfortunate monstrosity, because it depresses the mind, by a consciousness of imperfect formation in a very important part of the body. Such cases have been often miltaken for hermaphrodites.

a birt operation beautiful or a middle of the

#### The Nymphæ enlarged.

The nymphæ are not unufually enlarged beyond their natural fize. This fometimes happens to one only, and at other times to both. When the nymphæ are very much enlarged, they pass confiderably beyond the furface of the body, and have the same fort of covering with the labia, losing by their exposure the sine, vascular, sensible covering of natural nymphæ. This is a monstrous formation of no great consequence, unless the nymphæ be very large, and even then they can be extirpated by art.

## The external Labia growing together.

The two external labia are sometimes united together by a fine line of junction, at the upper end of which are situated the mentus urinæ, and the head of the clitoris. This fort of monstrous formation is not at all common, and is very easily remedied. When the external labia are separated by a slight operation, all the parts behind are found perfect.

The two labia are fometimes joined together by a continuation of the common fkin, fo that the appearance of labia is loft entirely. This defect may also be remedied by art, and the parts within will be found to be well formed.

The external parts, particularly the infide of the nymphæ, and the vestibulum, are subject to inflammation and ulcers from common causes, and especially from the application of the venereal poison. These diseases, although they are very often the subject of solicitude during life, yet are seldom examined after death, and therefore we shall omit them here altogether.

ate appearant to winer it incindenced, as he

#### CHAP. XXIV.

mater during a flate of in

would be with that of any other partielle he

Diseased Appearances of the Brain and its Membranes.

Inflammation of the Dura Mater.

The dura mater is sometimes sound in a state of inflammation. When this is the case, there are to be seen in the inflamed portion of it many extremely sine vessels, filled with florid blood, which are passing between the dura mater and the cranium. These sine vessels are seldom so crowded as in many other parts of the body when inflamed, which arises from the nature of the membrane itsels. In its natural state there are sew blood vessels ramifying through it, and therefore when it is inflamed, it does not appear so much crowded with vessels as other parts do which are naturally more vascular. Still, however, a person well acquainted with the natural appearance of the dura mater, would be as much struck with the differ-

ence of its appearance when it is inflamed, as he would be with that of any other part of the body.

The dura mater during a state of inflammation sometimes forms a layer of coagulable lymph, which adheres upon its inner surface like an adventitious membrane; but this is very uncommon. On such occasions, although the inflammation of the dura mater may subside, yet the membrane will remain, and may become the cause of a fatal disease.

When the dura mater is inflamed, adhesions are fometimes formed between it and the other membranes of the brain, so that for a considerable extent of surface they adhere together; but this appearance of disease is also very rare.

It is not uncommon when the dura mater has been inflamed, especially in consequence of some external violence, that suppuration takes place, and pus is sound covering a portion of the membrane.

The dura mater is likewise sometimes eroded by ulceration, but this is by no means frequent: it is more common, in violent injuries of the heads for a portion of it to become mortified. Scrofulous Tumours connected with the Dura Mater.

Scrofulous tumours are fometimes formed which are connected with the dura mater, but this happens very rarely. These resemble precisely the structure of a scrofulous absorbent gland, and occasionally there is to be found in them a curdly pus.

Spongy Tumours growing from the Dura Mater.

Spongy tumours also grow from the dura mater, but they are very uncommon. Such tumours, as far as I have had an opportunity of examining them, are pulpy to the feeling, and of a distinct fibrous structure.

Bony Matter formed in the Dura Mater.

One of the most common diseased appearances of the dura mater is the formation of a bony laminæ in some part of it. These are usually very small, being not larger than the nail of a singer, but they are also occasionally of a much larger

fize. They are thin, and frequently very irregular in their edge. They are not to be found indifferently in every part of the dura mater, but are almost always adhering at the superior longitudinal sinus, or its falciform process. In some of them the proportion of the earth to the animal part is larger than the common bone.

There is often but one of these offisications; at other times there are more of them. The falciform process has been said to be occasionally found almost entirely converted into bone; but this last appearance is very rare.

Very strong Adhesion of the Dura Mater to the Cranium.

There is at all times a strong adhesion between the dura mater and the inside of the cranium. This adhesion is principally formed by small blood vessels which pass from the one to the other, and likewise by a close application of the sibrous structure of the membrane to the bone. In a natural state, however, the dura mater can be perfectly separated from the cranium; yet it sometimes happens that the adhesion is so strong, as to render it impossible to separate the two completely. The dura mater in such an attempt is

torn in different parts into two laminæ, one of which adheres to the bone, and the other lies upon the pia mater. Whether this preternatural strength of adhesion arises from a previous state of inslammation in the dura mater, or from some other cause, I cannot determine; but it is not at all an uncommon appearance.

Diseased Appearances of the Tunica Archnoides.

Diseased appearances of structure are very rare in the tunica archnoides, and have almost been entirely overlooked by writers. The only diseased appearance of this coat which I have observed, is that of its becoming a good deal thicker than it is naturally, fo as to be a pretty firm membrane. In this, as well as in its natural state, there are to be feen no blood veffels ramifying upon it; or at least they are extremely few. It is also separated at fome distance from the pia mater, a gelatinous fluid being interposed between the one and the other. This diseased appearance does not in general take place equally over the whole furface of the brain, but is to be found chiefly on the upper part of the two hemispheres. It is not an uncommon appearance of difease, particularly after fovers where the brain has been a good deal affected.

Diseased Appearances of the Pia Mater.

Veins of the Pia Mater turgid with Blood.

The most common diseased appearance of the pia mater is that of its veins being turgid with blood. This depends upon some impediment to the free return of the blood from the head towards the heart, which may arise from a variety of causes, and is very different in its appearance from an inflamed state of the pia mater. The smaller branches of its arteries, filled with a florid blood, are not more numerous in this state than is natural, but its veins are much more distended with a dark blood.

## The Pia Mater inflamed.

When the pia mater is inflamed, it is upon the whole more difficult to distinguish it from its natural appearance than any other part of the body. This depends upon the great number of very small vessels which ramify upon it in its healthy state. When the pia mater is inflamed, these small vessels are much more numerous than in a healthy state,

are filled with a florid blood, and form by their anastomosis a beautiful net-work. It does not frequently occur, when the pia mater is inflamed, that it becomes so uniformly red as to shew no interstices between its vessels, a circumstance which happens in the inflammation of some other parts. The processes arising from the under surface of the pia mater are also more crowded with vessels than usual, and there is a stronger adhesion between them and the substance of the brain.

It very rarely happens that any layer of coagulable lymph is formed in the inflammation of the pia mater, which is fo very common in inflammation of the pleura and the peritonæum. When the pia mater is inflamed to a high degree, pus is formed. I have feen pus which had been formed during an inflammation of the pia mater, and which was effused over the whole upper furface of the brain.

Close adhesions, for some considerable extent of surface, have been seen between the pia and dura mater, which are probably the consequence of inflammation; but these are very rare, and have not fallen under my observation.

Scrofulous Tumours adhering to the Pia Mater.

I have feen a number of fcrofulous tumours adhering upon the infide of the pia mater. They exhibited the true scrosulous structure, which has been often explained. This diseased appearance is very uncommon.

It is not unufual to find fome of the veffels of the pia mater filled with air. This may be generated by putrefaction, but it is also fometimes seen when no process of this kind appears to have taken place. Under such circumstances, it is probable that air is extricated by some new arrangement in the particles of the blood, somewhat analogous to the change in secretion.

## Hydatids.

Little cyfts,\* containing water (which are generally called hydatids) have been feen adhering to the pia mater; but this is a very rare appearance of difease.

Diseased Appearances in the Substance of the Brain.

## Inflammation.

The substance of the brain, in which I include both the cerebrum and cerebellum, is liable to inslammation, although it is not very common, when no external injury has been applied to the head.

<sup>\*</sup> Vid. Lieutaud, Tom. 2, p. 145.

When inflammation takes place, it is rarely extended over any large portion of the brain, but is rather confined to one or more distinct spots. In this state of disease the inflamed portion becomes of a red colour, although this is seldom very intense. When cut into the colour is found to arise from a great many small vessels, which are silled with blood. If the inflamed portion be upon the surface of the brain, the membranes in the neighbourhood are also commonly inflamed. The part which is inflamed has no peculiar hardness, but yields nearly the same sensations to the touch, as it would do in a healthy state.

# Abfcesses.

Inflammation of the brain frequently advances to suppuration, and abscesses are formed in it. When these are of a large size, the weight of the pus breaks down the structure of the neighbouring parts, and they look simply as if they had been destroyed, or very much injured by the pressure. When the abscesses are small, there is an ulcerated appearance of the cavity in which the pus is contained.

## Gangrene.

Portions of the brain occasionally become gangrenous, especially after violent injuries of the head; but I believe this appearance of disease is extremely rare, where an inflammation of the brain has taken place from any other cause. I have met, however, with one instance of this; a portion of the brain at the instance of a very dark brown colour, and as soft as the most rotten pear.

# The brain very foft.

It is extremely common, when a brain is examined in a person who has been dead for several days, to find its substance so soft that it can hardly be cut so as to leave a smooth surface, and the smallest pressure of the singers breaks it down into a pultaceous mass. The brain, however, will sometimes retain, for several days, the sirmness and resistance which it had during life; yet this is by no means common. Neither of these appearances is to be considered as produced by disease.

## The Brain very firm.

The brain is fometimes found to be confiderably firmer than in a healthy state, to be tougher, and to have some degree of elasticity; it will

Mr. Hunter is the only person whom I have heard to remark this property of elasticity in the brain of maniacs. It was very remarkable in the only case of this kind which I have had an opportunity of examining. bear to be pulled out with some force, and will re-act so as to restore itself, or when pressed will recover its former shape. Under such circumstances the ventricles are sometimes found enlarged in size, and full of water. The brain has even been said to become so hard and dry as to be friable between the singers; and the medullary substance in these cases, is often much lighter than in a natural state. It has been remarked that the cerebellum is very often unaffected. When these changes take place in the brain, the mind is at the same time deranged: there is either mania or lethargy; or the person is much subject to convulsive paroxysms.

A white firm Substance formed in the Brain.

It is not a very unufual appearance of difease in the brain, to see a part of it changed into a white substance, of an uniform smooth texture, and with a considerable degree of sirmness. Around this substance the brain is frequently a little inslamed. Such a substance is generally considered as scrosulous, and it has certainly somewhat that appearance, but it is a good deal sirmer than a scrosulous absorbent gland, even where pus has not at all been formed. There are frequently more than one of these substances formed in the brain at a time. It is also not unusual to find round masses

# [ 240 ]

of the same fort of substance, lying as it were imbedded in the brain; some of these I have seen as large as a walnut.

## Encysted Tumours.

Encysted tumours containing a serous sluid thave sometimes been found in the substance of the brain; but they have never come under my own observation, and are very uncommon.

# Fungus and Schirrous Tumours.

Fungus ‡ and schirrous | tumours are also described by authors, as occasionally growing in the brain; but these too are extremely rare. After the operation of the trepan, it is not very uncommon for a fungus tumour to grow from the brain, which is sometimes of a very large size; but what is its particular structure I am unacquainted.

## Hydrocephalus.

One of the most common appearances of disease in the brain, is the accumulation of water in its ventricles: this generally takes place when a child is very young, and even sometimes before

<sup>†</sup> Vid. Lieutaud, Tom. 2, p. 194, 195.

<sup>‡</sup> Vid. Lieutaud, Tom. 2, p. 197.

<sup>|</sup> Vid. Lieutand, Tom. 2, p. 198, 199.

birth. The water is accumulated in greater or less quantity in different cases. It sometimes amounts only to a few ounces, and occasionally to many pints. The water is of a purer colour, and more limpid, than what is found in dropfy of the thorax or abdomen. It appears, however, to be of the fame nature with the water that is accumulated in both of those large cavities. In some trials which I have made, it partly coagulated upon the application of the common acids, exactly like the water in hydrothorax and afcites, or like the ferum of the blood. This is not what one would expect a priori; it is natural to think, that as the water which is accumulated in the ventricles of the brain in a healthy state does not contain any coagulable matter at all, or at least in a very small proportion, there should be the same property belonging to the water when it is accumulated in a larger quantity, fo as to form a disease. This, however, does not necessarily follow, because the water in hydrocephalus may not only be supposed to be increased in quantity beyond what is confistent with health, but also to be altered in its properties. How far this is always the case in the water of hydrocephalus, I cannot positively determine. We have no means likewife of afcertaining whether this makes a part of a general law, with regard to the accumulation of water in the other circumferibed cavities of the body, or whether it is an exception to it. In the cavities of the abdomen, thorax, &c. there is just enough of moisture, in the healthy state, to lubricate the surfaces of parts, and it cannot become an object of chemical examination, till it is accumulated beyond the healthy proportion, so as to form a disease. We can make no comparative trials, therefore, between the one sluid and the other.

When water is accumulated in the ventricles to a very large quantity, the fubstance of the brain, especially upon the fides and at the upper surface, appears almost to be a fort of pulpy bag, containing a fluid. The fcull too upon occasions is very much enlarged in fize, and altered in its shape. The cranium is exceedingly large in proportion to the fize of the face. The projections are very confiderable at the centres of offification, from whence the frontal, parietal, and occipital bones were originally formed, and the membranous divisions between these several bones are very wide. When the scalp is removed, so as to give an opportunity of looking immediately upon the cranium, the bones are found to be very thin, often not thicker than a shilling, and there are frequently broad fpots of membrane in the bone. The reason of this last appearance is, that offification takes place in many points of the membrane in fuch cases in

order to make a quicker progress, but the water accumulates too rapidly for it, so that spots of membrane are left not converted into bone. When such appearances take place in hydrocephalus, the disease has been of long continuance, occasionally for some years.

Water is also sometimes formed under the pia mater, and upon the surface of the brain, but very rarely in any considerable quantity. There is, generally, at the same time greater quantity than natural in the ventricles.

Water is likewise found in small quantity between the dura and pia mater.

It is related by authors, that water has been formed occasionally between the dura mater and the cranium.\* From the nature of the adhesion between the cranium and this membrane one would not easily be led to suspect an accumulation of water between them, and such cases are at least to be considered as very uncommon.

# Blood effused or extravasated.

Blood is frequently found effused within the cavity of the cranium in various fituations. It may either be poured out by the rupture of some vessel into the substance of the brain itself, or into some

¶ Vide Lieutaud, Tom. 2, p. 229, 230.

of the ventricles. It is frequently effused upon the furface of the brain under the pia mater, and likewise between the dura and the pia mater. This is most apt to happen where the effusion is in confequence of external violence. The quantity of blood which is effused from the rupture of vessels in the brain is frequently very confiderable. It is commonly found in a coagulated state, and the texture of the brain in the neighbourhood is often very much hurt from the pressure. When blood is extravalated within the cavity of the cranium from fome external injury, the vascular system is usually found, except for the rupture which may have happened. But when blood is extravafated within the cavity of the cranium where there has been no external injury, the vascular system of the brain will be almost always found diseased. It is very common in examining the brain of perfons who are confiderably advanced in life, to find the trunks of the internal carotid arteries upon the fide of the fella turcica very much difeafed, and this difease extends frequently more or less into the finall branches. The difease confists in a bony or earthy matter being deposited in the coats of the arteries, by which they lofe a part of their contractile and distensile powers, as well as of their tenacity. The fame fort of diseased structure is likewife found in the bafillary artery and its branches. The veffels of the brain, under fuch circumflances of difeafe, are much more liable to be ruptured than in a healthy flate. Whenever blood is
accumulated in unufual quantity, or the circulation is going on in them with unufual vigour, they
are liable to this accident, and accordingly in either of thefe flates ruptures frequently happen.
Were the internal carotid arteries and the bafillary
artery not fubject to the difeafed alteration of flructure which we have defcribed, effusions of blood
within the cavity of the cranium, where there has
been no previous external injury, would be very
rare.

# Difeased Appearances of the Plexus Choroides.

The plexus choroides feems to be subject to some diseases to which the vascular system of the brain generally is very little or not at all exposed.

Its vessels become sometimes a good deal enlarged and varicose.

Little bags refembling hydatids fometimes adhere to it. These are commonly small, viz. not larger than a garden pea, but sometimes they have been seen as large as a gooseberry.

Small fcrofulous tumours fometimes adhere to the plexus choroides; but this is a much rarer appearance of difease than the former.

## Earthy Matter in the Pineal Gland.

There is but one fort of disease to which the pineal gland is much exposed, and this is very frequent: it consists in a deposition of earthy matter in its substance. It very rarely happens that a pineal gland is entirely free from earth, but it is occasionally almost entirely converted into it. There is then a very thin covering of the natural substance of the pineal gland surrounding an earthy mass, and the mass itself is always very easily divisible into small particles, by pressure between the fingers.

The pineal gland has been mentioned by some authors as being sometimes schirrous. I have felt it on some occasions a little sirmer than on others; but it has never occured to me to observe that alteration of structure in it which could be properly called schirrous, and I believe it to be a very rare disease. The pineal gland has been sound to be very much distended with a limpid water;\* but this too is very uncommon.

That particular substance called the glandula pituitaria is liable to very few diseases; I have never seen in it any decided or obvious alteration from its natural structure. It is somewhat firmer in one

<sup>\*</sup> Vid. Morgagni, Epist. LXII. Art. 15.

person than another, but this never struck me to be disease.

It rarely happens that any of the nerves within the cavity of the cranium appear diseased. I have, however, sometimes seen a nerve a good deal smaller in its size than it ought to be, softer in its texture, and of a less opaque colour; this I recollect to have been particularly the case with one of the optic nerves in a person who was blind of one eye.

The nerves vary a good deal in their fize in different persons, as a part of their original formation, without there being any disease whatever.

These are the principal diseased changes which take place in the brain, and its appendages. I have just to add, that the brain is subject to great variety from original monstrous formation. A great part of what is usually called the cerebrum is sometimes wanting, while the cerebellum, and the medulla spinalis are entire; sometimes there is hardly any vestige of either the cerebrum or cerebellum, and the medulla spinalis is very much diminished in size; at other times there is a total want of the brain, and there is no appearance of the medulla spinalis. In this case one should expect a want of nerves through the whole body. It is, however, not so; nerves are found distributed in the common way, through the limbs, and the

dorsal nerves can be seen arising from a membrane somewhat resembling the dura mater in the canal behind the vertebræ. When there is a total want of brain, it sometimes happens that there is a medulla spinalis, which, however, is of a very diminished size. In cases of desiciency in the brain, the cranium is nearly upon a level with the two eyes, and there is often upon the scalp a soft spongy excrescence. This is generally divided into distinct protuberant masses, and is covered with a sine skin, capable of being rendered very vascular by injection. When cut into, the spongy excrescence consists of pretty large cells, which are filled with a fort of grumous matter.

There is also frequently, instead of this excrescence, a bag growing from the skin of the scalp, and covering more or less the back of the trunk. This bag sometimes consists of a sine membrane, with little strength; at other times it is pretty thick, and has very considerable sirmness. It sometimes communicates with the cavity of the cranium by a considerable opening; and at other times the communication is very small. It is filled with an aqueous sluid, and in some instances there is also in it a portion of the brain.

FINIS.

## CHAPTER I.

PAGE

Diseased Appearance of the Pericardium

Ser.

Inflammation of the Pericardium—Adhefions of the Pericardium to the Heart—
Dropfy of the Pericardium—Scrofulous
Tumours growing from the Pericardium—The Pericardium very dry.

#### CHAPTER II.

Diseased Appearances of the Heart

O

Instammation of the Heart—Abscess—Gangrene—White Spot upon the Heart—Polypus—Aneurysm of the Heart—Aneurysm of the Arch of the Aorta—The semilunar Valves bony and earthy—Semilunar Valves thickened—A semilunar Valve burst—Valve between the Auricles and Ventricles bony and earthy—Valve between the Auricles and Ventricles and Ventricles thickened—Rupture of the Heart—Blood effused into the

Cavity of the Pericardium without Rupture—Mal-formation of the Heart— Heart unufually large—Hydatids adhering to the Heart—A portion of the Heart bony—A portion of the Heart earthy.

#### CHAPTER III.

Diseased Appearances in the Cavity of the Thorax

25

Inflammation of the Pleura—Adhesions— Empyema—Hydrothorax—Ossification of the Pleura.

### CHAPTER IV.

Diseased Appearances of the Lungs

33

Inflammation—Abscesses—Common Tubercles—Large brown Tubercle—Lungs incapable of collapsing—Cells of the Lungs
very large—Lungs changed into a Substance like the Liver—Lungs bony—Lungs
compressed by a large porous Tumour—
Earthy Concretions in the Lungs—Hydatids in the Lungs.

#### CHAPTER V.

Diseased Appearances in the Posterior Mediastinum.

43

Inflammation of the inner Membrane of the

Trachea—Polypus—A Portion of the Trachea Schirrous—Rings of the Trachea bony—Ulcer of the Trachea.

Esophagus lined with a layer of Coagulable Lymph—Spasmodic Stricture—Stricture arising from a narrow puckering of the inner Membrane of the Esophagus —Stricture attended with a cancerous Ulcer—Portion of the Esophagus cartilaginous—Fungus of the Esophagus— A Pouch formed by the Pharynx lying behind the Esophagus.

The descending Aorta aneurysmal—The Vena Azygos enlarged and varicose—The Vena Azygos ruptured—The Thoracic Dust enlarged and varicose—The Thoracic Dust obstructed by earthy Matter—Rupture of the Thoracic Dust.

The Absorbent Gland Scrofulous—Schirrous—Converted into an earthy and bony Matter—Some diseased Appearances of the Anterior Mediastinum.

#### CHAPTER VI.

Diseased Appearances within the Cavity of the Abdomen.

Ascites—Inflammation of the Peritonaum
—Adhesions—A white soft granulated

Matter adhering to the Peritonaum—Cancerous Tumours growing from the Peritonaum—Hydatids in the Cavity of the Abdomen—Air.

#### CHAPTER VII.

Diseased Appearances of the Stomach

67

Inflammation—Ulcers—Schirrus and Cancer—A Pouch formed in the Stomach—
The Orifice at the Pylorus extremely contracted—The whole of the Stomach very much contracted—The Stomach distended with Air—A Portion of the Coats of the Stomach dissolved by the Gastric Juice—
Tumour of a suetty Substance in the Stomach—Calculi in the Stomach—Papillæ—Small-pox Pushules said to have been observed in it.

#### CHAPTER VIII.

Diseased Appearances in the Intestines

77

Inflammation—Ulcer—Mortification—Intus Sufceptio—Ruptures—Schirrus and Cancer—The great Intestine thickened, and its inner Membrane formed into white and yellowish Tubercles—Folds of the inner Membrane of the great Intestine enlarged and loaded with Blood—Little

Processes of the inner Membrane a little above the Anus—Piles—Fistulæ in Ano—
The Rectum terminating in a Cul-de-sac—
The Rectum terminating in the Bladder—
Worms—Lumbricus Teres—Tænia—Ascaris—Air in the Intestines—A Ring formed in the Intestine by one of the Valvulæ Conniventes being much enlarged—
Calculous Matter in the Intestines—Small-pox Pustules said to have been observed.

Mefentery inflamed—Abscess—Gangrene—
Absorbent Glands of the Mesentery Scrofulous—Cancerous—Filled with bony and
earthy Matter—Hydatids adhering to the
Mesentery—Fatty Tumours adhering to the
Mesentery.

#### CHAPTER IX.

Diseased Appearances of the Liver

107

Inflammation of the Membrane of the Liver—Adhesions—Inflammation of the Substance of the Liver—Abscess—Mortification—Common Tubercle of the Liver—
Large white Tubercle—Soft brown Tubercle—Scrofulous Tubercle—Liver very
foft—Liver very hard—Hydatids—A Cyst
in the Liver containing earthy Matter—

Worms in the Cyst of the Liver, and in the Biliary Ducts.

#### CHAPTER X.

Difeased Appearances in the Gall-bladder

Inflammation of its Coats-Adhefion-Coats of the Gall-bladder schirrous—Coats of the Gall-bladder bony-Difeafed State of the Ducts-Gall-stones-Bile-Gall-bladder much distended-Much contracted-Hydatids in the Gall-badder.

#### CHAPTER XI.

Difeased Appearances of the Spleen

131

Inflammation of the Coats of the Spleen-Adhesions-Inflammation of the Substance of the Spleen-The Spleen extremely foft-The Spleen very hard-Tubercles in the Spleen-Coats of the Spleen cartilaginous -Spleen very large-Stony Concretions in the Spleen-Spleen wanting.

#### CHAPTER XII.

Diseased Appearances of the Pancreas 138

Pancreas hard-Calculi of the Pancreas-Abscesses-Worms in its Ducts-Pancreas wanting.

#### CHAPTER XIII.

Diseased Appearances of the Kidneys, and the Renal Capsules 1.

141

Abscesses of the Kidneys—Kidneys schirrous

—Kidneys very soft—Hydatids of the
Kidneys—Calculi of the Kidneys—Kidney
converted into an earthy Substance—
Converted into a bony Substance—Kidneys much subject to variety in Situation
and other natural Circumstances without
Disease—A Kidney on one side wanting.
Renal Capsules containing a black sluid—
Containing Pus—Little granules of Stone
in them.

### CHAPTER XIV.

Diseased Appearances of the Bladder

152

Inflammation of the Peritonæat Covering—
Inflammation of the inner Membrane—
Ulcers—Schirrus and Cancer—Fungous
Excrescences—Elongations of the inner
Membrane—Muscular Coat thickened—
Calculi—Bladder filled with a Substance
like Mortar—Bladder much distended—
Bladder much contracted—Anterior part
of the Bladder wanting—Bladder and
Rectum communicating with each other



the Urethra-Stricture-Caruncle-Preternatural Orifice of the Urethra.

#### CHAPTER XVIII.

Diseased Appearances of the Testicles, and the Spermatic Chord 183

Hydrocele—Hydatids—Adhesions—Testicle
inflamed—Testicle scrosulous—Testicle enlarged and pulpy—Schirrus and Cancer
of the Testicle—Testicle cartilaginous—
Testicle bony—Empidydimis ending in
a Cul-de-sac—Stricture of the Vas Deserens—Testicles very small and wasted—
One and sometimes both Testicles in the
Cavity of the Abdomen—The Spermatic
Chord Schirrous—Scrosulous—Its veins
varicose—Water in the Cellular Membrane of the Spermatic Chord—Water in
a Cyst of the Spermatic Chord.

#### CHAPTER XIX.

Diseased Appearances in the Female Organs
193

Inflammation of the Uterus—Schirrus of the Uterus—Tubercles of the Uterus—Polypus—Inversion of the Uterus—Prolapfus Uteri—Stricture in the Cavity of the Uterus—Uterus bony—Uterus chang-

ed into an earthy Substance—A bony Mass in the Cavity of the Uterus—Stones in the Cavity of the Uterus—Water in the Cavity of the Uterus—Hydatids in the Uterus—Rupture of the Uterus—Two Uteri—A Variety in the natural Appearances of the Uterus without Disease.

#### CHAPTER XX.

Difeased Appearances of the Ovaria

207

Inflammation of the Peritonæal Covering of
the Ovaria—Inflammation of the Subflance of the Ovaria—Schirrus of the
Ovaria—Dropfy of the Ovaria—The Ovaria converted into a fatty Substance
with Hair and Teeth—A Fætus in the
Ovarium—Shrinking of the Ovaria—An
Ovarium wanting.

#### CHAPTER XXI.

Difeased Appearances of the Fallopian Tubes 215

Inflammation of the Fallopian Tubes—Adhesions—Dropsy of the Fallopian Tubes—
The Fallopian Tubes terminating in a Culde-sac—An Ovum in the Fallopian Tube
—Hard Tumour growing from the Fallopian Tube.



rachnoides-Veins of the Pia Mater turgid with Blood-The Pia Mater inflamed-Adhesions-Scrofulous Tumours adhering to the Pia Mater-Some of its Blood Veffels containing Air-Hydatids-Inflammation of the Substance of the Brain-Abscesses-Gangrene-The Brain very foft-The Brain very firm-A white Subflance formed in the Brain-Encysted Tumours-Fungous and Schirrous Tumours -Hydrocephalus-Blood effufed or extravafated-Blood Veffels of the Brain Difeafed-Diseased Appearances of the Plexus Choroides-Earthy Matter in the Pineal Gland-Water in it-Pituitary Gland very little subject to Disease-Diseased Appearances of Nerves-Mal-formation of the Brain.

ERRATUM.—Page 233, 8th and 10th lines from the top, instead of Tunica Archnoides, read Tunica Arachnoides.

Districted Absorbing of the Turing A.

Primingline of the Dura Merevandele

but Trungs of action and the Pers



