

**A short description of the human muscles, chiefly as they appear on dissection. Together with their several uses, and the synonyma of the best authors / By John Innes.**

### **Contributors**

Innes, John, 1739-1777

Monro, Alexander, 1733-1817

### **Publication/Creation**

London : Printed by E. Hodson ... for J. Murray ... and J. Cuthell ..., 1791.

### **Persistent URL**

<https://wellcomecollection.org/works/zsyz6w4h>


### **License and attribution**

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection  
183 Euston Road  
London NW1 2BE UK  
T +44 (0)20 7611 8722  
E [library@wellcomecollection.org](mailto:library@wellcomecollection.org)  
<https://wellcomecollection.org>





A.E.3.

29,967/A

B. 48. d. 34 -

~~A  
a~~

✓

1

2

4

(A.  
a.2.)











8354.3

*Alexander Lyon Emerson*

**MEDICAL STAFF**

SHORT DESCRIPTION  
**LIBRARY.**

OF THE

HUMAN MUSCLES,

CHIEFLY

AS THEY APPEAR ON DISSECTION;

TOGETHER WITH

THEIR SEVERAL USES, AND THE SYNONYMA  
OF THE BEST AUTHORS.

---

BY JOHN INNES.

---

A NEW EDITION.

Greatly improved by ALEX. MONRO, M. D.

---

L O N D O N :

PRINTED BY E. HODSON, BELL-YARD,  
FOR J. MURRAY, NO. 32, FLEET-STREET; AND  
J. CUTHELL, NO. 4, MIDDLE-ROW, HOLBORN.

M.DCC.XCI.



TO  
ALEXANDER MONRO, M.D.

PROFESSOR OF ANATOMY AND MEDICINE  
IN THE UNIVERSITY OF EDINBURGH.



HAVING been highly solicited  
for several years past by many of  
your Pupils to publish a short Descrip-  
tion of the Muscles as they appear up-  
on Dissection of the Human Body. I  
have at last ventured to comply with  
their requests. With anxiety to pro-  
mote the Science of Anatomy, and to  
encourage every thing that may be  
useful to the Gentleman who attend  
your Theatre, was the principal mo-  
tive which induced me to undertake  
this task. I have no knowledge of the  
subject but what I derived from you.  
It, therefore, this Treatise, which you  
have



MEDICAL STAFF.

LIBRARY.

TO

ALEXANDER MONRO, M. D.

PROFESSOR OF ANATOMY and MEDICINE  
in the University of Edinburgh.

S I R,

**H**AVING been repeatedly solicited, for several years past, by many of your Pupils, to publish a short Description of the Muscles, as they appear upon Dissection of the Human Body, I have at last ventured to comply with their request. Your anxiety to promote the Science of Anatomy, and to encourage every thing that may be useful to the Gentlemen who attend your Theatre, was the principal motive which induced me to undertake this task. I have no knowledge of the subject but what I derived from you. If, therefore, this Treatise, which you

A 2

have



have never seen till I now present it to you, should communicate any advantage to the Gentlemen under your care, it is to you alone they are indebted for the obligation. I am,

S I R,

With respect,

Gratitude, and esteem,

Your much obliged,

And very humble servant,

JOHN INNES.

EDINBURGH,

1776.



---

## P R E F A C E.

**S**EVERAL full and accurate descriptions of the Muscles have already been published. But their size and prolixity have rendered them of less value to the dissector than the small Treatise of Dr. Douglas, which was first published about the beginning of this century, and, since that time, has undergone various impressions, without receiving any improvement, excepting the addition of the synonyma from Albinus. It is therefore presumed, that a simple and concise description of the muscles, which should contain all the improvements of the moderns, is still wanting.

To class the muscles according to their uses, may do very well in a large work, or in describing their compound actions: But this method can never answer the purposes of dissection. To remedy this inconvenience, the muscles in the following Treatise are described chiefly as they appear in dissecting the human body.



The describing of the muscles according to their origins and insertions prevents much circumlocution. This is the method pursued by Dr. Douglas; and wherever his descriptions seemed tolerably accurate, they have been followed with little alteration. But Dr. Douglas's book is peculiarly defective with regard to the muscles of the abdomen, back, and neck: in describing these, therefore, the method of Albinus has been preferred.

Those who have not opportunity, or are averse from undergoing the labour of dissecting, may derive considerable advantages from comparing the descriptions now given, with the beautiful and correct tables of Albinus; and, to facilitate still more the study of these intricate organs, I have caused to be published eight of Albinus's tables, with concise explanations, on a small scale, adapted for the pocket.

For the benefit of those who wish to examine the history of the muscles more minutely, the synonyma of the best authors are added; and, for the sake of brevity, the compound action of the muscles, and the origin and insertion of several inconsiderable fibres are omitted.

The

The reader will observe, that, in general, the muscles of one side only are described; because all the muscles of the body, with very few exceptions, have correspondent ones on the opposite side.





A

SHORT ACCOUNT OF THE AUTHOR,  
AND OF HIS WRITINGS.

MR. JOHN INNES, Author of the following Treatise, was born at *Callart*, an obscure village in the Highlands of Scotland. He came to Edinburgh at an early period of life, where he obtained the patronage and protection of Dr. Alex. Monro, Professor of Anatomy; who instructed him in the knowledge of the human body, and in the art of dissection. When about the age of eighteen years, Dr. Monro appointed him dissector to the Anatomical theatre. The functions of this important and difficult office he continued to perform with much reputation for near twenty years.

But his abilities were not confined to the dexterity of dissecting the most minute parts of the human frame. He described the various organs with ease and with perspicuity. This happy talent attracted the notice of the students; and, at their solicitation, and by the approbation of Dr. Monro, he opened an evening course of anatomical demonstrations.

The



The number of pupils who annually attended these demonstrations afforded the best evidence of his abilities, and of the advantages derived from his labours. During his last course, he was attended by near two hundred students.

For some time before his death, he was troubled with an affection of the lungs, which terminated in a phthisis pulmonalis, and proved fatal to him on the 12th of January 1777.

On the 15th of the same month, the following account of him appeared in the public papers:

“Mr. JOHN INNES, at an early period of life, had been educated in the dissecting art: He made a rapid progress in his profession; and his genius and industry were rewarded with the privilege of giving private lectures for his own emolument. The utility of his lessons was soon perceived. Numbers of students resorted to him for instruction; and all of them acknowledged the advantages they had received. At that stage of life when men are most capable of benefiting themselves, and of being useful to the public, death hurried him out of the world. He has given two small specimens of what was to be expected from his anatomical skill. The year before  
his



his death, he published a short description of the Human Muscles as they appear on dissection, together with their several uses, and the synonyma of the best authors. The merit of this work was universally acknowledged. Some months after, he published, as a *vade mecum* for students, eight anatomical tables, containing the principal parts of the skeleton and muscles represented in the large tables of Albinus, with accurate explanations. These are all the monuments he has left by which the public is to judge of his ability. To his numerous friends and acquaintance, it is unnecessary to mention the warmth of his heart, or the integrity of his disposition."



## ADVERTISEMENT.

**D**URING the illness of which Mr. INNES died, he put into my hands the first edition of his Description of the Muscles, with a few, chiefly verbal corrections of it.

ON perusing that Work, lately, at the request of the Bookseller, I have found it necessary to make a very considerable number of alterations in what relates to the description, as well as to the uses, of the Muscles.

ALEX. MONRO.

EDINBURGH,  
September 5, 1777.

---

A

DESCRIPTION

OF THE

HUMAN MUSCLES.

---

CHAP. I.

MUSCLES *of the* TEGUMENTS *of the*  
CRANIUM.

THE skin that covers the cranium is moved by a single broad digastric muscle, and one small pair.

1. OCCIPITO FRONTALIS,

*Arises* fleshy from the transverse protuberant ridge near the middle of the os

B

occipitis



occipitis laterally, where it joins with the temporal bone; and tendinous from the rest of that ridge backwards, opposite to the lateral sinus; it arises after the same manner on the other side: From thence it comes straight forwards, by a broad thin tendon, which covers the upper part of the cranium at each side, as low down as the attollens aurem, to which it is connected, as also to the zygoma, and covers a part of the aponeurosis of the temporal muscle; when it comes as far forwards as near the hair of the front, it becomes fleshy, and descends with straight fibres.

*Inserted* into the orbicularis palpebrarum of each side, and into the skin of the eye-brows, sending down a fleshy slip between them, as far as the compressor naris and levator labii superioris aëque nasi.

*Use.* Pulls the skin of the head backwards; raises the eye-brows upwards; and, at the same time, it draws up and wrinkles the skin of the fore-head.

*Epi-*



*Epicranius*, Albinus.

*Frontalis et occipitalis*, Winflow.

## 2. CORRUGATOR SUPERCILII,

*Arises* fleshy from the internal angular process of the os frontis, above the joining of the os nasi, and nasal process of the superior maxillary bone; from thence it runs outwards, and a little upwards.

*Inserted* into the inner and inferior fleshy part of the occipito-frontalis muscle, where it joins with the orbicularis palpebrarum, and extends outwards as far as the middle of the superciliary ridge.

*Use.* To draw the eye-brow of that side towards the other, and make it project over the inner canthus of the eye: When both act, they pull down the skin of the forehead, and make it wrinkle, particularly between the eye-brows.

*Musculus supercilii*, Winflow.

*Musculus frontalis verus, seu Corrugator*,  
Douglas.



## C H A P. II.

*Of the MUSCLES of the EAR.*

THE muscles of the ear may be divided into three classes, viz. the common, proper, and internal. The common may move the whole ear; the proper only affect the particular parts to which they are connected; and the internal, the small bones within the tympanum.

The common muscles are,

## 1. ATTOLLENS AUREM,

*Arises* thin, broad, and tendinous, from the tendon of the occipito-frontalis, from which it is almost inseparable, where it covers the aponeurosis of the temporal muscle.

*Inserted* into the upper part of the ear, opposite to the antihelix.

*Use.*

*Use.* To draw the ear upwards, and make the parts into which it is inserted tense.

*Superior auris*, Winflow.

## 2. ANTERIOR AURIS,

*Arises* thin and membranous near the posterior part of the zygoma.

*Inserted* into a small eminence on the back of the helix, opposite to the concha.

*Use.* To draw this eminence a little forwards and upwards.

## 3. RETRAHENTES AURIS,

*Arises*, sometimes by three, but always by two distinct small muscles, from the external and posterior part of the root of the mastoid process, immediately above the insertion of the sterno-cleido mastoid muscle.

*Inserted* into that part of the back of the ear which is opposite to the septum that divides the scapha and concha.



*Use.* To draw the ear back, and stretch the concha.

*Posterior auris, Winslow.*

The proper muscles are,

1. HELICIS MAJOR,

*Arises* from the upper and acute part of the helix anteriorly.

*Inserted* into its cartilage a little above the tragus.

*Use.* To depress that part from which it arises a little downwards and forwards.

2. HELICIS MINOR,

*Arises* from the inferior and anterior part of the helix.

*Inserted* into the crus of the helix, near the fissure in the cartilage opposite to the concha.

*Use.* To contract the fissure.

3. TRA-

## 3. TRAGICUS,

*Arises* from the middle and outer part of the concha, at the root of the tragus, along which it runs.

*Inserted* into the point of the tragus.

*Use.* Pulls the point of the tragus a little forwards.

## 4. ANTITRAGICUS,

*Arises* from the internal part of the cartilage that supports the antitragus; and, running upwards, is

*Inserted* into the tip of the antitragus, as far as the inferior part of the antihelix, where there is a fissure in the cartilage.

*Use.* Turns the tip of the antitragus, a little outwards, and depresses the extremity of the antihelix towards it.



## 5. TRANSVERSUS AURIS,

*Arises* from the prominent part of the concha on the dorsum of the ear; the fibres not so fleshy as in the former.

*Inserted* opposite to the outer side of the antihelix.

*Use.* Draws the parts to which it is connected towards each other, and stretches the scapha and concha.

The muscles of the internal ear are three:

## 1. LAXATOR TYMPANI,

*Arises* by a small beginning from the extremity of the spinous process of the sphenoid bone, behind the entry of the artery of the dura mater; then running backwards, and a little upwards, along with the nerve called *chorda tympani*, in a fissure of the os temporis near the fossa that lodges the condyle of the lower jaw.

*In-*



*Inserted* into the long process of the malleus, within the tympanum, where it rests upon the edge of the fissure between the pars squamosa and petrosa.

*Use.* To draw the malleus obliquely forwards towards its origin, consequently the membrana tympani, by which that membrane is made less concave, or is relaxed.

*Externus mallei*, Albinus.

*Anterior mallei*, Winflow.

*Obliquus auris*, Douglas.

## 2. TENSOR TYMPANI,

*Arises*, by a very small beginning, from the cartilaginous extremity of the Eustachian tube, just where it begins to be covered by the pars petrosa, and spinous process of the sphenoid bone, near the entry of the artery of the dura mater; from thence running backwards near the osseous part of the Eustachian tube, forms a very distinct fleshy belly, below a thin osseous plate, between



and the pars squamosa and labyrinth; and sends off a slender tendon, which makes a turn into the tympanum along with the nerve called *chorda tympani*.

*Inserted* into the posterior part of the handle of the malleus, a little lower than the root of its long process.

*Use.* To pull the malleus and membrana tympani inwards towards the pars petrosa, by which the membrane is made more concave and tense.

*Internus mallei*, Winslow.

*Internus auris*, Douglas.

### 3. STAPEDIUS,

*Arises*, by a small fleshy belly, from a little cavern in the pars petrosa, near the cells of the mastoid process, before the inferior part of the passage for the portio dura of the auditory nerve; its tendon passes straight through a small round hole in the  
same

same cavern, enters the anterior part of the tympanum, and is

*Inserted* into the posterior part of the head of the stapes.

*Use.* To draw the stapes obliquely upwards towards the cavern, by which the posterior part of its base is moved inwards, and the anterior part outwards.

*Musculus stapedis*, Winflow.

*Stapidæus*, Douglas.



## C H A P. III.

*Of the MUSCLES of the EYE-LIDS.*

THE palpebræ, or eye-lids, have one muscle common to both, and the upper eye-lid one proper to itself.

## 1. ORBICULARIS PALPEBRARUM.

*Arises*, by a number of fleshy fibres, from the outer edge of the orbital process of the superior maxillary bone, and from a tendon near the inner angle of the eye: these run a little downwards, then outwards, over the upper part of the cheek, below the orbit, covering the under eye-lid, and surround the external angle, being loosely connected only to the skin and fat; run over the superciliary ridge of the os frontis, towards the inner canthus, where they intermix



termix with those of the occipito-frontalis and corrugator supercilii; then covering the upper eye-lid, descend to the inner angle opposite to the inferior origin of this muscle, firmly adhering to the internal angular process of the os frontis, and to the short round tendon which serves to fix the palpebræ and muscular fibres arising from it.

*Inserted*, by the short round tendon, into the nasal process of the superior maxillary bone, covering the anterior and upper part of the lachrymal sac; which tendon can be easily felt at the inner canthus of the eye.

*Use*. To shut the eye, by drawing both lids close together, the fibres contracting from the outer angle towards the inner, press the eye-ball, squeeze the lachrymal gland, and convey the tears towards the puncta lachrymalia.

The *ciliaris* of some authors is only a part of this muscle covering the cartilages of the eye-lids called *cilia* or *tarfi*.

There



There is often a small fleshy slip which runs down from the outer and inferior part of this muscle above the zygomaticus minor, and joins with the levator labii superioris alæque nasi.

## 2. LEVATOR PALPEBRÆ SUPERIORIS,

*Arises* from the upper part of the foramen opticum of the sphenoid bone, through which the optic nerve passes, above the levator oculi, near the trochlearis muscle.

*Inserted*, by a broad thin tendon, into the cartilage that supports the upper eye-lid, named *tarsus*.

*Use*. To open the eye, by drawing the eye-lid upwards; which it does completely, by being fixed to the tarsus, pulling it below the eye-brow, and within the orbit.

*Aperiens palpebram rectus*, Douglas.



## C H A P. IV.

MUSCLES *of the* EYE-BALL.

THE muscles which move the globe of the eye are six, viz.

Four *straight* and two *oblique*.

The four straight muscles very much resemble each other; all

*Arising*, by a narrow beginning, a little tendinous and fleshy, from the bottom of the orbit around the foramen opticum of the sphenoid bone, where the optic nerve enters, so that they may be taken out adhering to this nerve; and all having strong fleshy bellies.

*Inserted* at the fore-part of the globe of the eye into the anterior part of the tunica sclerotica, and under the tunica adnata, at opposite sides, which indicates both their names and *Use*; so that they scarcely require



quire any further description, but to name them singly.

1. LEVATOR OCULI,

*Arises* from the upper part of the foramen opticum of the sphenoid bone, below the levator palpebræ superioris; and runs forwards to be

*Inserted* into the superior and fore-part of the tunica sclerotica, by a broad thin tendon.

*Use.* To raise up the globe of the eye.

*Atollens*, Albinus.

*Elevator*, Douglas.

2. DEPRESSOR OCULI,

*Arises* from the inferior part of the foramen opticum.

*Inserted* opposite to the former.

*Use.* To pull the globe of the eye down.

*Deprimens*, Albinus.

3. A D-

## 3. ADDUCTOR OCULI,

*Arises*, as the former, between the obliquus superior and depressor, being, from its situation, the shortest.

*Inserted* opposite to the inner angle.

*Use*. To turn the eye towards the nose.

## 4. ABDUCTOR OCULI,

*Arises* from the bony partition between the foramen opticum and lacerum, being the longest from its situation; and is

*Inserted* into the globe opposite to the outer canthus.

*Use*. To move the globe outwards.

The oblique muscles are two:

## 1. OBLIQUUS SUPERIOR, SEU TROCHLEARIS,

*Arises*, like the straight muscles, from the  
edge



edge of the foramen opticum at the bottom of the orbit, between the levator and adductor oculi; from thence runs straight along the pars plana of the ethmoid bone to the upper part of the orbit, where a cartilaginous trochlea is fixed to the inside of the internal angular process of the os frontis, through which its tendon passes, and runs a little downwards and outwards, inclosed in a loose membranous sheath.

*Inserted*, by a broad thin tendon, into the tunica sclerotica, about half-way between the insertion of the attollens oculi and optic nerve.

*Use*. To roll the globe of the eye, and to turn the pupil downwards and outwards, so that the upper side of the globe is turned inwards, and the inferior part of the outside of the orbit, and the whole globe drawn forwards towards the inner canthus.

*Obliquus major*, Winflow.



## 2. OBLIQUUS INFERIOR,

*Arises*, by a narrow beginning, from the outer edge of the orbital process of the superior maxillary bone, near its juncture with the os unguis; and, running obliquely outwards, is

*Inserted* into the sclerotica, in the space between the abductor and optic nerve, by broad thin tendon.

*Use*. To draw the globe of the eye forwards, inwards, and downwards; and, contrary to the superior, to turn the pupil upwards, towards the inner extremity of the eye-brow; at the same time, the external part of the globe is turned towards the inferior side, and the internal rolls towards the upper part.

*Obliquus minor*, Winflow.



## C H A P. V.

*Of the MUSCLES of the NOSE.*

THERE is only one muscle on each side that can be called proper to the nose, though it is affected by several muscles of the face.

## COMPRESSOR NARIS,

*Arises*, by a narrow beginning, from the root of the ala nasi externally, where part of the levator labii superioris alæque nasi is connected to it; it spreads into a number of thin disgregated fibres, which run up along the cartilage in an oblique manner, towards the dorsum of the nose, where it joins with its fellow, and is

*Inserted* slightly into the anterior extremity of the os nasi and nasal process of the superior maxillary bone, where it meets  
with



with some of the fibres descending from the occipito-frontalis muscle.

*Use.* To compress the ala towards the septum nasi, particularly when we want to smell acutely; but, if the fibres of the frontal muscle which adhere to it act, the upper part of this thin muscle assists to pull the ala outwards. It also corrugates the skin of the nose, and assists in expressing certain passions.

*Rincæus, vel nasalis, Douglas.*

C H A P.



## C H A P. VI.

MUSCLES *of the* MOUTH *and* LIPS.

THE mouth has nine pair of muscles, which are inserted into the lips, and a common one formed by the termination of these, viz. three *above*, three *below*, three *outwards*, and the common muscle surrounds the mouth.

The three above are,

## 1. LEVATOR ANGULI ORIS,

*Arises*, thin and fleshy, from the hollow of the superior maxillary bone, between the root of the socket of the first dens molaris and the foramen infra-orbitarium.

*Inserted* into the angle of the mouth and under-lip, where it joins with its antagonist.

*Use.* To draw the corner of the mouth  
up-



upwards, and make that part of the cheek opposite to the chin prominent, as in smiling.

*Elevator labiorum communis*, Douglas.

*Caninus*, Winflow.

2. LEVATOR LABII SUPERIORIS ALÆ-  
QUE NASI,

*Arises* by two distinct origins; the first broad and fleshy, from the external part of the orbital process of the superior maxillary bone which forms the lower part of the orbit, immediately above the foramen infra orbitarium; the second portion arises from the nasal process of the superior maxillary bone, where it joins the os frontis at the inner canthus, descending along the edge of the groove for the lachrymal sac. The first and shortest portion is

*Inserted* into the upper lip and orbicularis labiorum; the second and longest, into the upper lip and outer part of the ala nasi.

*Use.* To raise the upper-lip towards the  
or-



orbit, and a little outwards; the second portion serves to draw the skin of the nose upwards and outwards, by which the nostril is dilated.

*Elevator labii superioris proprius*, Douglas.

*Incisivus lateralis*, First portion; *Pyramidalis*, Second portion; Winslow.

### 3. DEPRESSOR LABII SUPERIORIS ALÆ- QUE NASI,

*Arises*, thin and fleshy, from the os maxillare superius, immediately above the joining of the gums with the two dentes incisivi, and the dens caninus; from thence it runs up under part of the levator labii superioris alæque nasi.

*Inserted* into the upper lip and root of the ala nasi.

*Use*. To draw the upper lip and ala nasi downwards and backwards.

*Depressor alæ nasi*, Albinus,

*Incisivus medius*, Winslow.



*Depressor labii superioris proprius*, Douglas.

The three below are,

1. DEPRESSOR ANGULI ORIS,

*Arises*, broad and fleshy, from the lower edge of the maxilla inferior, at the side of the chin, being firmly connected to that part of the platysma myoides, which runs over the maxilla to the angle of the mouth, to the depressor labii inferioris within, and to the skin and fat without, gradually turning narrower ; and is

*Inserted* into the angle of the mouth, joining with the zygomaticus major and levator anguli oris.

*Use*. To pull down the corner of the mouth.

*Triangularis*, Winslow.

*Depressor labiorum communis*, Douglas.



## 2. DEPRESSOR LABII INFERIORIS,

*Arises*, broad and fleshy, intermixed with fat, from the inferior part of the lower jaw next the chin; runs obliquely upwards; and is

*Inserted* into the edge of the under lip, extends along one half of the lip, and is lost in its red part.

*Use*. To pull the under lip and the skin of the side of the chin downwards, and a little outwards.

*Quadratus*, Winslow.

*Depressor labii inferioris proprius*, Douglas.

## 3. LEVATOR LABII INFERIORIS,

*Arises*, from the lower jaw, at the roots of the alveoli of two dentes incisivi and of the caninus; is

*Inserted* into the under-lip and skin of the chin.

*Use*.



*Use.* To pull the parts into which it is inserted upwards.

*Levator menti*, Albinus.

*Incisivus inferior*, Winflow.

*Elevator labii inferioris proprius*, Douglas.

The three outward are,

1. BUCCINATOR,

*Arises*, tendinous and fleshy, from the lower jaw, as far back as the last dens molaris and fore-part of the root of the coronoid process; fleshy from the upper jaw, between the last dens molaris and pterygoid process of the sphenoid bone; from the extremity of which it arises tendinous, being continued between both jaws to the constrictor pharyngis superior, with which it joins; from thence proceeding with straight fibres, and adhering close to the membrane that lines the mouth, it is

*Inserted* into the angle of the mouth within the orbicularis oris.



*Use.* To draw the angle of the mouth backwards and outwards, and to contract its cavity, by pressing the cheek inwards, by which the food is thrust between the teeth.

*Retractor anguli oris*, Albinus.

## 2. ZYGOMATICUS MAJOR,

*Arises*, fleshy, from the os malæ, near the zygomatic future.

*Inserted* into the angle of the mouth, appearing to be lost in the depressor anguli oris and orbicularis oris.

*Use.* To draw the corner of the mouth and under-lip towards the origin of the muscle, and make the cheek prominent, as in laughing.

*Zygomaticus*, Douglas.

## 3. ZYGOMATICUS MINOR,

*Arises* from the upper prominent part of the os malæ, above the origin of the former  
mer



mer muscle ; and, descending obliquely downwards and forwards, is

*Inserted* into the upper lip, near the corner of the mouth, along with the levator anguli oris.

*Use.* To draw the corner of the mouth obliquely outwards, and upwards, towards the external canthus of the eye.

The common muscle is the

#### ORBICULARIS ORIS.

This muscle is, in a great measure, formed by the muscles that move the lips ; the fibres of the superior descending, those of the inferior ascending, and decussating each other about the corner of the mouth, run along the lip to join those of the opposite side, so that the fleshy fibres appear to surround the mouth like a sphincter.

*Use.* To shut the mouth, by contracting and drawing both lips together, and to counteract all the muscles that assist in forming it.



*Sphincter labiorum*, Douglas.

*Semi-orbicularis*, Winslow.

*Constrictor oris*, Cowper.

There is another small muscle describe by Albinus, which he calls *Nasalis lab. superioris*; but it seems to be only some fibres of the former connected to the septum nasi.



## C H A P. VII.

MUSCLES *of the* LOWER JAW.

THE lower jaw has four pair of muscles for its elevation or lateral motions, viz. two, which are seen on the side of the face, and two concealed by the angle of the jaw.

## 1. TEMPORALIS,

*Arises*, fleshy, from a semicircular ridge of the lower and lateral part of the parietal bone, from all the pars squamosa of the temporal bone, from the external angular process of the os frontis, from the temporal process of the sphenoid bone, and from an aponeurosis which covers it: from these different origins the fibres descend like radii towards the jugum, under which they pass; and are



*Inserted*, by a strong tendon, into the upper part of the coronoid process of the lower jaw; in the duplicature of which tendon this process is inclosed as in a sheath, being continued down all its forepart to near the last dens molaris.

*Use*. To pull the lower jaw upwards, and press it against the upper, at the same time, drawing it a little backwards.

*N. B.* This muscle is covered by a tendinous membrane, called its *aponeurosis*, which arises from the bones that give origin to the upper and semicircular part of the muscle; and, descending over it, is inserted into all the jugum, and the adjoining part of the os frontis.

The use of this membrane is to give room for the origin of a greater number of fleshy fibres, to fortify the muscle in its action, and to serve as a defence to it.

*Crotaphite muscle*, Winflow.



## 2. MASSETER.

*Arises*, by strong, tendinous, and fleshy fibres, which run in different directions, from the superior maxillary bone, where it joins the os malæ, and from the inferior and interior part of the zygoma, its whole length, as far back as the tubercle before the socket for the condyle of the lower jaw; the external fibres slanting backwards, and the internal forwards.

*Inserted* into the angle of the lower jaw, and from that upwards to near the top of its coronoid process.

*Use*. To pull the lower to the upper jaw, and, by means of its oblique decussation, a little forwards and backwards.

## 3. PTERYGOIDEUS INTERNUS,

*Arises*, tendinous and fleshy, from the inner and upper part of the internal plate  
c 5 of



of the pterygoid process, filling all the space between the two plates; and from the pterygoid process of the os palati between these plates.

*Inserted* into the angle of the lower jaw internally.

*Use.* To draw the jaw upwards, and obliquely towards the opposite side.

*Pterygoideus major*, Winflow.

#### 4. PTERYGOIDEUS EXTERNUS.

*Arises* from the outer side of the external plate of the pterygoid process of the sphenoid bone, from part of the tuberosity of the os maxillare adjoining to it, and from the root of the temporal process of the sphenoid bone.

*Inserted* into a cavity in the neck of the condyloid process of the lower jaw; some of its fibres are inserted into the ligament that connects the moveable cartilage and that process to each other.

*Use.*

*Use.* To pull the lower jaw forwards, and to the opposite side ; and to pull the ligament from the joint, that it may not be pinched during these motions : when both external pterygoid muscles act, the fore-teeth of the under-jaw are pushed forwards beyond those of the upper jaw.

*Pterygoideus minor*, Winflow.



## C H A P. VIII.

*The MUSCLES which appear about the anterior part of the NECK.*

**O**N the side of the neck are two muscles or layers.

1. MUSCULUS CUTANEUS,  
VULGO  
PLATYSMA MYOIDES.

*Arises*, by a number of slender disgregated fleshy fibres, from the cellular substance that covers the upper parts of the deltoid and pectoral muscles; in their ascent they all unite to form a thin muscle, which runs obliquely upwards along the side of the neck, adhering to the skin.

*Inserted* into the lower jaw, between its angle and the origin of the depressor, angulari

guli oris, to which it is firmly connected, and but slightly to the skin that covers the inferior part of the masseter muscle and parotid glands.

*Use.* To assist the depressor anguli oris in drawing the skin of the cheek downwards; and when the mouth is shut, it draws all that part of the skin, to which it is connected, below the lower jaw, upwards.

*Platysma myoides*, Galen.

*Musculus cutaneus*, Winslow.

*Quadratus genæ, vel Latissimus colli*, Douglas.

*Latissimus colli*, Albinus.

## 2. STERNO-CLEIDO-MASTOIDEUS;

*Arises* by two distinct origins; the anterior, tendinous and a little fleshy, from the top of the sternum near its junction with the clavicle; the posterior, fleshy, from the upper and anterior part of the clavicle; both unite a little above the anterior articulation



lation of the clavicle, to form one muscle, which runs obliquely upwards and outwards, to be

*Inserted*, by a thick strong tendon, into the mastoid process, which it surrounds; and, gradually turning thinner, is inserted as far back as the lambdoid suture.

*Use.* To turn the head to one side, and bend it forwards.

*Sterno-mastoideus* and *Cleido-mastoideus*, Albinus.

*Mastoideus*, Douglas.

## C H A P. IX.

MUSCLES *situated between the* LOWER  
JAW *and* Os HYOIDES.

THERE are four layers before, and two muscles at the side.

The four layers are,

## 1. DIGASTRICUS,

*Arises*, by a fleshy belly, intermixed with tendinous fibres, from the fossa at the root of the mastoid process of the temporal bone, and soon becomes tendinous; runs downwards and forwards: the tendon passes generally through the stylo-hyoideus muscle; then it is fixed by a ligament to the os hyoides; and, having received from that bone an addition of tendinous and muscular fibres, runs obliquely forwards, turns fleshy again, and is

*Inserted,*



*Inserted*, by this anterior belly, into a rough sinuosity at the inferior and anterior edge of that part of the lower jaw called the *chin*.

*Use*. To open the mouth, by pulling the lower jaw downwards and backwards; and, when the jaws are shut, to raise the larynx, and consequently the pharynx, upwards, as in deglutition.

*Biventer maxillæ inferioris*, Albinus.

## 2. MYLO-HYOIDEUS,

*Arises*, fleshy, from all the inside of the lower jaw between the last dens molaris and the middle of the chin, where it joins with its fellow.

*Inserted* into the lower edge of the basis of the os hyoides, and joins with its fellow.

*Use*. To pull the os hyoides forwards, upwards, and to a side.



## 3. GENIO-HYOIDEUS,

*Arises*, tendinous, from a rough protuberance in the middle of the lower jaw internally, or inside of the chin.

*Inserted* into the basis of the os hyoides.

*Use*. To draw this bone forwards to the chin.

## 4. GENIO-HYO-GLOSSUS,

*Arises*, tendinous, from a rough protuberance in the inside of the middle of the lower-jaw; its fibres run, like a fan, forwards, upwards, and backwards; and are

*Inserted* into the tip, middle, and root of the tongue, and base of the os hyoides, near its cornu.

*Use*. According to the direction of its fibres, to draw the tip of the tongue backwards into the mouth, the middle downwards, and to render its dorsum concave;

to



to draw its root and os hyoides forwards, and to thrust the tongue out of the mouth.

The two muscles at the side are,

1. HYO-GLOSSUS,

*Arises*, broad and fleshy, from the base, cornu, and appendix of the os hyoides; the fibres run upwards and outwards, to be

*Inserted* into the side of the tongue, near the stylo-glossus.

*Use*. To pull the tongue inwards and downwards.

*Basio-cerato-chondro-glossus*, Albinus.

*Cerato-glossus*, Douglas.

2. LINGUALIS,

*Arises* from the root of the tongue laterally; runs forwards between the hyo-glossus and genio-glossus, to be

*Inserted* into the tip of the tongue, along with part of the stylo-glossus.

*Use*. To contract the substance of the tongue, and bring it backwards.

## C H A P. X.

MUSCLES *situated between the* Os Hy-  
OIDES *and* TRUNK.

THESE may be divided into two layers.

The first layer consists of two muscles.

## 1. STERNO-HYOIDEUS,

*Arises*, thin and fleshy, from the cartilaginous extremity of the first rib, the upper and inner part of the sternum, and from the clavicle where it joins with the sternum.

*Inserted* into the base of the os hyoides.

*Use*. To pull the os hyoides downwards.

## 2. OMO-HYOIDEUS,

*Arises*, broad, thin, and fleshy, from the superior



superior costa of the scapula, near the femi-lunar nitch, and from the ligament that runs across it; thence ascending obliquely, it becomes tendinous below the sterno-cleido-mastoid muscle; and, growing fleshy again, is

*Inserted* into the base of the os hyoides, between its cornu and the insertion of the sterno hyoideus.

*Use.* To pull the os hyoides obliquely downwards.

*Coraco-hyoideus* Albinus and Douglas.

The second layer consists of three muscles.

#### 1. STERNO-THYROIDEUS,

*Arises*, fleshy, from the whole edge of the uppermost bone of the sternum internally, opposite to the cartilage of the first rib from which it receives a small part of its origin.

*Inserted*, into the surface of the rough line at the external part of the inferior edge of the thyroid cartilage.

*Use.* To draw the larynx downwards.

#### 2. THY-



## 2. THYREO-HYOIDEUS,

*Inserted* into part of the basis, and almost all the cornu of the os hyoides.

*Arises* from the rough line, opposite to the former.

*Use.* To pull the os hyoides downwards, or the thyroid cartilage upwards.

*Thyro-hyoideus, vel Hyo-thyroideus, Win-flow.*

## 3. CRICO-THYROIDEUS,

*Arises* from the side and fore-part of the cricoid cartilage, running obliquely upwards.

*Inserted* by two portions; the first, into the lower part of the thyroid cartilage; the second, into its inferior cornu.

*Use.* To pull forwards and depress the thyroid, or to elevate and draw backwards the cricoid cartilage.



## C H A P. XI.

MUSCLES *situated between the* LOWER JAW *and* Os HYOIDES *laterally.*

THEY are five in number. Three proceed from the styloid process of the temporal bone, from which they have half of their names; and two from the pterygoid process of the sphenoid bone.

The three from the styloid process are,

## 1. STYLO-GLOSSUS,

*Arises*, tendinous and fleshy, from the stylo process, and from a ligament that connects that process to the angle of the lower jaw.

*Inserted* into the root of the tongue, runs along its side, and is insensibly lost near its tip.

*Use.*

*Use.* To draw the tongue laterally and backwards.

## 2. STYLO-HYOIDEUS,

*Arises*, by a round tendon, from the middle and inferior part of the styloid process.

*Inserted* into the os hyoides at the junction of the base and cornu.

*Use.* To pull the os hyoides to one side, and a little upwards.

N. B. Its fleshy belly is generally perforated by the tendon of the digastric muscle, on one or both sides. There is often another accompanying it, called *stylo-hyoideus alter*; and has the same origin, insertion, and use.

## 3. STYLO-PHARYNGEUS,

*Arises*, fleshy, from the root of the styloid process.

*Inserted* into the side of the pharynx and back part of the thyroid cartilage.

*Use.*



*Ufe.* To dilate and raife the pharynx and thyroid cartilage upwards.

The two from the pterygoid procefs are,

1. CIRCUMFLEXUS, OR TENSOR-PALATI,

*Arifes* from the spinous procefs of the fphenoid bone, behind the foramen ovale, which tranfmits the third branch of the fifth pair of nerves ; from the Eufiachian tube, not far from its offeous part : it then runs down along the pterygoideus internus, paffes over the hook of the internal plate of the pterygoid procefs by a round tendon, which foon spreads into a broad membrane.

*Inferted* into the velum pendulum palati, and the femilunar edge of the os palati, and extends as far as the future which joins the two bones. Generally fome of its pofterior fibres join with the constrictor pharyngis fuperior, and palato-pharyngeus.

*Ufe.* To ftretch the velum, to draw it down-



downwards, and to a side towards the hook. It has little effect upon the tube, being chiefly connected to its osseous part.

*Circumflexus palati*, Albinus.

*Spheno-salpingo-staphilinus*, seu *Staphilinus externus*, Winflow.

*Musculus tubæ novus*, Valsalva; vel *Palato-salpingeus*, Douglas.

## 2. LEVATOR PALATI,

*Arises*, tendinous and fleshy, from the extremity of the pars petrosa of the temporal bone, where it is perforated by the Eustachian tube, and also from the membranous part of the same tube.

*Inserted* into the whole length of the velum pendulum palati, as far as the root of the uvula, and unites with its fellow.

*Use*. To draw the velum upwards and backwards, so as to shut the passage from the fauces into the mouth and nose.

*Levator palati mollis*, Albinus.



*Petro-salpingo-staphilinus*, vel *Salpingo-staphilinus internus*, Winflow.

*Salpingo-staphilinus*, Valsalva. *Pterigo-staphilinus externus*, vulgo, Douglas.

*Spheno-staphilinus*, Cowper.

Previous to the description of the muscles situated about the passage into the throat, it will be necessary to mention the principal parts to which they are connected.

Upon looking into any person's mouth, when wide opened, we see a soft curtain hanging from the palate-bones, named *velum pendulum palati*. In the middle of which, we likewise observe a papilla projecting from the velum, named *uvula*, or *pap of the throat*. From each side of the uvula, at its root, two arches, or columns, are sent down; the interior to the root of the tongue, the posterior to the pharynx. Between these arches, on each side, the cellular glands called *amygdalæ*, or *almonds of the ears*, are situated.

The



The common opening behind the anterior arch may be named *fauces*, or *top of the throat*, from which there are six passages, viz. two upwards, being one to each nostril: two at the sides, or one to each ear, called the *Eustachian tubes*: two downwards; the anterior is the passage through the *glottis* and *larynx*, into the trachea, which terminates in the lungs; the posterior is the largest, named *pharynx*, or top of the *æsophagus*, which leads to the stomach.



## C H A P. XII.

MUSCLES *situated about the entry to the*  
FAUCES.

THERE are two on each side, and a single one in the middle.

The two on each side are,

## 1. CONSTRICTOR ISTHMI FAUCIUM,

*Arises*, by a slender beginning, from the side of the tongue, near its root; from thence running upwards, within the anterior arch, before the amygdala, it is

*Inserted* into the middle of the velum pendulum palati, at the root of the uvula anteriorly, being connected with its fellow, and with the beginning of the palato-pharyngeus.

*Use.* Draws the velum towards the root of  
the



the tongue, which it raises at the same time, and, with its fellow, contracts the passage between the two arches, by which it shuts the opening into the fauces.

*Glossopharynx*, Winslow and Douglas.

## 2. PALATO-PHARYNGEUS.

*Arises*, by a broad beginning, from the middle of the velum pendulum palati, at the root of the uvula posteriorly, and from the tendinous expansion of the circumflexus palati. The fibres are collected within the posterior arch behind the amygdala, and run backwards to the top and lateral part of the pharynx, where the fibres are scattered, and mix with those of the stylo-pharyngeus.

*Inserted* into the edge of the upper and back part of the thyroid cartilage; some of its fibres being lost between the membrane of the pharynx and the two inferior constrictors.

*Use*. Draws the uvula and velum down-



wards and backwards ; and at the same time pulls the thyroid cartilage and pharynx upwards, and shortens it : with the constrictor superior and tongue, it assists in shutting the passage into the nostrils ; and, in swallowing, it thrusts the food from the fauces into the pharynx.

*Thyro-staphilinus*, Douglas.

*Thyro-pharyngo-staphilinus*, Winflow.

SALPINGO-PHARYNGEUS of Albinus is composed of a few fibres of the muscle, which

*Arise* from the anterior and lower part of the cartilaginous extremity of the Eustachian tube ; and are

*Inserted* into the inner part of the last-mentioned muscle.

*Use.* To assist the former, and to dilate the mouth of the tube.

The

The one in the middle is the

AZYGOS UVULÆ.

*Arises*, fleshy, from the extremity of the future which joins the palate-bones, runs down the whole length of the velum and uvula, resembling a small earth-worm, and adhering to the tendons of the circumflexi.

*Inserted* into the tip of the uvula.

*Use*. Raises the uvula upwards and forwards, and shortens it.

*Palato-staphilinus*, Douglas.

*Staphilinus*, or *Epistaphilinus*, Winslow.



## C H A P. XIII.

MUSCLES *situated on the posterior part*  
*of the PHARYNX.*

**O**F these there are three pair.

## 1. CONSTRICTOR PHARYNGIS INFERIOR,

*Arises* from the side of the thyriod cartilage, near the attachment of the sterno-hyoideus and thyreo-hyoideus muscles; and from the cricoid cartilage, near the crico-thyroideus. This muscle is the largest of the three, and is

*Inserted* into the white line, where it joins with its fellow; the superior fibres running obliquely upwards, covering nearly one half of the middle constrictor, and terminating in a point; the inferior fibres run more transf-



transversely, and cover the beginning of the œsophagus.

*Use.* To compress that part of the pharynx which it covers, and to raise it with the larynx a little upwards.

*Thyro-pharyngeus, Crico-pharyngeus, Douglas.*

## 2. CONSTRICTOR PHARYNGIS MEDIUS.

*Arises* from the appendix of the os hyoides, from the cornu of that bone, and from the ligament which connects it to the thyroid cartilage; the fibres of the superior part running obliquely upwards, and, covering a considerable part of the superior constrictor, terminate in a point.

*Inserted* into the middle of the cuneiform process of the os occipitis, before the foramen magnum, and joined to its fellow at a white line in the middle back part of the pharynx. The fibres at the middle part



run more transversely than those above or below.

*Use.* To compress that part of the pharynx which it covers, and to draw it and the os hyoides upwards.

*Hyo-pharyngeus, Syndesmo-pharyngeus, Douglas.*

### 3. CONSTRUCTOR PHARYNGIS SUPERIOR,

*Arises,* above, from the cuneiform process of the os occipitis, before the foramen magnum, near the holes where the ninth pair of nerves passes out ; lower down, from the pterygoid process of the sphenoid bone ; from the upper and under jaw, near the roots of the last dentes molares ; and between the jaws, it is continued with the buccinator muscle, and with some fibres from the root of the tongue, and from the palate.

*Inserted* into a white line in the middle of the pharynx, where it joins with its fellow,

low, and is covered by the constrictor medius.

*Use.* To compress the upper part of the pharynx, and draw it forwards and upwards.

*Cephalo-pharyngeus, Pterygo-pharyngeus,*  
*Mylo-pharyngeus, Glossopharyngeus,*  
Douglas.



## C H A P. XIII.

MUSCLES *situated about the* GLOTTIS.

THEY consist generally of four pair of small muscles, and a single one.

## 1. CRICO-ARYTÆNOIDEUS POSTICUS,

*Arises*, fleshy, from the back part of the cricoid cartilage, and is

*Inserted* into the posterior part of the base of the arytenoid cartilage.

*Use*. To open the rima glottidis a little ; and, by pulling back the arytenoid cartilage, to stretch the ligament so as to make it tense.

## 2. CRICO-ARYTÆNOIDEUS LATERALIS,

*Arises*, fleshy, from the cricoid cartilage,  
la-



laterally, where it is covered by part of the thyroid, and is

*Inserted* into the side of the base of the arytenoid cartilage near the former.

*Use.* To open the rima glottidis, by pulling the ligaments from each other.

### 3. THYREO-ARYTÆNOIDEUS,

*Arises* from the under and back part of the middle of the thyroid cartilage; and, running backwards and a little upwards, along the side of the glottis, is

*Inserted* into the arytenoid cartilage, higher up and farther forwards than the crico-arytænoides lateralis.

*Use.* To pull the arytenoid cartilage forwards nearer to the middle of the thyroid, and consequently to shorten and relax the ligament of the larynx or glottis vera.

### 4. ARY-



## 4. ARYTÆNOIDEUS OBLIQUUS,

*Arises* from the base of one arytenoid cartilage; and, crossing its fellow, is

*Inserted* near the tip of the other arytenoid cartilage.

*Use.* When both act, they pull the arytenoid cartilages towards each other.

*N. B.* Very often one of these is wanting.

*Arytænoides minor*, Douglas.

The single muscle is the

## ARYTÆNOIDEUS TRANSVERSUS,

*Arises* from the side of one arytenoid cartilage, from near its articulation with the cricoid to near its tip. The fibres run straight across; and are

*Inserted*, in the same manner, into the other arytenoid cartilage.

*Use.* To shut the rima glottidis, by bringing



ing these two cartilages, with the ligaments, nearer one another.

*Arytænoides major*, Douglas.

Besides these, there are a few disgregated muscular fibres on each side; which, from their general direction, are named,

1. THYREO-EPIGLOTTIDEUS,

*Arises*, by a few pale disgregated fibres, from the thyroid cartilage; and is

*Inserted* into the epiglottis laterally.

*Use*. To draw the epiglottis obliquely downwards, or, when both act, directly downwards; and, at the same time, it expands that soft cartilage.

2. ARYTÆNO-EPIGLOTTIDEUS,

*Arises*, by a number of small fibres, from the lateral and upper part of the arytenoid cartilage; and, running along the outer side of the external rima, is

*In-*



*Inserted* into the epiglottis along with the former.

*Use.* To pull that side of the epiglottis towards the external rima; or, when both act, to pull it close upon the glottis. It is counteracted by the elasticity of the epiglottis.

## C H A P XV.

MUSCLES *situated on the anterior part*  
*of the ABDOMEN.*

THEY consist of three broad layers on each side of the belly ; always a long one, and generally also a short one, on each side of the linea alba.

The three layers are,

1. OBLIQUUS DESCENDENS EXTERNUS,

*Arises*, by eight heads, from the lower edges of an equal number of inferior ribs, at a little distance from their cartilages : it always intermixes, in a serrated manner, with portions of the serratus major anticus ; and generally coheres to the pectoralis major, intercostals, and latissimus dorsi ; which last covers the edge of a portion of it extended



tended from the last rib to the spine of the os ilium. From these origins the fibres run down obliquely forwards, and terminate in a thin broad tendon, whose fibres are continued in the same direction.

*Inserted* into the whole length of the *linea alba*\*, becomes thicker towards the lower part of the abdomen, and is perforated in the middle by the umbilicus†. On the outside of the rectus muscle, the tendon of the external oblique appears whiter than elsewhere, by its being there connected with the tendons of the internal oblique and

\* The *linea alba* is formed by the tendinous fibres of the two oblique and transverse muscles, interlaced with those of the opposite side, the whole way from the cartilago ensiformis to the os pubis; so that some think they should be called three digastric muscles, with a broad middle tendon and two fleshy bellies.

† The *umbilicus* was originally the passage for the vessels that connected the foetus to the secundines; and is really a hole through the teguments and tendons, filled up only by a cellular substance, and covered within by the peritoneum.

trans



transverse muscles; so that this part has been called *Linea semilunaris*, from its curved shape. The under part of the tendon divides into two columns, which leaves an oval space between them, named the *ring* § of the external oblique muscle; for the passage of the spermatic cord in the male or round ligament of the womb: The anterior superior column passes over the cartilage between the ossa pubis, and is fixed to the opposite os pubis; the other is fixed to the os pubis of the same side. It is also inserted, tendinous and fleshy, into the middle of the spine of the ilium.

From that part, which is named its *anterior superior spinous process*, it is stretched tendinous to the os pubis, and is named

§ The *ring* of the external oblique muscle is made somewhat circular, by a thin tendinous or rough cellular substance, which helps to fill it up; and though a few muscular fibres of the internal are separated, yet the stricture in herniæ only happens in the tendon of the external.

Pou-



*Poupart's, or Fallopius's ligament* \*. From this ligament it sends a tendinous layer, which is lost in the membranous fascia of the thigh.

*Use.* Supports and compresses the peritoneum and abdomen; assists the evacuation of fæces and urine, and likewise in the exclusion of the foetus; thrusts the diaphragm upwards, and draws down the ribs in expiration; bends the body obliquely when the ribs are fixed, and raises the pelvis obliquely.

*Obliquus externus abdominis, Albinus.*

*Obliquus descendens, Douglas.*

\* *Poupart's, or Fallopius's ligament*, is the inferior part of the tendon of the external oblique, extending from the anterior superior spinous process of the ilium to the os pubis, where it is thickest, in order to strengthen the inferior part of the abdomen: here it is not inserted into any bone, but passes over the blood-vessels of the inferior extremity; and in women, from the greater size of the pelvis, is longer and looser, by which they are more subject to crural herniæ; but, by the size of the spermatic cord, men are more liable to the inguinal.



## 2. OBLIQUUS ASCENDENS INTERNUS,

*Arises* from the spine of the ilium, the whole length between the posterior and superior anterior spinous process; from the os sacrum and the three undermost lumbar vertebræ, by a tendon common to it and to the serratus posticus inferior muscle; from Poupart's ligament, at the middle of which it sends off the beginning of the cremaster muscle; and the spermatic cord in the male, or round ligament of the womb, passes under its thin edge, except a few detached fibres.

*Inserted* into the cartilago ensiformis, into the cartilages of the seventh and those of all the false ribs; but, at the upper part, it is extremely thin, resembling a cellular membrane, and only becomes fleshy at the cartilage of the tenth rib. Here its tendon divides into two layers\*; the anterior layer, with

\* To obtain a proper view of the two layers of the  
ten-



with a great portion of the inferior part of the posterior layer, joins the tendon of the external oblique, and runs over the rectus to be inserted into the whole length of the linea alba. The posterior layer joins the tendon of the transversalis muscle as low as half-way between the umbilicus and os pubis ; but, below this place, only a few fibres of the posterior layer are seen, and the rest of it passes before the rectus muscle, and is inserted into the linea alba ; so that the whole tendon of the external oblique muscle, with the anterior layer of the inter-

tendon of the internal oblique muscle, both the oblique muscles should be raised as far forwards as their joining near the linea semilunaris ; then the tendon before the rectus must be cut parallel to the linea alba, and turned outwards as far as the outer edge of the rectus ; by which the whole of the rectus is brought into view, and the tendons are preserved. But Douglas directs to cut the posterior layer of the internal oblique, where it joins with the transversalis : by this method the rectus is laid bare ; but the structure of the tendinous sheath, which incloses it, is destroyed.



nal oblique, passes before the rectus muscle; and the whole posterior layer of the internal oblique, together with the whole tendon of the transversalis muscle, excepting at the inferior part, pass behind the rectus, and are inserted into the linea alba. At its undermost part it is inserted into the forepart of the os pubis.

*Use.* To assist the former; but it bends the trunk in the reverse direction.

*Obliquus internus abdominis*, Albinus and Winslow.

*Obliquus ascendens*, Douglas.

### 3. TRANSVERSALIS,

*Arises* tendinous, but soon becoming fleshy from the inner or back part of the cartilages of the seven lower ribs, where some of its fibres are continued with those of the diaphragm and the intercostal muscles; by a broad thin tendon, connected to the transverse processes of the last vertebra of the back



back and the four superior vertebræ of the loins; fleshy, from the whole spine of the os ilium internally, and from the tendon of the external oblique muscle, where it intermixes with some fibres of the internal oblique.

*Inserted* into the cartilago enfiformis, and into the whole length of the linea alba, excepting its lowermost part.

*Use.* To support and compress the abdominal bowels; and it is so particularly well adapted for the latter purpose, that it might be called the *proper constrictor* of the abdomen.

*Transversus abdominis*, Albinus.

The long muscle in the middle is named,

### RECTUS ABDOMINIS,

*Arises*, by two heads, from the ligament of the cartilage which joins the two ossa pubis to each other; runs upwards the whole length of, and parallel to, the  
linea



linea alba, growing broader and thinner as it ascends.

*Inserted* into the cartilages of the three inferior true ribs, and often intermixes with some fibres of the pectoral muscle.

It is generally divided by three tendinous interfections; the first is at the umbilicus, the second where it runs over the cartilage of the seventh rib, the third in the navel between these; and there is commonly a half interfection below the umbilicus. These seldom penetrate through the whole thickness of the muscle; they adhere firmly to the anterior part of the sheath, but very slightly to the posterior layer.

*Use.* To compress the fore-part, but more particularly the lower part of the belly; to bend the trunk forwards, or to raise the pelvis. By its tendinous interfections, it is enabled to contract at any of the intermediate spaces; and, by its connection with the tendons of the other muscles, it is prevented from changing place, and from ris-



ing into a prominent form when in action.

The short muscle in the middle is named

PYRAMIDALIS,

*Arises* along with the rectus; and, running upwards within the same sheath, is

*Inserted*, by an acute termination, near half-way between the os pubis and umbilicus, into the linea alba and inner edge of the rectus muscle.

As it is frequently wanting in both sides, without any inconvenience, its

*Use* seems to be, to assist the inferior part of the rectus.



## C H A P. XVI.

MUSCLES *about the Male Organs of*  
GENERATION.

THE *testicles* are said to have a thin muscle common to both, and have one proper to each.

The supposed common muscle is called the

## DARTOS.

This appears to be no more than a condensation of the cellular membrane lining the scrotum ; yet the skin here is capable of being corrugated and relaxed in a greater degree than in other places.



The muscle proper to each testicle is the

### CREMASTER.

*Arises* from the internal oblique, where a few fibres of that muscle intermix with the transversalis, near the junction of the os ilium and pubis, over which part it passes, after having pierced the ring of the external oblique; and then it descends upon the spermatic cord.

*Inserted* into the tunica vaginalis of the testicle, upon which it spreads, and is insensibly lost.

*Use.* To suspend and draw up the testicle, and to compress it in the act of coition.

The penis has three pair of muscles.

### 1. ERECTOR PENIS,

*Arises*, tendinous and fleshy, from the tuberosity of the os ischium, and runs upwards, embracing the whole crus of the penis.

*Inserted* into the strong tendinous membrane



brane that covers the corpora cavernosa penis, near as far up as the union of these bodies.

*Use.* To compress the crus penis, by which the blood is pushed from it into the fore-part of the corpora cavernosa ; and the penis is by that means more completely distended. The erectores seem likewise to keep the penis in its proper direction.

*Ischio-cavernosus*, Winslow.

2. ACCELERATOR URINÆ, SEU EJACULATOR SEMINIS,

*Arises*, fleshy, from the sphincter ani and membranous part of the urethra ; and tendinous from the crus, near as far forwards as the beginning of the corpus cavernosum penis ; the inferior fibres run more transversely, and the superior descend in an oblique direction.

*Inserted* into a line in the middle of the bulb, where it joins with its fellow, by which the bulb is completely inclosed.



*Use.* To drive the urine or semen forwards; and, by grasping the bulb of the urethra, to push the blood towards its corpus cavernosum and the glans, by which they are distended.

*Bulbo-cavernosus*, Winslow.

### 3. TRANSVERSUS PERINEI,

*Arises* from the tough fatty membrane that covers the tuberosity of the os ischium; from thence it runs transversely inwards, and is

*Inserted* into the accelerator urinæ, and into that part of the sphincter ani which covers the bulb.

*Use.* To dilate the bulb, and draw the perineum and verge of the anus a little outwards and backwards.

*Transversalis urethræ*, Winslow.

*Transversus perinei*, Albinus.

*Levator parvus, seu externus*, Douglas.

There



There is often a fourth muscle, named

TRANSVERSUS PERINEI ALTER,

*Arises* behind the former, runs more obliquely forwards, and is

*Inserted* into that part of the accelerator urinæ which covers the anterior part of the bulb of the urethra.

*Use.* To assist the former.

*Inferior prostate, Winslow.*

*Transversus perinei alter, Albinus.*



## C H A P. XVII.

MUSCLES *of the* ANUS.

**T**HE *anus* has a single muscle, and one pair.

The single muscle is

## SPHINCTER ANI,

*Arises* from the skin and fat that surround the verge of the anus on both sides, near as far out as the tuber of the os ischium: the fibres are gradually collected into an oval form, and surround the extremity of the rectum.

*Inserted*, before, by a narrow point, into the perineum, acceleratores urinæ, and transversi perinei; behind, by an acute termination, into the extremity of the os coccygis.

*Use.*



*Use.* Shuts the passage through the anus into the rectum; pulls down the bulb of the urethra, by which it assists in ejecting the urine and semen.

*Sphincter externus*, Albinus and Douglas.

*Sphincter cutaneus*, Winflow.

N. B. The sphincter internus of Albinus and Douglas, is only that part of the circular fibres of the muscular coat of the rectum which surrounds its extremity.

#### LEVATOR ANI,

*Arises* from the os pubis within the pelvis, as far up as the upper edge of the foramen thyroideum, and joining of the os pubis with the os ischium; from the thin tendinous membrane that covers the obturator internus and coccygeus muscles; from the spinous process of the os ischium; and its fibres run down like rays from a circumference to a centre.

*Inserted* into the sphincter ani, acceleratores



tores urinæ, and anterior part of the two last bones of the coccygis; surrounds the extremity of the rectum, neck of the bladder, prostate gland, and part of the vesiculæ feminales; so that its fibres behind and below the os coccygis joining it with its fellow, they, together, very much resemble the shape of a funnel.

*Use.* To draw the rectum upwards after the evacuation of the fæces, and to assist in shutting it; to sustain the contents of the pelvis, and to help in ejecting the semen, urine, and contents of the rectum; and, perhaps, by pressing upon the veins, to contribute greatly to the erection of the penis.



## C H A P. XVIII.

MUSCLES of the Female Organs of  
GENERATION.

**T**HE *clitoris* has one pair.

## ERECTOR CLITORIDIS,

*Arises* from the crus of the os ischium internally, and in its ascent covers the crus of the clitoris as far up as the os pubis.

*Inserted* into the upper part of the crus and body of the clitoris.

*Use.* Draws the clitoris downwards and backwards; and may serve to make the body of the clitoris more tense, by squeezing the blood into it from its crus.

*First muscle of the clitoris, Douglas.*



The vagina has one pair,

### SPHINCTER VAGINÆ,

*Arises* from the sphincter ani, and from the posterior side of the vagina near the perineum; from thence it runs up the side of the vagina, near its external orifice, opposite to the nymphæ, and covers the corpus cavernosum vaginæ.

*Inserted* into the crus and body, or union of the crura clitoridis.

*Use.* Contracts the mouth of the vagina, and compresses its corpus cavernosum.

*Constrictor cunni*, Albinus.

*Second muscle of the clitoris*, Douglas.

The perineum has one pair.

### TRANSVERSUS PERINEI,

*Arises*, as in the male, from the fatty cellular membrane which covers the tuberosity of the os ischium.

*In-*



*Inserted* into the upper part of the sphincter ani, and into a white hardish tough substance in the perineum, between the lower part of the pudendum and anus.

*Use.* To sustain and keep the perineum in its proper place.

The anus, as in the male, has a single muscle, and one pair,

#### SPHINCTER ANI,

*Arises*, as in the male, from the skin and fat surrounding the extremity of the rectum.

*Inserted*, above, into the white tough substance of the perineum; and below into the point of the os coccygis.

*Use.* To shut the passage into the rectum; and, by pulling down the perineum, to assist in contracting the mouth of the vagina.

#### LEVATOR ANI,

*Arises*, as in the male, within the pelvis



vis, and descends along the inferior part of the vagina and rectum.

*Inserted* into the perineum, sphincter ani, extremity of the vagina, and rectum.

*Use.* To raise the extremity of the rectum upwards, to contract the inferior part of the rectum, and to assist in contracting and supporting the vagina; and, perhaps, by pressing on the veins, to contribute to the distention of the cells of the clitoris and corpus cavernosum of the vagina.



## C H A P. XIX.

MUSCLES *situated within the* PELVIS.

OF these there are two pair.

## 1. OBTURATOR INTERNUS,

*Arises* from more than one half of the internal circumference of the foramen thyroideum, formed by the os pubis and ischium: its inside is covered by a portion of the levator ani; and appears to be divided into a number of fasciculi, which unite and form a roundish tendon, that passes out of the pelvis, between the posterior sacro-ischiatic ligament and tuberosity of the os ischium, where it passes over the capsular ligament of the thigh-bone; it is inclosed, as in a sheath, by the gemini muscles.

*Inserted*, by a round tendon, into the large pit at the root of the trochanter major.

*Use.*



*Ufe.* To roll the os femoris obliquely outwards.

*Marfupialis, feu Obturator internus, Douglas.*

*N. B.* The infertion of this mufcle fhould not be profecuted, until the mufcles of the thigh, to which it belongs, are diffefted. *Vid.* Chap. xxix.

## 2. COCCYGEUS.

*Arifes,* tendinous and flefhy, from the fpinous procefs of the os ifchium, and covers the infide of the posterior facro-ifchiatic ligament; from this narrow beginning, it gradually increafes, to form a thin flefhy belly, interfperfed with tendinous fibres.

*Inferted* into the extremity of the os facrum, and near the whole length of the os coccygus laterally.

*Ufe.* To fupport and move the os coccygis forwards, and to tie it more firmly to the facrum.



## C H A P. XX.

MUSCLES *situated within the Cavity of the ABDOMEN.*

THESE consist of a single muscle, and four pair.

## DIAPHRAGMA,

This broad thin muscle, which makes a complete septum between the thorax and abdomen, is concave below and convex above; the middle of it on each side reaching as high within the thorax of the skeleton as the fourth rib, and is commonly divided into two portions.

1. The Superior or Greater Muscle of the DIAPHRAGM,

*Arises*, by distinct fleshy fibres, from the  
car-



cartilago enfiformis, from the cartilages of the seventh, and of all the inferior ribs on both sides. The fibres from the cartilago enfiformis, and from the seventh and eighth ribs, run obliquely upwards and backwards; from the ninth and tenth, transversely inwards and upwards; and from the eleventh and twelfth, obliquely upwards.—From these different origins the fibres run, like radii from the circumference to the centre of a circle; and are

*Inserted* into a cordiform tendon, of a considerable breadth, which is situated in the middle of the diaphragm; and in which, therefore, the fibres from opposite sides are interlaced. Towards the right side the tendon is perforated by a triangular hole, for the passage of the vena cava inferior; and to the upper convex part of it the pericardium and mediastinum are connected.

2. The



2. The Inferior, Lesser Muscle, or Appendix of the  
DIAPHRAGM,

*Arises* from the second, third, and fourth lumbar vertebræ, by eight heads; of which two in the middle, commonly called its *crura*, are the longest, and begin tendinous. Between the crura, the aorta and thoracic duct pass; and, on the outside of these, the great sympathetic nerves and branches of the vena azygos perforate the shorter heads. The muscular fibres run obliquely upwards and forwards, and form in the middle two fleshy columns, which decussate and leave an oval space between them for the passage of the œsophagus and eighth pair of nerves.

*Inserted*, by strong fleshy fibres, into the posterior part of the middle tendon.

*Use*. The diaphragm is the principal agent in respiration, particularly in inspiration: for when it is in action, the fibres, from  
their



their different attachments, endeavour to bring themselves into a plane towards the middle tendon, by which the cavity of the thorax is enlarged, particularly at the sides, where the lungs are chiefly situated; and as the lungs must always be contiguous to the inside of the thorax and upper side of the diaphragm, the air rushes into them, in order to fill up the increased space. This muscle is assisted by the two rows of intercostals, which elevate the ribs, and the cavity of the thorax is more enlarged. In time of violent exercise, or whatever cause drives the blood with unusual celerity towards the lungs, the pectoral muscles, the *ferrati antichi majores*, the *ferrati postici superiores*, and *scaleni* muscles, are brought into action. And in laborious inspiration, the muscles which arise from the upper part of the thorax, when the parts into which they are inserted are fixed, likewise assist. In expiration, the diaphragm is relaxed and pushed up by the pressure of the abdominal



minal muscles upon the viscera of the abdomen; and at the same time that they press it upwards, they also, together with the sterno costales and serrati postici inferiores, pull down the ribs, and are assisted in a powerful manner by the elasticity of the cartilages that join the ribs to the sternum; by which the cavity of the thorax is diminished, and the air suddenly pushed out of the lungs: and, in laborious expiration, the quadrati lumborum, sacrolumbales, and longissimi dorsi, concur in pulling down the ribs.

The four pair are,

1. QUADRATUS LUMBORUM,

*Arises*, pretty broad, tendinous and fleshy, from the posterior part of the spine of the os ilium.

*Inserted* into the transverse processes of all the vertebræ of the loins, into the last rib near



near the spine, and by a small tendon into the side of the last vertebra of the back.

*Use.* To move the loins to one side, pull down the last rib, and, when both act, to bend the loins forwards.

*Quadratus, seu Lumbaris externus, Winflow.*

## 2. PSOAS PARVUS,

*Arises,* fleshy, from the sides of the two upper vertebræ of the loins, and sends off a small long tendon which ends thin and flat, and is

*Inserted* into the brim of the pelvis, at the junction of the os ilium and pubis.

*Use.* To assist the psoas magnus in bending the loins forwards; and, in certain positions, to assist in raising the pelvis.

*N. B.* This muscle is very often wanting.

## 3. PSOAS MAGNUS

*Arises,* fleshy, from the side of the body,  
and



and transverse process of the last vertebra of the back; and, in the same manner, from all those of the loins, by as many distinct slips.

*Inserted*, tendinous, into the trochanter minor of the os femoris; and fleshy into that bone, a little below the same trochanter.

*Use*. To bend the thigh forwards; or, when the inferior extremity is fixed, to assist in bending the body.

*Psoas*, seu *Lumbaris internus*, Winslow.

#### 4. ILIACUS INTERNUS,

*Arises*, fleshy, from the transverse process of the last vertebra of the loins, from all the inner lip of the spine of the os ilium, from the edge of that bone between its anterior superior spinous process and the acetabulum, and from most of the hollow part of the ilium. It joins with the psoas magnus, where it begins to become tendinous; and is

*Inserted* along with it.

*Use*.



*Use.* To assist the psoas in bending the thigh, and to bring it directly forwards.

*N. B.* The infertion of the two last muscles should not be prosecuted till the muscles of the thigh are dissected.

P. A. H. C.



## C H A P. XXI.

MUSCLES *situated on the Anterior*  
*Part of the* THORAX.

THESE may be divided into two layers.  
The first layer consists of one muscle named

## PECTORALIS MAJOR.

*Arises* from the cartilaginous extremitie of the fifth and sixth ribs, where it always intermixes with the external oblique muscle of the abdomen; from almost the whole length of the sternum, and from near half of the anterior part of the clavicle: The fibres run towards the axilla in a folding manner.

*Inserted*, by two broad tendons, which cross each other at the upper and inner part



## 110 DESCRIPTION OF

of the os humeri, above the insertion of the deltoid muscle, and outer side of the groove for lodging the tendon of the long head of the biceps.

*Use.* To move the arm forwards and obliquely upwards, towards the sternum.

*Pectoralis, Albinus.*

The second layer consists of three muscles;

### 1. SUBCLAVIUS.

*Arises* tendinous from the cartilage that joins the first rib to the sternum.

*Inserted*, after becoming fleshy, into the inferior part of the clavicle, which it occupies from within an inch or so of the sternum, as far outwards as to its connection, by ligament, with the coracoid process of the scapula.

*Use.* To pull the clavicle downwards and forwards.

### 2. PEC.



## 2. PECTORALIS MINOR,

*Arises*, tendinous and fleshy, from the upper edge of the third, fourth, and fifth ribs, near where they join with their cartilages.

*Inserted*, tendinous, into the coracoid process of the scapula; but soon grows fleshy and broad.

*Use*. To bring the scapula forwards and downwards, or to raise the ribs upwards.

*Serratus anticus*, Albinus.

*Serratus minor anticus*, Douglas.

## 3. SERRATUS MAGNUS,

*Arises* from the nine superior ribs, by an equal number of fleshy digitations, resembling the teeth of a saw.

*Inserted*, fleshy, into the whole base of the scapula internally, between the insertion of the rhomboid and the origin of the sub-



scapularis muscles, being folded about the two angles of the scapula.

*Use.* To move the scapula forwards; and, when the scapula is forcibly raised, to draw upwards the ribs.

*Serratus major anticus, Douglas.*

## CHAP.



## C H A P. XXII.

MUSCLES *situated between the Ribs,*  
*and within the THORAX.*

**B**ETWEEN the ribs on each side, there are eleven double rows of muscles, which are therefore named *intercostals*. These decussate each other like the strokes of the letter X.

*Arise* from the inferior acute edge of each superior rib, and run obliquely forwards, the whole length from the spine to near the joining of the ribs with their cartilages; from which, to the sternum, there is only a thin membrane covering the internal intercostals.

*Inserted* into the upper obtuse edge of each inferior rib, as far back as the spine into which the posterior portion is fixed.



## INTERCOSTALES INTERNI,

*Arise* in the same manner as the external: but they begin at the sternum, and run obliquely backwards, as far as the angle of the rib; and from that to the spine they are wanting.

*Inserted* in the same manner as the external.

*Use.* By means of these muscles, the ribs are equally raised upwards during inspiration. Their fibres being oblique, give them a greater power of bringing the ribs near each other, than could be performed by straight ones. But, by the obliquity of the fibres, they are almost brought contiguous: and, as the fixed points of the ribs are before and behind, if the external had been continued forwards to the sternum, and the internal backwards to the spine, it would have hindered their motion, which is greatest in the middle, though the obliquity of the  
ribs



ribs renders it less perceptible; and, instead of raising the fibres fixed to the sternum and spine, would have depressed the ribs.

*N. B.* The portions of the external intercostals which arise from the transverse processes of the vertebræ, where the ribs are fixed to them, and other portions that pass over one rib and terminate in the next below it, Albinus calls *Levatores costarum longiores et breviores*.

The portions of the internal that pass over one rib, and are inserted into the next below it, Douglas calls *Costarum depressores proprii Cowperii*.

These portions of both rows assist in raising the ribs in the same manner as the rest of the intercostals.

*Supra-costales*, and *Infra-costales*, Winslow.



The muscles within the thorax are one pair, viz.

TRIANGULARIS, or STERNO COSTALIS,

*Arises*, fleshy and a little tendinous, from all the length of the cartilago enfiformis laterally, and from the edge of the lower half of the middle bone of the sternum, from whence its fibres ascend obliquely upwards and outwards.

*Inserted*, generally by three triangular terminations, into the lower edge of the cartilages of the third, fourth, and fifth ribs, near where these join with the ribs.

*Use*. To depress these cartilages, and the extremities of the ribs; and consequently to assist in contracting the cavity of the thorax.

This muscle often varies; and is sometimes inserted into the cartilage of the second rib, sometimes into the cartilage of the sixth rib.



## C H A P. XXIII.

MUSCLES *situated on the Anterior Part of the NECK, close to the VERTEBRÆ.*

THESE consist of one layer, formed by four muscles.

## 1. LONGUS COLLI,

*Arises*, tendinous and fleshy, from the bodies of the three superior vertebræ of the back laterally; and from the transverse processes of the third, fourth, fifth, and sixth vertebræ of the neck, near their roots.

*Inserted* into the fore-part of the bodies of all the vertebræ of the neck, by as many small tendons, which are covered with flesh.

*Use.* To bend the neck gradually forwards, and to one side.



## 2. RECTUS CAPITIS INTERNUS MAJOR,

*Arises*, from the anterior points of the transverse processes of the third, fourth, fifth, and sixth vertebræ of the neck, by four distinct beginnings.

*Inserted* into the cuneiform process of the os occipitis, a little before the condyloid process.

*Use*. To bend the head forwards.

*Rectus anterior longus*, Winslow.

## 3. RECTUS CAPITIS INTERNUS MINOR,

*Arises*, fleshy, from the fore-part of the body of the first vertebra of the neck, opposite to the superior oblique process.

*Inserted* near the root of the condyloid process of the os occipitis, under, and a little farther outwards than, the former muscle.

*Use*. To nod the head forwards.

*Rectus anterior brevis*, Winslow.

## 4. REC-



## 4. RECTUS CAPITIS LATERALIS,

*Arises*, fleshy, from the anterior part of the point of the transverse process of the first vertebra of the neck.

*Inserted* into the os occipitis, opposite to the foramen stylo-mastoideum of the temporal bone.

*Use*. To bend the head a little to one side.

*Transversalis anticus primus*, Winflow.



## C H A P. XXIV.

MUSCLES *situated on the Posterior Part*  
*of the TRUNK.*

THESE may be divided into four layers, and a single pair.

The first layer consists of two muscles, which cover almost the whole posterior part of the trunk.

1. TRAPEZIUS, SEU CUCULARIS,

*Arises*, by a strong round tendon, from the lower part of the protuberance in the middle of the os occipitis behind; and, by a thin membranous tendon, which covers part of the splenius and complexus muscles, from the rough curved line that extends from the protuberance towards the mastoid process of the temporal bone; runs down



down along the nape of the neck, where it seems to arise from its fellow, and covers the spinous processes of the superior vertebræ of the neck; but arises from the spinous processes of the two inferior, and from the spinous processes of all the vertebræ of the back; adhering, tendinous, to its fellow, the whole length of its origin.

*Inserted*, fleshy, into the posterior half of the clavicle; tendinous and fleshy, into the acromion, and into almost all the spine of the scapula.

*Use*. Moves the scapula according to the three different directions of its fibres; for the upper descending fibres draw it obliquely upwards, the middle transverse straight fibres draw it directly backwards, and the inferior ascending fibres draw it obliquely downwards and backwards.

*N. B.* Where it is inseparably united to its fellow in the nape of the neck, it is named *Ligamentum Nuchæ* or *Colli*.



## 2. LATISSIMUS DORSI.

*Arises*, by a broad thin tendon, from the posterior part of the spine of the os ilium, from all the spinous processes of the os sacrum and vertebræ of the loins, and from the seven inferior ones of the vertebræ of the back; also, tendinous and fleshy, from the extremities of the three or four inferior ribs, a little beyond their cartilages, by as many distinct slips. The inferior fibres ascend obliquely, and the superior run transversely, over the inferior angle of the scapula, towards the axilla, where they are all collected, twisted, and folded.

*Inserted*, by a strong thin tendon, into the inner edge of the groove for lodging the tendon of the long head of the biceps.

*Use*. To pull the arm backwards and downwards, and to roll the os humeri.

*N. B.* The insertion of this muscle should not be prosecuted till the muscles of the os hu-



humeri, to which it belongs, are dissected.

The second layer consists of three pair, two on the back, and one on the neck.

On the back,

1. SERRATUS POSTICUS INFERIOR,

*Arises*, by a broad thin tendon, in common with that of the latissimus dorsi, from the spinal processes of the two inferior vertebræ of the back, and from the three superior of the loins.

*Inserted* into the lower edges of the four inferior ribs, at a little distance from their cartilages, by as many distinct fleshy lips.

*Use*. To depress the ribs into which it is inserted.

2. RHOMBOIDEUS,

This muscle is divided into two portions.

1. *Rhomboideus major*, *arises*, tendinous,  
from



from the spinous processes of the five superior vertebræ of the back.

*Inserted* into all the basis of the scapula below its spine.

*Use.* To draw the scapula obliquely upwards, and directly inwards.

2. *Rhomboideus minor, arifes*, tendinous, from the spinous processes of the three inferior vertebræ of the neck, and from the ligamentum nuchæ.

*Inserted* into the base of the scapula, opposite to its spine.

*Use.* To assist the former.

On the neck,

### 3. SPLENIUS.

*Arifes*, tendinous, from the four superior spinous processes of the vertebræ of the back; tendinous and fleshy, from the five inferior of the neck, and adheres firmly to the ligamentum nuchæ. At the third vertebra of the neck, the splenii recede from  
each



each other, so that part of the complexus muscle is seen.

*Inserted*, by as many tendons, into the five superior transverse processes of the vertebræ of the neck; and tendinous and fleshy, into the posterior part of the mastoid process, and into the os occipitis, where it joins with the root of that process.

*Use*. To bring the head and upper vertebræ of the neck backwards laterally; and, when both act, to pull the head directly backwards.

*N. B.* Albinus divides this muscle into two; viz. That portion which arises from the five inferior spinous processes of the neck, and is inserted into the mastoid process and os occipitis, he calls *Splenius Capitis*; and that portion which arises from the third and fourth of the back, and is inserted into the five superior transverse processes of the neck, is called by him *Splenius Colli*.

The



The single pair,

SERRATUS SUPERIOR POSTICUS,

*Arises*, by a broad thin tendon, from the spinous processes of the three last vertebræ of the neck, and the two uppermost of the back.

*Inserted* into the second, third, fourth, and fifth ribs, by as many fleshy slips.

*Use*. To elevate the ribs, and dilate the thorax.

The third layer consists of three pair on the back, and three on the neck.

On the back,

1. SPINALIS DORSI,

*Arises* from the spinous processes of the two uppermost vertebræ of the loins, and the three inferior of the back, by as many tendons.

*Inserted* into the spinous processes of the  
nine.



nine uppermost vertebræ of the back, except the first, by as many tendons.

*Use.* To erect and fix the vertebræ, and to assist in raising the spine.

## 2. LONGISSIMUS DORSI,

*Arises*, tendinous without, and fleshy within, from the side, and all the spinous processes of the os sacrum; from the posterior spine of the os ilium; from all the spinous processes; and from the roots of the transverse processes of the vertebræ of the loins.

*Inserted* into all the transverse processes of the vertebrae of the back, chiefly by small double tendons; also, by a tendinous and fleshy slip, into the lower edge of all the ribs, except the two inferior, at a little distance from their tubercles.

*Use.* To extend the vertebræ, and to raise and keep the trunk of the body erect.

*N. B.* From the upper part of this muscle there runs up a round fleshy portion which joins with the cervicalis descendens.

## 3. SA-



## 3. SACRO-LUMBALIS,

*Arises*, in common with the longissimus dorsi.

*Inserted* into all the ribs, where they begin to be curved forwards, by as many long and thin tendons ; and,

From the upper part of the fix or eight lower ribs, arise as many bundles of thin fleshy fibres, which soon terminate in the inner-side of the muscle, and are named *Musculi ad Sacro-lumbalem Accessorii*.

*Use*. To pull the ribs down, and assist to erect the trunk of the body.

*N. B.* There is a fleshy slip which runs from the upper part of this muscle into the fourth, fifth, and sixth transverse processes of the vertbræ of the neck, by three distinct tendons : it is named *Cervicalis Descendens* ; and its *use* is to turn the neck obliquely backwards, and to one side.

On



On the neck,

1, COMPLEXUS,

*Arises* from the transverse processes of the seven superior vertebræ of the back, and four inferior of the neck; by as many distinct tendinous origins; in its ascent it receives a fleshy slip from the spinous process of the first vertebra of the back: from these different origins it runs upwards, and is every where intermixed with tendinous fibres.

*Inserted*, tendinous and fleshy, into the inferior edge of the protuberance in the middle of the os occipitis, and into a part of the curved line that runs forwards from that protuberance.

*Use*. To draw the head backwards, and to one side; and, when both act, to draw the head directly backwards.

*N. B.* The long portion of this muscle that is situated next the spinous processes,  
lies



lies more loose, and has a roundish tendon in the middle of it; for which reason Albinus calls it *Biventer cervicis*.

## 2. TRACHELO-MASTOIDEUS,

*Arises* from the transverse processes of the three uppermost vertebræ of the back, and from the five lowermost of the neck, where it is connected to the transversalis cervicis, by as many thin tendons, which unite into a belly, and run up under the splenius.

*Inserted* into the middle of the posterior side of the mastoid process, by a thin tendon.

*Use.* To assist the complexus; but it pulls the head more to a side.

*Complexus minor, seu Mastoideus lateralis,*  
Winflow.

*Trachelo-mastoideus, seu Capitis par terminium Fallopii,* Douglas.



## 3. LEVATOR SCAPULÆ,

*Arises*, tendinous and fleshy, from the transverse processes of the five superior vertebræ of the neck by as many distinct slips, which soon unite to form a muscle that runs downwards and outwards.

*Inserted*, fleshy, into the superior angle of the scapula.

*Use*. To pull the scapula upwards, and a little forwards.

*Angularis, vulgo Levator proprius*, Winslow.

*Elevator, seu Musculus patientiæ*, Douglas.

The fourth layer consists of two pair on the back, two on the posterior part of the neck, four small pair situated immediately below the posterior part of the occiput, and three on the side of the neck.

On



On the back,

1. SEMI-SPINALIS DORSI,

*Arises*, from the transverse processes of the seventh, eighth, ninth, and tenth vertebræ of the back, by as many distinct tendons, which soon grow fleshy, and then become tendinous again ; and are

*Inserted* into the spinous processes of all the vertebræ of the back above the eighth, and into the two lowermost of the neck, by as many tendons.

*Use*. To extend the spine obliquely backwards.

*Semispinalis externus*, seu *Transverso-spinalis dorsi*, Winflow.

1. MULTIFIDUS SPINÆ,

*Arises* from the side and spinous processes of the os sacrum, and from the posterior part of the os ilium, where it joins with  
the



the sacrum; from all the oblique and transverse processes of the vertebræ of the loins; from all the transverse processes of the vertebræ of the back, and from those of the neck, except the three first, by as many distinct tendons, which soon grow fleshy, run in an oblique direction, and are

*Inserted*, by distinct tendons into all the spinous processes of the vertebræ of the loins, of the back, and of the neck, except the first.

*Use*. When the different portions of this muscle act on one side, they extend the back obliquely, or move it laterally; but, if they act together on both sides, they extend the vertebræ backwards.

*Transverso-spinalis lumborum, veterib. Sacer.*

*Semi-spinalis internus, five Transverso-spinalis dorsi.*

*Semi-spinalis, five Transverso-spinalis colli.*

*Pars interna, Winslow.*

*Transversalis lumborum, vulgo Sacer.*



*Transversalis dorsi.*

*Transversalis colli*, Douglas.

On the posterior part of the neck,

### 1. SEMI-SPINALIS COLLI,

*Arises*, from the transverse processes of the uppermost six vertebræ of the back, by as many distinct tendons, ascending obliquely under the complexus.

*Inserted* into the spinous processes of all the vertebræ of the neck, except the first and the last.

*Use.* To extend the neck obliquely backwards.

*Semi-spinalis*, five *Transverso-spinalis colli*, Winflow.

*Spinalis cervicis*, Albinus.

*Spinalis*, Douglas.

### 2. TRANSVERSALIS COLLI,

*Arises* from the transverse processes of the five uppermost vertebræ of the back, by as many



many tendinous and fleshy origins; runs between the trachelo-mastoideus, and splenius colli and cervicalis descendens.

*Inserted* into the transverse processes of all the cervical vetebæ, except the first and the last.

*Use.* To turn the neck obliquely backward, and a little to one side.

Below the posterior part of the occiput,

#### 1. RECTUS CAPITIS POSTICUS MAJOR,

*Arises*, fleshy, from the external part of the spinous process of the second vertebra of the neck; and grows broader in its ascent, which is not straight, but obliquely outwards.

*Inserted*, tendinous and fleshy, into the os occipitis, near the rectus capitis lateralis, and the insertion of the obliquus capitis superior.



*Use.* To pull the head backwards, and to assist a little in its rotation.

*Rectus major*, Winslow and Douglas.

## 2. RECTUS CAPITIS POSTICUS MINOR,

*Arises*, by a narrow beginning, close by its fellow, from a little protuberance in the middle of the back-part of the first vertebra of the neck, its outer edge being covered by the rectus major.

*Inserted*, pretty broad, into the sides of a dimple in the os occipitis, near its foramen magnum.

*Use.* To assist the rectus major in moving the head backwards.

*Obliquus minor*, Winslow and Douglas.

## 3. OBLIQUUS CAPITIS SUPERIOR,

*Arises* from the transverse process of the first vertebra of the neck.

*Inserted*, tendinous and fleshy, into the

os



os occipitis behind the back-part of the mastoid process of the temporal bone, and under the insertion of the complexus muscle.

*Use.* To draw the head backwards.

*Obliquus major*, Winslow.

*Obliquus superior*, Douglas.

#### 4. OBLIQUUS CAPITIS INFERIOR,

*Arises*, fleshy, from the spinous process of the second vertebra of the neck, its whole length; and, forming a thick fleshy belly, is

*Inserted* into the transverse process of the first vertebra of the neck.

*Use.* To give a rotatory motion to the head.

On the side of the neck,

#### 1. SCALENUS ANTICUS,

*Arises* from the fourth, fifth, and sixth transverse processes of the first vertebra of the neck, by as many tendons.

*Inserted* tendinous and fleshy, into the



upper side of the first rib, near its cartilage.

*Scalenus prior*, Albinus.

*Anterior portion of the first scalenus*, Winflow.

*First scalenus*, Douglas.

## 2. SCALENUS MEDIUS,

*Arises*, from all the transverse processes of the vertebræ of the neck, by as many strong tendons; the nerves to the superior extremity pass between it and the former.

*Inserted* into the upper and outer part of the first rib, from its root, to within the distance of an inch from its cartilage.

*Posterior portion of the first scalenus*, Winflow.

*Second scalenus*, Douglas.

## 3. SCALENUS POSTICUS,

*Arises* from the fifth and sixth transverse processes of the vertebræ of the neck.

*In-*



*Inserted* into the upper edge of the second rib, not far from the spine.

*Posterior portion of the second scalenus*, Winflow.

*Third scalenus*, Douglas.

*Use* of the three scaleni: To bend the neck to one side; or, when the neck is fixed, to elevate the ribs, and to dilate the thorax.

There are a number of small muscles situated between the spinous and transverse processes of contiguous vertebræ, which are accordingly named,

#### 1. INTERSPINALES COLLI.

The space between the spinous processes of the vertebræ of the neck, most of which are bifurcated, is filled up with fleshy portions; which

*Arise*, double, from the spinous process of the inferior vertebræ of the neck; and ascend to be

*Inserted*, in the same manner, into the



spinous process of the superior vertebra. They are five in number.

*Use.* To draw these processes nearer to each other.

## 2. INTERTRANSVERSALES COLLI:

They begin from the transverse process of the first vertebra of the back, and fill up the spaces between the transverse processes of the vertebræ of the neck, which are likewise bifurcated; and, consequently, there are six distinct double muscles, which

*Arise* from the inferior transverse process of each vertebra of the neck, and first of the back, and are

*Inserted* into the superior transverse processes.

*Use.* To draw these processes towards each other, and turn the neck a little to one side.



3, 4, 5. INTERSPINALES DORSI et LUMBORUM, and the INTERTRANSVERSALES DORSI.

Are rather small tendons than muscles, serving to connect the spinal and transverse processes.

6. INTERTRANSVERSALES LUMBORUM,

Are four distinct small bundles of flesh, which fill up the spaces between the transverse processes of the vertebræ of the loins, and serve to draw them towards each other.



## C H A P. XXV.

MUSCLES *of the* SUPERIOR EXTRE-  
MITIES.

**T**HESE may be divided into the muscles that are situated on the scapula, on the os humeri, on the cubit or fore-arm, and on the hand.

Muscles situated on the scapula.

These are called muscles of the *os humeri*; and are three behind, one along its inferior costa, two before and one beneath it.

Behind,

## 1. SUPRASPINATUS,

*Arises*, fleshy, from all that part of the base of the scapula that is above its spine; also from the spine and superior costa; passes

un-



under the acromion, and adheres to the capsular ligament of the os humeri.

*Inserted*, tendinous, into that part of the large protuberance on the head of the os humeri that is next the groove for lodging the tendon of the long head of the biceps.

*Use*. To raise the arm upwards; and, at the same time, to pull the capsular ligament from between the bones, that it may not be pinched.

## 2. INFRASPINATUS,

*Arises*, fleshy, from all that part of the base of the scapula that is between its spine and inferior angle; from the spine, as far as the cervix of the scapula. The fibres ascend and descend obliquely towards a tendon in the middle of the muscle, which runs forwards, and adheres to the capsular ligament.

*Inserted*, by a thick and short tendon, into the upper and middle part of the large



protuberance on the head of the os humeri.

*Use.* To roll the humerus outwards; to assist in raising, and in supporting it when raised; and to pull the ligament from between the bones.

*N. B.* These two muscles are covered with a tendinous membrane, from which a number of their fleshy fibres arise. It serves besides to strengthen their actions, and keeps them from swelling too much outwardly when in action.

### 3. TERES MINOR,

*Arises,* fleshy, from all the round edge of the inferior costa of the scapula, and runs forwards along the inferior edge of the infraspinatus muscle, and adheres to the ligament.

*Inserted,* tendinous, into the back part of the large protuberance on the head of the

os



os humeri, a little behind and below the termination of the last-named muscle.

*Use.* To roll the humerus outwards; to draw the humerus backwards; and to prevent the ligament from being pinched between the bones.

Along the inferior costa of the scapula,

#### TERES MAJOR,

*Arises*, fleshy, from the inferior angle of the scapula, and from all that portion of its inferior costa that is rough and thicker than the rest; its fleshy fibres are continued over part of the infraspinatus muscle, to which they firmly adhere.

*Inserted*, by a broad, short, and thin tendon, into the ridge at the inner side of the groove for lodging the tendon of the long head of the biceps, along with the latissimus dorsi.

*Use.* To roll the humerus inwards, and to draw it backwards and downwards.

The



The two before the scapula.

1. DELTOIDES,

*Arises*, fleshy, from all the posterior part of the clavicle that the pectoralis major does not possess; tendinous and fleshy, from the acromion, and lower margin of almost the whole spine of the scapula opposite to the insertion of the cucullaris muscle: from these origins it runs in three different directions, i. e. from the clavicle outwards and downwards; from the spine of the scapula outwards, forwards, and downwards; and from the acromion straight downwards; and is composed of a number of fasciculi, which form a strong fleshy muscle that covers the anterior part of the joint of the os humeri.

*Inserted*, tendinous, into a rough protuberance in the outer side of the os humeri, near its middle, where the fibres of this muscle intermix with some part of the brachialis externus.

*Use.*



*Use.* To pull the arm directly outwards and upwards, and a little forwards or backwards, according to the different directions of its fibres.

2. CORACO-BRACHIALIS,

*Arises*, tendinous and fleshy, from the fore-part of the coracoid process of the scapula; adhering, in its descent, to the short head of the biceps.

*Inserted*, tendinous and fleshy, about the middle of the internal part of the os humeri, near the origin of the third head of the triceps, called *brachialis externus*, where it sends down a thin tendinous expansion to the internal condyle of the os humeri.

*Use.* To raise the arm upwards and forwards.

*N. B.* There passes a nerve through this muscle, called *Musculo-cutaneus*.

The



The one beneath the scapula.

SUPSCAPULARIS,

*Arises*, fleshy, from all the base of the scapula internally, and from its superior and inferior costæ, being composed of a number of tendinous and fleshy fasciculi, which make prints on the bone; they all join together, fill up the hollow of the scapula, and pass over the joint, adhering to the capsular ligament.

*Inserted*, tendinous, into the upper part of the internal protuberance at the head of the os humeri.

*Use*. To roll the humerus inwards, and to draw it to the side of the body; and to prevent the capsular ligament from being pinched.



## C H A P. XXVI.

MUSCLES *situated on the Os HUMERI.*

THESE are called

*Muscles of the Cubit or Fore-arm,*

They consist of two before, and two behind.

Before,

## 1. BICEPS FLEXOR CUBITI,

*Arises*, by two heads. The first and outermost, called *longus*, begins tendinous from the upper edge of the glenoid cavity of the scapula; passes over the head of the os humeri within the joint; and, in its descent without the joint, is inclosed in a groove near the head of the os humeri, by



a membranous ligament that proceeds from the capsular ligament and adjacent tendons. The second, or innermost head, called *brevi*, arises, tendinous and fleshy, from the coracoid process of the scapula, in common with the coraco-brachialis muscle. A little below the middle of the fore-part of the os humeri, these heads unite.

*Inserted*, by a strong roundish tendon, into the tubercle on the upper end of the radius internally.

*Use*. To turn the hand supine, and to bend the fore-arm.

N. B. At the bending of the elbow, where it begins to grow tendinous, it sends off an aponeurosis, which covers all the muscles on the inside of the fore-arm, and joins with another tendinous membrane, which is sent off from the triceps extensor cubiti, and covers all the muscles on the outside of the fore-arm; and a number of the fibres, from opposite fibres, decussate each other. It serves to strengthen the muscles,



muscles, by keeping them from swelling too much outwardly, when in action; and a number of their fleshy fibres take their origin from it.

*Biceps brachii*, Albinus.

*Coraco-radialis*, seu *biceps*, Winslow.

*Biceps internus*, Douglas.

## 2. BRACHIALIS INTERNUS,

*Arises*, fleshy, from the middle of the os humeri, at each side of the insertion of the deltoid muscle, covering all the inferior and fore part of this bone, runs over the joint, and adheres firmly to the ligament.

*Inserted*, by a strong short tendon, into the coronoid process of the ulna.

*Use*. To bend the fore-arm, and to prevent the capsular ligament of the joint from being pinched.

*Brachialis*, Winslow.



Behind,

1. TRICEPS EXTENSOR CUBITI,

*Arises*, by three heads; the first called *longus*, pretty broad and tendinous, from the inferior costa of the scapula, near its cervix. The second head, called *brevis*, arises by an acute, tendinous, and fleshy beginning, from the back part of the os humeri, a little below its head, outwardly. The third, called *brachialis externus*, arises by an acute beginning, from the back part of the os humeri. These three heads unite lower than the insertion of the *teres major*, and cover the whole posterior part of the humerus, from which they receive addition in their descent.

*Inserted* into the upper and external part of the process of the ulna, called *olecranon*, and partly into the condyles of the os humeri, adhering firmly to the ligament.

*Use.*



*Use.* To extend the fore-arm.

*Anconeus major, Anconeus externus, and  
Anconeus internus, Winflow.*

*Biceps externus and Brachialis externus,  
Douglas.*

### 3. ANCONEUS,

*Arises,* tendinous from the posterior part of the external condyle of the os humeri; it soon grows fleshy, and is continued from the third head of the triceps.

*Inserted,* fleshy and thin, into a ridge on the outer and posterior edge of the ulna, being continued some way below the olecranon, and covered with a tendinous membrane.

*Use.* To assist in extending the fore-arm.

*Anconeus minor, Winflow.*

*Anconeus, vel Cubitalis Riolani. Douglas.*



## C H A P. XXVII.

MUSCLES *situated on the* CUBIT *or*  
FORE-ARM\*.

THESE may be divided into three classes: *first*, flexors and extensors of the whole *hand*; *second*, flexors and extensors of the *fingers*; and, *third*, supinators and pronators, or those that roll the *radius* on the *ulna*.

*First class* consists of three flexors, and three extensors.

\*In the following description, the arm is supposed to hang by the side with the palm turned forwards; so that the radius and thumb are upon its outer side, and the ulna and little finger upon its inner side; whereas, when the muscles are described in the less straining posture of pronation, as has been generally done by authors, the utmost confusion is necessarily introduced in the application of the terms *outer* and *inner*, from the decussation of the radius and ulna.

Flexors:



Flexors :

1. PALMARIS LONGUS,

*Arises*, tendinous, from the internal condyle of the os humeri, soon grows fleshy, and, after a short progress, sends off a long slender tendon.

*Inserted* into the ligamentum carpi annulare, and into a tendinous membrane that is expanded on the palm of the hand, named *aponeurosis palmaris* ; which, above, begins at the transverse or annular ligament of the wrist, and, below, is fixed to the roots of the fingers.

*Use*. To bend the hand, and to stretch the membrane that is expanded on the palm.

*Ulnaris gracilis*, Winslow.

*N. B.* This muscle is sometimes wanting ; but the *aponeurosis palmaris* is always to be found, and a small muscle named

PAL-



## PALMARIS BREVIS,

*Arises* from the ligamentum carpi annulare, and tendinous membrane that is expanded on the palm of the hand.

*Inserted*, by small bundles of fleshy fibres, into the skin and fat that covers the abductor minimi digiti, and into the os pisiforme.

*Use*. To assist in contracting the palm of the hand.

*Palmaris cutaneus*, Winslow.

## 2. FLEXOR CARPI RADIALIS,

*Arises*, tendinous and fleshy, from the internal condyle of the os humeri, and from the anterior part of the upper end of the ulna, where it firmly adheres to the pronator radii teres.

*Inserted*, by a flat tendon, into the fore and upper part of the metacarpal bone that sustains



sustains the fore-finger, after running thro' a fossa in the os trapezium.

*Use.* To bend the hand, and to assist in its pronation.

*Radialis internus*, Albinus and Winflow.

### 3. FLEXOR CARPI ULNARIS,

*Arises*, tendinous, from the internal condyle of the os humeri. It has likewise a small fleshy beginning from the outer side of the olecranon; between which and the condyle the ulnar nerve passes to the forearm; and a number of its fleshy fibres arise from the tendinous membrane which covers the forearm.

*Inserted*, by a short strong tendon, into the os pisiforme; at a little distance from its insertion, a small ligament is sent off to the metacarpal bone that sustains the little finger.

*Use.* To assist the former in bending the hand.

*Ulnaris internus*, Albinus and Winflow.



## Extensfors :

## 1. EXTENSOR CARPI RADIALIS LONGIOR,

*Arises*, broad, thin, and fleshy, immediately below the supinator radii longus, from the lower part of the external ridge of the os humeri, above its external condyle.

*Inserted* by a round tendon, into the posterior and upper part of the metacarpal bone that sustains the fore-finger.

*Use*. To extend and bring the hand backwards.

*Radialis externus longior*, Albinus.

*Radialis externus primus*, Winslow.

## 2. EXTENSOR CARPI RADIALIS BREVIOR,

*Arises*, tendinous, from the external condyle of the os humeri, and from the ligament



ment that connects the radius to it, and runs along the outside of the radius.

*Inserted*, by a round tendon, into the upper and back part of the metacarpal bone that sustains the middle finger.

*Use*. To assist the last mentioned muscle.

*Radialis externus brevior*, Albinus.

*Radialis secundus*, Winflow.

#### EXTENSOR CARPI ULNARIS,

*Arises*, tendinous, from the external condyle of the os humeri ; and, in its progress, fleshy from the middle of the ulna, where it passes over it. Its round tendon is inclosed by a membranous sheath, in a groove which is situated at the extremity of the ulna.

*Inserted*, by its round tendon, into the posterior and upper part of the metacarpal bone that sustains the little-finger.

*Use*. To assist the former in extending the hand.

*Ulnaris externus*, Albinus and Winflow.



*Second Class.*

The flexors and extensors of the four fingers are, two long, and one small flexor to each finger, and one extensor.

## 1. FLEXOR SUBLIMIS PERFORATUS,

*Arises*, tendinous and fleshy, from the internal condyle of the os humeri; tendinous from the coronoid process of the ulna, near the edge of the cavity that receives the head of the radius; fleshy from the tubercle of the radius; and membranous and fleshy from the middle of the fore-part of the radius, where the flexor pollicis longus arises. Its fleshy belly sends off four round tendons before it passes under the ligament of the wrist.

*Inserted* into the anterior and upper part of the second bone of each finger, being, near the extremity of the first bone, divided for the passage of the perforans.

*Use.*



*Use.* To bend the second joint or phalanx of the fingers.

*Sublimis*, Albinus.

*Perforatus*, Douglas.

2. FLEXOR PROFUNDUS PERFORANS,

*Arises*, fleshy, from the external side, and upper part of the ulna, for some way downwards, and from a large share of the interosseous ligament. It splits into four tendons, a little before it passes under the ligamentum carpi annulare; and these pass through the slits in the tendons of the flexor sublimis.

*Inserted* into the fore and upper part of the third or last bone of all the four fingers.

*Use.* To bend the last joint of the fingers.

*Profundus*, Albinus.

*Perforans*, Douglas.



The four small flexors are named

LUMBRICALES,

*Arise*, thin and fleshy, from the outside of the tendons of the flexor profundus, a little above the lower edge of the ligamentum carpi annulare.

*Inserted*, by long slender tendons, into the outer sides of the broad tendons of the interossei muscles, about the middle of the first joint.

*Use*. To increase the flexion of the fingers while the long flexors are in full action.

Extensors :

EXTENSOR DIGITORUM COMMUNIS,

*Arises*, by an acute, tendinous, and fleshy beginning, from the external condyle of the os humeri, where it adheres to the supinator radii brevis. Before it passes under the ligamentum carpi annulare externum, it splits into four tendons ; some of which  
may



may be divided into several smaller; and about the fore-part of the metacarpal bones they remit tendinous filaments to each other.

*Inserted* into the posterior part of all the bones of the four fingers, by a tendinous expansion.

*Use.* To extend all the joints of the fingers.

### *Third Class.*

Consists of four muscles, viz. two *supinators*, and two *pronators*.

Supinators :

#### 1. SUPINATOR RADII LONGUS,

*Arises*, by an acute and fleshy origin, from the external ridge of the os humeri, above the external condyle, near as far up as the middle of that bone.

*Inserted* into the outer-side of the inferior extremity of the radius.



*Use.* To roll the radius outwards, and consequently the palm of the hand upwards.

*Supinator longus*, Albinus, Winflow, and Douglas.

## 2. SUPINATOR RADII BREVIS,

*Arises*, tendinous, from the external condyle of the os humeri ; tendinous and fleshy, from the external and upper part of the ulna, and adheres firmly to the ligament that joins these two bones.

*Inserted*, into the head, neck, and tubercle of the radius, near the insertion of the biceps, and ridge running from that downwards and outwards.

*Use.* To roll the radius outwards, and so bring the hand supine.

Pronators :

## 1. PRONATOR RADII TERES,

*Arises*, fleshy, from the internal condyle  
of



of the os humeri, and tendinous from the coronoid process of the ulna.

*Inserted*, thin, tendinous, and fleshy, into the middle of the posterior part of the radius.

*Use*. To roll the radius, together with the hand, inwards.

## 2. PRONATOR RADII QUADRATUS,

*Arises*, broad, tendinous, and fleshy from the lower and inner part of the ulna ; the fibres run transversely, to be

*Inserted* into the lower and anterior part of the radius, opposite to its origin.

*Use*. To turn the radius, together with the hand, inwards.



## C H A P. XXVIII.

MUSCLES *situated on the HAND chiefly.*

THESE may be divided into four classes, viz. muscles of the *thumb, fore-finger, little-finger, and metacarpal bones.*

*Muscles of the Thumb.*

These consist of *three flexors, three extensors, one abductor, and one adductor.*

Flexors:

1. FLEXOR LONGUS POLLICIS MANUS,

*Arises,* by an acute fleshy beginning, from the upper part of the radius, immediately below its tubercle, and is continued down for some space on the fore-part of this bone. It has likewise generally another origin from the internal condyle of the



the os humeri, which forms a distinct fleshy slip that terminates near the upper part of the origin from the radius.

*Inserted* into the last joint of the thumb, after having passed its tendon under the ligament of the wrist.

*Use.* To bend the last joint of the thumb.

*Flexor tertii internodii*, Douglas.

## 2. FLEXOR BREVIS POLLICIS MANUS,

*Arises*, from the os trapezoides, magnum, and unciforme of the carpus, and is divided into two portions by the tendon of the flexor pollicis longus.

*Inserted* into the ossa sesamoidea and first bone of the thumb.

*Use.* To bend the first joint of the thumb.

*Flexor secundi internodii*, Douglas.

## 3. FLEXOR OSSIS METACARPI POLLICIS, OR OPPONENS POLLICIS,

*Arises*, fleshy, from the os trapezium and



ligamentum carpi annulare, lying under the abductor pollicis.

*Inserted*, tendinous and fleshy, into the under and anterior part of the metacarpal bone of the thumb.

*Use*. To bring the thumb inwards, opposite to the other fingers.

*Flexor primi internodii*, Douglas.

Extensors :

1. EXTENSOR OSSIS METCARPI POLLICIS MANUS,

*Arises*, fleshy, from the middle and posterior part of the ulna, immediately below the insertion of the anconæus muscle, from the posterior part of the middle of the radius, and from the interosseous ligament.

*Inserted*, generally, by two tendons, into the os trapezium, and upper back-part of the metacarpal bone of the thumb, and often joins with the abductor pollicis.

*Use*.



*Use.* To extend the metacarpal bone of the thumb outwardly.

*Abductor longus pollicis manus*, Albinus.

*Extensor primi internodii*, Douglas.

## 2. EXTENSOR PRIMI INTERNODII,

*Arises*, fleshy, from the posterior part of the ulna near the former muscle, and from the interosseous ligament.

*Inserted*, tendinous into the posterior part of the first bone of the thumb; and part of it may be traced as far as the second bone.

*Use.* To extend the first bone of the thumb obliquely outwards.

*Extensor minor pollicis manus*, Albinus.

This and the preceding muscle is called

*Extensor pollicis primus*, Winflow.

*Extensor secundi internodii*, Douglas.

## 3. EXTENSOR SECUNDI INTERNODII,

*Arises*, by an acute, tendinous, and fleshy  
be-



beginning, from the middle back part of the ulna, and from the interosseous ligament; its tendon runs through a small groove at the inner and back part of the lower end of the radius.

*Inserted* into the last bone of the thumb.

*Use.* To extend the last joint of the thumb obliquely backwards.

*Extensor major pollicis manus*, Albinus.

*Extensor pollicis secundus*, Winflow.

*Extensor tertii internodii*, Douglas.

#### ABDUCTOR POLLICIS MANUS,

*Arises*, by a broad, tendinous, and fleshy beginning, from the ligamentum carpi annulare, and from the os trapezium.

*Inserted*, tendinous, into the outer side of the root of the first bone of the thumb.

*Use.* To draw the thumb from the fingers.

N. B. Albinus names the inner portion of this muscle *abductor brevis alter*.

*Abductor, Thenar Riolani*, Douglas.



## ADDUCTOR POLLICIS MANUS,

*Arises*, fleshy from almost the whole length of the metacarpal bone that sustains the middle finger; from thence its fibres are collected together.

*Inserted*, tendinous, into the inner part of the root of the first bone.

*Use*. To pull the thumb towards the fingers.

*Adductor ad minimum digitum*, Douglas.

Fore-finger,

## INDICATOR,

*Arises*, by an acute fleshy, beginning, from the middle of the posterior part of the ulna; its tendon passes under the same ligament with the extensor digitorum communis, with part of which it is



*Inserted* into the posterior part of the fore-finger.

*Extensor secundi internodii indicis proprius,*  
*vulgo Indicator, Douglas.*

#### ABDUCTOR INDICIS MANUS,

*Arises*, from the os trapezium, and from the superior part and inner side of the metacarpal bone of the thumb.

*Inserted*, by a short tendon, into the outer and back part of the first bone of the fore-finger.

*Use.* To bring the fore-finger towards the thumb.

*Semi interosseus, Winslow.*

Little-finger.

#### ABDUCTOR MINIMI DIGITI MANUS,

*Arises*, fleshy, from the os pisiforme, and from that part of the ligamentum carpi annulare next it.

*In-*



*Inserted*, tendinous, into the inner side of the upper end of the first bone of the little finger.

*Use*. To draw this finger from the rest.

*Hypothenar minor*, Winflow.

*Extensor tertii internodii minimi digiti*,  
Douglas.

#### ADDUCTOR METACARPI MINIMI DIGITI MANUS,

*Arises*, fleshy, from the thin edge of the os unciforme, and from that part of the ligament of the wrist next it.

*Inserted*, tendinous, into the inner side and anterior part of the metacarpal bone of this finger.

*Use*. To bend and bring the metacarpal bone of this finger towards the rest.

*Metacarpus*, Winflow.

*Flexor primi internodii minimi digiti*,  
Douglas.

#### FLEXOR



## FLEXOR PARVUS MINIMI DIGITI,

*Arises*, fleshy, from the outer side of the os unciforme, and from the ligament of the wrist which joins with that bone.

*Inserted*, by a roundish tendon, into the inner and anterior part of the upper end of the first bone of this finger.

*Use*. To bend the little finger, and assist the adductor.

*Abductor minimi digiti, Hypothenar Riolani, Douglas.*

Between the metacarpal bones, there are *four internal* and *three external* muscles, named *interossei*.

Interossei interni :

## 1. PRIOR INDICIS,

*Arises*, tendinous and fleshy, from the upper and outer part of the metacarpal bone that sustains the fore-finger.

*In-*



*Inserted* into the outside of that part of the tendinous expansion from the extensor digitorum communis, which covers the posterior part of the fore-finger.

*Use.* To draw the fore-finger inwards towards the thumb, and extend it obliquely.

*Extensor tertii internodii indicis*, Douglas.

## 2. POSTERIOR INDICIS,

*Arises*, tendinous and fleshy, from the root and inner part of the metacarpal bone that sustains the fore-finger.

*Inserted* into the inner side of the tendinous expansion which is sent off from the extensor digitorum communis, along the posterior part of the fore-finger.

*Use.* To extend the fore-finger obliquely, and to draw it outwards.

*First interosseus*, Douglas.

## 3. PRIOR



## 3. PRIOR ANNULARIS,

*Arises*, from the root of the outside of the metacarpal bone that sustains the ring-finger.

*Inserted* into the outside of the tendinous expansion of the extensor digitorum communis which covers the ring-finger.

*Use*. To extend and pull the ring-finger towards the thumb.

*Fourth interosseus*, Douglas.

## 4. INTEROSSEUS AURICULARIS,

*Arises*, from the root and outerside of the metacarpal bone of the little finger; and is

*Inserted*, into the outside of the tendinous expansion of the extensor digitorum communis, which covers the posterior part of the little finger.

*Use*.



*Use.* To extend and draw the little finger outwards.

*Sixth interosseus, Douglas.*

Interossei externi, seu bicipites :

### 1. PRIOR MEDII,

*Arises*, by two origins, from the roots of the metacarpal bones that sustain the fore and middle fingers externally, and next each other : Runs along the outside of the middle-finger ; and, being conspicuous on both sides of the hand, is

*Inserted* into the outside of the tendinous expansion from the extensor digitorum communis, which covers the posterior part of the middle-finger.

*Use.* To extend, and to draw the middle finger inwards.

*Second interosseus, Douglas.*

### 2. PO-



## 2. POSTERIOR MEDII,

*Arises*, by two origins, from the roots of the metacarpal bones, next each other, that sustain the middle and ring-fingers.

*Inserted* into the inside of the tendinous expansion from the extensor digitorum communis, which runs along the posterior part of the middle-finger.

*Use*. To extend and draw the middle-finger outwards.

*Third interosseus*, Douglas.

## 3. POSTERIOR ANNULARIS,

*Arises*, by two origins, from the roots of the metacarpal bones that sustain the ring and little fingers next each other.

*Inserted* into the inside of the tendinous expansion of the extensor digitorum communis, which runs along the posterior part of the ring-finger.

*Use*.



*Use.* To extend and draw the ring-finger inwards.

*Fifth interosseus, Douglas.*

*N. B.* The internal interossei are only conspicuous on the palm of the hand; but the external are apparent on both the palm and back of the hand.



## C H A P. XXIX.

MUSCLES *of the Inferior* EXTREMITIES.

THESE may be divided into the muscles *situated on the outside of the pelvis, on the thigh, on the leg, and on the foot.*

Muscles on the outside of the *pelvis*, which are called *muscles of the thigh.*

These are composed of *one layer before and three layers behind.*

The layer before consists of five muscles:

- |                      |                                  |
|----------------------|----------------------------------|
| 1. PSOAS MAGNUS.     | } These were formerly described. |
| 2. ILIACUS INTERNUS. |                                  |

*Vid. p. 106, 107.*

## 3. PECTINALIS,

*Arises, broad and fleshy, from the upper and*



and anterior part of the os pubis or pectinis, immediately above the foramen thyroideum.

*Inserted* into the anterior and upper part of the linea aspera of the os femoris, a little below the trochanter minor, by a flat and short tendon.

*Use.* To bring the thigh upwards and inwards, and to give it a degree of rotation outwards.

*Pectineus*, Albinus.

#### 4. TRICEPS ADDUCTOR FEMORIS,

Under this appellation are comprehended three distinct muscles.

##### 1. ADDUCTOR LONGUS FEMORIS,

*Arises*, by a pretty strong roundish tendon, from the upper and anterior part of the os pubis, and ligament of its synchondrosis, on the inner-side of the pectinalis.

*Inserted*, tendinous, near the middle of  
I the



the posterior part of the linea aspera, being continued for some way down.

*Adductor femoris primus*, Douglas.

*Triceps minus*, Winflow.

## 2. ADDUCTOR BREVIS FEMORIS,

*Arises*, tendinous, from the os pubis near its joining with the opposite os pubis below and behind the former.

*Inserted*, tendinous and fleshy, into the inner and upper part of the linea aspera, from a little below the trochanter minor, to the beginning of the insertion of the adductor longus.

*Adductor femoris secundus*, Douglas.

*Triceps secundus*, Winflow.

## 3. ADDUCTOR MAGNUS FEMORIS,

*Arises*, a little lower down than the former, near the symphysis of the ossa pubis; tendinous and fleshy, from the tuberosity of the  
the



the os ischium; the fibres run outwards and downwards.

*Inserted*, into almost the whole length of the linea aspera; into a ridge above the internal condyle of the os femoris; and, by a roundish long tendon, into the upper part of that condyle, a little above which the femoral artery takes a spinal turn towards the ham, passing between this muscle and the bone.

*Use* of these three muscles or triceps. To bring the thigh inwards and upwards, according to the different directions of their fibres; and, in some degree, to roll the thigh outwards.

*Adductor femoris tertius*, and

*Adductor femoris quartus*, Douglas.

*Triceps tertius*, Winflow.

## 5. OBTURATOR EXTERNUS.

*Arises*, fleshy, from the lower fore-part of the os pubis, and fore-part of the inner



crus of the ischium; surrounds the foramen thyroideum; a number of its fibres, arising from the membrane which fills up that foramen, are collected like rays towards a centre, and pass outwards around the root of the back-part of the cervix of the os femoris.

*Inserted*, by a strong tendon, into the cavity at the inner and back part of the root of the trochanter major, adhering in its course to the capsular ligament of the thigh-bone.

*Use*. To roll the thigh-bone obliquely outwards, and to prevent the capsular ligament from being pinched.

Behind :

*First Layer.*

GLUTEUS MAXIMUS,

*Arises*, fleshy, from the posterior part of the spine of the os ilium, a little higher up than the joining of the ilium with the os sacrum, from the whole external side of the os  
sa-



8101  
sacrum, below the posterior spinous process of the os ilium; from the posterior sacro-ischiatic ligament, over which part of the inferior edge of this muscle hangs in a folded manner; from the os coccygis. All the fleshy fibres run obliquely forwards and a little downwards, to form a thick broad muscle, which is divided into a number of strong fasciculi. The upper part of it covers almost the whole of the trochanter major, between which and the tendon of this muscle there is a large bursa mucosa, and where it is inseparably joined to the broad tendon of the *tensor vaginae femoris*.

*Inserted*, by a strong, thick, and broad tendon into the upper and outer part of the linea aspera, which is continued from the trochanter major, for some way downwards.

*Use*. To extend the thigh, by pulling it directly backwards, and a little outwards.

*Gluteus magnus*, Albinus.

*Gluteus major*, Cowper.



*Second Layer.*

## GLUTEUS MEDIUS,

*Arises*, fleshy, from the anterior superior spinous process of the os ilium, and from all the outer edge of the spine of the ilium, except its posterior part, where it arises from the dorsum of that bone.

*Inserted*, by a broad tendon, into the outer and posterior part of the trochanter major.

*Use*. To draw the thigh-bone outwards, and a little backwards; to roll the thigh-bone outwards, especially when it is bended.

N. B. The anterior and upper part of this muscle is covered by a tendinous membrane, from which a number of its fleshy fibres arise, and which joins with the broad tendons of the gluteus maximus, tensor vaginae femoris, and latissimus dorsi.

*Third*



*Third Layer consists of four Muscles.*

1. GLUTEUS MINIMUS,

*Arises*, fleshy, from a ridge that is continued from the superior anterior spinous process of the os ilium, and from the middle of the dorsum of that bone, as far back as its great nitch.

*Inserted*, by a strong tendon, into the fore and upper part of the trochanter major.

*Use*. To assist the former in pulling the thigh outwards and backwards, and in rolling it.

*Gluteus minor*, Albinus.

2. PYRIFORMIS,

*Arises*, within the pelvis, by three tendinous and fleshy origins, from the second, third, and fourth pieces of the os sacrum; from thence growing gradually narrower, it passes out of the pelvis along with the posterior crural nerve, below the nitch in the



posterior part of the os ilium, where it receives a few fleshy fibres.

*Inserted*, by a roundish tendon, into the upper part of the cavity at the inner side of the root of the trochanter major.

*Use*. To move the thigh a little upwards, and roll it outwards.

*Pyriformis, seu iliacus externus, Douglas.*

### 3. GEMINI,

*Arises*, by two distinct origins; the superior from the spinous process, and the inferior from the tuberosity of the os ischium; also, from the posterior sacro-ischiatic ligament. They are both united by a tendinous and fleshy membrane, and form a purse for the tendon of the obturator internus muscle, which was formerly described.

*Inserted*, tendinous and fleshy, into the cavity at the inner side of the root of the trochanter major, on each side of the tendon of the obturator internus, to which they firmly adhere.

*Use*.



*Use.* To roll the thigh outwards, and to preserve the tendon of the obturator internus from being hurt by the hardness of that part of the ischium over which it passes; also, to hinder it from starting out of its place, while the muscle is in action.

*Gemelli, Winslow.*

#### 4. QUADRATUS FEMORIS,

*Arises,* tendinous and fleshy, from the outside of the tuberosity of the os ischium; and, running transversely, is

*Inserted,* fleshy, into a rough ridge, continued from the root of the large trochanter, to the root of the small one.

*Use.* To roll the thigh outwards.



## CHAP. XXX.

MUSCLES *situated on the* THIGH.

THESE are called *muscles* of the *leg*; and consist of *one*, on the *outside*; *two*, on the *inside*; *four*, *before*; and, *four* *behind*.

Previous to the description of the muscles that are situated on the thigh and leg, it is necessary to take notice of a broad tendinous fascia or sheath, which is sent off from the back and from the tendons of the glutei and adjacent muscles.

It is a strong thick membrane on the outside of the thigh and leg; but, towards the inside of both, it gradually turns thinner, and has rather the appearance of a cellular substance, than a tendinous membrane. A little below the trochanter major, it is firmly  
fixed



fixed to the linea aspera; and, farther down, to that part of the head of the tibia that is next the fibula; where it sends off the tendinous expansion along the outside of the leg.

It serves to strengthen the action of the muscles, by keeping them firm in their proper places while in action, particularly the tendons that pass over the joints, where this membrane is thickest, and it gives origin to a number of the fleshy fibres of the muscles.

Outside,

TENSOR VAGINÆ FEMORIS,

*Arises*, by a narrow, tendinous, and fleshy beginning, from the external part of the anterior superior spinous process of the os ilium.

*Inserted*, a little below the trochanter major, into the inner side of the membranous fascia which covers the outside of the thigh.

*Use*. To stretch the membranous fascia, to assist



assist in the abduction of the thigh, and somewhat in its rotation inwards.

*Musculus aponeurosis, vel Fasciæ latae,*  
Winflow.

Inside,

1. SARTORIUS,

*Arises*, tendinous, from the anterior superior spinous process of the os ilium, soon grows fleshy, runs down for some space upon the rectus, and, going obliquely inwards, it passes over the vastus internus, and, about the middle of the os femoris, over part of the triceps, it runs down further between the tendon of the adductor magnus, and that of the gracilis muscle.

*Inserted*, by a broad and thin tendon, into the inner side of the tibia, near the inferior part of its tubercle.

*Use*. To bend the leg obliquely inwards, or to bring one leg across the other.

2. GRACILIS,

*Arises*, by a thin tendon, from the os pubis



bis near the symphysis of these two bones ; soon grows fleshy ; and, descending by the inside of the thigh, is

*Inserted*, tendinous, into the tibia under the sartorius.

*Use*. To assist the sartorius.

*Gracilis internus*, five *Rectus internus*, Winslow.

Before,

1. RECTUS,

*Arises*, fleshy from the inferior anterior spinous process of the os ilium, and tendinous from the dorsum of the ilium, a little above the acetabulum ; runs down over the anterior part of the cervix of the os femoris, the fibres not being straight, but running down like the plumage of a feather obliquely outwards and inwards, from a tendon in the middle.

*Inserted*, tendinous, into the upper part of the patella, from which a thin tendon runs down on the fore-part of this bone, to ter-



terminate in a thick strong ligament, which is sent off from the inferior part of the patella, and inserted into the tubercle of the tibia.

*Use.* To extend the leg, and in a powerful manner, by the intervention of the patella, like a pulley.

*Rectus five Gracilis anterior, Winflow.*

## 2. VASTUS EXTERNUS,

*Arises,* broad, tendinous, and fleshy, from the root of the trochanter major, and upper part of the linea aspera, its origin being continued from near the insertion of the gluteus minimus, the whole length of the linea aspera, by fleshy fibres which run obliquely forwards to a middle tendon, where they terminate.

*Inserted,* into a large share of the upper part of the patella; and part of it ends in an aponeurosis, which is continued down to the leg, and, in its passage, is firmly fixed to the head of the tibia.

*Use.*



*Use.* To extend the leg.

3. VASTUS INTERNUS,

*Arises*, tendinous and fleshy, from between the fore-part of the os femoris and root of the trochanter minor, and from almost all the inside of the linea aspera, by fibres running obliquely forwards and downwards.

*Inserted*, tendinous, into the upper and inside of the patella, continuing fleshy lower than the vastus externus. Part of it likewise ends in an aponeurosis continued down to the leg, and fixed in its passage to the upper part of the tibia.

*Use.* To extend the leg.

4. CRURALIS,

*Arises*, fleshy from between the two trochanters of the os femoris, but nearer the minor, and firmly adhering to most of the fore-



fore-part of the os femoris, and connected to both vasti muscles.

*Inserted*, tendinous, into the upper part of the patella, behind the rectus.

*Use*. To assist in the extension of the leg.

*Cruralis*, Albinus.

N. B. These four muscles *before*, being inserted into the patella, have the same effect upon the leg, as if they were immediately inserted into it, by means of the strong tendon, or rather ligament, which is sent off from the inferior part of the patella to the tibia.

Behind,

#### 1. SEMITENDINOSUS,

*Arises*, tendinous and fleshy, in common with the long head of the biceps, from the posterior part of the tuberosity of the os ischium; and sending down a long roundish tendon, which ends flat, is

*Inserted* into the inside of the ridge of the tibia, a little below its tubercle.

*Use*.



*Use.* To bend the leg backwards, and a little inwards.

*Seminervosus*, Winslow and Douglas.

## 2. SEMIMEMBRANOSUS,

*Arises*, tendinous, from the upper and posterior part of the tuberosity of the os ischium, sends down a broad flat tendon, which ends in a fleshy belly, and, in its descent, runs at first on the fore-part of the biceps, and, lower, between it and the semitendinosus.

*Inserted*, tendinous, into the inner and back part of the head of the tibia.

*Use.* To bend the leg, and bring it directly backwards.

N. B. The two last form what is called the *inner ham-string*.

## 3. BICEPS FLEXOR CRURIS,

*Arises*, by two distinct heads. The first,  
called



called *longus*, arises, in common with the semitendinosus, from the upper and posterior part of the tuberosity of the os ischium. The second, called *brevis*, arises from the linea aspera, a little below the termination of the gluteus maximus, by a fleshy acute beginning, which soon grows broader as it descends to join with the first head, a little above the external condyle of the os femoris.

*Inserted* by a strong tendon, into the upper part of the head of the fibula.

*Use.* To bend the leg.

*Biceps cruris*, Albinus.

*Biceps*, Winflow and Douglas.

N. B. This muscle forms what is called the *outer ham-string*; and between it and the inner, the nervous popliteus, and arteria and vena poplitea, are situated.

#### 4. POPLITEUS,

*Arises*, by a round tendon, from the lower  
and



and back-part of the external condyle of the os femoris; then runs over the ligament that involves the joint, firmly adhering to it, and part of the femilunar cartilage. As it runs over the joint, it becomes fleshy, and the fibres run obliquely inwards, being covered with a thin tendinous membrane.

*Inserted*, broad, thin, and fleshy, into a ridge at the upper and internal edge of the tibia, a little below its head.

*Use*. To assist in bending the leg, and to prevent the capsular ligament from being pinched. After the leg is bent, this muscle serves to roll it inwards.



## C H A P. XXXI.

MUSCLES *situated on the* LEG.

**T**HESE are called *Muscles* of the *Foot*;  
 and may be divided into *two classes*,  
*viz. Extensors and Flexors* of the *Foot*.  
 2. *Common Extensors and Flexors* of the *Toes*.

*First Class.**Extensors.*

These consist of three:

1. GASTROCNEMIUS EXTERNUS, SEU  
 GEMELLUS,

*Arises*, by two distinct heads. The first head arises from the upper and back-part of the internal condyle of the os femoris, and from that bone, a little above its condyle, by two distinct tendinous origins. The second



cond head arises tendinous from the upper and back-part of the external condyle of the os femoris. A little below the joint, their fleshy bellies unite in a middle tendon; and, below the middle of the tibia, it sends off a broad thin tendon, which joins a little above the extremity of the tibia with the tendon of the following.

2. SOLEUS, SEU GASTROCNEMIUS INTERNUS,

*Arises* by two origins. The first is from the upper and back-part of the head of the fibula, continuing to receive many of its fleshy fibres from the posterior part of that bone for some space below its head. The other origin begins from the posterior and upper part of the middle of the tibia; and runs inwards along the inferior edge of the popliteus towards the inner part of the tibia, from which it receives fleshy fibres for some way down. The flesh of this muscle,



covered by the tendon of the gemellus, runs down near as far as the extremity of the tibia; a little above which the tendons of both gastrocnemii unite, and form a strong round chord, which is called *tendo Achillis*.

*Inserted* into the upper and posterior part of the os calcis; by the projection of which the tendo Achillis is at a considerable distance from the tibia.

*Use.* To extend the foot, by bringing it backwards and downwards.

*Gemellus and Soleus, Albinus.*

*Gastrocnemii and Soleus, Winflow.*

*Extensor tarfi suralis, vel Extensor magnus, Douglas.*

### 3. PLANTARIS,

*Arises*, thin and fleshy, from the upper and back-part of the root of the external condyle of the os femoris, near the inferior extremity of that bone, adhering to the ligament that involves the joint in its descent.

It



It passes along the second origin of the soleus, and under the gemellus, where it sends off a long, slender, thin tendon, which comes from between the great extensors, where they join tendons; then runs down by the inside of the tendo Achillis.

*Inserted* into the inside of the posterior part of the os calcis, below the tendo Achillis.

*Use.* To assist the former, and to pull the capsular ligament of the knee from between the bones. It seems likewise to assist in rolling the foot inwards.

*Tibialis gracilis, vulgo Plantaris, Winflow.*

*Extensor tarfi minor, vulgo Plantaris, Douglas.*

N. B. This muscle, though seldom, has been found wanting on both sides.

**Flexors:**

These consist of four; two that belong to the tibia, and two to the fibula.



## 1. TIBIALIS ANTICUS,

*Arises*, tendinous and fleshy, from the middle of that process of the tibia, to which the fibula is connected above; then it runs down fleshy on the outside of the tibia; from which, and the upper-part of the interosseous ligament, it receives a number of distinct fleshy fibres; near the extremity of the tibia, it sends off a strong round tendon, which passes under part of the ligamentum tarfi annulare near the malleolus internus.

*Inserted*, tendinous, into the inside of the os cuneiforme internum, and posterior end of the metatarsal bone that sustains the great toe.

*Use*. To bend the foot by drawing it upwards, and, at the same time, to turn the toes inwards.



## 2. TIBIALIS POSTICUS,

*Arises*, by a narrow fleshy beginning, from the fore and upper part of the tibia, just under the process which joins it to the fibula; then passing through a perforation in the upper part of the interosseus ligament, it continues its origin from the back-part of the fibula next the tibia, and from near one half of the upper-part of the last-named bone; as also, from the interosseous ligament, the fibres running towards a middle tendon, which sends off a round one that passes in a groove behind the malleolus internus.

*Inserted*, tendinous, into the upper and inner part of the os naviculare, being further continued to the os cuneiforme internum and medium; besides, it gives some tendinous filaments to the os calcis, os cuboides, and to the root of the metatarsal bone that sustains the middle toe.



*Use.* To extend the foot, and to turn the toes inwards.

### 3. PERONEUS LONGUS,

*Arises*, tendinous and fleshy, from the fore-part of the head of the perone, or fibula, the fibres running straight down; also from the upper and external part of the fibula, where it begins to rise into a round edge; as also from the hollow between that and its anterior edge, as far down as to reach within a handbreadth of the ankle, by a number of fleshy fibres, which run outwards towards a tendon, that sends off a long round one, which passes through a channel at the outer ankle, in the back-part of the inferior extremity of the fibula; then, being reflected to the sinuosity of the os calcis, it runs along a groove in the os cuboides, above the muscles in the sole of the foot.

*Inserted*, tendinous, into the outside of  
the



the root of the metatarsal bone that sustains the great toe, and by some tendinous fibres in the os cuneiforme internum.

*Use.* To move the foot outwards, and to extend it a little.

*Peroneus maximus*, vulgo *Peroneus posterior*, Winflow.

*Peroneus primus*, feu *Posticus*, Douglas.

#### 4. PERONEUS BREVIS,

*Arises*, by an acute fleshy beginning, from above the middle of the external part of the fibula; from the outer side of the anterior spine of this bone; as also, from its round edge externally, the fibres running obliquely outwards towards a tendon on its external side: it sends off a round tendon which passes through the groove at the outer ankle, being there included under the same ligament with that of the preceding muscle; and a little further, it runs through a particular one of its own.



*Inserted*, tendinous, into the root and external part of the metatarsal bone that sustains the little toe.

*Use.* To assist the former in pulling the foot outwards, and extending it a little.

*Peroneus medius*, vulgo *Peroneus anticus*, Winslow.

*Peroneus secundus*, seu *Anticus*, Douglas.

### *Second Class.*

#### Common Extensors.

These consist of two.

#### 1. EXTENSOR LONGUS DIGITORUM PEDIS,

*Arises*, tendinous and fleshy, from the upper and outer part of the head of the tibia, and from the head of the fibula where it joins with the tibia, and from the interosseous ligament; also from the tendinous fascia, which covers the upper and outside of the leg by a number of fleshy fibres; and

ten-



tendinous and fleshy, from the anterior spine of the fibula, almost its whole length, where it is inseparable from the *peroneus tertius*. It splits into four round tendons, under the ligamentum tarfi annulare.

*Inserted*, by a flat tendon, into the root of the first joint of each of the four small toes; and is expanded over the upper-side of the toes, as far as the root of the last joint.

*Use*. To extend all the joints of the four small toes.

*Extensor longus*, Douglas.

*N. B.* A portion of this muscle, which

*Arises* from the middle of the fibula, continues down to near its inferior extremity, and sends its fleshy fibres forwards to a tendon, which passes under the annular ligament, and is

*Inserted* into the root of the metatarsal bone that sustains the little toe: it is called by Albinus, *Peroneus tertius*; and by others, the *Nonus Vesalii*.

*Use*. To assist in bending the foot.



2. EXTENSOR BREVIS DIGITORUM  
PEDIS,

*Arises*, fleshy and tendinous, from the fore and upper part of the os calcis; and soon forms a fleshy belly, divisible into four portions, which sends off an equal number of tendons that pass over the upper part of the foot, under the tendons of the former.

*Inserted*, by four slender tendons, into the tendinous expansion from the extensor longus, which covers the small toes, except the little one; also into the tendinous expansion from the extensor pollicis, that covers the upper part of the great toe.

*Use*. To extend the toes.

*Extensor brevis*, Douglas.

Flexors:

These may be reckoned three.

1. FLEXOR BREVIS DIGITORUM PEDIS  
PERFORATUS, SUBLIMIS,

*Arises*, by a narrow fleshy beginning,  
from



from the inferior and posterior part of a protuberance of the os calcis, between the abductors of the great and little toes; and soon forms a thick fleshy belly, which sends off four tendons that split for the passage of the flexor longus.

*Inserted* into the second phalanx of the four lesser toes. The tendon of the little toe is often wanting,

*Use.* To bend the second joint of the toes.

*Perforatus, seu Sublimis, Douglas.*

## 2. FLEXOR LONGUS DIGITORUM PEDIS, PROFUNDUS, PERFORANS,

*Arises,* by an acute tendon, which soon becomes fleshy, from the back part of the tibia, some way below its head, near the entry of the medullary artery; which beginning, is continued down the inner edge of this bone by short fleshy fibres, ending in its tendon; also, by tendinous and fleshy fibres, from the outer edge of the tibia;



and between this double order of fibres the tibialis posticus muscle lies inclosed. Having passed under two annular ligaments, it then passes through a sinuosity at the inside of the os calcis; and about the middle of the sole of the foot, divides into four tendons, which pass through the flits of the perforatus; and, just before its division, it receives a considerable tendon from that of the flexor pollicis longus.

*Inserted* into the extremity of the last joint of the four lesser toes.

*Use.* To bend the last joint of the toes.

This muscle is assisted by the

FLEXOR DIGITORUM ACCESSORIUS, SEU  
MASSA CARNEA JACOBI SYLVII,

*Arises*, by a thin fleshy origin, from most part of the sinuosity at the inside of the os calcis, which is continued forwards for some space on the same bone; also, by a thin tendinous beginning, from before the tuberosity



rofity of the os calcis externally; and, soon becoming all fleshy, is

*Inserted* into the tendon of the flexor longus, just at its division into four tendons.

*Use.* To assist the flexor longus.

### 3. LUMBRICALES PEDIS,

*Arise*, by four tendinous and fleshy beginnings, from the tendon of the flexor profundus, just before its division, near the insertion of the massa carnea.

*Inserted*, by four slender tendons, into the inside of the first joint of the four lesser toes, and are lost in the tendinous expansion that is sent from the extensors to cover the upper part of the toes.

*Use.* To increase the flexion of the toes, and to draw them inwards.



## C H A P. XXXII.

MUSCLES *which are chiefly situated on*  
*the FOOT.*

THESE may be divided into the *muscles*  
of the *great toe*, of the *little toe*, and  
of the *metatarsal bones*.

Muscles of the great toe.

These are five :

1. EXTENSOR PROPRIUS POLLICIS  
PEDIS,

*Arises*, by an acute tendinous, and fleshy  
beginning, some way below the head and  
anterior part of the fibula, along which it  
runs to near its lower extremity, connected  
to it by a number of fleshy fibres, which  
descend obliquely towards a tendon.

*In-*



*Inserted*, tendinous, into the posterior part of the first and last joint of the great toe.

*Use*. To extend the great toe.

*Extensor longus*, Douglas.

## 2. FLEXOR LONGUS POLLICIS PEDIS,

*Arises*, by an acute, tendinous, and fleshy beginning, from the posterior part of the fibula, some way below its head, being continued down the same bone, almost to its inferior extremity, by a double order of oblique fleshy fibres; its tendon passes under an annular ligament at the inner ankle.

*Inserted* into the last joint of the great toe, and generally sends a small tendon to the os calcis.

*Use*. To bend the last joint of this toe.

*Flexor longus*, Douglas.



## 3. FLEXOR BREVIS POLLICIS PEDIS,

*Arises*, tendinous, from the under and fore-part of the os calcis, where it joins with the os cuboides, from the os cuneiforme externum, and is inseparably united with the abductor and adductor pollicis.

*Inserted* into the external os sesamoideum and root of the first joint of the great toe.

*Use*. To bend the first joint.

## 4. ABDUCTOR POLLICES PEDIS,

*Arises*, fleshy, from the inside of the root of the protuberance of the os calcis, where it forms the heel; and tendinous from the same bone, where it joins with the os naviculare.

*Inserted*, tendinous, into the internal os sesamoideum, and root of the first joint of the great toe.

*Use*.



*Use.* To pull the great toe from the rest.

*Thenar*, Winflow.

5. ADDUCTOR POLLICIS PEDIS,

*Arises*, by a long thin tendon, from the os calcis, from the os cuboides, from the os cuneiforme externum, and from the root of the metatarsal bone of the second toe.

*Inserted* into the external os sesamoideum, and root of the metatarsal bone of the great toe.

*Use.* To bring this toe nearer the rest.

*Antithenar*, Winflow.

Muscles of the little toe.

These, besides the common extensors and flexors, are two, viz.

1. ABDUCTOR MINIMI DIGITI PEDIS,

*Arises*, tendinous and fleshy, from the semicircular edge of a cavity on the inferior part of the protuberance of the os calcis,  
and



and from the root of the metatarsal bone of the little toe.

*Inserted*, into the root of the first joint of the little toe externally.

*Use*. To draw the little toe outwards from the rest.

*Parathenar major*, and *Metatarseus*, Winflow.

## 2. FLEXOR BREVIS MINIMI DIGITI PEDIS,

*Arises*, tendinous, from the os cuboides, near the fulcus or furrow for lodging the tendon of the peroneus longus; fleshy from the outside of the metatarsal bone that sustains this toe, below its protuberant part.

*Inserted* into the anterior extremity of the metatarsal bone, and root of the first joint of this toe.

*Use*. To bend this toe.

*Parathenar minor*, Winflow.

Muscles



Muscles from the metatarsal bones.

These are four external and three internal interossei, and one muscle which is common to all the metatarsal bones.

*Interossei Pedis externi, Bicipites.*

1. ABDUCTOR INDICIS PEDIS,

*Arises*, tendinous and fleshy, by two origins, from the root of the inside of the metatarsal bone of the fore-toe, from the outside of the root of the metatarsal bone of the great toe, and from the os cuneiforme internum.

*Inserted*, tendinous, into the inside of the root of the first joint of the fore-toe.

*Use*. To pull the fore-toe inwards from the rest of the small toes.

2. ADDUCTOR INDICIS PEDIS,

*Arises*, tendinous and fleshy, from the roots



roots of the metatarsal bones of the fore and second toe.

*Inserted*, tendinous, into the outside of the root of the first joint of the fore-toe.

*Use*. To pull the fore-toe outwards towards the rest.

3. ADDUCTOR MEDII DIGITI PEDIS,

*Arises*, tendinous and fleshy, from the roots of the metatarsal bones of the second and third toes.

*Inserted*, tendinous, into the outside of the root of the first joint of the second toe.

*Use*. To pull the second toe outwards.

4. ADDUCTOR TERTII DIGITI PEDIS,

*Arises*, tendinous and fleshy, from the roots of the metatarsal bones of the third and little toe.

*Inserted*, tendinous, into the outside of the root of the first joint of the third toe.

*Use*.



*Use.* To pull the third toe outwards.

*Interossei Pedis interni.*

1. ABDUCTOR MEDII DIGITI PEDIS,

*Arises*, tendinous and fleshy, from the inside of the root of the metatarsal bone of the middle toe internally.

*Inserted*, tendinous, into the inside of the root of the first joint of the middle toe.

*Use.* To pull the middle toe inwards.

2. ABDUCTOR TERTII DIGITI PEDIS,

*Arises*, tendinous and fleshy, from the inside and inferior part of the root of the metatarsal bone of the third toe.

*Inserted*, tendinous, into the inside of the root of the first joint of the third toe.

*Use.* To pull the third toe inwards.

3. ADDUCTOR MINIMI DIGITI PEDIS,

*Arises*, tendinous and fleshy, from the inside



side of the root of the metatarsal bone of the little toe.

*Inserted*, tendinous, into the inside of the root of the first joint of the little toe.

*Use*. To pull the little toe inwards.

The muscle which brings the extremities of the metatarsal bones towards each other, is named

#### TRANSVERSALIS PEDIS,

*Arises*, tendinous, from the under part of the anterior extremity of the metatarsal bone of the great toe, and from the internal os sesamoideum of the first joint, adhering to the adductor pollicis.

*Inserted*, tendinous, into the under and outer part of the anterior extremity of the metatarsal bone of the little toe, and ligament of the next toe.


*Use*. To contract the foot, by bringing the great toe and the two outermost toes nearer each other.

N. B.



*N. B.* The muscles situated on the sole of the foot are covered by a strong tenacious aponeurosis, which is extended from the os calcis to the first joints of all the toes, and serves to preserve the subjacent parts from being compressed in standing and walking.







# I N D E X.

A	Page
<b>A</b> BDOMEN, muscles situated on the anterior part of it	77
— within its cavity	101
Abductor indicis manus	172
— indicis pedis	219
— minimi digiti manus	172
— minimi digiti pedis	217
— oculi	92
— pollicis manus	170
— pollicis pedis	216
— medii digiti pedis	221
— tertii digiti pedis	ib.
Accelerator urinæ	89
Adductor brevis femoris	182
— longus femoris	181
— medii digiti pedis	220
— magnus femoris	182
— metacarpi minimi digiti manus	173
— minimi digiti pedis	221
— oculi	29
— pollicis manus	171
— pollicis pedis	217
— tertii digiti pedis	220
Amygdalæ	62
Anconeus	153
Angularis	131
Ani spincter	92. 97
— levator	93
Anterior auris	71



Antithenar	-	-	-	217
Antitragicus	-	-	-	19
Anus, muscles of the	-	-	-	91
Arytenoideus minor	-	-	-	73
———— obliquus	-	-	-	72
———— transversus	-	-	-	73
Arytæno-epiglottideus	-	-	-	74
Attollens aurem	-	-	-	16
———— oculi	-	-	-	28
Aurem attollens	-	-	-	16
Auris anterior	-	-	-	17
———— retrahentes	-	-	-	ib.
———— posterior	-	-	-	18
———— internus	-	-	-	22
———— transversus	-	-	-	20
———— obliquus	-	-	-	21
Azygos uvulæ	-	-	-	67

## B

Basio-cerato-chondro-glossus	-	-	54
Biceps brachii	-	-	151
———— flexor cubiti	-	-	149
———— flexor cruris	-	-	197, 198
———— internus	-	-	151
———— externus	-	-	153
Biventer maxillæ inferioris	-	-	52
Brachialis internus	-	-	151
Buccinator	-	-	39
Bulbo-cavernosus	-	-	90

## C

Capitis obliquus superior	-	-	136
———— obliquus inferior	-	-	137



Capitis par tertium	-	-	-	156
—— rectus	-	-	-	135
—— posticus	-	-	-	ib.
Cephalo-pharyngeus	-	-	-	71
Cerato-glossus	-	-	-	54
Cervicalis descendens	-	-	-	128
Circumflexus, or Tensor palati	-	-	-	60
Clitoridis erectior	-	-	-	95
Coccygeus	-	-	-	100
Colli latissimus	-	-	-	49
—— longus	-	-	-	117
—— transversalis	-	-	-	134
—— semispinalis	-	-	-	ib.
—— interspinales	-	-	-	139
—— intertransversales	-	-	-	140
Complexus	-	-	-	129
—— minor	-	-	-	130
Compressor naris	-	-	-	32
Constrictor oris	-	-	-	42
—— isthmi faucium	-	-	-	64
—— pharyngis	-	-	-	68, 69, 70
Coraco brachialis	-	-	-	147
—— hyoideus	-	-	-	56
—— radialis	-	-	-	151
Corrugator supercilii	-	-	-	15
Cranium, muscles of its teguments	-	-	-	13
Cremaster	-	-	-	88
Crico-arytenoideus lateralis	-	-	-	72
Crico-arytenoideus posticus	-	-	-	ib.
Crico-thyroideus	-	-	-	57
Crico pharyngeus	-	-	-	69
Crotaphyte muscle	-	-	-	44



Cruralis	-	-	195
Cubit, or fore-arm, muscles of	-	-	149, 154
Cueularis	-	-	120
Cutaneus	-	-	48
D			
Dartos	-	-	87
Deltoides	-	-	146
Depressor anguli oris	-	-	37
———— labii inferioris	-	-	38
———— labii superioris alæque nasi	-	-	36
———— labiorum communis	-	-	37
———— oculi	-	-	28
Diaphragma	-	-	101
Digastricus	-	-	51
Dorsi interspinales	-	-	141
———— latissimus	-	-	122
———— spinalis	-	-	126
———— longissimus	-	-	127
———— semispinalis	-	-	132, 133
E			
Ear, muscles of the	-	-	16
Elevator oculi	-	-	28
———— labii superioris	-	-	36
———— labii inferioris	-	-	39
Epicranius	-	-	15
Epiptaphilinus	-	-	67
Erector clitoridis	-	-	95
———— penis	-	-	88
Extensor brevis digitorum pedis	-	-	210
———— carpi radialis brevior	-	-	158
———— carpi radialis longior	-	-	ib.
———— carpi ulnaris	-	-	159



Extensor digitorum communis	- -	162
—— longus digitorum pedis	- -	208
—— ossis metacarpi pollicis manus	- -	168
—— primi internodii pollicis manus	- -	169
—— proprius pollicis pedis	- -	214
—— secundi internodii pollicis manus	- -	169
—— major pollicis manus	- -	170
—— pollicis secundus	- -	ib.
—— tertii internodii	- -	ib. 173, 175
—— tarfi suralis	- -	202
—— tarfi minor	- -	203
Extremities, superior, muscles of	- -	142
—— inferior, muscles of	- -	180
Eye-ball, muscles of	- -	27
Eye-lids, muscles of	- -	24

## F

Fallopian capitis par tertium	- -	130
Fauces, muscles about the	- -	64
Faucium isthmi constrictor	- -	ib.
Flexor accessorius digitorum pedis	- -	212
—— brevis digitorum pedis	- -	210
—— brevis pollicis manus	- -	167
—— brevis pollicis pedis	- -	216
—— longus digitorum pedis	- -	211
—— primi internodii	- -	168, 173
—— tertii internodii	- -	167
—— secundi internodii	- -	ib.
—— carpi radialis	- -	156
—— carpi ulnaris	- -	157
—— longus digitorum pedis	- -	211
—— longus pollicis manus	- -	166

## L



Flexor longus pollicis pedis	-	215
— ossis metacarpi pollicis	-	167
— parvus minimi digiti manus	-	174
— profundus perforans	-	161
— sublimis perforatus	-	160

Foot, muscles on the	-	214
----------------------	---	-----

Frontalis	-	12
-----------	---	----

## G

Gastrocnemius externus	-	200
------------------------	---	-----

— internus	-	201
------------	---	-----

Gemellus	-	200
----------	---	-----

Gemini	-	188
--------	---	-----

Genae quadratus	-	49
-----------------	---	----

Generation, organs of; muscles about them	87, 95
---	--------

Genio-hyo-glossus	-	53
-------------------	---	----

Genio-hyoideus	-	52
----------------	---	----

Glossopharyngeus	-	65
------------------	---	----

— pharyngeus	-	71
--------------	---	----

Glottis, muscles situated about it	-	72
------------------------------------	---	----

Gluteus maximus	-	184
-----------------	---	-----

— medius	-	186
----------	---	-----

— minimus	-	187
-----------	---	-----

Gracilis	-	192
----------	---	-----

## H

Hand, muscles of the	-	166
----------------------	---	-----

Helicis major	-	18
---------------	---	----

— minor	-	ib
---------	---	----

Humeri os, muscles situated on	-	149
--------------------------------	---	-----

Hyoides os, muscles about it	-	51, 55, 58
------------------------------	---	------------

Hyo-glossus	-	54
-------------	---	----

Hyo-pharyngeus	-	70
----------------	---	----



Hyo-thyroideus	-	-	-	57
Hypothenar minor	-	-	-	173
———— Riolani	-	-	-	174
I				
Jaw, lower, muscles of	-	-	-	43
—— muscles about it and the os hyoides	-	-	-	51, 55, 58
Iliacus internus	-	-	-	107
Incisivus inferior	-	-	-	39
—— lateralis	-	-	-	36
—— medius	-	-	-	ib.
Indicator	-	-	-	171
Infra Spinatus	-	-	-	143
—— costales	-	-	-	115
Intercostales externi	-	-	-	113
—— interni	-	-	-	114
Interosseus auricularis	-	-	-	176
—— pedis externi	-	-	-	219
—— pedis interni	-	-	-	221
Interspinales colli	-	-	-	139
—— dorfi et lumborum	-	-	-	141
Intertransversales colli	-	-	-	140
—— dorfi	-	-	-	141
—— lumborum	-	-	-	ib.
Ischio-cavernosus	-	-	-	89
Isthmi faucium constrictor	-	-	-	64
L				
Labii inferioris depressor	-	-	-	38
—— Inferioris levator	-	-	-	ib.
—— superioris, alaeque nasi levator	-	-	-	35
—— elevator	-	-	-	36
—— depressor	-	-	-	37



Labiorum communis depressor	37
——— sphincter	42
Latissimus colli	49
——— dorsi	122
Laxator tympani	20
Leg, muscles situated on	200
Levatores costarum	115
Levator anguli oris	34
——— ani	93
——— ani feminæ	97
——— labii inferioris	39
——— labii superioris alæque nasi	35
——— menti	39
——— oculi	28
——— palati	61
——— palpebræ superioris	26
——— parvus	93
——— proprius	131
——— scapulæ	ib
Lingualis	54
Lips, muscles of	34
Longissimus dorsi	127
Longus colli	117
Lumbaris externus	106
——— internus	107
Lumborum intertransversales	141
——— quadratus	105
Lumbricales manus	162
——— pedis	214
M	
Mallei anterior	21



Mallei externus	-	-	-	22
—— internus	-	-	-	21
Massa carnea Jacobi Sylvii	-	-	-	212
Masseter	-	-	-	45
Menti levator	-	-	-	39
Marsupialis	-	-	-	100
Mastoideus	-	-	-	50, 130
Metacarpeus	-	-	-	173
Metatarseus	-	-	-	218
Mouth, muscles of	-	-	-	34
Multifidus spinæ	-	-	-	132
Musculus cutaneus	-	-	-	48
Mylo-hyoideus	-	-	-	52
—— pharyngeus	-	-	-	71

## N

Nasalis	-	-	-	33
—— labii superioris	-	-	-	42
Naris compressor	-	-	-	32
Nasi alæ levator	-	-	-	35
—— depressor	-	-	-	37
Neck, muscles about its anterior part			48, 117	
Nonus Vesalii	-	-	-	209
Nose, muscles of	-	-	-	32

## O

Obliquus ascendens internus	-	-	81
—— capitis inferior	-	-	137
—— capitis superior	-	-	136
—— descendens	-	-	80
—— descendens externus	-	-	77
—— externus abdominis	-	-	86
—— internus abdominis	-	-	83



Obliquus inferior oculi	- - -	31
Obliquus superior, seu trochlearis	- - -	29
Obturator internus	- - -	100
Oecipito-frontalis	- - -	13
Oculi adductor	- - -	29
—— abductor	- - -	ib.
—— attollens	- - -	28
—— depresso	- - -	ib.
—— deprimens	- - -	ib.
—— elevator	- - -	ib.
—— levator	- - -	ib.
—— obliquus inferior	- - -	31
—— obliquus major	- - -	30
—— obliquus minor	- - -	31
Omo-hyoideus	- - -	55
Opponens pollicis	- - -	167
Orbicularis oris	- - -	41
Orbicularis palpebrarum	- - -	24
Oris anguli depresso	- - -	37
—— anguli retractor	- - -	40
—— constrictor	- - -	42
—— levator anguli	- - -	34
—— orbicularis	- - -	41
P		
Palati circumflexus	- - -	61
—— levator	- - -	ib.
—— tensor	- - -	60
Palato-pharyngeus	- - -	65
—— salpingeus	- - -	61
—— staphilinus	- - -	6
Palmaris brevis	- - -	156
—— cutaneus	- - -	ib.



Palmaris longus	-	155
Palpebræ superioris levator	-	26
Palpebram aperiens rectus	-	ib.
Palpebrarum orbicularis	-	24
Patientiæ musculus	-	131
Parathenar major	-	218
———— minor	-	ib.
Pectinalis	-	180
Pectineus	-	181
Pectoralis major	-	109
———— minor	-	111
Pedis interossei externi	-	219
———— interossei interni	-	221
Pelvis, muscles about it	-	99
Penis erector	-	81
Perforans	-	161, 211
Perforatus	-	161, 211
Perinei transversus	-	90, 91
Peroneus brevis	-	207
———— longus	-	206
———— maximus	-	207
———— medius	-	208
———— primus	-	207
———— secundus	-	208
———— tertius	-	209
Pharynx, muscles on the posterior part of it	-	68
———— constrictor	-	68, 69, 70
Plantaris	-	202
Platysma myoides	-	48
Popliteus	-	198
Posterior annularis	-	178



Posterior auris	-	-	18
—— indicis	-	-	175
—— medii	-	-	178
Prior annularis	-	-	176
—— indicis	-	-	174
—— medii	-	-	177
Pronator radii quadratus	-	-	165
—— radii teres	-	-	164
Prostrate inferior	-	-	90
Psoas magnus	-	-	106
—— parvus	-	-	ib.
Pterygoideus externus	-	-	46
—— internus	-	-	45
—— major	-	-	46
—— minor	-	-	47
Pterygo-pharyngeus	-	-	71
Pyramidalis	-	-	36, 86
Pyriformis	-	-	188

## Q

Quadratus	-	-	38
—— femoris	-	-	189
—— genæ	-	-	49
—— lumborum	-	-	105

## R

Radialis externus longior	-	-	158
Radialis externus brevior	-	-	159
—— primus	-	-	158
—— internus	-	-	157
—— secundus	-	-	157
Rectus	-	-	193
—— abdominis	-	-	84



Rectus capitis internus major	-	-	118
capitis internus minor	-	-	ib.
capitis lateralis	-	-	119
capitis posticus major	-	-	195
capitis posticus minor	-	-	136
Retractor anguli oris	-	-	40
Retrahentes auris	-	-	17
Rinæus	-	-	33
Rhomboideus	-	-	123
S			
Sacer	-	-	133
Sacro-lumbalis	-	-	128
Salpingo-pharyngeus	-	-	66
staphilinus	-	-	62
Sartorius	-	-	190
Scalenus anticus	-	-	137
medius	-	-	138
posticus	-	-	ib.
prior	-	-	ib.
Scapulæ levator	-	-	131
Semi-interosseus	-	-	172
Semi-nervosus	-	-	197
Semi-orbicularis	-	-	42
Semimembranosus	-	-	197
Semispinalis colli	-	-	134
dorsi	-	-	132
Semitendinosus	-	-	196
Seminis ejaculator	-	-	89
Serratus anticus	-	-	111
magnus	-	-	ib.
posticus inferior	-	-	123



Serratus posticus superior	126
Soleus	201, 202
Spheno-falpingo-staphilinus externus	61
Sphincter ani	92, 97
—— cutaneus	93
—— externus	ib.
—— labiorum	42
—— vaginae	96
Spinæ multifidus	132
Splenius	126
—— capitis	125
—— colli	ib.
Stapedius	22
Staphilinus	64, 67
Sterno cleido-mastoideus	49
Sterno-costalis	116
Sterno-hyoideus	55
Sterno-thyroideus	56
Stylo-glossus	58
Stylo-hyoideus	59
Stylo-pharyngeus	ib.
Subclavius	110
Sublimis	161
Subscapularis	148
Supercilii corrugator	15
Supinator radii brevis	164
—— radii longus	163
Supra costales	115
—— spinatus	142
Sylvii Jacobi massa carnea	212
Syndesmo-pharyngeus	70



## T

Temporalis	-	-	-	43
Tensor tympani	-	-	-	21
—— palati	-	-	-	61
—— vaginæ femoris	-	-	-	191
Teres major	-	-	-	145
—— minor	-	-	-	144
Thenar Riolani	-	-	-	170
—— Winfl.	-	-	-	217
Thorax, muscles situated about its anterior part				119
—— between it and the ribs	-	-	-	113
Thigh, muscles situated on	-	-	-	190
Thumb, muscles of	-	-	-	166
Thyreo arytenoideus	-	-	-	73
Thyreo-epiglottideus	-	-	-	75
Thyro-pharyngeus	-	-	-	69
Thyro-hyoideus	-	-	-	57
Thyro-pharyngo-staphilinus	-	-	-	66
Thyro-staphilinus	-	-	-	66
Tibialis anticus	-	-	-	204
—— gracilis	-	-	-	203
—— posticus	-	-	-	205
Trachelo-mastoideus	-	-	-	130
Tragicus	-	-	-	19
Transversalis	-	-	-	83
—— anticus	-	-	-	119
—— colli	-	-	-	134
—— lumborum	-	-	-	133
—— pedis	-	-	-	222
—— urethræ	-	-	-	90
Transverso-spinalis	-	-	-	132



Transversus dorsi	132
—— abdominis	84
—— auris	20
—— perinei	90, 91
Trapezius, seu Cucullaris	120
Triangularis	37
—— or Sterno-costalis	116
Triceps adductor femoris	181
—— extensor cubiti	152
—— primus	182
—— secundus	ib.
—— tertius	183
Trochlearis	29
Trunk, muscles situated on its posterior part	120
Tympani laxator	20
—— tensor	21
V	
Vaginæ sphincter	96
Vastus externus	194
—— internus	195
Ulnaris externus	157
—— gracilis	155
—— internus	157
Urethræ transversalis	90
Urinæ accelerator	89
Uvulæ azygos	67
Z	
Zygomaticus major	40
—— minor	ib.



