The modern practice of the London hospitals. Viz. St. Bartholomew's, St. Thomas's, Guy's, St. George's, the Portuguese, and the Lock, at Hyde-Park-Corner. - Copy 2

Contributors

St. Thomas' Hospital (London, England) St. George's Hospital (London, England) Guy's Hospital. St. Bartholomew's Hospital (London, England) Lock Hospital (London, England) Portuguese Hospital (London, England)

Publication/Creation

London : Printed for S. Crowder, and J. Coote ... : W. Griffin ... and : W. Nichols ..., MDCCLXX. [1770]

Persistent URL

https://wellcomecollection.org/works/ep5476xk


License and attribution


This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.




Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org




















THE MODERN PRACTICE

OFTHE

LONDON HOSPITALS.

VIZ.

St. Bartholomew's, | The Portuguese, St. Thomas's, Guy's, St. George's,

and The Lock, at Hyde-Park-Corner.

THE THIRD EDITION.


WITHAN

Useful INDEX of Diseases and their Remedies.

LONDON:

Printed for S. CROWDER, and J. COOTE, in Pater-nofter Row; W. GRIFFIN, in Catharine-Street in the Strand; and W. NICOLL, in St. Paul's Church Yard.

M DCC LXX.



OF DIET. TABLE

Freeking, A P c:

FULL

Sunday and Thursday.

Reakfast. A Pint of Water Gruel. Dinner. Half a Pound of boiled Beef with Greens.

Supper. A Pint of Broth.

Tuesday and Saturday.

Breakfast. A Pint of Water Gruel. Dinner. Half a Pound of boiled Mutton with Greens.

Supper. A Pint of Broth.

Monday.

Breakfaft. A Pint of Milk Pottage. Dinner. A Pint of Rice Milk. Supper. Two Ounces of Cheefe, or Butter.

Wednesay.

Breakfast. A Pint of Milk Pottage. Dinner. Half a Pound of boiled Pudding. Supper. A Pint of Water-Gruel. A 2 Friday

Friday.

Breakfaft. A Pint of Milk Pottage. Dinner. A Pint of Plumb Broth. Supper. Two Ounces of Cheefe, or Butter

The Patients upon Full Diet, shall have one Loaf of Bread per Day.

Three Pints of Small Beer per Day, from Ladyday to Mickaelmas.

One quart per Day, from Michaelmas to Lady-

Day. N. B. The Loaf of Bread weighs fourteen Ounces.

ballet. Little Later ball . 1906

Tralling wind rather

Enter & Plate of Witness Grund

Supper. A Pleased Britis ?!

Supper, A Pire of Breek



Sec. 2. 19 198 48 303 23854

+12150 L CO

toin actual of herical de benefit a that are

LOW

LOW DIET.

Sunday.

B Reakfaft. A Pint of Water-Gruel. Dinner. Two Ounces of Roafted Veal with a Slice of Bread Pudding. Supper, A Pint of Broth.

Tuesday and Saturday.

Breakfast. A Pint of Water-Gruel. Dinner. Two Ounces of boiled Mutton with Greens, and a Pint of Broth. Supper. A Pint of Broth.

Monday.

Breakfast. A Pint of Milk Pottage. Dinner. A Pint of Rice Milk. Supper. Two Ounces of Cheefe or Butter.

Wednesday.

Breakfast. A Pint of Milk Pottage. Dinner. A Slice of boiled Pudding. Supper. A Pint of Water Gruel.

Thursday.

Breakfaft. A Pint of Water Gruel. Dinner. Two Ounces of roafted Veal, and a Pint of Rice Milk. Supper. A Pint of Broth. A 3 Friday.

Friday.

Breakfaft. A Pint of Milk Pottage. Dinner. A Pint of Plumb Broth. Supper. Two Ounces of Cheefe or Butter.

The Patients upon Low Diet, shall have one Loaf of Bread per Day.

One Quart of Beer per Day, from Lady-day to Michaelmas.

One Pint per Day, from Michaelmas to Lady-Day.

Treplay and Samerica

Light The ATAT TO PATE A

A Pau of Broth.

Breaking, A Philler Milling & Bridger

Tot of Water Grad.

The Dintes of Cheers on Dates.

Two Chartes of holded Mutton Wills

MILK

(vii)

MILK DIET.

Sunday, Tuesday, Thursday and Saturday.

B Reakfast. A Pint of Milk Pottage or Water-Gruel.

Dinner. A Pint of Plumb Pottage, and four Ounces of Bread Pudding.

Supper. A Pint of Milk Pottage or Water-Gruel.

Monday, Wednesday, and Friday.

Breakfast. A Pint of Milk Pottage or Water-Gruel.

Dinner. A Pint of Rice Milk.

Supper. A Pint of Milk Pottage or Water-Gruel.

The Patients upon Milk Diet, shall have one Loaf of Bread per Day.

Three Pints of Drink per Day; one Pint whereof shall be Milk, and two Water.

The Patients upon Fish Diet, shall have Fish for Dinner on Mondays, Wednesdays, and Fridays, if it can conveniently be had; if not the Low Diet.

The Patients upon Dry Diet, fhall have two Ounces of Butter or Cheefe for Breakfast, and the fame for Supper every Day in the Week; and the Low Diet for Dinner, but without Broth or Rice.

Milk on Tuesdays, Thursdays, and Saturdays. Bread and Beer, as those upon Low Diet. The Patients upon Raifin Diet, fhall have Half a Pound of Raifins per Day, as much Bread as they can eat, a Quart of Decoct. Guaic. Fort. and as much of the Decoct. Guaic. Tenue, as they can Drink.

The Patients under Salivation, shall have one Quart of Milk per Day, and Half a Pound of Mutton to be boiled for Broth.


with A Fire of Mille Pottage or Water

A P. he of Mult Pottage of Water-

St. A. Diab of Ride Mills

This Post of the Star Star

inc of Mills Pottage on Wert-




more of Buger or Cheere lee Meabrieff, Med the

The ball of Ther Thereffinger, and Sander The

fiere far Supper every Day is the Wiede a and the Law Diet for Dimner, Sur without Prother Kice.

ind theer in those upon Low Lifes.



Powder of in Forty { Contains one Grain of Opium. Compound Amber, grains,

(x)

Scordium Species with Opium, in Forty-five grains, Opium.

The Simple Colocynth Pill, in Half a Drachm Troy, Contains almost Ten Grains of Scammony, and Ten Grains of Colocynth.

Aloes, in Half a Drachm Troy,

Coargound

Contain almost Eight The Colocynth Pill, with Grains of Aloes, Eight Grains of Scammony, Four Grains of Colocynth.

The Mercurial Pill, in } { Contains Fifteen Grains Twenty-eight Grains, } { of Quick filver.

The Saponaceous Pill, ¿ S Contains almost a Grain 52 of Opium. in Ten Grains,

The Storax Pill, in Five] [Contains One Grain of Opium. Grains and 4ths,

The Scammony Elec- { Contains Fifteen Grains tuary, in a Drachm { Scammony, of Scammony. and an half Troy,

The

The Scordium Electuary, in Three Contains a Grain of O-Drachms Troy,

The Confection called Paulina, in Thirtytwo Grains, Scottains a Grain of Opium.

Mithridate, in Half an Contains a Grain of O-

Venice Treacle, in Seven } { Contains a Grain of Oty-five Grains, } { pium.

London Philonium, in } { Contains a Grain of O-Thirty-fix Grains, } { pium.


In the Ammoniacum Plaister with Mercury, the Quickfilver, is almost a Fifth Part of the whole.

In the common Mercurial Plaister, the Quickfilver, is almost a Fifth Part of the whole.

In the Strong blue Ointment, the Quickfilver, is almost a Third Part of the whole.

In the weaker blue Ointment, the Quickfilver, is almost a Fifth Part of the whole.

In the Mercurial Cerate, the Quickfilver is almost a Fifth Part of the whole.



THE

FOITOANT

PRACTICE

OFTHE

LONDON HOSPITALS.

St. BARTHOLOMEW'S HOSPITAL.

WineGAR of fquills, Æthiops mineral, fimple alexiterial water, fpirituous alexiterial water, lime water, compound lime water, fimple pepper-mint water, fimple mintwater, fimple pennyroyal water, fpirituous pennyroyal water, nutmeg water, vitriolic camphorated water: are all prepared according to the Difpenfatory of the College of Phyficians of London, as directed in the Supplement to this work.

Aluminous bolus.

Take a fcruple of conferve of roles, half a cruple of roch-allum, fimple fyrup, as much as is ufficient to make a bolus, which is to be taken wice in a day.—This is ufually given in cafes which require ftrong aftringents.

Guaiacum

2.

Guaiacum bolus.

Take half a drachm of gum guaiacum powdered, a fcruple of conferve of rofes, and a fufficient quantity of fimple fyrup to make a bolus, which is to be taken every morning.—This is reckoned to promote infenfible perfpiration greatly, and on that account is good in fuch cutaneous diforders as proceed from obftructions of the perfpirable matter: in gouts and rheumatic complaints it is alfo very ferviceable.

Stomachic bolus.

Take powder of ginger, and powder of long pepper, of each fifteen grains; conferve of orangepeel, a fcruple; as much fimple fyrup as is fufficient to make a bolus.—This bolus is given three times in a day, in all complaints arifing from indigeftion, for which it is an excellent remedy.

Rhubarb bolus.

Take of toafted rhubarb, fifteen grains; and half a drachm of the fcordium electuary; mix them together: to be taken as often as occasion requires.—This is an efficacious remedy in loofenefles; the patient at the fame time abstaining from malt liquors, and making use of hartshorn drink, or the chalk decosion, for his common liquor.

Strengthning bolus.

Take of yellow colophony, one fcruple; of rhubarb, five grains; conferve of rofes, half a fcruple; fimple fyrup, a fufficient quantity to make a bolus. —This is given twice in a day, in weakneffes and gleets of eitner fex.

Saponaceous bolus.

Take of the best white soap, half a drachm; oil of juniper, five drops; fimple syrup, as much

States and

as

St. Bartholomew's Hospital.

as is required to form a bolus, which is to be taken twice a day.—This is very efficacious, not only in nephritic diforders, but alfo in the jaundice and afthma; in the most obstinate cases of which, this bolus is directed with the greatest certainty of fucces.

The faponaceous bolus with rhubarb, is made by adding ten grains of rhubarb to the foregoing bolus, and is directed for the fame complaints when attended with coffiveness.

Specific bolus.

Take of calcined mercury, one grain and an half; of London philonium, a fcruple; mix them together: to be taken every night going to reft.— This is an excellent mercurial alterative, and very much promotes the cure of ftubborn ulcers, as well as venereal complaints.

Turpentine bolus.

Take of fpirits of turpentine, twenty-five drops; conferve of wormwood, one fcruple; mix them into a bolus.—To be given in nephritic and fciatic diforders every night and morning; but fhould be cautioufly administered.

Discutient pultice.

Take of oatmeal and stale beer grounds, of each fufficient to make a pultice; —which is to be applied in mortifications and bruises, every night and morning; for which purposes it is far preferable to any other application.

Saponaceous pultice.

Take of foap lees, three ounces; fweet oil, half a pint; of warm water, a gallon; of wheat flour, a fufficient quantity to make the whole of a proper confiftence.—This is alfo an excellent difcutient.

3

4

Cordial confection, conferves, of wormwood, of orange-peels, of hips, and of red rofes, are made exactly as directed in the Supplement.

White decoction

Is prepared by boiling two ounces of burnt hartfhorn, and a quarter of an ounce of gum arabic, in three pints of water, till there remains only a quart.—This is prefcribed for common drink in diarrhæas, and all fluxes of the belly.

Decostion of marshmallows

Is made by boiling two ounces of marfhmallow roots in three pints of water, till one pint has boiled away; then add two ounces of fyrup of marfhmallows, and drink half a pint three times in a day.—It is ufually directed to be taken at the fame time with other medicines, in gravelly complaints, dyfuries, &c.

Decoction of the Peruvian bark.

Take four ounces of the beft Peruvian bark; boil in eight pints of water till two quarts only remain. The dole, two ounces, three times in a day, in intermitting fevers, and other diforders proceeding from lax fibres.

Decostion of oak bark.

Take oak bark and pomegranate shells, of each four ounces; water and milk, of each a gallon; let one half boil away; then strain and sweeten the remainder.—To be taken, two ounces three times in a day, in every case where astringents are proper.

Pectoral decoction

Is prepared as directed in the Supplement, and is a good auxiliary remedy in diforders of the breaft. See page 168.

Decoction

Sto. Barthoiomew's Hospital.

Decoction of Sarsaparilla.

Take twelve ounces of farfaparilla, and fix quarts of water, boil the water away to two quarts.—Half a pint of this decoction, drank every fix hours, has been found to relieve those pains which frequently remain after a mercurial course, and makes an excellent diet-drink during the use of mercurials.

Decoction of elm bark.

Take of the fresh inner bark of the elm, four ounces; and boil it in a sufficient quantity of water, till there remains no more than a quart.— This is directed to be taken as a vulnerary, the quantity of half a pint three times in a day: it is also sometimes used as a gargle for fore mouths, occasioned by heat and sharp humours.

Opening electuary. .

Take of lenitive electuary, two pounds; cream of tartar, four ounces; powder of jalap, two ounces; oil of carraway-feeds, a quarter of an ounce; fyrup of rofes, a fufficient quantity to make an electuary. The dole is the quantity of a nutmeg, and to be repeated as often as occasion requires.

Aromatic electuary.

Take fix ounces of the aromatic fpecies; conferve of orange-peel, one pound; and as much fimple fyrup as is fufficient to make an electuary.

Aromatic electuary, with Reel.

Take of the foregoing electuary, half a pound; of prepared fteel, half an ounce; mix them together: the dole of both these electuaries, the quantity of a nutmeg three times in a day.—They are very warm, ftrengthen the ftomach, and are of very great service in flatulencies arising from B 3 indigestion, indigeftion, as well as in complaints which arife from a languid circulation, and impoverished state of the blood.

Aftringent electuary.

Take of toasted nutmeg and diascordium, each four ounces; of prepared chalk, one ounce; fimple syrup, a sufficient quantity to form an electuary.— A drachm of this electuary is given three times a day in diarrhæas, after proper evacuations.

Cinnabarine electuary.

Take two pounds of vermillion, one pound of gum guaicum, half a pound of Venice foap, and as much fimple fyrup as is fufficient to make an electuary.— The dofe is the quantity of a nutmeg twice a day, in obstinate rheumatisms.

Electuary for the piles.

Take lenitive electuary, and flower of brimflone, of each equal parts; mix them into an electuary with fimple fyrup.—A drachm taken night and morning, will keep the belly moderately lax, and greatly eafe the piles.

Hydragogue electuary.

Take gamboge, two ounces; vitriolated tartar and powder of ginger, of each half an ounce; oil of juniper, two drachms; fyrup of buckthorn, a fufficient quantity.—The dofe is half a drachm twice a week, and is intended for dropfical patients, where they have ftrength to undergo its operation.

Myrrh electuary.

Take compound powder of myrrh, a pound; a quarter of a pound of conferve of wormwood; and make into an electuary, with a fufficient quantity of the fimple fyrup. The dofe, two fcruples, three times in a day.—This is prefcribed in diforders of the female fex, with fuccels.

6

St. Bartholomew's Hofpital.

Nephritic electuary.

Take of lenitive electuary, two ounces; juice of liquorice, one ounce and an half; Venice turpentine, half an ounce; powder of rhubarb, two drachms; mix them into an electuary: a drachm of which is to be taken every morning.——This gently purges, and is directed in all obftructions and foulneffes of the kidneys and ureters.

Cathartic electuary.

Take powder of jalap, one pound; one ounce of powder of ginger; and mix them into an electuary, with fyrup of buckthorn.— Half a drachm of this electuary may be given twice in a weck, in all cafes where cathartics are required.

Specific electuary.

Take one ounce of crude quick filver; extinguish it in an ounce and an half of conferve of roles; add two drachms of powder of ginger; and a fufficient quantity of the fimple fyrup.—'The dole is half a drachm twice in a day, in all cafes where mercurials are proper.

Tin electuary.

Take prepared tin, a pound; conferve of wormwood, two ounces; mix into an electuary, with a fufficient quantity of the fimple fyrup.——Half an ounce of this electuary is given every morning, as a certain fpecific against worms.

Elaterium, the paregoric elixir; acid elixir of vitriol, drawing plaister, common plaister, mercurial plaister, cephalic plaister, cummin-feed plaister, galbanum plaister, strengthning plaister, stop plaister, stomachic plaister, and blister plaister: are all prepared as directed in the Supplement.

Ischiatic

8

Ischiatic plaister.

Take Burgundy pitch, four ounces; euphorbium, one drachm: mix them together. — This is a good plaister for the purpose its title express.

Anodine clyfter.

Take two ounces of jelly of flarch; one ounce of the flyptic tincture; and two grains of extract of opium; mix them together.—This is an excellent clyfter in the bloody-flux.

Common clyfter.

Take of fat mutton broth, half a pint; of linfeed oil, three ounces; of common falt, half an ounce: mix them for use.

Oily clyfler.

Take three ounces of fweet oil, and forty drops of liquid laudanum: mix them for use.

Turpentine clyster.

Take half an ounce of Venice turpentine; mix it with a fufficient quantity of the yolk of an egg; and add thereto half a pint of fat broth. This clyfter is generally preferibed in fits of the gravel and ftone.

Purging extract; the extracts of, campeachy wood, jalap, the bark, opium; and the common fomentation, are made as directed in the Supplement.

Emollient fomentation.

Take mallow-leaves, marsh-mallow leaves, and elder-flowers, of each an handful; linfeeds, fenugreek, and cummin-feeds, of each two ounces: boil them in a fufficient quantity of water till there remains only a quart; which strain for use. Its title expresses its virtues.

oppy

St. Bartholomew's Hospital.

Poppy fomentation.

Take two ounces of white poppy heads; boil them in a quart of water to a pint.—This is an excellent fomentation for painful wounds or ulcers.

Restringent fomentation.

Take of oak bark, an ounce and an half; of fmith's forge water, three pints : boil them together to one quart; to which add a quarter of an ounce of roch-allum. — This is a very powerful aftringent.

Flores martiales

Are prepared as directed in the Supplement.

Common gargle.

Take a pint of barley-water; two ounces of white wine vinegar; one ounce of honey of rofes; and half an ounce of tincture of myrrh: mix them together for use.

.The contrayerva gargle.

Take one ounce of figs; half an ounce of contrayerva-root bruifed; boil them in a fufficient quantity of water to ftrain off twelve ounces: it is fometimes made with the addition of an ounce of vinegar.—This gargle is highly recommended by Dr. Fothergill, in the putrid fore throat.

Palfey gargle.

Take half an ounce of pellitory of Spain; boil it in a fufficient quantity of water, to one pint: then, add half an ounce of fpirits of fal armoniac.

Cordial draught.

Diffolve half a drachm of the cordial confection, in two ounces of the common julap; this draught is to be repeated every fix hours, in cafes where cordials are required.

B 5

Anodine

g.,

Anodine draught.

Take of fpring water, an ounce and an half; fpirituous alexiterial water, and fyrup of white poppy-heads, of each two drachms; of liquid laudanum, fifteen drops: to be taken every night going to reft.

Oily draught.

Take fweet oil, and the common julap without fyrup, of each one ounce; fyrup of marfhmallows, two drachms; powder of gum-arabic, half a drachm; mix them together. One of thefe draughts to be taken three times in a day. Syrup of white poppy-heads is fometimes ufed, inftead of fyrup of marfhmallows.

Oily draught with manna,

Is made by adding three drachms of manna.— These draughts are given in nephritic cases, and diforders of the breast.

The fimple bitter infusion, purging bitter infufion, and infusion of sena, are made as directed in the Supplement.

Laxative infusion.

Take two ounces 'of the infusion of sena; three drachms of Glauber's salt, and half an ounce of solutive syrup of roses: to be taken twice in a week.

Infusion of cascarilla.

Take of the bark of cafcarilla, one ounce; pour upon it a fufficient quantity of boiling water, to ftrain off a pint. The dofe, three ounces twice in a day.—This is a very powerful expectorant.

Infusion of mint.

Take an ounce of the dried leaves of mint, and pour on them a pint of boiling fimple mint water; when

St. Bartholomew's Hofpital.

when cold, ftrain off the liquor, and take a large fpoonful every hour.——This is an excellent ftomachic, and is of great fervice in removing reachings to vomit, and in ftopping fluxes of the belly.

Paralytic infusion.

Take horfe-radifh fliced, and muffard-feed bruifed, of each two ounces; boiling water, a quart: let them infufe for twelve hours, in a veffel well ftopped, and fet in a warm place: to the ftrained liquor add two ounces of fpirituous peppermint water.—The dofe, a quarter of a pint, twice in a day.

Common julap.

Take fimple alexiterial water, and fpring water, of each four ounces; brandy, one ounce and an half; the fimple fyrup, half an ounce.

Chalk julap.

Hydragogue julap.

Take of the common julap without fyrup, fix ounces; of tartar emetic, ten grains; of fyrup of buckthorn, two ounces.—The dofe, a large fpoonful every hour.

Mint julap.

Take fimple mint-water, half a pint; conferve of rofes, one ounce; acid elixir of vitriol, one drachm.—This is an excellent ftomachic; the dofe, three large spoonfuls every fix hours.

ReAringent

12

Restringent julap.

Take of the common julap without fyrup, a pint; of diafcordium, fix drams; mix them together.—The dofe, three large fpoonfuls after every loofe ftool.

Volatile julap.

Take of falt of hartfhorn, two drachms; of fpring water, a pint; of fpirituous alexiterial water, one ounce; and half an ounce of loaf fugar. —The dofe, four large fpoonfuls, three times in a day.

Volatile julap diluted,

Is made, by adding to the former, half a pint of fpring water. — Thefe are generally directed in rheumatic cafes, and in fevers where the pulfe is deprefied.

Ammoniacum milk.

The dose, two large spoonfuls, three times a day, in fits of the asthma. It is prepared as directed in the Supplement.

Ammoniacum milk with oil,

Is made, by adding three ounces of fweet oil, to fix ounces of the ammoniacum milk.—Its use and dose is the fame as the former.

Gommon linetus.

Take conferve of hips, fix ounces; fweet oil, and fyrups of red poppies, of each one pint and an half; add a fufficient quantity of fpirit of vitriol, to make it agreeably tart.—The dofe is a fmall fpoonful, whenever the cough is troublefome.

Common linetus, with fyrup of white poppy-heads.

Take conferve of hips, fix ounces; fweet oil, a pint and an half; fyrups, of red poppies, and of white poppy-heads, of each twelve ounces; fpirit

of

St. Bartholomew's Hofpital.

of vitriol, a fufficient quantity to make it agreeably tart.—The dofe, a fmall fpoonful when the cough is troublefome.

Loboch.

Take spermaceti, and conferve of roses, of each one ounce; of the pectoral syrup, two ounces: to which may be sometimes added, half an ounce of myrrh, or three drachms of nitre — The dose, a tea-spoonful when the cough is troublesome.

Loboch, with Syrup of white poppy-heads.

Take spermaceti, conferve of roles, pectoral fyrup, and syrup of white poppy-heads, of each one ounce. The dose the same as the former. If a cough does not yield to these remedies in a few days, it will be to no purpose to rely on pectorals, especially if it is attended with a fever, or proceeds from a pleurify, or peripneumony; for then it must be cured by plentiful bleeding and purging.

Oleaginous liniment.

Take neat's foot oil, a pint; Venice turpentine, two ounces; Castile soap, three drachms; mix them together for use.

The faponaceous liniment, and the volatile liniment, are both made as directed in the Supplement. Sometimes a fourth part of liquid laudanum is added to the volatile liniment.

Diuretic lixivium.

Take falt of tartar, and loaf fugar, of each a pound; diffolve them in a gallon of fimple peppermint water : to which add a quarter of a pint of fpirituous pepper-mint water.—Two ounces of this lixivium are given three times in a day, in dropfical habits; it is an attenuating and warm diuretic, which acts very powerfully.

Magnesia alba,

Is recommended, for children, by Dr. Cadogan; in diforders arifing from acidities, inflead of the common abforbent powders, on account of their being apt to make the body coffive.

Alkalized mercury, calcined mercury, calomel, honey of rofes, mithridate, are prepared as directed in the Supplement.

Ammoniacum mixture.

Take twelve ounces of ammoniacum milk; fyrup of fquills, four ounces.—A large fpoonful is to be taken when the fhortnefs of breath is troublefome.

Antiemetic mixture.

Take of the faline mixture, half a pint; of liquid laudanum, a drachm.—The dofe, two large spoonfuls every fix hours.

Mixture for the venereal disease.

Take of corrofive fublimate, twelve grains; of brandy, twenty-four ounces; let the mercury diffolve of itfelf, and take a large fpoonful, twice in a day, in water.— This medicine has been ufed with fuccefs in the military hofpitals, but is rather uncertain in its effect, the quantity of mercury introduced into the conflitution being fo extremely fmall. It is of excellent ufe in old ulcers, and in beginning cataracts.

Mixture to promote expectoration.

Take of the infusion of cascarilla, half a pint; of oxymel of fquills, one ounce.——The dose, three large spoonfuls, three times in a day: by adding four ounces of linseed oil, it becomes the expectorating mixture with oil; by adding an ounce of the paregoric tincture, the anodine expectorating mixture. Hysteric

St. Bartholomew's Hofpstal.

Hysteric mixture.

Take a pint of ammoniacum milk; and half an ounce of tincture of affafætida. The dofe, two fpoonfuls as often as occasion requires.——This is a powerful remedy in hysteric cases.

Oil of juniper mixture.

Take oil of juniper, one drachm; loaf-fugar, a quarter of an ounce; of the common julap, half a pint.——Two fpoonfuls are to be taken when the hiccough is troublefome: fometimes forty drops of liquid laudanum are added to this mix ture.

Olibanum mixture.

Take gum olibanum, one ounce; diffolve it in a pint and an half of the common julap.—Three large fpoonfuls of this mixture are given twice in a day, in the fame cafes for which the olibanum bolus is prefcribed.

The oily mixture.

Take of the volatile julap diluted, twelve ounces; fweet oil, fix ounces; fpirit of fal armoniac, one drachm; of the fimple fyrup, two ounces and an half.—The dofe, three large fpoonfuls three times in a day.

Saline mixture.

Take falt of tartar, one ounce and an half; lemon juice, eighteen ounces; fpirituous alexiterial water, half a pint; fpring-water, a quart; loaf-fugar, three ounces.—The dofe, three fpoonfuls, three times in a day.

Saline mixture with rhubarb,

Is made, by adding fix grains of rhubarb to every dofe.——The faline mixture is very efficacious in removing vomitings, and is generally given for a few days in intermitting fevers, before the bark is administred.

Spermaceti mixture.

Take two drachms of fpermaceti; diffolve in the yolk of an egg; fix ounces of fpring water, one ounce of fpirituous alexiterial-water, and half an ounce of the fimple fyrup.—The dofe, two fpoonfuls when the cough is troublefome: fometimes two fcruples of nitre, and fometimes fyrup of white poppy-heads, inftead of the fimple fyrup, are added to this mixture.

Purified nitre, oil of juniper berries, oil of turpentine, oxymel of fquills, and London philonium, are made as directed in the Supplement.

Anodine pill.

Take extract of opium, one grain; Venetian foap, five grains.——This is a very fafe opiate, and may be given every night going to reft, in all cafes where opiates are required.

Strengthning pills.

Take of rhubarb one ounce; of boiled turpentine, three ounces; mix them together.—The dofe is a fcruple twice in a day. These pills are of great fervice in the whites of women, and in stubborn gleets.

Ecphractic pills,

Are prepared as directed in the Supplement.— The dofe is a fcruple twice in a day. This composition is a powerful deobstruent, and is therefore of use in almost all chronic diforders from obstructions.

St. Bartholomew's Hospital.

tions. They keep the body moderately open, and do great fervice in hypochondriacal as well as hyfterical affections.

Alfafætida pills.

Take affafœtida, one pound; fuccotrine aloes, four ounces: powder of ginger, half a pound; and a fufficient quantity of fimple fyrup to make into a mass for pills.—The dose is a scruple twice in a day. These are given in hysteric complaints.

Gum pills and mercurial pills.

Are prepared as directed in the Supplement.-The dose of each is a scruple twice in a day.

Purging pills with mercury.

Take of the purging extract, a scruple; of calomel, ten grains; syrup of buckthorn, a sufficient quanti y.—These are frequently preseribed in old ulcers, and are to be repeated twice in a week.

Squill pills.

Take fresh squills, gum ammoniacum, and black pepper, of each an equal quantity; make twelve pills out of each drachm, two of which pills are to be taken twice in a day.—There is scarce any medicine more effectual in asthmas, and they powerfully promote the urinary discharge.

Capivi mixture.

Take three drachms of balfam of capivi, thirty drops of oil of juniper, a fufficient quantity of mucilage of gum arabic, fix ounces of fpring water, proof fpirit, and fyrup of marfhmallows, of each an ounce.—The dofe, two large fpoonfuls three times in a day, in all diforders of the urinary paffages; it is likewife a powerful balfamic, and good in most diftempers of the breaft.

17

Balfam

Balfam of Peru mixture.

Take of balsam of Peru, one scruple; of mucilage of gum arabic, a sufficient quantity; of the common julap, one ounce and an half;—to be taken twice in a day, in nephritic complaints, inward ulcerations, and diforders of the breast.

Æthiopic powder with rhubarb.

Take of Æthiop's mineral, a pound; rhubarb, fix ounces; ginger, two ounces.—The dofe, from ten grains to half a drachm, twice in a day. This may be given to young children for the worms, all crudities, and an acrimonious tafte of the humours.

Compound powder of armenian bole, with or without opium,

Is prepared as directed in the Supplement.

Dr. Dover's powder.

Take powder'd opium, ipecacoanha, and liquorice powder, of each one ounce; purified nitre, and vitriolated tartar, of each four ounces; mix them well.—The dofe is half a drachm every night going to reft.

This powder is the noftrum, which Mr. Ward used in rheumatic diforders, with great fucces. The sweating should be encouraged by lying between the blankets, and drinking plentifully of the faline mixture, as this powder occasions great heat.

Compound powder of myrrh,

Is prepared as in the Supplement.

Pacific powder.

Take half a scruple of rhubarb, and one grain of the extract of opium : to be taken every night going to reft, in cases where opiates are apt to render the body costive.

St. Bartholomew's Hospital.

Purging powder.

Take scammony, jalap, and sena, of each equal parts: by adding a fourth part of calomel, is made the purging powder with mercury.—The dose is half a drachm three times in a week.

Sneezing powder,

Is made as directed in the Supplement.---This is prefcribed in the head-ach, giddinefs, deafnefs, and other diforders of the head. It is generally adminiftred at night going to reft, and in the morning.

Worm powder.

Take the flowers of tanley, and worm-feed, of each three drachms; falt of fteel, a drachm.---The dofe is a fcruple, three times in a day.---This is a very efficacious remedy against the diforder its title expreffes.

Salt of wormwood, falt of fteel, falt of tartar, bitter purging falt, diuretic falt, volatile falt of hartfhorn, volatile falt armoniac, and alum-whey, are made as directed in the Supplement.

Mustard whey.

Take milk, and water, of each a pint; bruifed muftard-feed, one ounce; boil them together, till the whey is perfectly feparated from the curd.--The dofe is half a pint twice in a day: it is useful in afthmatic, dropfical, and rheumatic cafes.

Sinapism.

Take mustard-feed, and the crumb of a new loaf, of each equal parts; make into a poultice, with a fufficient quantity of the best vinegar.--This is applied to paralytic members; and in fevers, is fometimes applied to the feet, by way of causing a revulsion from the head.

Aromatic

Aromatic Species.

Take Winter's bark, and fweet-cane, of each two pounds; ginger, and long pepper, of each one pound; mix them together.

Lavender drops, spirit of fal armoniac, volatile spirit of association of vitriol, sprups of marshmallows, of white poppy-heads, of red poppies, of roses solutive, simple, of squills, troches for the head-burn, and soluble tartar, are made as directed in the Supplement.

Tinclure of wood-foot,

Is made as directed in the Supplement.--The dofe is a quarter of an ounce in a glafs of water, twice in a day, in hysterical and nervous diforders.

Tinctures, of myrrh, of rhubarb with wine, of rhubarb spirituous, of roses, of hiera picra, of snakeroot, volatile, of guaiacum, and liquid laudanum, are prepared as directed in the Supplement.

Tincture of Sena.

Take of the leaves of fena, one pound; of Jamaica pepper, two ounces; of proof fpirit, a gallon; and half a pint of folutive fyrup of rofes. The dofe, two ounces, in cholicky complaints, as often as occafion may require.

Alcaline aloetic wine,

Is prepared as in the Supplement.---Sixty drops of this, taken every morning, will do infinite fervice in diforders of the female fex, and this medicine is alfo a good ftomachic and alterative.

Ointments, ægyptiacum, white, camphorated, and of marshmallows, are prepared as directed in the Supplement.

Blue ointment.

Take hog's-lard and quickfilver, of each equal parts; turpentine, a sufficient quantity to extinguish the

20

St. Bartholomew's Hospital.

the quickfilver entirely.---This is the common mercurial ointment, which is used to raise a falivation by unction.

Yellow ointment.

Take of quickfilver, one ounce; of ftrong spirit of nitre, two ounces; digest in a fand heat till the mercury is dissolved; to which, while very hot, mix a pound of sweet oil; and stir all well together, till the ointment is quite cold.--- This ointment is particularly efficacious in removing tetterous eruptions, and all diforders of the skin.

Ointment for the leprofy.

Take of the roots of tharp pointed dock, four pounds; boil in eight pounds of mutton fuet; ftrongly prefs out the liquor; and add two pounds of tar, three pounds of tweet oil, and half a pound of brimftone; mix them together for ufe.---An admirable ointment for the ufe its title express.

Tar ointment, ointment of elder, and ointment of brimftone, with or without effence of lemons, are made as directed in the Supplement,

a farmin of chuba by tour marinesto

fars assiding has swallandista one

hand all generative providers and clocklash

or analysis average or and a start of the

From the SURGEONS Books.

laine (22)

Eye water.

TAKE half a drachm of white vitriol, and half a fcruple of camphor, and pour on them fix ounces of boiling water.

Another.

Diffolve ten grains of white vitriol, in an ounce of fpring water.—Either of these will very fasely cool and repel those sharp humours which sometimes affect the eyes, and will clear them of beginning films and specks. If too sharp they may be further diluted by the addition of a little spring water.

An epulotic lotion, by Mr. Potts.

Take fix drachms of white vitriol, and three drachms of camphor, and pour on them a quart of boiling water.

A digestive balfam.

Take twelve ounces of oil of St. John's wort, and four ounces of Arceus's liniment, and mix them together.—This is an excellent application for all kinds of green wounds, efpecially when attended with great pain.

A restringent bolus.

Take a fcruple of rhubarb, four grains of ipecacoanha, and half a drachm of the London philonium.—This does amazing fervice in fluxes of the belly, if taken every night going to reft.

The emollient powder.

Take marshmallows and mallow-leaves, each eight handfuls; chamomile, melilot, and elder flow-

ers,

St. Bartholomew's Hospitals.

ers, each four handfuls; linfeeds and fenugreek feeds, each one pound; and grind them into a fine powder.

Mr. Nourse's anodine cataplasm.

Take one ounce of linseeds, and boil them in a quart of water to a pint, ftrain off the liquor, and diffolve in it two drachms of ftrained opium, then add twelve ounces of the emollient powder, and make the whole of a proper confistence for a pultice. —This is useful to affuage violent pains in any part of the body, particularly in diforders of the eyes.

The suppurative cataplasm.

Take white lilly roots, onions and figs, each one ounce; linfeed flower, one ounce; boil all together in a fufficient quantity of water, and when they are boiled tender, add an ounce of Burgundy pitch.—An excellent fuppurant.

The astringent cataplasm.

Take bean flower and barley meal, each three ounces; Cyprefs nuts, balauftines and roch allum, each one ounce; red rofe leaves, fix drachms; oil of rofes, three ounces; the whites of two eggs; and a fufficient quantity of vinegar, to give the whole, a proper confiftence.

The mustard pultice.

Take flower of mustard and oatmeal, of each equal parts; and a fufficient quantity of vinegar to give a proper confistence. It has been customary to apply these kind of acrid medicines to the soles of the feet with a view of making a revulsion from the head; the service which these irritating applications are of where the head is affected, can arise only from their impressing a stimulus; which, though at first partial, by degrees becomes universal.

The

The opthalmic pultice.

Take half a pint of the allum curd, and mix therewith a fufficient quantity of red rofe leaves powdered, to give it a proper confiftence. This is an ufeful aftringent application for fore moift eyes, and excellently cools and repreffes their defluxions.

Another opthalmic pultice.

Take half a pint of linfeed mucilage, and as much flower of linfeed, as is requifite to make it of a proper confiftence.—This pultice lies as foft as that of white bread and milk, upon inflamed eyes, and is far preferable, in as much as the latter is apt to become four and acrid; inconveniencies from which this is free.

The bougie.

Take three drachms of quickfilver, and two drachms of lead; make an amalgama: then add an ounce of crude antimony in fine powder, and one pound of melted bees-wax.

Another.

Take a pound of yellow bees-wax; of Strafbourgh turpentine, four ounces ; cinnabar of antimony, one ounce; mix them together .- Bougies are of fervice in diforders of the urethra, by diftending the part as well as procuring a suppuration. The properties therefore requifite in the bougie, are a sufficient degree of firmness, that it may be introduced with some force; a suppleness and tenacity that it may conform to the motions of the body, without breaking; a lenient suppurative disposition to bring on a discharge without pain; and a finoothnels of furface that it may lastly, not

St. Bartholomew's-Hofpital.

not only be introduced with more eafe, but that it may lie eafy in the paffage, till it begins to diffolve.

The Green Cerate.

Take diapalma plaister, and ointment of elder, each equal parts, and melt them together.

The Yellow Cerate.

Take three pints of linseed oil, two pounds of yellow wax, and melt the wax in the oil.

The Common Cerate.

Take two pints of the common plaister and unguentum nutritum; and one part of black basilicon; and melt them together.

The Saponaceous Cerate.

Take a pound of litharge in fine powder, a gallon of the fharpeft vinegar, half a pound of caftile foap, ten ounces of wax, a pint of olive oil. Diffolve two ounces of the foap in a quart of the vinegar, mix with the litharge, and evaporate the vinegar over a gentle fire; then add the fame quantity of foap and vinegar, and evaporate again, till the whole of the vinegar is evaporated. —Repeat this four times, then add the oil and wax, and fhake them well together till they are thoroughly melted; then remove the mixture from the fire, and when cold it will have the confiftence of a cerate.

This is contrived for the purpose of a discutient, and seems by its subtility to be able to carry almost any thing off through the pores of the skin.

Sky coloured Collyrium.

Take half a pint of lime water, and half a drachm of crude fal armoniac, and let them ftand together twelve hours in a copper veffel.—This is useful in most diforders of the eyes, particularly in specks and films thereof.

The Specific Electuary.

Take an ounce of quickfilver an ounce and an half of conferve of red rofes, half an ounce of the fimple fyrup, rub them together in a marble mortar, till the globules of the quickfilver entirely difappear

Another Specific Electuary.

Take lenitive electuary, and alkalized mercury, each an ounce; gum olibanum half an ounce; balfam copaiva, and fyrup of rofes, each half an ounce; mix them together. The dofe of either of these electuaries is the quantity of a nutmeg every night and morning: they are usually given in complaints arising from a veneral cause.

Sticking Plaister.

Diffolve two ounces of ifinglass in a pint of brandy or melasses spirits, and add one ounce of tincture of Benjamin.—This is the original receipt for preparing the lady's black sticking plaister.

An Opthalmic Epithem.

Take one ounce of allum in fine powder, the white of an egg, of armenian bole and conferve of roles, each fix drachms, mix them together.— A powerful reftringent, and proper in all diforders of the eyes, arifing from relaxion.

The Discutient Leniment.

Take three drachms of opodeldoc, two ounces of oil of earthworms, and a few drops of the oils of origanum and lavender.

Another

26

St. Bartholomew's Hofpital.

Another Discutient Liniment.

Take half an ounce of Barbadoes tar, half a drachm of oil of origanum, half a drachm of volatile fprits of fal armoniac, compound fpirits of lavender a drachm, powder of euphorbium half a drachm.—Thefe are ufeful in paralytic complaints, being rubbed on the part affected, till the part changes colour, and looks red.

The Opthalmic Fomentation.

Take fix drachms of white poppy heads bruifed, with their feeds, and boil them in milk and water, each half a pint, till one half is confumed away; then diffolve in the ftrained liquor a drachm of white vitriol, and a fcruple of fugar of lead.

An Aftringent Fomentation.

Take oak bark and pomegranate shells each an ounce, balaustine flowers half an ounce, red rose leaves an handful, boil them in sour red wine and lime-water equal parts of each.

The Detergent and Aftringent Gargle.

Take a pint of the decoction of oak bark, two ounces of rectified fpirits of wine, one ounce of tinclure of myrrh, mix them together. — This was used very successfully, to wash the mouth with, ten or twelve times a day, after removing a large fungus from the under jaw.

The Expression of Millepedes.

Take four ounces of live millepedes, and bruife them in a marble mortar, with a pint and an half of the common julap. Then fqueeze out the liquor, and take two ounces, with two drachms of fal polychreft, and a drachm of honey, twice in a day.—This is of fervice in the jaundice, an I C 2 almost

PRACTISE OF

almost all chronic diforders, which foul the glands.

The Injection for the Fluor Albus.

Take two drachms of fugar of lead, an ounce and an half of the fcordium electuary, and a quart of lime-water.

An Injection for a gleet in Men.

Take half a pint of lime-water, a drachm of the foordium electuary, and eight grains of fugar of lead.

A Lotion for the Itch.

Take half a drachm of mercury fublimate, and three drachms of white vitriol, diffolve them in a pint of warm water; and with this lotion wash the parts affected, two or three times a day.

The Discutient Mixture, or Mindererus's Spirit.

Take as much of crude fal armoniac, as is fufficient to thoroughly faturate a quart of the beft white wine vinegar, and mix them together. — This is a most admirable difcutient; it dispersed a tumour in a girl's knee, which appeared from the fluctuation, to contain at least an ounce of a glairy fluid.

An Æthiopic Pill.

Take quick-filver, and golden fulphur of antimony, each four ounces; gum guaicum two ounces; mix them well together in a marble mortar; and then add four ounces of caffile foap, and a fufficient quantity of the balfamic fyrup to give it a proper confiftence.—Thefe pills refemble those of Doctor Plumber, described in the Medical Eflays, to which they are preferrable in one respect, they are lefs apt to run off by ftool.—This medicine is an excellent

28

St. Bartholomew's Hofpital.

29

cellent alterative in cutaneous and venereal diforders; three or four middle fized pills are to be taken every night and morning, the patient keeping moderately warm during the whole time that this course is " continued.

A Mercurial Pill. P. P.

Take two ounces of quickfilver, thirty drops of spirit's of turpentine, half an ounce of fyrup of buckthorn, and as much liquorice powder as is necessary to give it a proper confistence.- The dofe is from ten grains to half a drachm every night going to reft.

Another.

Take quickfilver seven grains; turbith mineral one grain ; extract of opium barely a grain : and mix into a pill with the fimple fyrup; which is to. be taken every night going to reft.

Another. Ed. N.

Take quickfilver, fyrup of buckthorn, and clarified honey, each one ounce; powder of jalap, ten drachms; rub the fyrup and quickfilver together, till the globules entirely disappear; then add the jalap and honey, and make fix pills out of each drachm .- The dose is one pill every night going to reft.

All these mercurial pills are capable of doing great fervice in the venereal and chronic diforders. Belloste's pills are somewhat similar to these last, if the analysis which has been made of them is just.

Restringent Pill.

Take of the peruvian bark in fine powder, and gum olibanum, each half an ounce; rhubarb two drachms; and make them into pills with Strafburgh turpentine. - These are good in all. weakneffes

weakneffes of the reins, and will be of great fervice in the whites, and gonorrhæas, after the virulence is removed. Three or four pills are a proper dofe, every night and morning.

The Olibanum Pill.

Take gum olibanum and alkalized mercury, each half an ounce; powder of rhubarb, two drachms; fyrup of rofes folutive, as much as is neceffary to give a proper confiftence, and divide, the mafs into pills of a middling fize—Two or three of which may be taken twice in a day in gleets or weakneffes of either fex, where there is a fufpicion the virulence is not entirely removed.

Aftringent Drink.

Take pomegranate shells, balaustine flowers, and red role leaves, each two drachms; boil in two quarts of water till one half is wassed away; to the restrained liquor add a quart of new milk, and use it for common drink.

The Gaussic Powder.

Boil a pint of foap leys to one half, and add feven ounces of quick lime in fine powder; mix them together, and keep the powder in a glass bottle, well ftopped from the air.

This, mixed with fost soap, is the caustic usually made use of to open abscesses, &c.

Epulotic Powder.

Take prepared calamine, and powder of myrrh, each equal parts, and mix them together.

Tooth Powder.

Take myrrh, roch allum, dragons blood, and cream of tartar, each half an ounce; and make them into a very fine powder; — This, tho' fimple,

18

30

St. Bartholomew's Hospital.

is an excellent dentrifice; but nothing of this kind fhould be applied too frequently to the teeth, for fear of hurting their enamel.

The Sarsaparilla Bolus.

Take half a drachm of farfaparilla in fine powder, half a fcruple of gum guaiacum, a fcruple of gum arabic, and as much balfam capivi as is fufficient to give it the confiftence of a bolus. This is frequently ordered by Mr. P. P. to be taken every morning in a gleet remaining after a falivation.

A successful Method of Cure in the Dead Palsy, by J. F.

Take an ounce of tincture of hiera picra, half a drachm of spirits of turpentine, and fix drachms of syrup of buckthorn; mix them into a draught -to be taken every night going to reft.

Take fpirits of turpentine and oil of origanum, each one ounce; oil of amber two drachms; ointment of marshmallows two ounces; mix them into a liniment;—to be rubbed on the part affected as occasion may require.

Paralytic Tincture.

Take of the peruvian bark, and valerian root, each half an ounce; fpirits of fal volatile five ounces; let them digeft together for four days, in a glafs vefiel closely ftopped, then ftrain off the tincture; —The dofe is from half a drachm to a drachm, three times in a day, in a glafs of water; it is an excellent medicine in all nervous and cephalic diforders.

Tincture of Myrrh.

Take myrrh, fuccorrine aloes, dragons blood, farcocol, olibanum, each half an ounce; euphorbium powdered three drachms; digeft them for C 4 three three weeks, in three pints of brandy or melaffes fpirits.—This tincture is contrived principally for chirurgical uses.

Yellow Basilicon.

Take yellow rozin, bees-wax and mutton fuet, each half a pound; linfeed oil ten ounces, venice turpentine three ounces; melt all the ingredients together, except the turpentine; and when they are all melted, take it off the fire, add the turpentine, and keep ftirring it till the ointment is quite cold.

Another by P. P.

Take bees-wax and mutton fuet, each half a pound; linfeed oil a pint; venice turpentine four ounces; melt them together, and ftir the ingredients, till the ointment is quite cold.

Tar Ointment.

Take Barbadoes tar and mutton fuet, equal parts of each, and melt them together. — This is ufed as a plaifter for children's fcald heads; but care must be taken to give proper medicines internally, during the use of this external application.

Another.

Take tar, and the yellow mercurial ointment, each equal parts; mix them together. — This is used for the same purposes as the former, and requires the same caution.

Opthalmic Ointment.

Take prepared tutty, and lapis calaminaris each a drachm; camphor a scruple; white vitriol half a drachm; ointment of elder two ounces. —This receipt is taken from the experience

32

St. Bartholomew's Hofpital.

ence of a very eminent person, who used it many years with great success, in disorders of the eyes.

Lead Ointment.

Take calomel finely levigated two drachms, fresh lime-water half a pint : let them stand together for two days, then strain off the liquor andkeep the powder.

Take of this powder, and black lead levigated, each two fcruples, and mix with an ounce of ointment of elder flowers.

This application was once made use of in a noli me tangere, when it succeeded happily; many things had been tried before, but met with no fucces.

It also once cured a very bad cutaneous diforder of the scorbutic leprous kind, which was spread a over both arms in great blotches.

The following method cured a girl of a contraction of the muscles in the calf of her left leg, under which she had laboured some months, and by means whereof her soot was drawn quite inwards, so that she could not put it to the ground.

A blifter was applied to the calf of the left leg.

A vomit of half a drachm of ipecacoanha., was administered twice a week.

Half a drachm of valerian was given every. fix hours.

The warm bath was used twice in a week.

In scrophulous tumours of the glands the following method bids fairest to effect a cure.

Take e

33-

Take crude antimony in fine powder, and burnt fponge, each a fcruple; calomel prepared, a grain; mix them together: — to be taken every night and morning, drinking half a pint of fea-water after each dofe.

But to fay the truth, the fuccess depends chiefly on the sea-water, which succeeds as well when used fingly.

the set with the set of the set o



they a starting company manufacture and and

and the set of the set of the state of the set of the

hairse in grant store

(35).

St. THOMAS's HOSPITAL.

Vinegar of Squills.

DOSE from twenty to fixty drops twice a day.

Antimonial Æthiops.

Take of crude antimony, and quick-filver, of each one ounce; flour of brimftone, half an ounce: mix them well together.—The dofe is a fcruple twice in a day. This is an admirable medicine, much preferable to the æthiops of the fhops, and of great efficacy in all cutaneous diforders.

Æthiops Mineral

Is prepared according to the Supplement.-Its dofe is a fcruple twice in a day

Vegetable Æthiops.

Take of the quercus marinus calcined, a drachm twice in a day.—Many furprifing cures have been effected in fcrophulous cafes, by means of this æthiops.

Allum Water and Simple Lime-water

Are prepared as directed in the Supplement, and are used for the same purposes.

Barley-water, fimple alexiterial, fimple cinnamon, fimple pepper-mint, fimple common mint, fimple penny-royal, and rofe-water are prepared as directed in the Supplement.

Spirituous alexiterial-water, fpirituous alexiterial-water with vinegar, fpirituous water of orange-peal, fpirituous water of juniper, fpirituous water of common mint, fpirituous water of nutmeg, fpirituous water of carraways, are prepared as directed in the Supplement,

Barley-Water with Gum Arabic,

Is prepared by diffolving an ounce and an half of gum arabic in a quart of barley-water,—to be taken a quarter of a pint three times in a day. This is to be used as common drink in gravelly complaints.

Barley-Water with Nitre.

Diffolve a quarter of an ounce of nitre, in a quart of barley-water.—The dose, four ounces three times in a day: it is used in all diforders of the urinary passages.

Lambeth-Well's Water.

The dose is half a pint three times in a day.

Aqua Liberans.

Take of the fhavings of faffafras, four ounces; guaiacum, one ounce; liquorice-root, two ounces; coriander-feeds bruifed, one ounce; infufe them in one gallon of lime-water. The dofe, fix ounces twice in a day.—This is good in all cafes that require abforbents and fweetners, efpecially in venereal affections, after falivation.

From four ounces to a pint of fea-water, may be every morning in fcrophulous cafes.

Purging Water with Manna.

Boil two quarts of Sydenham-wells water to one quart, and diffolve therein one ounce of manna; to be taken in the morning early, and repeated as often as occasion may require, in cases where cooling physic is proper.

Waters, Sapphire coloured, Sulphurated, Vitriolic, Camphorated, Blue, are prepared according to the Supplement.

Balfams of Lucatellus, and of Sulphur, are prepared according to the Supplement.

Alexiterial

St. Thomas's Hospital.

Alexiterial Bolus.

Take of the alexiterial powder, twenty-four grains; fimple fyrup, a fufficient quantity to make into a bolus.—This is frequently directed to be taken to promote a diaphorefis, after any operation which is attended with great pain, and in feverifh diforders; and is to be repeated every fix hours, drinking after each bolus a few fpoonfuls of the common julap.

Alexterial and Epileptic Bolus.

Take of the alexiterial and epileptic bolus, of each a drachm; mix them together,—to be taken every fix hours.

Alexiterial Bolus, with Caftor.

Take of the alexiterial bolus one drachm; of Ruffia-caftor, fix grains; mix them together,—to be taken every fix hours, in nervous fevers.

Alexiterial Bolus, with the Cordial Confection.

Take of the alexiterial bolus, one drachm; of the cordial confection, half a fcruple; mix them together: —to be taken every fix hours, to promote a difcharge by the fkin.

Alexiterial Bolus, with Nitre.

Take of the alexiterial bolus, one drachm; of purified nitre, eight grains; mix them together:to be taken every fix hours. This is lefs heating than the foregoing.

Alterative Bolus.

Take of gum guiacum, half a fcruple; æthiops mineral, half a drachm; lenitive electuary, one drachm; fimple fyrup, a fufficient quantity; mix them together:—to be taken twice in a day. This is a most efficacious prescription, in all foulness of blood whatsoever.

Alze

Aluminous Bolus.

Take conferve of roles, one fcruple; allum, half a fcruple; fimple fyrup, a fufficient quantity.— This is very aftringent, and is directed to be taken, twice in a day, in obstinate fluxes.

Antimonial Bolus.

Take of the æthiops mineral, and crude antimony, a fcruple each; conferve of elder, half a fcruple; fimple fyrup, a fufficient quantity.—This is to be taken twice in a day.—Vide antimonial æthiops, for its virtues.

Antimonial Bolus, with the Aromatic Species,

Is made, by adding four grains of the aromatic fpecies to the foregoing bolus.—Its dofe is the fame, and is used for the fame intentions, as the antimonial æthiops.

Aromatic Bolus.

Take of toafted nutmeg, electuary of bole armoniac with opium, each four ounces; chalk prepared, one ounce; fimple fyrup, a fufficient quantity; mix them together:—a drachm of this bolus is directed to be taken three times in a day, with a draught of the white drink, in diarrhœas, after a few dofes of rhubarb have been previoufly given.

Bolus for the Gout.

Take of the best honey, three ounces; spirit of turpentine, two ounces: mix them together. The dose is a drachm twice in a day.—This is of great efficacy in the sciatica, or hip gout: but the use of spirits of turpentine requires caution.

Astringent Bolus.

Take of conferve of roles, twelve ounces; japan earth, three ounces; elixir of vitriol, and alum, of each an ounce; fyrup of red poppies, a fufficient quantity; mix them together.—This is a powerful

St. Thomas's Hospital.

powerful aftringent, and is given, a drachm three times a day, in obstinate fluxes, and a relaxed state of the bowels.

Balfamic Bolus.

Take Lucatellus's balfam, one ounce; conferve of rofes, two ounces; fyrup of red poppies, a fufficient quantity.—This is given in diffempers of the breaft and lungs, and in all fuspicions of inward ulcerations, as also upon any accidental bruifes; the quantity of a drachm three times in a day, with a draught of the pectoral decoction after each dofe.

Balfamic Bolus, with Balfam of Sulphur.

Take of the balfamic bolus, fix ounces; balfam of fulphur, one ounce; mix them together. —The dofe of this is the fame as the former, and is used for the fame purposes.

Calomel Bolus.

Take of calomel, ten grains; conferve of rofes, half a drachm; mix them together.—This is given over night twice in a week, to be carried off by the common purging draught, in any cafes that require brifk purging.

Steel Bolus.

Take of conferve of wormwood, twenty-four grains; powder of steel, twelve grains; ginger and winter's bark, of each three grains; simple fyrup, a sufficient quantity. This is directed in a chlorofis, and all menstrual obstructions; likewise in all decays of the constitution from chronic diseases. —It is to be taken twice in a day, using as much exercise as is consistent with the condition of the patient.

Chamomile Bolus.

Take of the chamomile flowers, half a drachm; fimple fyrup, a fufficient quantity; to be taken three three times in a day. This is given to firengther the flomach, in the heart-burn, and in agues.

40

Compound chamomile bolus.

Take of chamomile flowers, half a drachm; allum, and myrrh, of each five grains; fimple fyrup, a fufficient quantity.—Its dofe is the fame as the former, and they are both fometimes given in intermitting fevers.

Copaiva bolus.

Take lenitive electuary, and balfam copaiva, of each one ounce; powdered rhubarb, thre drachms; mix them together.—A drachm may be given twice a day, with fuccefs, in gleets remaining after a falivation, and to compleat the cure in veneral gonorrhæas, after the virulence is removed.

Strengthening bolus.

Take of yellow colophony, a fcruple; rhubarb, and conferve of rofes, of each ten grains; fimple fyrup, a fufficient quantity.—To be taken twice in a day.

Cleanfing bolus.

Take of gum guaicum, and loaf fugar, each half a fcruple; flour of brimstone, two fcruples; fimple fyrup, a fufficient quantity.—To be taken twice in a day, in diforders of the skin, or rheumatic cases.

Bolus for the Diarrhæa.

Take of rhubarb, fifteen grains; electuary of armenian bole with opium, half a drachm; fimple fyrup, a fufficient quantity.—This may be given in all fluxes of the belly that require to be fropped as often as the urgency of the fymptoms indicate; but it is generally directed only at night going to reft.

Epileptic

St. Thomas's Hofpital

Epileptic bolus.

Take two ounces of the powder of valerian; fimple fyrup, fix ounces; mix them together.—The dofe is a quarter of an ounce twice in a day. It is a good medicine in nervous or paralytic diforders.

Epileptic Bolus, with Powder of Steel.

Take of the epileptic bolus, a quarter of an ounce; powder of steel, ten grains.—This is to be taken twice in a day, in epileptic complaints, accompanied with an obstruction of the menstrual discharge

Epileptic Bolus, with Nitre.

Take of the epileptic bole, a quarter of an ounce; purified nitre, ten grains.—To be taken twice in a day, where epileptic complaints are attended with feverish symptoms.

Epileptic Bolus, with Sal Armoniac.

Take of the epileptic bole, a quarter of an ounce; fal armoniac five grains.—To be taken twice in a day.

Febrifuge Bolus.

Take powder of the peruvian bark, two ounces; fimple fyrup, fix ounces; mix them together.— The dofe is a quarter of an ounce every two hours in intermitting fevers.

Febrifuge bolus, with allum.

Take of the febrifuge bolus, a quarter of an ounce; allum, ten grains.—To be taken every fix hours, in obstinate agues.

Febrifuge bolus with Myrrh.

Take of the febrifuge bolus, a quarter of an ounce; myrrh, ten grains.—To be taken every fix hours, in the fame cafes.

Febrifuge

Febrifuge Bolus, with Sal Armoniac.

42

Take of the febrifuge bolus, a quarter of an ounce; fal armoniac, three grains.—To be taken every fix hours, in the fame cafes.

Febrifuge Bolus with Salt of Steel.

Bolus for a Gonorrhæa.

Take of powdered jalap, three ounces; liquorrice powder, and fcammony, each one ounce; Venice turpentine, ten ounces; mix them together.—The dofe is a drachm every morning. This is a brifk purge, and is directed in a gonorihæa, to be taken till the matter becomes well coloured; when it is left off, for things more ftrengthening.

Guaicum Bolus.

Take of the gum guiacum, and conferve of elder, each half a drachm; fimple fyrup, a fufficient quantity.——To be taken every morning in rheumatic diforders.

Guaicum and Arthritic Bolus.

Take of the guaicum bolus, one drachm; of the arthritic bolus, half a drachm; mix them together.—To be taken every morning, in rheumatic and gouty complaints.

Bolus for the piles.

Take of the lenitive electuary, and flour of brimftone, each half a drachm; folutive fyrup of rofes, a fufficient quantity.—To be taken twice in a day. This is very good for what the title expresses, especially where perfons are subject to be costive, and also in diforders of the skin.

Jalap

St. Thomas's Hospital.

Jalap Bolus ...

Take of powdered jalap, half a drachm; powder of ginger, fix grains; fyrup of buckthorn, a fufficient quantity.—This is a very fafe and brifk cathartic, and is directed to be taken twice in a week, in all cafes where purging is required.

Tin Bolus.

Take of prepared tin, a drachm; conferve of rue, half a drachm; fimple fyrup, a fufficient quantity.—This is a specific against worms, and is directed to be taken twice in a day.

Nephritic Bolus.

Take of lenitive electuary, twelve ounces; Strafburgh turpentine, four ounces; cream of tartar, one ounce; mix them together.—A drachm is directed to be taken twice in a day, in all obstructions and foulness of urinary passages.

Paralytic Bolus.

Take of mustard-feed, two ounces; carrawayfeeds, and winter's bark, of each two drachms; ginger, one drachm; spirits of turpentine, one drachm and an half; simple syrup, a sufficient quantity.—An excellent medicine for the use its title expresses, and is to be taken a drachm twice in a day.

Itch Bolus.

See the bolus for the piles. Page 42.

Pepper Bolus.

Take of conferve of rue, three ounces; long pepper powdered, one ounce and an half; fal gem, fix drachms; fimple fyrup, a fufficient quantity.— The dofe is a drachm twice in a day. It is precribed in diforders of the ftomach, and bad digeftion proceeding from a cold caufe.

Rhubarh

44

Rhubarb Bolus.

Take of powder of rhubarb, half a drachm; fimple fyrup, a fufficient quantity;—to be taken twice in a week, as often as occasion may require.

Rhubarb Bolus, with Jalap.

Take of the rhubarb bolus, a drachm; powder of jalap, ten grains; mix them together: — to be taken as the former.

Saponaceus Bolus.

Take hard foap, one ounce; oil of anifeeds, and carraway-feeds, of each half a drachm; fimple fyrup, a fufficient quantity.—The dofe, a drachm three times in a day.

Saponaceous bolus, with hiera picra.

Take of the faponaceous bolus, one drachm; hiera picra, four grains; mix them together:--to be taken twice in a day.

Saponaceous bolus, with Rhubarb.

Take of the faponaceous bolus, one drachm; powder of rhubarb, ten grains; mix them together: --to be taken twice in a day.

These saponaceus bolusses are prescribed with fuccess, not only in the stone and gravel, but also in the jaundice and asthma; in the most obstinate cases of which they are of the greatest fervice.

Squill bolus.

Take of elecampane, and fquills powdered, each one ounce ; oil of anifeeds, half a drachm ; fyrup of marshmallows, a sufficient quantity:—The dose is one drachm twice in a day.

In affhmatic and dropfical habits, this is an admirable medicine : but its use must be complied with for some time; and if so continued, the most obstinate obstructions of the small vessels, that cause

St. Thomas's-Hospital.

cause both dropfies and afthma's will give way to its efficacy.

Laxative bolus.

Take of lenitive electuary, one ounce; powder of jalap, two drachms; mix them together: the dofe is one drachm and an half, twice in a week. —This is a fafe and effectual family purge.

Laxative bolus, with mercury.

Take of the laxative bolus, one drachm; calomel, ten grains; mix them together; it is to be taken twice in a week.

Bolus for a Scrophula, or King's-Evil.

Take of burnt fponge, one drachm; conferve of roles, half a drachm; fimple fyrup, a fufficient quantity.

Bolas for the Scrophula, or King's-Evil, with Nitre.

Take of the bolus for the fchrophula, two drachms; nitre, ten grains; mix them together.—These bolusses are directed frequently in scrophulous habits; to be taken every night and morning, with a draught of sea-water, after each dose.

Tartar bolus.

Take of conferve of wormwood, half a drachm; cream of tartar, and tartar of vitriol, of each half a fcruple; falt of steel, five grains; fimple fyrup, a fufficient quantity.—This is directed to be taken twice in a day.

Mercurial emetic bolus.

Take of turbith mineral, five grains; emetic tartar, three grains; make into a bolus, with a fcruple of conferve of rofes.—This is fometimes given to caufe a revulfion, in an obftinate Hernia Humouralis; but as it will frequently operate with great violence, it is unfafe in infirm and weakly conftitutions.

Bolus for the Small-pox.

Take of prepared crab's-claws, eighteen grains ; purified nitre, fix grains; fimple fyrup, a fufficient quantity.—This is given every fix hours, with a few spoonfuls of the common julap, during the eruption of the small-pox.

A drachm of the powder of calamus aromatic we directed to be taken three times in a day, in all fuch like complaints arifing from indigestion, and a cold ftomach.

From ten grains to two scruples of Ruffia caftor are prescribed three times a day in nervous cafes.

The Cummen-feed cataplasm, and ripening cataplasm, are prepared according to the Supplement.

Cataplasm against the Rheumatism.

Take of fresh elder-flowers, fixteen pounds; the best vinegar, four pints; French sea-falt, one pound; mix them together :---apply a sufficient quantity to the part affected, every other day.

The White, the Yellow, and Mercurial Cerates are prepared as directed in the Supplement.

Common Cerate.

Take of yellow wax, and oil of olives, each one ounce; mix them together.

Vitriolic Collyrium.

Take ten grains of white vitriol, and diffolve in two ounces of water; if a double quantity of water is made use of, it is called the Vitriolic Collyrium diluted.—In all hot humours the eyes may be washed with a little of either of these collyriums, at discretion.

A scruple of the cordial confection of the London Dispensatory, is ordered to be taken every fix hours, to promote a diaphorefis.

2

St. Thomas's Hospital.

Mithridate, conferve of red rofes, conferve of elder, conferve of rue, conferve of hips, and conferve of wormwood, are prepared according to the Supplement.

White Drink.

Take of burnt hartshorn, fix ounces; gumarabic, one ounce; boil them in a sufficient quantity of water, to strain off a gallon.—This is used for common drink in fluxes of the belly, and in the small-pox and measles, when apprehensive of a looseness.

White Drink, with Gum Arabic,

Is make by diffolving five ounces of gum arabic in a gallon of the white drink :—its dofe is a quarter of a pint three times in a day. This is used in the fame cafes as the former.

White Drink, with Nitre.

Diffolve one ounce of nitre in a gallon of the white drink : its dofe is the fame as the former— This is given in those diarrhœas which are accompanied with a fever.

Aftringent Decostion.

Take of burnt hartfhorn, fix ounces; gum arabic, and pomegranate-fhells, each one ounce; tormentil root, four ounces; boil them in a fufficient quantity of water to ftrain off a gallon.—A quarter of a pint of this decoction, is to be taken three times in a day, in diarrheeas.

Decoction of Logwood.

Boil eight ounces of logwood, in a fufficient quantity of water, to ftrain off a gallon:—its dofe is the fame as the former, and it is used for the fame intentions.

Common Decoction for a Glyster Is prepared according to the Supplement.

Decoction

47

Decoction of the peruvian bark.

Take of the peruvian bark, grossly powdered, half a pound; boil it in a fufficient quantity of water to strain off a gallon:—Dose three ounces, three times in a day,

Decoction of the Peruvian bark, with Winter's Bark.

Take of the peruvian bark grossly powdered, two ounces; boil it in four pints of water to a quart; when almost boiled, add an ounce of Winter's bark :- The dose is the fame as the former.

Decoction of the Peruvian Bark, with Tincture of Bark.

Take of the decoction of the bark, two ounces and an half; volatile tincture of the bark, a drachm; mix them together.—This dose is to be taken three times in a day. These are prescribed in intermitting fevers, to persons of a delicate constitution, whose store and the bark in start of the bark in fubflance.

Dietetic Decoction.

Take fhavings of faffafras, and guaicum, each one ounce; liquorice root, two ounces; coriander feeds bruifed, half an ounce; boil them in a fufficient quantity of water to ftrain off a gallon.— This is used as a fweetner of the blood, in all obftinate cutaneous eruptions, and pocky complaints. Its dose is half a pint twice in a day.

Decostion of Sarsaparilla.

Take farfaparilla, eight ounces; boil it in a fufficient quantity of water to ftrain off a gallon :-----The dofe is half a pint four times in a day. This is very efficacious in removing those pains which frequently remain after falivation.

Decostion

St. Thomas's Hofpital.

Decoction of Simarouba.

Take of the bark of Simarouba, two ounces; boil it in a gallon and a half of water, to a gallon: its dole is a quarter of a pint, three times in a day.—This is a very powerful remedy in all kinds of fluxes of the belly.

Vulnerary Decoction.

Take of ground-ivy, colts-foot, and liquoriceroot, each two ounces; elecampane, one ounce; boil them in nine pints of water, to a gallon.—This is directed to be taken for common drink; or a quarter of a pint three times in a day, in all diftempers of the breaft.

Decoction of Elm-bark.

See this article in the Difpensatory of St. Bartholomew's Hospital. Page 5.

Bay-berry Electuary,

Is prepared according to the Supplement,

Armenian Bole Electuary, with Opium.

Take of the compound powder of armenian hole, with opium, one ounce; fyrup of white poppy heads, three ounces; mix them together.—It is a reftringent, and is given in diarrhæas, rhubard having been previoufly administred.

A drachm of the lenitive electuary, prepared-according to the Supplement, is given in collive habits, as often as occasion requires.

Scammony Electuary, of the London Difpensatory, Is given, a drachm and an half twice in a week, as a brifk purge in rhumatic diforders.

Electuary of Scordium, with Opium, Is prepared according to the Supplement.

Volatile

PRACTICE OF

Volatile Acid Elixir.

Take of the volatile aromatic spirit, fix drachms; fyrup of violets, a drachm; dropt in spirits of vitriol, till the fermentation ceases, and the liquid begins to look red. —Its dose is a drachm twice in a day; sometimes a scruple of camphor is added.

Elixir of Aloes,

Of the London Difpenfatory, is given from twenty to fixty drops every morning.—This is an excellent ftomachic.

A drachm of the Elexir Paregoricum, is given every night in afthmatic cafes, which require the use of opiates.

Twenty drops of the Acid Elixir of Vitriol, of the London Difpenfatory, are given to ftrengthen digeftion, brace up the fibres, and ftop those sweats, which so much weaken the constitution, in confumptive diforders.

The following plaisfers, Drawing, Cephalic, Common, and Cummin-feed, are made according to the Supplement.

Cummin-seed Plaister, with Opium.

Take twelve ounces of cummin-seed plaister; ftrained opium, one ounce; mix them together.

Cummin-feed Paifter, with Soap.

Take three ounces of the cummin-feed plaister, and one ounce of Castile soap, mix them together -These are both excellent discutients.

Strengthning Plaister for the Back.

Take eight pounds of the ftrengthning plaister; of the London Dispensatory, and half a pound of yellow wax; mix them together.

Str. ngth

St. Thomas's Hofpital.

Strengthning Plaister for the Back, with Opium.

Take of the above ftrengthning plaifter, twelve ounces; ftrained opium, one ounce; mix them together.—Thefe are directed to be applied to the loins, in all weakneffes of those parts, from what cause soever; they are serviceable in the whites, and for children who are ricketty; they should be spread to cover quite down the spine, from the nape of the neck, to the bottom of the back.

Galbanum Plaister.

Take ftrained galbanum, one ounce and an half; yellow wax, and myrrh, each half an ounce; Venice turpentine, two drachms; mix them together.— This admirably warms, foftens, and difcuffes indurations of every kind.

Ischiatic Plaister.

Take of Burgundy pitch, four ounces; euphorbium, a drachm; mix them together.—A good penetrating application for the purpose its title expreffes.

Plaister for the Feet.

Take blifter-plaister, and cephalic-plaister, of each an equal quantity.—In fevers attended with a dilirium, this is frequently ordered; because, by its irritation of the feet, it is supposed to increase the blood's velocity that way, and of course diminish it in the head.

Soap and Blifter Plaisters,

Are made as directed in the Supplement.

Oily Emulfion.

Take of the volatile julap, and water, each fix ounces; oil of almonds, four ounces; mix them together.—This is a good medicine for a cough, and is to be given, three large fpoonfuls, three times in a day.

Anodin?

Anodine Clyfter.

Take of the white drink, four ounces; and diffolve in it two grains of strained opium.—This is used particularly, in fluxes attended with violent griping pains in the bowels.

Aftringent Clyfter.

Take of the white drink, four ounces; and diffolve in it three drachms of the fcordium electuary, with opium.—Either of these clysters may be injected in obstinate fluxes, as often as occasion may require.

Common Clyfter.

Take of the common clyfter decoction, half a pint; honey, two ounces; lenitive electuary, half an ounce; mix them together.

Common Clyfter, with Oil.

Take of the common clyfter, ten ounces and an half; oil of olives, two ounces; mix them together for use.

Emolient Clyster.

Take half a pint of milk, coarfe fugar, and oil of olives, of each two ounces; mix them for ule,

Oily Clyfter,

Take five ounces of warm oil of olives.

large ipoonsu's, pitese

Ansaize

100

Any of these may be injected as often as occasion requires, to procure stools in costive habits.

Clyfter for the Piles.

Take fix ounces of the aftringent fomentation, and inject it warm. Excellent for the blind piles, but must not be used where the patient has been accustomed to an hemorrhoidal discharge.

Tur-

Turpentine Clyfter.

Take of the common decoction, eight ounces; Venice turpentine, (diffolved in the yolk of an egg) and honey, half an ounce each : mix them together. —This gives great relief in nephritic diforders.

Expression of Hog-lice.

Take live hog-lice, three ounces; water, a pint and an half; brandy, half a pint; mix thefe together: — Two ounces of the ftrained liquor are to be taken twice in a day, and this medicine is found to be very beneficial in dropfies, the jaundice, the afthma, and gravel.

Green Expression.

Take of the juice of artichoaks, eight ounces; compound juniper-water, two ounces; mix them together.—Four fpoonfuls of this mixture are fometimes given in the jaundice, on account of its diuretic quality, twice in a day.

Extracts, carthatic, of the Peruvian bark, of liquorice, of logwood, and of opium;

Flowers of Benjamin, flowers of steel, and flowers of fulphur; are made as directed in the Supplement.

Astringent Fomentation.

Take gauls bruifed, one ounce; boil them in a fufficient quantity of water, to strain off a pint : its title expresses its use.

Common Fomentation

Is made as directed in the Supplement, and used to foment old ulcers, wounds, &c.

Fomentation of Poppy-heads.

Take four ounces of white poppy-heads, boil them in two quarts of water, to one quart; fometimes two ounces of vinegar is added to this fomen-

 D_3

tation.

53

54

tation.—This wonderfully affuages, and is therefore directed in ulcers which are very painful, and in all inflammatory tumours.

A fcruple of factitious cinnabar, is used for a fumigation; to be repeated as occasion requires.

Aluminous Gargle.

Take of the common gargle, eighteen ounces; allum, half an ounce; mix them together.—This is used in ulcerations of the tongue and mouth.

Common Gargle.

Take tinclure of roles, a pint; honey of roles, two ounces; mix them together.

Contrayerva Gargle.

Take one ounce of figs, and half an ounce of contrayerva root, boil them in a fufficient quantity of water, to firain off twelve ounces; fometimes two ounces of vinegar, and fometimes half an ounce of tincture of myrrh, are added to this gargle, which is excellent in the putrid fore throat.

Jelly of Starch.

Boil one ounce of flarch in a pint of water, to a proper thickness; then add two ounces of tincture of cinnamon, and half an ounce of the simple syrup. —Four ounces of this jelly are frequently ordered to be taken, three times in a day, in the bloody-flux.

Hydragogue Draught.

Take of fimple cinnamon-water, fix drachms; oxymel of fquills, one drachm and an half; compound fpirits of lavender, half a drachm; falt of tartar, half a fcruple; mix them together.—This is given every night, in all cafes where diuretics are required.

15 Mantell 10 Marsh

Oily

Oily Draught.

Take of the common julap, and oil of almonds, each one ounce; mix them together: fometimes this is made with linfeed oil, or olive oil;—it is to be taken every fix hours.

Oily Draught with Diacodium.

Take two ounces of the oily draught, and one drachm of fyrup of white poppies; mix them together: Its dole is the fame as the former.— These are useful in diforders of the breast and coughs, &c.

Oily Draught, with Manna.

Take of the oily draught, two ounces; manna, three diachms; mix them together.—This is often directed in the gravel and ftone, to be taken every fix hours, and is a noble medicine.

Paregoric Draught.

Take one ounce of water, two drachms of fpirituous alexiterial water, and half an ounce of fyrup of white poppy-heads; mix them together, to be taken whenever occasion requires.—It may be used in all cases where opiates are proper.

Common purging Draught.

Take three ounces of the infusion of fenna, one ounce of fyrup of buckthorn, and half an ounce of carraway water; mix them together.—This is to be taken twice in a week, and is a very fafe purge.

Solutive Draught.

Take three onces of water, Glauber's falt fix drachms, and half an ounce of the tincture of fenna; mix them together. This is directed to be taken as the former, where a gentle purge only is required.

55

Hiera picra,

Is made according to the Supplement.

Bitter Infusion.

Take of gentian root, half an ounce; orangepeel, two drachms; zedoary, one drachm; pour on these ingredients a sufficient quantity of boiling water, to strain off twelve ounces; then add fix drachms of spirituous orange water.—Four large spoonfuls of this infusion are to be taken twice in a day.

Bitter Infusion with Salt of Wormwood.

Take of the bitter infusion, a pint, and diffolve in it a drachm of falt of wormwood. This is to be taken in the same manner as the former, and promotes digestion greatly.

Purging bitter Infusion.

Take fenna, three drachms; gentian root, orange-peel, and carraway feeds, of each half a drachm; infuse these ingredients in sour ounces of boiling water, and, when strained, add half an ounce of carraway-water.—This is to be taken twice in a week.

Bitter Infusion, with the diuretic Salt.

Take of the fimple bitter infufion, a pint; and diffolve in it half an ounce of the diuretic falt. Its dofe is four ounces twice in a day.—Thefe are all excellent ftomachics, and are directed to mend digestion, strengthen the stomach, procure an appetite, and in windy diforders of that organ.

Cascarilla Infusion.

Infuse half an ounce of cascarilla bark in a sufficient quantity of boiling water, to strain off a pint. Its dose is fix spoonfuls twice in a day. See page 14.

Droply

Dropfy Infusion.

Take equal parts of the bitter and paralytic infusions, mix them together.—Take a quarter of a pint twice in a day.

Laxative Infusion.

Take of the infusion of senna, three ounces; manna, one ounce; carraway-water, two drachms; mix them together.—This is to be taken twice in a week: sometimes three drachms of Glauber's salt are added to this infusion.

Laxative Infusion, with Tincture of Senna.

Take three ounces of the laxative infusion, and one ounce of the tincture of fenna; mix them together :-It is to be taken as the former.

Infusion of Senna

Is made according to the London Difpenfatory, except carraway-feeds are used instead of cardamomfeeds.

Buck-bean Infusion.

Take of dried buck-bean, one ounce; infuse it in a sufficient quantity of boiling water, to strain off a quart.—It is directed to be taken a quarter of a pint three times in a day, in rheumatic, scorbutic and scrophulous cases.

Paralytic Infusion.

Take of horfe-rhadish, and mustard-seed, each one ounce; winter's-bark, three drachms; infuse in a pint and half of boiling water; when strained off, add two ounces of carraway-water.—Its dose is a quarter of a pint twice in a day.

A sctuple of ipecacoanha is directed to be taken, with a proper regimen, as an emetic, in all cases where emetics are required.

AAringent

Astringent julap.

Take of the Armenian bole electuary, with opium, half an ounce; fimple alexiterial water, half a pint; fpirituous alexiterial water, two ounces; mix them together, and take three large fpoonfuls after every loofe ftool, or as occafion requires, rhubarb having been previoufly exhibited.

Camphor julap of the London Dispensatory

Is directed to be taken, two ounces every fix hours, to encourage a diaphorefis, in epidemic, deprefied and malignant fevers.

Cordial julap acidulated.

Take of fimple alexiterial water, half a pint; treacle water, two ounces; fyrup of red poppies, half an ounce; mix them together for use.

Common julap.

Take of fimple alexiterial water, half a pint; fpirituous alexiterial water, two ounces; fimple ivrup, half an ounce; mix them into a julap.— A few spoonfuls of either of these julaps are ordered to be taken to wash down the alexiterial bolusses.

Chalk julap of the London Dispensatory,

Ts fometimes made with barley cinnamon water, inftead of fpring water; and fometimes two ounces of nutmeg-water, or tincture of cinnamon, are added to a pint of the julap.

Scordium julap.

Take of the common julap without fyrup, ten ounces; electuary of fcordium with opium, half an ounce; mix them together into a julap.—Three fpoonfuls of either of thefe julaps may be given in diarrhæas, after every loofe ftool. In diarrhæas, rhubaib thould always precede the ufe of reftringents.

Musk

Musk julap of the London Dispensatory,

Is given, four spoonfuls every fix hours, in nervous and other fevers, attended with a low deprest pulse, or convulsions. Musc is also useful in meniacal affections.

Mint julapt.

Take of fimple mint-water, eight ounces; fpirituous mint-water, one ounce; loaf fugar, a drachm; mix them into a julap.—Four large fpoonfuls taken frequently are of great fervice in removing naufeas and reachings to vomit.

Mithridate julap.

Take of the common julap without fyrup, ten ounces; mithridate, half an ounce.—Mix them into a julap.

Venice treacle julap.

Take of the common julap without fyrup, ten ounces; Venice treacle, a quarter of an ounce; mix them into a julap.—Three large fpoonfuls of either of these julaps are to be taken every fix hours. Either of these julaps are very good diaphoretics.

Volatile julap.

Take of the fimple alexiterial water, ten ounces; fpirituous alexiterial water, two ounces; loaf fugar, three drachms; volatile falt of hartfhorn, a drachm and an half; mix them into a julap.—Its dofe, is four fpoonfuls three times in a day, in rheumatic, paralytic or nervous cafes.

Volatile julap diluted.

Take of the volatile julap, and fpring water, each fix ounces; — mix them into a julap; fometimes three drachms of nitre, or four ounces of fweet oil, are added. Its dofe, is three fpoonfuls three times in a day.

Two

Two fpoonfuls of Ammoniacum milk of the London Difpenfatory, are to be taken, three times in a day, in afthmatic complaints.

Ammoniacum milk, with oil.

Take fix ounces of ammoniacum milk, and three ounces of fweet oil; mix them together.—To be taken, three fpoonfuls, three times a day.

Ammoniacum milk, with the afthmatic mixture.

Take of ammoniacum milk, and the afthmatic mixture, each equal parts; mix them together.— Two fpoonfuls to be taken three times a day.

Ammoniacum milk, with the volatile julap.

Take of ammoniacum milk, and the volatile julap, each equal parts; mix them together.—Three large fpoonfuls to be taken three times in a day. Any of these mixtures may be taken in asthmatic diforders, according as particular circumstances may indicate, and in all cases where powerful expectorations are wanted.

Affafætida milk.

Diffolve an ounce of affafætida in a quart of water. This is an excellent medicine in hyfteric diforders, and it is to be taken a large fpoonful frequently.

Restringent milk.

Take oak bark, an ounce; pomegranate-bark, half an ounce; cinnamon, two drachms: bruife them and boil them in milk and water, of each a quart, till half is boiled away.—Two ounces of this are directed to be taken three times in a day: it is an efficacious prefeription.

Pestoral linetus.

Take of conferve of hips, four ounces; pectoral fyrup, and linfeed oil, each four pints; mix them together.

together.—A large fpoonful of this linctus is to be taken when the cough is troublefome.

White liniment

Is made according to the Supplement.

Borax Liniment.

Take one ounce of oil of almonds, the yolk of one egg, and a drachm of borax: mix them into a liniment.

Saponaceous Liniment

Is made according to the London Difpenfatory.

Saponaceous liniment, with the Spirit of Sal armoniac.

Take fixteen ounces of the faponaceous liniment, and one ounce of fpirits of fal armoniac: mix them together. These are both admirable discutients.

Viper liniment.

Take colcothar of vitriol, one drachm; viper's fat, one ounce: mix them together.

Volatile liniment.

Take two drachms of bees-wax, and one ounce and an half of fweet oil, melt them together, and then add an ounce and an half of fpirit of fal armoniac.—This is used in paralytic cafes, and to discuss extravalated blood, or humours, in ftrains, &c.

Lixivium of Tartar

May be taken, from twenty to fixty drops, twice in a day. This is a good aperient, and is given in diforders which have their origin from obstructions.

Loboch.

Take of fperma ceti, and conferve of rofes, each an ounce; of the pectoral fyrup, two ounces: mix mix them together. Sometimes half an ounce of myrrh, or three drachms of nitre, or half an ounce of the teftaceous powder, is added.—It is a good pectoral, and is given, the quantity of a nutmeg, whenever the cough is troublefome.

Sulphur lotion.

Boil fix ounces of brimftone powdered, in a gallon of lime water, till a pint is wafted away.— This is a very cleanly application in the itch, but not fo certain in its effect as the itch ointment.

Egyptian Honey, and Honey of Roses, Are made as directed in the Supplement.

Honey of Roses, with burnt Allum.

Take honey of roles, one ounce; and burnt allum, one drachm: mix them together.—This is used to gargle the mouths of children, in the thrush.

Mixture for Deafness.

Take oil of almonds, three drachms; fpirits of fal armoniac, one drachm; mix them together. —A few drops are to be dropt in the ear, which is afterwards to be ftopped with a little wool, every night going to bed, and generally affords great relief.

Saline Mixture.

Take falt of wormwood, a quarter of an ounce; lemon juice, three ounces; cinnamon-water, fix ounces; mix them together.—Three large fpoonfuls are to be taken every fix hours, in fickness of the ftomach or feverish complaints.

Afthmatic Mixture.

Take oxymel of fquils, half an ounce; fimple pennyroyal - water, one ounce; mix them together.—This is very good for what its title expresses, and

and this dofe is to be repeated twice in a day. Sometimes half an ounce of fweet oil is added to this mixture

Cordial Mixture.

Take of the common julap without fyrup, ten ounces; of the cordial confection, a quarter of an ounce: mix them together. — Three or four large fpoonfuls are to be taken every fix hours, where cordials are proper.

Emetic Draught.

Take a fcruple of ipecacoanha, fix drachms of oxymel of fquills, and an ounce of water : mix them -A very fafe and efficacious emetic.

Febrifuge Mixture.

Take two drachms of falt of wormwood, three ounces of lemon juice, and fix ounces of the common julap: mix them together.—Three fpoonfuls are to be taken every fix hours.

To the Febrifuge Mixture,

Sometimes two drachms of red coral, or two drachms of elixir of vitriol, or two drachms of alixiterial powder, or two drachms of lavender drops, are added, as particular circumftances require.

Oily Mixture.

Take oil of almonds, and pectoral fyrup, of each four ounces: mix them together.—The dole is two spoonfuls three times in a day, in coughs, diforders of the breast, pleurisies, &c.

Olibanum Mixture.

Take gum olibanum, half a drachm; of loaf fugar, a drachm; fpirituous orange-water, two drachms; fpring water, an ounce. To be taken every night and morning.

Vide Guy's Hospital.

Mucilage

63

Mucilage of Quince Seeds

64

Is made by boiling a drachm of quince-feeds in half a pint of water. This is an excellent gargle in foreneffes of the mouth, and to blunt fharp humours.

Nephrytic mucilage.

Take gum arabic, and gum tragacanth, of each one ounce; diffolve in fix pints of water, and add four ounces of fyrup of marshmallows. —The dose is three spoonfuls three times in a day.

Half a drachm of myrrh may be taken, twice in a day, in hyfteric diforders.

Nitre may be given, to the quantity of a fcruple, three times in a day, in inflammatory diforders, gravel, fore throat, and heat of urine, as alfo to prevent mercury from effecting the mouth.

Oil of Sulphur.

Take of the flour of brimftone, one ounce; fweet oil, half a pint; boil them together over a flow fire. This medicine is commended in coughs, afthmas, and confumptions.—Its dofe is from twenty to forty drops three or four times in a day.

Oxymel of Squils

Is made according to the Supplement, and is of great use in phthisicky and asthmatic diforders.

London Philonium

Is a warm opiate, and may be given, to the quantity of a fcruple, at night going to reft.

Barbadoes tar, and rectified spirits of wine, of each equal quantities, mixt together, are used as an embrocation, with success, in white swellings of the joints.

A scruple of the affascetida pill of the London Dispensatory, may be taken, twice in a day, in nervous and hysteric cases.

Squill

Squill pills.

Take fixteen grains of affafœtida, and four grains of fquills powdered ;—mix them into three pills, which are to be taken twice in a day, in the nervous afthma, for which they are excellent.

Balfamic pills.

Take of prepared hog lice, three drachms; gum ammoniacum, a drachm and an half; flowers of benjamin, a drachm; balfam of Peru, and extract of faffron, each fifteen grains; balfam of fulphur, a fufficient quantity to make into a mafs of a due confiftence, which is to be divided into pills of a middle fize, four of which are to be taken twice in a day.—Thefe will do great fervice where the lungs begin to be fluffed with vifcid or acrimonious juices, and will wear off thofe fhort hufky coughs, which are figns of tubercles and beginning ulcerations.

Calomel pills.

Take of the fimple colocynth pill, one ounce; calomel prepared, half an ounce : mix them together.—Half a drachm may be taken twice in a week, in all cafes where mercurial purges are judged proper.

Pills for a catarrh.

Take twelve grains of Rufus's pill, add fix grains of the ftorax pill, and divide into three pills, which are to be taken every night going to reft.—This prefcription is intended to carry the phlegm downwards by ftool, in afthmatic cafes, and catarrhs.

Cinnabar pills.

Take a scruple of native cinnabar, and four grains of conferve of roses, mix them into a pill.— This is to be taken every night and morning. in obstinate, nervous, and cephalic diforders; and, to fay the truth, cinnabar fucceeds better in diforders of the nerves, than any other medicine.

66

Strengthening pills.

Take an ounce of rhubarb, and two ounces of boiled turpentine: mix them well together, and form into middle fized pills; three of which are to be taken, twice in a day, in gleets, fluor albus, running of the reins, and diforders of the kidneys.

Ephrastic pills.

Take of Rufus's pill, and falt of steel, each eight ounces; strained galbanum, four ounces; make them into a mass; a scruple of which is to be divided into three pills, and taken every night.—This composition is a powerful deobstruent, and is of use in all chronic diforders from obstructions.

Three middle fized pills of the gum pills of the London Difpenfatory, are fometimes directed to be taken every night in hyfterical and nervous complaints, with fuccels.— They are rendered purgative by the addition of four grains of fcammony to each fcruple of the gum pill.

Mercurial pills.

Take two ounces of quickfilver, extinguish the mercury with two drachms of ballam of fulphur, then add a drachm of extract of liquorice, and an ounce of gum guaicum : divide this mais into 144 pills, three of which may be taken every night.

Purging mercurial pills.

Take half a drachm of the mercurial pill of the London Dispensatory, every morning.—Either of these may be taken in cases which require the administration of mercury, as circumstances may dictate.

Anodine pills.

Take one grain of strained opium; extract of liquorice, thee grains; mix into a pill,—which may be taken every night, or as occasion requires, where opiates are wanted.

Two middle fized pills of Rufus's pill of the London Difpenfatory, given every night, will purge gently, and prove peculiarly ferviceable in cold conflictutions, indigeftions, and hypocondriacal diforders.

Rufus's pill with Steel.

Take Rufus's pill, and powder of fteel, of each half a fcruple; mix them together, and divide into three pills, which may be taken every night, in a fupprefiion of the menfirual difcharge.

Six grains of the ftorax pill of the London Difpenfatory are frequenly given every night, to procure reft in violent coughs.

Pills for the venereal difease.

Take of calcined mercury, one grain; crumb of new bread, two grains; mucilage of gum tragacanth, a fufficient quantity, to make into a pill, which is to be taken every night. By adding four grains of aloes, is made the purging pill for the venereal difeafe. This is reckoned one of the most efficacious mercurial medicines.

Tragacanth pills.

Take gum tragacanth, and gum arabic, of each eight ounces; diffolve them in half a pint of water, and with eight ounces of liquorice-powder, make into a mafs for pills, of which, take three pills three times in a day.—Thefe are ufed with fuccefs in tickling coughs.

White

White potion.

Take an ounce of balfam copaiva, diffolve it in the yolk of an egg; fyrup of marfhmallows, four ounces; fpring water, a pint; brandy, half a pint.— Four fpoonfuls of this mixture are directed to be taken three times in a day, in all diforders of the urinary paffages : it is alfo a good balfamic, and is preferibed for many diforders of the breaft.

Potion of balfam of Peru.

Take of balfam of Peru, a fcruple; yolk of egg, a fufficient quantity; water, an ounce; brandy, two drachms; fyrup of marfhmallows, a drachm and an half: mix them together for a draught, which may be taken twice in a day, in all diforders of the breaft.

Alexiterial powder.

Take of the teftaceous powder, a pound and an half; powder of contrayerva root, five ounces: mix them together:—Its dole is half a drachm every fix hours, to promote a discharge by the skin.

Powder against the bite of a mad dog, and compound powder of aron-root, are made as directed in the Supplement. The dose of both, is a scruple twice in a day.

Compound powder of armenian bole, with or without opium,

Is given, from three to fifteen grains, in cafes which require aftringents, as often as occasion requires.

Diuretic powder.

Take of calcined egg fhells, a fcruple; falt of wormwood, and nitre, each five grains; mix them together.—This powder is a very forcible diuretic, and may be given, three times in a day, in a proper quantity of any diluting liquid.

A fcruple

St. Bartholomew's Hospital.

A fcruple of the compound powder of myrrh, of the London Difpenfatory, is given, three times in a day, in hyfterical diforders, and obftructions of the menfes; fometimes ten grains of powder of fteel are added to each dofe.

Opthalmic powder.

Take of glass finely levigated, as much as you please.—This is recommended to be blown into the eye, to remove specks on the cornea.

Digestive powder.

Take of cinnamon, nutmegs, and prepared amber, of each ten grains; white pepper, five grains; mix them together.—It is given twice in a day, to mend digeftion, in cold weak ftomachs.

Powder for a bearing down of the anus.

Take of balaustine flowers, and armenian bole, each one ounce, mix them together, and sprinkle a little upon the part affected. An useful prescription for the purpose.

From five grains to a scruple, of the compound fcammony powder of the London Dispensatory, may be given twice in a week.—A good purge for children.

Scammony powder with mercury.

Take of the compound powder of fcammony, three ounces; calomel, one ounce; mix them well together:—The dofe is from ten grains to half a drachm, twice in a week.—This is chiefly used for children in diforders proceeding from worms.

Sweating powder.

Take strained opium, and powder of ipecacoanha, of each two grains; tartar of vitriol, and nitre, each eight grains:—This may be taken every other night, in rheumatic cases, and to promote a diaphores diaphorefes.—This is the fame with Dr. Dover's powder, and is the noftrum used by Dr. Ward for the cure of rheumatic diforders. See page 18.

The fneezing powder is made as directed in the Supplement.

Testaceous powder.

Take of crab's claws prepared, half a pound; oyfter fhells, and red coral, prepared, an ounce and an half each; mix them together.—This is used in the heart-burn, and in all diforders of children arising from acidities.

Compound tragacanth powder, of the London Difpensatory,

May be taken, from a drachm to two drachms, three times in a day.---This is intended for an emollient to foften and heal any internal injuries; for which purpofe it is prefcribed in fpitting of blood, bruifes, and diforders of the kidneys; it cools the urine, and is of great fervice in ftranguries.

A drachm of the diuretic falt of the London Difpenfatory, is to be taken twice in a day, in anafarcous and dropfical cafes.

Allum whey of the London Dispensatory, (See Page 170,) is given, a quarter of a pint twice in a day.---This is highly recommended in the diabetes, and pissing of blood. The method of preparing compound tragacanth powder, diuretic falt, and allum whey, is shewn in the Supplement.

Mustard whey.

Boil an ounce of mustard-feed in a quart of milk:---of the strained whey, half a pint may be taken every night going to rest, in rheumatic, gouty or paralytic cases.

Sinapifm

70

Sinapism.

Take of wheat flower, and flower of muftardfeed, each an equal quantity, and make into a proper confiftence for a poultice, with a fufficient quantity of vinegar.---This is used in paralytic diforders, and is fometimes applied to the feet, in fevers attended with a delirium, to cause a revulsion.

Mercurial Solution.

Take of corrofive fublimate, eight grains; brandy, a pint; mix them together :--- The dofe is half an ounce, twice in a day, in venereal cafes; drinking plentifully of barley-water, or fome other diluting liquor after each dofe.--- This medicine has been attended with great fuccefs in the Military Hofpitals. See page 14.

Aromatic Species

- Is prepared according to the Supplement, and is a good warm ftomachic.

Scordium species of the London Dispensatory, with or without opium,

Is given, from three to fifteen grains, as often as occafion requires, in diforders which require reftringent medicines.

Scordium species, spirits of sal armoniac; of hartshorn, of lavender compound, volatile aromatic, of affascetida, and of turpentine, are made as directed in the Supplement.

Scorbutic juices.

Take of fcurvy-grafs, a pound; brook lime, and water-creffes, each half a pound; Seville orangejuice, ten ounces; mix them together, and take four ounces, twice in a day, in the fpring time, against fcorbutic complaints, and to cleanse the blood from impurities.

Precipitated

Precipatated fulphur of the London Dispensatory,

May be taken, twice in a day, in difeases of the breast, and cutaneous distempers, to the quantity of half a drachm each dose.

Four grains of the precipitated fulphur of antimony, taken every night, is a very efficacious alterative.

Suppositories

Are made of honey boiled to a proper confiftence. They are used only where clysters cannot be injected.

Syrups of marshmallows, of faffron, of white poppy-heads, of red poppies, pectoral, of roses folutive, simple, and of violets, are made according to the Supplement.

Syrup of buckthorn.

Take of buckthorn juice, a gallon; Jamaica pepper, and ginger, each an ounce and an half; coarfe fugar, feven pounds; boil them together into a fyrup. This is used only to mix with other cathartics, to quicken their operation.

Troches for the heart-burn.

Take of prepared chalk, two ounces; prepared crab's claws, an ounce; Armenian bole, half an ounce; nutmeg, ten grains; loaf fugar, an ounce and an half: make them into troches, by dropping on them a fufficient quantity of water.—An excellent remedy for that complaint.

Soluble tartar

Is made according to the Supplement :- Its dofe is a drachm every morning; in obstructions, jaundice, dropfies, &c.

Half a drachm of the aromatic tincure of the London Difpenfatory, may be taken twice in a day, in flatulent complaints.

Equal

Tinctures aromatic, of cantharides, of cinnamon, and of myrrh, are made as directed in the Supplement.

Simple tincture of the bark,

Is made according to the Supplement :- Three drachms are to be taken three times in a day, in intermitting fevers, an impoverished state of the blood, and fome nervous diforders.

Volatile tincture of the bark,

Is made according to the Supplement :--- Its dole is a drachm three times in a day.

Tincture of Aslafætida,

Is made according to the Supplement :-- Its dose is a drachm twice in a day, in nervous diforders.

Tincture of wood-foot,

Is made according to the Supplement :- Two drachms are to be taken twice in a day, in hysterical cases.

Tincture of guaiacum.

Take gum guaiacum, an ounce; melaffes spirit, a pint; digest them together for some days :-- Two spoonfuls are to be taken every morning, in rheumatic diforders.

Volatile tineture of guaiacum.

Take of gum guaiacum four ounces; digeft for fome days in a glafs well ftopped, in a pint and an half of the volatile aromatic fpirit:—Its dofe is a drachm every morning, and it is used for the fame purposes as the other tincture.

Tincture

Tinctures of jalap, and Japan earth,

Are made according to the Supplement :- Two drachms are the proper dole of each.

Tincture of flowers of fleel.

Take flowers of steel, four ounces; digest them a fufficient time in a pint of melasses spirit.—From five to thirty drops are given in histerical complaints, and female obstructions, twice in a day.

Tincture of Steel, with Spirit of Salt.

Take of filings of iron, half a pound; Glauber's fpirit of falt, three pints; rectified fpirit of wine, three pints; macerate the filings of iron in the fpirit of falt, without heat, till the fpirit has corroded the iron; when the fæces have fubfided, the liquor must be evaporated to a pint, and the fpirit of wine be added :- The dose, twenty drops twice in a day, in histeric disorders, and obstructions of the menstrual discharge, and is a much fafer medicine in fanguine habits.

Tincture of black hellebore.

Take of black hellebore, four ounces; cochineal, two fcruples; digeft them a fufficient time in a quart of melaffes fpirit: — The dole is a drachm twice in a day, in the fame cafes as the former.

Tincture of rhubarb.

Take three ounces of rhubarb; fix drachms of carraway-feeds; melaffes fpirit, a pint; fpring water, a quart; digeft them together :- Two ounces are to be taken twice in a week, or as often as occasion requires, in diforders of the bowels and ftomach.

Tinclure of rhubarb, with jalap.

Take tincture of rhubarb, an ounce and an half; powder of jalap, eight grains; mix them together:—To be taken as the former. This is vaftly more purgative.

74

Tincture of rhubarb, with Venice treacle.

Take tincture of rhubarb, an ounce and an half; Venice treacle, a fcruple; mix them together; to be taken every other night.

Tincture of roses.

Take half an ounce of red role buds; oil of vitriol, twenty drops; boiling water, two pints and an half; loaf fugar, an ounce and an half; when cold ftrain off the liquor, and take three ounces, three times in a day. — This is a very grateful julap in all cafes that require coolers and fubaftringents, and nothing is better fuited for washing down boluffes or electuaries of the Peruvian bark.

Tinclure of aloes.

Take fuccotrine aloes, eight ounces; Winter's bark, two ounces; melaffes fpirit, four pints; fpringwater, fix ounces; digeft them together :- The dofe, two ounces as often as occasion may require; fometimes a drachm of compound lavender drops, or half an ounce of fyrup of buckthorn, is added to each dofe. This is a warm cathartic, and is given in nervous and paralytic cafes, and diforders of the female fex.

Saturnine tincture.

Take sugar of lead, and green vitriol, of each two ounces; digest them a sufficient time in a quart of rectified spirit of wine :- The dose, twenty drops twice a day.

Tincture of Inake-root.

Take three ounces of Virginian fnake-root; digeft with two pints of melaffes fpirit.—This is a powerful alexipharmic : two drachms are to be taken every fix hours.

Styptic tinEture.

Take one drachm of calcined vitriol; infuse it in E 2 a quart a quart of brandy in a wooden cafk :- The dofe, two drachms as often as occafions requires. This can be but little depended on, and is used only among other articles for the fame intention.

Thebaic tincture,

Is prepared according to the Supplement ----Its dole is from twenty to thirty drops, every night going to reft, in all cales where opiates are proper.

Tincture of valerian

Is prepared according to the Supplement :- The dofe is three drachms three times in a day, in nervous cafes.

Volatile tincture of valerian.

Take four ounces of valerian; infuse in a quart of the volatile armoniac spirit:—The dose is half a drachm, three times in a day, in the same cases as the former, and where the spirits are much depressed.

Troches of nitre.

Take two ounces of nitre, and half a pound of loaf fugar; make into troches, with the mucilage of gum tragacanth.—These are good in inflammations of the throat, and are more effectual than most gargles contrived for those purposes.

Two drachms of the powder of valerian root, are often prescribed in nervous disorders, to be taken three times in a day, with great success.

Alkaline aloetic tincture of the London Dispensatory

Is directed to be taken, to the quantity of a drachm, every night, as an alterative in all cachexies, and is to be continued for a long time together.

Antimonial wine.

Take faffron of antimony, an ounce; white wine, a pint and an half; digest them together.—Twenty drops

drops may be given twice in a day; and, if continued for fome time, prove an efficacious alterative in chronic diffempers: or two ounces may be given as an emetic.

Ointments of marshmallows and basilicon,

Are made according to the Supplement; and are used, the first as an emollient, the last as the common digestive, in almost all kinds of green wounds.

Emollient and discutient ointment.

Take four ounces of the marshmallow ointment; and one ounce of spirits of turpentine; mix them together.

Emollient ointment with opium.

Take of the emollient ointment, fix ounces; ftrained opium, half an ounce; mix them together.—Either of these ointments may be applied to any part bruised or indurated by obstructed humours; because they attenuate and warm the stagnant matter, and help to breathe it out through the poses of the skin.

Pile ointment.

Take of the white liniment, an ounce and an half; ftrained opium, a drachm; mix them into an ointment.—This will immediately give eafe.

Blue or mercurial ointment.

Take one pound of hog's-lard; quickfilver, half a pound; balfam of fulphur, two drachms; rub them well together until the quickfilver is intirely extinguifhed. This is ufed to raife a falivation, by rubbing two drachms of it, every night, over the hands, arms, legs and thighs of the patient, before a good fire. Sometimes three or four unctions will raife a fpitting, though in fome conftitutions half a pound, nay a pound, of this medicine has been rubbed in, and yet no falivation has followed. E 3 Sometimes the body's being over coffive, will prevent a falivation rifing, in which cafe a gentle laxative will promote it.

Opthalmic ointment.

Take of goole fat, an ounce; prepared tutty, half an ounce; mix them together.—This is to be used every night going to bed, or oftener if occafion requires. Vide Guy's hospital. Page 108.

Pitch ointment, and bliftering ointment, Are prepared according to the Supplement.

Itch ointment.

Take hog's-lard, a pound; flour of brimftone, four ounces; lixivium of tartar, one ounce; effence of lemons, half a drachm; mix them together into an ointment.—This is the common, and a very efficacious ointment for the itch. It hardly ever fails of anfwering its end, if continued long enough, and applied every night.

* ** EN *** **

HERE STATESTATE FILST IN ALL STATES STATES

ANTYNYERS SET IN THE REPORT OF THE PARTY OF THE

source an aligitation or such a

From the SURGEONS Books.

The sharpest white wine vinegar is used as adiscutient.

Lime Water.

AKE two pounds and an half of quick lime, and pour on it a gallon of boiling water; ftir the mixture well, and, when cold, ftrain it off for use .- This is an excellent water for internal as well as external uses; it is not only approved of for inflammations, burns, cutaneous eruptions, the eryfipelas, old ulcers, &c. but it is alfo famous for its virtues in the phthific, dropfy, diabetes, and fcurvy in nothern climates; but how it comes to be remarkably ferviceable against the fcurvy in England, Holland, &c. and as remarkably pernicious in the fame diffemper in Italy, the fouthern parts of France, &c. has perplexed fome eminent members of the academy of fciences at Paris; but is well accounted for by Boerhaave, in his Chymistry, page 192. In the first volume of the Medical Observations of London, is a remarkable cafe of fcorbutic ulcers, in the legs of an old clergyman, being removed by the use of lime-water, after almost every other application had been tried for feveral years without fuccefs.

Phagedenic water.

Take a pint of lime-water, and a fcruple of white corrofive fublimate; make a folution thereof. —This is a good lotion for old eating ulcers or venereal fores, it may be occafionally diluted by the addition of more lime-water, or mixed with a little fpirits of wine, as circumftances may indicate.

Vitriolio

Vitriolic water.

Diffolve two ounces of blue vitriol, in a quart of fpring water.—This is dabbed on fores to deftroy fungous flesh, and is generally preferred to harsher methods.

Bougie.

Take quickfilver, and lead, each two drachms; make them into an amalgama; then add of crude antimony in fine powder, and Venice turpentine, each three ounces; white wax, fixteen onuces.— Vide St. Bartholomew's Hofpital, page 24.

The hemlock pultice.

Take of the hemlock fomentation, a pint; and boil it with a fufficient quantity of oatmeal, to a proper confiftence for a pultice; then add two or three ounces of oil of olives.—Vide the hemlock f.mentation, page 82.

The common pultice.

Take of the common fomentation a pint; boil it with a fufficient quantity of oatmeal, till it is of a proper conditionce; then add fix ounces of fallad oil, and two ounces of rectified fpirits of wine.

The discutient pultice.

Take a pint of stale beer grounds, a fufficient quantity of oatmeal, and boil them to the confistence of a pultice; then add fix ounces of oil of olives.---This is infinitely preferable to any other application, in bruises, strains, mortifications, and all cases that require discutients.

The emollient pultice.

Take the crumb of ftale rolls, and, with a fufficient quantity of milk, boil it to a proper confiftence; then add fome fallad oil, to prevent its growing hard.

In using this pultice, care must be taken not to leave it on the part long enough to grow four; it should therefore be changed every night and morning, otherwise it will do more mischief than good.

The repellent pultice.

Boil a pint of the sharpest white wine vinegar, with a sufficient quantity of oatmeal to give it the proper confistence; then add fix ounces of oil of olives.

This, tho' fimple, is efficacious, and is the pultice ufually applied to diflocated limbs, to reduce the fwelling, and abate the inflammation.

The common cauftic is prepared as directed in the Supplement. Its use requires no explanation.

The lunar caustic.

Take of the chrystals of filver, one ounce; put them into a crucible which is capable of containing five or fix times their quantity, by reason they are apt to boil and swell: you must give a very gentle heat till the ebullition is over; then increase the fire a little till the filver finks and flows like oil, and cast it into a warm greased ingot mould. It must be kept in a glass well stopped from the air. See page 187

The yellow cerate.

Take a pint and an half of oil of olives, and a pound of yellow wax, and melt them together over a gentle fire.

The epulotic and mercurial cerates, are prepared as directed in the Supplement.

Plaisters, adhæsive, common, and mercurial, are prepared, according to the London Dispensatory, as directed in the Supplement.

Strengthning

Strengthning plaister.

Take five pounds of the common plaister, and a pound of colcothar of vitriol in fine powder, and mix them together with a small proportion of olive oil. This, tho' a simple composition, answers the intention well.

The common fomentation.

Take of wormwood, bay leaves, and rolemary, each half a pound; boil them in a fufficient quantity of water to strain off a gallon; fometimes spirits of wine, or spirits of wine camphorated, are added to this fomentation.

The hemlock fomentation.

Boil four ounces of dried hemlock, in a gallon and an half of water, till it comes to a gallon.---Hemlock has been greatly recommended by Dr. Storck of Vienna, for its virtues in cancerous cafes. This fomentation and the hemlock pultice are used externally, during the time the patient takes the extract of this plant inwardly.---The method of taking the extract of hemlock is to begin with taking only two grains night and morning, which may be gradually increased to the quantity of thirty grains in the twenty-four hours.

A fumigation of a fcruple of native cinnabar, repeated as occasion may require, is of fervice in ulcers of the throat, and other obstinate fores from a venereal cause.

The Ægyptian ointment of the London Difpenfatory, is used externally to keep down fungous flesh, to remove excressences, &c. as is also red precipitate, for the like purposes.

The angelic powder.

Take red precipitate and burnt alum, each equal parts; mix them into a fine powder.--- This is an excellent escharotic preparation.

Oils of Neats feet — Olives — Turpentine.

Tincture of myrrh.

Take myrrh in powder, two ounces; melasses fpirits, and rectified spirits of wine, each half a pint; digest them together.—This, as an external application, is used generally to carious bones, to promote exfoliation.

Detergent ointment.

Take hog's-lard, a pound; red precipitate, an ounce and an half, mix them for use

Another.

Take a pound of the yellow cerate, and mix therewith an ounce and an half of red precipitate in fine powder.—Either of these may be used to procure a good discharge from wounds which afford a thin fanious discharge, as they act without giving the least pain, and are used in all the hopitals for that purpose.

Digestive ointment.

Take yellow wax, rozin and turpentine, each five pounds; oil of olives, fix pounds; and melt them together.---Vide yellow bafilicon, in all the hofpitals.

Green digestive ointment.

Take a pound of the digeftive ointment, two ounces of verdigreafe in fine powder, and two ounces of oil of olives, mix them together.---This is a warm digeftive and good detergent, and is therefore used to cleanse old ulcers, and wear away fungous flesh.

Ointment

83

Ointment of gum elemi.

Take three pounds of mutton fuet, one pound and an half of gum elemi, one pound of Venice turpentine, and fix ounces of oil of olives; melt them together over a gentle fire.---This is the digeftive ufually made use of in wounds of the head and brain, being imagined to posses peculiar virtues in those cases.

The ointment of three ingredients, is prepared according to the London Difpenfatory, as directed in the Supplement.

VIEW MENTION DAY STRAND



GUY's

S. 2. 4. 1 2.

(85)

GUY'S HOSPITAL.

Purging water.

T AKE of Dog and Duck water three pints; powder of ginger, a drachm; boil them together, till one half is wafted away, and then add one ounce of manna.

This quantity may be taken two or three times in a week, and is extremely gentle in its operation.

Rose water, sapphire coloured water, and balfam of sulphur, are prepared as directed in the Supplement.

Aromatic bolus.

Take of toafted nutmeg a fcruple; of the fcordium electuary, two fcruples; and mix into a bolus, with a fufficient quantity of fyrup of white poppy heads.

This bolus is to be taken every night and morning, and is directed in old diarrhæas and flatulent diforders.

Balfamic bolus.

Take balm of gilead one fcruple; liquorice powder, and conferve of hips, each half a fcruple; and mix into a bolus with a fufficient quantity of the fimple fyrup.---This is recommended to be taken night and morning, with a draught of the pectoral decoction, page 90, in inward ulcerations, and diforders of the breaft.

Camphor bolus.

Take conferve of hips a fcruple; camphor, ten grains; mix them together.---To be repeated every eight hours.

This

This has been frequently found to promote a diaphorefis, in fevers of the low deprest kind, when every other medicine has been tried without fucces.

Cordial bolus with Venice treacle.

Take of the compound powder of crabs claws, Virginia fnake root, and contrayerva root, each ten grains; faffron, five grains; and mix into a bolus, with a fufficient quantity of fyrup of clove gilly flowers.

Cordial bolus with Venice treacle.

Take of the cordial powder, and Venice treacle, each one fcruple; mix into a bolus.

Either of these bolusses may be taken every fix hours, with a sew spoonfuls of the cordial julap after every dose, to promote perspiration.

- rool and lo Cephalic bolus. The salat

Take conferve of rofemary, a fcruple; powder of caftor, fifteen grains; and a fufficient quantity of the fimple fyrup.—This is a good medicine for the purpofes its title expresses, and is to be taken three times in a day, with a few spoonfuls of the hysteric julap, page 100.

Saffron bolus.

Take conferve of roles fifteen grains; faffron, one fcruple; mix them into a bolus with fimple fyrup: which may be taken every fix hours, to promote a diaphorefis.

Cinnabarine bolus.

Take conferve of orange peels, fifteen grains; cinnabar of antimony, half a drachm; mix them together.

This bolus is directed to be taken twice in a day in cephalic and nervous diforders, and in obstinate rheumatic complaints.

Bolus

86

2.11.2

Guy's Hospital.

Bolus for the piles.

Take of lenitive electuary two drachms; flowers of brimftone, two fcruples; mix them together. This, though fimple, is a very efficacious remedy, and is to be taken night and morning.

Epileptic bolus.

Take conferve of rofemary, half a drachm; flowers of steel, fifteen grains; mix them into a bolus:—Which may be given twice in a day, in epileptic diforders, with a cup-full of valerian infusion. See page 99.

Emetic bolus.

Take conferve of roses, two scruples; turbith mineral, eight grains; mix them into a bolus.

This operates very roughly, and is therefore feldom prescribed but in very robust constitutions.

Calomel bolus.

Take half a drachm of conferve of rofes, and fifteen grains of calomel; mix them into a bolus.

Mercurial bolus.

Take conferve of orange peels, a fcruple; alkaliz'd mercury, half a drachm; mix them together.—This is an excellent alterative, and may be taken twice in a week.

Guaiacum bolus.

Take conferve of orange peels, and gum guaiacum, of each fifteen grains; mix them into a bolus with the fimple fyrup. This may be taken night and morning, in all difeafes from obstructed perspiration.

Guaiacum bolus, with mithridate.

Take ten grains of gum guaiacum, and two fcruples of mithridate; mix them into a bolus with the fimple fyrup.--- fyrup.—This is to be taken at night going to reft, and is used for the fame purposes as the former.

Musk bolus.

Take a fcruple of conferve of rofemary, and ten grains of musk; mix them into a bolus.

This may be taken two or three times in a day, and has been frequently experienced of fervice, in many nervous and hifterical complaints, as well as in depress to promote a diaphorefis.

Myrrh bolus.

Take conferve of orange peels, and myrrh in powder, of each a fcruple; falt of wormwood, five grains; and make them into a bolus with the fimple fyrup.

This fhould be taken three times in a day, is good in hifterical affections, and is of manifeft fervice in ripening the fmall-pox, especially that fort where the pustules rife with a pellucid humour.

Pectoral bolus.

Take half a drachm of spermaceti, a scruple of tragacanth powder, and mix them into a bolus with the simple syrup, which may be taken three times in a day.

Astringent pectoral bolus.

Take of the fcordium electuary, and tragacanth powder, each half a drachm; and with fyrup of white poppy heads, make into a bolus, which may be taken twice in a day.

Purging bolus.

Take fifteen grains of gamboge, ten grains of cream of tartar, five grains of powdered ginger, and make into a bolus with the fimple fyrup.

In dropfical habits this is a very common, and a very good cathartic; for it thoroughly purges off watry

watry and viscous humours, and greatly promotes the discharge by urine: but to render the cure of a dropsy lasting, there is afterwards a necessifity of restoring and strengthning the tone of the parts, by proper medicines for such purposes. See page 11, 56, 92.

Tin bolus.

Take conferve of orange peels, a fcruple; of prepared tin, half a drachm; and with the fimple fyrup, make into a bolus, which may be taken twice in a day.

This is particularly deftructive to worms, and is with fome a fecret against that species called the joint-worm.

Bolus for the king's evil.

Take conferve of orange peels, and burnt fponge, of each a drachm; and make into a bolus with the fimple fyrup.—To be taken three times in a day.

This is given in all kinds of fcrophulous diforders, at the fame time with the water.

Turpentine bolus.

Take of clarified honey, and the ætherial oil of turpentine, each half a drachm: liquorice powder, as much as will give it the confiftence of a bolus.— This is to be taken twice in a day, and will effectually diflodge those viscidities which clog the joints, and occasion ischiatic pains, but its use requires caution.

Aluminous collyrium.

Diffolve a drachm of alum in fix ounces of rofewater.—This is applied to repel a defluction of fharp humours on the eyes.

Confection of alkermes, cordial confection, conferves of orange-peels, hips, rofemary, rofes, the white decoction, and clyfter decoctions, are prepared as directed in the Supplement.

Restringent

Restringent decostion.

Take of the Peruvian bark, one ounce; balaustine flowers, pomegranate shells, and tormentil roots, each half a drachm; bruise them, and boil in a sufficient quantity of water, to strain off a quart .- Four spoonfuls may be taken two or three times in a day, in those diforders which require the affistance of restringents.

Dietetic decostion.

Take of guaiacum fhavings, and bark, each a pound; liquorice root, half a pound; coriander feeds, two ounces; boil them in fix gallons of fpring water to four gallons, and while boiling add two ounces of mazerion roots:

Half a pint is given three times a day, as the ordinaty sweetner in any course of alteratives, for cutaneous eruptions, leprofies, and venereal diforders. It is very much like the diet drink with which Dr. Wall cured his venereal patients; the great fuccefs from it is supposed to arise from the mazerion root, which feems peculiarly adapted to diffolve nodous fwellings.

Febrifuge decoction.

Take of the Peruvian bark powdered, two ounces; boil it in three pints of water to one pint : strain and add a tea-cup full of brandy :- four large fpoonfuls of the liquor may be, taken three or four times in a day. A very good method of administering the Peruvian bark, in delicate conftitutions, where the ftomach is apt to recoil at the bark in fubstance.

Pectoral decoction.

Take of liquorice root bruifed, two ounces; boil it in a quart of barley-water, to a pint and an half .--- A quarter of a pint may be taken at any time as common drink, in all diforders of the breaft and

Decle theest

lungs.

TIN .

Decoction of elm bark. See Batholomew's Hospital, page 5.

Restringent electuary.

Take of the fcordium electuary, and dragon's blood, each fix drachms; alum, Japan earth, and cinnamon, each two drachms and an half; make into an electuary with the fimple fyrup.

The quantity of a nutmeg is directed to be taken two or three times in a day, in the most obstinate fluxes of the belly, after proper doses of rhubarb have been administred.

Lenitive electuary.

The dose is a quarter of an ounce as often as occasion may require.

Alterative electuary.

Take of lenitive electuary, one ounce; gum guaiacum, and Æthiop's mineral, each half an ounce; mix into an electuary with the fimple fyrup.—The dofe is a drachm twice in a day, drinking after it the dietetic decoction, or fome other fuitably medicated liquor.—It is a most efficacion prefeription, in all foulness of blood whatsoever.

Chalybeate electuary.

Take of the flomachic electuary, two cunces and an half; prepared fleel, half an ounce; make into an electuary: The dose is a drachm twice in a day, and is an excellent medicine in a cachectic habit of body, and suppressions in the female sex.

Bark electuary, with alum.

Take one ounce of the Peruvian bark in powder, a quarter of an ounce of alum, and make into an electuary with the fimple fyrup:—The dofe is the quantity of a nutmeg every three or four hours.

Scammony electuary

Is prepared as directed in the Supplement :- The dofe is a drachm night and morning, twice in a week, in rheumatic cafes.

Barbadoes tar electuary.

Take of virgin honey, four ounces; gum olibanum, elecampane, and Barbadoes tar, each twoounces; make into an electuary with the fimple fyrup.—A quarter of an ounce may be taken twice in a day; and where the ftomach will difpense with it, will do great fervice in obstinate tickling coughs, and will succeed where more elegant medicines have failed.

Scorbutic electuary.

Take of preferved aron root, and Winter's bark in powder, each one ounce; mix them into an electuary with the fimple fyrup: The dofe is a drachm twice in a day.

Stomachic electuary.

Take of fweet case, zedoary, galangals, and orange peel in powder, each half an ounce; fimple fyrup, as much as will make an electuary:— The dofe is a drachm and an half twice in a day. It is a very good medicine in cholics, and all complaints arifing from indigeftion, and a cold weak ftomach.

Elixirs, of aloes, stomachic, and of vitriol, are prepared as in the Supplement.—Forty drops of each may be taken twice in a day.

Mynsichts elixir of vitrol.

Take cinnamon, ginger, and cloves, of each three drachms; calamus aromaticus, one ounce; galangals, one ounce and an half; fage and mint dried, each half an ounce; cubebs and nutmeg, each two ounces; aloes wood, and citron peel, each

each a drachm; powder them together, and add to them, fugar candy, three ounces; fpirits of wine, a pint and an half; and a pint of oil of vitriol: digeft together thirty days, and then pour off the liquor and filter it for ufe.—The dofe, from ten to forty drops in any fuitable vehicle, two or three times in a day, obferving to take it when the ftomach is most empty. It is taken to promote an appetite, strengthen digestion, and in relaxed habits; being much preferable to the elixir of vitriol of the shops.

Strengthning plaister.

Take of diachylon plaister, five pounds; Burgundy pitch, and colcothar of vitriol, finely powdered, each half a pound; olive oil, two ounces; melt them together over a flow fire, flirring the whole till the ingredients are entirely mixed.—A good application for strained limbs.

Anodine plaister.

Take cummin-feed plaister, and yellow wax, each one ounce; opium, and camphor, each two drachms; mix them together.—This is calculated to give ease in arthritic pains, and all such as arise from acrimonious humours, whether from venereal or scorbutic causes.

Ischiatic plaister.

Take Burgundy pitch, four ounces; euphorbium in powder, a drachm and an half; and make into a plaister with a sufficient quantity of Venice turpentine. Vide, St. Thomas's Hospital, page 51.

Volatile plaister.

Take of volatile fal-armoniac, a drachm; camphor, two drachms; and mix into an emplaister, with Venice turpentine. A warm discutient.

93

Restingent clyster.

Take of the reftringent decoction without the Peruvian bark, four ounces; of the fcordium electuary, half an ounce; mix them together.

Restringent clyster, with opium.

Take of the reftringent clyfter, fix ounces and an half; and diffolve in it two grains of pure opium.— Either of these may be used in obstinate fluxes of the belly.

Common clyfter.

Take of the clyfter decoction, ten ounces; fweet oil, three ounces; fyrup of buckthorn, one ounce; mix them together.

Cholic clyfter.

Take of the clyfter decoction, ten ounces; fweet oil, three ounces; Glauber's falt, one ounce; and powder of hiera picra, a drachm; mix for use.

Oleaginous clyster.

Take of the clyfter decoction, and fweet oil, each fix ounces; mix them for use.

Turpentine clyfter.

Take of the clyfter decoction, ten ounces; and mix therewith half an ounce of turpentine, incorporated with the yolk of an egg; and half an ounce of lenitive electuary.—This is the clyfter generally ufed in fits of the itone and gravel.

Restringent fomentation.

Take of oak bark, two pounds; of balaustine flowers, half a pound; boil them in three pints of water, till one half is wasted away; then diffolve in the remaining liquor fix ounces of common alum. This is the usual fomentation after cutting for the store, and it will greatly aftringe the fibres, not only

I

only fo as to reduce a relaxed part to its proper tone, but also leffen any aperture of the veffels, made by incifion, in fuch a manner as to haften their reunion. It ought to be used as hot as possible without scalding.

Common fomentation.

Take of wormwood, bay, and rolemary leaves, each four handfuls; boil them in a fufficient quantity of water to firain off a gallon, to which add a quart of malt or melaffes fpirits.

Poppy fomentation.

Take two ounces of white poppy heads, and boil them in a fufficient quantity of water to firain off a quart, to which add an ounce and an half of the beft vinegar.—This is ufed in inflammations of the eyes, and in wounds attended with great pain.

Alexipharmic gargle.

Take of the pectoral decoction, and alexipharmic tincture, equal parts; and mix them together. —This gargle does wonderful fervice in the putrid fore throat.

Common gargle.

Take barley-water, one pint; honey of rofes, three ounces; and make the mixture agreeably tart with fpirits of vitriol.—This is the beft gargle that can be used for a cooler and detergent. Where the mouth has white specks, as is common in young children, it may be made more sour, if necessary, and rubbed hard upon those places, with a rag tied on a spoon, or skewer.

Volatile gargle.

Take of the paralytic infusion, a pint; and mix with it an ounce of spirits of fal-armoniac.—This is very pungent and spirituous, and well contrived to warm and ftimulate the fibres which have loft their tone in paralytic cafes.

Anodine draught.

Take of London philonium, thirty grains; fimple mint-water, ten drachms; Jamaica pepper water, a quarter of an ounce: mix them together for one dofe.—This is a warm opiate, and is prefcribed in diforders of the bowels, to be repeated at difcretion.

Diuretic draught.

Take of the ftomachic draught, two ounces; and diffolve in it two drachms of the diuretic falt.—One of these draughts may be given night and morning in dropfical habits, during the use of other medicines calculated for that diforder.

Emetic draught.

Take powder of ipecacoanha, fifteen grains; oxymel of fquills, fix drachms; and fimple alexiterial water, one ounce; mix them for one dofe.— A fafe and efficacious emetic, if warm water is drank plentifully during its operation.

Epileptic draught.

Take a drachm and an half of valerian finely powdered; loaf fugar, a drachm; and one ounce and an half of fimple alexiterial water; mix them together.---One of these draughts may be taken three times in a day, in nervous, paralytic, or epileptic cases.

Nephritic draught.

Take half an ounce of Venice turpentine, and incorporate it with honey; to which add three drachms of Jamaica pepper-water, and an ounce and an half of fimple mint-water.---One of thefe draughts fhould be taken every night and morning.

Oily

96

Oily draught.

Take of cold drawn linfeed oil, half an ounce; fimple alexiterial water, one ounce; fimple fyrup, two drachms; fpirit of fal armoniac, ten drops; mix them into a draught—To be taken every night and morning.

These are in great esteem, and much used in common practice, not only in inward ulcerations and distempers of the breast, but also in obstructions of the urinary passages, and some cholicky disorders.

Purging oily draught.

Take oil of fweet almonds and fimple alexiterial water, each one ounce; Jamaica pepper water, three drachms; Epfom falt, one ounce; mix into a draught to be repeated as occasion requires.

Olibanum draught.

Take powder of gum olibanum, half a drachm; mix it with a little honey; then add fyrup of balfam, and Jamaica pepper water, each two drachms; fimple alexiterial water, one ounce; mix together for one dofe—To be taken twice in a day.

This is ufually perfcribed for the whites, and to remove weakneffes of the urinary parts.

Purging Draught.

Take three ounces of the infusion of fenna; half an ounce of Jamaica pepper water, and one ounce of fyrup of buckthorn; mix together, to be taken three times in a week, or as often as occasion may require.—A fafe cathartic in all cafes where purging is required.

Refrigerant draught.

Take nitre and loaf fugar, each two fcruples; fimple alexiterial water, one ounce and an half; nutmeg water, two drachms.—One of these draughts is to be taken every fix hours.

Saponaceous draught.

Take Venice foap, one drachm; fimple alexiterial water, two ounces; lavender drops, one drachm; fimple fyrup, threedrachms; mix into a draught.— To be taken twice in a day, in the jaundice, ftone, or gravel.

Stomachic draught.

Take falt of tartar, one scruple; lemon juice, half an ounce; simple mint water, an ounce; Jamaica pepper water, half an ounce; mix into a draught—To be taken three times in a day in naufeas, &c.

Nervous draught.

Take Ruffian caftor in powder, one drachm; oil of amber, twelve drops; fal volatile, twenty drops; fimple cinnamon water, two ounces; fyrup of cloves; two drachms; mix them into a draught.—If taken twice in a day, it is of amazing efficacy against the diforders its title express.

Sulphurated draught.

Take of fimple alexiterial water, two ounces; fulphurated water, half an ounce; fimple fyrup, two drachms; mix into a draught—To be taken three times in a day.

This is prescribed with fuccess, in fits of the convulsive althma.

Volatile draught.

Take volatile fal armoniac, fifteen grains; fimple alexiterial water, two ounces; and two drachms

of the fimple fyrup; mix into a draught.—To be taken every fix hours, to promote a diaphorefis in low nervous fevers, or in the rheumatic fever.

Chamomile infusion.

Take an handful of chamomile flowers; infufe them in a fufficient quantity of boiling water to ftrain off a quart; to which add a quarter of a pint of malt or melaffes fpirits.

The dose is a quarter of a pint twice in a day, to create an appetite and promote digestion.

Paralytic infusion.

Take horfe-rhadifh, and muftard-feed bruifed, each three ounces; infufe them in a quart of boiling water; when cold ftrain off the liquor, and add a quarter of a pint of malt, or melaffes fpirits.— Three ounces may be taken two or three times in a day.,

Infusion of valerian.

Take one ounce of bruifed valerian root, infuse it in twelve ounces of boiling water; when cold Arain off the liquor, and add fix drachms of compound lavender drops.

Three ounces may be taken two or three times in a day in nervous and paralytic diforders.

Cordial julap.

Take half a pint of the fimple alexiterial water; two ounces of Jamaica pepper water; and fix drachms of fyrup of clove-gilly-flowers; mix them into a julap.

Common julap.

Take of fimple alexiterial water, eight ounces; treacle water, two ounces; of the fimple fyrup, half an ounce; and mix them into a julap.

F 2

Hyfteri

Hysteric julap.

Take half a pint of fimple penny royal water, two ounces of fpirituous penny royal-water, half an ounce of the fimple fyrup, and mix them into a julap.

Mint julap.

Take half a pint of fimple mint water, two ounces of fpirit of mint, and half an ounce of the fimple fyrup; mix them into a julap. Vide St. Thomas's hofpital, page 59.

Styptic julap.

Take half a pint of fpring water, two ounces of the flyptic tincture, and half an ounce of fyrup of quinces; mix them into a julap.

Three or four spoonfuls of these julaps, are prefcribed to wash down medicines in a more folid form, calculated for the purposes their respective titles express.

Two large fpoonfuls of ammoniacum milk, prepared as directed in the Supplement, are directed to be taken in the afthma, whenever the fhortnefs of breath is troublefome, and in all cafes, where expectoration is wanted.

Restringent milk.

Take oak bark, pomegranate shells, and tormentil roots, bruised, of each two ounces; boil them in a pint of milk, and a pint of water, till one half is wasted away; towards the end add two ounces of cinnamon, and strain off the liquor when cold.

Two ounces may be taken frequently. This is prefcribed in those diarrhæas which accompany the last stage of confumptive diforders.

The white mercurial liniment.

Take a quarter of a pound of hogs-lard, and two

two drachms of white precipitate; mix them together.

This is a very neat and efficacious ointment for the itch, and all outward foulneffes of the fkin; but the body must be kept open during the use of it; left, if the patient gets cold, a falivation might unexpectedly ensue.

Volatile liniment.

Take half a pint of neat's foot oil, two ounces of fpirits of wine and camphor, and two ounces of fpirits of fal armoniac; mix them together.

This is a very powerful difcutient, and is directed to be rubbed in upon any induration, as by its warmth and fubtility, it attenuates the flagnant matter, and renders it fit to be taken up again by the abforbing veffels, or helps to breath it out by the pores of the fkin.

Sulphur lotion.

Take half a pound of flower of brimftone; of any fixt alkaline falt, two ounces; boil them in fix pints of water, to four pints; when cold strain the liquor off for use.—For its use see St. Thomas's Hospital. page 62.

Alcalized mercury.

Take four ounces of quickfilver; and fix ounces of crabs eyes, finely levigated; rub them together in a marble mortar till the quickfilver entirely difappears, and the powder is changed to a grey colour.—From five grains to two foruples may be given in any vehicle twice in a day.

This is a fafe medicine, and may be given to young children for the worms, and all crudities and acrimony of the humours.

Mixture against deafness.

Take one ounce and an half of oil of bitter F 3 almonds almonds; and half a drachm of spirit of fal-armoniac; mix them together.

A few drops are to be poured into each ear every night, ftopping them afterwards with cotton or black wool.

Pectoral mixture.

Take cold drawn linfeed oil, and fyrup of balfam, each two ounces: mix them together.—Two large fpoonfuls to be taken three times in a day.

Pestoral mixture with diacodium.

Take linfeed oil and fyrup of white poppy heads, each two ounces; mix them together:—The dofe is the fame as the former. Both are excellent medicines in pleurifies, inflammations of the lungs, coughs, and all diforders of the breaft.

Afthmatic mizture.

Take fimple mint water, four ounces; Jamaica pepper water, and oxymel of fquills, each two ounces; mix them together.—Two fpoonfuls are to be taken frequently. This is a powerful expectorant.

Restringent mixture.

Take fimple alexiterial water, fix ounces; Jamaica pepper water, two ounces; of the fcordium electuary, half an ounce; mix them together— Take two large fpoonfuls after every loofe ftool.

Alterative pills.

Take one drachm of calomel, two drachms of the fcordium electuary, and, with a fufficient quantity of liquorice powder, make the mais into ixty pills—One of which may be taken every night and morning. These are used in venereal, fcorbutic, and leprous diforders.

el anticia de la las des bas opares sao sela la Ammoniacum

102

Guy's Hospital. Ammoniacum pills.

Take a scruple of gum ammoniacum, and fix grains of the ftorax pill; mix them together for one dofe.—To be taken every night going to reft. This is a very good pectoral medicine.

Pills for a cough.

Take Rufus's pill, fifteen grains; of the faponaceous pill, eight grains; divide into three pills, which are to be taken at night going to reft.—This is well contrived to ftop the defluxion of rheum which continually provokes coughing, and to divert it by a proper difcharge.

Strengthning pills.

Take rhubarb in powder, one ounce; and makeit into fifty pills, with a fufficient quantity of Venice turpentine:—The dofe is half a drachm twice in a day. Thefe are directed with fuccefs in the fluor albus, and old gleets, free from virulence.

Ecphractic pills.

Take hiera picra, gum ammoniacum, and prepared steel, each half an ounce; make into pills with the simple syrup.—Half a drachm divided into sour pills, may be taken twice in a day.

This composition is a powerful deobstruent, and is therefore of use in almost all-chronic difordersfrom obstructions. They keep the body moderately open, and will do great service in hypochondriac affections.

Quicksilver pills ...

Take quickfilver killed with Venice turpentine, two drachms; fcammony, jalap, and gamboge, two drachms each; make into a mass with syrup of buckthorn.—The dose is half a drachm every morning. It is certainly a good medicine in venereal, scrophulous, and leprous cases.

Hysteric pills.

Take galbanum and gum ammoniacum, half an ounce of each; make into a mafs for pills, with the fimple fyrup:---The dole is a fcruple, twice in a day.---This pill is excellently contrived to answer the purposes of an hysteric, being as effectual as it is fimple and easy of preparation.

Stomachic pills.

Take Virginia fnake root and hiera picra, each two drachms; extract of gentian, half an ounce; make into pills, with the white fyrup.—The dofe is half a drachm, night and morning. These continued for some time together, warm a cold weak ftomach, that is filled with crudities and flatulencies, and by degrees give new strength to the fibres, and procure a good digestion.

White potion.

Take balfam of capivi diffolved with the yolk of an egg, half a drachm; two drachms of the fimple fyrup, and one ounce and an half of fimple alexiterial water; mix them together—To be taken twice in a day. Vide St. Thomas's Hofpital, page 67.

Alexiterial powder.

Take of the compound powder of crabs claws, one fcruple; Virginia fnake root, four grains; contrayerva, fix grains; mix for one dofe.—To be repeated every fix hours, with a few fpoonfuls of the cordial julap, to promote a diaphorefis in fevers of the low depreft kind.

Antimonial powder.

Take crude antimony in fine powder, two fcruples; Æthiop's mineral, one fcruple; mix for one dofe.-

304

dofe.—To be taken twice in a day. This is an admirable remedy in all cutaneous foulneffes; as fcabs, itch, herpes, leprofy, and the like, efpecially if the patient be first bled, and purged with calomel, drinking at the fame time the dietetic decoction, and anointing the eruptions with the itch ointment.

The powder for the bite of a mad dog is prepared as directed in the Supplement, and used according to the directions given by Dr. Mead for its adminiftration.

Worm powder.

Take prepared coral, and scammony, finely powdered, each fix drachms; rozin of jalap, one drachm; mix together.—Fiscen grains arethe proper dose, two or three times in week.

This purges very brifkly, and is extremely effectual against worms, especially in young persons.

The opthalmic powder is white vitriol very finely powdered.

Common purging powder.

Take half a drachm of powder of jalap, twice in a week, in any liquid.

Purging mercurial powder.

Take rhubarb, one fcruple; calomel, ten grains; mix them together.—To be taken twice in a week.

Tin powder ..

Take of prepared tin, one ounce and an half; wormfeed, half an ounce; mix them together.--The proper dofe is from ten grains to half a drachm night and morning. See the Tin Bolus, page 89.

Compound lavender drops, spirits of nitre, of fal armoniac, of wine camphorated, of vitriol, of fal volatile; Syrups, of white poppy heads, of buckthorn, and simple, are prepared as directed in the Supplement.

Syrup of cloves.

Take one pound of cloves, and one ounce of cochineal, bruife them and infufe them, in a veffel clofe ftopt, in two gallons and a quart of boiling water, for twenty-four hours; then add twenty eight pounds of loaf fugar, and boil to a proper confiftence.—This is used as a substitute for the fyrup of clove gilly flowers.

Alexipharmic tincture.

Take of contrayerva and inake root bruiled, each two drachms; cochineal, a fcruple; infufe them in a fufficient quantity of boiling water to ftrain off twelve ounces; to which add four ounces of Jamaica pepper water.—Four ipoonfuls may be taken every fix hours, or oftener, according to the exigence of the cafe, in the beginning of acute diftempers. It will, with proper warm diluters, raife a diaphorefis fooner than almost any other memedicine.

Tinctures, bitter, aromatic, of antimony, of cantharides, of cardamoms, of caftor, of faffron, of affafœtida, of guaiacum volatile, of fteel with spirit of salt, of myrrh, of sena, of roses, of lead, of fnake-root, styptic, and thebaic, are prepared as directed in the Supplement.

Tincture to provoke the menses.

Take black hellebore, großsly powdered, four ounces; rectified spirits of wine, twelve ounces; digest them together for three weeks.

This is an admirable medicine for many purpofes, but particularly it excels in removing uterine obstructions; and in fanguine conflications, where steel is hurtful, this never fails of forcing the menstrual discharge.—From twenty to fixty drops may be taken twice in a day in any fuitable vehicle.

Tincture of Japan earth.

Take of Japan earth, one ounce; Peruvian bark, and balaustines, each half an ounce; bruise them, and infuse in half a pint of rectified spirits of wine, and four ounces of spirituous cinnamon water.

This is of good fervice in all fluxes of the belly, dyfenteries, and immoderate flux of the menfes, and even in a gonorrhæa, and old gleets, where the virulence has been already removed :--Its dofe is from a tea fpoonful to two fpoonfuls, in a glafs of red port wine, two or three times in a day.

Ticture of rhubarb.

Take rhubarb, großiy powdered, four ounces and an half; coriander or cardamon feeds, two drachms; infuse in a quart of water and a pint of melasses spirits.

This is a mighty mild and innocent tincture of rhubarb, and may with the utmost fafety be given to young children without danger of inflaming their tender viscera, or raising a fever, by the burning spirit, with which tincture of rhubarb is commonly extracted.

Yellow basilicon.

Take yellow rozin, and linfeed oil cold drawn, each five pounds; yellow wax, two pounds and an half; common turpentine, ten ounces; diffolve the other ingredients in the oil, ftir them well together, and then ftrain the whole for an ointment.—This is the common digeftive in all green wounds, &c.

Ointment for a perpetual blifter.

Take half a pound of yellow bafilicon, and mix with it one ounce of Spanish flies in fine powder. This ointment seems principally intended for dreffing blifters, in order keep them running during F 6 pleasure, pleasure, as may be commodiously affected, by fpreading a little thereof upon a piece of linen and applying it occasionally to the part.

Emollient ointment.

Take four ounces of ointment of marshmallows, and mix with it one ounce of oil of turpentine. Vide St. Thomas's hospital, page 77.

Mercurial ointment.

Take fix ounces of quickfilver, and extinguish in an ounce of Venice turpentine, then mix with it a quarter of a pound of hog's-lard. See page 77.

Ditto camphorated.

Is made by adding half an ounce of camphor, to two ounces of the mercurial ointment.

Opthalmic liniment.

Take four ounces and an half of hog's-lard; bees-wax and prepared tutty, each half an ounce; camphor, two drachms; mix them together, and make thereof a liniment.

For fimplicity and excellence few opthalmic ointments can compare with this.

Itch ointment.

Take two pounds of hog's-lard, and four ounces of common brimftone, mix them together into an ointment. See St. George's hofpital, page 78.

Refrigerent ointment.

Take fweet oil and bees-wax, each four ounces; fpermaceti, one ounce and an half; and make them into an ointment according to the rules of art. This is used in the eryfipelas, and to prevent pits in the face from the small-pox.

Pale coloured mercurial ointment.

Take hog's-lard, one pound; and two ounces of mercury sublimate; mix them together into an ointment.

Ointment for a scald head.

Take one pound of tar, four ounces of common brimstone, and one ounce of yellow wax; mix together into an ointment.

During the use of this ointment, proper alteratives should be taken internally, and the ohild's head should be covered with an hog's bladder.

Green ointment.

Take half a pound of yellow basilicon, and a drachm of verdigrease; mix them together into an ointment. See St. George's hospital, page 134.



ALL STREET STREET, STR

St. GEORGE's HOSPITAL.

Camphorated Water.

A K E Roman vitriol, and Armenian bole, each four ounces; camphor, one ounce; and powder them together: of this mixture fprinkle one ounce at a time, into two quarts of boiling water, in which flir it well about; then take the liquor off the fire, and let it fettle.

This is an excellent medicine for many purpofes externally; it cleanfes ulcers, by washing them frequently with it warm; it is good against all inflammations; and almost infallibly cures fore eyes; if it is too sharp it may be diluted with a little spring water; and in the itch it is both fase and efficacious.

Shell lime water.

Take calcined oyfter-fhells a pound, and pour on them a gallon of boiling water; the water fhould ftand four hours or longer on the fhells, and it fhould be made in an earthern veffel.

This, according to Dr. Whytt, poffess the whole lithontriptic power of Mrs. Stephens's medicines.—From three pints to two quarts must be taken every day, and continued till the cure is effected.

The difagreeable tafte of the lime-water may be mitigated by adding a very finall quantity of new milk to each dofe; and if it occasions costiveness, it will be necessary now and then to take an ounce of manna diffolved in whey, or barley-water; the patient ought to drink no more of any liquor than is sufficient to quench his thirst, and he should retain

St. George's Hospital.

retain his urine as long as he can without uneafinefs, that it may have the longer time to act on the ftone.

Mercurial bolus.

Take half a drachm of conferve of rofes, and half a fcruple of quickfilver; mix them well together; rub till the mercurial globules are extingushed. To be taken every night going to rest.—This is used as an alterative in cases which require the affistance of mercury.

Rhubarb bolus.

Take fifteen grains of torrified rhubarb, and one fcruple of the fcordium electuary; mix them together for one dofe: to be repeated as occasion may require, in diforders of the bowels.

Discutient cataplasm.

Take of stale beer grounds a quart; oatmeal, as much as is fufficient to give a proper confistence.—This cataplasm cannot be too much recommended in violent bruises and even in beginning mortifications. See page 80.

Medicated Ale.

Take fcurvy-grafs, water creffes, and brook lime, each three handfuls; broom afhes, a pound; horferhadifh, and muftard feed bruifed, each three ounces; juniper berrries, two ounces; filings of fteel, two pounds; infuse them in old ftrong beer for four days, then decant the liquor off for use.—This is a good antifcorbutic, and in dropfical and cachectic habits, will be of great fervice if continued for fome time.

Vitriolic collyrium.

Diffolve fifteen grains of white vitriol in two ounces and an half of fpring water.—Of great efficacy in removing fpecks and films on the eyes. Decolion

Decoction of gum arabic.

Diffolve half an ounce of gum arabic, in a quart of barley-water. This is given for common drink in the ftrangury arifing from the application of blifters.

Scarlet decoction.

Take nitre one ounce; loaf fugar, two ounces; and cochineal, fixteen grains; boil them in a quart of water till the nitre is intirely diffolved; then ftrain off the liquor for use.

Two ounces may be taken two or three times in a day, to promote urine, and cool the urinary paffages in gonorrhæas, ftranguries, &c.

Decoction of chalk.

Take two ounces of prepared chalk; and half an ounce of gum arabic; boil them in a fufficient quantity of water to strain off a gallon.—A good restringent in diarrhæas, after a few doses of rhubarb, or an emetic of ipecacoanha; and may be drank at pleasure.

Compound hartshorn drink.

Take fix drachms of burnt hartfhorn, three drachms of prepared crabs eyes; comfrey, and tormentil roots, each two drachms; boil them in three pints of water to a quart; to which add one ounce of fimple pepper-mint water, and half an ounce of fyrup of white poppy heads.

Either this or the former may be taken as common drink in diarrhæas, and all acute difeafes attended with a loofenefs, obferving the foregoing caution.

Bran water.

Take four handfuls of bran, and boil in fix quarts of water to four quarts; then strain off the liquor, and add a quarter of a pound of the best honey.—

St. George's Hofpital.

honey.—This is used as common drink in coughs and diforders of the breast.

Strong guaiacum decostion.

Take guaiacum shavings eight pounds, and macerate them in fixteen gallons of warm water for fix hours; then boil for fix hours, and strain off for use ten gallons.

The weak decoction of ditto is made by adding a pint of warm water to every quart of the ftrong decoction.—This decoction is of fervice in venereal and fcorbutic complaints, drank plentifully.

Bark decostion.

Take of the peruvian bark großly powdered, one ounce; water, three pints; boil them together over a flow fire for three hours; then increase the fire, and boil for three or four hours longer, fo as to pour off clear about one pint, to which add two ounces of melasses fpirits.—The dose is a tea-cupfull every two hours, in intermitting fevers. Vide Guy's hospital, page 109.

Decoction of elm bark.

Take of the inner bark of the elm one ounce, and boil it in a quart of water to a pint. Then strain off the liquor for use.

This makes an admirable gargle in ulcerations of the mouth, and is fuppofed to cure cutaneous eruptions, and even the leprofy itfelf. See page 5.

Wormwood electuary.

Take conferve of wormwood, a pound ; winter's bark in powder, half a pound; and mix into an electuary, with fyrup of ginger.—The dofe is the quantity of a nutmeg, two or three times in a day, in all complaints arifing from indigeftion.

Acid

Acid electuary.

Take conferve of roles, and ftoned raifins, each three ounces; oil of vitriol, half a'drachm; and mix into an electuary with two ounces of the fimple fyrup.

Pectoral electuary.

Take conferve of hips, one ounce and an half; fpermaceti, three drachms; prepared crabs eyes, one drachm; mix into an electuary with the fimple fyrup.

Elecampane electuary.

Take elecampane, and Spanish liquorice, diffolved in water, and gum ammoniacum disfolved in vinegar, each an equal quantity; and mix into an electuary, with fyrup of garlick.

The quantity of a nutmeg of either of these electuaries, may be taken to promote expectoration, three or four times in a day.

Chalybeate electuary.

Take conferve of wormwood, and prepared fteel; each two ounces; powder of ginger, half an ounce; and mix into an electuary with the fimple fyrup.— The dofe is the quantity of a nutmeg twice in a day; ufing exercise greatly affists the efficacy of this medicine, in obstructions and other diforders incident to women.

Guaiacum electuary.

Take conferve of wormwood, cinnabar of antimony, and gum guaiacum, each an equal quantity, and mix into an electuary with the fimple fyrup.— The dofe is the quantity of a nutmeg twice in a day, and is furprizingly efficacious in removing rheumatic diforders.

Compound

St. George's Hofpital.

Compound lenitive electuary.

Take half a pound of lenitive electuary, one ounce of powder of jalap, a quarter of an ounce of falt prunella, and mix into an electuary with the fimple fyrup.

This purges very gently, and is generally given in the morning, the quantity of a nutmeg, to work off any mercurials which may have been taken over night.

Bark electuary.

Take one ounce of the Peruvian bark in powder, and mix it into an electuary with the fimple fyrup.—The dofe is the quantity of a large nutmeg three or four times a day. See St. Thomas's hofpital, page 44.

Valerian electuary.

Take two ounces of powder of valerian, and one ounce of cinnabar of antimony; mix into an electuary with fyrup of ginger.—The dole is the quantity of a large nutmeg three or four times in a day. An efficacious medicine in nervous complaints. See page 66 and 86.

Bate's pepper electuary.

Take long-pepper, one ounce and an half; rue, half an ounce; fal-gemm, five drachms; and make into an electuary with twelve ounces of treacle.— A warm ftomachic, and of great fervice in flatulencies proceeding from a cold weak ftomach.

Mustard electuary.

Take whole muftard feeds, and conferve of wormwood, each equal parts; and mix into an electuary with fyrup of ginger.

I his is very efficacious in stimulating the fibres and discussing viscidities, and is therefore excellent

in

116

in paralytic, dropfical, and rheumatic diforders.— The dofe is the quantity of a large nutmeg, three or four times in a day.

Balfamic electuary.

Take spermaceti, and conferve of hips, each four ounces; prepared oyster shells, one ounce; balfam of Peru, fix drachms; and make into an electuary with syrup of white poppy heads.—The dose is the quantity of a large nutmeg, three or four times in a day.

Care must be taken during the use of balfamic medicines, that they do not pall and relax the stomach too much, which they are apt to do; for which reason they are often acidulated with spirits of vitriol, or such like things, which help to keep the fibres of the stomach in their due tension.

Elixir of vitriol.

Take Winter's bark, orange peels, and fweet cane, großly powdered, each an ounce; cloves, and Jamaica pepper, each half an ounce; melaffes spirits, five pints; digest them together for four days; then strain off the tincture, and pour on it, by degrees, a pint and four ounces of oil of vitriol.

For the dofe and virtues, see Guy's hospital, page 92.

Anodyne embrocation.

Take spirits of fal armoniac, and liquid laudanum, each half an ounce; spirits of wine camphorated, three ounces; mix them together for use. —This is very penetrating, and in rheumatic and arthritic swellings, will do great service, by relaxing the irritated fibres, at the fame time that it promotes the perspiration of the obstructed parts.

St. George's Hospital.

Strengthning plaister.

Take of diachylon plaister, one pound; melt it over a flow fire, and ftir into it three ounces of armenian bole in fine powder.

Galbanum plaister.

Take ftrained galbanum, half a pound ; frankincenfe, camphor, and oil of amber, each half an ounce; mix them together.—This is intended to promote fuppuration, for which purpofe it is very efficacious.

Plaister for the scrophula.

Take of the mercurial plaister, and the hemlock plaister, of the Edinburgh Dispensatory, each four ounces; camphor, three drachms; mix them together.

Blister plaister.

Take two ounces of Burgundy pitch, fourteen ounces of melilot plaister, and half a pound of Spanish flies in fine powder; mix them together.

Common glyfter.

Infuse an handful of chamomile flowers, in two ounces of boiling bran water; when cold, strain off the liquor for use.

Oily glyAer.

Add two ounces of fweet oil to the common glyfter.

Purging glyster

Is made by adding to the oily glyfter, an ounce and an half of fyrup of buckthorn, and half an ounce of common falt.

Starch glyfter.

Diffolve a quarter of an ounce of extract of logwood, in fix ounces of starch jelly.

117

Anodyne flarch glyfler

Is made by adding a drachm of London philenium to the ftarch glyfter.—Thefe two laft are excellent glyfters in the bloody flux.

Restringent glyster.

Take of the chalk decoction, fix ounces; and diffolve in it half an ounce of the fcordium electuary.

This is generally used in obstinate diarrhæas, after the use of rhubarb or an emetic of ipecacoanha.

Turpentine glyfter.

Take fix drachms of turp entine diffolved in the yolk of an egg, and mix it with ten ounces of the oily glyfter. See Guy's hofpital, page 94.

Emollient fomentation.

Take of the dried leaves of mallows, and marfhmallows, with the flowers, each an handful; boil them flightly in a quart of water, and strain off the liquor for use.

Discutient fomentation.

Take wormwood and chamomile-flowers, each one ounce; rofemary and bay-leaves, each half an ounce; boil them flightly in fix pounds of water, and ftrain off the liquor for use.

Compound fomentation

Is made by adding a quart of vinegar, and a pint of rectified spirit, to two quarts of the discutient fomentation.

Volatile fomentation.

Sprinkle a flannel wetted in the discutient fomentation, with spirits of fal armoniac, immediately before you apply it to the part affected.

ntiseptic

St. George's Hospital.

Antiseptic gargle.

Take barley-water, a pint; white wine vinegar, two ounces; tincture of myrrh, half an ounce; and two drachms of the aromatic tincture; mix together.—This is the gargle ufually prefcribed in the putrid fore throat, in which it is of infinite fervice.

Detergent gargle.

Take half a pint of bran-tea; and half an ounce of honey of roles; mix them together.—This is frequently directed for ulcers, and inflammations of the mouth, &c.

Sometimes two fcruples of oil of vitriol are added to this gargle, to render it ftill more deterigent.

Volatile gargle.

Take half a pint of bran-tea, four ounces of melaffes fpirits, and one drachm of fpirit of fal armoniac; mix them together.

Palfey gargle.

Boil one ounce of pellitory of Spain, in a fufficient quantity of water, to ftrain off a pint; to which add half an ounce of fpirit of fal armoniac.— Thefe are both excellent in the palfy of the tongue.

Cordial draught.

Take one ounce and an half of fimple alexiterial water, half an ounce of compound juniper water, and two drachms of the fimple fyrup.

Anodyne draught.

Take one ounce and an half of the cordial draught, and twenty drops of the thebaic tincture; mix them together.—To be taken at night going to reft.

Camphor ated

Camphorated draught.

Take ten grains of camphor; honey, one fcruple; fimple penny-royal-water, two ounces; mix together.—These draughts may be repeated every fix hours; and when a diaphores is to be encouraged, nothing will sooner raise one, provided the patient drinks plentifully of small diluting liquors.

Mithridate draught.

Take of the cordial draught, two ounces; mithridate, one fcruple; mix them together for one dofe, to be repeated every fix hours.

Valerian draught.

Take two ounces of the cordial draught, and half a drachm of powder of valerian; mix them for one dofe, to be repeated every fix hours.

Sudorific draught.

Take mithridate, one fcruple; the beft white wine vinegar, half an ounce; fimple alexiterial water, one ounce; nutmeg-water, two drachms; fimple fyrup one drachm; mix together for one dofe, which is to be repeated as often as may judged neceffary.

Oily draught.

Take fix drachms of oil of almonds, one ounce of fimple penny-royal-watert nutmeg water, and fyrup of marfhmallows, each two drachms; mix together.—One of these draughts may be taken three or four times in a day.

Oily draught with rhubarb.

Take one ounce and an half of tincure of rhubarb; half an ounce of oil of almonds; two drachms of fyrup of white poppy-heads; and fifteen drops of the thebaic tincture; mix together.

St. George's Hofpital.

together. — In dyfenteries this is a most excellent medicine, fince at the fame time, it both aftringes, composes, and helps forward the evacuation of such humours as vellicate the bowels, abrade their mucus, and lay open the capillaries in such a manner as to cause the blood to flow at every stool.

Common purging draught.

Take three ounces of the infusion of sena; one ounce of fyrup of buckthorn, and three drachms of compound juniper water; mix them together.

Saline draught.

Take half a drachm of fait of wormwood, thirty drops of spirit of vitriol, one ounce and an half of simple mint water, two drachms of treacle water, and one drachm of the simple syrup. See page 15.

Diaphoretic draught.

Take two ounces of the faline draught, and a fcruple of the diaphoretic powder; mix them together.

Diaphoretic draught with mithridate.

Take two ounces of the faline draught and one fcrup'e of mithridate; mix them together.

Saline draught, with oxymel of squills.

Take two ounces of the faline draught, and two drachms of oxymel of fquills mix them together.

Ditto with valerian.

Take two ounces of the faline draught, and a fcruple of valerian; mix them together.

Ditto with rhubarb.

Take two ounces of the faline draught, and ten grains of rhubarb; mix them together.

Saponaceous

Saponaceous draught.

Take Castile soap, one drachm; diffolve it in two ounces and an half of warm water; and add to it compound juniper water, and syrup of marshmallows, of each a quarter of an ounce; mix together. See page 3, 44.

Epileptic draught.

Take two ounces of the faline draught, falt of amber, and Ruffia caftor, each ten grains; mix together.

Squill draught.

Take vinegar of squills, one drachm; compound juniper water, three drachms; fyrup of marshmallows, one drachm; and ten drachms of springwater; mix together. An efficacious diuretic in dropfical diforders.

Volatile draught.

Take ten grains of falt of hartfhorn, two ounces of penny-royal-water, and a drachm of the fimple fyrup; mix them together.

The oleaginous draught.

Take ten grains of falt of hartfhorn, ten drachms of penny-royal water, half an ounce of oil of almonds, and two drachms of fyrup of marfhmallows; mix them together. -- These draughts are to be taken every fix hours, in those disorders for which they are respectively calculated.

Bitter Infusion.

Take gentian root fliced, and dried orange-peel, each one ounce and an half; infuse them in five quarts of boiling water; when cold, strain off the liquor, and add to it one pint of melasses spirit...-An excellent stomachic bitter.

Garlick

St. George's Hofpital.

Garlick infusion.

Take elecampane, and garlick fliced, each one ounce; liquorice root, three drachms; anifeeds and carraway-feeds, half an ounce each; infufe in fix pints of boiling water, and when cold strain off the liquor for use.

This powerfully deterges the lungs, whence it gives great relief in afthmas, and difficulties of breathing.

Horse-radish infusion.

Take horfe-radifh, and muftard feeds bruifed, each two ounces; dried orange-peels, one ounce; winter's bark, half an ounce; infuse in a quart of water over a very flow fire; when cold strain off the liquor, and add to it a quarter of a pint of melass spirits:—The dose of these infusions is from a quarter, to half a pint, every night and morning. —This is ferviceable in paralytic and rheumatic cafes.

Arabic injection.

Diffolve three, drachms of gum arabic, in a pint of barley-water; and add one ounce and an half of honey of rofes.

Gamphorated injection.

Take of the camphorated water, page 110, a pint; Ægyptian honey, three drachms; and mix them together.

Terebinthinate injection.

Take one pint of barley-water, and half an ounce of turpentine, diffolved in the yolk of an egg; mix them together.

Acid julap.

Take a pint of spring water; four ounces of treacle water, and one ounce and an half of syrup of vinegar; mix them together.

Chalk julap

Is made by adding five drops of oil of carraways, mixt with a lump of fugar, to a quart of the chalk julap as prepared in the Supplement. Three or four table fpoonfuls may be taken at diferention.

Penny-royal Julap.

Take a pint of fimple penny-royal water; four ounces of ftrong penny-royal water; and an ounce of loaf fugar: mix them together.—These are used to wash down other medicines of greater efficacy.

Common linetus.

Take three ounces of fweet oil; fyrup of white poppy heads and marfhmallows, each an ounce and an half; loaf-fugar, as much as is fufficient to give it a proper confiftence.—A tea-spoonful is to be frequently taken in coughs, &c.

Bituminous liniment.

Take Barbadoes tar, and oil of tartar, each equal parts.—This liniment is of furprizing fervice in white iwellings.

Campborated liniment.

Take-half a pound of rofemary leaves, and infufe them three or four days in two gallons of rectified spirits of wine.

Take a quart of this tincture, a quarter of a pound of Caftile foap, and two ounces of camphor: digest them together with a gentle heat.

Oily liniment.

Take a pint of neat's-foot oil, two ounces of oil of turpentine, and three ounces of Caftile foap; mix them together.—Thefe are both excellent difcutient.

Ammoniacum

St. George's Hospital.

Ammoniacum mixture.

Take ten ounces of ammoniacum milk, and ten drachms of oxymel of fquills; mix them together.—Three or four table fpointuls, are to be taken whenever the cough or fhortness of breath is troublesome.

Restringent mixture.

Take fix ounces of fimple alexiterial water; one ounce of nutmeg water, and three drachms of the foordium electuary; mix them together.—Three large fpoonfuls are to be taken after every loofe ftool; rhubarb or ipecacoanha having been previoufly administred.

Laxative mixture.

Take three ounces of fena; cream of tartar, and carraway feeds, each three quarters of an ounce; infuse them in a quart of boiling water; strain off the liquor when cold, and add four ounces of manna, and fix ounces of tincture of sena.—A large cup full of this mixture is to be taken at discretion.

Spermaceti mixture.

Take two drachms of spermaceti, diffolved in the yolk of an egg; fix ounces of alexiterial water, one ounce of nutmeg water, and a drachm and an half of loaf-sugar; mix together.

Compound spermaceti mixture

Is made by adding one ounce of fyrup of white poppy heads, and forty drops of balfam of Peru, to feven ounces of the fpermaceti mixture.—Thefe are preferrised in coughs, ulcerations of the lungs, and other diforders of the breaft.

Volatile Spermaceti mixture.

Take two drachms of spermaceti, diffolved in the yo'k of an egg; eight ounces of fimple alexiterial G 3 water, water, two foruples of falt of hartfhorn, and half an ounce of loaf-fugar; mix them together.

Calomel pills.

Take five grains of calomel, and mix into a pill with conferve of rofes. This is generally prefcribed to be taken over night, and worked off with the common purging draught the next morning.

Chalybeate pills.

Take prepared steel, one ounce ; extract of gentian, as much as is fufficient to make it into a mass; which is to divided into forty-eight pills. Two pills are to be taken night and morning, in obstructions of the menses.

Affafætida pills.

Take affafætida, camphor, and myrrh, each a drachm; Caftile foap, two drachms; make into a mass with tincture of wood-foot.—A good nervous and antihisteric medicine. Two middle fizes pills may be taken three times in a day.

Mercurial pills.

Take three ounces of quickfilver, one ounce of powder of jallap, two drachms of powdered ginger; and make them into a mais, with fyrup of marshmallows. The dose, from ten grains to a scruple, night and morning.

Pectoral pills.

Take gum ammoniacum, one ounce and an half; elecampane powder, and balfam of fulphur, half an ounce each; make into a mass with fyrup of marshmallows.

Three moderate fized pills, taken two or three times in a day, will do great fervice in afthmas, wheezing coughs, and all infarctions of the lungs.

Sagapenum pills.

Take sagapenum, one ounce ; powder of valerian, two ounces; Castile soap, two drachms; make

.St. George's Hospital.

into a mafs with elixir of aloes.--- Two middle fized pills may be taken two or three times in a day. This pill is excellently contrived to answer the purpose of an antihysteric, in all distempers of the head, epilepsies, hypochor driacal, hysteric, and nervous cases; it is as effectual, as it is simple and easy of preparation.

Saponaceous pill.

Take Caftile foap, half an ounce; gum guaiacum, four fcruples; and make them into a mafswith the fimple fyrup. — Two middle fized pills may be taken, night and morning. These much promote infensible perspiration, and are good in such cutaneous diforders as proceed from an obstruction of the perspirable matter, and very much relieve rheumatic and arthritic pains.

Squill pills ...

Take four scruples of fresh squills; two scruplesof powder of ginger; make them into a mass with fyrup of garlic, and divide into twenty-four pills: two of which, taken three times a day, as they cause a large expectoration, cannot but be extremely ferviceable in phthisical and afthmatic cases.

Turpentine pills.

Take two ounces of liquorice powder; make into a mafs with Venice turpentine; and divide the mafs into pills of a moderate fize: two or three of which, taken night and morning, are very ufeful in weakneffes of either fex.

Aluminous powder.

Take crude alum, and japan earth, each one ounce; and reduce them together into a fine powder.---This is a very powerful aftringent, and is used with success in old fluxes of the belly, after the use of rhubarb or an emetic of ipecacoanha.

Dr. Dover's powder.

Take vitriolated tartar, and mitre, each four ounces; powder them together and put them into a red hot crucible; keep the mixture flirring about till the red fumes ceafe to rife: then powder it and add one cunce of fliced opium; then powder again, and add one cunce of liquorice, and as much ipecacoanha finely powdered; mix them well together.— Its proper dofe is from twenty to forty grains, in any liquid, going to bed between the blankets, and drinking plentifully now and then fomething warm, fuch as white-wine whey, balm tea, the faline mixture, &c. The fweating is not to be checked, but encouraged by lying ftill and keeping warm.

Chamomile powder.

Take chamomile flowers, three drachms; myrth, and alum, each one drachm; beat all together into a powder.

This is sometimes used in intermitting fevers, instead of the Peruvian bark.—The dose a scruple, or half a drachm, every fix hours.

Cochineal powder.

Take loaf-fugar and nitre, each four ounces; cochineal, one drachm; beat all together into powder.

Worm powder.

Take fea mols, and worm feed, each one ounce; powder them together.--- This is an effectual medicine in worm cafes where purging is not required, or after it has been fruitlefsly tried : a fcruple may be taken, night and morning.

Diaphoretie powder.

Take four ounces of the compound contrayerva powder, and one ounce of faffron; mix them together.---This is given to promote a difcharge by the fkin.--The dofe from fifteen grains to half a drachm, every fix hours.

+

St. George's Hospital.

Purging powder.

Take fix drachms of jalap, and two drachms of ginger powder; mix them together.

Purging powder with nitre.

Mix two drachms of nitre, with an ounce of the purging powder.---The dofe of either of these powders, is from half a drachm to two scruples, three times in a week, in all cases where purging is required.

Nitrous powder.

Take nitre, one ounce; Rhenish tartar, and brimstone, each two drachms; beat them together into a powder. A cooling diuretic.

Sal-prunell powder.

Take sal-prunell, fix drachms; oyster-shells, prepared, two drachms; mix them together.

Compound fal-prunell powder.

Mix two drachms of the diaphoretic powder, with one ounce of the fal-prunell powder.

Plummer's alterative powder.

Take calomel, and golden fulphur of antimony, each equal parts; rub them together in a marble mortar, till the powder is changed to a dufky brown colour: feven grains and an half may be taken twice in a day, in cutaneous and leprous diforders, with the dietetic decoction. See page 90, 113.

Rhubarb powder.

Take rhubarb, in fine powder, half an ounce; of the diaphoretic powder, two drachms; mix them together.

Red powder.

Take loaf-fugar, spermaceti, and nitre, each half a jound; cochineal, two drachms; mix them into G 5 a powder. a powder.--- This is generally prefcribed in pleuritic diforders, and those of the urinary paffages.

Sponge powder.

Take burnt sponge, half an ounce; nitre, two drachms; mix them together. See Guy's hospital, page 89.

Tin powder.

Take three ounces of prepared tin, and one ounce of fea mofs; mix them together.---An efficacious medicine in diforders proceeding from worms.

Brimstone powder.

Take flour of brimstone, one ounce and an half; oyster shell powder, two drachms; cochineal, two scruples; mix them together.

Alkaliz'd mercury.

Take quickfilver, four ounces; prepared chalk, fix ounces; rub them together in a marble mortar till the quickfilver entirely difappears, and the powder looks of a greyifh colour.---This is a good alterative for children, whofe ftomachs generally abound with acidities. Its dofe from five grains to a fcruple twice a day.

Mindererus's spirit.

Take diftilled vinegar, two ounces; drop into it by degrees fpirits of fal armoniac, till the effervefcence entirely ceafes. When intended to promote a diarefis, half an ounce may be given twice in a day, mixt with an equal quantity of fyrup of marfhmallows, and it will feldom fail; but if you defign to provoke plentiful fweating, as in rheumatic difeafes, &c. half an ounce must be taken every two or three hours, till the fweat breaks out; repeating the dofe as occafion requires, when warm diluting liquors are not fufficient to keep the fweating up. Syrup

St. George's Hospital.

Syrup of vinegar.

Take five pounds of loaf-fugar, and diffolve over r a gentle fire in two pints of vinegar.

Cerated glass of antimony.

Take glass of antimony in powder, one ounce ; : bees-wax, one drachm; melt the wax in an iron ladle, then add the powder; fet them on a flow fire without flame, for the space of half an hour, continually flirring them with a spatula; then take the mixture from the fire, pour it upon a piece of clean white paper, powder it and keep it for use.—The dose is from fix grains to a scruple; according to the ftrength of the patient.

This medicine has been used with great fuccess for the dysentery, and the preparation of it kept as a secret for many years.

Antimonial tincture.

Take fix drachms of antimonial wine, and two drachms of liquid laudanum; mix them together. From twenty drops to a drachm, may be taken as. an alterative twice a day.

Golden tincture.

Take two ounces of elixir of aloes and fix drachms of tincture of fteel in fpirit of falt; mix them together.---The dofe, from twenty to forty drops, two or three times in a day.

This is a good ftomachic, and is of fervice in the jaundice, green-ficknefs, and obstructions of the menstrual discharge.

Tincture of wood foot.

Take affafœtida, and volatile fal armoniac, each one ounce; wood foot, two ounces; digeft together with a gentle heat, in a quart of rectified fpirit.

From

132

From twenty to forty drops may be taken, two or three times in a day, in a glass of water, in all nervous and hysteric cases.

Tincture of rhubarb.

Take rhubarb grossly powdered, three ounces; coriander seeds, and winter's bark, each two drachms; digest them with a gentle heat, in a quart of melasses spirits, and a pint of water. Vide St. Thomas's hospital, page 74.

Tinsture of Sena.

Take raifins ftoned, and fena, each eight ounces; carraway-feeds bruifed, one ounce and an half; digeft them for three weeks, in two quarts of melaffes fpitits, without heat; then ftrain off the tincture for ufe.—This is ufually known by the name of Daffy's Elixir, and is a warm carminative purge.

Tincture of hiera picra.

Take one ounce and an half of the fpecies hiera picra of the old London Dispensatory; and digest with a gentle heat, in one pint of melasses spirits, and half a pint of water.

This is prescribed from two to three ounces, either over night, or early in the morning, as a stomachic, or a preparative to alterative courses, to cleanse the first passages, and enable the stomach the better to digest what it receives.

Yellow bafilicon.

Take of oil olive, fixteen ounces; bees-wax, and yellow rozin, each fourteen ounces, melt them together over a gentle fire; and while hot ftrain the mixture for use. Vide Guy's Hospital, page 107.

St. George's Hospital.

Tar ointment.

Take Barbadoes tar, one ounce; fulphur vivum, three ounces; flour of brimftone, and yellow wax, each three drachms; oil of olives as much as is required to give the whole the confiftence of a foft ointment.

This is in mighty esteem for curing scald heads in children. See page 109.

Sir Hans Sloane's opthalmic liniment.

Take prepared tutty, one ounce; prepared blood-stone, two scruples; aloes, in fine powder, twelve grains; mix them well together in a marble mortar, with as much prepared viper's fat as is requisite to bring the whole to the confistence of a fost ointment.

This liniment is to be applied with an hair pencil, the eyes winking or a little opened; it has cured many whole eyes were covered with opake films, and cicatrices, left by inflammations and abfceffes of the cornea. It is to be obferved, that cathartics, especially with the addition of mercury, are prejudicial to the discases of the eyes, which are cured by this medicine.

Turner's cerate.

Take oil of olives, eight pints; hog's-lard, and yellow wax, each feven pounds; prepared calamine, five pounds and an half; melt the wax and hog'slard with the oil; and when the mixture begins to congeal, forinkle in the calamine, and flir the mixture well till it is quite cold.—This ointment is particularly ferviceable in burns and fcalds.

Itch ointment.

Take fulphur vivum, half a pound; hog's-lard, one pound; oil of tartar, one ounce; mix together for use.

Vitrialis

Vitriolic ointment.

Take half a pound of hog's-lard; two ounces of flour of brimstone; and one ounce of white vitriol, in powder; mix them into an ointment.

Either of these are very safe and very efficacious in removing the most obstinate itchy, or scorbutic : eruptions.

Yellow Mercurial Ointment. .

Take an ounce of quickfilver, and diffolve it in as much fpirit of nitre as will ferve for that purpofe; then add by degrees a pound of melted hog's-lard or butter; and keep the mixture flirring till it is quite cold.—For the purpofes of a detergent this. feems to be a fine contrivance.

Perpetual bliftering ointment.

Take yellow basilicon, half a pound; Spanish flies in fine powder, one ounce; mix them well together. See Guy's Hospital, page 107, 108.

Green ointment.

Take half a pound of yellow bafilicon; prepared verdigrease, a drachm; and mix them together for use. See page 109.

This is a good detergent, and is used to cleanse Id foul ulcers, and wear away fungous flesh.

THE

THE

LOCK HOSPITAL,

Near HYDE-PARK CORNER.

The brown collyrium.

T A K E of depurated white vitriol, half a drachm; boiling water, fix ounces; mix and make into a collyrium.

In all inflammations of the eyes, this cooling walk may be used at difcretion.

The white collyrium.

Take of the white troches of Rhafi's, one drachm; ten grains of camphor, diffolved in a little rectified spirits of wine; and spring-water, fix ounces; mix and make into a collyrium: this may be made use of in any defluxions of the organs of fight, as it fafely cools and repels those sharp humours which affect the eyes, and will clear them of beginning films and specks.

The collyrium with falt of tartar and lime-water.

Take of fimple lime-water, half a pint; falt of tartar, one drachm. This is likewife useful in most diforders of the eyes, particularly in specks and films thereof.

A decostion

A decostion against vomitting and fevers.

Take fix drachms of falt of wormwood, boil in three pints of fpring-water, ftrain the liquor and drop into it gradually of fpirit of vitriol, as much as will fufficiently impregnate it: then add three ounces of ftrong alixiterial water, and the like quantity of refined fugar.

Fevers, from too lax a ftate of the fibres of the ftomach, are often attended with nauseas, and frequent reachings to vomit. In this case, three spoonfuls given now and then, particularly after every return of that complaint, feldom fail to give relief.

Bark decoction.

Take Peruvian bark bruised, two drachms; nitre, half an ounce; boil the liquor in three pints of water to half the quantity: strain it off for use.

This form is contrived for those whose stomachs, perhaps, would be apt to nauseate the bark in subitance. Two or three ounces are to be taken every fix hours, as occasion offers, in agues, or intermitting fevers, and a general relaxation of the solids.

Arabic decoction.

Take gum arabic, one ounce; boil it in two pints of barley-water: when ftrained, add nutmegwater, three ounces; refined fugar, a fufficient quantity. In acidities of the ftomach, in a ftrangury, arifing either from blifters, or any other cause, this decoction is a foft, eafy, yet effectual remedy. A draught of it is to be drank at discretion.

Aftringent decoction.

Take of the arabic decoction, two pints and an half; boil in it tormentil root bruised, half an ounce; then add to the strained liquor, conferve of red roses, three ounces.

In

The Lock Hospital.

In all kinds of fevers, the small-pox, the measles, and acute difeases, where the intestines, from to violent an irritation of their fibres, by acid, or acrid humours, are two much follicited to an evacuation of the ferous matter of the blood by this canal; or are in a state of too great relaxation this astringent decoction, taken by way of common drink, will be found endued with falutary qualities, equal perhaps to those of more pompous compositions.

The astringent electuary.

Take compound powder of bole (either with or without opium, as you find occasion) for which see the London Pharmacopœia; compound species of tragacanth, of the London Pharmacopœia, of each two ounces; simple syrup, a sufficient quantity to form an electuary.

This is recommended in violent loofeneffes. Half a drachm or two fcruples of it, at a dole, generally abates those immoderate evacuations of the lower belly; and repeated, between whiles, for a day or two, will give an effectual check to the diforder.

Bleeding, a vomit, and gentle purging with rhubarb, ought to precede the use of it.

An electuary for a virulent gonorrhæa.

Take lenitive electuary, three ounces; powder of jalap, one ounce; nitre half an ounce; fyrup of roles, a sufficient quantity to make into an electuary.

A violent inflammation and tenfion in the urinary parts and paffages, being ever attendant on a virulent gonorrhœa, nothing can be better adapted to the urgency of these symptoms than this electuary; which, by its cooling, diuretic, and laxative properties, is every way qualified to afford relief. The dose is a drachm, to be taken night and morning. Should Should this quantity give more than three ftools, the dose must be lessened : if not fo many, increased.

An electuary for a gonorrhæa, after the inflammation.

Take a pound of lenitive electuary; balfam of copaiva, half a pound; powder of rhubarb, gum guaiacum, and nitre, of each four ounces; make into an electuary.

At this ftage of the diftemper, the heat and inflammation of the parts being over, this medicine is of very great use.

To be taken, a drachm night and morning.

A Arengthning electuary.

Take conferve of roles, one pound; gum olibanum in powder, half a pound; myrrh in powder, two ounces; fimple fyrup, a fufficient quantity to make into an electuary. The dole is two fcruples, three times a day.

This is prefcribed in diforders of the female fex, fuccefsfully.

An alterative electuary.

Take conferve of roles, four ounces; cinnabar of antimony prepared, three ounces; gum guaiacum in powder, two ounces; ginger in powder, two drachms; fimple fyrup, a fufficient quantity to make into an electuary.

As cinnabar, and gum guaiacum, are powerful attenuants, and plentifully promote perfpiration, this electuary is very rationally directed in rheumatic pains.

The dofe is a drachm night and morning.

A mercurial electuary.

Take quickfilver, one ounce; conferve of roses, three ounces; make into an electuary.

In all foulnefs of blood, productive of cutaneous eruptions, fuch as the itch, tetters, &c. this is a very efficacious remedy.

The Lock Hospital.

It may be taken twice a day, to the quantity of an ordinary nutmeg, at each dose.

A Romach electuary.

Take conferve of orange-peel, and fea wormwood, each two ounces; gum guaiacum in powder, winter's bark in powder, nutmegs, cinnamon, and carraway-feeds, powdered, of each two drachms; fyrup of ginger, a fufficient quantity to make an electuary.

This is a very warm and ftrengthning electuary to the ftomach, and is of very great fervice in flatulencies arifing from indigestion.

The dose is the quantity of a nutmeg, three times a day.

Blister plaister.

Take Burgundy pitch, eight ounces; Venice turpentine, and catharides, in powder, of each two ounces and an half; mix and make into a plaifter.

Blifters are useful in nervous fevers, and other cases where the pulse is low and languid, and the spirits depressed.

Common glyfter.

Take decoction of bran, twelve ounces; lenitive electuary, one ounce; common falt, three drachms; olive-oil, one ounce; make into a glyfter.

The use of this glyster, is well known to every one, therefore, need not be enlarged upon.

A glyster for a dysentery.

Take chicken broth, fix ounces; Venice turpentine, diffolved, three drachms; make into a glyfter. This glyfter is to be given, immediately after every ftool, in a dyfentery; and as occasion may require, add thebaic extract, two grains.

ar Masthuo wa

Cammon

Gonmon gargle.

Take tincture of roles, fix ounces; honey of roles, two ounces; fpirit of vitriol, thirty drops: mix and make into a gargle.

This gargle is of a cool and deterging kind. It is used, to cleanse and scour the glands of the mouth, from the phlegmatic matter, that stuffs and swells them, and may be used at discretion.

Common julap.

Take spring-water, twelve ounces; strong alexiterial-water, three ounces; simple syrup, half an ounce; make into a julap.

The intention of this julap, is as a vehicle for boles, powders, &c.

A restringent Milk.

Take of pomegranate bark, balaustin flowers, and tormentil-roots, of each half an ounce; cinnamon, two drachms; brune, and boil them in three pints of water, to two pints; towards the end of boiling, add a quart of milk; and conferve of roses, three ounces; then strain it.

Two ounces of this milk may be taken, three times a day, in every cafe where aftringents are proper.

A Mercurial lotion.

Take corrofive sublimate mercury, two drachms; fpring-water, one pint; boil together till the mercury is diffolved.

This is calculated for the itch; the parts lightly to be washed with it two or three times a day.

An astringent mixture.

Take of the astringent electuary, half an ounce; spring-water, fix ounces; strong cinnamon-water, two ounces; make into a mixture.

This

The Lock Hofpital.

This is of excellent fervice in all fluxes whatfoever, and a great strengthner of the stomach and bowels.

Its dose is a spoonful or two after every stool.

Saline mixture.

Take falt of wormwood, one drachm; fpirits of vitriol, eighty drops; fpring-water, fix ounces; fpirituous mint-water, two ounces; fyrup of balfam, an ounce and a half; mix and make into a mixture. Three fpoonfuls of this mixture to be taken every three hours, in inflammatory fevers, and naufeas of the ftomach.

White mixture.

Take spermaceti in powder, three drachms; olive-oil, half an ounce; yolk of egg, a sufficient quantity; spring-water, fix ounces; strong cinnamon-water, two ounces; make into a mixture.

This mixture is peculiarly calculated for freeing the lungs from that oppreffion which is effentially confequent to an inflammation of them.

The dose two spoonfuls, to be taken when the cough is troublesome.

The red pill.

Take red precipitate deflagrated, one grain; conferve of roles, a fufficient quantity; mix and make a pill, to be taken morning and evening.

This is an excellent alterative, in cutaneous and venereal difeases.

Specific pills.

Take calcined mercury, one fcruple; red coral prepared, two fcruples; thebaic extract, ten grains; conferve of rofes, a fufficient quantity. Mix and form the mass into twenty pills. These pills likewise are capable of doing great fervice in venereal and chronic diforders, and are at present supposed to be the most effectual mercurial medicine.

The dose is from one pill to four in a day, encreasing the number by degrees.

A purging draught.

Take infusion of sena, two ounces and an half; solutive syrup of roses, tinclure of sena, and Glauber's salt, of each two drachms; mix and make a draught.

In all cafes, where moderate purging is neceffary, this potion will be very properly prefcribed.

A diaphoretic powder.

Take compound powder of contrayerva, and nitre, of each a fcruple; mix and make a powder.

This is given to promote a discharge by the skin, every four orfix hours.

The nitrous powder.

Take purified nitre, and fine fugar, of each half a drachm; mix and make a powder.

Pectoral Syrup.

Take green liquorice, one pound; cochineal, half an ounce; boil in a fufficient quantity of water, to ftrain off two gallons; then add of fine fugar, twenty-five pounds; and boil it to the confiftence of a fyrup.

This is very properly contrived for all diforders of the breaft.

Troches of red lead.

Take crumbs of white bread, four ounces; corrofive fublimate mercury, one ounce; powder of red lead, half an ounce; rofe-water, a fufficient quantity to make into troches.

These are contrived for external use only, and are powerful eschartics.

The Lock Hospital.

An emollient ointment, with mercury.

Take ointment of marshmallows, two ounces; elder ointment, quickfilver, one ounce of each; mix and make an ointment.

A digestive ointment.

Take yellow wax, twelve ounces ; yellow rozin, frankincense, of each one pound ; hog's-lard, two pounds; mix and make into an ointment.

The green digestive ointment.

Take of arcæus's liniment, three ounces; balfam of copaiva, one ounce; green ointment of elder, half an ounce; mix and make into an ointment. A most excellent digestive.

A mercurial digestive ointment.

Take the digeftive ointment, and crude mercury, of each two ounces; mix and make into an ointment.

A very efficacious application in venereal ulcers, where the use of digestives is required.

Brown ointment.

Take olive-oil, two ounces; white wax, frefh butter, mutton-fuet, and litharge of gold, of each three ounces; boil the litharge with the oil till it becomes brown, after that add the other ingredients.

THE

PORTUGUESE HOSPITAL.

The diuretic and refrigerant apozem.

A K E of grass-roots, two ounces; boil in three pints of barley-water, till one third is wasted away; add a few minutes before it is removed from the fire, an handful of forrel leaves; one ounce and an half of tamarinds; and two drachms of nitre. The dose, a quarter of a pint whenever agreeable to the patient. Its virtues are fufficiently expressed by its title.

The aluminous bolus.

Take of alum, fifteen grains; nutmeg, and extract of the Peruvian bark, of each half a foruple; fimple fyrup, a fufficient quantity to make a bolus. This botus may be taken every fix hours, where the ftomach will bear fo large a dofe of alum; in which case it will warm and invigorate a cold relaxed ftomach; and in old agues, which have eluded the force of the bark alone, will feldom fail to effect a cure, if taken an hour before the access of the fit.

The Portuguese Hospitai.

The folutive anodyne bolus.

Take of aloes, and the London philonium, each one fcruple; mix them together for one dofe. This bolus may be taken going to reft, or at any other time when required. It is excellent in cholicky diforders.

The borax bolus.

Take of borax, myrrh, and nutmeg, in powder, each half a fcruple; and make into a bolus, with fyrup of orange-peels. This medicine may be repeated occafionally, and has long been held in efteem, as an effectual remedy to promote childbirth.

The camphor bolus.

Take of gum arabic, half a drachm; camphor, one fcruple; and mix into a bolus with fyrup of marfhmallows. For its virtues, and dofe, fee the fame article in the foregoing hospital dispensatories.

The chalk bolus.

Take of prepared chalk, a foruple; nutmeg, half a foruple; and mix into a bolus with the fimple fyrup. The dofe may be repeated every fix hours. An excellent abforbent and a mild reftringent, in diarrhœas; but its most experienced virtue, is in removing that uneafiness at the stomach, commonly called the heart-burn.

The emetic bolus.

Take of yellow emeric mercury, in fine powder, fix grains; and mix into a bolus with conferve of rofes. A rough and violent emetic, feldom ufed but to procure a ftrong revultion from the genital parts, in an hernia humouralis, phymofis or paraphymofis, when from the vaft fluxion of humours, it is feared a mortification will enfue.

The anodyne fætid bolus.

Take of afafœtida, one scruple; the confection called paulina, fifteen grains; mix into a bolus. It is of confiderable efficacy in flatulent cholics, and hifteric complaints; and may be repeated according to the urgency of the symptoms.

The gamboge bolus.

Take gamboge, and cream of tartar, of each a fcruple; and mix together into a bolus with fyrup of buckthorn. A powerful hydragogue, and therefore good in dropfical diforders; but its roughnefs requires the utmost caution in its exhibition.

The ipecacoanha bolus.

Take of toasted ipecacoanha, a scruple; winter's bark in powder, ten grains; and mix into a bolus with syrup of orange-peels. An useful reftringent in diarrhœas and disenterics. To be repeated as occasion requires.

The myrrh bolus with seel.

Take of compound powder of myrrh, half a fcruple; flowers of steel, and extract of favine, of each a scruple; mix into a bolus with syrup of orange-peel: to be taken night and morning. A powerful emmenagogue.

The musk bolus.

Take musk, and zedoary-root in powder, of each fifteen grains; and mix into a bolus with fyrup of fugar: if necessary, add a grain of thebaic extract. This bolus without the opium, may be taken every for hours; it never fails producing a mild diaphorefis. without heating; and has extraordinary effects in convulsive diforders, even when attended with the worft fymptoms.

The

The Portuguese Hospital.

The olibanum bolus.

Take olibanum, and spermaceti, of each fifteen grains; nitre, ten grains; mix into a bolus with syrup of maidenhair. For the virtues of this medicine refer to pages, 15, 30, 63, 97.

The pepper bolus.

Take of long pepper, a fcruple; London philonium, half a fcruple; mix into a bolus with fyrup of fugar. It gives great relief in the flatulent cholic.

The rhubarb bolus.

Take of rhubarb powdered, half a drachm; London philonium, ten grains; mix into a bolus with the fimple fyrup. An elegant and efficacious remedy in all cholicky complaints.

The tin bolus.

Take of prepared tin, a drachm; mix into a bolus with conferve of orange-peel. For its victues &c. fee pages, 7, 43, 89, 130.

The contrayerva bolus.

Take of contrayerva-root in powder, half a drachm; nitre, fifteen grains; mix into a bolus with the fimple fyrup. This bolus is to be taken every four or fix hours; and is an useful diaphoretic.

The aftringent decoction.

Take of the Brazilian bark in powder, one ounce and an half; boil in three pints of water to a quart, then strain off the liquor for use. A quarter of a pint may be drank, two or three times in a day.

147

The bark decoction.

Take two ounces of powdered Jesuits bark, and boil in three pints of water, to a quart; then strain off the remainder for use.

The diuretic decoction.

Take of pareira brava root in powder, two ounces; water, two quarts; boil to three pints; add towards the end of the coction, two drachms of fennel-feeds bruifed; and firain off the liquor for ute. The dofe three or four ounces every four or fix hours. This is greatly extolled against fuppressions of urine and nephritic pains, and has been found useful with the addition of balfam copaivæ in healing ulcers of the kidnies and bladders. Its attenuating quality renders it also of fervice in humoural asthmas, and jaundices proceeding from a viscidity of the bile.

The sudorific decoction.

Take of the bark of guaiacum-wood, and faffafras-fhavings, each one ounce; water, two quarts; boil to three pints; add, a few minutes before the liquor is removed from the fire, half an ounce of ftoned raifins; and two drachms of liquorice-root: ftrain off the liquor for ufe.

The aromatic and anodyne electuary.

Take of the aromatic species of the London Difpenfatory, and pareira-brava root in powder, each fix drachms; London philonium, three drachms; mix into an electuary with syrup of orange-peel. The dose, the quantity of a nutmeg night and morning. For its victues see the above diuretic decodion.

The afringent electuary.

Take of the Brazilian bark in powder, an ounce. and an half; conferve of red rofes, an ounce; mix into

The Portuguese Hospital.

into an electuary with fyrup of orange-peels. The dofe, the quantity of a imall nutmeg every fix hours, observing the usual cautions.

The balfamic electuary.

Take of St. Thomas's balfam, first rubbed with loaf-fugar, and afterwards diffolved in the yolk of an egg, half an ounce; elecampane in powder, an ounce and an half; mix into an electuary with fyrup of maidenhair. The dose, the same as the former. A good detergent pectoral.

The chalybeate electuary.

Take of ruft of fteel, fix drachms; preferved ginger, an ounce; conferve of orange-peels, three ounces; mix into an electuary with fyrup of orangepeels. The quantity of a nutmeg to be taken two or three times in a day. This is adapted for those who labour under obstructions of the menses; it also gives relief in hypochondrincal diforders. Brisk exercise is of great advantage in the use of this medicine.

The cinnabarine electuary.

Take of cinnabar of antimony prepared, one ounce; gum guaiacum, three drachms; conferve of orange-peels, two ounces; mix into an electuary with fyrup of orange-peels. The quantity of a nutmeg to be taken night and morning. This is usually given in nervous and rheumatic diforders.

The bark electuary

Take of the Peruvian bark in powder, three ounces; the eleutherian bark, half an ounce; mix into an electuary with fyrup of orange-peels.

The guaiacum electuary.

Take of gum guaiacum in powder, compound powder of aron-root, of the London Dispensatory, I 3 and and winter's bark, each fix drachms; conferve of garden fourvy-grafs, two ounces; mix into an electuary with fyrup of orange-peels. The dofe, the quantity of a nutmeg twice in a day. This promotes greatly natural transpiration, and is therefore of tervice in cutaneous eruptions, and rheumatic pains.

The black bellebore electuary.

Take of black hellebore root in powder, extract of favine, and compound powder of myrrh, each half an ounce; winters-bark, two drachms; mix into an electuary with fyrup of orange-peels. The quantity of a nutmeg, may be taken night and morning. In fanguine conflictutions, it is preferable to freel; and powerfully breaks through uterine obftructions.

The faponaceous electuary.

Take of Castile soap, two ounces; pareira brava root in powder, one ounce; rhubarb, and aloes, each three drachms; mix into an electuary with syrup of orange-peels. The dose the quantity of a nutmeg twice or thrice in a day. This is particularly ealculated for the jaundice. See page 2, 44, 68, 122, 148.

The brimstone electuary.

Take of flowers of brimftone washed, half an ounce; lenitive electuary, two ounces; mix into an electuary with fyrup of marshmallows. The quantity of a nutmeg is to be taken discretionally. It does great service in the piles, especially where perfons are subject to be costive.

The anodyne discutient plaister.

Take of the cummin feed plaister, two ounces; camphor (rubbed down with a few drops of fweet oil) three drachms; extract of opium, a drachm and an half; mix together for a plaister. A warm discu-

The Portuguese Hospital.

tient, and is applied to ftrengthen the viscera, expel flatulencies, and appeale rheumatic and ischiatic pains.

The arabic emulsion.

Take of gum-arabic, one ounce; diffolve it in a pint of barley-water. This may be drank diferetionally. In heat of urine and ftranguries either from epifpaftics, or acrimonious humours, it gives immediate relief.

The balfamic emulfion.

Take of St. Thomas's balfam, three drachms; rub it well with loaf-fugar, and mix with the yolk of an egg; then add fyrup of maidenhair, an ounce and an half; barley water, a pint. The dofe, three or four table fpoonfuls, three times a day. Its title expresses its virtues.

The oily emulfion.

Take of fimple penny-royal water, twelve ounces; oil of olives, four ounces; fyrup of maidenhair, one ounce and an half; hartfhorn drops, two drachms; mix into an emulfion. The dofe, the fame as the former.

The common glyster.

Take of the common glyfter decoction, of the London Difpenfatory, twelve ounces; lenitive electuary, one ounce; falt, half an ounce; oliveoil, two ounces; mix together for use.

The anodyne glyster. .

Take of sheeps-head broth, five ounces; extract. of opium, three grains; diffolve the opium thoroughly in the broth. A very powerful remedy against the dysentery; to be injected at discretion.

The demulcent glyfter. .

Take white decoction, five ounces; ftarch, and Portuguese earth, of each two drachms; the yolk of an egg; mix together for use. In a tenesmus or a dysentery, when the mucus is abraded from the bowels, this glyster may be administered with confiderable advantage.

The turpentine glyster.

Take of the common glyster decoction, of the London Dispensatory, fix ounces; Venice turpentine, half an ounce; diffolve the turpentine in the yolk of an egg; then add four ounces of olive oil. See pages, 8, 53, 94, 123, 139.

The expression of hog-lice.

Take of live hog-lice, one ounce and an half; bruife them in a marble mortar, and add gradually half a pint of fimple penny-royal water, and an ounce of the fimple fyrup. The dofe, two or three ounces night and morning. It is a great detergent, and of fervice in the jaundice, dropfy, afthma, and all chronic diforders that foul the glands.

Extract of the Brazilian bark.

Take of the Brazilian-bark in powder, one pound; boil in ten pints of water, for an hour or two; then pour off the liquor, and boil the remaining bark in ten pints of fresh water for the same length of time; once more pour off the liquor, and add more water; repeat the coction a fourth time; lastly mix these several liquors together, and evaporate the whole to the confistence of an extract, over a gentle fire.

The common gargle.

Take of tincture of roles, a pint; honey of roles, two ounces; mix together,

The

The Portuguese Hospital.

The detergent gargle.

Take of the emollient decoction, a pint; tincture of myrrh, half an ounce; honey, one ounce and an half; mix for use. An excellent gargle in the putrid fore throat.

The aromatic anodyne draught.

Take of fimple pepper-mint water, one ounce and an half; aromatic tincture, forty drops; London philonium, a fcruple; fimple fyrup, a drachm. Its title expresses its virtues. It contains nearly half a grain of opium.

The diuretic draught.

Take of fpring-water, one ounce and an half; diuretic falt, two scruples; oxymel of squills, a drachm.

A Aronger diuretic draught.

Take of fimple penny-royal water, one ounce and an half; lemon-juice, fix drachms; falt of wormwood, half a drachm; tincture of cantherrides, fifteen drops; fimple fyrup, two drachms. Thefe draughts are to be repeated difcretionally, where the use of powerful diuretics feems indicated.

The emetic draught.

Take of carduus tea, two ounces; ipecacoanba in powder, half a drachm; oxymel of fquills, two drachms.

A Stronger emetic draught.

Take of ipecacoanha wine, of the London Difpenfatory, two ounces; emetic tartar, two grains.

The faline draught.

Take of fpring-water, one ounce; falt of wormwood, half a drachm; lemon-juice, a fufficient quantity nuantity to faturate the falt perfectly; tincture of cinnamon, two drachms; loaf-fugar, a fmall lump. This draught may be taken every four or fix hours. For its uses, see pages 16, 62, 141.

The spermaceti draught.

Take of barley-water, one ounce and an half; fpermaceti(diffolved in the yolk of an egg) half a drachm; nitre, a fcruple; fimple fyrup, a drachm. To be taken every four or fix hours. An elegant pectoral. Half a fcruple of volatile falt of hartfhorn may be added occafionally.

The volatile draught.

Take of spring-water, three ounces; volatile falt of hartshorn, a scruple; spermaceti, a drachm and an half; syrup of maidenhair, two drachms; to be repeated discretionally.

The aloetic infusion.

Take of fuccotrine aloes, two drachms; pareira brava-root, four fcruples; boiling water, nine ounces; infuse together, and when cold add half an ounce of compound spirits of lavender. See page, 148.

The bitter infusion.

Take zedoary, the yellow rind of lemons carefully feparated from the white inner rind; of each half an ounce; the yellow rind of Seville oranges alfo carefully feparated from the white inner rind, a drachm and an half; boiling water, twelve ounces; infuse together an hour or two; and then strain off the liquor through flannel without preffure for use. The dose, three ounces two or three times in a day. An elegant and agreeable bitter.

The physicians of this hospital wholly reject gentian-root, not because it possesses any noxious quality

154

The Portuguese Hospital.

lity itfelf; but left the apothecary, through miltake, fhould use a poisonous root, which very nearly refembles it. They therefore in every composition wherever this root is an ingredient, substitute zedoary in its room; and instead of extract of gentian, use the extract of rue.

The diuretic infusion.

Take of zedoary root, two drachms; dried fquills, rhubarb, and juniper berries bruifed, of each a drachm; cinnamon in powder, three drachms; falt of wormwood, a drachm and an half: infufe in a pint and an half of old hock wine, and when fit for use filtre the liquor. The dose, about a gill two or three times in a day. A powerful diuretic and of excellent use in hydropic diforders, and obfiructions of the urinary passages from viscid humours.

Infusion of Inake-root with vinegar.

Take of inake-root powdered, half an ounce; cochineal, half a drachm; boiling water, twelve ounces; let these ingredients infuse together a sufficient time: then strain off the liquor and add thereto of distilled vinegar two ounces. The dose, four spoonfuls every four or fix hours. A noble alexipharmic in contagious, petechial and eruptive fevers.

The mustard infusion.

Take of mustard-seed bruised, one ounce; winter's bark in gross powder, half an ounce; infuse cold in a pint of Lisbon wine: when fit for use strain off the liquor, and add two ounces of compound spirit of laves der. The dose, the same as the former. It is an useful prescription in rheumatic and paralitic cases.

Aftringent

Astringent milk.

Take of pomegranate shells bruised, half an ounce; cinnamon bruised, two drachms; boil in a pint of new-milk; add by little and little, a pint of spring water; continue the coclion till there remains but a pint of liquor. A quarter of a pint may be drank discretionally in hectic cases attended with a diarhœa; or where a soft nutriment is required, and milk alone is apt to purge the patient.

Chalybeate milk.

Take of falt of steel, one ounce; gum arabic, a drachm and an half; boiling water, a pint; fimmer together a few minutes over a gentle fire. A go d medicine in cachectic habits, and obstructions of the menses.

The acidulated lineaus.

Take of conferve of roles, two ounces; weak fpirit of vitriol, four scruples; or rather a sufficient quantity to render the conferve agreeably tart. A tea-spoonful now and then is very proper to cleanse the mouth when parched in fevers.

The common linetus.

Take of conferve of marshmallows, one ounce; oil of olives, and fyrup of marshmallows, of each four ounces.

The folutive linetus.

Take of conferve of hips, one ounce; oil of olives, and folutive fyrup of roles, of each four ounces.

Oily linctuffes are prefcribed in coughs, afthmas, and pleuritic diforders; to heal excoriations, and foften and relax the crifpy tone of the fibres.

The borax liniment.

Take borax in powder, and oil of olives, of each equal parts; rub together in a marble mortar till

The Portuguese Hospital.

they are perfectly incorporated, and become of the confiftence of a liniment. This is used as a difcutient application.

The astringent mixture.

Take of fimple cinnamon-water, ten ounces; extract of the Brazilian-bark, three drachms. Thee or four fpoonfuls may be taken, three times a day, or after every loofe ftool; obferving the cautions laid down in the former part of this work, concerning aftringents. See page 137.

The Peruvian bark mixture.

Take of the decoction of the Peruvian bark, ten ounces; extract of the Peruvian bark, two drachms; fyrop of orange-peels, one ounce. The dofe, two large spoonfuls, to be repeated every four hours, or discretionally, in intermittents and disorders from a relaxed state of the solids.

The purging mixture.

Take of the infusion of fena of the London Difpenfatory, ten ounces; tincture of fena, two ounces; tincture of jalap, three drachms. Three or four table spoonfuls are to be taken at a time, and repeated every hour or two, till a fufficient number of ftools are procured.

The nitrous mixture.

Take of fpring-water, fix ounces; crabs-claws prepared, and nitre, of each a drachm and an half; fyrup of elder, fix drachms. An excellent refrigerant. Two table spoonfuls are to be taken, every four or fix hours.

The squill mixture.

Take of fimple cinnamon-water, eight ounces; vinegar of fquills, and fyrup of marshmallows, of each

PRACTICE OF

each three ounces. An efficacious diuretic if administred in very small doses, frequently repeated, and is of vast fervice in dropsical habits.

The valerian mixture.

Take of fimple pepper-mint water, twelve ounces; powder of valerian-root, one ounce; compound fpirit of lavender, half an ounce; fyrup of orangepeels, one ounce. Three or four table fpoonfulls may be taken every fix or eight hours. A noble medicine in every species of nervous diforders.

The misletoe mixture.

Take of spring-water, twelve ounces; misletoe in fine powde, fix drachms; syrup of orange-peels, one ounce. Its dose, is the same as the former, and it stands recommended as a great antiepeleptic.

Æthiopic pills.

Take of Æthiops mineral, two ounces; gumaloes, half an ounce; fyrup of buckthorn a fufficient quantity to form a mass for pills. The dose, three or four middle fized pills twice in a day. A good alterative.

Compound aloetic pills.

Take of fuccotrine-aloes, two ounces; fcammony in powder, half an ounce; falt of wormwood, three drachms; fyrup of buckthorn, a fufficient quantity to make the whole into a mass. A brisk warm purgative, fit for phlegmatic habits. The dose, from a scruple to two scruples.

Antimonial pills.

Take of precipitated fulphur of antimony, two drachms; calomel finely levigated, and gum guaiacum in powder, of each one ounce; fyrup of buckthorn, a fufficient quantity to form a mafs. Make

158

The Portuguese Hospital.

Make out of each drachm twelve pills ; two of which may be taken night and morning with a draught of the fudorific decoction (See page 148,) the patient keeping himfelf moderately warm. This medicine is a very-ufeful alterative both in cutaneous and venereal diforders. See page 129.

Astringent pills.

Take extract of the Brazilian bark, and Japan earth, of each an ounce; St. Thomas's baliam, a fufficient quantity to form a mass. Twelve pills may be made out of each drachm; two or three of which may be taken discretionally in the fluor albus, old gleets, diarrhæas, &c. observing the usual cautions when given for a diarrhæa.

Hiera picra pills with Aeel.

Take of hiera-picra, one ounce; falt of fteel, fix drachms; compound powder of aron-root, and extract of rue, of each one ounce and an halt; folutive fyrup of rofes, a fufficient quantity to form a mass. Two middle fized pills may be taken night and morning, to promote digestion in languid habits, and in all cases where warm aperient and deobstruent medicines are proper.

Mercurial pills.

Take of quickfilver, five drachms; Venice turpentine, two drachms; gum guaiacum, three drachms; extinguifh the quickfilver perfectly with the turpentine, which if too vifeid may be rendered thinner with a little oil of olives; then add the guaiacum and beat the whole together into a mafs for pills. Form twelve pills out of each drachm, two of which may be taken night and morning with a draught of diet drink, and will be of fervice in all cafes that require mercurial alteratives.

159

Myrrh pills with Steel.

Take of myrrh in powder, fix drachms; falt of fteel, two drachms; gum ammoniacum, and extract of favin, of each three drachms; beat into a maßs with the fimple fyrup. Two middling fize pills may be taken every night and morning in chlorotic cafes, and obstructions of the menses, for which they are very efficatious.

Soap pills.

Take of Castile foap, two ounces; gum ammoniacum, one ounce and an half; fquills in powder, and rhubarb, of each three drachms; form into a mass with the fimple fyrup. Two middle fized pills may be taken night and morning in the jaundice, asthma, and all obstructions of the viscera, for which they are a very powerful remedy.

Storaz pills.

Take of strained storax, two ounces; extract of elecampane, one ounce; strained opium, one drachm; beat together into a mass. Half a drachm of this mass contains a full grain of opium. This compofition is a powerful expectorant.

The antimonial powder.

Take of crude antimony in fine powder, one ounce; Æthiops mineral, fix drachms; rub them well together. The dofe, half a drachm, two or three times a day. This is an admirable medicine in all cutaneous foulneffes, even the worft leprofies. A vomit should be previously administred.

The astringent powder.

Take of the Brazilian bark, three ounces; Japan earth, cinnamon, and olibanum, of each five drachms; reduce them into a fine powder.

The

The Portuguese Hospital.

The aftringent powder with opium,

Is made by adding a drachm of dried opium, to the above ingredients before they are reduced to powder.

Compound powder of Portuguese earth.

Take prepared Portuguese earth, and gum-arabic, of each sour ounces; cinnamon, two ounces; reduce the whole into a fine powder.

Compound powder of Portuguese earth,

Is made by adding a drachm of dried opium, to the above ingredients before they are powdered. The proper dofe, of the above powders, is from a fcruple to half a drachm; to be repeated occafionally.

Purging whey.

Take of Glauber's falt, fix drachms; diffolve in half a pint of water; and add a pint of new milk; then boil this mixture till one half has evaporated; and as foon as the curd has feparated, ftrain off the whey, and add one ounce and an half of manna. A quarter of a pint may be taken difcretionally.

Mindererus's Spirit.

Take of volatile spirit of sal armoniac, four ounces; distilled vinegar, a sufficient quantity to perfectly neutralize the spirit. The vinegar should be added by degrees. See pages 28, 130.

Syrup of elder.

Take of the juice of elder-berries depurated, one pint; loaf-fugar, two pounds; boil together over a flow fire to the confiftence of a fyrup.

Troches

16.2 Practice of the Portuguese Hospital.

Troches of burnt Sponge.

Take of burnt sponge, one ounce; loaf-sugar, three ounces; make into troches with a fufficient quantity of the mucilage of gum tragacanth. These are contrived for scrophulous disorders in children.

Yellow basilicon with red precipitate:

Take of yellow bafilicon, half an ounce; oliveoil, two drachms; red corrolive mercury finely levigated, a scruple; mix together into an ointment. The detergent application commonly ufed in all the hospitals.

All the medicines of the London Difpenfatory; in those compounds, wherein gentian-root is prefcribed, substituting zedoary root in its room. See page 154.

and Thornes at the second base to many the short of a

in a paint of the second second different in the restored by

A GRADE STREAM STREAM

.051 No. 1991 30

man wedge and the a state of the state of the second bear and the

or gringer, a futurent quertile to

Traches

anticar his had no been added and Los this weather

is a light at the a to be all as

od bluost myentie for T. Ante foreite file utheren facult be

the most is the effetite with them in the

Inche cates freehous

"移居"的"五百星"的"五百星"。

Weights and Measures.

I N this book, Troy Weight is used in the composition of the medicines, divided in the following manner:

| The pound - |) a statistic in the | twelve ounces. |
|---------------|----------------------|-----------------|
| The ounce | contains } | eight drachms. |
| The drachm | | three scruples. |
| The fcruple J | Para Barra C | twenty grains. |

For Liquids, wine measure

Is used, and is divided as follows:

The pint } contains { fixteen ounces. The ounce } contains { eight drachms.

A gallon contains eight pints.

A spoonful is equal to the measure of half an ounce.

Alum curd.

Take of the white of an egg at pleafure, and ftir it in a pewter veffel with a fufficient lump of alum, till it is coagulated.

Suppurating cataplasm.

Take of dried figs, four ounces; of yellow bafilicon, one ounce; of strained galbanum, half an ounce. Beat well the figs with a little wine, or strong

164

ftrong stale beer; then carefully mix them in the oin ment first melted with the galbanum.

The preparation of terrestrious and such other bodies as will not dissolve in water.

These bodies are first to be pounded in a mortar, then levigated with a little water upon a bard and smooth marble into an impalpable powder, afterwards dried upon a chalk-stone, and then set by for a few days in some warm, or, at least, a very dry place.

After this manner are to be reduced into powder, amber; antimony; bezoar, which fhould be levigated with fpirit of wine inftead of water; bloodftone; calamine, first calcined for the use of the makers of brass; chalk; coral; crabs claws; crabs eyes; egg-fhells, first separated from the membrane adhering to them by boiling in water; oyster-shells, first cleansed; pearls, verdigrease, tutty.

In antimony, calamine, and tutty, fingular care ought to be taken to reduce them into the most fubtle powder possible.

Cummin cataplasm.

Take of cummin feeds, half a pound; bay berries, the leaves of water germander dried, Virginia fnake root, of each three ounces; of cloves, one ounce. With honey equal to thrice the weight of the fpecies powdered make a cataplafm.

The purification, or trying of bog's-lard, and mutton suet.

Melt them by a gentle fire with the addition of a little water, after they have been first chopt into fmall pieces; and then strain them from the membranes intermixed.

Volatile well and the with a low said Volatile

Volatile epithem.

Take equal weights of common turpentine, and of spirit of fal armoniac. Stir the turpentine in a mortar, and gradually drop in the spirit, till the whole is reduced to a white mass.

Blistering epithem.

Take of cantharides reduced to a very fine powder, and of wheat flower, equal weights; with a fufficient quantity of vinegar, make them into a pafte.

The purification of viper's fat.

Melt the fat, first separated from the intestines, with a gentle heat, and then strain it out through, a thin cloth.

Mercurial cerate.

Take yellow wax, tried hog's-lard, of each half a pound; of quickfilver, three ounces; of the fimple balfam of fulphur, a drachm. Melt the wax with the lard; then add them gradually to the quickfilver first well divided by the balfam of fulphur.

Epulotic cerate.

Take of oil olive, a pound; yellow wax, and prepared calamine, of each half a pound. Melt the wax with the oil, and, as foon as the mixture begins to congeal, fprinkle in the calamine, and ftir all well, till the cerate is quite cold.

The clarifying of boney.

Liquify the honey, by fetting the veffel containing the honey into hot water; and let the fcum, that rifes, be taken off.

Yellow

Yellow cerate.

Take of yellow bafilicon, half a pound; of yellow wax, an ounce. Melt them together.

White cerate.

Take of oil olive, four ounces in measure; of white wax, four ounces in weight; of spermaceti, half an ounce in weight. Melt all together, and ftir them well, till the cerate is quite cold.

Baking of Squills.

Inclose the fquill in passe of wheat flower, having first separated the outer skin, and the hard part from which the fibrous roots grow; then bake the squill in an oven till the passe is dry, and the squill is rendered soft and tender throughout.

Volatile liniment.

Take of oil of almonds, one ounce in measure; of the spirit of sal armoniac, the weight of two drachms. Shake them together in a wide mouthed vial, till they perfectly unite.

The drying of squills.

Cut the fquill, after the outer skin has been taken off, tranversely into thin slices, and dry it with a very gentle heat.

Liniment of three ingredients.

Take of the common plaister, four ounces in weight; of oil olive, four ounces in measure; of vinegar, the measure of one ounce. Set them over a gentle fire, continually flirting them till the liniment has acquired its due confistence.

The roafting of rubarb, and nutmeg.

Roaft them with a gentle heat, till they become eafily friable into powder.

Saponaceous liniment.

Take of the fpirit of rolemary, a pint; of hard Spanish foap, three ounces; of camphor, one ounce. Digest the foap with the spirit of rolemary, till it is diffolved; then add the camphor.

White liniment.

Take of oil olive, three ounces in measure; of spermaceti, the weight of fix drachms; of white wax, two drachms. Melt all together with a gentle fire, briskly stirring without intermission, till the mixture is become quite cold.

The burning of Sponge.

Heat the sponge in a covered vessel, till it becomes black, and is easily friable; then reduce it to powder in a glass or marble mortar.

Green ointment.

Take of the green oil, three pounds; of yellow wax, ten ounces. Melt the wax with the oil over a gentle fire, continually ftirring till the mixture is cold.

The calcination of hartshorn.

Burn pieces of hartshorn in a potter's furnace, till they become perfectly white; then reduce them to powder after the fame manner as other terrestrious substances.

Ointment for bisters.

Take of tried hog's-lard, and of the bliftering plaifter, equal weights. Melt them together with a very

a very gentle heat, and ftir them well, till fully cold.

The extraction of pulps.

Pulpy fruits, that are unripe, and those which are ripe, if dry, are to be boiled in a small quantity of water, till they become toft; then the pulp is to be pressed through a strong hair-fieve, and afterwards boiled over a gentle fire, and continually stirred to avoid burning, till it is brought to a due confistence.

Caffia is also to be boiled out from the pod or cane bruifed, and reduced afterwards to a just confistence by evaporating the water.

The pulps of fruits, which are both ripe and fresh, are to be pressed out without any previous boiling.

Ointment of tutty.

Take any quantity of prepared tutty, and mix with it as much purified viper's fat, as is requifite to bring it to the confiftence of a foft ointment.

Ointment of three ingredients.

Take of the common plaister, four ounces in weight; of oil olive, two ounces in measure; of vinegar, one ounce in measure. Set them together over a gentle fire, continually stirring them, till they are brought to the confistence of an ointment.

Ointment of sulphur.

Take of the fimple ointment, half a pound; of flowers of fulphur unwafhed, two ounces; of effence of lemons, a fcruple. Mix all together.

Lead ointment.

Take of oil olive, half a pint; of white wax, an ounce and an half; of fugar of lead, two drachms. Rub

Rub the fugar of lead, first brought to a very fubtle powder, with some part of the oil; then add this to the wax melted with the rest of the oil, and stir the mixture till it is fully cold.

Ointment of elder.

Take of elder flowers full blown, four pounds; of tried mutton fewet, three pounds; of oil olive, one pound. Boil the flowers, till they become almost crifp, in the fewet and oil first melted together; then prefs them out.

Ointment of tar.

Take of tar, and of tried mutton fewet, equal weights. Melt them together, and ftrain, while hot.

Ointment with precipitate of mercury.

Take of the fimple ointment, an ounce and an half; of precipitated fulphur, two drachms; of white precipitate of mercury, two fcruples. Mix all together, and moisten them with the ley of tartar, to bring the whole to the confistence of an ointment.

Ointment of gum clemi.

Take of tried mutton fewet fresh, two pounds; of gum elemi, one pound; of common turpentine, ten ounces. Melt the gum with the fewet, and all being removed from the fire, add forthwith the turpentine, and, while the mixture is fluid, strain it.

The weaker blue ointment.

Take of tried hog's-lard, four pounds; of quickfilver, one pound; of common turpentine, an ounce. Rub the quickfilver in a mortar with the turpentine, till the quickfilver appears no longer; then add by degrees the lard warmed, and mix them diligently.

The Aronger blue ointment.

Take of tried hog's-lard, two pounds; of quickfilver, one pound; of the fimple balfam of fulphur, half an ounce. Rub the quickfilver with the balfam of fulphur, till the quickfilver no longer appears; then add by degrees the lard warmed, and carefully mix them.

Green basilicon.

Take of yellow bafilicon, eight ounces in weight; of oil olive, three ounces in measure; of prepared verdigreafe, one ounce. Mix all into an ointment.

The Araining of Aorax.

Boil ftorax in water till it becomes foft, then prefs it out between warm iron plates, and feparate the ftorax now cleared of its dregs, from the water.

Opium Arained.

Take of opium cut into fmall bits, one pound weight; diffolve it into a pulp with one pint or lefs of boiling water, with care to avoid burning; and while it remains quite hot, prefs it ftrongly through a linen cloth from its dregs; then reduce the ftrained opium by a water balneum, or other fmall heat, to its first confistence.

Opium fostened in this small quantity of water, passes the strainer unaltered in its substance, and freed only from dregs; but if it be diffolved in a large quantity of water, the gummy and refinous parts will divide from each other.

After the fame manner the reft of the gums may be purified, fuch as gum ammoniacum, affatœtida, galbanum, and the like. But a greater quantity of water may be fafely used. If the refinous part fubfides, let it be taken out, and added towards the conclusion of the infpifiation, that it may unite with the reft into one uniform mais.

Any gum (as galbanum,) which eafily melts may be purified by including the gum in a bullock's bladder, and retaining it in warm water, till the gum becomes foft enough to be feparated from its dregs, by preffing through a canvas ftrainer.

Preparation of millepedes.

Let millepedes be inclosed in a thin canvas cloth, and fuspended within a covered veffel, over the fleam of hot spirit of wine; and they will soon be killed by the vapour, and be rendered friable.

Black basilicon.

Take of oil olive, a pint; yellow wax, yellow rofin, and common pitch, of each nine ounces. Melt all together, and firain the mixture while hot.

Yellow basilicon.

Take of oil olive, a pint; yellow wax, yellow rofin, and Burgundy pitch, of each a pound; of common turpentine, three ounces. Melt the wax, rofin, and pitch, with the oil, over a gentle fire; then take them off, add the turpentine, and ftrain the mixture while it remains hot.

Ointment of marshmallows.

Take of the oil of mucilages, three pints; of yellow wax, one pound; of yellow rofin, half a pound; of common turpentine, two ounces. Melt the rofin and wax with the oil; then, thefe being taken off the fire, add the turpentine, and firain the mixture while it is hot.

Pomatum.

Take of tried hog's-lard, two pounds; of rofewater, three ounces. Pound the lard with the rofe-water, till they are well mixt; then melt the lard with a very gentle fire, and fet it by a little while, that the water may fubfide; afterwards 1 2 pour pour out the lard, and leave the water; then flir and beat the lard without ceafing, while it is growing cold, that it may be broke into as light and yielding a mafs, as may be; and then add as much effence of lemons, as fhall be requifite to give it an agreeable fcent.

Whate ointment.

Take of oil olive, one pint; of white wax, four ounces; of fpermaceti, three ounces. Melt all together with a gentle heat, and ftir them very brifkly without ceafing, till they are fully cold.

Camphorated white ointment.

This is made by adding to the former a drachm and an half of camphor, first beat with a few drops of oil of almonds.

Conferves of the leaves of garden scurvy grass, spearmint, rue, and wood-forrel, of the tops of sea-wormwood; of the flowers of lavender, mallows, rosemary, and red roses, while in bud; and of the outer yellow part of Seville orange peel.

The leaves are to be plucked from their ftalks, and the flowers from their calix's; the outer rind of the orange-peel fhould be foraped off with a rafp, or grater, every one of them when thus prepared, is to be pounded in a mortar with a wooden peftle, first by itfelf, and then with the addition of three times its weight of double refined fugar, till they are well incorporated together.

Conferve of hips.

Take of the pulp of ripe hips, one pound; of double refined fugar, twenty ounces; and mix them into a conferve.

Conferve of floes.

Scald the floes in water to fosten them, taking care their skins are not broken; then take them

173

boil

out and express their pulp, which mix with thrice its weight of double refined fugar.

Blistering plaister.

Take of the drawing plaister, two pounds; of canthalides, one pound; of vinegar, half a pint. The plaister being melted, a little before it hardens, sprinkle in, and mix the cantharides, reduced to a very fine powder; then add the vinegar, and beat all well together.

The scorbutic juices.

Take of the juice of garden fcurvy-grafs, a quart; the juice of brook-lime, and of water-creffes, of each one pint; of the juice of Seville oranges, a pint and a quarter. These being mixed, let them ftand till the dregs subfide; then let the juice be poured off clear, or strained.

Rob of elder-berries.

Let the depurated juice of elder-berries be inspiffated, with a gentle heat, to a proper confistence.

Elaterium.

Slit ripe wild-cucumbers, and pass the juice, gently prefied out, through a very fine hair fieve into a glazed veffel; fet it by some hours, till its thicker part shall have subsided: then pour off as much of the thin part of the juice, as can conveniently be done, by inclining the veffel, and draw away the rest by the filtre: let the thicker part which remains, be covered over with a linen cloth and dried, either in the fun or by a gentle fire.

Extracts, of the roots of elecampane, gentian, and black hellebore; and of the leaves of rue and favin.

Boil them in water, strain and press out the decoction, and set it by, till its dregs are subfided; then

174

boil it to the confiftence of a pill, with care towards the end to avoid burning.

Extract of liquorice.

Boil the roots of liquorice lightly in water, ftrain and prefs out the decoction; then, after its dregs have fubfided, boil it away, till it will not flick to the fingers, using due care towards the end to avoid burning.

Extract of logwood.

Take of logwood, in powder, one pound. Boil it four times, or oftener, in a gallon of water to half; then boil all the liquors mixt together and firained, to a just confistence.

Extract of Peruvian bark, both fift and hard.

Take of Peruvian bark, reduced to powder, one pound; of water, ten or twelve pints. Boil for an hour or two, and pour off the liquor, which will be red and transparent; but as soon as it grows cold, becomes yellow and turbid; boil the bark again in the same quantity of fresh water as before, repeating these boilings till the liquor remains transparent when cold: then evaporate all these decoctions strained and mixt together, to the proper confistence, over a gentle fire, with due care to avoid burning.

This extract is to be prepared under a double form; one, of the confiftence of a pill; the other, hard enough to be reduced to a powder.

Extract of lignum vitæ, soft and hard.

Take of the fhavings of lignum vitæ, one pound. Boil them four times, or oftener, in a gallon of water to half; then infpiffate the liquors, after they have been ftrained: but when the water is near all dried away, add a fmall portion of rectified fpirit,

spirit, by which the extract will be brought to an uniform and tenacious mass.

This extract is also to be prepared under two forms, one solter and the other harder.

Extract of jalap.

Pour upon jalap root powdered, rectified spirit of wine; and with a due heat draw a tincture; and boil the refidue several times in water: after straining, draw off the spirit from the first tincture, till it begins to thicken; inspissate also the strained decoctions; then mix the two extracts, and, with a gentle fire, reduce them to the confissence of a pill.

The cathartic extract.

Take of focotorine aloes, an ounce and an half; of the pith of coloquintida, fix drachms; feammony, and the leffer cardamom feeds hufked, of each half an ounce; of proof fpirit, a pint. The fpirit being poured upon the coloquintida cut fmall, and the feeds bruifed, draw a tincture with a gentle heat continued four days; then, to the tincturepreffed out, add the aloes and feammony, firft feparately reduced to powder; and, thefe being diffolved, draw off the fpirit, and reduce the mais to the confiftence of a pill.

The gum and refin of aloes.

Take of focotorine aloes, four ounces ; of water, a quart. Boil the aloes till it is diffolved as much as may be, and fet all by for a night: the refin will be precipitated to the bottom of the veffel: the liquor poured off or ftrained, being evaporated, will leave the gum.

Oil of almonds.

Let either fweet or bitter almonds that are fresh, be pounded in a stone mortar, and then the oil forced out with a press not heated. 176

After the fame manner fhould the oil be preffed from linfeed and mustard-feed.

Effential oils, from the root of fassfafras; from the leaves of fweet marjoram, wild marjoram, peppermint, spear mint, penny-royal, rosemary, rue, savine, and wormwood; from the flowers of chamomile, and lavender; from the seeds of anise, carraway, cummin, and dill; from juniper berries; and from the spices, cloves, nutmeg, and others.

These oils are obtained by diffillation, with an alembic and large refrigeratory. Water must be added to the materials, in sufficient quantity, to prevent their burning, and the subject be macerated in that water a little time before the diffillation. The oil comes over with the water, and either swims on the top, or finks to the bottom, according as it is heavier or lighter.

Oil of box.

Diftil pieces of box in a retort with a fire gradually raifed. The oil will come over with an acid fpirit, from which the oil is to be feparated by a funnel.

Oil of bricks.

Let bricks heated red hot, be plunged into oil olive, till the whole is imbibed: then the bricks being fufficiently broke, are to be put into a retort: and, by a fand heat the oil will afcend, with a fpirit, which is to be feparated from the oil.

Oil of Barbadoes tar.

Let Barbadoes tar be distilled in a fand heat, and oil will afcend with a spirit.

Oil of turpentine.

Turpentine is to be diffilled with water in a copper flill, like the effential oils of vegetables.

After

After the diffillation, remains in the still yellow rofin:

This oil is often, though improperly, called fpirit of turpentine.

The etherial oil, and the balfam of turpentine.

Let oil of turpentine be distilled in a retort with a very gentle heat, till what remains is become of the confistence of a balfam.

Balfam of turpentine may also be diffilled from yellow rofin, whence, after a portion of oil, which must be removed in time, will come a thick balfam; a blackish rofin remaining in the retort, which is also called colophony.

Compound oil of balfam of copaiva.

Take of balfam of copaiva, two pounds; of gum guaiacum, four ounces. Distil them together in a retort.

Purified nitre.

Boil nitre in water, and ftrain the water through bloffom paper; then, after due evaporation, fet it by in a cold place, that the nitre may fhoot.

In the fame manner is purified fal-ammoniac.

Burnt alum.

Let alum be put into an iron or earthen veffel, and calcined as long as it rifes up and fwells.

Calcined vitriol.

Put green vitriol into an earthen veffel, and calcine it with an open fire as long as it exhales any moifture; then take it out by breaking the veffel, and fet it by for ufe, well clofed from the air. The vitriol is most perfectly calcined, if, at the bottom and fides of the veffel containing it, it is become red.

Salt

Salt of wormwood.

Let the afhes of wormwood be put into an iron pot, and kept red hot by a ftrong fire for tome hours, often ftirring them, that all remains of oil may be burnt out; then boil them in water; ftrain the water, which will be impregnated with the falt, through paper, and evaporate it to drynefs.

In this manner is to be prepared the fixt alkaline falt of any plant, whose ashes will yield that kind of falt.

Salt of tartar.

Wrap up any species of tartar in strong brown paper, first made wet, or inclose it in a proper vessel, and expose it to the fire, that its oil may be burnt out; then boil it in water, and collect its falt, as before.

Ley of tartar.

When the tartar is calcined white, let it be put in a damp place, that it may liquify by the moifture of the air.

Soluble tartar.

Take of any alkaline fixt falt, a pound; of water, a gallon. The falt being diffolved in the water boiling, throw in chryftals of tartar in powder, as long as any fermentation is raifed, which ufually ceafes before thrice the weight of the alkali is thrown in. Then firain the liquor through paper; and after due evaporation fet it by, for the falt to chryftallize, or elfe evaporate the liquor wholly away, that the falt may be left dry.

Soap leys.

Take equal weights of Russia pot-ash, and quicklime, and throw water upon them by degrees, till the lime is flaked; then throw on more water, and ftir

ftir all together, that the falt of the afhes may be diffolved; after fome time pour the liquor, (filtered through paper, if needful) into another veffel. A true ftandard wine pint of this liquor, meafured with the greateft exactnefs, ought to weigh juft fixteen ounces; if it is heavier, for every drachm it exceeds that weight, an ounce and an half of water in meafure is to be added to each pint of the liquor; but, if it is lighter, it must be boiled, till the like quantity of water is carried off, or elfe must be thrown upon fresh lime and ashes.

Almond Joap.

Take any quantity of fresh oil of almonds, and thrice its measure of the foap-leys. Digest them together for some time, in such a heat wherewith the mixture shall but just boil, and within a few hours the oil and leys will be united ; after which the liquor in boiling will foon become ropy, and in a good degree transparent, and will cool into the confistence of a jelly; then throw in fea falt, till the boiling liquor has loft its ropinefs; continue the boiling, till drops of the liquor being received upon a tile, the water is seen to separate freely from the coagulated foap; then remove the fire, and the foap will gradually rife to the top of the liquor, which is to be taken out, before it is cold, and put into a wooden frame, which has a cloth for its bottom : in the last place, being taken out, it is to be fet by, till it acquires its just confistence.

After the same manner may soap be made with oil olive; in which the finest oil ought to be employed, that the soap may be as little ungrateful, either to the palate or stomach, as possible.

I6

The common Aronger caustic.

Boil to a fourth part any quantity of the foapleys above defcribed; then fprinkle in, while boiling, lime that has been kept in a vefiel pretty clofe ftopt for feveral months; continue to add this lime, till all the liquor is abforbed, and the whole reduced to a pafte, which is to be kept in a vefiel well ftopt.

The common milder caustic.

Take of foft foap, and of fresh quick lime, equal parts; and mix them at the time of using.

Spirit of vinegar.

Let vinegar be distilled with a gentle heat as long as the drops fall free from any empyreuma.

If fome part of what comes first off, be thrown away, what is referved will be stronger.

Diuretic falt.

Take of any alkaline fixt falt, one pound; and boil it in four or five pints of diffilled vinegar with a very gentle heat; when the fermentation ceases, add more diffilled vinegar; and when the fermentation arising from the addition is over, pour on another quantity of the like vinegar; and proceed thus till the vinegar being near all evaporated, frefh vinegar will not excite any fermentation; which will generally happen by the time about ten quarts of vinegar shall have been used; then gently evaporate to drynefs. The falt left will be impure, which is to be melted for a time, but not too long, with a gentle heat, afterwards diffolved in water, and transcolated through paper. If the meling has been rightly performed, the ftrained liquor will be limpid and colourlefs, like water, but otherwife brownish. Lastly, the water is to be evaporated with a very gentle heat, in a fhallow veffel; the falt,

falt, as it dries, being frequently flirred, that the humidity may the fooner be difcharged. This falt must be kept in a close veffel, that it may not run by the moisture of the air.

The falt ought to be very white, and fhould diffolve wholly, either in water, or fpirit of wine, without leaving any fæces. If the falt, though ever fo white, leave in fpirit any fæces, after it is diffolved in this fpirit, it is to be filtred through paper, and dried again.

The weak and Arong Spirit of vitriol, alfo the colcothar.

Vitriol, first calcined, is to be diffilled in earthen veffels for three days without intermission, in a reverberatory heat; the liquor drove off being diftilled over again in a glass retort, with a fand heat, the weak spirit will ascend, the strong remaining behind, which is usually, though improperly, called oil of vitriol.

What remains in the earthen veffels after the first distillation, is called the colcothar of vitriol.

Glauber's spirit of nitre.

Take of nitre, three pounds; of the firong fpirit of vitriol, one pound. Let them be mixt with caution, and gradually, under a chimney; afterwards let them be diffilled, first with a gentle heat, and then with a fironger.

Glauber's spirit of Sea-falt.

Take fea-falt, and the ftrong spirit of vitriol, of each two pounds; of water, one pint. The oil and water being first mixt together, add the mixture gradually to the falt, under a chimney; then distil, first with a small, and asterwards with a stronger fire.

Aqua-

182

Aqua-fortis.

Take nitre, and green vitriol, not calcined, of each three pounds; of the fame vitriol, calcined, a pound and an half. Mix all well together, and diftii with a very ftrong fire, as long as red fumes arife.

Compound aqua-fortis.

Take of aqua-fortis, fixteen ounces in weight; of falt, one drachm. Distil to drynes.

Salt of vitriol.

Take of white vitriol, a pound; of the ftrong fpirit of vitriol, an ounce in weight; of water, as much as is fufficient. Diffolve the vitriol by boiling; then ftrain the decoclion through paper, and after proper exhalation fet it in a cold place, that the falt may fhoot.

Vitriolated tartar.

Take of green vitriol, the weight of eight ounces; of water, two quarts. The vitriol being diffolved in the water boiling, throw in falt of tartar, or any other fixt alkali, till all fermentation ceafes; which ufually happens after throwing in four ounces, or fomething more, of the alkaline falt. Then ftrain through paper, and evaporate duly, that the falt may chryftallize.

Vitriolated nitre.

Diffolve the cake left after the diffillation of Glauber's fpirit of nitre, as defcribed above, in hot water; and after purifying through paper, evaporate, that the falt may fhoot.

Glauber's cathartic falt.

Diffolve in water, the cake which remains after the diffillation of Glauber's fpirit of fea-falt; purify the folution through paper, and then duly evaporate it, that the falt may christallize.

Spirit

SUPPLEMENT. IST

Spirit of Sea-Salt coagulated.

Pour gradually upon spirit of sea-falt, the lixivium of any fixt alkali, till all fermentation ceases; and then evaporate to drynes.

Dulcified Spirit of vitriol.

Take of the ftrong spirit of vitriol, called the oil, one pound; of rectified spirit of wine, one pint.

Dulcified Spirit of nitre.

Take of rectified spirit of wine, one quart; of Glauber's spirit of nitre, half a pound. Mix them by pouring the spirit of nitre on the other, and distil the mixture with a gentle heat, as long as what comes off will not raise any fermentation with a lixivial falt.

Spirit of fulphur by the bell.

Let fulphur be fet on fire under a glass veffel fitted for this purpose, which is usually called a bell, and the acid spirit will drop from it, which is to be received into a dish placed underneath.

Spirit, falt, and oil of amber.

Let amber be diftilled by a fand heat, gradually increased. From it will arise a spirit, oil, and falt, foul by a mixture of the oil.

The oil diffilled again will part into a thinner oil, which will afcend, and a thicker part remain, called the balfam of amber.

The falt is to be boiled either in the fpirit, or water, and fet by to fhoot: thus it will be freed from its oil, and the oftner this process is repeated, the purer will the falt be.

Spirit, Salt, and oil of bartshorn.

Distil fragments of hartshorn with a fire gradually raised to a great height. A spirit, falt, and oil will ascend. If the oil be feparated, and the fpirit and falt mixt together, be diffilled again with a very gentle heat, they will both rife more pure. If this is carefully repeated feveral times, the falt will become very white, and the fpirit as limpid as water, with a grateful fmell.

If the falt be feparated from the fpirit, and fublimed, first from an equal weight of fine chalk, and then again from a small quantity of rectified spirit of wine, it will become sooner pure.

Calcined hartfhorn is, for the most part, made by burning the horns, after they have passed through the preceding operation.

After the fame manner a fpirit, falt and oil, may be diftilled from any animal substance.

The spirit, falt, and oil of foot.

Diftil wood foot in the fame manner as hartfhorn; but here more labour is required to render the fpirit and falt pure.

Volatile falt of fal-armoniac.

Take of the finest chalk, or whiting, two pounds; of fal-armoniac, one pound. Sublime the volatile falt in a retort with a strong fire.

Spirit of Sal-armoniac.

Take of any fixt alkaline falt, a pound and an half; of fal-armoniac, a pound; of water, two quarts. With a gentle fire diftil off one quart.

Dulcified spirit of fal-armoniac.

Take of any fixt alkaline falt, half a pound; of fal-armoniac, four ounces; of proof fpirit, three pints. Diftil off, with a gentle fire, a pint and an half.

Fetid

2

Fetid volatile Spirit.

Take of any fixt alkaline falt, a pound and an half; of fal-armoniac, a pound; of affafœtida, four ounces; of proof fpirit, three quarts. Diftil off with a gentle heat, five pints.

Aromatic volatile Spirit.

Take effence of lemons, and effential oil of nutmegs, of each two drachms; of effential oil of cloves, half a drachm; of dulcified fpirit of falarmoniac, a quart. Diftil with a very gentle fire.

Flowers of benjamin.

Put powdered benjamin into an earthen-pot placed in fand, and with a fmall heat the flowers will rife, and may be caught by a paper cone placed over the pot.

Or else the benjamin may be put into a retort, and the flowers will ascend into, and fasten themfelves upon its neck.

The flowers, if tinged yellow, are to be mixt with tobacco-pipe clay, and fublimed again.

Flowers of Sulphur.

Let fulphur be fublimed in a fit veffel; and any part of the flowers, which may have concreted, are to be reduced to powder by a wooden mill, or in a marble mortar with a wooden peftle.

Flowers of fulphur washed.

Pour water on the flowers, to the height of three or four fingers above them, and boil them for a time: then pour off this water, and with fresh cold water wash the remains of this away; then dry the flowers for use.

The

The simple balfam of fulphur.

Boil flowers of fulphur in four times their weight of oil olive, in a pot lightly covered, till the oil and fulphur are joined into the confiftence of a balfam.

In the fame manner is a balfam of fulphur alfo prepared with Barbadoes tar.

Precipitated fulphur.

Boil flowers of fulphur with thrice their weight of quick lime, till the fulphur is diffolved, and filtre the folution through paper; then with weak fpirit of vitriol make a precipitation, which is to be often washed, till it is become quite infipid.

Sulphurated water.

Take of water, a quart; of fulphur, half a pound. Let fome portion of the fulphur, fet on fire in an iron ladle, be fufpended over the water in a close vefiel: and let this be repeated, as often as the fumes from the last fulphur fubfide, till the whole is burnt away.

The ruft of steel prepared.

Expose filings of steel to the air, and moisten them sometimes with water or vinegar, till they are turned into rust; then rub them in a mortar, and by pouring on water, wash off the finest powder; the residue, which by moderate rubbing was not brought to a powder fine enough to be washed off, is again to be exposed moist to the air, and, when farther rusted, is to be treated as before. The powder thus washed off, is to be dried and kept for use.

Steel prepared with fulphur.

Touch the steel heated to a white heat, with a roll of brimstone, that the steel may melt, and drop

drop into water placed under it. 'Then let it be feparated from the fulphur, which has dropt along with it into the water, and be reduced into the fineft powder.

Martial flowers.

Take of washed colcothar of green vitriol, or of iron filings, one pound; of fal-armoniac, two pounds. Mix and sublime them in a retort, and mixing again the bottom with the flowers, renew the sublimation, till the flowers acquire a beautiful yellow colour.

To the refidue may be added half a pound of fresh fal-armoniac, and the sublimation repeated; and the same process may be thus continued on, as long as the flowers rise duly coloured.

The ley of iron.

Set by the refidue after the foregoing fublimation, in a damp place, that it may liquify by the air.

Salt of iron.

Take of the ftrong spirit, or oil of vitriol, the weight of eight ounces; of filings of iron, four ounces; of water, a quart. Mix them, and when the ebullition has ceased, set the mixture some time upon a fand heat; then filtre the liquor through paper, and evaporate it, that the salt may chrystallize.

The lunar caustic.

Diffolve pure filver by a fand heat, in about twice its weight of aqua-fortis; then dry away the humidity with a gentle fire; afterwards melt it in a crucible, that it may be poured into proper moulds, carefully avoiding over much heat, left the matter fhould grow too thick. See page 81.

Sugar

Sugar of lead.

Boil ceruffe in diffilled vinegar, in a leaden veffel, till the vinegar is fufficiently fweet; then filtre the vinegar through paper, and after due evaporation fet it by, that the falt may fhoot.

The medicinal some.

Take alum, litharge, and bole-armeniac, or French bole, of each half a pound; of the colcothar of green vitriol, three ounces; of vinegar, a quarter of a pint. Dry the whole mixt together over a fire, till it grows hard.

Powdered tin.

Let melted tin be poured into a wooden-box chalked within; and while the tin grows cold, let the box be brickly fhaken, and part of the tin will be reduced to powder. The remainder, by being treated in the fame way, may also be reduced to powder.

The purification of quickfilver.

Diftil the quickfilver in a retort, and then wafh it well with water and falt, or vinegar.

Mofaic gold.

Take of tin, one pound; of flowers of fulphur, feven ounces; fal-armoniac, and purified quickfilver, of each half a pound. Add the quickfilver to the tin melted; when the mixture is cold, reduce it to powder; mix well with it the fulphur and falarmoniac, and fublime the compound in a matras. The mofaic gold will be found under the part fublimed, with a fmall quantity of foulnefs at the bottom.

Æthiops mineral.

Take equal parts of quickfilver purified, and of flowers of fulphur unwashed. Rub them together

in

in a mortar of glafs or marble, till the quickfilver perfectly difappears, and the union is perfected.

Artificial cinnabar.

Take of purified quickfilver, twenty-five ounces; of fulphur, feven ounces: flir the quickfilver into the fulphur melted, and if the mixture takes fire, it is to be extinguished by covering the vefiel. Then let the matter be reduced to powder, and fublimed.

Corrosive Sublimate.

Take of purified quickfilver, forty ounces; of fea falt, thirty-three ounces; of nitre, twentyeight ounces; of calcined green vitriol, fixty-fix ounces. Rub the quickfilver firft with about an ounce or more of corrofive fublimate in a wooden or ftone veffel, till it breaks into fmall grains; then mix it with the nitre, afterwards with the fea-falt, till the quickfilver quite difappears; laftly, add the calcined vitriol, but do not rub the mixture too long with it, left the quickfilver fhould begin to part again. Sublime the mixture in a matras, to which may be fitted an alembic head, that a fpirit which will afcend in a fmall quantity, may be faved.

Dulcified mercury Jublimate.

Take of corrofive fublimate, one pound; of purified quickfilver, nine ounces. Add the quickfilver to the fublimate reduced to powder, and in a glafs matras digeft them together in a gentle fand heat, often fhaking the glafs, till they are united. Then augmenting the heat, fublime the mixture. After an acrid part on the top of the fublimation is fcraped off, and if any globules of quickfilver chance to appear, they likewife being feparated, the mafs fublimed is to be reduced to powder, and fublimed

190

fublimed again; the fublimation is to be fix times repeated.

Calcined quickfilver.

Set quickfilver purified, upon a fand-heat for feveral months, in a glafs veffel with a broad bottom, and opening to the air by a finall hole, till it is reduced to a red powder.

White precipitate mercury.

Take of fal-armoniac, and of corrofive fublimate, equal weights. Diffolve them together in water, filtre the folution through paper, and with a folution of fome alkaline fixt falt, make a precipitation; then wafh off all acrimony from the precipitated powder.

The mercurial red corrofive.

Take of quickfilver purified, and of the compound aqua-fortis, equal weights. Set them together in a flat bottomed glafs, upon a fand-heat, till all humidity is exhaled, and the dry mafs has acquired a red colour.

Coralline mercury.

Pour upon the mercurial red corrofive, thrice its weight of rectified fpirit of wine, and digeft them together two or three days in a gentle heat, often fhaking the vefiel: then fet fire to the fpirit, flirring the powder continually till the fpirit is quite burnt away.

The yellow mercurial emetic.

Pour upon purified quickfilver in a glafs veffel, double its weight of the firong fpirit of vitriol. Let the liquor heat gradually, and then boil till in the bottom of the glafs there remains a white mafs, which is to be perfectly dried with a firong heat. This, upon the effution of warm water, will turn yellow, and fall into powder. Rub this powder

powder and the warm water diligently together in a glafs mortar. Then, when the powder has fubfided, pour off this water, and wafh the powder often with fresh water, till it is become perfectly free from all acrimony.

The precipitated sulphur of antimony.

Take of antimony, fixteen ounces; of tartar, a pound; of nitre, half a pound. Being feparately reduced to powder, mix them well, and throw them by degrees into a crucible, red hot, and melt the mixture with a flrong fire; then pour it out into a conical mould, that the metallic part, commonly called the regulus of antimony, may fall to the bottom, while the fcoriæ float at top. Diffolve thefe fcoriæ in water, and filtre the folution through paper; then precipitate the fulphur, by dropping in fpirit of fea-falt; and afterwards wafh it with water from the falts adhering.

Crocus of antimony.

Take of antimony and nitre, equal weights. Being feparately reduced to powder, let them be well mixt, and then gradually thrown into a hot crucible, to melt; the matter being poured out, is to be feparated from its fcoriæ. It will not always appear of the fame colour; it is the more yellow, the longer it has been melted.

Washed crocus of antimony.

Boil the crocus of antimony, reduced to a very fine powder, in water; and this water being poured way, wash the powder often with hot water, till the water comes off infipid.

Emetic tartar.

Take washed crocus of antimony, and chrystals of tartar, of each half a pound; of water, three pints.

pints. Boil them together for half an hour, then filtre the water through paper, and after due evaporation fet it by, that the falt may chryftallize.

The calx of antimony.

Let antimony in powder be well mixed with thrice its weight of nitre, and thrown by degrees into a crucible moderately heated. Then being removed from the fire, let it be washed in water, both from the adhering falts and from what coarser parts may have been less perfectly calcined than the rest.

The antimonial caustic.

Take of antimony, one pound; of corrofive fublimate, two pounds. Being reduced feparately into powder, mix them well, and diffil them in a retort with a wide neck, in a gentle heat of fand. Let what afcends into the neck of the retort be expofed to the air, that it may run into a liquor.

Cinnabar of antimony.

Sublime the remains of the former process in a coated bolt-head with an open fire.

Simple alexiterial water.

Take of the green leaves of fpear-mint, a pound and an half; the tops of fea wormwood, likewife green, the green leaves of angelica, of each a pound; of water, as much as is fufficient to prevent burning. Diftil off three gallons.

Water of dill seed.

Take of dill feed, one pound; of water, as much as is fufficient to prevent burning. Diftil off a gallon.

The simple distilled water of orange-peel.

Take of the outer yellow rind of fresh Seville oranges, four ounces; of water, as much as is sufficient to prevent burning. Distil off a gallon. Water

Water of castor.

Take of Ruffia castor, one ounce; of water, as much as is sufficient to prevent burning. Distil off a quart.

Simple cinnamon water.

Take of cinnamon, one pound; of water, as much as is fufficient to prevent burning. Diftil off a gallon.

Fennel water.

Take of the feeds of sweet fennel, a pound; of water, as much as is sufficient to prevent burning. Diftil off a gallon.

Simple pepper-mint water.

Take of the leaves of pepper-mint, dried, a pound and an half; of water, as much as is fufficient to prevent burning. Diftil off a gallon.

Simple Spear-mint water.

Take of the leaves of spear-mint, dried, a pound and a half; of water, as much as is sufficient to prevent burning. Distil off a gallon.

Water of Jamaica pepper.

Take of Jamaica pepper, half a pound; of water, as much as is fufficient to prevent burning. Diftil off a gallon.

Simple penny-royal water.

Take of the leaves of penny-royal, dried, 2 pound and a half; of water, as much as is sufficient to prevent burning. Diftil off a gallon.

Damask-rose water.

Take of fresh damask-roses, fix pounds; of water, as much as is sufficient to prevent burning. Distil off a gallon.

Spiri-

Spirituous alexiterial water.

Take of the green leaves of fpear-mint, half a pound; the leaves of angelica green, the tops of fea-wormwood green, of each four ounces; of proof-fpirit, one gallon; of water, as much as is fufficient to prevent burning. Diffil off one gallon.

Spirituous alexiterial water with vinegar.

Take the green leaves of spear-mint, the leaves of angelica likewise green, of each half a pound; of the tops of sea-wormwood green, four ounces; of proof-spirit, one gallon; of water, as much as is sufficient to prevent burning. Distil off one gallon, and then add one pint of vinegar.

Compound anifeed-water.

Take anifeeds, and angelica feeds, of each half a pound; of proof-fpirit, a gallon; of water, as much as is fufficient to prevent burning. Diftil off a gallon.

The spirituous water of orange-peel.

Take of the outer yellow rind of fresh Seville oranges, half a pound; of proof-spirit, a gallon; of water, as much as is sufficient to avoid burning. Distil off a gallon.

Water of cardamon Jeeds.

Take of the leffer cardamon feeds, hufked, four ounces; of proof-fpirit, a gallon; of water, as much as is fufficient to prevent burning. Diftil off a gallon.

Spirituous cinnamon-water.

Take of cinnamon, a pound; of proof-spirit, a gallon; of water, as much as is sufficient to prevent burning. Distil off a gallon.

Compound

Compound juniper water.

Take of juniper-berries, a pound; carrawayfeeds, fweet fennel-feeds, of each an ounce and a half; of proof-fpirit, a gallon; of water, as much as is fufficient to prevent burning. Diftil off a gallon.

The spirituous water of pepper-mint.

Take of the leaves of pepper-mint, dried, a pound and a half; of proof-spirit, a gallon; of water, as much as is sufficient to prevent burning. Distil off a gallon.

Spirituous Spear-mint water.

Take of the leaves of fpear-mint, dried, a pound and a half; of proof-fpirit, a gallon; of water, as much as is fufficient to prevent burning. Diftil off a gallon.

Nutmeg water.

Take of nutmeg, two ounces; of proof-spirit, a gallon; of water, as much as is sufficient to prevent burning. Diltil off a gallon.

Spirituous penny-royal water.

Take of the leaves of penny-royal, dried, a pound and a half; of proof-fpirit, a gallon; of water, as much as is fufficient to prevent burning. Diftil off a gallon.

Compound water of borfe-radifb.

Take of the fresh leaves of garden scurvygrafs, four pounds; fresh horse-radish root, the outer yellow rind of fresh Seville oranges, of each two pounds; of nutmegs, nine ounces; of prooffpirit, two gallons; of water, as much as is fufficient to prevent burning. Distil off two gallons.

Simple

Simple Spirit of lavender.

Take of fresh lavender flowers, a pound and a half; of proof-spirit, a gallon. Distil off in a water balneum five pints.

Spirit of rosemary.

Take of fresh rosemary tops, a pound and a half; of proof-spirit, a gallon. Distil off in a water balneum five pints.

Compound Spirit of lavender.

Take of fimple spirit of lavender, three pints; of spirit of rosemary, one pint; cinnamon, nutmegs, of each half an ounce; of rod fanders, three drachms. Digest them together, and then strain off the spirit.

The common decostion of glyfters.

Take of mallow leaves, dry, one ounce; dried chamomile flowers, sweet fennel seeds, of each half an ounce; of water, one pint. After boiling strain it off.

Pectoral decoction.

Take of common barley, raifins ftoned, and figs, of each two ounces; of liquorice root, half an ounce; of water, two quarts. Boil the water first with the barley, then add the raifins, and afterwards toward the latter end of the decoction, the figs and liquorice; the decoction being fully ended when one quart only of liquor will be left after ftraining.

Barley water.

Take of pearl-barley, two ounces; of water, two quarts. Wash the barley first well with some cold water; then pouring on half a pint of water, boil

boil it a little while; and this water, which will be coloured, being thrown away, put the barley into the quantity of water above directed, first made builing hot, and boil away to half.

The common fomentation.

Take the leaves of fouthernwood, or of lavender-cotton, dry, the tops of fea wormwood alfo dry, dried chamomile flowers, of each one ounce; of bay leaves, dry, half an ounce; of water, three quarts. After a light boiling ftrain the water off.

Viper broth.

Take a viper of middle fize without the fkin, head, or entrails; of water, a quart. Boil to about a pint and a half; remove all from the fire, and when the water is cold, if the viper be not a dried one, take away the congealed fat. Then take a chicken of a middle fize, drawn, and the fkin with all the fat taken off, and put it whole into this decoction while cold, fet it upon the fire till it boils; then remove it from the fire, take out the chicken, cut the flefh of it into fmall pieces, which put again into the water, and fet it over the fi e, but as foon as it begins to boil up, pour it off; firft having taken away whatever foum may have arifen.

Mucilage of quince feed.

Take of quince feeds, a drachm; of water, fix ounces. Boil with a gentle fire, till the water becomes roapy, refembling the white of an egg; then ftrain it through a linen cloth.

Alum whey.

Take of cow's milk, one pint; of alum in powder, two drachms. Boil, till the whey is formed, which is to be well feparated from the curd.

Scorbutic

Scorbutic whey.

Take of cow's milk, one pint; of the fcorbutic juices, a quarter of a pint. Boil till a whey is formed, which is to be well feparated from the curd.

The simple bitter infusion.

Take gentian root, the yellow rind of lemonpeel, fresh, carefully separated from the inner white part, of each half an ounce; of the yellow rind of Seville orange-peel, also carefully separated from its inner white part, but dried, a drachm and an half; of boiling water, three quarters of a pint. After infusing for an hour or two, strain it, either through paper or a cloth, without any preffing out.

The purging bitter infusion.

Take the leaves of fena, the yellow rind of fresh lemon-peel, of each three drachms; gentian root, the yellow part of Seville orange-peel, dried, the lesser cardamon feeds husked, of each half a drachm; of boiling water, five ounces. After infusing till the liquor is cold, strain it off.

The common infusion of fena.

Take the leaves of fena, an ounce and an half; of crystals of tartar, three drachms; of the leffer cardamon feeds husked, two drachms; of water, one pint: Boil the chrystals of tartar in water, till they are diffolved, then pour the water, while boiling hot, upon the fena and the reft. When the liquor is cold, strain it off.

The infusion of Sena with lemon.

Take of the leaves of fena, one ounce and an half; of the yellow of fresh lemon-peel, an ounce in

in weight; of lemon juice, an ounce in measure; of boiling water, one pint. Infuse till cold, and then ftrain.

Simple lime water.

Take of quick-lime, one pound; of water, a gallon and an half. Pour the water on gradually; and after the ebullition is over, let the lime fubfide, and the liqour be filtred throught the paper.

The less compound lime-water.

Take of liquorice, one ounce; of faflafras bark, half an ounce; of fimple lime-water, three quarts. Infuse two days without heat, and then strain off the liquor.

The more Compound lime-water.

Take of the raspings of lignum-vitæ, half a pound; of liquorice, one ounce; of saffafras bark, half an ounce; of coriander seeds, three drachms; of simple lime-water, three quarts. Infuse as before, and then strain off.

Tinsture of roses.

Take of red rofe-buds, the white heels being cut off, half an ounce; of the ftrong spirit of vitriol, called the oil, one scruple; of boiling water, two pints and an half; of double refined sugar, an ounce and an half. First add the spirit of vitriol to the water, in a vessel of glass or earth glazed; and then infuse the roses: ftrain the liquor when cold, and add the sugar.

Vinegar of Squills.

Take of dried fquills, one pound; of vinegar, three quarts. Infuse the fquills in the vinegar with a small heat, then press it out, and set it by, K 4

200

till the dregs are fubfided; afterwards, to the depurated vinegar, add about a twelfth part of prooffpirit, that it may be preferved from contracting dregs by time.

Aloetic alkaline wine.

Take of any fixt a'kaline falt, eight ounces; focotorine aloes, faffron, myrrh, of each one ounce; of purified fal-armoniac, fix drachms; of white wine, a quart. Infuse them together without heat for a week, or longer. Then filtre the wine through paper.

Bitter wine.

Take gentian root, the yellow part of fresh lemon-peel, of each one ounce; of long pepper, two drachms; of white wine, a quart. Infuse without heat, and strain.

Antimonial wine.

Take of the crocus of antimony, washed, one ounce; of white wine, a pint and an half. Infuse without heat, and then strain the wine off through paper.

Chalybeate wine.

Take of filings of iron, four ounces; cinnamon, and mace, of each half an ounce; of Rhenish wine, two quarts. Insufe a month without heat, often stirring; then strain it off.

Saffron wine.

Take of faffron, one ounce; of canary, one pint. Infuse without heat, and firain.

Wine with ipecacoanha.

Take of the root ipecacoanha, two ounces; of the yellow part of Seville orange-peel, dried, half an ounce; of canary, a quart. Infuse without heat, and firain.

Viper

Viper wine.

Take of dried vipers, two ounces; of white wine, three pints. Infuse with a gentle heat for a week, and then strain the wine off.

Tincture of rhubarb in wine.

Take of rhubarb, two ounces; of the leffer cardamon feeds, hufked, half an ounce; of faffron, two drachms; of white wine, a quart. Infuse three days without heat, and strain.

Tinctura sacra.

Take of focotorine aloes, eight ounces; of winter's bark, two ounces; of white wine, five quarts. Pulverize the aloes and bark feparately, then mix them and pour on the wine; infufe for a week or longer without heat, the glafs being often fhook; and laftly, ftrain the wine off.

It is convenient to mix fome clean white faud with the powders, that the aloes, when they become moift, may not cling into a lump.

Thebaic tincture.

Take of opium strained, two ounces; cinnamon, and cloves, of each a drachm; of white wine, a pint. Intuse without heat for a week; and then strain off the wine through paper.

The bitter tincture.

Take of gentian root, two ounces; of the outer yellow rind of Seville orange-peel, dried, one ounce; of the leffer cardamon feeds, hufked, half an ounce; of proof spirit, a quart. Digest without heat, and then strain.

ton of will want to the section of the states to a

Tinsture

Tinclure of antimony.

Take of any fixt alkaline falt, a pound; of antimony, half a pound; of rectified spirit of wine, a quart. Mix the antimony reduced to powder with the falt, and melt them together for an hour in a strong fire; then pour all out, and being pulverized, pour on the spirit of wine; digest for three or four days, and afterwards strain off.

Aromatic tin&ure.

Take of cinnamon, fix drachms; of the leffer cardamon feeds, hufked, three drachms; long pepper, ginger, of each two drachms; of proof fpirit, a quart. Digeft without heat, and ftrain the fpirit off.

Tincture of cantharides.

Take of cantharides, bruised, two drachms; of cochineal, half a drachm; of proof-spirit, a pint and an half. After digestion filtre the spirit through paper.

Tincture af cardamon seeds.

Take of the leffer cardamon feeds, freed from their hufks, half a pound; of proof-fpirit, a quart. Digeft without heat, and strain off the spirit.

Tinsture of castor.

Take of Ruffia caftor, powdered, two ounces; of proof-spirit, a quart. Digest for ten days without heat, and then strain the spirit off.

Tincture of cinnamon.

Take of cinnamon, an ounce and an half; of proof-spirit, a pint. Digest without heat, and strain the spirit off.

The simple tincture of the Peruvian bark.

Take of the Peruvian bark, four ounces; of prooffpirit, a quart. After digestion strain the spirit off.

Volatile tincture of the Peruvian bark.

Take of the Peruvian bark, four ounces; of fpirit of fal-armoniac, a quart. Digest without heat in a close vessel, and then strain the spirit off.

The fetid tincture.

Take of affafœtida, four ounces; of rectified spirit of wine, a quart. After digestion strain the spirit off.

Tincture of foot.

Take of wood-foot, two ounces; of affafœtida, one ounce; of proof-spirit, a quart. After digestion strain the spirit off.

Volatile tineture of gum-guaiacum.

Take of gum-guaiacum, four ounces; of the aromatic volatile fpirit, a pint and a half. Digest without heat in a well closed vessel, and then strain the spirit off.

Tineture of jalap.

Take of the root of jalap, eight ounces; of proof-fpirit, a quart. After digestion strain off the spirit.

Tincture of Japan earth.

Take of Japan earth, three ounces; of cinnamon, two ounces; of proof-spirit, a quart. After digestion strain the spirit off.

Tincture of martial flowers.

Take of martial flowers, four ounces; of proof-spirit, a pint. After digestion strain the spirit off.

K6

Tincture

Tinsture of iron in Spirit of Salt.

Take of the filings of iron, half a pound; of Glauber's spirit of sea-salt, three pounds; of rectified spirit of wine, three pints. Digest the filings in the spirit of salt without heat, as long as the spirit will work on them; then, after the fæces have subsided, evaporate the liquor poured off clear to one pound; and to this add the spirit of wine.

Tincture of black hellebore.

Take of the root of black hellebore, four ounces; of cochineal, two fcruples; of prooffpirit, a quart. After digeftion filtre through paper.

Tincture of myrrh.

Take of myrrh, three ounces; of proof-spirit, a quart. Digest them together, and then strain the spirit off.

Tinclure of rhubarb in Spirit.

Take of rhubarb, two ounces; of the leffer cardamon feeds, freed from their hufks, half an ounce; of faffron, two drachms; of proof fpirit, a quart. Digeft without heat, and firain the fpirit off.

The faturnine tincture.

Take fugar of lead, green vitriol, of each two ounces; of rectified spirit of wine, a quart. Reduce the falts separately to powder, and put them into the spirit; then digest without heat, and filtre the spirit through paper.

Tineture of Sena.

Take of floned raifins, fixteen ounces; of the leaves of fena, a pound; of carraway-feeds, an ounce and an half; of cardamon feeds, husked, half

half an ounce; of proof-spirit, a gallon.-Digest without heat, and strain off the spirit.

Tincture of Inake-root.

Take of Virginia snake-root, three ounces; of proof-spirit, a quart. Digest without heat, and strain off the spirit.

Stomachic tincture.

Take of ftoned raifins, four ounces; of cinnamon, half an ounce; caraway-feeds, the leffer cardamon feeds, freed from their hufks, and cochineal, of each two drachms; of proof fpirit, a quart. Digeft without heat, and ftrain off the fpirit.

The Aiptic tincture.

Take of calcined green vitriol, one drachm; of French brandy tinctured by the cafk, a quart. Mix them, that the fpirit may turn black, and then ftrain it off.

The simple tincture of valerian.

Take of wild valerian root, four ounces; of proof-spirit a quart. After digestion strain of the spirit.

Volatile tincture of valerian.

Take of the root of wild valerian, four ounces; of the volatile aromatic fpirit, a quart. Digeft them together in a close veffel without heat, and then ftrain the tincture off.

Tincture of white heilebore.

Take of the root of white hellebore, eight ounces; of proof-spirit, a quart. After digestion filtre through paper.

Balfam of guaiacum.

Take of gum-guaiacum, a pound; of balfam of Peru, three drachms; of rectified spirit of wine, two

3

two pints and an half. Digest them together, that the gum may be disfolved, and then strain off the spirit.

Vulnerary balfam.

Take of Benjamin, three ounces; of ftrained ftorax, two ounces; of balfam of Tolu, one ounce; of focotorine aloes, half an ounce; of rectified fpirit of wine, a quart. Digeft them together, till as much as may be of the gums are diffolved; then ftrain the tincture off.

Elixir of aloes.

Take of the tincture of myrrh, a quart; faffron, focotorine aloes, of each three ounces in weight. After digeftion ftrain off the spirit.

The Paregoric elixir.

Take flowers of Benjamin, opium, strained, of each a drachm; of camphire, two scruples; of the effential oil of anifeeds, half a drachm; of rectified spirit of wine, a quart. After digestion strain off the spirit.

Acid Elixir of vitriol.

Take of the aromatic tincture, a pint; of the ftrong fpirit or oil of vitriol, the weight of four ounces. Mix them gradually, and when the fæces are fublided, filtre through paper.

Dulcified elixir of vitriol.

Take of the aromatic tincture, a pint; of dulcified fpirit of vitriol, eight ounces in weight. Mix them.

The compound elixir of myrrh.

Take of the extract of favine, one ounce; of the tincture of caftor, a pint; of the tincture of myrrh, half a pint. After digeftion ftrain off the tincture.

The camphorated julap.

Take of camphor, one drachm; of double refined fugar, half an ounce; of boiling water, a pint. First grind the camphor, with a little rectified spirit of wine, till it is softened; then with the fugar, till they are perfectly united; lastly, add the water by degrees; and, when the mixture has stood in a covered vessel till it is cold, strain it off.

The chalk julap.

Take of the whiteft chalk, prepared, one ounce; of double refined fugar, fix drachms; of gum-arabic two drachms; of water, a quart. Mix all together.

The musk julap.

Take of damafk-role water, the measure of fix ounces; of musk, twelve grains; of double refined sugar, one drachm. Grind the musk and sugar together, and gradually add the role-water.

The common emulfion.

Take of fweet almonds, blanched, one ounce; of gum-arabic, half an ounce; of double refined fugar, fix drachms; of barley-water, a quart. Diffolve the gum in the barley-water hot, and when the water is quite cold, pour it gradually upon the almonds, pounded with the fugar, rubbing them together, that the liquor may grow milky; then ftrain it off.

Milk of gum-ammoniac.

Take of gum-ammoniac, two drachms; of fimple penny-royal water, half a pint. Rub the gum in a mortar with the water, till it is diffolved.

Camphorated .

Camphorated Spirit of wine.

Take of camphor, two ounces; of rectified spirit of wine, a quart. Mix them, that the camphor may be diffolved.

Syrups.

Wherever the weight of the fugar is not specified, it is to be understood, that to each pint of liquor are to be allowed twenty-nine ounces of fugar. The fugar should be double refined, reduced to powder, and melted in the heat of a balneum, unless it be ordered otherwise: and the syrup, as foon as made, is to be set by, till the next day, when any faccharine cruft, that may swim on the top, is to be taken off.

Syrup of garlic.

Take of the roots of garlic fliced, one pound; of boiling water, a quart. Steep the garlic in the water twelve hours in a close veffel, and in the liquor ftrained diffolve a fufficient quantity of fugar, fo as to make the fyrup.

Syrup of marshmallows.

Take of the fresh roots of marshmallows, a pound; of double refined sugar, four pounds; of water, one gallon. Boil the water with the roots, till it is half wasted; after it is quite cold, pour it off and press it out; let the liquor stand by for a night, that its faces may subside; in the morning pour off the clear, and adding the sugar, boil all down to the weight of fix pounds

Syrup of orange-peel.

Take of the outer yellow rind of fresh Seville orange-peel, eight ounces; of boiling water, five pints. Steep the peel in the water for a night in a close vessel, and in the morning dissolve in the liquor strained, of double refined sugar beaten to powder, as much as is sufficient to make a syrup.

Syrup of balfam.

Take of balfam of Tolu, eight ounces; of water, three pints. Boil the balfam in the water in a circulatory veffel, or at least in a matras with a tall neck, and the orifice lightly covered, for two or three hours. When the water is cold and strained off, add double refined fugar to make it into a syrup.

Syrup of clove july-flowers.

Take of clove july-flowers, fresh, and their heels cut off, three pounds; of boiling water, five pints. Steep the flowers in the water for a night in a vessel of glass, or of earth glazed; and in the liquor strained dissolved as much double refined sugar as is required to make a syrup.

After the fame manner is prepared the fyrup of cowflips.

Syrup of Saffron.

Take of faffron wine, a pint; of double refined fugar, twenty-five ounces; which diffolve in the wine, fo as to make a fyrup.

Syrup of quinces.

Take of the depurated juice of quinces, three pints; of cinnamon, one drachm; cloves, and ginger, of each half a drachm; of red wine, one pint; of double refined fugar, nine pounds. Digeft the juice with the aromatics fix hours in a heat of afhes; then add the wine, and ftrain the liquor off; and laftly add the fugar to make the fyrup.

Syrup of lemon-juice.

Take of lemon-juice, after it has ftood till its fæces are subfided, and it has been strained off,

off, a quart; of double refined fugar, fifty ounces. Diffolve the fugar in the juice fo as to make the fyrup.

After the fame manner are made the fyrups of mulberries and rafberries.

Syrup of white poppy-heads.

Take of the heads of dried white poppies without their feeds, three pounds and an half; of water, fix gallons. Slice the heads, and boil them in the water, often flirring them, that they may not burn, till about a third only of the liquor is left, which will be almost all imbibed by the poppy heads; then take all from the fire, and prefs the liquor flongly out from the heads: in the next place, boil the liquor by itself to about two quarts, and then ftrain it, while hot, first through a fieve, and then through a thin flannel: fet it by for a night, that what fæces have passed the ftrainers may fubfide; next morning pour off the clear liquor, and boil it with fix pounds of double refined fugar, till the whole comes to the weight of nine pounds, or a little more, that it may become a fyrup of a just confistence.

Syrup of wild poppies.

Take of the fresh flowers of wild poppies, four pounds; of boiling water, four pints and an half. Set the water poured on the flowers over the fire, and flir the flowers in, till they are all thoroughly wet; and, as soon as ever the flowers are sunk, let them steep for a night; next day pour off, and press out the liquor, setting it by for another night, that its seces may subside; then with a proper addition of double refined sugar make the syrup.

Pettoral

Pectoral Syrup.

Take of the leaves of English maiden-hair dried, five ounces; of liquorice, four ounces; of boiling water, five pints. Steep the ingredients for fome hours; and when the liquor is strained off, diffolve in it a proper quantity of double refined sugar to make a syrup.

Solutive Syrup of rofes.

Take the decoction left after the diffillation of fix pounds of damafk rofes; and five pounds of double refined fugar. Boil down the decoction preffed out to three pints, and fet it by for a night, that its faces may fubfide; next morning pour off the clear liquor, and adding the fugar, make it into a fyrup by boiling it away to the weight of feven pounds and a half.

Syrup of Squills.

Take of vinegar of fquills, a pint and an half; cinnamon, ginger, of each an ounce; of double refined fugar, three pounds and an half. Steep the fpices for three days in the vinegar, and, when ftrained, make the fyrup by adding the fugar.

The Simple Syrup.

Diffolve in any quantity of water, the proper weight of double refined fugar to make a fyrup.

Syrup of buckthorn.

Take of the juice of buckthorn-berries, ripe and fresh, one gallon; cinnamon, ginger, nutmeg, of each one ounce; of double refined sugar, seven pounds. Set the juice by, a few days, that its fæces may separate; then strain it, and in a small quantity of it infuse the spices. Boil down the the reft, towards the end adding that wherein the fpices have been infused, but strained from them, that the whole may be reduced to two quarts. Then add the sugar, and make the syrup.

Syrup of violets.

Take of violets, fresh and well coloured, two pounds; of boiling water, five pints. Steep the flowers a whole day in a glass, or earthen vessel glazed, then pour off the liquor, and strain it through a fine linen cloth, with caution not to press at all the flowers; afterwards, with a proper quantity of double refined sugar, make it into a syrup.

Syrup of ginger.

Take of ginger fliced thin, four ounces; of boiling water, three pints. Let the ginger fleep fome hours, and then ftrain off the liquor; to which add the proper quantity of double refined fugar to make a fyrup.

Confection of kermes.

Take of the juice of kermes, warmed and ftrained, three pounds; of damafk-rofe water, fix ounces in measure; of oil of cinnamon, half a fcruple; of double refined fugar; one pound. Melt the fugar by the heat of a balneum, into a fyrup with the rofe-water; then add the kermes juice, and, after it is cold, the oil of cinnamon.

Egyptian honey.

Take of verdigrease powdered very fine, five ounces; of honey, the weight of fourteen ounces; of vinegar, the measure of seven ounces. Boil all together over a gentle fire, till the mixture acquires a proper confistence and reddish colour; after a time, a groffer part will subside from this mixture;

mixture; the upper and more liquid part of which is called the Egyptian honey.

Honey of fluellin.

Take of the depurated juice of female fluellin, four pints; of clarified honey, four pounds. Boil them together to a proper confiftence.

Honey of hellebore.

Take of the roots of white hellebore dried and fliced, one pound; of clarified honey, three pounds; of water, four pints. After fleeping the roots three days in the water, boil them a little while; then boil the liquor well preffed out and flrained, with the honey, to a due confiftence.

Honey of roses.

Take of red-rofe buds quick dried, and their heels cut off, four ounces; of boiling water, three pints; of clarified honey, five pounds. Steep the rofes fome hours in water; then to the ftrained liquor add the honey, and boil to a proper confiftence.

Solutive honey.

Take the decoction remaining after the diftillation of fix pounds of damalk-roles; take alfo of cummin-feed a little bruifed, an ounce; of coarfe fugar, four pounds; of honey, two pounds. Boil the decoction preffed out to three pints, adding towards the end the feeds tied up in a cloth; then gently boil it with the fugar and honey into the confiftence of a liquid honey.

Oxymel with garlic.

Take of garlic fliced, an ounce and an half; carraway-feeds, fweet-fennel feeds, of each two drachms; of clarified honey, ten ounces; of vinegar, half a pint. Boil the vinegar a little while

in

in a glazed earthen veffel with the feeds bruifed; then add the garlic, and cover the veffel; after all is cold prefs out the liquor, and with the heat of a balneum diffolve it in the honey.

Oxymel of squills.

Take of clarified honey, three pounds; of vinegar of fquills, a quart. Boil them together in a glazed earthen veffel, with a gentle fire, to the confiftence of a fyrup.

Simple oxymel.

Take of clarified honey, two pounds; of vinegar, a pint. Boil them in a glazed earthen veffel, with a gentle fire, to the confiftence of a fyrup.

Powder against the bite of a mad dog.

Take of ash-coloured ground liverwort, two ounces; of black pepper, one ounce. Beat them together into a powder.

Compound powder of cuckow-pint.

Take of the root of cuckow-pint fresh dried, two ounces; the root of the yellow water-flag, the root of burnet faxifrage, of each one ounce; prepared crabs-eyes, cinnamon, of each half an ounce; of falt of wormwood, two drachms. Let all be beat into a powder, which must be kept in a very close vessel.

Compound powder of bole without opium.

Take of bole armeniac, or of French bole, half a pound; of cinnamon, four ounces; tormentil root, gum-arabic, of each three ounces; of long pepper, half an ounce. Make them into a powder.

Compound

Compound powder of bole with opium.

Take of opium ftrained, three drachms. Then let it be a little dried, that it may be commodioufly reduced to powder, and add it to the fpecies of the preceding composition, before they are pulverized, that they may be all beat together into a powder.

Compound powder of cerusse.

Take of ceruffe, five ounces; of farcocol, an ounce and an half; of gum-dragant, half an ounce. Make all into a powder.

Compound powder of crabs claws.

Take of the tips of crabs claws prepared, one pound; prepared pearls, red coral, prepared, of each three ounces. Mix all together.

Bezoardic powder.

Take of the compound powder of crabs claws, a pound; of oriental bezoar, prepared, an ounce. Make them together into a powder.

Compound powder of contrayerva.

Take of the compound powder of crabs claws, a pound and an half; of contrayerva root, five ounces. Make them into a powder.

Compound powder of myrrb.

Take the dried leaves of rue, dittany of Crete, myrrh, of each an ounce and an half; affafœtida, fagapenum, Ruffia caftor, opopanax, of each an ounce. Beat all together into a powder.

Compound powder of Scammony.

Take of fcammony, four ounces; of burnt hartfhorn, prepared, three ounces. Grind them carefully together into a powder.

Compound

Compound powder of Sena.

Take leaves of fena, chrystals of tartar, of each two ounces; of fcammony, half an ounce; cloves, cinnamon, ginger, of each two drachms. Powder the fcammony by itfelf, the reft all together, and then mix them.

Sneezing powder.

Take the dried leaves of afarabacca, of marjoram, of Syrian mastich thyme, dried lavender flowers, of each equal weights; and rub all into a powder.

Compound powder of amber.

Take prepared amber, gum-arabic, of each ten drachms; juice of the rape of ciftus, baluftines, japan earth, of each five drachms; of olibanum, half an ounce; of strained opium, a drachm. Reduce all into a powder.

Compound powder of gum-dragant.

Take of gum-dragant, gum-arabic, marfh-mallow root, of each an ounce and an half; ftarch, liquorice, of each half an ounce; of double refined fugar, three ounces. Reduce all together into a powder.

Hiera-picra.

Take of the gum extracted from focotorine aloes, one pound; of winter's bark, three ounces; powder them feparately, and then mix them.

Aromatic Species.

Take of cinnamon, two ounces; the leffer cardamom feeds freed from their hufks, ginger, long depper, of each one ounce. Make all into a powder by beating them together.

Species

Species of fcordium without opium.

Take of bole-armeniac, or of French bole, four ounces; of fcordium or water germander, two ounces; of cinnamon, an ounce and an half; ftorax ftrained, roots of tormentil, biftort, gentian, leaves of dittany of Crete, galbanum ftrained, gum-arabic, and red rofes, of each one ounce; long pepper, and ginger, of each an ounce. Beat all into a powder.

Species of scordium with opium.

Take of strained opium three drachms; and add it to the former species, while they are pounding together, it being first a little dried, that it may the more commodiously be beaten to powder.

Sugar of roses.

Take of red rofe-buds, quick dried, and their white heels cut off, one ounce; of double refined fugar, a pound. Reduce the rofes and fugar to powder feparately; then mix them, and with a little water form lozenges to be dried with a gentle heat.

White pectoral troches.

Take of double refined fugar, a pound and an half; of ftarch, an ounce and a half; of liquorice, fix drachms; of Florentine orris, half an ounce. All the ingredients being reduced to powder, with the mucilage of gum-tragacanth form troches.

Black pectoral troches.

Take extract of liquorice, and double refined fugar, of each ten ounces; of gum-dragant, half a pound. By moistening with water make troches.

Troches

and the

218

Troches of nitre.

Take of purified nitre, four ounces; of double refined fugar, a pound. Make them into troches with the mucilage of gum-tragacanth.

Troches of Squills.

Take of baked fquills, half a pound; of wheatflower, four ounces. Pound them together, and form them into troches to be dried with a fmall heat.

Troches of Julphur.

Take of washed flowers of fulphur, two ounces; of double refined fugar, four ounces. Beat them together, and form troches, by gradually adding the mucilage of quince-feeds.

Troches of Japan earth.

Take Japan earth, and gum-arabic, of each two ounces; of fugar of rofes, fixteen ounces. Beat them together, and with a little water make into troches.

Cardialgic lozenges.

Take of prepared chalk, four ounces; of prepared grabs claws, two ounces; of bole-armeniac, or French bole, half an ounce; of nutmeg, a fcruple; of double refined fugar, three ounces. Make all into a powder, and then with a little water form it into lozenges.

Aromatic pills.

Take of focotorine aloes, an ounce and an half; of gum-guaiacum, an ounce; the aromatic fpecies, and balfam of Peru, of each half an ounce. Let the aloes and gum-guaiacum be powdered feparately, then mixt with the reft, and formed into a mafs with the fyrup of orange-peel.

The more simple pills of coloquintida.

Take the pith of coloquintida, and scammony, of each two ounces; of oil of cloves, two drachms. Let the dry species be reduced to powder separately, the oil be mixt with them, and the whole be formed into a mass with syrup of buckthorn.

Pills of coloquintida with aloes.

Take focotorine aloes and fcammony, of each two ounces; of the pith of coloquintida, one ounce; of oil of cloves, two drachms. Let the dry fpecies be reduced to powder feparately, the oil mixt among them, and the whole formed into a mass with fyrup of buckthorn.

Deobstruent pills.

Take of the aromatic pill, three ounces; rhubarb, extract of gentian, and falt of iron, of each one ounce; of falt of wormwood, half an ounce. With folutive fyrup of rofes beat them diligently into a mafs.

Gum pills.

Take galbanum, opopanax, myrrh, and fagapenum, of each an ounce; of affafœtida, half an ounce. With fyrup of faffron make them into a mafs.

Mercurial pills.

Take of quickfilver, five drachms; of Strafbourgh turpentine, two drachms; of the cathartic extract, four fcruples; of rhubarb in powder, one drachm. Firft grind the quickfilver with the turpentine; till it appear no longer; then beat them up with the reft into a mass. If the turpentine chance to be too thick, it is to be thinned with a little oil of olives.

1.2

Rufus's

Rufus's pills.

Take of focotorine aloes, two ounces; myrrh, and faffron, of each one ounce. Make them into a mass with fyrup of faffron. of cach two o

Soap pills.

Take of almond foap, four ounces; of strained opium, half an ounce; of effence of lemons, a drachm. Beat the opium foftened with a little wine, along with the reft of the ingredients, till they are perfectly mixt.

Storax pills.

Take of ftrained ftorax, two ounces; of faffron one ounce; of strained opium, five drachms. Beat them diligently together, till they are per-Electuary of bay-berries. fectly mixed.

Take the leaves of rue dried, carraway-feeds, common parfley-feeds, and bay-berries, of each an ounce; of fagapenum, half an ounce; black pepper, and Ruffia caftor, of each two drachms; of clarified honey, thrice the weight of the species, when powdered. Mix the species with the honey into an electuary.

Electuary of caffia.

Take solutive fyrup of roses, and pulp of cassia fresh extracted, of each half a pound ; of manna, two ounces; of the pulp of tamarinds, one ounce. Rub the manna in a mortar, and with a very gentle heat diffolve it in the fyrup; then add the pulps, and the heat being continued, reduce the whole to a proper confiftence.

Lenitive electuary.

Take of dried figs, one pound; of the leaves of sena, eight ounces; the pulps of tamarinds, ot

of caffia, and of French prunes, of each half a pound; of coriander feeds, four ounces; of liquorice, three ounces; of double refined fugar, two pounds and an half. Reduce the fena with the coriander feeds to powder, and feparate by the fieve ten ounces; boil the reft with the figs and liquorice, in two quarts of water, till it is boiled half away; then ftrain and prefs it out; let the ftrained liquor be evaporated to the weight of a pound and an half, or a little lefs; afterwards add the fugar to make a fyrup; this fyrup mix gradually with the pulps; and laftly ftir in the powder before feparated by the fieve.

Electuary of Scammony.

Take of fcammony, an ounce and an half; c'oves, and ginger, of each fix drachms; of the effential oil of caraway feeds, half a drachm; of honey, half a pound. Reduce the fcammony to powder by itfelf; mix the aromatics, first pounded together, with the honey; then add the fcammony, and in the last place the oil.

Electuary of Scordium ..

Take of any quantity of the species of scordium, or water-germander, with opium; and thrice their weight of diacodion, boiled to the thickness of honey. Mix the species with the syrup into an electuary.

Locatellus's balfam.

Take of oil olive, a pint; Strafburgh turpentine, and yellow wax, of each half a pound; of red fanders, fix drachms. Melt the wax with fome part of the oil over a gentle fire, then add the reft of the oil, and the turpentine; in the laft place mix in the fanders, and ftir the whole well together, till it is nearly cold,

The:

The cordial confection.

Take fresh rosemary tops, and juniper berries, of each a pound; the lesser cardamom feeds freed from their husks, zedoary, and fassron, of each half a pound. Draw a tincture with about a gallon and an half of proof spirit; reduce by a gentle heat, this tincture strained, nearly to the weight of two pounds and an half; then finish the electuary by adding the following species very finely powdered; viz. of the compound powder of crabs claws, fixteen ounces; cinnamon, and nutmegs, of each two ounces; of cloves, an ounce; of double refined sugar, two pounds.

The confection called paulina.

Take coftus, or in its flead zedoary, cinnamon, long pepper, black pepper, ftrained ftorax, ftrained galbanum, ftrained opium, and Ruffia caftor, of each two ounces; of the fimple fyrup boiled to the confiftence of honey, an equal weight to thrice the fpecies. Mix carefully the opium, first diffolved in wine, with the fyrup, warmed; then to the ftorax and galbanum melted together, add by degees the fyrup, while it remains warm; afterwards the in the other fpecies reduced to powder.

Mitbridate.

Take of cinnamon, fourteen drachms; of myrrh, eleven drachms; agaric, fpikenard, ginger, faffron, feeds of treacle muftard, or of mithridate muftard, frankincenfe, and chio turpentine, of each ten drachms; camel's hay, coftus, or in its ftead zedoary, Indian leaf, or in its ftead mace, French lavender, long pepper, feeds of hartwort, juice of the rape of ciftus, ftrained ftorax, opopanax, ftrained galbanum, balfam of Gilead,

or in its stead, expressed oil of nutmegs, and Russia castor, of each an ounce; poley mountain, water-germander, the fruit of the balfamtree, or in its stead cubebs, white pepper, seeds of the carrot of Crete, and bdellium ftrained, of each feven drachms; celtic nard, gentian root, leaves of dittany of Crete, red rofes, feeds of Macedonian parsley, the lesser cardamon feeds, freed from their hufks, fweet fennel feeds, gum-arabic, and opium ftrained, of each five drachms ; root of the fweet flag, root of wild valerian, anife-feed, and fagapenum ftrained, of each three drachms; fpignel, St. John's wort, juice of acacia, or in its ftead Japan earth, the bellies of feinks, of each two drachms and an half; of clarified honey, thrice the weight of all the reft. Diffolve the opium first in a little wine, and then mix it with the honey made hot; in the mean time melt together in another veffel, the galbanum, ftorax, turpentine, and the balfam of Gilead, or the expressed oil of nutmeg, continually ftirring them round, that they may not burn; and as foon as these are melted, add to them the hot honey, first by spoonfuls, and afterwards more freely: laftly, when this mixture is near cold, add by degrees the reft of the fpecies reduced to powder.

The London philonium.

Take white pepper, ginger, carraway feeds, of each two ounces; of opium firained, fix drachms; of diacodion boiled to the confiftence of honey, thrice the weight of all the reft. Mix carefully the opium, diffolved first in wine, with the fyrup warmed, and then add the other species reduced to powded.

L4

Venice

Venice treacle.

Take of the troches of fquills, half a pound ; long pepper, opium strained, and dried vipers, of each three ounces; cinnamon, balfam of Gilead, or in its stead, expressed oil of nutmeg, of each two ounces; agaric, the root of Florentine orrice, water-germander, red rofes, feeds of navew, and extract of liquorice, of each an ounce and an half; fpikenard, saffion, amomum, myrrh, coftus, or in its flead zedoary, and camel's hay, of each an ounce; the root of cinquesoil, rhubarb, ginger, Indian leaf, er in its stead mace, leaves of dittany of Crete, of horehound, and of calamint, French lavender, black pepper, feeds of Macedonian parfley, olibanum, Chio turpentine, root of wild valerian, of each fix drachms; gentian root, Celtic nard, spignal, leaves of poley mountain, of St. John's wort, of ground-pine, tops of creeping germander with the feed, the fruit of the balfam tree, or in its stead cubebs, anife-feeds, sweet sennel feeds, the lesser cardamon feeds, freed from their hufks, feed of hishop's weed, of hartwort, of treacle mustard, or mithridate muftard, juice of the rape of ciftus, acacia, or in its stead Japan earth, gum-arabic, storax itrained, sagapenum strained, Lemnian earth, or in its ftead bole-armeniac or French bole, and green vitriol calcined, of each half an ounce; roots of creeping birthwort, or in its fread of the long birthworr, tops of the leffer centaury, feeds of the carrot of Crete, opopanax, galbanum strained, Russia castor, Jew's pitch, or in its stead white amber prepared, and root of the fweet flag, of each two drachms; of clarified honey, thrice the weight of all the reft. The ingredients are to be mixed in the fame manner as in the mithridate.

Bate's alum water.

Take alum, and white vitriol, of each half an ounce; of water, a quart. Diffolve the falts by boiling them in the water, and, when the fæces have fubfided, filtre the liquor through paper.

The Sapphire coloured water.

Take of lime-water, a pint; of fal-ammoniac one drachm. Let them fland together in a copper veffel, or with a few bits of copper, till the water has acquired a blue fapphire colour.

The blue vitriolic water.

Take of blue vitriol, three ounces; alum, and the firong fpirit or oil of vitriol, of each two ounces; of water, a pint and an half. Boil the falts in the water, till they are diffolved; then add the oil of vitriol, and ftrain the mixture through paper.

Camphorated vitriolic water.

Take of white vitriol, half an ounce; of camphor, two drachms; of boiling water, a quart. Mix them, that the vitriol may be diffolved; and after the fæces have fubfided, filtre the water through paper.

The Saponaceous lotion.

Take of damafk rofe water, three quarters of a pint; of oil olive, a quarter of a pint; of the ley of tartar, the measure of half an ounce. Rub the ley of tartar and oil together, till they are mixed; then gradually add the water.

Oil of St. John's wort.

Take of the flowers of St. John's wort full blown, fresh, and carefully picked from their calyxes, four ounces; of oil olive, a quart. The

oil

226

oil being poured on the flowers, let them fland together, till the oil is fufficiently tinged.

Oil of mucilages.

Take of the roots of marshmallows fresh, half a pound; linseed, and senugreek feeds, of each three ounces; of water, a quart; of oil olive, two quarts. Boil gently the roots and feeds bruised, in the water for half an hour; afterwards add the oil, and renew the boiling, till the water is quite wasted; then pour the oil cautiously off.

Oil of Elder.

Take of elder flowers, one pound; of oil olive, a quart. Boil the flowers in the oil, till they are almost crisp; then press out the oil, and set it by, that the fæces may subfide.

Green oil.

Take bay-leaves, leaves of rue, of marjoram, of fea wormwood, and of chamomile, of each three ounces; of oil olive, a quart. The herbs being bruifed boil them lightly in the oil, till they are become crifp, then prefs out the oil; and after the fæces have fubfided, pour it off.

The ammoniac plaister with quickfilver.

Take of gum-ammoniac ftrained, a pound; of quickfilver, three ounces; of the fimple balfam of fulphur, a drachm. Rub the quickfilver with the balfam of fulphur, till it no longer appears; then add by degrees the gum-ammoniac melted, a little before it is cold, and mix them carefully.

Drawing plaiser.

Take yellow rozin, and yellow wax, of each three pounds; of tried mutton fewet, one pound. Melt all together, and firain the mixture while it remains fluid.

Cephalic plaister.

Take of Burgundy pitch, two pounds; of foft labdanum, one pound; yellow rozin, and yellow wax, of each four ounces; of the expressed oil of mace, one ounce. The pitch, rozin, and wax, being melted together, add first the labdanum, and then the oil of mace.

The common plaister.

Take of oil olive, one gallon; of litharge finely powdered, five pounds. Boil them together with about a quart of water over a gentle fire, continually flirring, till the oil and litharge are united, and they acquire the due confiftence of a plaister; and if the water is wasted, before the operation is over, more water must be poured on hot.

The common Sticking plaister.

Take of the common plaister, three pounds; of yellow rozin, half a pound. Throw the rozin, first reduced to powder, that it may the sooner melt, into the common plaister melted with a very gentle heat, and stir them well together.

The common plaister with gums.

Take of the common plaifter, three pounds; of galbanum strained, eight ounces; common turpentine, and frankincense, of each three ounces. To the galbanum and turpentine melted together with a gentle heat, sprinkle in the frankincense reduced to powder; and then gradually add to them the plaister first melted likewise with a very gentle heat.

The common plaister with quickfilver.

Take of the common plaister, one pound; of quickfilver, three ounces; of the fimple balfam of L 6 fulphur,

fulphur, a drachm. Mix them together after the fame manner as in the ammoniac plaister with quickfilver.

The cummin plaister.

Take of Burgundy pitch, three pounds; yellow wax, cummin feeds, carraway feeds, and bayberries, of each three ounces. The pitch and wax being melted together, fprinkle into them the reft reduced to powder, and ftir all well together.

Red lead plaister.

Take oil olive, two quarts; of red-lead finely powdered, two pounds and an half. This plaifter is to be prepared in the fame manner as the common plaifter; only here more water is required, and more caution, that the plaifter may not be burnt and turn black.

Plaister of mucilages.

Take of yellow wax, forty ounces; of the oil of mucilages, eight ounces in measure; of gumammoniac strained, half a pound; of common turpentine, two ounces. The gum-ammoniac being melted with the turpentine, add to them gradually the wax melted with the oil in another vessel.

Strengthening plaister.

Take of the common plaister, two pounds; of frankincense, half a pound; of dragon's blood, three ounces. To the common plaister melted add the rest reduced to powder.

Soap plaiser.

Take of the common plaister, three pounds; of hard foap, half a pound. To the common plaister liquified add the foap, then melt all to the confistence of a plaister, and take particular care, that it


226

AN

it does not grow too cold, before it is formed into rolls.

Stomach plaister.

Take of foft labdanum, three ounces; of frankincenfe, one ounce; cinnamon, and expressed oil of mace, so called, of each half an ounce; of effential oil of mint, one drachm. Add to the frankincense melted, first the labdanum a little heated, till it is become soft, and then the oil of mace; afterwards mix in the cinnamon with the oil of mint; and beat them together in a warm mortar into a mass, which is to be kept in a vessel well closed.



INDEX

OF

er Bart A.A.Powelere Stantorald

DISEASES

AND THEIR

REMEDIES.

A

A BORTION to prevent. Decoction of Oak Batk, 4. Strengthning Pills, 16. Strengthning Plaister, 50. Bark Electuary, 91. Aluminous Powder, 127.

ABSCESS to cleanse. Yellow Basilicon, 32. Detergent Ointment, 83, and various others. Digestive Balsam, 22. Common Fomentation, 82. Turner's Cerate, 133.

ACHES from cold. Discutient Liniment, 26, and 27. Saponaceous Liniment, 61. Guaiacum Bolus, 2. Ischiatic Plaister, 7. Volatile Tincture of Guaiacum, 73. Turpentine Bolus, 89.

ACIDITIES in the first passages. Magnefia Alba, 13. White Drink, 47. Chalk Julap, 58. Teftaceous ceous Powder, 70. Troches for the Heart-burn, 72. All fixt Alkaline Salts and Powders. Chamomile Bolus, 40.

AFTER-PAINS. Alexiterial Bolus with Caftor, 37. Balfamic Bolus, 39. Mithridate and Venice Treacle Julaps, 59. Alexiterial Powder, 69. Cordial Boluffes, 86. Hyfteric Julap, 100. Mithridate, and Sudorific Draughts, 120. Penny-royal Julap, 124.

APPETITE, Want of. Stomachic Bolus, 2. Bitter Infusions various, 56, 154. Tincture of Rhubarb, 74. Elixir of Vitriol, 92. Chamomile Infusion, 99. Tincture of Hiera-picra, 132.

ASTHMA. Ammoniacum Mixture, 14, 60 and 62. Squill Pills, 17, 65, and 127 Squill Bolus, 44. Expression of Millepedes, 53, 152. Diuretic Decoction, 148. Asthmatic Mixture, 102. Garlic Infusion, 123.

AGUE See Intermitting Fever.

- St. ANTHONY's Fire. Emollient and Poppy Fomentations, 8. Anodyne Cataplasm, 23. Limewater, 79. Emollient Pultice, 81. Refrigerant Ointment, 108. And all cooling Purges.
- APOPLEXY. Sinapifms, 19 and 23. Epileptic Bolus, 41. Blifter Plaifter, 117. Volatile Julap, 59. Preparations of Valerian, 76 and 99. Nervous Draught, 98. Purging Clyfter, 117. And all Volatiles.

MTROPHY: Aromatic Electuary, 5. Saponaceous Bolus with Rhubarb, 44. Elixir of Aloes, 50. Bitter Infusions, and Tinctures, 56 and 75. Ecphractic Pills, 66. Tincture of Rhubarb, 107. Tincture of Hiera-picra, 132.

BALD-

TI.

INDEX.

BALDNESS. Tincture of Cantharides, 72.

BELLY-ACH, Dry. Infusion of Sena, and Tincture of ditto, 57 and 132. Fomentations, 8, 82, 95, 118. Clysters, 8, 52, 94, 117.

BITE of a mad Dog. Dr. Mead's Powder, 105. Musk Bolus, 88, 146.

BLISTERING PLAISTER, 51, 117, 139.

BLOOD, Spitting of. Vulnerary Decoction, 49. Tincture of Roles, 75. Saturnine ditto, 75. Bark Electuary, 115. Balfamic Electuary, 116.

BRUISES, Internal. Capivi, and Balfam of Peru Mixtures, 17. Balfamic Bolus, 39. Vulnerary Decoction, 49. Balfamic Bolus, 85. Oily Draught, 97.

BRUISES, External. Saponaceous Cerate, 25. Ditto Liniment, 61.

BILIOUS Cholic. Vide Belly-Ach dry.

BUBOES. Suppurative Cataplasm, 23. Mercurial Plaister, 82. Specific Pills, 141.

BURNS and Scalds. Turner's Cerate, 133. Ointment of Elder, 21. Camphorated Spirits of Wine. Brown Ointment, 143.

C

CACHEXY. Aromatic Electuary with Steel, 5. Steel Bolus, 39. Hiera-picra, 55. Bitter Infusion, 56. Elixir of Vitriol, 116. Tincture of Rhubarb, 132. Chalybeate Milk, 156. Golden Tincture, 131.

CANCER.

CANCER. Hemlock Fomentation, Pultice, and Extract, 80, 82.

iv

CARBUNCLES. Suppurative Cataplasm, 23. Green Cerate, 25. Emollient Pultice, 81. Refrigerant Ointment, 108. Yellow Basilicon, 132.

- CATALEPSY. Sneezing Powder, 18. Epileptic Bolus, 41. Blifters, 51. Tinctures of Valerian, 76, 139. Emetic Draught, 96.
- CATARRHS. Pectoral Decoction, 4. Oily Draughts, 10. Linctuffes, 12. Sperma Ceti Mixture, 15, 125, 141, 154. Balfamic Bolus, 39. Oily Emulfion, 51, 151. Oily Draughts, 55. Pectoral Linctus, 60, 124, 156. Balfamic Pills, 65. Balfamic Bolus, 85. Pectoral Bolus, 88. Pectoral Decoction, 90. Pectoral Mixture, 102. Pills for a Cough, 103. Elecampane Electuary, 114. Pectoral Syrup, 142. Syrup of Elder, 161.
- CATHARTICS. 7, 10, 17, 19, 43, 49, 55, 74, 85, 88, 97, 105, 117, 121, 129.
- CHAPS. Yellow Cerate, 81. Refrigerant Ointment, 108.
- CHLOROSIS. Steel Bolus, 39. Ecphractic Pills, 66, and 67, 105. Tincture of Hellebore, 74, 106. Chalybeate Electuary, 91, 114. Chalybeate Pills, 126. Golden Tincture, 131.

CLYSTERS. 8, 52, 94, 117, 118, 139, 149, 151.

COLDS. Mindererus's Spirit, 130, 161.

CHOLIC. Rhubarb Bolus, 2, 44, 111, 147. Various Clyfters, 8, 52, 94, 117. Tincture of Rhubarb, 74* 74, 107. Stomachic Electuary, 92. Anodyne Draught, 119, 153. Tinctures of fena, and Hierapicra, 132. Solutive Anodyne Bolus, 145, Anodyne Fettd Bolus, 146.

- CONSUMPTIONS. All Pectorals and Balfamics. Vide Catarrins. Balfam of Capivi, and Peru Mixtures, 17. Vulnerary Decoction, 49. Febrifuge Decoction, 90. Elixir of Vitriol, 93. Reftringent Milk, 100, 140, 156. Conferve of Rofes, 89. Chalk Julap, 124. Decoction of Chalk, and Hartshorn Drink, 112.
- CONVULSIONS. Tincture of Wood Soot, 19, 33. Epileptic Bolus, 41, 87. Volatile Julap, 56. Cinnabar Pills, 65. Tincture of Valerian, 96. Mufk Bolus, 88, 146. Galbanum Plaister, 51.
- CORPULENCY. Expression of Hog-lice, 53. Diuretic Salt, 70. Medicated Ale, 111. Ecphractic Pills, 16. Saponaceous Pills, 127. And all Acids.
- COSTIVENESS. Opening Electuary, 5. Laxative Infusion, 10, 57. Saponaceous Bolus, 44. Tincture of Hiera-picra, 132. Magnefia Alba, 13. Purging Draught, 142, 157. Solutive Linctus, 156.

COUGHS. Vide Catarrhs.

650%

- CRAMPS. Saponaceous Liniment, 61. Ointment of Marshmallows, 77. Neat's Foot Oil, Hungary Water, &c.
- CRAPULA. Emetic Draught, 96. Infusion of Mint, 10. Of Chamomile, 99. Bitter Infusions, 56, 154. Laxative ditto, 10, 57. Tincture of Hiera-picra, 132.

CUTANEOUS Diforders. Ointment for the Leprofy, 20. Yellow ditto, Lead ditto, 33. Scorbutic Juices, 71. Precipitated Sulphur, ibid. Decoction of Elm Bark, 113. Cleanfing Bolus, 40. Buck Bean Infution, 57. Sulphur Lotion, 62. Antimonial Wine, 76. Lime-water, 79. Dietetic Docoction, 90, 148. Quickfilver Pills, 103. Vitriolic Ointment, 133. Antimonial Powder, 105, 160. Plummer's alterative Powder, 129. Mercurial Electuary, 138. Red Pill, 141. Æthiopic Pills, 158. Antimonial Pills, 158.

D

DEAFNESS. Mixture for, 62, 101.

- DELIVERI, to promote. Alexiterial Bolus with Caftor, 37. Anodyne Pills, 67. Myrrh Bolus, 88. Nervous Draught, 98. Borax Bolus, 145.
- DECAYS, Nervous. Mustard Whey, 19. Steel Bolus, 39, 154. Bitter Infusions, 56. Tincture of Bark, 73. Chalybeate Electuary, 114. Emetic Draught, 96.
- DEFLUXIONS. Anodyne Draught, 9. Anodyne Pill, 16. Pacific Powder, 18. Squill Bolus, 44. Expression of Hog-lice, 53. Pills for a Catarrh, 65. Pills for a Cough, 103. Purging Draught, 121.
- DIABETES. Aqua Liberans, 36. Alum Bolus, 38. Lime Water, 110. Bark Electuary, 115. Alum Whey, 19, 70. Elixir of Vitriol, 116.

DIGESTION, to promote. Chamomile Bolus, 39. Pepper Bolus, 43. Saponaceous Bolus with Hierar picra, 44. Elixir of Aloes, 50. Bitter Infusions, 56, 154. Digestive Powder, 69. Stomachic Electuary, 92. Chamomile Infusion, 99. Stomachic Pills, 104. 104. Bates's Pepper Electuary, 115. Elixir of Vitriol, 116. Golden Tincture, 131. Tincture of Hiera-picra, 132. Hiera-picra Pills with Steel, 159.

DIGESTIVES. Digeftive Balfam, 22. Yellow Bafilicon, 32. Digeftive Ointments, 83, 84, 107, 132, 134, 143.

- DIARRHOEA. Rhubarb Bolus, z. Aftringent Electuary, 5, 137, 148. Ipecacoanha Bolus, 146. Extract of the Brazilian Bark, 152. Aluminous Bolus, 38. Aftringent Bolus, ibid. Aftringent Drink, 47. Decoction of Simarouba, 49. Jelly of Starch, 54. Reftringent Decoction, 90. Emetic Draught, 96. Reftringent Mixture, 102, 140, 157. Decoction of Chalk, 112. Reftringent Clyfters, 118. Aluminous Powder, 127. Aftringent Decoction, 136, 147. Arabic Decoction, 136. Chalk, 145. Aftringent Powders, 161.
- DROPSY. Hydragogue Electuary, 6. Hydragogue Julap, 11. Diuretic Lixivium, 13. Mustard Whey, 19. Jalap Bolus, 43. Squill Bolus, 44. Expression of Hog-lice, 53 152. Hydragogue Draught, 54. Purging Bolus, 88. Diuretic Draught, 96. Chamomile Infusion, 99. Mustard Electuary, 115, Gamboge Bolus, 146. Diuretic Infusion, 155.
- DYSENTERY. Anodyne Clyflers, 8, 52, 139. Rhubarb Bolus, 44. White Drink, 47. Jelly of Starch, 54. Emetic Draught, 96, Decoction of Chalk, 112. Oily Draught with Rhubarb, 120. Cerated Glafs of Antimony, 131. Ipecacoanha Bolus, 146. Reftringent Decoction, 147.

DYSURY. Barley Water, 36. Nephritic Mucilage, 64. Diuretic Powder, 98. Oily Draught, 97. Refrigerant Draught, 98. Scarlet Decoction, 112. Nitrous Powders, 129, 142. Camphor Bolus, 145. Arabic Emulfion, 151.

EMETICS.

E

SEMICE CONTRACT & BOOM & STREET

Triok red. Colde

Bates's Perman E

EMETICS. 45, 63, 88, 96, 145, 143.

EPILEPSY. Epileptic Bolus, 41. Volatile Julap, 59. Cinnabar Pills, 65. Tinclure of Valerian, 76. Epileptic Draught, 96, 99. Nervous Draught, 98. Valerian Electuary, 115. Sagapenum Pills, 126. Misletoe Mixture, 158. Valerian Mixture, 158.

E.

ERUCTATIONS. Stomachic Bolus, 2. Aromatic Species, 19. Pepper Bolus, 43. Stomachic Electuary, 92. Wormwood Electuary, 113. Pepper Electuary, 115:

- ESCHAROTICS. Troches of Red Lead, 142.
- EXCORIATIONS. Yellow Cerate, 25. Refrigerant Ointment, 108. Turner's Cerate, 133. Brown Ointment, 143.
- EXPECTOR ATION, to promote. Pectoral Decoction, 4. Infusion of Cascarilla, 10. Expectorant Mixture, 14. Squill Bolus, 44. Ammoniacum Mixtures, 60. Althmatic Mixture, 62. Pectoral Mixture, 102. Elecampane Electuary, 142. Pectoral Pills, 126. Pectoral Syrup, 142. Storax Pills, 160. Balfamic Electuary, 149. Balfamic Emulsion, 151. Squill Mixture, 157.

EYES, Diseases of. All Purgatives. Eye Waters, 22, 135, 110. Opthalmic Pultice, 24. Sky coloured Collyrium, 25. Opthalmic Epithem, 26. Opthalmic Fomentation, 27, 46. Opthalmic Ointments, 32, 78, 108, 133.

FAINTING. Cordial Julap, 58. Tincture of Wood-foot, 131. And all Volatiles.

F

FEVERS, Inflammatory. Saline Mixture, 15. Cordial Julap acidulated, 58. Common Clyfter, 94. Emetic Draught, 96. Scarlet Decoction, 112. Saline Draught, 121. Salprunel Powder, 129, 142. Mindererus's Spirit, 130. Diuretic and Refrigerant Apozem, 144. Nitrous Mixture, 157.

Nervous. Alexiterial Bolus's, 37. Mufk Bolus, 88, 146. Cordial Julaps, 58, 59. Cordial Mixture, 63. Cordial Bolus's, 86. Nervous Draught, 98. Alexiterial Powder, 104. Alexipharmic Tincture, 106. Camphorated Draught, 120. Diaphoretic Powder, 128. Blifters. Contrayerva Bolus, 147.

Putrid. Camphorated Draught, 120. Infusion of Snake-Root with Vinegar, 155. Cordial Julap acidulated, 58. Musk Julap, 59. Bark Decoction, 148;

Intermitting. Decoction of the Bark, 4, 48, 115, 136, 148. Compound Chamomile Bolus, 40. Febrifuge Bolus's, 41, 42. Febrifuge Mixture, 63, 157. Tincture of the Bark, 73. Bark Electuary, 91. Elixir of Vitriol, 92. Chalybeate Electuary, 114. Tincture of Hiera-picra, 132. Saline Mixture 15, 141. Aluminous Bolus, 144.

FLATULENCY. Stomachic Bolus, 2. Aromatic Electuary, 5. Purging Bitter Infusion, 56. Aromatic Species, 71. Tincture of Rhubarb, 74. Stomachic Pills, 104. Stomachic Electuary, 139. Wormwood Electuary, 113. Bate's Pepper Electuary, 115. And all Carminatives.

FOMENTATIONS. 27, 53, 82, 95, 118.

FLUOR Albus. Capivi Mixture, 17, 159. Strengthening Pills, 16, 66. Injection, 28. Reftringent Pills, 29, 30, 31, 159. Aqua Liberans, 36. Strengthening Bolus, 40. Tincture of Rhubarb, 74. Oliba. num Draught, 97. Tincture of Japan Earth, 107: Strengthening Electuary, 138. Olibanum Bolus, 147.

FUNDAMENT, Bearing down of. Powder for, 69+

G

GANGLES. Decoclion of Elm Bark, 5, 54, 95, 119, 140, 152, 153.

GANGRENE. See Mortification.

GLEET. Injection, 28. See Fluor Albus.

- GONORRHOEA, Virulent. Specific Electuary, 7, 137, 138. Laxative Infusion, 10. Purging Pills with Mercury, 17. Mercurial Pills, 29. Calomel Bolus, 39, 63, 67, 126. Bolus for a Gonorrhœa, 42. Quickfilver Pills, 103. Arabic Injection, and others, 123. Turpentine Pills, 127. And all cooling Purges, particularly the Compound Lenitive Electuary, 115.
- GOUT. Guaiacum Bolus for the Gout, 2, 32, 42. Bitter Infusion, 56 154. Muslard Whey, 70. Volatile Tincture of Guaiacum, 73. Turpentine Bolus, 89. Anodyne Embrocation, 116. Bitter Infusion, 122. Dr. Dover's Powder, 128.
- GRAVEL. Saponaceous Bolus, 2, 44. Turpentine Bolus, 3. Nephritic Electuary, 7. Turpentine Clyfter, 8, 52. Oily Draught with Manna, 10, 55. Capivi Mixture, 17. Laxative Infusion, 57. Nephritic Mucilage, 64. Nephritic Draught, 96.

GRIPES. See Cholic.

HARDNESS

HARDNESS of any part. Oleaginous Liniment, 13. Galbanum Plaister, 51.

HOEMOPTOE. See Blood, spitting of.

HOEMORRHAGE, to restrain. Acid Elixir, 50. Tincture of Roses, 75. Stiptic Tincture, ibid. Scarlet Decoction, 112. Elixir of Vitriol, 116.

HIP-Gout. Bolus for the Gout, 38, 89. Sinapifm, 19. Difcutient Liniment, 26. Ifchiatic Plaister, 7, 51, 93. Mustard Whey, 70. Tinctures of Guaiacum, 73. Anodyne Difcutient Plaister, 150.

- HYPOCHONDRIACAL Distempers. Aromatic Electuary with Steel, 5. Ecphractic Pills, 16, 66. Bitter Infusions, 56. Tincture of Aloes, 75. Chalybeate Electuary, 91, 114, 149 Elixir of Vitriol, 92, 116. Infusion of Valerian, 99. Stcmachic Pills, 104. Tincture of Wood Soot, 131. And all Preparations of the Peruvian Bark.
- HYSTERIC Passion. Myrrh Electuary, 6. Hysteric Mixture, 14. Asside Pills, 16, 126. Tincture of Wood Soot, 19, 131. Galbanum Plaister, 51. Musk Julap, 59. Asside Milk, 60. Tincture of Steel, 74. Myrrh Bolus, 88. Infusion of Valerian, 99. Hysteric Julap, 100. Hysteric Pills, 104. Valerian Draught, 120. Epileptic Draught, 122. Penny-royal Julap, 124. Sagapenum Pills, 126.
- HOARSENESS. Pectoral Decoction, 4. Oily Draughts, 10, 54, 97, 120. Vulnerary Decoction 49. Linctuffes and Lohochs, 12, 13, 61. Sperma Ceti Mixture, 15, 125. Balfamic Bolus, 39, 85. Tragacanth Pills, 67. Ditto Powder, 70. Pectoral Bolus, 88. Bran Water, 112. Pectoral Electuary, 114, 116.

HICCOUGH.

HICCOUGH. Oil of Juniper Mixture, 15. Lavender Drops, 19. Mulk Julap, 59.

HEART-BURN. Chalk Julap, 11. Magnefia Alba, 13. Chamomile Bolus, 39. Bitter Infusion, 56. Testaceous Powder, 70. Troches for the Heartburn, 72. Stomachic Electuary, 92.

JAUNDICE. Saponaceous Bolus, 2, 44. Aloetic Wine, 20. Calomel Bolus, 39. Cummin Seed Plaifter with Soap, 50. Expression of Hog-lice, 53, 152. Bitter Insufions, 56. Emetic Draught, 63. Chalybeate Electuary, 91. Saponaceous Draught, 98, 122. Tincture of Rhubaib, 107, 132. Tincture of Hiera-picra, 132. Aloetic In suffon, 154. Soluble Tartar, 72. Soap Pills, 160. Diuretic Decoction, 148. Saponaceous Electuary 150

ILIAC PASSION. See Belly-Ach dry-

INFLAMMATION. Anodyne Cataplaim, 23. Saline Mixture, 62. Fomentation of Poppy Heads, 8, 53. Purging Draught, 55. Ointment of Marihmallows, 77. Emollient Pultice, 81. Common Fomentation, 82. Refrigerant Ointment, 108. Nitrous Powder, 129, 142.

INDIGESTION. Emetic Draught, 63. Stomachic Bolus, 2. Arcmatic Electuary, 5. Bitter Infufions, 56. Saline Mixture, 62. Rufus's Pill, 67.
Tincture of Aloes, 75. Stomachic Electuary, 92, 139. Elixir of Vitriol, 92. Stomachic Pills, 104.
Wormwood Electuary, 113. Bates's Pepper Elecuary, 115. Tincture of Rhubarb, 132. Chamomile Infution, 99.

INJECTIONS. 28, 123.

JOINTS,

Survey Decembral, Star Street and Street

JOINTS, white Swelling of. Difcutient Mixture, 28. Saponaceous Cerate, 25. Embrocations of Barbadoes Tar, 64, 124. And Sea Water taken internally.

ISCHURY. See Dyfury.

ITCH. Itch Ointment, 78, 108, 133. Sulphur Lotion, 62, 101. Mercurial Liniment, 101. Camphorated Water, 110. Antimonial Powder, 104. Dietetic Decoction, 90. Precipitated Sulphur, 72. Lotion for the Itch, 28, 140. Itch Bolus, 42.

water Kennelik

KIDNEYS, weak. Strengthening Plaister, 50.

ULCERATED. cerated Diuretic Decoction, 143.

KING'S EVIL. Æthiopic Powder with Rhubarb, 17. Method of Cure, 34. Vegetable Æthiops, 35, 45. Sea Water, 36. Decoction of the Bark, 48. Buckbean Infusion, 57. Dietetic Decoction, 90. Bolus for the King's Evil, 89. Plaister for the Scrophula, 117. Troches of Burnt Sponge, 162. Antimonial Powder, 160.

L

LEPROSY. Ointment for the Leprofy, 20. Æthiopic Pill, 28. Lead Ointment, 33. Antimonial Æthiopic, 35. Aqua Liberans, 36. Alterative Bolus, 37. Mercurial Solutions, 71. Alterative Pills, 102, 103. Antimonial Powder, 104. Decoclion of Elm Bark, 113. Plummer's Powder, 129.

LUNGS, Diseases of. Capivi Mixture, 17, 67. Vulnerary Decoction, 49. Oily Draught, 55. Balsamic M 2 Pills,

Pills, 65. Potion Balfam of Peru, 68. Balfamie Bolus, 85. Pectoral Decoction, 90. Pectoral Mixtures, 102. Pectoral Pills, 126.

-LETHARGY. Volatile Julap, 12. Sneezing Powder, 18. Sinapism, 19. Blifters, Preparations of Valerian, and all Stimulants in general. See Apoplexy.

LOOSENESS. Rhubarb Bolus, 2. White Decoction, 4, 112. Aftringent Electuary, 6, 91. Infusion of Mint, 10. Chalk Julap, 11, 58. Reitringent Julap, 11, 102. Reftringent Bolus, 22, 38. Aftringent Drink, 30. Aluminous Bolus, 38. Bolus for a Diarrhœa, 40. Astringent Decoctions, 47, 90. Decoction of Simarouba, 49. Jelly of Starch, 54. Emetic Draught, 63. Tincture of Rhubarb, 74. Tincture of Japan Earth, 107. Reftringent Clyfter, 118. Aluminous Powder, 127.

LOWNESS of Spirits. Cordial Mixture, 63. Spirit of Sal Armoniac, Hartshorn, Lavender, Volatile, Aromatic, of Affafætida, &c. 71.

M

MADNESS. Emetic Draught, 63. Camphor Julap, 58, 120. Tincture of Black Hellebore, 74. Antimonial Wine, 76. Camphor Bolus, 85. And all Chalybeates.

MEASLES. Alexiterial Powder, 68. Saline Draught, 112. Sal Prunel Powder, 129. Mindererus's Spirit, 130. Purging Draught, 121. (This Draught is a Dofe for a grown perion)

MENSES

MENSES to promote. Aromatic Electuary with Steel, 5, 91, 114. Myrrh Electuary, 6. Steel Bolus, 39. Rufus's Pill with Steel, 67. Tincture of Steel, 74. Of Black Hellebore, ibid. Tincture to promote the Menfes, 106. Sagapenum Pills, &c. 126. Golden Tincture, 131. Myrrh Pills with Steel, 160. Chalybeate Electuary, 149. Hellebore Electuary, 150.

Excefive Discharge of, to restrain. Aluminous Bolus, 1. Restringent Drink, 30, 47, 90. Decoction of the Bark, 48. Tincture of Roses, 75. Styptic Tincture, ibid. Bark Electuary, with Alum, 91. Refrigerant Draught, 91. Styptic Julap, 800. Tincture of Japan Earth, 107. Anodyne Draught, 119. Aluminous Powder, 127.

MISCARRIAGE. See Abortion.

MORTIFICATION. Febrifuge Bolus, 41. Cordial Mixture, 63. Difcutient Pultice, 80. Common Fomentation, 82, 118. Digeftive Balfam, 22. Yellow Bafilicon, 132.

OBSTRUCTIONS of the Viscera. Soap Pills, 150.

Phananast

Profes Wester with Mine 196 W value

P A I N. Poppy Fomentation, 8, 53. Anodyne Cataplasm, 23. Hemlock Fomentation, 82. Emollient Pultice, 81. Anodyne Plaister, 93. Emollient Fomentation, 118.

PALSY. Palfy Gargle, 9. Paralytic Infusion, 11, 57. Volatile Julap, 12, 59, 154. Sinapism, 19, M 3 Discutient Discutient Liniment, 27 Method of Cure, 31. Epileptic Bolus, 41. Volatile Liniments, 61. Muftard Whey, 70. Volatile Tinctures of Bark, 73. Volatile Tinctures of Valerian, 76. Cinnabarine Bolus, 86. Volatile Gargle, 95, 119. Infusion of Valerian, 99. Valerian Electuary, 115. Blifter Plaister, 117. Horse-radish Infusion, 123. Tincture of Hiera-picra, 132. Tincture of Wood Soot, 131. Mustard Infusion, 155.

- PERSPIRATION to promote. Alexiterial Bolus, 37. Camphor Julap, 58, 59. Alexiterial Powder, 68, 104. Cordial Bolus, 86. Alexipharmic Tincture, 106, Sudorific Draught, 120. Dr. Dover's Powder, and Diaphoretic Powder, 128, 142. Mindererus's Spirit, 130, 161. Contrayerva Bolus, 147.
- PHLEGM to cut. Ammoniacum Mixture, 14, 60. Squill Bolus, 44. Oily Emulfion, 51. Afthmatic Mixtures, 62. Balfamic Bolus, 85. Barbadoes Tar Electuary, 92. Ammoniacum Pills, 102. Elecampane Electuary, 114. Oleaginous Draught, 122. Volatile Sperma Ceti Mixture, 125. Acid Electuary, 114.
- PILES. Electuary, 6, 42, 87. Emollient Fomentation, 8. Clyfters for the Piles, 52. Pile Ointment, 77. The Brimftone Electuary, 150.
- PLEURISY. Barley Water with Nitre, 36. Vulnerary Decoction, 49. Oily Draughts, 55. Saline Mixtures, 62. Pectoral Decoction, 4, 90. Refrigerant Draught, 98. Pectoral Mixtures, 102, 141. Sal Prunel Powder, 129. Mindererus's Spirit, 130.

o a Paralyner Infedian, 1 m.

20 main Jack

. 21 . al goard . + 21 . 22 . 4. gelat site 2UINSY

RUP-

Q

Printer, Siz. Andreal monthly in a start

5 WITTER A. Maintenater Francisco States

QUINSY, common. Gargles, 9, 54. Purging Draught, 55. Mindererus's Spirit. 130. Troches of Nitre, 76.

yerva Gargle, 9, 54. Alexiterial Bolus, 37. Decoction of the Bark, 48. Cordial Mixture, 63. Alexipharmic Gargle, 95, 119. Detergent Gargle, 153.

R

Con Breat and

a company (or invester) walker

RHEUMATISM. Guaiacum Bolus, 2, 42. Volatile Julap, 12, 59. Muftard Whey, 19, 70. Alterative Bolus; 37. Ditto Electuary, 138. Cataplafm, 46. Sweating Powder, 69. Tincture of Guaiacum, 73. Turpentine Bolus, 86. Scammony Electuary, 92. Guaiacum Electuary, 114, 149. Muftard Electuary, 115. Anodyne Embrocation, 116. Saponaceous Pill, 127. Mindererus's Spirit, 130. Doctor Dover's Powder, 128. Cinnabarine Electuary, 149.

RICKETS. Magnefia Alba, 13. Aromatic Species, 19. Flores Martiales, 9. Saponaceous Liniment, 61. Soluble Tartar, 72. Common Fomentation, 82. Spirits of Wine and Camphor, 105. Volatile Fomentation, 114. Strengthning Plaister, 51.

3

The XLYCE

RUPTURE. Reftringent Fomentation, 9. Strengthening Plaister, 82. And all internal Strengthening Medicines.

S

SCALD Head. Tar Ointment, 32, 132. Antimonial Æthiops, 35. Ointment for a Scald Head, 109, 133.

SCIATICA. See Hip-gout.

SCURVY. Yellow Ointment, 20. Æthiopic Pill, 28. Lead Ointment, 33. Alterative Bolus, 37. Dietetic Decoction, 48. Buck-bean Infusion, 57. Mercurial Pills, 66. Scorbutic Juices, 71. Lime Waters, 79. Guaiacum Bolus, 87. Alterative Electuary, 91, 92. Alkalized Mercury, 101. Antimonial Powder, 104. Medicated Ale, 111. Plummer's Powder, 129. And Sea Water.

SHINGLES. Purging Powder, 18. Common Cerate, 25. Alterative Bolus, 37. Cleanfing Bolus, 40. Purging Draught, 55. Sulphur Lotion, 62. Scorbutic Juices, 71. Precipitated Sulphur, ibid. White Liniment, 61. Yellow Cerate, 81. Alkalized Mercury, 130. Dietetic Decoction, 90.

SMALL POX. Alexiterial Bolus with Nitre, 37. Bolus for the Small Pox, 46. Tincture of Roles, 75. Camphor Bolus, 86. Bark Electuary with Alum, 91. See Fever.

52

STITCHES

STITCHES in the Side. Cummin Seed Plaister, 50. Saponaceous Liniment, 61. Volatile Liniment, 101.

STONE. Nephritic Electuary, 7. Oily Draught with Manna, 10. Turpentine Clyfter, 52. Shell Lime Water, 110.

STRAINS. Difcutient Potion, 3. Saponaceous Cerate, 25. Volatile Liniment, 61. Strengthening Plaister, 82, 93, 117. Camphorated Liniment, 124.

SWEAT, to promote. See Perspiration, to promote.

Acid Elixir, 50, 92. Bark Electuary, 115. Saturnine Tincture, 75.

T

TEETH to cleanse. Tooth Powder, 30. Tincture of Bark, 72. Tincture of Myrrh, 83.

TENESMUS. Anodyne Clyfter, 8. Anodyne Draught, 9. Cordial Mixture, 63. Venice Treacle Julap, 59. London Philonium, 64. See Dyfentery.

TETTARS. Yellow Ointment, 20. See Cutaneous Diforders.

T'HRUSH.

THRUSH. Decoction of Elm Bark, 5. Aluminous Gargle, 54. Honey of Roses, with burnt Alum, 62.

TUMOURS, to suppurate. Suppurative Cataplasm, 23. Galbanum Plaister, 51, 117. Common and Pultice, 80, 81. Emollient

to discuss. Saponaceous Cerate, 25. Discutient Pultice, 80. Common Fomentation, 82. Discutient Mixture, 28. Volatile Liniment, 101. Camphorated Liniment, 124. Mercurial and Soap Plaister,

THROAT, sore. See Quinfy.

XX

TYMPANY. Aromatic Species, 19. Scammony Electuary, 49. Tincture of Aloes, 75. Purging Bolus, 88. Common Clyfter, 94. Common Fomentation, 95. Muftard Electuary, 115. Tincture of Sena, 132.

V

VENEREAL Difeafe. Specific Bolus, 3. Specific Electuary, 7, 26. Mixture for, 14. Purging Pills with Mercury, 17. Blue Ointment, 20. Æthiopic Pill, 29. Mercurial Pills, 66, 67, 159. Mercurial Ointment, 77. Ditto Digeftive, 143. Fumigations of Cinnabar, 82. Dietetic Decoction, 90. Alterative Pills, 102, 103. Guaiacum Decoction, 113, 148. Sarfaparilla Decoction, 48. Specific Pills, 141.

VERTIGO. See Epilepfy.

ULCERS.

ULCERS. Emollient Fomentation, 8, 82. Yellow Bafilicon, 32. Detergent Ointment, 83, 84, 162. Alterative Electuary. 91. Green Ointment, 109, 134. Digeftive Ointments, 83, 84, 107, 132, 134, 143.

ULCERATIONS internal. Balfam of Peru Mixture, 17. Balfamic Bolus, 39, 85. Vulnerary Decoction, 49. Balfamic Pills, 65. Pectoral Decoction, 90. Oily Draught, 97. Balfamic Electuary, 149.

VOMITING to reftrain. Infusion of Mint, 10. Saline Mixture, 15. Paregoric Draught, 55. Bitter Infusion, 56. Mint Julap, 59. Elixir Vitriol, 92. Anodyne Draught, 96. Stomachic Draught, 98. Chamomile Infusion, 99. Saline Decoction, 136.

URINE, Heat of. See Dyfury.

to promote. Saponaceous Bolus, 2. Diuretic Salt, 19. Hydragogue Draught, 54. Diuretic Powder, 68. Soluble Tartar, 72. Turpentine Bolus, 89. Diuretic Draught, 96, 153. Nitrous Powder, 129, 142. Mindererus's Spirit, 130. Diuretic Apozem, 144. Ditto Decoction, 148.

-involuntary Discharge of. See Diabetes.

Bloody. Aluminous Bolus, 1. Febrifuge Bolus with Alum; 41. Tincture of Rofes, 75. Elixir of Vitriol, 92. Alum Whey, 70. And all ftrong Aftringents.

WHITES

. of. Erress Oin

Washington Washington

WHITES. See Fluor Albus

WIND in the Bowels. All Carminatives. Anodyne Fetid Bolus, 146. Pepper Bolus, 147, Anodyne Difcutient Plaister, 150.

WORMS. Tin Electuary, 7. Æthiopic Powder with Rhubarb, 17. Worm Powder, 18, 128. Tin Bolus, 43, 89, 147. Alkalized Mercury, 101. Tin Powder, 105.

WOUNDS. See Ulcers.

FINIS.

