Publication/Creation

4 July 1933

Persistent URL

https://wellcomecollection.org/works/y6pqudgy

License and attribution

Conditions of use: it is possible this item is protected by copyright and/or related rights. You are free to use this item in any way that is permitted by the copyright and related rights legislation that applies to your use. For other uses you need to obtain permission from the rights-holder(s).



Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org





FIG. 7.—This illustrates the various forms of curious bodies found in silicoanthracosis. The measurements of the spicules forming the nuclei of the bodies are:—(1) $85 \cdot 0\mu$; (2) $27 \cdot 5\mu$; (3) $35 \cdot 0\mu$; (4) $35 \cdot 0\mu$; (5) $22 \cdot 5\mu$; (6) $17 \cdot 5 \times 12 \cdot 5\mu$; (7) $35 \cdot 0\mu$; (8) $42 \cdot 5\mu$; (9) $30 \cdot 0\mu$; (10) $50 \cdot 0\mu$; (11) the longest spicule in the group is $77 \cdot 5\mu$; (12) $37 \cdot 5\mu$; (13) $44 \cdot 0\mu$; (14) $37 \cdot 5\mu$; (15) a curious body from a case of asbestosis. This form is unusual in asbestosis and is included here for comparison. The length of the nucleus is $40 \cdot 0\mu$. Photomicrographs by C. F. Hill and W. E. Cooke.