

M0005146: Richard Lower's blood transfusion experiment, page from an account of 1666

Publication/Creation

14 April 1937

Persistent URL

<https://wellcomecollection.org/works/bzfv7kmd>

License and attribution

Wellcome Library; GB.

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>

there be occasion to open them.

All things being thus prepared, ty^e of Dogs
on their sides towards one another so convenient,
if quills may goe into each other (for if Dogs necks
cannot be brought soe near, but if you must
put 2 or 3 severall quills more into y^e first
two, to convey y^e blood from one to another) After
y^e unstop y^e quill going downe into y^e Dogs jugular veine
& y^e other quill coming out of y^e other Dogs artery,
& by y^e help of 2 or 3 other quills, put into each other
according as there shall be occasion, put them into one
another, then slip y^e running knot, & immediately y^e blood
runs through y^e quill, as through an artery very impetuously
& immediately as blood runs into y^e Dogs, out of y^e other
quill coming out of y^e upper part of his jugular veine (a
ligature being first made about his neck, or else his other
jugular veine being compr^{es}sed by ones finger) & let his
owne blood run out at y^e same time into dishes (yet
not constantly, but according as you perceive him to
bare it) till other Dog begins to cry & faint, & fall
into convulsions, & at last dye by his side.

Then take out both y^e quills out of y^e Dogs jugular
veine & ty^e y^e running knot fast, & cut y^e veines as
if you may doe without any harme to y^e Dog, one
jugular veine being sufficient to convey all y^e blood
from y^e head & upper parts by reason of a large anastomosis
wherby both y^e jugular veines meet about y^e Larynx.
This done sow up y^e skin, & dismisse him & y^e Dog
will leape from y^e table, & shake himselfe & run away
as if nothing ailed him. And this I have tryed severall
times before severall in y^e University, but never yet
upon more then one dog at a time, for want of
times & convenient supplies of severall dogs
at once. But when I returne (wh^o I hope may be
this Autumme) I doubt not but to give you
a fuller Acc^t, not only by bleeding severall
Dogs into one, but severall other Creatures
into one another, as you did propose to me
before you left Oxford. It will be very easy to
performe, & will afford many pleasant & perhaps not
unusefull Experiments.