Institutions of medicine. Part I, Physiology. For the use of the students in the University of Edinburgh ... / [William Cullen].

Contributors

Cullen, William, 1710-1790

Publication/Creation

Edinburgh : Charles Elliot; London : T. Cadell, 1785.

Persistent URL

https://wellcomecollection.org/works/apmnmcva


License and attribution

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.




Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org













INSTITUTIONS

OF

43239

MEDICINE. PARTI. PHYSIOLOGY.


For the use of the STUDENTS in the University of Edinburgh.

By WILLIAM CULLEN, M.D. PROFESSOR OF THE PRACTICE OF PHYSIC, &c. &.

THE THIRD EDITION, CORRECTED.

E D I N B U R G H: Printed for CHARLES ELLIOT, EDINBURGH; And T. CADELL, Strand, LONDON.

M, DCC, LXXXV.



ADVEERTISMENT.

HE following sheets were originally printed as a text-book for the use of Students of Medicine in the University of Edinburgh; it of courfe contains only fketches of what was more fully delivered in the Professor's lectures. Though the impression formerly made has been long ago difposed of, the Author did not choose to renew it till he should be at leifure to give it to the public in a more complete form. But Dr Gregory, who now teaches this branch of medicine, thinking those sketches might be useful to his Students, the author has confented to the republication: and it is now offered to the public in a more correct manner.

Just published by CHARLES ELLIOT Edinburgh, and T. CADELL London,

FIRST LINES

OFTHE

PRACTICE OF PHYSIC.

By WILLIAM CULLEN, M.D.

Professor of the Practice of Physic in the University of Edinburgh; First Physician to his Majesty for Scotland; Fellow of the Royal College of Physicians of Edinburgh, of the Royal Societies of London, of Edinburgh, &c. &c.

> NOW FIRST COMPLETED IN FOUR VOLUMES OCTAVO. Price 11. 45. in boards.

> > ALSO,

DR CULLEN ON THE RECOVERY OF

PERSONS DROWNED AND SEEMINGLY DEAD. Price One Shilling.

INSTITUTIONS

MEDICINE.

O F

MEDICINE is the art of preventing and of curing difeafes.

onitBoob lensing any

II.

Before confidering the application of this art to particular difeafes, certain general doctrines are neceffary to be premifed, which are called THE INSTITU-TIONS OF MEDICINE.

A

III.

(6)

ANSTITUTIONS

The Inftitutions of Medicine are divided into three parts.

The first treats of life and health.

The fecond delivers the general doctrine of difeafes.

The third delivers the general docctrine concerning the means of preventing and curing difeafes.

PART

PART I.

PHYSIOLOGY.

N. IV.

THE doctrine which explains the conditions of the body and of the mind neceffary to life and health, is called PHY-SIOLOGY, or the Doctrine of the Animal Oeconomy.

V. /

The functions of the animal ∞ complicated are many and various; and fo complicated with each other, that it is difficult to find the most proper order in which they may be delivered. That, however, seems the A 2 best

8

best which confiders them as nearly as may be according to the feries of causes and effects.

PHYSLIVEOG

Upon this plan we fhall treat,

1. Of the folid matter, of which a great part of every organ of the body confifts.

2. Of the nervous fystem, in which the motions of the body for the most part begin, and upon which the motions produced in it chiefly depend.

3. Of the motion and circulation of the blood, and of the feveral organs and actions employed in fupporting it.

4. Of the functions employed in fupporting and repairing the feveral folid and fluid matters of the body; and, on this occasion, of the nature of the feveral fluids themfelves.

52

5. Of

5. Of the organs employed in receiving and modifying the impreflions of external bodies neceffary to fenfation; and of their several functions.

6. Of the motions of the whole body, or of its feveral parts which depend on the action of muscles, and not before explained.

7. Of the functions peculiar to the fexes, and of generation.

firft are called the SIMPLE SOLIDS. Of

ins, being the fundamental part of the

-Y H 9 ace, in opt A ion to which, the

SECT. I.

OF THE SIMPLE SOLIDS.

VII.

THE folid parts of the body feem to be of two kinds; one whofe properties are the fame in the dead as in the living, and the fame in the animate as in many inanimate bodies; the other, whofe properties appear only in living bodies. In the laft, a peculiar organization, or addition, is fuppofed to take place; in oppofition to which, the firft are called the SIMPLE SOLIDS. Of thefe only we fhall treat here; and of the others, which may be called VITAL SO-LIDS, being the fundamental part of the ner-

nervous fystem, we shall treat under that title in the following section.

VIII.

The fimple folids are fuited to the purpofes of the animal œconomy by a certain force of cohefion, joined with a certain degree of flexibility and elafticity.

IX.

Thefe properties of the fimple folids, in different parts of the body, in different bodies, and on different occasions in the fame body, are neceffarily in different degrees; and this feems to depend upon the difference of the mixture, aggregation, or organization of the folid.

The matter of the fimple folid every where

X. called the ANT-

where, except in the bones, appears to be an homogeneous aggregate; and there is no proper evidence of its being formed of certain parts naturally difcrete and incoherent, which are cemented by others of a different nature.

force of cohefion, joined with a certain IX. degree of flexibility and elafficity.

Of the fimple folid confidered as an homogeneous aggregate, the integrant parts are a mixt, which feems to be nearly of the fame kind in all the different parts of the human body, and perhaps in most of the parts of every animal: So far as we yet know, the variety of it is very inconfiderable.

organization of the folid. IIX

This, which may be called the ANI-MAL MIXT, is found, by chemical experiment,

riment, to be confiderably different from every kind of vegetable or foffil matter; but the fame experiments hardly teach us any thing exact or useful with respect to the conflituent parts of this mixt.

XIII.

defletent perfona are under the

The only particular relative to this, which we exactly know, is, that the animal mixt is formed of water, and of fome other matter concreting with it; that, on different occafions, the flate of it is varied by the proportion which the water bears to the other concreting matter; and that, efpecially by a different proportion in this refpect, the fimple folid differs in its force of cohefion, flexibility, and elafticity (VIII.)

The proportion of water to the other matter

13

matter in the animal mixt of different perfons feems to depend, in the first place, upon the nature of the original stamina in each; as the different state of the simple folids, which appears early to diffinguish fex and temperament, continues respectively the same through the whole of life, even though the different perfons are under the fame external circumstances.

XV.

But, in every particular perfon, that proportion is conftantly changed by the progrefs of life; and this happens more or lefs as other caufes concur.

XVI.

The caufes that can affect the mixture of the fimple folid, are either the flate of the nutritious fluid conveyed by the ordinary channels,

channels, or fome matters from without infinuated into the folid.

XVII.

The flate of the nutritious fluid may be varied by the quantity and quality of the aliment taken in, by the powers of concoction and affimilation, by the circumflances of application and concretion, or by certain preternatural matters carried along with it.

XVIII.

The external matter that may be infinuated into the fimple folid is various, but for the most part is only aqueous moisture in greater or lefs quantity.

XIX.

That these several causes may affect the pro-

16

proportion of water in the fimple folid, and thereby give a different flate of it, is fufficiently obvious: That the fame caufes may also affect the other concreting matter, we can, in general, perceive to be poffible; but in what manner, or upon what occasions they do fo, is not eafily difcerned.

of application and XX cretion, or by cer-

tion and allimitation, by the circumitances

The properties of the fimple folid (VIII.) may be also varied by its state of aggregation; and this again may be varied, I. By the temperature of the atmosphere to which the body is long exposed. 2. By the preffure, external or internal, which is applied to the folid. 3. By the degree of extension of the folid beyond its natural state, which, in every living body, is given more or lefs to every part of the foft or flexible folids; bra hat thefe foreral cantes may sugel th

17

and, *laftly*, by the motion or reft to which the folid is accustomed.

XXI.

The properties (VIII.) of the folid parts are alfo varied by the ftate of their organization. This every where depends upon an arrangement of fibres, the ftate of cellular texture, or upon a texture of veffels; and therefore, to explain the different ftates of organization, it will be enough to mention the caufes of the differences which occur in thefe fundamental parts.

XXII.

Fibres may differ in fize by the feveral caufes (XIV.—XXI.) affecting the mixture and aggregation of the matter of which they are formed, and by these causes alone; but how far the organization of any

18

any part depends upon an arrangement of fibres, we cannot diffinctly perceive; and, if it does, we cannot perceive that the flate of fuch parts differs otherwife than by the flate of the cellular texture every where interpofed between the fuppofed fibres.

XXIII. and I double stat

The flate of cellular texture is the moft important circumflance in all organized parts; and it may be varied by many different caufes. 1. The texture may be more denfe, and thereby firmer, as it has been more preffed by the actions of life or external force; by which means efpecially it is changed in the progrefs of life. 2. The cellular texture may be increafed in bulk, and rendered firmer by a new growth taking place in it, as frequently happens in membranes which are flowly and gradually ftretched out. 3. The fame texture may be-

become weaker by fome part of it being eroded by acrid matters generated in the body, or externally applied. 4. It is analogous to this, that, when any part is fustained by several layers of cellular texture or membranes, fuch fupport is weakened by one or more of thefe layers being cut through; and the fame weaknefs is induced when any external compression, which, for fome time, had been applied, is taken away. 5. The flate of the cellular texture is varied by the matter contained in its cells; which is fometimes a matter concreting into a folid mafs, and fometimes a preternatural quantity of an aqueous inelaftic fluid. The bones formed in the first manner may again become foft by the hardened matter's being diffolved and reabforbed. 6. When the mobility of parts on one another depends upon the extent of cellular texture connecting them, that mobility is diminished or destroyed by a great part Ā

19

part of the cellular texture being eroded or cut away, and the remaining parts being united together; fo that the parts are now connected by a fhorter portion of cellular texture than before. 7. Parts naturally feparate may lofe their mobility by being joined together by a cellular texture formed between them, as happens when any two furfaces are for fome time kept clofely applied to each other.

is waried by the VIXX contained in its

cells ; which is fometimes a

away: 5. The flate of the cellular texture

In fo far as a folid part is formed by a texture of veffels, its properties (VIII.) may be varied by the different flates of thefe veffels; which, 1. may be more or lefs full of fluids. 2. They may be changed into a folid mafs, by the fluid, contained and flagnating in them, concreting into a folid. 3. They may be changed into a folid, if the fluids that flould pafs through

through them are intercepted, and the cavity is filled with a cellular texture; or, 4. They may be changed into a folid, if, by collapse or preffure, the fides of the veffels are applied to each other and concrete together.

XXV.

The pathology of the fimple folids cannot be properly feparated from their phyfiology; and therefore many different flates of thefe folids, though fuch as are always morbid are mentioned above. We think it proper to fubjoin here a flort view of the whole of that pathology.

XXVI.

The difeases of the fimple folids are,

- I. Those of the naturally fost parts.
- 1. Debility with flexibility.

Debile tenerum, gracile, Gaub. Pathol. 161. 1.

B

Debile

Debile tabidum Gaub. ibid. 161. 2. 2. Debility with fragility. Debile fiffile Gaub. 161. 3. 3. Laxity. Debile laxum, flaccidum, Gaub. 160. 1. 4. Flaccidity. Debile iners Gaub. 160. 2. 5. Rigidity diminishing flexibility. Rigidum tenax Gaub. 165. 1. CT ST & 6. Rigidity deftroying flexibility. Rigidum durum Gaub. 165. 2. II. Those of the naturally hard parts. 1. Flexibility. Debile flexile Gaub. 160. 3. 2. Fragility. Fragile Spongiofum Gaub. 161. 4. Fragile vitreum Gaub. 165. 3. Of all these morbid affections, we fuppofe the remote and proximate caufes may be understood from what is delivered above

(XIV.-XXIV.)

SECT.

22

SECT. II.

OF THE NERVOUS SYSTEM.

XXVII.

THE nervous fystem, as the organ of fense and motion, is connected with so so many functions of the animal æconomy, that the study of it must be of the utmost importance, and a fundamental part of the study of the whole æconomy.

A general view of the Nervous System.

XXVIII.

The nervous fystem confists of the medullary substance of the brain, cerebellum, B 2 medulla

medulla oblongata, and fpinalis; and of the fame fubstance continued into the nerves, by which it is diffributed to many different parts of the body.

XXIX.

The whole of this fystem seems to be properly distinguished into these four parts.

1. The medullary fubftance contained in the cranium and vertebral cavity; the whole of which feems to confift of diftinct fibres, but without the feveral fibres being feparated from each other by any evident enveloping membranes.

N. When we fpeak of functions, which are or may be in common to every part of this portion of the nervous fystem, we shall speak of the whole under the title of the BRAIN; but, when it is necessary to distinguish the particular parts, we shall take care to avoid ambiguity.

2. Connected with one part or other of \mathbb{N}° 1. are, the NERVES, in which the fame medullary fubftance is continued, but here more evidently divided into fibres, each of which is feparated from the others by an enveloping membrane derived from the pia mater.

3. Parts of the extremities of certain nerves (2.), in which the medullary fubftance is divefted of the enveloping membranes from the pia mater, and fo fituated as to be exposed to the action of certain external bodies, and perhaps fo framed as to be affected by the action of certain bodies only: These we name the SEN-TIENT EXTREMITIES of the nerves.

4. Certain extremities of the nerves (2.) fo framed as to be capable of a peculiar contractility: and, in confequence of their fituation and attachments, to be, by their contraction, capable of moving most of the folid and fluid parts of the body. These

we

B 3

we name the MOVING EXTREMITIES of the nerves: They are commonly named MOVING or USCULAR FIBRES.

N. That the mulcular fibres are a continuation of the medullary fubftance of the brain and nerves, has not been shown by the anatomists, nor univerfally admitted by the physiologists; but we now suppose it, and hope afterwards to render it fufficiently probable.

Are the ganglions of the nerves to be confidered as a part of the nervous fystem distinguished by a peculiar function?

XXX.

Thefe feveral parts of the nervous fyftem are every where the fame continuous medullary fubftance which we fuppofe to be the vital folid of animals, fo conftituted in living animals, and in living fyftems only, as to admit of motions being readily propagated from any one part to every other part

part of the nervous fystem, so long as the continuity and natural living state of the medullary substance remains.

N. It is obferved, that the compression of any part of the medullary substance prevents the communication of motion between the parts that lie on different fides of the part compressed; and it is probable, there are other causes befides compresfion, which may also affect the medullary substance, so as to interrupt in it the communication of motion; but they are not diffictly known. In the mean time, we use the expression, that a nerve, or other portion of the nervous system, is *free*, to denote its being free, not only from compression, but from every other supposed cause interrupting the communication of motion.

The condition fitting the medullary fubftance for having motion propagated in it, we fuppofe to be the prefence of a certain fluid; which we therefore name the *nervous fluid*, without meaning however at prefent to determine any thing with regard to its fource, nature, or mannner of acting.

B 4

XXXI.

27

28

XXXI.

In the living man, there is an immaterial thinking fubstance, or MIND, constantly prefent; and every phenomenon of thinking is to be confidered as an affection or faculty of the mind alone. But this immaterial and thinking part of man is fo connected with the material and corporeal part of him, and particularly with the nervous fystem, that motions excited in this give occasion to thought; and thought, however occasioned, gives occasion to new motions in the nervous fystem. This mutual communication or influence we affume with confidence as a fact : But the mode of it we do not understand, nor pretend to explain; and therefore are not bound to obviate the difficulties that attend any of the fuppofitions which have been made concerning it.

XXXII.

XXXII.

The phenomena of the nervous fyftem occur commonly in the following order. The impulfe of external bodies acts upon the fentient extremities of the nerves; and this gives occafion to preception or thought, which, as firft arifing in the mind, we term SENSATION. This fenfation, according to its various modification, gives occafion to VOLITION, or the willing of certain ends to be obtained by the motion of certain parts of the body; and this volition gives occafion to the contraction of mulcular fibres, by which the motion of the part required is produced.

N. This is an example of the most ordinary cafe; but we do not mean to fay it is the only cafe of communication between the different parts of the nervous fystem.

XXXIII.

Unable to display this page

there, and propagated along the nerves, produces the contraction of muscles.

XXXIV.

From what is now faid, we perceive more diffinctly the different functions of the feveral parts of the nervous fystem, as diftinguished in (XXIX.), 1. The fentient extremities (XXIX. 3.) feem to be particularly fitted to receive the impreffions of external bodies; and, according to the difference of these impressions, and of the condition of the fentient extremity itfelf, to propagate along the nerves motions of a determined kind, which, communicated to the brain, give occasion to fensation. 2. The Bain (XXIX. 1.) feems to be a part fitted for, and fusceptible of, those motions with which fenfation, and the whole confequent operations of thought, are connected; and thereby is fitted to form

form a communication between the motions excited in the fentient, and those in confequence arifing in the moving extremities of the nerves, which are often remote and diftant from each other. 3. The moving extremities (XXIX. 4.) are fo framed as to be capable of contraction, and of having this contraction excited by motion propagated from the brain, and communicated to the contractile fibre. 4. The nerves, more strictly fo called (XXIX. 2.), are to be confidered as a collection of medullary fibres, each inveloped. in its proper membrane, and thereby fo feparated from every other, as hardly to admit of any communication of motion from any one to the others, and to admit only of motion along the continuous medullary fubstance of the fame fibre, from its origin to the extremities, or contrariwife.

XXXV.

33

XXXV.

From this view of the parts of the nervous fystem, of their feveral functions and communication with each other, it appears, that the beginning of motion in the animal acconomy is generally connected with fenfation; and that the ultimate effects of fuch motion are chiefly actions depending immediately upon the contraction of moving fibres, between which and the fentient extremities the communication is by means of the brain. Wherefore, in fludying the nervous fystem, we judge it proper to confider, 1. Senfation, and with that the general function of the fentient extremities. 2. The action of the moving fibres. 3. The function of the brain. In confidering thefe three heads, the function of the nerves, more firicity fo called, will of courfe be fufficiently explained.

CHAP.

2

CHAP. I.

OF SENSATION.

XXXVI.

O UR fenfations may be be confidered as of two kinds: 1. Thofe which arife from the impulfe or imprefion of external bodies, which we therefore name SENSATIONS OF IMPRESSION. 2. Thofe which arife from the mind's being confcious of its own action, and of the motions it excites; and thefe we name SENSATIONS OF CONSCIOUSNESS.

Senfations of Impression.

XXXVII.

The fenfations of impression are very z various;

35

various; but have been generally referred to five heads or kinds, commonly called the five fenfes; that is, those of fight, hearing, fmell, tafte, and touch.

XXXVIII.

The four first of these are each of them properly confidered, as forming one genus of sensations: 1. As the particular sensations comprehended under each head (XXXVIII.), though very various, are, however, perceived to have somewhat common to all of them. 2. As those of the fame genus all arise from impressions made upon one part of the body only, and that of a peculiar organization. 3. As those of the fame genus all arise from the action of external bodies of one kind only, or of one and the fame quality, by means of which they act upon our organs.

XXXIX.

XXXIX.

No fuch characters concur in eftablifhing one genus of the fenfations referred to the fifth head of touch, which are various in all thofe refpects (XXXVIII.); and phyfiologifts feem to have referred to this head of touch every fenfation that does not manifeftly belong to the other four, and among the reft many of the fenfations of confcioufnefs. It might perhaps be useful to diftinguish into genera, the feveral fenfations referred to touch; but it is not necessary to be done here.

From certain fenfations referred to touch, it appears, that not only the extremities (XXIX. 3.), but that every part of the nervous fystem (XXVIII.), is fentient with respect to certain impressions.

Sen-

37

fen-

Senfations of Consciousness.

XL.

The fenfations of confcioufness may be referred to the following heads: 1. Those of apperception, by which we are in general confcious of thinking, of perceiving, judging, and willing, and thereby of our existence and identity. 2. The senfations arifing from the particular state of thinking, as perception, memory, and judgment, are more or lefs clear, ready, or exact. 3. The fenfations arising from the particular state of volition, and its various modes. 4. The fenfations arifing from the general flate of action, as vigorous or weak, eafy or difficult. 5. The fensations arising from particular actions, or a confcioufness of the actions excited, and of the motion of the different parts of the body. 6. The

C

fenfations arifing from the diminution or absence of impressions.

Under each of these heads, a great number of particular sensations are comprehended, but not necessary to be farther specified here.

Laws or general Circumstances of Senfation.

XLI.

Of the four first genera (XXXVII.), the fensations arising give no indication of the nature of the bodies acting on our organs, or of the mode of their action; and when we otherwise learn these circumstances, we can perceive no necessary connection between them and the fensations which they produce. But, from certain fensations of touch and confcious fers, we acquire the notions of folid figure, of motion, impulse,

pulfe, impenetrability, and the communication of motion, and confider the fenfations as exactly correspondent to the circumftances of external bodies. At the fame time, as we know of no other action of bodies on each other but that of impulfe; and as, in the cafe of the fenfations of the four first genera, we learn, that an impulse takes place, we have comprehended the whole under the title of Senfations of Impression, and confider all of them as perceptions of impulse.

XLII.

To produce any fenfation of imprefion, a certain force of imprefion is neceflary; and from a leffer force, no fenfation arifes. The degree of force is likewife fo limited on the other hand, that, in a high degree, it deftroys the organ; and, in degrees approaching to this, a general fenfation of C_2 pain,

"Ilv correliondence to the cir-

pain, rather than the fensation of any particular object, is produced.

ade 14 .collocd XLIH. to soons

Within these limits, however, our fenfations are not exactly correspondent to the force of impression, nor do they make any exact estimate of that force. Usually fensation is relative to the change that is produced in the nervous system; and a fenfation proves strong or weak, only as it is stronger or weaker than that which had immediately preceded it, or than that degree of force to which the nerves had been immediately before accustomed. For this reason too the limits (XLII.) are very variable.

the other hand, th.VIJX a high degree, it

Different senfations do not always imply

0 3

2

a different kind of action in the bodies producing them; for fometimes different fenfations arife merely from a different degree of force in the fame kind of action, as is manifest in the case of heat and cold.

XLV.

To fenfation from impression, a certain duration of impression is necessary.

NJX mprehion or ac-

for fome time

The mind's refting for fome time upon one fenfation, is called ATTENTION. This, like the duration (XLV.), is neceffary to give an impression its full effect.

XLVII.

The mind feems to be determined to C 3 . attention

attention by the force of imprefiion; by the pleafure or pain arifing from it; by the degree of emotion or paffion produced by thefe; and, laftly, by the emotion's being more or lefs related to the perfon feeling,

XLVIII.

If the force and duration of impreffion, and the attention of mind, are all in the due degree, the fenfation often remains for fome time after the imprefion or action of the external body has ceafed.

XLIX.

The mind admits of, or can attend to, one fenfation only at one time.

L.

L.

Though the mind admits but of one fenfation at one time, feveral imprefiions may act at the fame time, if they be fuch as can unite in producing a fingle fenfation; and fuch is the cafe of many of the imprefiions which produce the particular fenfations of the fame genus, as in thofe efpecially of colour, found, fmell, and tafte.

LI.

entre wel service years

In each of these genera, many impreffions, which separately produce particular species, can unite in producing a single sensation, which is always a neutral, or one different from either of the separate sensations.

C 4

LII.

This union of impressions may take place, either when the impressions are exactly fynchronous, or when the one fucceeds the other before the fensation of the first (XLVIII.) has ceased.

LIII.

preditions which produce the particul

Though the motion excited in the fentient extremities by impression remains fome time, as in (XLVIII.), it must be supposed to become continually weaker, till at length it ceases altogether, and with it the fensation.

LIV.

The fame impression foon repeated, does not produce the fame strength of sensation as before. Hence, all new impressions are,

are, *cæteris paribus*, ftrongeft; and moderate impressions frequently repeated, produce no sensation, unless their force is considerably increased.

LV.

Actions which at first produced a fensation of confciousness, as accompanied with volition, come, by repetition, to be performed without any fensation; or they produce it only when they are performed with uneafiness, pain, or unufual force.

edi lo stall od LVI.

Imprefions being given, their effects in producing fenfation are different in different perfons, and in the fame perfon at different times. This must arife from fome difference in the state of the bodies acted upon, which may perhaps be referred to the

the following heads: 1. The state of the common teguments, or other parts interpofed between the impreffing body and the medullary fubstance of the fentient extremity. 2. The different state of the medullary fubstance of the fentient extremities, as given to it in the original stamina. 3. The different state of tension in the medullary fubstance of the fentient extremities, as given to it by the state of the blood-veffels conftantly connected with it. 4. The ftate of the fame medullary fubstance, as affected by heat or cold. 5. The ftate of it as produced by former impreffions (XLIII. LIV.) 6. The state of the nerves along which the motion is propagated. 7. The state of the brain or sensorium. 8. The state of attention (XLVI. and XLVII.) and and mi but scooled and

of many pushage be reader to

-

LVII.

47

LVII.

Different parts of the body are fenfible, and fenfible only by means of nerves prefent in them: but anatomy does not always determine certainly with regard to the prefence or abfence of nerves; and, therefore, the fenfibility of feveral parts can be determined by experiment only; which, however, is alfo fallacious.

LVIII.

Particular fenfations arife from impreffions on certain parts only :. 1. Becaufe the fentient extremities in thefe parts are fo fituated as to be exposed to the action of certain external bodies only. 2. Becaufe the fentient extremities are connected with an organ that increases the force of the external agent, or modifies its action in the manner

ner neceffary to a determined imprefion. 3. Becaufe the fibres of certain fentient extremities are, by their fize or tenfion, fitted to be acted upon by certain external bodies only. 4. Becaufe certain fentient extremities are fo conftantly preferved in a certain flate, as to render them more fenfible to a change.

These circumstances determine the mode of impulse, but do not account for the senfation arising from it.

LIX.

Different fenfations are accompanied with different judgements concerning the bodies making imprefiion, and the part of the human body upon which it is made. Some fenfations are referred to bodies at a diftance; others, to external bodies in contact; and others to the feeling body itfelf.

LX.

When fenfations are referred to our own bodies, it is in three feveral ways: They' are most commonly referred to the part on which immediately the impression is made: and this, with regard to the external parts, very accurately; but, with regard to the internal, much lefs fo: And commonly the fensations arising from internal parts, are referred to the incumbent external part, with some obscure distinction between fuperficial and more deep. 2. Senfations are fometimes referred, not to the part upon which the impreffion is immediately made; but to a diftant more sensible part, to which a motion is propagated from the part impressed. 3. As fensations usually arife from impreffions made upon the extremities of the nerves, and are referred to these; so impressions made on the nerves

in

49

50

in their courfe, are fometimes referred to the extremities from whence they had commonly arifen.

LXI.

The fenfations of confcioufnefs (XL.1.2.) are referred to the encephalon. So are thofe of XL. 3. if they are moderate; but, if more vehement, they are often referred to thofe parts in which their effects are exerted, as the heart and organs of refpiration. The fenfations (XL. 4. and 5.) are feldom, with accuracy, referred to particular parts, but indiffinctly to a whole member. We are not confcious of the action of particular mufcles, except when their contraction is fpafmodic.

LXII.

We are disposed to combine our senfa-

1

tions

tions as united in one object; and thus form what are called COMPLEX IDEAS.

LXIII.

We compare our feveral fenfations, and from thence acquire new fensations of **RELATION.**

LXIV.

When fenfations formerly received are again renewed by the fame objects, it is, for the most part, with a confcioufness of their having been formerly received; and this faculty we call REMINISCENCE.

LXV.

Perceptions formerly received can be renewed without the prefence or action of the object which formerly gave occasion to them

them: and if this is attended with the confcioufnefs of a difference between the vividity of the two perceptions, and particularly of the abfence of the original objects, fuch a renewed perception is called an IDEA; and the faculty by which this renewal is made, is called MEMORY.

LXVI.

Perceptions formerly received, can, without the prefence of the original object, be renewed alfo in fuch a manner, that the mind does not perceive any difference between the original and the renewed perception; and therefore, fuch renewal is always attended with the perfuafion of the prefence of the object. The faculty by which fuch renewal is made, we call IMA-GINATION, more ftrictly.

the object which ferrearly gave occation to

LXVII.

Unable to display this page

to the different states of the brain, very little known.

LXIX.

Imagination feems always to depend upon internal caufes, that is, upon caufes acting in the brain.

LXX.

Memory and imagination renew diftinctly the ideas of feeing and hearing only. All others are renewed imperfectly, or not at all; but all others may be affociated (LXVIII.) with the fenfations or ideas of feeing and hearing, fo that thefe become figns of the others. The memory, in renewing thefe figns, fo far renews the idea belonging to them, as to renew their feveral affociations and relations; to renew, in fome degree, the pleafure or pain which formerly attended

ed the fenfations themfelves; and particularly to renew the emotions of mind, or motions of the body, which the fenfations formerly produced.

LXXI.

Most of our sensations, perhaps all of them, are either pleasant or painful.

LXXII.

The words *pleafant* and *painful* are commonly generic terms, each of them comprehending a great many fpecies, which feem to require being afforted under feveral different genera. Thus, in the first place, our fenfations may be divided into those we defire, and those we are averse to. Of those we defire, we may diffinguish those which arise from qualities we refer to other bodies, from those we refer en-D 2 tirely

tirely to our own. The first may be named more firifly the AGREEABLE, the last the PLEASANT. In like manner, of -the fenfations we are averse to, we may diflinguish the DISAGREEABLE and the PAINFUL. But, farther, the laft muft be diftinguished from the fense of averfion, which accompanies certain fenfations of confcioufnefs, as the fenfe of debility, laffitude, difficulty, &c. and particularly from that which is referred obfcurely to internal parts, and this we name ANXIETY. Thefe fenfations may be called the UNEASY; and every one diftinguishes this kind from that of the PAINFUL, more firicity fo called. Thefe laft feem to be always fenfations of impreflion, referred pretty accurately to a particular part.

There is thus a foundation for eftablifhing different genera of the fenfations we defire, and of those we are averse to; as also, for greater precision in the employment

ment of terms: but the fixing the limits of thefe genera, and afforting the feveral fpecies, may be still difficult; fo that we cannot be certain of applying the terms every where with strict propriety.

LXXIII.

The enumeration of the agreeable or difagreeable, and even of the pleafant fenfafations, would not be of much ufe here; and the enumeration of the uneafy and painful, though much more interefting, belongs to the pathology. However, we think it proper to deliver here the few following propositions.

LXXIV.

Senfation and action, within certain limits, are always defired; and the want of fenfation, or imperfect and indiffinct fen-2 D 3 fations,

58

fations, are always uneafy. In action of every kind, the fenfations of debility and difficulty are alfo uneafy.

LXXV.

In fenfations of impression, their being pleafant or painful often depends on the degree of force in the impression, allowance being made for the fensibility of the fystem.

LXXVI.

As imprefions, by being repeated, produce weak fenfations (LIV.), imprefions, at firft painful, may, by repetition, be changed into pleafant, and the pleafant into infipid and uneafy. Hence arifes, with regard to moderate imprefions, the pleafure of novelty, the defire of variety, and the the defire of increasing the force of pleasant impressions.

PHYSIOLOGY.

59

LXXVII.

There is a condition of imprefilons, rendering them objects of defire or averfion, that cannot with certainty be referred to their force. This condition we call the quality of imprefilons.

LXXVIII.

Impreffions are often rendered objects of defire or averfion, by combination, fucceffion, and relation.

LXXIX.

No fenfations arife originally in the mind, without a previous change in the state of the body.

D4 LXXX.

LXXX.

Certain impreffions, and certain flates of the body, like to those which produce the fensations of consciousness, may both of them act upon the nervous system, without producing any fensation.

SECT.

PHYSIOLOGY. 6r

SECT. II.

CHAP. II.

OF THE ACTION OF MOVING FIBRES.

LXXXI.

THE moving fibres (XXIX. 4.), fo far as yet known, are of one kind only, and the fame every where, as in the most commonly known muscles. Hence, the terms moving and muscular fibres are of the fame import.

LXXXII.

LXXXII.

A mulcular fibre is fuppofed to have a peculiar organization, different both from that of the fimple folid fibre, and from that of the medullary fibres in every other part of the nervous fystem; but wherein that peculiarity of organization confist, is not yet afcertained.

LXXXIII.

A mufcular fibre is endued with a contractility, which is different from that of the fimple folids, or of any inanimate elaftics, efpecially in this, that the contraction of a mufcular fibre is excited by caufes which do not affect thefe others. For, the contraction of a mufcular fibre is excited by being extended ; and a contraction is produced, whilft the ftretching power

power continues to be applied. The fame contraction is alfo excited by various applications, whofe mode of action we do not perceive; but we know them to be fuch as do not affect inanimate elaftics.

In refpect of these causes by which it may be excited, the contractility of mufcular fibres hath been called IRRITABI-LITY.

LXXXIV.

The force of contraction in mulcular fibres is often much greater than that of the caufes exciting it.

LXXXV.

The contractility of mulcular fibres (LXXXIII). appears especially in living bodies, ceases with life, or soon after, and is probably never produced but with life.

life. Hence it is, that, by fome writers, it is called the VITAL POWER; and the mulcular fibre endued with it, the LIVING SOLID. *Gaub. Patholog.* 169, 170.

LXXXVI.

The contractility (LXXXIII. LXXXIV. and LXXXV.) hath been fuppofed to belong to mufcular fibres, independently of their connection with the other parts of the nervous fyftem; and, upon that fuppofition, it hath been called the vis infita. We fhall call it the INHERENT POWER; Haller. Prim. lin. 400.

LXXXVII.

The contraction of mulcular fibres can be excited by applications made to other parts of the nervous fystem, as well as to the mulcles; and, as the effects of those ap-

applications made to the other parts of the nervous fyftem can be prevented by ligatures made upon the nerves between the place of application and the mufcle to be moved, it is concluded, that the contraction of mufcular fibres can be excited by a power communicated to them by a motion propagated along the nerves. This power is called the NERVOUS POWER.

LXXXVIII.

The nervous power (LXXXVII.) is commonly determined to motion by the will. This we fuppofe to act in the brain only (XXXIII.), and to depend upon fenfation, and other modifications of thought; and this power, which is to be chiefly referred to the mind, and acts in the brain only, we name the ANIMAL POWER.

LXXXIX.

LXXXIX.

The facility with which the contraction of mufcular fibres can be excited, and the force with which it can be performed, are to be diftinguifhed. The firft we name the MOBILITY, the laft the VIGOUR, of mufcular fibres. Both have been confounded under the name of Irritability.

XC.

The mobility and vigour of mufcular fibres (LXXXIX.) can both of them be increafed or diminifhed by various means. Whatever can excite the contraction of mufcular fibres is called a STIMULUS; and, in general, the means of exciting contraction are called STIMULANT POW-ERS. The means diminifhing the mobility and vigour of mufcular fibres are called SEDATIVE POWERS.

I

XCL.

XCI.

The inherent power (LXXXVI.) is fuppofed to be more vigorous, moveable, and permanent, in certain muscular fibres than in others.

XCII.

The inherent power, or the contraction dependent upon it, can be excited by certain applications made either to the mufcles themfelves, or to the nerves connected with them; and, in either cafe, the effects of fuch application are fo exactly the fame as to allow us to conclude, that the matter in the nerves, and in the mufcular fibres, is of the fame kind.

XCIII.

The muscular fibres are sensible to various impressions, and are otherwise organs of

of the fenfations of confcioufnefs (XL.4.5.) From this alfo, it is prefumed, that the mufcular fibres confift of the fame matter which is the fubject of fenfe in other parts of the nervous fyftem (XXXIX.)

XCIV.

From XCII. XCIII. and other confiderations, we think it probable, that the mufcular fibres are a continuation of the medullary fubftance of the brain and nerves, as alleged XXIX.

XCV.

Though the muscular fibres confift of the fame kind of matter as that in the nerves, the latter show no contractility, because they have not the peculiar organization (LXXXII.) of the former.

XCVI.

XCVI.

The nervous power (LXXXVII.), and the inherent (LXXXVI.), may fubfift for fome time without any connection of the nerves or mufcles with the brain; and they fubfift alfo in entire bodies for fome time after life has feemingly ceafed. Both powers, however, are feemingly of equal duration in thefe refpects; and neither power feems to fubfift long but in entire and living bodies.

XCVII.

From what is faid (XCII.—XCVI.), it is probable, that the nervous and the inherent powers are fomewhat of the fame nature; and it is alfo probable, that, in entire and living bodies, both the nervous and inherent powers have a conftant dependence upon the animal (LXXXVIII.) E XCVIII.

XCVIII.

The contraction of mulcular fibres does not depend immediately on the motion of the blood, as it fubfifts in many animals after all motion of the blood has ceafed.

XCIX.

The contraction of muscular fibres does not depend on the inflation of vesicles, or other such analogous structure, as the shortening of the fibres in contraction is often greater than take place in such structure.

C.

As the force of cohefion in the mufcular fibres of living animals is much greater than in those of dead ones, it is probable from this and other confiderations, that the

71

the caufe of mulcular contraction is an increafe only of that fame power which gives the contractility of the fimple folids, and of other inanimate elaftics; *Haller. Prim. Lin.* 407 408.

If this is true, it will alfo explain why the force of cohefion in mufcular fibres is greater than that of the medullary fibres in any other part of the nervous fystem, though both kinds of fibres (by XCIV.) confift of the fame kind of matter.

CI.

In living and healthy animals, the mufcular fibres have a conftant tendency to contract; and this tendency we call their TONE, or TONIC POWER.

CII.

The tonic power of muscular fibres ne-E 2 ceffarily

ceffarily fuppofes their being conftantly in a ftate of extension beyond their natural or most contracted ftate; and in this state they are constantly kept by the action of antagonist muscles, by the weight of the parts they fustain, by fluids distending the cavities they furround, and by their connection with fuch distended cavities, particularly the blood-vessel.

CIII.

As the differition of mulcular fibres (by LXXXIII.) proves a stimulus (XC.), we conclude, that the tonic power in them will *cæteris paribus* be in proportion to the degree of tension (CII.)

CIV.

If the inherent power (as in XCVII.) is in dependenence upon the nervous and ani-

animal powers, and thefe may be increafed or diminished by various means, the tonic, as a part of the inherent power, must in fome measure be in proportion to the state of the nervous and animal powers.

CV.

If the tonic power of any mulcular fibre depends more upon its flate of tenfion (CIII.) than upon the flate of the nervous and animal powers (CIV.), fuch fibres will be more affected by changes in the flate of tenfion, than by changes in the flate of the nervous and animal powers; and on the contrary, &c.

CVI.

The force of contraction, or the vigour of mufcular fibres, will be always as the force of ftimulus, and the vigour of the E 3 ani-

animal, nervous, and inherent powers taken together.

CVII.

The mobility of mufcular fibres (LXXXIX.) feems often to be increafed by caufes which weaken their vigour; and, therefore, it is induced by the diminution of tenfion, and by caufes weakening the animal, nervous, or inherent powers.

CVIII.

The ordinary contraction of mufcular fibres is difpofed fpontaneoufly to alternate with a relaxation or extension of the fame.

CIX.

In the ftraight mufcles and in the heart, the alternate contractions and relaxations readly take place; and that though a ftimulus is conftantly applied : but in mufcular fibres furrounding cavities, as in the alimentary canal, bladder of urine, &c. the alternate motions do not appear, unlefs a portion of the fibres is cut out, and feparated from the reft.

CX.

From a difference in the ftate of a mufcle contracted by inherent power, while the member it fuftains is moved by external force, and that of the fame mufcle contracted by the power of the will, we perceive that in the mufcles there may be a ftate of relaxation without their extenfion.

CXI.

75

CXI.

There is a flate of the contraction of mufcles that is not difpofed fpontaneoufly to alternate with relaxation, and in which too the fibres do not eafily yield to extending powers applied: Such a flate of contraction is called a SPASM.

CXII.

When mufcles are excited to contraction by preternatural cafes, and are contracted with unufual velocity and force, and efpecially when the contractions, alternating with relaxation, are frequently and preternaturally repeated; fuch motions are called CONVULSIONS.

CXIII,

If the contraction of muscles are exert-

ed

77

ed with unufual force, and fuch contractions are often repeated, they in a fhort time become uneafy and weaker; and though coutractions are not exerted with unufual rorce, if they are often repeated, and for a long time, without an interval of reft, they alfo become uneafy and weaker.

CXIV.

Within certain bounds, with refpect to force, frequency, and duration, the contraction of muscles, by being repeated, is performed with more facility and force.

CXV.

Are not the contractions of muscles produced by the action of the animal power; those which are more especially liable to become uneasy and weak by frequent repetition?

SECT. II.

CHAP. III.

OF THE FUNCTIONS OF THE BRAIN.

CXVI.

FROM the effects of ligatures made upon the nerves, and of the deftruction of their continuity, it appears, that, in their entire ftate, motions may be communicated from the brain to the other parts of the nervous fyftem, and from the latter to the former; and, from the fame experiments it appears, that the brain (XXIX. 1.)

is

is the organ of fenfation and volition, and of the feveral intellectual operations intervening between thefe: All which is confirmed by the effects of the organic affections of the brain upon the intellectual faculties.

CXVII.

The brain is thus the fenforium or corporeal organ, more immediately connected with the mind; and, fo far as a corporeal organ is employed, all the operations of thought arifing in confequence of fenfation are operations of the brain, and are modified by its various condition; *Boerb. Inft. Med.* 581. *Haller, Prim. Lin.* 570. *Gaub. Bath. Med.* 523. See afterwards (CXXII.)

CXVIII.

As certain impressions act on the nervous

So PHYSIOLOGY.

vous fystem, without producing any fenfation (LXXX.); and as, at the fame time, there is hardly any communication between the different parts of the nervous fystem, but by the intervention of the brain; it is from hence also probable, that the brain, by its organization, is fitted to propagate the motions arifing in one part, to the other parts of the nervous fystem; and, as these mechanical communications produce different effects, according to the different state of the brain itself, we, upon the whole, conclude, that the brain is a corporeal organ fusceptible of various conditions, and thereby of confiderable influence in most of the phenomena of the nervous fystem.

CXIX.

The action of the brain, in moving the feveral parts of the body, is excited by various

PHYSIOLOGY. Si

various causes, or by the same causes in different circumstances.

1. It is effectially excited by the WILL, directing the motion of certain parts as means to an end.

As the motion of certain parts is adapted to various purposes, we are confcious of willing these purposes, as they occasionally occur, and fo far alfo of the motion of the parts concerned in them: but, where the motion of the parts is connected with one fenfation, or a few only, the motions required follow these fensations without our being confcious of fpecially willing them; and, unlefs we have continued the practice of adapting the motions to different purpofes, we lofe the power of doing fo, and the motions become unavoidably connected with those fenfations which, for a long time, had alone given occasion to them. In most of the inftances of what are called VOLUN-TARY

TARY MOTIONS, we are confcious of willing the end propofed, more than the motions excited; and, of the motions produced, we are confcious chiefly of those of a whole member, or of the general effect, and very little of the many particular motions that concur to produce it. We are never confcious of the particular muscles employed.

2. The action of the brain is excited by the more general and vehement volitions named EMOTIONS and PASSIONS. Upon occasion of these, the confcious field of willing the particular motions produced is always much less diffinct, and in many cases is not at all perceived. Of the last kind are, particularly, most of the expresfions of the passions in the countenance and gesture.

3. By the difposition of human nature to IMITATION. This imitation is fometimes involuntary, often without confciousnefs;

nefs; and the confcioufnefs which takes place is often of the general purpofe only, without that of the particular motions produced; or, at leaft, it is of these only as a general effect.

4. By APPETITES or defires, directed to certain external objects, and arifing from fenfation, without any reafoning directing to an end; at leaft, without any other end in the first instance butthat of the gratification of the defire.

5. By certain PROPENSITIES or defires to remove an uneafy or painful fenfation, in confequence of which motions are excited, which are not directed to any external object, but confined to the body itfelf.

Thefe motions are not forefeen; nor are we ever confcious of willing any thing, but the general effect. Of this kind, the chief are the motions of fneezing, coughing, fighing, hiccuping, vomiting, voiding urine and feces, yawning, ftretching 2 (pan-

(*pandiculatio*), and those motions of restlessness and inquietude which pain and uneasiness produce. Weeping and laughing are expressions of emotion and passion.

In all thefe, as well as in the motions of N° 4. fome volition is concerned, not only as they can often be prevented by another volition prefenting itfelf; but, befides, as the feveral motions which occur in executing thefe propenfities are more or fewer, and more or lefs forcible, according to the vehemence of the propenfity or effort. Very often the ftimulus to thefe propenfities is irrefiftible; and, unlefs the peculiar ftimulus is prefent, the motions cannot be produced by any volition.

6. By certain internal impressions arising from the exercise of the functions of the body itself, which produce no sensation, nor produce motions of which we are confcious, except when exercised in an unusual manner. Such are the causes of the motions

85

tions of the heart and arteries, of the organs of refpiration, of the ftomach, inteftines, and perhaps of many other parts. With regard to most of these motions, it may be fupposed, that they are the mechanical effects of their feveral causes, acting upon the inherent power of muscular fibres (LXXXVI.): but it is sufficiently certain, that they also depend upon an action of the brain; and the effects of pasfions, as well as the effects of destroying or compressing the nerves of the organs concerned, are proofs of it.

The motions mentioned in this article are commonly fuppofed not to be accompanied with any volition of which we are diffinctly confcious. This perhaps is not ftrictly true with regard to most of them; and, fo far as it is, it may be imputed to that repetition which deftroys confcioufnefs (LV.): But neither can we entirely adopt this explanation; as these F motions,

motions, which are intended to follow one ftimulus only, may be fuppofed to require no exercife of volition, as in the cafe of the heart, arteries, and alimentary canal, while the action of refpiration, as adapted to various purpofes, continues to be a voluntary motion.

7. By various occafional imprefions of external bodies, and by various occafional flates of the fyftem, or of its particular parts which excite motions, not only in the parts to which the imprefions are immediately applied, but alfo in diftant parts, on which they can operate only by the intervention of the brain. Some of these causes operate with, others without, sensation or volition.

CXX.

In all, or any of thefe cafes in which the action of the brain takes place, we do not per-

perceive the manner, that is, the mechanical means, by which the feveral caufes produce their effects; and we perceive only an inftitution of our Maker, eftablishing a connection between the feveral caufes and the motions which enfue. At the fame time we, for the most part, perceive, that the connections established are fuited. to the purpose of the animal œconomy; and particularly, to the purpose of supporting the fystem in a certain condition for a certain time, and of averting what might hurt or deftroy it. This conftitution of of the animal ceconomy we call NATURE; and every where in the æconomy we perceive the vires confervatrices and medicatrices naturæ, fo juftly celebrated in the fchools of phyfic.

CXXI.

It is in consequence of this constitution, It is in confequence of this constitution, It is in confequence of this constitution,

that not only impulfe, and other caufes which may be fuppofed to produce motion, do accordingly excite motions in the animal œconomy; but that alfo many caufes which feem to diminifh motion, do, however, produce an increase of motion in animal bodies. Thus feveral passions which, in their first tendency, diminish motion, feveral propensities arising from debility and difficulty of action, the absence of usual impressions, evacuations, and other caufes of relaxation, cold and narcotic powers, are all of them caufes of confiderable motions arising in the animal fystem.

CXXII.

As the mechanism of the brain, fuited to its feveral functions, is not at all perceived (CXX.); and, at the fame time, as very few of these functions are carried on without sensation and volition, it must appear

pear from this, and many other confiderations, that the mechanifm of the brain (CXVII.) would not be fufficient for the purpofe, without being united with a fentient principle, or mind, that is conflantly prefent in the living fyftem. But, at the fame time, it is with little probability alleged, that the administration of the corporeal functions is entirely directed by the mind acting independently of the body, and with intelligence perceiving the tendency of imprefions, and exciting fuch motions as may favour the beneficial, or obviate the hurtful tendency of all caufes acting upon the body.

We are certainly confcious of no fuch administration. Many impressions have their effects without fensation or volition. In most cases where volition takes place, it is very general, with little confcious field for the motions excited, and none at all of F 3 the

the organs employed. The force of impreffion is every where abfolute; and it is according to the force of impreffion, and other mechanical conditions of the fystem, that the motions excited prove either falutary or pernicious. The general principle, therefore, is ill-founded; it is not neceffary, (vide Stabl. Præf. ad Junker. Confp. Med.); it can be of no use, and may be hurtful to the fystem of physic.

CXXIII.

The action of the brain is often determined and regulated by cuftom and habit; that is, by laws eftablifhed by frequent and uniform repetition. See above XLIII.LIV. LV. LVI. 5. LX. 3. LXVIII. and LXX. for the effects of cuftom on fenfation, and CXIV. for one effect of it on the action of moving fibres. It is now to be obferved further, 1. That cuftom determines the Unable to display this page

mined. Thefe laws, which may be eftablished by custom, are, many of them, with difficulty avoided; they are often rigidly fixed, have a confiderable influence on the action of the brain, and govern the revolutions of the animal fystem. Thus, any caufes producing a deviation from the ufualdegree of force and velocity (Nº 4.) are apt to deftroy the measure of it altogether; and, in like manner, caufes producing a deviation from the ufual order and velocity in the fucceffion of motions (N° 5.) are apt to deftroy the power of the mind in following that order, or in giving any measure to the feveral motions which should be performed; and perhaps, in this way, the effects of debility, of feveral passions, and of surprise, are in some meafure explained.

CXXIV.

CXXIV.

The brain feems, by its conflication, to be difpofed to the alternate flates of reft and activity; as appears in the alternate flates of fleep and waking, which conflantly take place in every animal: but wherein this conflication confifts, it is difficult to difcover.

CXXV.

The moft common opinion is, that the brain is a fecretory organ, which fecretes a fluid neceffary to the functions of the nervous fyftem; that this fluid is alternately exhaufted and recruited, and thereby gives occafion to the alternate flates of fleep and waking. But this fuppofition is attended with many difficulties. 1. It is probable, that the nevous fluid exifted in the animal embryo, before the action of the heart, or

any

any fecretory function, could take place. 2. In animals which during the winter fuffer a temporary death, fuch as bats, when by heat they are again reftored to life, the vital power of the folids is reftored before the fluidity of the blood. 3. The nervous fluid fubfifts in the nerves and muscular fibres long after they are feparated from the brain, and often when cut into many fmall parts. 4. Though it be true that the brain is a fecretory organ, the fluid fecreted may be deftined to another purpose; and fo far as we understand that purpose, the fluid fit for it, must be unfit for the purpose of sense and motion. 5. There is no appearance, in any part of the nervous fystem, of provision made for an occasional accumulation of the fecreted fluid; nor is there any evidence of its actually taking place. 6. The phenomena of fleep and waking do not correspond with fuch a fupposition; as fleep

theep often takes place when the fecreted fluid must be copiously present, and waking can be protracted when the fluid is exhausted much beyond its usual measure. 7. Both states are induced by many causes, which can hardly be supposed to act upon secretion.

CXXVI.

A certain compression of the brain can produce a state of the fystem refembling sleep; but that state is in some respects, different from that of ordinary sleep: And it does not by any means appear, that natural and ordinary sleep depends upon any compression of the brain.

CXXVII.

As it is therefore probable that fleep and waking do not depend upon a different quantity

quantity of the matter of the nervous fluid for the time prefent in the fyftem (CXXV.), or upon any caufes interrupting its motion, while the condition of the matter remains the fame (CXXVI.); we are difpofed to believe, that those flates of fleep and waking depend upon the nature of the nervous fluid itself capable of becoming more or less moveable; that it is chiefly in the brain fusceptible of these different conditions; and that, especially by its condition there, it has its more general effects on the whole fystem.

CXXVIII.

This may perhaps be confirmed by confidering the remote caufes of fleep and waking: And it appears, that cold, the abfence of impressions, attention to a fingle fensation, or to fensations that have no confequence in thought or action; the finished

finished gratification of all vehement defires, fedative fensations and impressions, evacuations, relaxation, and any violent, frequent, or long continued exercise of the animal power; are all of them, feverally or together, causes inducing fleep.

CXXIX.

On the other hand, it appears, that a certain degree of heat, all fenfations of impreffion, impreffions analogous to those which produce fensation, all fensations which lead to thought and action, and the increased impetus of the blood in the veffels of the brain, are all of them causes favouring or inducing a state of waking.

CXXX.

As most of the causes CXXVIII. are evidently such as diminish motion in the r brain,

brain, and those of CXXIX. are fuch as increase it; it is from thence probable, that the nervous fluid in the brain is truly capable of different flates or degrees of mobility, which we shall call its flates of EX-CITEMENT and COLLAPSE; but, without intending, by these terms, to express or determine any thing with regard to the nature of the nervous fluid, or wherein its different flates confist.

CXXXI.

This fubject may be further illustrated, by obferving, that the excitement of the brain appears to be in very different degrees on different occafions. It feems to be greatest in certain maniacs endued with uncommon strength, refisting the force of most impressions, and with the utmost difficulty admitting step.

CXXXII.

99

CXXXII.

A leffer degree of excitement occurs in the ordinary flate of waking men in health, when the excitement is total with refpect to the functions of the brain, but readily admitting of fleep. This excitement may be confidered as of two kinds; either as it refpects the vigour, or as it refpects the mobility of the fyftem : And thefe different flates of the brain are expressed in the body, by ftrength or debility, alacrity or fluggifhness; and in the mind, by courage or timidity, gaiety or fadness.

CXXXIII.

A degree of collapfe takes place in the cafe of natural fleep, when the collapfe prevails fo far as to fufpend very entirely the exercife of the animal functions; and, though the exercife of the vital and natural

ral continue, they are confiderably weakened. The partial collapse that may take place in the brain difcovers itself by the delirium which appears in a state that often occurs as intermediate between step and waking; and even in step the collapse with respect to the animal functions, takes place more less entirely; whence the step is with or without dreaming, and the dreaming is more or less active.

CXXXIV.

A ftill greater degree of collapfe takes place in the cafe of fyncope; in which it is fo great, as to fufpend the exercife of the vital functions concerned in the circulation of the blood, notwithftanding the force of habit in thefe, and their being exposed to conftant ftimuli. Here the collapfe may be very confiderable; but there ftill remains fome degree of excitement

ment while the brain can be acted upon by ftimuli, which act only on vital powers, and while its ufual excitement is ftill recoverable by fuch ftimuli.

CXXXV.

If the collapse is still more complete and irrecoverable, it is the state of DEATH.

CXXXVI.

From what is now faid of the excitement and collapse of the brain, it will appear, that we suppose LIFE, so far as it is corporeal, to confiss in the excitement of the nervous system, and especially of the brain, which unites the different parts, and forms them into a whole. But, as certain other functions of the body are necessary to the support of this excitement, we thence G learn,

learn, that the caufes of death may be of two kinds; one that acts directly on the nervous fystem, destroying its excitement; and another that indirectly produces the fame effect, by destroying the organs and functions necessary to its fupport. Of the first kind are chiefly the caufes of sleep operating in a higher degree; as cold, fedative passions, poisons, and all caufes of very violent excitement.

CXXXVII.

This fubject may receive fill further illuftration from confidering the flate of the other parts of the nervous fystem with respect to excitement and collapse. In the nerves strictly fo called (XXIX. 2.), we do not know that the nervous fluid fuffers any change, but what is exactly correspondent to its flate in the brain and extremities; and

and therefore the only difference of the ftate of the nerves to be taken notice of is their being more or lefs free (XXX.)

CXXXVIII.

In the fentient extremities of the nerves (XXIX. 3.), a difference of the flate of the nervous fluid arifes from the feveral caufes mentioned (LVI. 2. 3. 4. and 5.), which give a different degree of fenfibility; and it is probable, that thefe different flates of the fentient extremities are analogous to the different degrees of excitement in the brain.

CXXXIX.

The moving extremities, or mufcular fibres (XXIX. 4.), may also be in a different condition with respect to excitement. It is probable, that their constitution is G 2 fuch

fuch as to admit of a higher degree of ezeitement than any other portion of the nervous fystem; and that upon this their contractility depends. But, whatever is in this, we perceive very clearly, that the condition of mufcular fibres may be varied by caufes affecting their tonic power (CI.), or their vigour and mobility (LXXXIX.), and by the effect of cuftom (CXIV.); and it is probable, that the states produced by these causes are analogous to the different degrees of excitement in the brain (CXXX.) and in the fentient extremities (CXXXVIII); and thus the feveral parts of the nervous fystem (XXIX.), as they confist of the fame kind of matter (XCIV.), are alfo fubject to fimilar conditions.

CXL.

The beginning of motion in the nervous fystem is most commonly accompanied 2 with

with fenfation, and the force of this in producing its feveral effects is more or lefs, 1. According to the force (XLII.), quality (LXXVII.), and novelty (LIV.) of imprefion. 2. According to the fenfibility of the fentient extremity and brain (LVI.) 3. According to the ftate of attention (XLVII.) Thefe feveral caufes often concur, frequently balance one another, and are always to be confidered together.

CXLI.

The effect of fenfation is commonly that of exciting the action of the brain; and this is generally according to the degree of volition produced under the different circumftances mentioned in CXIX.

CXLII.

The action of the brain excited by vor G 3 lition

105

lition or other caufes, is determined to particular parts of the body, most constantly by the connections established in the fystem (CXX.), but also occasionally, by acquired habits, or the greater mobility of certain parts.

CXLIII.

With regard to the connections eftablifhed in the fyftem (CXX. CXXI.) it is to be obferved, as of great confequence in pathology, that certain parts of the body which have a common function and conflitution, have thereby a peculiar relation to the brain, fo as to be more liable to be affected by the different flates of it, and in their turn by the difference of their condition to affect the brain. Such are, efpecially, the organs of voluntary motion; the alimentary canal, and efpecially the flomach; the circulatory fyftem, and particularly

ticularly the extremities of the veffels on the furface of the body; the uterine and genital fystem in females; and some others to be mentioned in the pathology.

CXLIV.

The communications of motion between the feveral parts of the nervous fyftem which have been mentioned as inftances of a particular fympathy between thefe parts, are very feldom to be explained by any contiguity or contact, either in the origin or courfe of the nerves of the communicating parts; and more commonly they may be explained, by fuppofing the action of the imprefion to be general with refpect to the brain, and that the affection of particular parts depends upon the caufes of determination (CXLII. CXLIII.)

When the action of feveral parts, together or fucceffively, are neceffary to pro-G 4 duce

duce one effect, thefe concur, though the ftimulus exciting the action of the brain is applied to one fingle part only; and commonly no other caufe of communication can be affigned, but the feveral motions being neceffary to the execution of the volition, propenfity, &c. arifing from the ftimulus.

CXLV.

Thefe are the chief facts and laws relative to the nervous fyftem. The whole might perhaps be illuftrated, and more exactly afcertained, by a more particular inquiry into the nature of the nervous fluid; but we are not fo confident in our opinion on this fubject, or of the application it will admit of, as to deliver it here.

SECT.

SECT. III.

OF THE CIRCULATION OF THE BLOOD.

CXLVI.

THE circulation of the blood, by CXXIX. feems neceffary to the excitement of the brain; and for this, as well as other reafons, it is, next after the function of the brain itfelf, the most important of the animal œconomy.

On this fubject, we shall confider, 1. The course of the blood. 2. The powers by which

which the blood is moved. 3. The laws and general circumstances of the circulation. 4. Respiration as necessary to it.

On all these subjects we suppose the anatomy of the parts to be known.

SECT. III.

CHAP. I.

OF THE COURSE OF THE BLOOD.

CXLVII.

WOUNDS and hemorrhagies flow, that, in living bodies, the blood is in conftant motion, and flowing into any one part from many others.

CXLVIII.

PHYSIOLOGY. III

CXLVIII.

In man, and other analogous animals, which have once breathed for fome time, the courfe of the blood is very conftantly in the following manner.

From the left ventricle of the heart, the blood paffes into the trunk of the aorta, and fucceffively into the following veffels and cavities, viz. the branches of the aorta, the branches of the vena cava, the trunk of the fame, the right auricle of the heart, the right ventricle of the fame, the pulmonary artery, the pulmonary veins, the left auricle of the heart; from which laft it paffes into the left ventricle of the heart to return again into the fame courfe as before.

From all this, it appears, that in the arteries the ufual courfe of the blood is from the heart towards the extreme branches of thefe;

these; and that, in the veins, the course is in the contrary direction, from the extreme branches towards the heart.

CXLIX.

The courfe of the blood through the cavities of the heart, as above described, is not in a continued stream, but alternately interrupted and free during the contraction and dilatation of those cavities, which alternately happen. Thus, while the left ventricle of the heart is in a flate of contraction, the blood passes out of it into the aorta; but at the fame time no blood passes into it from the left auricle, which is then dilated and filled by the blood flowing into it from the pulmonary vein. It is only when the ventricle is emptied by contraction, and confequently relaxed, that the blood paffes into it from the auricle, urged by the contractions of the auricle and

and adjoining finus venofus, which fucceed immediately to that of the ventricle. During this contraction of the auricle and filling of the ventricle, no blood paffes from the ventricle into the aorta, nor till a contraction of the ventricle fucceeds in confequence of its being filled. The fame circumftances take place with regard to the right ventricle and auricle of the heart, and precifely at the fame times; for it appears, that the two ventricles of the heart are contracted and relaxed at the fame time; and, in like manner, the two auricles.

CL.

That the courfe of the blood, as defcribed CXLVIII. and CXLIX. is its ufual and conftant courfe, appears from infpection of the heart in living animals; from the fituation of the valves of the heart;

heart; from the fituation of the valves at the orifices of the aorta and pulmonary artery; from the fituation of the valves of the veins; from the effects of ligatures made upon the arteries and veins; from the effects of hæmorrhagies of the veins; from obfervations with the microfcope; and, laftly, from experiments of infufion and transfufion in living, and of injection in dead animals.

ore contracted an .I.I. azed at the house

pears, that the two ventricles of the heart.

This courfe of the blood is, however, in fome parts, and upon fome occafions, a little changed. 1. In the penis, and fome other parts, the blood does not pafs from the extreme arteries immediately into continuous veins; but is poured out into an intermediate cellular texture, from which it is afterwards received by the extreme veins. 2. In the fmall branches of the aorta,

aorta, the blood does not move constantly in a direction from the heart towards the extremities; but, in certain portions of them, is fometimes retrograde to that course. In this, it is favoured by the anastomoses frequent between the small vessels, which, at the same time, prevent this deviation from being considerable or durable, while the action of the heart continues.

CLII.

In the courfe of the venous blood, there is alfo fome variety. 1. In the fmall veins, the blood is liable to have a motion retrograde to its ufual direction (CXLVIII.), as in the arteries (CLI. 2.) 2. The blood, returning to the heart from most parts of the body, passes fucceffively from smaller into larger veins, by a feries of pretty regularly increasing vessels, till they form the vena cava entering the heart. But this

this is varied in the abdomen, where the veins carrying the blood returning from every vifcus included in that cavity, except the kidneys and genitals, unite in forming the vena portarum, whereby they undergo a peculiar distribution. 3. The veins returning the blood from the extreme arteries in the brain, do not carry it to the heart by a feries of regularly increasing vessels, but by the interposition of finuses, into which the fmall veins iffuing from the cortical part of the brain immediately pour their blood. 4. As the course of the blood through the veffels of the lungs is not at all times equally free, and particularly is confiderably interrupted at the end of exfpiration; fo at the fame time it is alfo interrupted at entering the right ventricle of the heart; and this often occafions fome regurgitation, or retrograde motion, in both the afcending and defcending cava.

CLIII.

CLIII,

The whole of the fluids carried in the aorta to its extreme branches, are not returned again by the continuous veins to the heart, as, by fecretory veffels, a part of them is constantly carried out of the course above described. Some of these fecreted fluids are thrown entirely out of the body, and others are poured into certain cavities, for various purpofes of the æconomy; and fome of these are again returned into the course of the circulation. Of these last, there is a peculiar fluid which, from the extremities of the arteries, is poured out in a liquid form, or exhaled in that of vapour, into perhaps every cavity and vacuity of the body. This, after having ferved the purpose of the effusion, seems intended to be regularly returned again into the courfe of the circulation; and ac-H cordingly,

cordingly, in all the feveral cavities into which it had been effufed, there are abforbents which again take it in. Thefe do not carry the fluid immediately into the veins; but uniting, form the veffels called LYMPHATICS, which, in their courfe, pafs through the conglobate glands, and at length terminate either in the receptacle of the chyle, in the thoracic duct, or in the left fubclavian vein; and, in this way, return the abforbed fluid into the courfe of the ordinary circulation.

CLIV.

There are abforbent veffels, not only in all the feveral cavities, but alfo on the external furface of the body, by which many extraneous matters may be introduced into it.

CLV.

CLV.

Moft of the fluids fecreted from the circulating mafs, and poured into cavities, may be abforbed from thefe, and returned again by the lymphatics, as in CLIII. to the courfe of the circulation. But the fame fecreted fluids feem often to be returned alfo into the courfe of the circulation by a regurgitation, or retrograde motion, in the excretory and fecretory veffels.

Ha

SECT.

it's

SECT. III.

CHAP. II.

OF THE POWERS MOVING THE BLOOD.

CLVI.

THE chief power by which the blood is moved, and the circulation carried on, is the action of the heart, or its repeated contractions occurring, as mentioned CXLIX. For this purpofe, the heart is a mufcular part; the action of which may depend upon an inherent power ftimulated by the dilatation of its cavities; but this in-

inherent power requires the conftant fupport of the nervous and animal powers, and is often actuated by thefe.

CLVII.

The contraction and relaxation of the heart, or, as thefe are called, its fyftole and diaftole, are neceffarily alternate by the general law (CVIII.) affecting all mufcles, and by the ftimulus from the influx of venous blood being alternately applied and removed.

CLVIII.

If we may be allowed to effimate the vigour of mufcles by the number of their fibres, we must suppose the force of the heart to be very confiderable; but it is very difficult to obtain any exact estimate of its absolute force. It is perhaps the re-H 3 lative

lative force only that we are concerned to know.

CLIX.

Do the arteries, by their contraction, contribute to promote the motion of the blood begun by the heart? They cannot do it by the elafticity of the fimple folid in their structure; and can do it only by their being endued with a muscular power, whereby they may, in their contraction, communicate to the moving blood more than was loft of the force of the heart in dilating them. That they are indued with a muscular contractility (LXXXII.), is probable from the appearance of the muscular fibres in their ftructure; from their irritability appearing in the experiments of Verschuir ; from their becoming flaccid on tying the nerves belonging to them; from the motion of the blood

blood being fupported, when the force of the heart is confiderably weakened; from the motion of the blood becoming languid, when the action of the arteries is deftroyed; from the velocity of the blood in the extreme arteries being greater than was to be expected from the velocity of the blood iffuing out of the heart, exposed to fo many causes of retardation as constantly occur; and, lastly, from the velocity and impetus of the blood in different parts of the body, and at different times, being unequal, while the action of the heart continues the fame.

It is probable, that the muscular fibres of the arteries become more irritable as the arteries are more diftant from the heart.

CLX.

CLX.

The tone and action of the arteries, as a mulcular part, may be increased by flimuli immediately applied to them, or by the increased force of the nervous and animal powers with regard to them; and they may be diminished by fedative powers applied, or by weakening the nervous and animal powers.

CLXI,

There does not appear to be any ofcillatory motion in the extreme arteries independent of the action of the heart.

CLXII.

There does not appear to be any operation of capillary attraction in the extreme arteries; nor does there feem to be any occafion

occasion for such a power in any part of the arterial system.

CLXIII.

The power of derivation (Vis derivationis Ill. Halleri 174.) in the fanguiferous fyftem, feems to be no other than that which arifes from the fulnefs of contractile veffels.

CLXIV.

The motion of the blood in the arteries of any particular part is promoted by the action of adjoining mufcles.

CLXV.

The blood in the vena cava, and its branches, is moved by the action of the heart and of the arteries. These powers are affisted by the action of muscles, which, in

in their contraction, prefs the veins lying between their feveral fibres; and alfo, by the fwelling of their whole mafs, prefs the adjoining veins. Thefe veins are commonly provided with valves, which determine the effect of all preffure upon them, to be the motion of the blood towards the heart.

The great trunks, both of the vena cava and pulmonary vein, are provided with muscular fibres, and manifestly endued with muscular contractility.

CLXVI.

In the abforbent veffels, the fluids are probably taken in by a capillary attraction.

CLXVII.

In the lymphatic veffels, provided with numerous valves, which necessarily determine

mine the motion of the contained fluid to be towards the heart, the fluid is moved by the preffure of the neighbouring muscles and arteries. But further, as the lymphatics are remarkably irritable, it is probable that the fluid in them is moved by a peristaltic motion begun by the action of their absorbent extremities.

CLXVIII.

The motion of the blood through the veffels of the lungs, depends upon refpiration, to be confidered hereafter.

or the left vent clear the heart in

to suttro palling light do with the

SECT.

127

SECT. III.

CHAP. III.

OF THE LAWS OF THE CIRCULATION.

CLXIX.

THE velocity of the blood paffing out of the left ventricle of the heart into the aorta, may be eftimated from knowing the quantity of blood paffing out at each fyftole, the area of the orifice of the aorta, and the time occupied by the fyftole; but none of these circumftances are exactly afcertained.

CLXX.

CLXX.

As the blood moves onwards through the arteries, the velocity (CLXIX.) fuffers a confiderable retardation from feveral caufes. 1. From the capacity of the arteries being enlarged as they are more diftant from the heart. 2. From the frequent flexures of the arteries. 3. From the angles which the branches make with the trunks from which they arife. 4. From anaftomofes. 5. From the vifcidity of the blood. 6. From the friction of adhefion. 7. From the weight and rigidity of the parts furrounding the arteries.

CLXXI.

The velocity (CLXIX.) and the caufes of retardation (CLXX.) being given, the velocity of the blood in the arteries will be

be as the frequency of the fystole of the heart.

CLXXII.

The frequency of the fystole of the heart will be more or less, 1. As the blood in the veins is more or less quickly returned to either ventricle of the heart. 2. As the ventricles of the heart are more or less entirely evacuated at each fystole. 3. As the muscular fibres of the heart are more or less moveable. 4. As the action of the nervous and animal powers are more or less increased with respect to the heart.

CLXXIII.

As the arteries of a healthy body are always full, the blood thrown out of the ventricles into the arteries during the fystole of the heart, can only find a place there

by

by pufhing on the blood with the velocity (CLXIX.), or by dilating the arteries; but as the refiftances CLXX. prevent the blood from flowing with the velocity CLXIX. the blood thrown out of the heart must, in some measure, dilate the arteries, and thereby form what is called the PULSE.

CLXXIV.

It appears, that, in the arteries to a certain length, the blood moves fafter during the fyftole than during the diaftole of the heart; but as the refiftances and caufes of retardation become greater in every portion of the arteries as it is more diftant from the heart, fo the acceleration of the blood during the fyftole of the heart muft be greater in any portion of the arteries nearer the heart than in the next adjoining that is more diftant; and fo far as this takes

takes place, a dilatation of the arteries will happen, even from a fmall quantity of blood thrown out of the ventricles.

shood right flowing with the vicery

CLXXV.

As the refiftances to the blood's motion in the blood-veffels increafe with the diftance from the heart, there may be a part of the fanguiferous fyftem in which the motion of the blood will not be accelerated during the fyftole of the heart, and in which, therefore, no pulfe can be difcerned. This happens in the extreme branches of the aorta; and no pulfe is ever obferved in the extreme branches of the vena cava.

CLXXVI.

from the heart, to the acceleration of the

The velocity and impetus of the blood in

in the whole fystem of blood-vessels will always be as the action of the heart and arteries taken together.

CLXXVII.

The velocity and impetus of the blood, in any particular part of the fyftem, will_ be, 1. As the part is more or lefs diftant from the heart. 2. As the circumftances (CLXX.) take place more or lefs in the part. 3. As the gravity of the blood concurs with, or oppofes its motion in the part. 4. As caufes increasing or diminishing the action of the arteries of the part are applied or removed.

CLXXVIII.

The quantity of blood distributed to any particular part of the fanguiferous fystem, will be greater or less according to I the Unable to display this page

different parts of the fystem, is in different proportion at different periods of life.

1. The capacity and force of the heart, in proportion to the fystem of vessels, is greater at the beginning of life than at any after period. Till the body arrives at its full growth, the capacity of the vessels increases in greater proportion than that of the heart; but from that period, the capacity of the vessels is constantly diminishing, while that of the heart suffers little change.

2. A greater quantity of blood is contained in the arteries, in proportion to that which is contained in the veins, at the beginning of life, than at any after period. From the time that the body has arrived at its full growth, the quantity of blood contained in the veins, in proportion to that which is contained in the arteries, is conftantly increafing.

3. The veffels of the head receive a I 2 greater

greater quantity of blood, in proportion to the reft of the fyftem, at the beginning of life than at any after period.

4. Any general increase of the action of the heart and arteries determines the blood more copiously to the extreme arteries on the furface of the body, than to those of the internal parts.

5. The equilibrium of the fanguiferous fyftem, with regard to the diftribution of the blood, may be changed by various caufes (CLXXVII. CLXXVIII.); and thefe caufes continuing to operate for fome time, induce a habit which renders the changed diftribution neceffary to the health of the fyftem.

6. The lymphatic fystem is fuller in young perfons than in old.

SECT.

SECT. III.

CHAP. IV.

OF RESPIRATION.

CLXXXI.

RESPIRATION confifts of the motion of infpiration, or the admiffion of air into the lungs; and of exfpiration, or the expulsion of air from the fame; alternately happening.

CLXXXII.

Refpiration takes place in man, and I 3 other

other analogous animals, foon after the infant is taken from the uterus of the mother, and is exposed to the air. After it has taken place for a little time, it is ever after neceffary to the continuance of life, as it is absolutely neceffary to the continuance of the circulation of the blood.

CLXXXIII.

The lungs are a hollow fpongy mafs, capable of confining air, and readily dilatable by it. By the wind-pipe, they are open to the atmosphere; and they are fo fituated in the thorax, that the air must enter into them, if the cavities of the thorax, in which they are placed, are enlarged. For, as there is no air in these cavities, and the external air cannot enter into them, the enlargement of the thorax must form a vacuum around the lungs, which the external heavy and elastic air will supply by enter-

entering into and dilating the lungs, while these do not allow the air to pass through them into the cavities of the thorax.

CLXXXIV.

Infpiration therefore depends upon the enlargement of the capacity of the thorax; and this is performed chiefly by the contraction of the diaphragm. This in its relaxed ftate is fufpended by the mediaftinum, and its middle tendinous part is raifed high in the thorax; wherefore, as this middle part, by the contraction of the mufcular, is moved downwards, the thorax is thereby confiderably enlarged.

CLXXXV.

The capacity of the thorax is alfo enlarged by the motion of the ribs upwards, whereby the curvatures of oppofite ribs

are

I 4

are fet at a greater distance from each other; and, by the fame motion, the sternum is moved outwards, and fet at a greater diftance from the vertebræ of the back. The motion of the ribs upwards is caufed by the contraction of both layers of intercoftal muscles. That the muscles called internal intercostals concur with the external in raifing the ribs, appears from the fituation of those muscles, from the greater mobility of the inferior ribs, from the infpection of those muscles in living animals, and from experiments imitating their action. In more violent and laborious infpirations, the raifing of the ribs is affifted by many muscles attached to the ribs, and arifing from the clavicle, humerus, fcapula, and vertebræ of the neck or back.

and the second s

CLXXXVI,

CLXXXVI.

By the enlargement of the thorax, a dilatation of the lungs is produced, in proportion to the bulk of air entering into them; but the dilatation may often be greater by the air that enters into the lungs being heated and rarified; and the greatest distension of the lungs is obtained by a constriction of the glottis confining the air that has already entered into the lungs.

CLXXXVII.

As infpiration, or the admiffion of air into the lungs, depends upon the enlargement of the thorax, the diminution of it must expel the air, or produce exspiration. The capacity of the thorax is diminished, while the muscles dilating it are spontaneously relaxed, by the elasticity of the

the ligaments connecting the ribs with the vertebræ, and by the elafticity of the cartilages and ligaments connecting the ribs with the sternum; both which powers, commonly affifted by the weight of the ribs themfelves, bring the ribs and fternum into the position they were in before infpiration. At the fame time, the elasticity of the mediastinum draws the diaphragm upwards; and the contraction of the abdominal muscles both preffes the diaphragm upwards, and pulls the ribs downwards; and, in the last, they are affifted by the fterno-coftal and infra-coftal muscles. While these powers concur in diminishing the capacity of the thorax, the expulsion of the air from the lungs is affifted by the elafticity of the lungs themfelves, and by the contraction of the mufcular fibres of the bronchiæ.

CLXXXVIII.

CLXXXVIII.

Thefe are the ordinary powers of exfpiration, which, depending upon the reaction of elaftic parts, is performed flowly, and with little force: but when it is neceffary to perform it with more velocity and force, fome other and very powerful mufcles, as the quadratus lumborum, facrolumbalis, and longiflimus dorfi, concur in pulling down the ribs; and, at the fame time, the abdominal mufcles, actuated by the animal power, are contracted with greater velocity and force than in fpontaneous exfpiration.

CLXXXIX.

The fituation of the blood-veffels of the lungs is fuch, that, in the contracted flate of this vifcus, thefe veffels must be much folded and ftraitened; and it appears, that, in

in the fœtus, where they are conftantly in a contracted ftate, their capacity is not fufficient to tranfmit, in the time required, the whole of the blood returning to the heart by the vena cava; but, after refpiration has been repeated for fome time by the dilatation of the lungs to a certain degree in infpiration, their blood-veffels are unfolded, lengthened, and enlarged, fo as to be capable of tranfmitting the whole blood of the cava.

CXC.

In the infant who has breathed for fome time, the whole blood of the vena cava paffes into the right ventricle of the heart, and from thence enters into the veffels of the lungs; but, in the contracted ftate of the lungs which occurs at the end of exfpiration, the blood cannot be proper-

ly

ly transmitted; and, for that purpose, an infpiration becomes absolutely necessary.

CXCI.

It is, however, under a certain degree of infpiration only, that the blood is freely tranfmitted through the veffels of the lungs; for, if the infpiration is full, and continued, fo that the lungs are thereby much diftended, we find that this ftate alfo interrupts the free paffage of the blood, and renders exfpiration neceffary.

Exfpiration becomes alfo neceffary; becaufe, perhaps, the air long retained in the lungs lofes a part of its elafticity, and becomes thereby unfit to keep the lungs diftended; but, more certainly, and more efpecially, becaufe, in an animal that has breathed for fome time, there is a noxious vapour conftantly arifing from the lungs, which, if not diffolved by the air, and carried

carried out of the lungs, proves pernicious to life.

CXCII.

From what has been faid, it appears, that the alternate motions of infpiration and exfpiration are neceffary to the circulation of the blood, and otherwife alfo to the health of the body; and it appears alfo, that the more frequent the alternate motions of refpiration are, the more quickly is the blood transmitted from the right to the left ventricle of the heart.

CXCIII.

We can now perceive alfo the caufes exciting thefe alternate motions; and we find no occafion for fuppoing them to arife from any caufes alternately interrupting the motions of the nervous fluid,

10

E

or of the blood, into the muscles concerned in these functions.

Infpiration, or the action of the mufcles producing it, is excited, in all cafes of general effort, to remove pain and uneafinefs; and it is, perhaps, a propenfity of this kind that gives the first beginning to respiration in the new-born infant, exposed to feveral new and uneafy impreffions.

For the continuance of refpiration, infpiration is efpecially excited by the fenfe of uneafinefs that attends any difficulty in the paffage of the blood through the veffels of the lungs; but this uneafinefs arifes, in fome meafure, at the end of every exfpiration, and is much increafed by any continuance of this ftate.

CXCIV.

Exfpiration, in fome meafure, necessarily fuc-

fucceeds infpiration, by the fpontaneous relaxation of the infpiratory mufcles (CVIII.), while the elafticity of the membranes, ligaments, and cartilages ftretched in infpiration, brings back the ribs and diaphragm into their former fituations; and the fame effects are alfo produced by the action of the abdominal mufcles, and of the mufcular fibres of the bronchia; both of which are ftretched, and thereby excited in infpiration.

In the cafe of ordinary infpiration, thefe caufes are fufficient to produce a fpontaneous exfpiration. But as it appears that a violent and long continued infpiration interrupts the paffage of the blood through the lungs, this creates an uneafinefs, and a propenfity, which muft produce a relaxation of the infpiratory, and excite a contraction of the exfpiratory, mufcles.

It is further to be fuppofed, that, in animals which have breathed for fome time,

cuftom

I

cuftom has affociated the feveral motions concerned both in infpiration and exfpiration; fo that an irritation applied to any part of them neceffarily excites the whole; and it may alfo be fuppofed, that habit determines thefe motions regularly to fucceed one another.

and svail link CXCV. Some hour out

ration on the animal fluids is delayed till

In this manner (CXCIII. CXCIV.) refpiration is continued for the general purpofes of the animal œconomy; but the feveral motions of which it confifts are alfo occafionally excited and varioufly modified by the will, intending particular effects to be produced by thefe motions. They are alfo excited, and varioufly modified, by certain emotions and paffions, and give particular expressions of thefe. They are often excited alfo by imitations; and they are particularly excited by propenfi-K

ties to remove pain and uneafinefs, which operate more frequently on refpiration than upon any other function.

CXCVI.

The confideration of the effects of refpiration on the animal fluids is delayed till the nature of these fluids shall have been more generally confidered.

in the second state of the

SECT.

SECT. IV.

OF THE NATURAL FUNCTIONS.

orusan sostellio CXCVII.

con the matter of the body itleff, or at

THE animal body from a fmall beginning, grows to a confiderable fize; and at the fame time, from the period of the birth, during the whole of after life, the body fuffers, by various means, a daily and confiderable wafte.

K 2

CXCVIII,

CXCVIII.

The increase of bulk, therefore, must be acquired, and the daily waste supplied, by matters taken into the body, the most part of which, from the prefumed purpose of them, we name ALIMENTS.

CXCIX.

A great part of thefe aliments, as taken into the body, are of a different nature from the matter of the body itfelf; or at leaft, are in fuch a flate as not to be fit for being immediately applied to the purpofes of it: they muft, therefore, be changed, and fitted to the purpofes of the œconomy, by powers within the body itfelf.

CC.

CC.

The conversion, or affimilation, of the aliments to the nature of the folids and fluids of the animal body; the farther changes of these fluids for various purposes, by secretion; and the application of some part of them in nutrition, or in increasing the growth of the body; make what are called the NATURAL FUNC-TIONS.

K3 SECT. todiw appuonite bill terra

SECT. IV.

CHAP. I.

OF DIGESTION.

CCI.

THE term digestion is commonly employed to fignify the function of the stomach alone in changing the aliments; but, in this chapter, we are to confider all the changes of these as they occur succeffively in the different stages through which the matters pass.

CCII,

CCII.

Animals are determined to take in aliment, by the appetites of hunger and thirst.

CCIII.

Hunger is an appetite depending upon a fenfation referred to the ftomach, and arifing from a particular ftate of it.

This flate feems to be in fome refpect the degree of emptinels, but more effecially the flate of contraction in the mufcular fibres which emptinels gives occasion to. This flate of contraction may alfo be excited by certain flimulants applied; but more commonly it depends upon, and is correspondent to, the flate of inanition, and therefore of contraction, in the veffels of the fkin emitting the matter of perspiration.

CCIV.

CCIV.

Thirft is an appetite for liquids, which depends upon a fenfation chiefly referred to the internal fauces, and arifing from the drynefs or heat of thefe parts; from acrimony applied to them, or exifting in the fluids poured out there; from the putrefcency or vifcidity of the contents of the ftomach; and from all increafed evacuations.

CCV.

These appetites determine men to take in a great variety of solid and liquid matters, directed by inftinctive likings and difgusts; in some inftances corrected by experience.

CCVI,

CCVI.

Of the matters chosen, it appears that fome of them are fuited to fupply the matters of the fluids or folids of the body, and therefore properly named Aliment; while others of them are fuited only to improve the relifh of aliment, or to obviate fome deviations ready to happen in the bufinefs of digeftion; and these we name CONDIMENTS.

CCVII.

The proper alimentary matters are animal or vegetable only.

CCVIII,

The animal aliments feem to be fo nearly of the fame nature with the matter of the

the body itfelf, that, to be rendered fit for the purposes of the æconomy, they seem to require no other change but that of being rendered fluid.

CCIX.

But the vegetable aliment is very different from the matter of the animal fluids or folids, and muft therefore be changed into the nature of thefe by the powers CXCIX; and as many animals are nourifhed by vegetable aliment alone, and as perhaps all animal matters may be ultimately traced to a vegetable origin, it will appear, that, to account for the production of animal matters, it is effectially, and in the first place, neceffary to show how vegetable matter may be converted into animal.

CCX.

CCX.

If we confider the many different odours, taftes, and colours, which are to be obferved in different vegetables, we fhould be ready to think that vegetable matter is of very great variety: but we know that the matter diffinguished by its fensible qualities makes but a small part of the whole of any vegetable; and that, besides the matter peculiar to each, there is in most, perhaps in all vegetables, a large proportion of common matter, which we prefume to be the matter adapted, and that very univerfally, to the aliment of animals.

CCX1.

It is this common matter of vegetables, therefore, that we are to confider here; and

and we think it may be confidered as of three kinds only; that is, oily, faccharine, and what feems to be a combination of thefe two.

CCXII.

The oily matter of vegetables, which makes part of the aliment of animals, is without any fenfible odour or tafte; and is not only very nearly the fame in the many different vegetables from which we take it, but is alfo in all of thefe fo nearly a-kin to the oil which appears in animals, that it is not neceffary to fuppofe any confiderable change to be made upon the vegetable oil on its being taken into the bodies of animals.

CCXIII.

It is the faccharine matter, and efpecially this

this when blended with oily matter in different proportion, that makes the greatest part of the common matter of vegetables, and is the chief part of the vegetable aliment of animals. It is this, therefore, that we have efpecially to confider here; and, as it lies in vegetables, it is different from the most part of animal matters in the following respects.

It is readily fusceptible of a vinous and acetous fermentation, and spontaneously enters into the one or the other of these; and, without undergoing more or less of these, it perhaps never enters into a putrefactive fermentation.

The fame matter treated by diffillation, without addition, gives out always, in the first part of the distillation, an acid, and only afterwards a volatile alkali in small proportion.

The fame vegetable matter, treated by calcination, leaves afhes, which contain a fixed

fixed alkali, and an earth that is or may be converted into a quick-lime.

CCXIV.

In all these respects, the common matter of animals is confiderably different.

This enters fpontaneoufly into a putrefactive fermentation, and that without paffing through the vinous or acetous: At leaft, thefe are not to be diffinctly perceived.

The fame animal matter, treated by diftillation, gives out always, in the first part of the distillation, a volatile alkali in large proportion, and only afterwards by a great force of fire it gives out an acid.

Animal matters, treated by calcination, leave afhes, in which no alkali is to be found; and the earth is not calcareous, nor convertible into a quick-lime, by any means yet known.

I

CCXV.

CCXV.

These differences are fufficiently marked; but it is proper to observe here, that the vegetable matter we treat of, by undergoing a putrefactive fermentation, is changed so, as to acquire very exactly most of those characters of animal matter we have just now mentioned.

CCXVI.

The aliment being thus confidered, we proceed to confider the changes it undergoes after being taken into the animal body; but, first, of the course it passes through, and of the motions it is subjected to in its progress.

CCXVII.

CCXVII.

The aliment is taken into the mouth; and there the more folid parts of it are commonly fubjected to a triture, or what is called manducation. At the fame time a quantity of faliva, and of the other fluids of the mouth, with fome portion of our drink, is intimately mixed with it, whereby the whole is reduced to a foft pulpy mass. In this state, by the action of deglutition, it passes through the fauces into the cefophagus, by which it is con-

body; but, fr.IIIVXOO COLVIE and ; what

through, and of the motions it

goes after being taken into the animal

Here the aliment is detained for some time, subjected to a constant agitation and fome preffure, both by the contractions of the different parts of the ftomach itfelf.

felf, and by the alternate preffure of the diaphragm and abdominal mufcles. After fome time, however, first the more fluid parts, and at length the most minute parts of the folid matter are pushed thro' the pylorus into the duodenum.

CCXIX.

The matters received from the flomach into the duodenum pafs on from thence fucceflively through the feveral parts of the inteftinal canal; and, in the whole of the courfe, the matters are ftill fubjected to the alternate preffure of the diaphragm and abdominal mufcles, and to the contractions of the inteftines themfelves.

CCXX.

Through the whole course of the inteftines, but especially in those named the I L fmall,

fmall, the more fluid part of the contents, and particularly the peculiar fluid we name chyle, is taken into the veffels named lacteals. Thefe, from imperceptible beginnings on the internal furface of the intestines, unite into larger veffels laid in the mefentery, and convey the chyle, and what accompanies it, first into the conglobate glands of the mefentery, and from thence to the receptaculum chyli, as it is called. From this the chyle paffes by the thoracic duct into the left fubclavian vein. In one or other part of this course of the chyle, the veffels carrying it are joined by lymphatics, returning the lymph from almost every part of the body.

CCXXI.

The matters contained in the inteftinal canal, not taken into the lacteals, are moved onwards in the courfe of the inteftines, becoming

coming by degrees of a thicker confiftence, efpecially in the colon, where their motion is confiderably retarded; but, at length, they are moved onwards to the extremity of the rectum, where their weight, bulk, and acrimony excite motions which throw them entirely out of the body.

CCXXII.

This is the course of the alimentary matters, so far as they can be confidered as any ways in a separate state. Of the motions of the several organs concerned in this course, we pass over those of manducation, deglutition, or others depending on the action of muscles, the functions of which are readily understood from a knowledge of their situation; and we are here to consider only the motions of the alimentary canal itself.

L 2

2

CCXXIII.

CCXXIII.

The motions in the cefophagus depend upon the action of its muscular fibres, which are chiefly those forming a chain and cirlarly furrounding it. This tube, by the morfel of food pushed into it by the action of deglutition, is neceffarily dilated, and its circular fibres are thereby excited to a contraction. But as these fibres are fucceffively dilated, fo are they also contracted, and pufh on their contents through the feveral portions of the tube, alternately and fucceffively dilated and contracted, giving the appearance of a vermicular motion, and what is commonly called periftaltic. This motion may be propagated either upwards or downwards; and the direction of it is in the one or the other way, as the motion happens to begin at the upper or lower extremity.

CCXXIV.

CCXXIV.

The motion of the ftomach is not fo fimple. Its muscular fibres are in like manner irritable by dilatation, and its circular fibres must therefore be in fome measure subjected to a successive dilatation and contraction. But, though the direction of fuch motions is from the left to the right, this does not immediately pufh the contents of the flomach into the intestines. It feems to be the purpose of the œconomy, to detain the aliment for fome time in the ftomach; and therefore, any confiderable dilatation of the circular fibres, efpecially that which occurs in a full ftomach, feems to have the effect of exciting the longitudinal fibres to a contraction, which draws the two orifices of the flomach nearer to one another. By this the pylorus is raifed up and rendered lefs L 3

lefs eafily paffable; and probably, at the fame time, the peculiar band of circular fibres which furround the pylorus, are more firmly contracted, and render it lefs pervious. Vide CXLIV. From hence it is, that the direction of the periftaltic motion of the ftomach is fometimes from the left to the right, and fometimes alfo the contrary way. It is, however, most constantly in the first manner; because it is commonly begun from the œfophagus, and becaufe, when it is inverted, the refistances on the left from the blind fac of the flomach, from the higher fituation of the cardia, and from the constriction of this by the diaphragm in infpiration, are commonly more confiderable than the refistance at the pylorus. The contents of the ftomach, therefore, are at length pufhed through the pylorus; in the first place, the more fluid contents, as these occupy the antrum pylori, while the more folid, having

having their air loofened by fermentation, are rendered fpecifically lighter, and float nearer the upper orifice. But at length, as the ftomach is in any meafure emptied, the pylorus is lefs raifed, is more relaxed, and allows matter to pafs more eafily; and, at the fame time, the empty ftomach contracted more, is efpecially towards the right extremity contracted to fuch a degree, as to embrace the fmalleft folid matters, now fallen down into it, and to pufh them through the pylorus.

This is an idea of the ordinary motions of the ftomach; but they are, upon fome occafions, fubject to other modifications, as in eructation, rumination, and vomiting, which, however, as morbid, we referve to be confidered in the pathology.

CCXXV.

The motions of the inteftines will be readily

readily understood, from what has been faid of those of the cefophagus. Any portion of the inteftinal canal being dilated, will in confequence be contracted, and will urge on its contents in the fame direction in which the motion was begun. But as the force here is gentle; and as, in the long course of the canal, there occur many flexures, different politions, and occafional irritations; it is obvious, that refistances and stronger contractions may frequently occur here, to change the direction of the motion: accordingly we find it frequently changed, and directed from below upwards, in fo far that the contents of the intestines frequently pass into the ftomach. But the motions of the intestines are, however, most constantly directed from above downwards, both becaufe they are commonly begun from the ftomach, and becaufe, when inverfions do occur, there is commonly still fo much refift-

refiftance at the pylorus, and more efpecially at the valve of the colon, as to turn the direction again into its proper courfe. In the colon, from its polition, ftructure, and the confiftence of its contents, the progrefs of thefe is more flow and difficult; and it is therefore here affifted by the longitudinal fibres peculiarly difpofed, fo as by their contraction to contribute more to the dilatation of every fucceeding portion of the inteftine.

CCXXVI.

The chyle is taken into the lacteals, and moved onwards in thefe in the fame manner (CLXVII.) as the lymph is in the feveral lymphatics in other parts of the body, to which the lacteals are in ftructure and fituation exactly fimilar.

CCXXVII,

CCXX II.

The course of the alimentary matters, and the motions by which they are carried on, being now explained, we return to confider the feveral changes which the aliment undergoes in this course.

CCXXVIII.

In the mouth, if the aliment taken in be of a folid confiftence, it is here, as we have faid, fubjected to a triture; and if our food is of a foft and moift kind, we are inftinctively directed to take in along with it fome dry matter, as bread, that the whole may be fubjected more certainly to a complete manducation. By this our aliment is not only more minutely broken down, but is alfo intimately mixed, with the liquids at the fame time taken in, with the

the faliva and other fluids of the mouth, and with a quantity of air intangled by thefe vifcid fluids.

CCXXIX.

In this divided and moiftened flate, the aliment is taken down into the flomach, where it is farther diffolved, the vegetable matter of it begins to be changed to the nature of animal, and the oily parts of the whole begin to be united with the watery. But thefe changes by folution, affimilation, and mixture, require to be feparately confidered.

CCXXX.

The folution here, as in other cafes, may be affifted by the mechanical division of the folid matter, by the agitation of the diffolving mass, and by the application of heat,

heat; and, with these affistances, the folution must be performed by the application of a proper menstruum.

CCXXXI,

The division of the folid is fometimes affifted by a previous cookery, and commonly by the manducation we have mentioned; but the human ftomach does not feem by any mechanical powers to contribute to this. It gives only a moderate agitation, which, in any cafe, contributes little to mechanical division.

CCXXXII.

The degree of heat applied here, being that of the common temperature of the human body, may affift the folution; but it is of no confiderable power, and no affiftance is got from any clofenefs of the veffel

veffel which occurs here. Upon the whole, the affiftances applied here are not confiderable, and the fpeedy folution that takes place must be chiefly owing to the power of the menstruum.

CCXXXIII.

The menftruum that appears here, is a compound of the liquid matters taken in, of the faliva, and of the gaftric liquors: but in all, or any of thefe, we do not readily perceive any confiderable folvent power; nor, by any artifice, in employing thefe out of the body, can we imitate the folutions performed in the ftomach.

CCXXXIV.

However, from what happens in the ftomachs of certain animals, there is ground to prefume, that indeed in every one there is

is a peculiar folvent. But whether this be a menftruum dividing the folid into integrant parts, and thereby reducing it to a fluid state, or if the folvent here be a peculiar fermentative power, refolving matters more or lefs into conftituent parts, is not clearly perceived.

CCXXXV.

The latter is the most probable, as the circumstances of fermentation very conftantly appear; and as the deviations which at any time appear in the course of digestion, appear always to be an excess of fermentation, either acescent or putrefactive.

CCXXXVI.

The bufiness seems to us to proceed in this manner. The fluids of the stomach have

have the power of fuddenly and powerfully loofening the fixed air of the alimentary matters, which is the first step towards putrefaction, and that which most effectually breaks down the texture, and perhaps the mixture of bodies. But we now know, that putrefcent bodies are very powerful in exciting an acefcent fermentation in vegetable fubstances, which the human ftomach is hardly ever without; and that this acefcency therefore, in the next place, very conftantly fucceeds, and an acid is produced in the ftomach. This acidity makes the effects of the putrefaction difappear; and the acidity in its turn difappears alfo, probably by its being abforbed by, or united with, the putrefcent and oily matters here prefent; and it is in this manner that we fuppofe that the animal fluid is produced, and daily renewed by the combination of a fresh portion of acid with putrefcent fluids previoufly exifting

in the body. The daily production of acid in the human ftomach, and its readily difappearing again, without flowing any morbid effects, renders our doctrine fufficiently probable.

CCXXXVII.

This is the affimilation of vegetables that I fuppofe to take place, and is begun in the ftomach, but is not completed there: for we obferve, that the long retention of the alimentary matters in the ftomach, whether from the infolubility of the matter, or from an obftruction of the pylorus, produces a greater degree of acidity; and, in general, the acidity which commonly prevails in the ftomach does not difappear but in the after courfe of the aliment.

CCXXXVIII.

CCXXXVIII.

It is efpecially the bile, added to the matters which have paffed from the ftomach into the duodenum, that is fitted to cover the acidity which appeared in the ftomach. It is probable alfo, that the pancreatic and inteftinal liquors contribute to the fame effect; and it is perhaps for the fame purpofe, that the lymph is conftantly added to the chyle in its courfe. But, after all, we must reft in the general idea, and own that we do not know exactly how this matter proceeds, nor what the feveral fluids, added to the aliment in the different parts of its courfe, truly contribute to the changes of it.

CCXXXIX.

It is probable, however, that, by the M mix-

mixture mentioned, the peculiar fluid which we name the chyle is produced; for, though it is certain that a variety of fluid matters may enter the lacteals and accompany the chyle there, it is still probable, that there is a peculiar fluid produced by the actions of the ftomach and inteffines, and fuch as becomes the principal ingredient in the animal fluids afterwards formed, that is strictly intitled to that apellation. This chyle does not appear in the ftomach; but first in the duodenum, and more copioufly still in the jejunum and first part of the ileum. It appears indeed in the whole of the ileum, cœcum, and colon, but in the last lefs copioufly: all which shows, that a particular mixture is necessary to it; and at the fame time that it is not made at once, but fucceffively in the courfe of the intestines.

CCXL.

CCXL.

It remains to speak of the mixture of the oily with the watery parts of the aliment. This we cannot well explain; but it is of confequence to observe here, that fuch a mixture is actually made. It is evident that a large quantity of oil in a separate state is taken in as a part of our aliment, but at the fame time no oil commonly appears in a feparate state in the mafs of blood; it must therefore be united. with the other parts of the mass in the way of mixture. Hitherto the phyfiologifts have hardly mentioned any other means for this union of oil but the application of viscid fluids; but these can occafion only a diffusion, and some means of mixture must necessarily be fupposed. What these however are, we do not certainly know. They do not produce their effect in the first passages; for in the chyle, M 2 till

till it enters the fubclavian vein, the oil appears to be only in a diffused state, and probably the perfect mixture is only made in the passage through the lungs.

CCXLI.

It may be proper here to take notice of another matter which conftantly enters' into the mixture of animal fluids. This is air, which, by different means, can be extracted in confiderable quantity from every kind of animal matter. What is properly the origin of this, when and where it is infinuated into the animal fluids, and by what means it is either fixed in thefe or loofened from them, are all queftions not yet refolved; but perhaps neceffary to be refolved, before we can fpeak with any confidence of the changes which the animal fluids undergo in different parts of the fyftem. We can obferve,

in

in the mean time, that a quantity of air is always prefent in the chyle in a very loofe ftate; that it becomes more fixed in the mafs of blood after this has paffed through the lungs; and that again, in the different fecreted fluids, the air appears to be in fome of them ftill fixed, and in others much more loofe; and it is probable, that all this has a particular relation to the production and properties of the different fluids of animals.

CCXLII.

We have now followed the courfe of the aliments, fo far as we can confider them as any ways in a feparate ftate; but we do not perceive, that, in any part of this courfe, the proper animal fluids are entirely formed: And it is very juftly fuppofed, that the proper mixture or affimilation is not finished till the chyle, mixed M 3 with

with the mass of blood, has undergone the action of the lungs, through the vessels of which it must almost immediately pass, after entering the subclavian vein, and seemingly before it is applied to any of the purposes of the animal œconomy.

CCXLIII.

What change the fluids undergo in paffing through the lungs, or by what means the fuppofed changes are produced, after all that has been faid, feems ftill to be very little known.

The mechanical powers of preffure, commonly fpoken of, do not in fact take place, nor are their fuppofed effects any ways confiftent with found philofophy; and, on the other hand, it is very probable, that the changes produced are the effects either of chemical feparation or mixture. What

What has been fuppofed to be performed in this way by an abforption of air, or of a particular matter from it, is very uncertain in fact, and has led to a still more uncertain reasoning.

It is now certain, that a quantity of · mephitic air, and perhaps fome other matters are conftantly exhaling from the lungs of living animals, and are carried off by the atmospherical air alternately entering and iffuing from the lungs. This is a pretty certain evidence that fome change of mixture is going on in the fluids paffing through the lungs; but from what particular portion of the fluids the mephitic air proceeds, or what is the effect of its feparation, we know not: And indeed, as we have faid before, what are the effects of the action of the lungs upon the state of the fluids, we are very uncertain. Upon the whole, we still know but little of

M 4

the

the production, or formation, of the animal fluids; and therefore from the confideration of their formation, we have learned little of their nature; but we muft now try to difcover what we can of it, by examining thefe fluids as they are found already formed in the blood-veffels.

fe air proceeds, or what is the effect of it.

teparation, we know not: And indeed as

on the win ic, we that have but the

sit sit mogu ignel bei to s SECT.

SECT. IV.

CHAP. II.

OF ANIMAL BLOOD.

CCXLIV.

THE red fluid paffing from the lungs to the left ventricle of the heart, and thence by the aorta and its branches to every part of the body, may be confidered as a mass containing, either formally or materially, every part of the animal fluids; and

and may therefore be called the common mafs of blood. This term, however, muft be ftrictly confined to the circulating fluids while they retain their red colour; for when they lofe this, it is always in confequence of fome feparation of parts. The fame red fluid, indeed, as it is found in the veins, has alfo fuffered fome feparation of parts; but as the blood in the veins is never entirely deprived of the whole of any matter that was prefent in the arteries, fo we think the venous blood may ftill be confidered as a part of the common mafs.

CCXLV.

This mass of blood we find to be an heterogeneous aggregate; and it will be proper to inquire into the feveral parts of this before we employ any chemical trials for

for difcovering the mixture of the whole, or of its parts.

CCXLVI.

We difcover the parts of this aggregate chiefly by the fpontaneous feparation of them, which takes place upon their being drawn out of the veffels of a living animal.

CCXLVII.

The feparation commonly proceeds in this manner. Immediately after the blood is drawn out, it exhales a fenfible vapour, and, after fome time, it is found by that exhalation to have loft a part of its weight, more or lefs, according to the degree of heat it is exposed to, according to the extent of furface by which it is exposed to the air, and probably alfo according to dif-

different conditions of the blood itfelf. The matter thus exhaling may be called the *balitus* or vapour of the blood.

CCXLVIII.

Soon after the blood has been drawn out of the veffels, it lofes its fluidity, and the whole of it concretes into one foft gelatinous mafs: but after fome time, there oozes out from this mafs a thin fluid; and as the feparation of this proceeds, the mafs contracts into a fmaller bulk, and, in proportion, becomes more denfe.

CXLIX.

This is the feparation which almost always takes place, and has at all times been obferved by physicians. The fluid part is called *ferum*; and the thicker confiscent

fistent part has been called cruor, but more properly the craffamentum.

CCL.

Both parts feeem homogeneous and fimple, but are not. For, if the craffamentum taken from the ferum be laid upon a linen cloth, and water is poured upon it, the water wafhes off a red coloured part, and carries it through the pores of the cloth; and there remains a whitifh, confiftent, but foft and tough mafs, not to be further diminifhed or feparated into parts by any ablution.

A like experiment flows always a like matter prefent in the mafs of blood; and upon feveral occafions, both while the blood remains within the veffels, whether of the living or dead body, and when it is drawn out of the veffels of the living, this matter fpontaneoufly feparates from the other

other parts of the blood. It is therefore a part conftantly prefent in the blood. It is what Gaubius, after Malpighi, calls the *fibra fanguinis*. Mr Senac names it the *coagulable lymph*; and we fhall fpeak of it under the title of the gluten of the blood. When it appears upon the furface of the blood drawn out of the veffels of living animals, it is called the *inflammatory cruft*.

CCLI.

When the blood is viewed with a microfcope, whether as moving in the veffels of a living animal, or when out of the veffels remaining ftill fluid, there are certain parts of it which appear of a round figure, and alfo of a red colour, while the reft is almost colourles. The parts thus diftinguishable by their figure are called the red globules; and it appears that the red colour of the whole mass depends up-

on

on the prefence of thefe only. It is chiefly thefe parts which are wafhed off from the craffamentum in the experiment abovementioned; and we now conclude, that, befides the globules, the gluten, and a portion of ferum that happens to be entangled in the pores of the concreting mafs, there is no other matter evident in the craffamentum.

CLII.

The ferum is a transparent fluid of very little colour, and feemingly fimple; but if it be exposed to a heat of 156 degrees of Fahrenheit's thermometer, it concretes into a firm and almost transparent gelly; and, if this be cut into minute pieces, there exudes from it a thin colourles fluid of a faline tafte. In proportion as this fluid is more carefully separated, the coagulated part becomes infipid, and in all its pro-

properties refembles the gluten feparated from the craffamentum. From hence we are ready to conclude, that the ferum, as obtained by fpontaneous feparation, confifts of a portion of gluten diffolved in a faline fluid, which we name the SERO-SITY.

CCLIII.

From the whole that has been faid from CCXLIV. to CCLII. it appears that there are three diffinct portions and kinds of matter in the common mafs of blood; that is, red globules, gluten, and ferofity. What other matters may alfo be there, we fhall confider afterwards; but, in the mean time, fhall fay a little more of each of the parts we have already mentioned.

CCLIV.

CCLIV.

The red globules have been confidered. as an oily matter, and from thence their diffinct and globular appearance has been accounted for; but there is no direct proof of their oily nature, and their ready union with and diffufibility in water renders it very improbable. As being microfcopical objects only, they have been reprefented by different perfons very differently. Some have thought them fpherical bodies, but divisible into fix parts, each of which in their feparate state were alfo fpherical; but other perfons have not obferved them. to be thus divifible. To many observers, they have appeared as perfectly fpherical, while others judge them to be oblate fpheroids or lenticular. To fome they have appeared as annular; and, to others, as containing a hollow veficle. All this, with feveral other circumstances relating to. them, N

them, very varioufly reprefented, flow fome uncertainty in microfcopical obfervations; and it leaves me, who am not converfant in fuch obfervations, altogether uncertain with refpect to the precife nature of this part of the blood. The chemical history of it is equally precarious; and therefore, what has been hitherto faid of the production, and changes happening to thefe red globules, we choofe to leave untouched. We shall afterwards fay fomething with respect to their general use in the animal fystem; and now we shall attempt to explain the caufe of fome changes, which in certain circumstances appear in the colour of the whole mafs of blood.

CCLV.

We fuppofe that the red globules, when viewed fingly, have very little colour; and that it is only when a certain number

of

of them are laid upon one another, that the colour appears of a bright red: but this alfo hath its limits; fo that, when the number of globules laid on one another is confiderable, the colour becomes of a darker red. Upon this fuppofition, the colour of the mafs of blood will be brighter or darker as the colouring part is more or lefs diffufed among the other parts of the mafs; and we think this appears to be truly the cafe, from every circumftance that attends the changes which have been at any time obferved in the colour of the blood.

CCLVI. what for a count of the former

The gluten of the blood, from its refemblance on the one hand to the *albumen ovi*, and on the other to the matter of the folids of animal bodies, we confider as the principal part of animal fluids, as N 2 that

that which is immediately formed of the aliment taken in, and as that which is employed in increasing the growth of the folids, or in repairing their wafte.

CCLVII.

But it is well known, that the animal fluids in general, and particularly the gluten, is prone to putrefaction; and that, even in the living body, if fresh aliment be not conftantly taken in, and alfo if certain excretions which carry off putrefcent matter be not conftantly fupported, a confiderable putrefaction certainly takes place. From hence we are led to think, that fome approach to putrefaction conftantly takes place, even in the most healthy bodies; and that it appears efpecially in an evolution of faline matter; and that this, taken up by the water conftantly prefent, forms the ferofity. We fuppofe it is this which

which affords the vapour of the blood, (CCXLVII.) and that it is the ferofity diffolving a portion of the gluten which forms the ferum that appears upon fpontaneous feparation (CCXLVII.)

CCLVIII.

The faline matters impregnating the ferofity, if we may judge from the analyfis of urine, are of various kinds; but particularly, there is prefent an ammoniacal falt, now well known under the name of the *effential falt of urine*, which, if not originally formed, is at leaft most copiously evolved in animal fluids.

CCLIX.

These are our conjectures concerning the parts of animal blood; and it remains to fay in what proportion each of them is N 3 pre-

prefent in it. This will perhaps be always difficult; and in the mean time we can perceive, that many estimates formerly made could not be exact, as the feveral parts were not properly known; and, while judging chiefly from the appearances upon spontaneous separation, physicians were not aware how much these are affected by the circumstances of extravalation, and by those in which the blood is placed after being drawn out. There are not yet indeed experiments made to afcertain, with any exactness, the proportion of the feveral parts mentioned: but it is probable, that the red globules make a fmall part of the whole; that the gluten, if we confider both what is in the craffamentum and in the ferum, is in much larger proportion; but that the watery portion is the largest of all, and at the same time that this has always a confiderable quantity of faline matter diffolved in it.

CCLX.

CCLX.

We would next put the queftion, By what means the parts of this heterogeneous mafs are kept fo equably diffufed among one another, and the fluidity of the whole fo conftantly preferved? This we suppose to be done chiefly by motion and heat, and by the parts difpofed to concrete being kept from the contact of any matters to which they might adhere more firmly than they do to the other parts of the blood. The diffused parts we suppose to be prefent only in those veffels in which a confiderable degree of agitation is conftantly kept up; and we fuppofe alfo, that the heat always here prefent, both diminishes the cohesion of the gluten, and increases the folvent power of the ferofity. Experiments made with neutral falts feem to confirm the latter; and it is also probable, that the N4

the fame folvent power may be increafed by a quantity of air that is conftantly intermixed with the mafs of blood while it remains in the veffels, and is under a conftant agitation. It is fuppofed, that an attention to thefe feveral circumftances will explain most of the cafes of fpontaneous feparation that occur either in the living or dead body, within the veffels or without them; but the detail would be too long for this place.

CCLXI.

We fhall add here a few words on the ufe of this fingular composition of animal blood which we have been confidering.

It appears evidently, from many circumftances of the animal œconomy, that its functions require a fystem of vessels conftantly filled, and even distended; but as, at the fame time, these vessels must be open

open by a multitude of their extremities. if all the fluids were fuch as could pass by thefe extremities, the fystem could not be kept filled for a few minutes. It is necesfary, therefore, that the fluids should be partly of fuch a fize as that they cannot pafs through the fmaller veffels, and partly in a diffused state only, which has commonly the fame effect. Hence it is, that the red globules, under the ordinary impetus of the heart and arteries, are firicily confined to certain veffels; and it is probable, that, in the like circumstances, the diffused gluten does not go much farther. This ferves to keep the larger veffels of the fystem constantly filled. But, on the other hand, the ferofity being fufficiently fluid, might be fuppofed to run off by the many outlets open to it, and thereby to leave the fluids in the larger veffels of a confistence unfit to circulate. This, however, feems alfo to be obviated by the vifcidity of the groffer

groffer parts of the blood, fufficient always to entangle fo much of the more fluid, as may be neceffary to preferve the due fluidity of the whole.

CCLXII.

The heat of the human body, fupported by powers within itfelf, is probably the effect of the motion of the blood, and might have been treated of when we were confidering that fubject. But as many perfons fuppofe it to depend in fome meafure on the nature of the fluids, we have referved it for this place; and here perhaps to fay only, that the queftion concerning the caufe of animal heat is not yet folved.

CCLXIII.

The opinion of animal-heat's being the effect

effect of mixture, is to be little regarded; as the matters fuppofed to be mixed, the place in which the mixture is made, and the other circumftances relating to it, are equally hypothetical, and the whole is ill fupported by any analogy.

CCLXIV.

More fpecioufly is animal-heat fuppofed to be the effect of putrefaction, towards which there is certainly fome approach in animal bodies; but the opinion is ftill very doubtful. For, *firft*, the effect of any degree of putrefaction in producing heat is not well afcertained. *Secondly*, It is not fupported by any analogy, that putrefaction, in the degree to which only it proceeds in living bodies, is capable of producing the heat appearing there. And, *laftly*, Whatever is the degree to which putrefaction proceeds in living bodies, it does

does not appear that there is any increase of heat correspondent to the increase of putrefaction, and rather the contrary.

CCLXV.

The fuppositions either of mixture, or of putrefaction, as the caufe of animal-heat, are both of them rejected by this, that the generation of heat in animal bodies is manifeftly dependent on another caufe, that is, the motion of the blood. For the power of generating heat in any animal is not perfect till the motion of the blood in it is fully established; and, when the generating power is established, we perceive the heat to be increased or diminished as various caufes increafe or diminish the motion of the blood. In dying animals, the heat grows lefs as the motion of the blood grows lefs; and when at death this ceafes altogether, the heat ceafes alfo, commonly,

at

at leaft, as foon after death as we can fuppofe a body of the fame bulk to lofe the heat it had acquired.

CCLXVI.

This connection between the heat and motion of the blood feems in general to be well proved; and, though it may be difficult to reconcile certain appearances to it, we would fo far admit of the fuppofition, as to inquire, in the next place, into the manner in which the motion of the blood may generate heat.

CCLXVII.

On this fubject, the most common opinion is, that the heat is produced by the attrition of the particles of the blood upon one another, or of these on the internal furface of the vessels in which they move. But

But we cannot find any analogy to fupport either the one or the other fuppofition.

The attempt made to fupport the latter fuppolition, by endeavouring to flow, that upon this the equality of heat in the different parts of the fame body is well explained, deferves little regard, as it is founded on doubtful principles and miftaken facts.

CCLXVIII.

The equality of heat in the different parts of the fame body feems to require the generating power to be very generally diffufed over the whole; but it does not feem to require its being precifely equal in every part, as the interpolition of pretty large veffels in every part of the body, and the fpeedy communication of the fluids from any one part to every other, will fufficiently account for the equality of heat, though

though the generating power fhould be in fome measure confined to certain parts only.

However, we take no notice of the fuppolitions which have been made of the generating powers being confined to certain fmall portions of the fystem only. These fuppositions give no relief in the general theory, and they are not fupported by any particular evidence. The breathing animals are the warmeft; but that they are warmer because they breathe, is not more probable than that they breathe becaufe they are warmer.

CCLXIX.

With respect to this theory, which deduces animal-heat from the motion of the blood, we must own, that it is attended with feveral difficulties. It will be difficult to show, in so many animals of different age,

age, fize, and temperament, in which the degree of heat is nearly the fame, that the motion of the blood, in all its circumstances, is also exactly the fame; or to show, in the different animals in which the degree of heat is confiderably different, that the motion of the blood in each is correfpondent to the difference of heat. May it not be fuppofed, that there is fome circumstance in the vital principle of animals which is in common to those of the fame class, and of like œconomy, and which determines the effect of motion upon the vital principle to be the fame, though the motion acting upon it may be in different circumftances?

CCLXX.

In all we have hitherto faid of animal fluids, we have confidered the common mafs of blood as confifting of three parts

or

or three kinds of matter only; but many more have been fuppofed to be prefent in it, and we fhall inquire upon what ground.

It is common to fuppofe, that the aliment or the chyle formed of it is not perfectly affimilated in paffing once only thro' the lungs; but that, for fome time after fuch passage, it continues to circulate with the blood under the fame form and of the fame qualities which it had when it first entered the fubclavian, and particularly in this state to furnish the milk which is fecreted in the breafts of females. There is, however, no proper evidence of the chyle's ever appearing in the blood-veffels, and the appearances of it alleged can be otherwife accounted for. The arguments for the fame opinion, which are drawn from the confideration of the fecretion of milk, are embarraffed with many difficulties.

0

CCLXXI.

CCLXXI.

It is probable, that the animal fluid (CCLV. CCLVI.) is in a conftant progrefs, and hardly for a moment flationary, or therefore uniformly the fame over the whole of the common mafs. Some part of it is that which was laft formed, and therefore the neareft to the vegetable matter from which chiefly it was produced; while another part of it is that which has remained longeft in the body, and is therefore the neareft to putrefaction. Between thefe two there may be feveral intermediate flates, which, however, like the neareft flades of the fame colour, are not diftinguifhable by our fenfes or experiments.

CCLXXII.

Befides the difference of matter arifing from

from the progrefs of the animal fluid, there have been other matters fuppofed prefent in the common mafs, and as commonly conflituent parts of it. Such are a mucous matter, like to the mucous matter of vegetables; and a gelatinous matter, like to that which is extracted by decoction from the folid parts of animals. But there is no evidence of either being formally prefent in the mafs of blood, and the fuppofition is founded on miftaken facts and falfe reafonings.

CCLXXIII.

But it is proper to be obferved here, that many extraneous matters may, by different ways, be introduced into the bloodveffels; and that many of the fecreted fluids, fometimes very different from any thing that exifted before in the mafs of blood, may, by abforption or regurgita- O_2 tion,

tion, be again taken into the blood-veffels: But, with regard to all of thefe, whether extraneous matters, or thofe produced in the body itfelf, it is probable that hardly any of them enter into the mixture of the animal fluid, and that they are only diffufed in the ferofity till they can be again thrown out of the blood-veffels by the readieft outlets. The oil of the adipofe membrane is frequently, and perhaps neceffarily reabforbed, and feems to be, befides the lymph, the only reabforbed matter which enters again into the mixture of the animal fluid.

SECT.

SECT. IV.

CHAP. III.

OF SECRETION.

CCLXXV.

A FTER thus confidering the parts of the mafs contained in the red veffels, we muft next confider the feveral fluids which appear in the other parts of the of the body.

03

CCLXXVI.

CCLXXVI.

All of thefe we fuppofe to be derived from the common mafs, as they appear in veffels continuous with those of the common mafs, and as their appearance ceases when the communication of the veffels containing them, with the fanguiferous veffels, is any how interrupted.

CCLXXVII.

The fluids thus derived from the common mass feem to be produced in confequence of a certain structure, with perhaps fome other condition in the extreme vessels through which the fluids pass; and a part having such a structure, is called a gland or fecretory organ, the function of which, from the most obviovs notion of the manner of it, is called fecretion.

CCLXXVIII.

CCLXXVIII.

The ftructure of the organ, and the manner of its function, feem to me for the most part unknown; at least, what we know or fuppose with regard to the structure hardly in any case applies to the explanation of the function.

CCLXXIX.

If it any how appeared that the feveral fecreted fluids were all of them previoufly exiftent in the fame forms in the mafs of blood, it would not perhaps be difficult to explain what might be ftrictly called a fecretion. But fuch previous exiftence does not appear; for, except the matter of exhalation into the feveral cavities of the body, and the matter of urine and of perfpiration, we find no O_4 proper

proper evidence of any other fecreted fluids prefent in the mafs of blood. We cannot find there, either milk, mucus, or oil, and much lefs the appearance of many other fluids which are only found after they have paffed through certain organs.

CCLXXX.

This being the cafe, the confiderations of the phyfiologifts with regard to the velocity of the blood, and other circumftances favouring the feparation of the parts of a fluid which are only diffufed among one another, deferve no attention. The effects of different apertures may go fome length; but we can perceive their particular application only in the few cafes of a fimple feparation. In most others, there appears to be a change of mixture; but we perceive neither the precife changes that are made, nor the the caufe of them.

CCLXXXI.

CCLXXXI.

Till we can difcover thefe more clearly, we may in the mean time obferve, that the action of the veffels of the fecretory organ has a confiderable fhare in determining both the quantity and quality of the fecreted fluid, and that both very often are very little affected by the general ftate of the circulation, or by the different conditions of the mafs of blood.

CLXXXII.

It would feem that no other fecretion but those of perspiration and sweat are manifestly increased by the increased action of the heart and arteries (CLXXXI.), and that most of the other secretions are increased only by stimulants applied to their organs. These stimulants may be either

either fuch as are immediately applied externally or internally to the excretory, or perhaps to the fecretory veffels; or they may be fuch as are applied to the fenforium, or to diftant parts of the nervous fyftem, which by the laws of the animal œconomy have a connection with the organs of fecretion. Thefe ftimulants, at the fame time that they act in either of thefe ways on the fecretory organs, for the most part have no fensible effect on the general state of the circulation of the blood.

CCLXXXIII.

With refpect to the influence of the condition of the common mafs of blood upon the feveral fecretions, we prefume that the flate of the quantity of the fluids in general will affect the quantity of every fecretion; but the effects of the quantity of

of the whole mass are very remarkable only, with respect to the fecretions of perspiration, urine, and milk.

The qualities of the common mafs may alfo be prefumed to affect the feveral fecretions: but the effect of thefe qualities appears most remarkable in the fame fecretions of perspiration, urine, and milk; and, even in these, the effect seems to depend upon the proportion of water more than upon that of any other matter in the common mass. With respect to the other fecretions, we cannot perceive that any of them are increased by a particular matter present in the mass of blood, except it be fuch a matter as stimulates the fecretory organ.

CCLXXXIV.

The feveral fecretions are frequently obferved to affect each other mutually, fo that

that the increase of one diminishes another, and vice versa. This seems to depend either upon a change of determination in the course of the blood (CLXXVIII.), or upon a change in the state of suidity of the common mass, or perhaps upon a connection established between the different organs of secretion as parts of the nervous system; and, except it be in the case of perspiration and urine, we cannot perceive that the effect of the state of one secretion upon that of another depends upon an increase or diminution of any particular matter in the mass of blood.

CCLXXXV.

After mentioning thefe generalities with refpect to fecretion, we fhould, perhaps, proceed in the next place to confider the application of them to the particular fecretions, and alfo to confider more particularly

cularly the feveral fecreted fluids: but we omit both thefe fubjects, as we prefume the former will be obvious from what is already faid; and with refpect to the latter, we have not yet a fufficient number of experiments to proceed any length in it.

SECT. IV.

CHAP. IV.

OF NUTRITION.

CCLXXXVI.

UNDER this title we might confider how the matter both of the fluids and folids of the body is fupplied: but, after what we have formerly faid of the taking in

in and affimilation of the aliment, we have nothing now to add with refpect to the fluids; and we therefore confine ourfelves here to confider in what manner the folid parts obtain their increase of matter and growth, or have their occasional waste repaired.

CCLXXXVII.

There is no doubt that the folids are formed of the fluid prepared from the aliment in the manner we have faid; but it is required now to fay what portion of the fluids is employed in nourifhing the folids, by what channels the nourifhment is conveyed to them, and, being applied there, how from fluid it becomes folid.

CCLXXXVIII.

With regard to the first question, we have no doubt in afferting, that in ovi-1 parous

parous animals, it is the albumen ovi that is employed in nourifhing the chick; and we prefume that it is an analogous fluid which is employed in nourifhing the bird during the whole time of its growth. We think the analogy may be fafely applied with refpect to all animals, the folid matter of which is of the fame kind with that of the oviparous.

CCLXXXIX.

This analogous fluid we take to be the gluten of the blood, properly diluted and freed from any adhering faline matter.

CCXC.

To determine in what manner this nutritious fluid is applied to the nourifhment of the folids, it is neceffary to confider what are the fimple fundamental folids, of which all the others are formed.

CCXCI.

CCXCI.

It feems to be the opinion of the greater part of modern anatomists, that the folid parts confift entirely of a cellular texture, of various denfity in the different parts; and indeed, the structure of the greatest part of the folids is evidently of this kind. But at the fame time, it is alfo true, that a fibrous structure is to be observed almost every where in the body. It appears in the medullary fubstance of the brain and nerves, in the mufcles and tendons, in the arteries, in the excretories of the glands, in the lymphatic veffels, in the alimentary canal, in the uterus and bladder of urine, in the ligaments, in most membranes; and it is to be feen in those membranes which are afterwards changed into bones, efpecially whilft this change is going on.

CCXCII.

CCXCII.

From this view of the univerfality of a fibrous ftructure in animal bodies we are difpofed to believe, that thefe fibres are the fundamental part of animal folids; that they are the primordial ftaminal part of animal bodies; and that the cellular texture is, for the most part, an accretion formed upon thefe fibres.

The confideration of the ftructure and growth of vegetables feems to illustrate and confirm this opinion.

CCXCIII.

At the fame time, from the fibrous parts (CCXCI.) being evidently, in most instances, parts of the nervous fystem, and from the gradual formation of the foetus in which the nervous fystem is first formed, we think it probable, that the P whole

whole of the fibres in the different parts of the body are a continuation of the nerves; and this again will lead to the conclusion, that the nourifhment of the foft and homogeneous folid every where is conveyed to it by the nerves.

CCXCIV.

This fuppofes alfo, what is otherwife probable, that the cortical part of the brain, or common origin of the nerves, is a fecretory organ, in which the gluten of the blood being freed from all faline matter before adhering to it, becomes fit for the nourifhment of the folids, and being poured in a fufficiently diluted ftate upon the organ of the nerves, it is filtrated along the fibres of thefe, and is thus conveyed to every ftaminal fibre of the fyftem. We fuppofe, at the fame time, that the medullary, or what may be called the folid matter

matter of the nerves, is in the living body conftantly accompanied with a fubtle elaftic fluid, which fits them for being the organs of fenfe and motion, and which probably is alfo the means by which the nutritious fluid is carried on in the fubftance of the nerves, from their origin to their extremities.

In what manner the nutritious fluid, thus carried to the feveral parts, is there applied, fo as to increafe the length of the nervous fibre itfelf, or to form a cellular texture upon its furface, and in what manner from fluid it becomes folid, we cannot explain; nor can thefe particulars be explained upon any other fuppofition that has been formed with refpect to nutrition.

CCXCV.

It is probable, that, for a certain time, P 2 at

at its first beginning, the growth of animal bodies proceeds in the fame manner as that of vegetables: but it is evident, that, at a certain period, in the growth of animals, a different œconomy takes place; and that, afterwards, the growth feems to depend upon an extension of the arteries in length and widenefs by the blood propelled into them by the powers CLVI. CLIX. It may be fuppofed, that this extenfion of the arteries is applied to every fibre of the body, and that by the extenfion of thefe it gives an opportunity to the application and accretion of nutritious matter; to the growth therefore of the fibre itfelf, and to the growth of cellular texture on its furface. Perhaps the fame extension of the arterial fystem gives occafion to the fecretion of fluids, which poured into the cellular texture already formed, according to the difpofition of thefe fluids to concrete more or lefs firmly, gives

gives the different degrees of denfity and hardnefs which appears in different parts of the body.

CCXCVI.

By this extension of the arterial fystem, the feveral parts of the body are gradually evolved, fome of them fooner, others later, as by the conflitution of the original stamina, or after occurrences, they are feverally put into the conditions CLXXVII. CLXXVIII. by which they are more or lefs exposed to the impetus of the blood, and fitted to receive a greater quantity of it. But as the parts by these causes first evolved will increase the most in the denfity of their folid parts, they will therefore more and more refift their further growth; and by the fame refistance, will determine the blood with more force, and in greater quantity, into the parts not then P 3

then fo far evolved. Hence the whole fyftem will be at length evolved, and every part of the folids will, in refpect of denfity and refiftance, be in balance with every other, and with the forces to which they are feverally exposed.

CCXCVII.

The extension of the arteries (CCXCV.) depends upon the refistances which occur to the free transmission of the blood through them, as in CLXX.; and further, from a refistance in the veins. For, as a confiderable portion of the blood (by CCLXI.) does not commonly pass into the smaller branches of the arteries, but must pass very entirely into the veins; fo these, by their capacity constantly diminissing as they approach nearer to the heart, and by their coats being of a density and firmness fufficient to prevent further dila-

dilatation, confiderably refift the free paffage of the blood from the arteries into them.

CCXCVIII.

While thefe refistances continue, the arteries, and with them almost every fibre of the body, must be extended at every fystole of the heart; and with this extenfion, the growth of every part will proceed: but, as every part, by its receiving an addition of folid matter, becomes more dense and rigid; so it is less eafily extended, and perhaps lefs readily receives an accretion of new matter, than before. Hence it is, that the more the body grows, it admits of any additional growth more flowly; and unlefs the extending powers increase in the fame proportion with the increasing denfity of the folids, there must be a period at which these two powers will balance each other,

other, and the growth will proceed no farther. But, as it is evident, that the bulk and weight of the heart, and probably therefore its force, does not increafe with the increafing bulk of the body, and that the action of the heart is the principal extending power in the fystem; it is alfo plain, that the extending power does not increafe in the fame proportion with the increafing density of the folids; and therefore, that thefe two powers will, at a certain period, come to balance each other.

CCXCIX.

But not only is the force of the heart thus conftantly diminishing, with respect to the resistance of the arteries; but the force of the heart, though it were still subfisting, has, from other causes, less effect in extending the arteries. The blood is more confined to the arteries, and extends them

them further in proportion to the refiftance in the veins, as in CCXCVII.; and this refistance in the veins, and the extenfion of the arteries depending upon it, will be more or lefs, according to the refpective denfity of these two sets of vessels. But it appears from the experiments of Sir Clifton Wintringham, that the denfity and firmnefs of the veins with refpect to their correspondent arteries, is much greater in young animals than in old; and thence it appears, that during the growth of animals, the arteries are acquiring an increase of denfity in a greater proportion than the veins are at the fame time; and therefore, that the refistance in the veins with refpect to the arteries, must be constantly diminifhing; that the veins will therefore receive a greater proportion of blood; that in the fame proportion the arteries will be lefs extended; and, laftly, that the diminifhed refistance in the veins, concurring with 2

with the diminished force of the heart, will the fooner bring the increasing rigidity of the arteries, and therefore of every fibre of the body, to be in balance with the extending powers; at least, fo far as to prevent their producing any further growth.

CCC.

This account of the change of the refiftances in the arteries and veins, with refpect to one another, is agreeable to phenomena, which flow that the arteries are larger, and contain more blood in proportion to the veins in young animals, than in old; that arterial hemorrhagies, occur most frequently in young perfons; and that congestions in the veins, with hemorrhagies, or hydropic effusions depending upon them, occur most frequently in old age.

CCCI.

CCCI.

It is probable, that the refistance both. of arteries and veins goes on increasing, while the force of the heart is not increafing at the fame time: but it appears alfo, that, from the diminished force of the heart, and the compression which the fmaller veffels are conftantly exposed to from the diftention of the larger, the action of the muscles and other causes, the number of fmall veffels, and therefore the capacity of the whole fystem, is constantly diminishing fo much, that the heart may still for some time be sufficient for the circulation of the blood. But, while the refiftances in the veffels are conftantly increasing, the irritability of the moving fibres, and the energy of the brain, are at the fame time constantly diminishing; and therefore the power of the heart must

I

at

at length become unequal to its tafk, the circulation must cease, and death ensue.

CCCII.

The unavoidable death of old perfons is thus in part accounted for; but it is, however, ftill probable, that the fame event proceeds chiefly from the decay and total extinction of the excitement or vital power (CXXXVI.) of the nervous fyftem, and that from caufes very independent of the circulation of the blood, and arifing in the nervous fyftem itfelf in confequence of the progrefs of life. This feems to be proved by the decay of fenfe, memory, intellect, and irritability, which conftantly takes place, as life advances beyond a certain period.

FINIS.



