Medical reports, on the effects of water, cold and warm, as a remedy in fever, and other diseases, whether applied to the surface of the body, or used internally ... with observations on the nature of fever; and on the effects of opium, alcohol, and inanition / [James Currie].

Contributors

Currie, James, 1756-1805.

Publication/Creation

London: T. Cadell, Jnr., and W. Davies; [etc., etc.], 1798.

Persistent URL

https://wellcomecollection.org/works/yyz5hqts


License and attribution

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection 183 Euston Road London NW1 2BE UK T +44 (0)20 7611 8722 E library@wellcomecollection.org https://wellcomecollection.org



19453/B

O T24RADFORD LIBRARY,

Saint Mary's Yospital, Manchester.

No. 45 F. Sec 30

This Book to be returned in _____ days.

Fine for overtime ____ per day.

Note.—No book can be renewed if wanted by another reader, nor unless brought to the Library for that purpose.


It is requested that the leaves of books may not be turned down,—that no person will write in them,—and that the greatest possible care may be taken of them.

EXTRACTS FROM THE RULES.

That each Medical Officer shall be allowed not more than two works out of the Library at one time, and not more than two volumes of each work.

That Registered Medical Students shall be allowed to take out books every Tuesday and Saturday, from eleven till one, or at such hours as may be ordered from time to time by the Board.

That each Registered Medical Student shall be allowed to have not more than one book out of the Library at the same time, unless the work consist of two or more volumes, and in no case more than two volumes.



19453/B

MEDICAL REPORT

ON THE

EFFECTS OF WATE

COLD AND WARM,

AS A REMEDY IN

FEVER AND OTHER DISEASES,

Whether applied to the Surface of the Body, or used Internally;

An Inquiry into the Circumstances that render Cold Drink, or the Cold Bath, dangerous in Health.

TO WHICH ARE ADDED

OBSERVATIONS ON THE NATURE OF FEVER;

AND ON THE EFFECTS OF

OPIUM, ALCOHOL, AND INANITION.

THE SECOND EDITION, CORRECTED AND ENLARGED.

BY JAMES CURRIE, M. D. F. R. S.

PHYSICIAN IN LIVERPOOL, AND FELLOW OF THE ROYAL COLLEGE OF PHYSICIANS, EDINBURGH.

Intentiones operationum, quas proposuimus (ut arbitramur) verissimæ sunt, remedia intentionibus sida. ** Rem ipsam experimentum et comprobavit et promovebit. ** Opera consilii cujusque prudentioris, sunt effectu admiranda, ordine quoque egregia, modis faciendi tanquam vulgaria.

BACON. Historia vitæ et mortis,

LIVERPOOL.

PRINTED BY J. M'CREERY,

FOR MESSRS. CADELL JUN. AND DAVIES, LONDON;
AND MR. CREECH, EDINBURGH.

1798.

MEDICAL REPORTS.

EFFECTS OF WATER

Contract and

FEVER AND OTHER DISC

HISTORIAL MEDIL

CREARLY NO WALLES CALLED OF LEVER OF LEVER CONTROL OF LEVER LIVE CONTROL OF LI

BY JAMES OURRIE, ALD F.R.S.

DATOS SET AN ANTICIDADE AND ANDRESS OF THE ROPAL OF

The contract of the contract o

POR MERSEN. CADRIL JUN. AND PAVIES, CORDORS.

1708.

RIGHT HON. SIR JOSEPH BAN

SIMARYS TIOS DEORD LIBRAN Baronet and Knight of the Bath, President of Royal Society, &c. &c. &c.

recorded such facts as related to the sol

IN presenting this volume to you, I beg leave to say a few words on the views with which it was written, and the circumstances that gave rise to the publication.

About eighteen years ago, when I was at Edinburgh, it fell to my lot to write a paper on the influence of cold on the living body, for one of the societies of students, of which I was a member. In defending my speculations against some ingenious opponents, a perpetual contradiction occurred as to facts, which a reference to original authorities, did not enable me to remove; for I discovered, that the accounts given of the temperature of the human body under disease, even by the most approved authors, are, with a few exceptions, founded, not on any exact measurement of heat, but on the sensations of the patient himself, or his attendants.

Impressed

Impressed with the belief, that till more accurate information should be obtained respecting the actual temperature in different circumstances of health, and disease, no permanent theory of vital motion could be established, nor any certain progress made in the treatment of those diseases in which the temperature is diminished or increased, I have occasionally since that time, observed and recorded such facts as related to the subject; intending, one day or other, to lay my observations before the public, if they acquired an importance that deserved attention. In the outset of this undertaking, nothing seemed wanting but accurate thermometers, and a moderate portion of time and attention; and I embraced in imagination the whole effects of temperature upon health and disease; -a range of inquiry which experience has convinced me it would be temerity and folly to hope to go through. In this general view of the subject, some valuable communications have however been made to me by my much respected friend Dr. Percival; which do not apply to the particulars treated of in the following volume, but which I am not without the hopes of specifying at some future period, when I may have occasion to employ them.

Though I have some time seen, that the delay

delay of publishing till my original plan was executed, was likely to render my labours wholly abortive, or to convert the imperfect product from a gift into a legacy; yet I should not have given to the world so detached and unfinished a work as the present, according to the views I had originally entertained, but for the circumstances I am about to relate.

By the accounts received at the beginning of the present year, the fever of the West Indies appeared to continue its desolating progress with little abatement, and in America to be beginning its ravages anew.-With this pestilence, science seemed hitherto to have contended, in a great measure, in vain, and new methods of opposing it, were not merely justifiable, but requisite. At this time an account of the success of the nitric acid in Lues and Hepatitis, as employed by your correspondent Mr. Scott, of Bengal, was through your means given to the public. His theory, suggested by the new chemistry, did not appear promising; but it was neither wise nor candid to reject his experience, on that account. The first trials which I made of his practice persuaded me, that, though the success of the nitric acid might have been exaggerated by a warm imagination and a benevolent heart, it assuredly did succeed in certain cases of the dis-

eases

eases in question, and that a remedy of considerable power, and of perfect safety, was introduced into medicine. These opinions subsequent experience has served to confirm. In the fever of the West Indies mercury had been much employed, and though different notions were entertained of its mode of operation, it seemed on the whole the most approved remedy. Since the nitric acid appeared to be a substitute for mercury in other cases, it seemed reasonable to try its effects in this fatal fever. I suggested this practice by letter to some practitioners in the West Indies, and I submitted my notions on the subject to you; sensible that your character and station might bring that into immediate notice, which the influence of a private individual could but slowly effect.-Whatever difference there might be in some of our opinions, I was confident there was none in our wishes for alleviating human misery, and mitigating the destruction of this desolating war. Your conduct even exceeded my expectation.

Our correspondence on this occasion turned my views to the other means of opposing this pestilence. Ablution with cold water in fever had been so long employed at the hospital here, and in private practice, by my friends and colleagues Dr. Brandreth and Dr. Gerard, as well as myself, that it was become general in Liverpool, and common in the county of Lancaster. So long ago as the year 1791, a general statement by Dr. Brandreth of its advantages had been published by Dr. Duncan, in the Medical Commentaries of that year. It had also been noticed by me in the philosophical transactions for 1792, (see Appendix, No. II. p. 39) and I had repeatedly mentioned it in private correspondence; it had often been recommended to the surgeons of African ships in those examinations required by the legislature, and which are chiefly made by the physicians and surgeons of our hospital. On different occasions likewise I had not only explained, but exhibited the practice, to practitioners from a distance, and particularly to one or two going to the West Indies. A method of treatment so bold and so contrary to common prejudices, made however, as it appears, slow progress. The mode of operation of our remedy has been misapprehended; the proper period for using it has not been understood; and on some occasions having been resorted to improperly, the consequences have brought it into disrepute. Reflecting on these circumstances, and exposed by situation to the re-iterated sounds of death from the Western world, my decision was speedily made: I resolved no longer to delay an account of our treatment of fever, in the expectation a 4

pectation of including it in a larger field of discussion, and of presenting it in a form more conducive to reputation; and the fruit of this determination is the work now presented to you. In treating my subjects, perspicuity has been studied rather than rigorous method; I have every where endeavoured to make my steps so plain, that they may be distinctly traced; the most important points are impressed again and again to guard against mistake; my thermometrical observations have enabled me to give a precision to the directions for the use of the affusion of cold water which otherwise they could not have had; and, if I do not flatter myself, have laid a foundation for my reasonings, which speculations on fever have seldom possessed. I have guarded against the unnecessary use of technical as well as of general expressions. It were better perhaps that medicine, like other branches of natural knowledge, were brought from its hiding-place, and exhibited in the simplicity of science and the nakedness of truth. If it had been in my choice I would not have adopted the language of theory, like Boerhaave, or Sydenham; but have exhibited a medical work in the phraseology that Bacon had he lived in our days, might have used. Unfortunately in the present state of medical knowledge, wholly to avoid the language of theory is impossible. The corruptions of false doctrines

must remain more or less, in our phraseology, after the doctrines themselves are exploded; since custom has rendered the expressions in which they are found, intelligible, and human sagacity has not yet discovered those first principles of living motion by which the doctrines and the language of physiology might at once be reformed. Hence the term re-action is applied to certain motions of life, though in a sense very different from that in which it is used in the science of inanimate motion from which it is borrowed; and such words as tone will still be found in the following pages, though the theory that introduced them into medicine be universally abandoned. The use of such expressions is however an evil, justified only by necessity; and I have endeavoured to avoid it as much as lay in my power.

Possibly this notice may procure me some readers among men of general science; and this I confess to be one of my objects in dedicating the work to you. It is naturally an author's wish that his book may be read by those who can appreciate it, and who from their situations may have it in their power to bring its precepts into practice. In both these points of view I appeal to you—to your scientific knowledge, and to your generous heart. The work that I address to you is in a great measure practical. A man of genius

at the head of a fleet or army, would probably find little difficulty in understanding it; and possibly, if he understood it, there might be occasions on which it would afford scope to his humanity and patriotism. But whatever be its fate with men professedly military, I trust it will not be overlooked by the medical practitioners of our fleets and armies; a most meritorious class of the profession, to whom a great part of the improvements in the modern practice of medicine is to be ascribed.

Conceiving that the circulation of this volume, as well as its usefulness, might be extended by connecting the history of the affusion of cold water in fever, with other views of the same remedy, and with a few observations on the other remedies in fever, I have entered upon these points, without any very strict regard to method; and have been insensibly led to speak of some of the operations of temperature on the body in health, a subject which I had reserved. Such as it is, this volume may serve as the first of a series on similar subjects, if I should ever write them; and it may, I hope, stand alone, if I should write no more.

I am sensible that some of these particulars would have appeared with more propriety in a professed

professed preface; but having entered on certain explanations in my address to you, I have given the whole of these preliminary observations in the same form; a freedom that I trust you will forgive.

I cannot conclude without declaring the sense I entertain of your candour and politeness. Accept the tribute of my respect.—May you live long to cultivate and protect the sciences—the sciences, whose utility is beyond dispute; whose progress is superior to obstruction; and which, of all the possessions of man, seem least to partake of the imperfection of his nature!

I have the honour to be,
Sir,
Your faithful

And very obedient Servant,

JAMES CURRIE.

Liverpool, 31st October, 1797.

ERRATA.

Page 17, line 6, for hour, read hours.

39. line 7, after difeafe infert bas.

77, note, line 3d from the bottom, for naufea, read naufea. 78, note, line 10th from the bottom, for mihe, read mihi.

90, for fimiliar read fimilar.

92, note, bottom line, for ancient, read anciente.

96, line 6th from the bottom, for frigidimiffimi, read frigidiffimi.

- line 5, for deficit, read defecit. 105, line 1, for drank, read drunk.

- line 3d from the bottom, the fame correction.

121, line 6, tor no, read not.

--- last word, for own, read now.

127, line 5th from the bottom, for refrigerat, read refrigeret.

201, line 2d from the bottom, for by for, read for by.

203, line r5, for they, read the.

260, line 4th from the bottom, after into, infert a.

268, last line, for edition, read edition.

Appendix, page 46, line 6th, for gentlemen, read gentleman.

N. B. Where perspiration is employed in the following volume without any adnoun, it is sensible perspiration or sweating that is meant, unless where the context evidently shews that both sensible and insensible perspiration are included.

TABLE OF CONTENTS.

TABLE OF CONTENTS!

		P
CHAP. I. Narrative of Dr. Wright,		1
In a voyage from Jamaica to Liverpool, he		
employed the affusion of cold water in fever,	I	to 5
CHAP. II. History of a fever which broke out		3
in the Liverpool Infirmary,	5020	6
The affusion of cold water first practised in this		1103
fever by the author, December, 1787,	6	8
CHAP. III. History of a fever which occurred		dás
in the 30th Regiment,		9
How produced,	9	10
Situation and symptoms of the fick,	nnes	10
The progress of the infection how stopped,	11	13
In what fituations this practice may be imitated,	13	14
CHAP. IV. The manner in which the affusion	3900	
of cold water should be used in fever,		15
The kind of fever in which it is to be used,	15	16
The time and manner of using it,		17
Not to be used in the cold stage, nor when the		
heat is below what is natural, nor in the fweating	PERSONAL PROPERTY.	
flage	19	20
CHAP. V. Cases in which the affusion of cold		
water was used in different stages of fever,	10/15/	21
Seven cases detailed,	21	30
Comparison of the advantages attending the		
affusion of cold water at different periods of fever.		30
CASE, VIII. The cold affusion used in inter-		
mittent fever,	31	32
	Cir	D

Not

TABLE OF CONTENTS.	P.	iii P.
Not so permanent in its effects as the affusion		
of cold water,		70
Sponging or wetting the furface,		71
Digression on the general action of cold, .	72	75
Affusion of cold water practised in a fever at		
Breflaw in Silefia, 1737,	76	83
Warm bath, how used by the Greeks in the he-		
	83	85
CHAP. XI. Of the internal use of cold water		
in fever, the bill of hand annual to the side!		86
Practifed among the ancients,	86	87
The Boerhaavean theory of fever brought it	H W	
into difrepute,		87
And the fatal accidents following its improper		
administration,		28
How to be used with advantage,		92
The fame rules apply to drinking cold water, as	A SHI	19:13
to its affusion on the skin,		93
CHAP. XII. Of the disease that arises from drinking		
cold liquids or using the cold bath after severe exerci		95
	96	98
Sufferings of the army of Alexander the Great,		-
in their march through the defarts of the Sog-		
diani, &c	00	100
Dr. Rush's treatment of this disease contro-	77	-
verted, Ole	102	108
When the cold bath may be used with safety,		
Observations on the custom of the Roman		-09
youth of plunging during exercise into the Tyber,	110	170
The danger of Alexander the Great from bath-	Donse	19
ing in the Cydnus, explained,	10	770
Dr. Franklin quoted and controverted,		
Temperature of springs in different climates,	Narr	

- Notices

TABLE OF CONTENTS.		v
Marine of Girls Indian China Division	P.	P.
Notices respecting their use in the Plague, &c		186
Communication on this subject with Mr. How	ALCO CONTRACTOR	187
CHAP. XV. Animal heat—its origin. Perf-		
piration—its cooling influence. Mode of operation		
of other remedies in fever—antimonials—opium, &c.		190
Origin of animal heat a recent chemical difco-		
very,	191	192
Evaporation from the furface regulates the liv-		
ing heat,	193	194
Quantity of the matter perspired uncertain-	9200	
Sanctorius and Mr. Cruikshanks referred to,	196	197
Cooling Effects of evaporation discovered by	ATTE	
Dr. Cullen, and applied to explain the coolness of	192	
the body in an instance of extraordinary external	(TETO)	
heat, by Dr. Franklin,	197	198
by Dr. Bell,	1000	199
Cooling effects of the fecretion of fweat,	9 9979	200
Observations on the diseases of Europeans in	1000	
the West Indies,	13900	202
Perspirable matter of the negroe, .		204
Unguents of the ancients,	. T 5 23	205
Practical remarks on the warm bath,	207	209
General view of the operation of opium,	o her	210
Cafe illustrating its connexion with the heat of	FINA	
the fystem,	1976	211
When opium produces fleep,	17/1	212
How and when to be administered in fever,	2000 H	213
General view of the operation of alcohol .		215
How and when to be administered in fever,		216
Note respecting the use of cold in inflammator	v	
fevers, &cc.	218	220
CHAP. XVI. Population of Liverpool. Pre-		9
valence of fever among the poor, with documents.		
A		enefit

P	•	P
Benefits derived from the Manchester House of Re-		
covery in febrile diseases. Propriety of the general		
establishment of similar institutions,	683	221
Population of Liverpool,	#ST	222
Charitable institutions in Liverpool for the fick		
and hurt poor, the		224
General table of diseases entered at the Dif-		
penfary for feventeen years, with the comparative		
healthiness of the different months, . 22	6	228
General table of fevers entered at the Difpen-		
fary for feventeen years, &c		230
Average number of patients in fever on the		Die
books of the Difpenfary,		231
Information on the means of prevention of fe-		2112
ver, in the writings of Dr. Percival, &c.		235
Account of the House of Recovery at Man-		-
chefter, by Dr. Ferriar,		236
History of the receptacles for fever at Liverpool,		238
Benefits refulting from hospitals for contagious		541
difeafes,		242
CHAP. XVII. Is there an inhalation by the		
Skin? Case of P. M. Esq. who died of inanition.		400
General reflections. Conclusion,		244
The inhalation of the fkin a doctrine universal		7
among the ancients, and general in our own days,		244
Experiments that render it improbable, 2.	45	246
Cafe of Mr. M		261
Reflections on this cafe	SAN	261
The author's opinion that there is no inhala-		
tion by the fkin, supported by the experiments of		
M. Séguin,		265
Phenomena supposed to depend on this absorp-		
tion explained differently, 2	67	269
	Th	cory

TABLE OF CONTENTS.		vii
	P.	P.
Theory of Dr. Rutherford on diabetes,		269
Conclusion,		273
APPENDIX, No. I. Of Tetanus and convulsive		
diseases,		1
CASE 1, Illustrating the effects of opium, mer-		
cury, and the cold bath,	2	6
CASE 2, 3, 4, 5, 6,	6	8
CASE 7, Tetanus cured by wine, .	9	12
Use of cold bath in other convultive diseases,	13	16
General remarks,	17	19
APPENDIX, No. II. An account of the remark-		
able effects of a Shipwreck on the mariners; with		
experiments and observations on the influence of im-		
mersion in fresh and salt water, hot and cold, on		
the powers of the living body.		20
Narrative of the shipwreck,	20	25
Experiments on immersion, &c	25	42
General conclusions,	42	43
Postscript respecting the manner of taking the		
heat of the living body,	44	45
APPENDIX, No. III. Extracts from more re-		
cent publications of Dr. Wright on the treatment of		
fever, &c. Also an extract from M. Fourcroy's ac-		
count of the experiments of M. Seguin on the inha-		
lation of the surface of the bodyt		46
Ablution with cold water in fever used by Dr.		
Gregory,	46	47
Recent practice of Dr. Wright, in the ship		71
fever, and the yellow fever, &c	48	53
New practice in the plague,		54
Experiments of Séguiu on cutaneous Inhalation	56	62

MEDICAL

TABLE OF CONTENTS.

	Theory of Dr. Ruther (ont on disherer, and
	Concloberation
	the state of the s
	Car I, Hubrain, in office of opium, mar-
	cury and the cold but.
	Cast 27 3 4 5 6.
	Care 7. Tetrant cared my place
	The alocal bearing other convenience defeates!
	depresentation and other receiver on the engineers of their
	distributed of the fields bedy to the training side
	Manufactured the dispurent, a control of
	Cicross conclutions and a second second
	Policy relacions the manner of taking the
	ous Adrives to springers and Superpolar of grant and
	Second profile of Dr. Weels, in the rise the
181	fever, and the galley fever, fee,
	maker process in the History of the contract of

MEDICAL

MEDICAL REPORTS, &c.

mailguity. I attended this person often, but could

not prevail with him to be removed from a dark

CHAP. I.

Narrative of Dr. Wright.

IN the London Medical Journal for the year 1786, Dr. William Wright,* formerly of the island of Jamaica, gave an account of the successful treatment of some cases of sever by the ablution of the patient with cold water.

"On the 1st of August, 1777," says Dr. Wright, "I embarked in a ship bound to Liverpool, and sailed the same evening from Montego Bay. The master told me he had hired several sailors on the same day we took our departure;

B one

^{*} Now I believe physician to the army in the West Indies.

one of whom had been at fick quarters on shore, and was now but in a convalescent state. On the 23d of August we were in the latitude of Bermudas, and had had a very heavy gale of wind for three days, when the above-mentioned man relapsed, and had a sever, with symptoms of the greatest malignity. I attended this person often, but could not prevail with him to be removed from a dark and confined situation, to a more airy and convenient part of the ship; and as he resused medicines, and even food, he died on the eighth day of his illness.

"By my attention to the fick man I caught the contagion, and began to be indisposed on the 5th of September, and the following is a narrative of my case, extracted from notes daily marked down: I had been many years in Jamaica, but, except being somewhat relaxed by the climate, and satigue of business, I ailed nothing when I embarked. This circumstance, however, might perhaps dispose me more readily to receive the infection.

"Sept. 5th, 6th, 7th, Small rigors now and then—a preternatural heat of the skin—a dull pain in the forehead—the pulse small and quick—a loss of appetite, but no sickness at stomach—the tongue white and slimy—little or no thirst—the belly regu-

lar—the urine pale, and rather scanty—in the night restless, with starting and delirium.

- "Sept. 8th, Every fymptom aggravated, with pains in the loins and lower limbs, and stiffness in the thighs and hams.
- "I took a gentle vomit in the second day of this illness, and next morning a decoction of tamarinds; at bed-time, an opiate, joined with antimonial wine, but this did not procure sleep, or open the pores of the skin. No inflammatory symptoms being present, a drachm of Peruvian bark was taken every hour for six hours successively, and now and then a glass of Port wine, but with no apparent benefit. When upon deck, my pains were greatly mitigated, and the colder the air the better. This circumstance, and the failure of every means I had tried, encouraged me to put in practice on myself what I had often wished to try on others, in severs similar to my own.
- "Sept. 9th, Having given the necessary directions, about three o'clock in the afternoon, I stripped off all my cloaths, and threw a sea cloak loosely about me till I got upon the deck, when the cloak also was laid aside: Three buckets full of salt water were then thrown at once on me; the shock was great, but I selt immediate relief. The head-ach

and other pains inftantly abated, and a fine glow and diaphorefis fucceeded. Towards evening, however, the febrile fymptoms threatened a return, and I had again recourse to the same method as before, with the same good effect. I now took food with an appetite, and for the first time had a sound night's rest.

- "Sept. 10th, No fever, but a little uneafiness in the hams and thighs—used the cold bath twice.
- "Sept. 11th, Every fymptom vanished, but to prevent a relapse, I used the cold bath twice.
- "Mr. Thomas Kirk, a young gentleman, paffenger in the same ship, sell sick of a sever on the 9th of August. His symptoms were nearly similar to mine, and having taken some medicines without experiencing relief, he was desirous of trying the cold bath, which, with my approbation, he did on the 11th and 12th of September, and, by this method, was happily restored to health. He lives at this time (Jan 1786) near Liverpool."

To this interesting narrative, Dr. Wright adds some general observations on the traces that are to be found of the use of cold water internally and externally in severs, in several works ancient and modern.

dern. But whether he himself pursued this practice any farther, I have not been informed.

Having before experienced that Dr. Wright was a fafe guide,* I immediately on reading this narrative, determined on following his practice in the present instance; and before an opportunity occurred of carrying my intention into effect, I was farther encouraged, by learning, that my respectable colleague, Dr. Brandreth, had employed cold water externally in some recent cases of sever with happy effects.

^{*} See a paper in the Memoirs of the London Medical Society, vol. iii. p. 147, to be found in a subsequent part of this volume.



CHAP. II.

Having before experienced that Dr. Wright was

History of a fever which broke out in the Liverpool Infirmary.

tive, electromed on tollowing his practice in the

prefere inflance a and before an opportunity occur-

ON the 9th of Dec. 1787, a contagious fever made its appearance in the Liverpool Infirmary. For fome time previously the weather had been extremely cold, and the discipline of the house, owing to causes which it is unnecessary to mention, had been much relaxed. The intenfity of the cold prevented the necessary degree of ventilation, and the regulations for the preservation of cleanliness had been in some measure neglected. These circumstances operated particularly on one of the wards of the eaftern wing, employed as a lock-hospital for females, where the contagion first appeared. The fever spread rapidly, and before its progress could be arrested, fixteen persons were affected, of which two died. Of these fixteen, eight were under my care. On this occasion I used for the first time the affusion of cold water, in the manner described by Dr. Wright. It was first tried in two cases only,

the one in the fecond, the other in the fourth day of fever. The effects corresponded exactly with those mentioned by him to have occurred in his own case; and thus encouraged, the remedy was employed in five other cases. It was repeated daily, and of these seven patients, the whole recovered .-In the eighth case, the aspersion of cold water seemed too hazardous a practice, and it was not employed. The ftrength of the patient was much impaired by lues, and at the time of catching the contagion, she laboured under ptyalism. I was not then aware that this last circumstance formed no objection against the cold affusion, and in a situation so critical, it was thought imprudent to use it. The usual remedies were directed for this patient, particularly bark, wine, and opium, but unfuccefsfully; she died on the 16th day of her difeafe. do not eready deceive mytelf, may affort

From this time forth, I have conftantly wished to employ the affusion of cold water in every case of the low contagious sever, in which the strength was not already much exhausted; and I have preserved a register of a hundred and sifty three cases, in which the cure was chiefly trusted to this remedy. Of these, ninety-sour occurred in the hospital in the four years subsequent to the period already mentioned, twenty-seven in private practice, and thirty-two in the 30th regiment of foot, when quartered in Liverpool in the year 1792. Of late I have

have not thought it necessary to register all the cases in which this remedy has been employed.-Having satisfied myself of its extraordinary efficacy, and of the precautions necessary in using it, I have found it the shorter method, as well as the more instructive, to record the instances in which it has proved unfuccessful. To relate the whole of my experience would be a tedious and an useless labour. I purpose to digest the results under a few distinct heads, supporting and illustrating each general proposition by an ample detail of cases. Before, however, this preliminary account is closed, it will be useful to enter more particularly into the history of the contagious fever which broke out in the 30th regiment, because the account of its rife, progress, treatment, and termination, will support in a striking manner the doctrines I wish to establish, and if I do not greatly deceive myfelf, may afford important instruction, as well as encouragement, to those whose duty may call them to oppose the progress of contagious fever in fimilar fituations.

CHAP. III.

or benefit show on box your dei loopsvil on

cover could adams no autor. The consacion con-

History of a fever which occurred in the 30th regiment.

THE 30th regiment, as is usual with troops in Liverpool, was billetted in the town, but paraded and mounted guard in the fort, situated north of the town, and on the banks of the river. The general guard-room had been used previous to the arrival of the 30th, as a place of consinement for deserters; it was extremely close and dirty, and under it was a cellar, which in the winter had been full of water. This water was now half evaporated, and from the surface issued offensive exhalations.

In a dark, narrow, and unventilated cell, off the guard-room, it was usual to confine such men as were sent to the guard for misbehaviour, and about the beginning of June (1792) several men had been shut up in this place on account of drunkenness, and suffered to remain there twenty-sour hours, under the debility that succeeds intoxication. The typhus, typhus, or jail fever, made its appearance in two of these men about the sirst of the month, and spread with great rapidity. Ten of the soldiers labouring under this epidemic, were received into the Liverpool infirmary, and the wards allotted to sever could admit no more. The contagion continuing its progress, a temporary hospital was sitted up at the fort, and I was requested to give my assistance there to the surgeon of the regiment, by Captains Brereton* and Torriano.

In two low rooms, each about fifteen feet square, were fourteen patients labouring under sever. They were in different stages of its progress: one was in the fourteenth day of the disease, two were in the twelfth, and the rest from the ninth to the fourth inclusive. The symptoms of the sever were very uniform. In every case there was more or less cough, with mucous expectoration: in all those who had sustained the disease eight days and upwards, there were petechiæ on the skin; in several there were occasional bleedings from the nostrils, and streaks of blood in the expectoration. The debility was considerable from the first, and it had been

^{*} Now Colonel Brereton of the 63d foot.

[†] The gallant and accomplished officer who fell soon after at the heights of Pharon, in the defence of Toulon.

been increased in several cases by the use of venesection, before the nature of the epidemic was understood. The pulse varied from 130 strokes in the
minute to 100; the heat rose in one case to 105°
of Fahrenheit, but was in general from 101° to
103°; and towards the latter stages of the disease it
was scarcely above the temperature of health.—
Great pain in the head, with stupor, pervaded the
whole, and in several instances there occurred a
considerable degree of the low delirium.

Our first care was to ventilate and clean the rooms, which were in a high degree foul and pestilential. Our second was to wash and clean the patients themselves. This was done by pouring seawater, in the manner already described, over the naked bodies of those whose strength was not greatly reduced, and whose heat was steadily above the temperature of health. In those advanced in the fever, whose debility was of course great, we did not venture on this treatment, but contented ourselves with spunging the whole surface of the body with tepid vinegar, a practice that in every stage of fever is salutary and refreshing.

Our next care was to stop the progress of the infection. With this view, the guard-house was at first attempted to be purified by washing and ventilation, the greatest part of its furniture having been burnt

burnt, or thrown into the sea. All our precautions and exertions of this kind were however found to be ineffectual. The weather was at this time wet. and extremely cold for the feafon; the men on guard could not be prevailed on to remain in the open air; and from paffing the night in the infected guard-room, feveral of the privates of the fucceffive reliefs, on the 10th, 11th, and 12th of the month, caught the infection. In several of these the fever ran through its course; and in others, it was immediately arrested by the affusion of sea-water as already described. No means having been found effectual for the purification of the guard-room, it was shut up, and a temporary shed erected in its stead. Still the contagion proceeded; on the morning of the 13th three more having been added to the lift of the infected. On that day, therefore, the whole regiment was drawn up at my requeft, and the men examined in their ranks: Seventeen were found with fymptoms of fever upon them .-It was not difficult to diffinguish them as they stood by their fellows. Their countenances were languid, their whole appearance dejected, and the adnata of their eyes had a dull red fuffusion. These men were carefully separated from the rest of the corps, and immediately subjected to the cold affusion, always repeated once, and fometimes twice a day .-In fifteen of the number the contagion was extinguished; but two went through the regular disease. On

On the same day, the commanding officer, at my defire, issued an order for the whole of the remaining part of the regiment to bathe in the sea; and for some time they were regularly mustered, and marched down at high water, to plunge into the tide.

These means were successful in arresting the epidemic: after the 13th of June no person was attacked by it. It extended to fifty-eight persons in all, of which thirty-two went through the regular course of the fever, and in twenty-fix the disease feemed to be cut short by the cold affusion. Of thirty-two already mentioned, two died. Both of these were men whose constitutions were weakened by the climate of the West Indies; both of them had been bled in the early stages of the fever; and the one of them being in the twelfth, the other in the fourteenth day of the disease when I first visited them, neither of them was subjected to the cold affusion. The water employed on this occasion was taken up from the river Mersey close by the fort. It was at that time of a temperature from 58° to 60° of Fahrenheit, and it contains in folution from a 32d to a 33d part of sea-salt.

In hospitals, manufactories, and prisons—situations in which the low contagious sever so frequently originates, the practice I have detailed may be followed followed with great ease, safety, and advantage; but it is in a more particular degree applicable to this contagion when it appears on ship-board, because in that situation the usual means of prevention or cure are necessarily limited, and the imminence of the danger requires a remedy that operates with speed as well as essicacy. The waters of the ocean assord this remedy; in every point of view a most happy one for mariners, since it can be applied almost as easily as it can be procured.

these were men whose, conflitutions were weak ned

them, acidies of them, was findedied to the cold al-

taken up from the ever Merfey close by the tire is

of Palesconoit, and it contains in ISlan Monorday to

band that I make should sale to vale and CHAP.

chap. Iv.

The manner in which the affusion of cold water ought to be used in fever.

" rus; peillus parvus, econocio percuaque frequens;

HAVING given this general account of my experience of this remedy in fever, it will now be necesfary to enter more particularly on the rules which ought to govern its application, and on the different effects to be expected from it, according to the different stages of the disease in which it is employed. It will be proper to premise, that when the term fever is used in the present work without any adjunctive, it is the low contagious fever that is meant. This is the Typhus of Dr. Cullen; the contagious fever of Dr. Lind; the Febris inirritativa of Dr. Darwin. In popular language, it is generally called the nervous fever, and where particular fymptoms appear, the putrid fever. It is usually produced in fituations where there is a want of cleanliness, and more especially of ventilation; and when produced, it is propagated by contagion. This is the common fever of England; its symptoms have been

been detailed with great minuteness in a variety of modern publications, and I have therefore declined repeating descriptions that are every where to be met with. Dr. Cullen has defined the disease as follows:-- "Morbus contagiofus; calor parum auc-" tus; pulsus parvus, debilis, plerumque frequens; " urina parum mutata; sensorii functiones pluri-" mum turbatae; vires multum imminutae." fixteen years practice I have found the contagious fever of Liverpool remarkably uniform, and in general to correspond exactly with this concise and perspicuous definition. This disease prevails chiefly among the poor, who from the nature of their diet and habits, are peculiarly exposed to the causes that produce it. Seldom extending itself in any confiderable degree among the other classes of the community, it has been supposed that Liverpool was little subject to fever; but this will be shewn, from authentic documents, to be a great and a pernicious error. Let us proceed at present to inquire into the rules that ought to govern the use of the affusion or aspersion of cold water in this disease.

Whoever has watched the progress of sever, must have observed the justness of the observation made by Cullen, Vogel, De Haen, and others, that even those genera which are denominated continued, are not strictly such, but have pretty regular and distinct exacerbations and remissions in each diurnal period.

period. In this space of time, Dr. Cullen contends, that an attentive observer may commonly distinguish two separate paroxysms.* My observations do not enable me to confirm his polition in its full extent-but one exacerbation, and one remission in the twenty-four hour, feem generally observable. The exacerbation usually occurs in the afternoon, or evening, the remission towards morning. These exacerbations are marked by increased flushing, thirst, and restlessness. If the heat of the patient be, at fuch times, taken by the thermometer, it will be found to have rifen one or two degrees in the central parts of the body, above the average heat of the fever, and still more on the extremities .-The fafest and most advantageous time for using the aspersion or affusion of cold water, is when the exacerbation is at its height, or immediately after its declination is begun; and this has led me almost always to direct it to be employed from fix to nine o'clock in the evening; but it may be fafely used at any time of the day, when there is no sense of chilliness present, when the heat of the surface is steadily above what is natural, and when there is no general or profuse perspiration.—These particulars are of the utmost importance.

rug blutorq vobnu si vi C

^{*} Paroxyímis quovis die binis.

- 1. If the aspersion of cold water on the surface of the body be used during the cold stage of the paroxysm of fever, the respiration is nearly suspended; the pulse becomes fluttering, feeble, and of an incalculable frequency; the furface and extremities become doubly cold and shrivelled, and the patient feems to struggle with the pangs of instant dissolution. I have no doubt, from what I have observed, that in fuch circumstances, the repeated affusion of a few buckets of cold water would extinguish life. This remedy should therefore never be used when any confiderable fense of chilliness is present, even though the thermometer, applied to the trunk of the body, should indicate a degree of heat greater The fafelt and most advantageous tim fairlibring the afpertion or affulion of cold water, is when the
 - 2. Neither ought it to be used, when the heat measured by the thermometer, is less than, or even only equal to the natural heat, though the patient should feel no degree of chilliness. This is sometimes the case towards the last stages of sever, when the powers of life are too weak to sustain so powerful a stimulus.
 - 3. It is also necessary to abstain from the use of this remedy when the body is under profuse perspiration, and this caution is more important in proportion to the continuance of this perspiration. In the commencement of perspiration, especially if it has

has been brought on by violent exercise, the affusion of cold water on the naked body, or even immerfion in the cold bath, may be hazarded with little risque, and sometimes may be reforted to with great benefit. After the perspiration has continued fome time and flowed freely, especially if the body has remained at rest, either the affusion or immersion is attended with danger, even though the heat of the body at the moment of using them be greater than natural.—Perspiration is always a cooling process in itself, but in bed it is often prolonged by artificial means, and the body is prevented from cooling under it to the natural degree, by the load of heated clothes. When the heat has been thus artificially kept up, a practitioner, judging by the information of his thermometer only, may be led into error. In this fituation, however, I have observed that the heat finks rapidly on the exposure of the furface of the body even to the external air, and that the application of cold water, either by affusion or immersion, is accompanied by a loss of heat and a deficiency of re-action, which are altogether inconfistent with fafety.-Each of these points will be illustrated more fully in the fequel.

Under these restrictions the cold affusion may be used at any period of sever; but its effects will be more salutary in proportion as it is used more early. When employed in the advanced stages of sever,

C 2

where

where the heat is reduced, and the debility great, fome cordial should be given immediately after it, and the best is warm wine. The general effects of the cold affusion will be more clearly illustrated by the following cases. They are a selection from a great number, the records of which have been preserved, and which lead to the same results. They are so arranged as to exhibit the salutary effects of this remedy in the different stages of sever, and illustrate the precautions laid down against using it improperly. If they should appear tedious after what has already been mentioned, this must be forgiven; on a subject so important and so little understood, it is better to incur the charge of tediousness than the hazard of being obscure.

formation of its shermoureter only, may be fed into

which are any period of level; had us effects will

this arming state to delice the CHAP.

CHAP. V.

CASE I.

Cases in which the Affusion of Cold Water was used in the different stages of sever.

lot's of colomba were ordered for nor, wall

January 1, 1790.

A Nurse in the sever ward of the Infirmary having several patients under her care, caught the infection. She was seized with violent rigors, chilliness, and wandering pains, succeeded by great heat, thirst, and head-ach. Sixteen hours after the first attack, her heat at the axilla was 103° of Faht. her pulse 112 in the minute and strong; her thirst great, her tongue furred, and her skin dry.

Five gallons of falt water, of the temperature of 44° were poured over her naked body, at five o'clock in the afternoon, and after being hastily dried with towels, she was replaced in bed: when the agitation and sobbing had subsided, her pulse

was found to beat at the rate of 96 strokes in the minute, and in half an hour afterwards it had fallen to 80. The heat was reduced to 98° by the ablution, and half an hour afterwards it remained stationary. The sense of heat and the head-ach were gone, and the thirst nearly gone. Six hours afterwards she was found perfectly free of sever, but a good deal of debility remained.

Small doses of colombo were ordered for her, with a light nourishing diet, and for several days the cold affusion was repeated at the same hour of the day as at first; the sever never returned.

Namenty T. 1750.

During the progress of fever when epidemic, a great number of cases similar to the above have occurred, in which the disease was suddenly cut short by the use of the cold affusion on the first and second day; twenty six of these cases were in the 30th regiment, as has already been stated. In such instances, the result was so precisely similar to what occurred in the case I have related, that it would be to no purpose to detail them.

When an epidemic fever is spreading, and the danger is known, patients will take the alarm on the first attack, and the power, as well as the utility of such a remedy, as the cold affusion, in such situations

tions of general danger, will be easily imagined.—
It cannot be employed too foon after the first attack, provided the original chill is over, and the hot stage is firmly established.

In cases in which the affusion was not employed till the third day of sever, I have seen several instances of the same complete solution of the disease. I have even seen this take place, when the remedy has been deserred till the sourth day; but this is not common.—The sollowing case will point out the usual effects of this remedy in the third and sourth days of the disease.

CASE II. and district aid

On the 18th at doon his pulle was on and Tole ;

Jan. 17, 1790. A. B. aged nineteen, a pupil of the Infirmary, caught the infection in attending the fever-ward. When I saw him, seventy-eight hours had elapsed since the first attack: he was of course in the fourth day of the disease. He had all the usual symptoms—head-ach, thirst, furred tongue, pain in the back and loins, with great debility. His heat was 101°, and his pulse 112 in the minute.

A bucket full of falt water was poured over him as usual, at noon on the 17th. His heat sunk to 99°, and his pulse to 98 in the minute. A profuse C4 perspiration

perspiration followed, with the cessation of all his feverish symptoms.—This intermission continued for several hours, during which he enjoyed some comfortable sleep: but at five in the afternoon he was again seized with severish rigors, followed by heat, thirst, and head-ach, as before. An hour afterwards, the hot stage was established; his heat was 100°, his pulse 100. The same quantity of cold water was again poured over him, and with similar effects. His pulse sell immediately to 80, and became more sull; his heat became natural. The following night he took twenty drops of laudanum and slept well.

On the 18th at noon his pulse was 96 and soft; his skin moist, but a little above the natural heat. His tongue was a little furred, and his head ached: he also complained of thirst; the heat at the axilla was 100°.—The same remedy was again applied. He was greatly refreshed by it. The pulse sell to 90, the skin became cool, the thirst went off, and all the severish symptoms vanished.

On the 19th his pulse was 88, his heat natural, the thirst and head-ach were gone, and his appetite improving. The ablution was repeated for the last time at fix o'clock in the evening.

On the 20th his pulse was 78 and soft, his tongue clean,

clean, and his appetite farther improved. He had still some remains of debility on the 21st, but on the 22d he was free of complaint. This patient during his fever took no medicine but the effervescing mixture, the dose of laudanum excepted.—The affusion was used four times.

beis camons on and CASE III. on belo source

and carried to be described and more. This

Dec. 8, 1791. A woman aged fifty-seven, who had caught the infection, in her attendance on a poor family labouring under sever, came under my care in the hospital, fifty hours after the first attack. She was of course in the third day of sever; she had the usual symptoms—head-ach, pain in the back and loins, and thirst: her tongue was surred and her eyes heavy: her pulse 96, her heat 101°. The affusion of cold water was performed at noon, as usual. In a few minutes afterwards, the heat under the tongue was 98°, the pulse 80. Towards evening however, the severish rigor returned, with all the usual consequences. As soon as the hot stage was established, the affusion was repeated, and with the usual happy effects.

Dec. 9.—Noon—pulse 90 and seeble—respiration easy—heat 100°.—The affusion was immediately employed, and again repeated in the evening—perspiration,

piration, coolness, quiet, and sleep, were the confequence.

Dec. 10. This day the affusion was twice repeated as yesterday, viz. at noon, and at six in the evening.—At eight in the evening the pulse was 76, the skin soft, the heat 97°. the respiration easy and natural.—The sever returned no more. This patient used no other remedies but an enema, and after it an opiate, every night. The affusion was used twice every day; in all six times.

care in the belief CASE IV. desigled advai one

poor family labouring under fever, come under my

four, about seventy-two hours after the first attack of sever, became my patient. His pulse 100—his heat 104°.—The other symptoms as usual—but the pain in the head and back particularly severe. Two minutes after the affusion—pulse 90, heat 101°; six minutes after—pulse 90 and weak;—ten minutes after, pulse 90, heat 100°.—This patient selt great refreshment, and was entirely relieved of the pain in the head and back.—In the evening however the exacerbation of sever was severe, and the headach returned with violence.—He passed a restless night. About sour o'clock in the ensuing morning the affusion was repeated by his own desire. At 9,

A. M.

A. M. a general and gentle perspiration covered the surface of the body; the pulse was 84, the tongue moist, the skin cool, and the pains of the head and back entirely gone off. In the afternoon, however, the sever returned, though in a slighter degree. The affusion was repeated the fourth time with the same happy effects, and after this he had no return of the disease.

Thus it appears, that the cold affusion used on the third and sourth days of sever, does not usually produce an immediate solution of the disease; but that it instantly abates it, and by a sew repetitions brings it to a happy termination in two or three days.

CASE V.

Oct. 25. M. S. came under my care on the 24th Dec. 1791, on the feventh day of typhus, with the usual symptoms—pulse 108, heat 100°.— The cold affusion was immediately directed—two minutes afterwards the heat was 96°—three minutes afterwards 98°—the pulse 98. This patient experienced great relief. The affusion was repeated on this and the following day— a gentle diaphoresis always

always succeeded it, with tranquil sleep, and on the third day the sever was completely removed.

CASE VI.

A young lady of 19, in the 7th or 8th day of typhus, became my patient, Sept. 26, 1794. Her pulse was 112 and feeble, heat 101°. She had great pain in the head, and much proftration of strength -her eyes were fuffused and dull-her tongue furred-her spirits greatly depressed. Saline medicines were used for this patient, with lemonade for her usual drink, and moderate quantities of wine were given mixed with water. The burning fensation in the palms of her hands and temples was affuaged by frequent spunging with vinegar, and every evening at fix P. M. three or four gallons of cold brine were thrown over her. The happy effects so frequently described, were in this case particularly striking :- The pulse fell almost immediately to 90, the heat to its natural standard, and the head-ach vanished-a gentle diaphoresis followed, with eafy fleep: -in a few hours, however, the feverish symptoms returned, and towards the hour of fix in the evening the fever was as its highest state of exacerbation. At this hour therefore the affusion was repeated with the same happy effectsthough the fever returned as before, it was in a milder

milder form; the same practice was continued, and on the second of October, she was entirely free from the disease.

CASE VII.

attution having always been performed in the even-

aved, its usual period, hast on the tash day, the F. G. a foldier of the 30th regiment, aged 33, fell under my care on the 9th June, 1792, during the prevalence of the epidemic in that regiment, of which I have already given an account. He was in the 9th day of the disease-his pulse 100 and feeblehis heat 1040—his thirst was very great—his tongue foul and black-frequent cough occurred, with streaks of blood in the expectoration-and petechiæ appeared all over his body. His mind was at all times confused, and occasionally he was completely delirious. I directed that his strength should be supported by administering a bottle of wine every day, with an equal quantity of gruel; -that every night he should take an opiate draught, and that a complete operation of his bowels should be procured by a clyster administered daily, and if this did not fucceed, by a few grains of calomel. I also directed that a bucket full of falt-water should be thrown over him immediately, and repeated according to circumstances. In a few minutes after the affusion, the heat was 98°—the pulse 98—his mind was more calm and collected: two hours afterwards

he had relapsed into nearly his former state, but the night was passed more tranquilly. The whole of this practice was continued, with nearly the same result, till the 12th day of the disease, the affusion having always been performed in the evening, and occasionally at noon. The sever continued its usual period, but on the 12th day, the heat having sunk to its natural standard, the cold affusion was thenceforth omitted; we however, spunged the whole body once or twice a day with vinegar.—The patient was in a state of convalescence on the 18th day from the first attack.

foul and black-frequent cough occurred, with I have related this case the more circumstantially, because it contains the particulars of my practice in the epidemic in which it occurred, at the same time that it affords an example of the effects of the cold affusion used in the more advanced periods of fever. In fuch instances, as might be expected, it does not procure the fame advantages, as in the earlier stages, when the strength is less impaired, and the morbid actions less firmly affociated; nevertheless, it is evidently advantageous while the heat of the patient exceeds the natural standard, though it ought to be employed with caution in the more advanced stages of the disease, and in such cases, according to my later experience, of a temperature from fifteen to twenty degrees only, below the human I have related, the water employed was the pumpwater of our hospital, saturated with sea-salt, and of a temperature from 40° to 50° of Fah^t.

of August, two bours hefore the

The cold affusion may also be employed with success in intermittent severs, as I have sound by repeated trials, and as the following case will demonstrate.

case vill.

return, the rook the basis in larger quantity, the

Ann Hall, aged 22, was admitted into the Infirmary, July 19, 1792, under an obstinate quotidian of three months standing-she had from time to time taken the bark, but as the great delicacy of her stomach, would not permit her to use it in sufficient quantities, she was become very feeble and much emaciated. A gentle emetic was administered to her in the first instance, and on the commencement of the hot stage of the paroxysm, twenty drops of the tincture of opium were directed, after the practice of Dr. Lind. During the intermiffion, the bark was ordered to be taken in fuch doses as her stomach would bear; she was put on a nourishing diet, and was ordered a pint of port wine every day. This plan was purfued for fourteen days, but without fuccess; the paroxysms returned daily, though with fome irregularity-her strength was however

however rather improved. Still the delicacy of her stomach continued, and the bark, except in very small doses, was constantly rejected.

a temperature from 40° to 50° of Fair.

On the 8th of August, two hours before the expected accession of the sever, sour gallons of brine were dashed over her, of the temperature of 66° of Faht and this day she escaped the attack. In the interval between this and the period of the next return, she took the bark in larger quantity, the power of her stomach being increased: but on the 10th, two hours later then usual, the paroxysm returned with unusual severity. After the hot stage was completely formed, the brine was poured over her, as before—the symptoms instantly abated; she fell into a gentle perspiration, with prosound sleep. She afterwards continued the bark as before, and from this time forwards was free of disease.

It would be easy to multiply these details, but their uniformity has already perhaps rendered them tedious; a sew general observations therefore shall conclude this division of the subject.

CHAP. VI.

course, citizer for the purpose of int-

General Observations.

- 1. THOUGH the patients were often startled at the first proposal of dashing the cold water over them, yet after one trial, there was feldom any difficulty in perfuading them to have it repeated. The effects were in general highly grateful and refreshing to their sensations, the extinction or abatement of fever was commonly followed by more or less diaphoresis, and this again by refreshing fleep.
- 2. At first I used fresh water-afterwards fresh water mixed with vinegar-and laftly, a faturated folution of fea-falt in water. In the instance of the 30th regiment I used the water of the river, which contains about a thirty-third part of falt, as has been already mentioned, and this I commonly D employ

employ in private practice.* I was led to prefer falt-water to fresh on account of the stimulating effect of sea-salt on the vessels of the skin, by which I apprehend the debilitating action of cold is prevented. Salt-water, either for the purpose of immersion or affusion, is more grateful to the patient than fresh water, and it is well known that it may be applied to the surface for a length of time, with much less hazard. Persons immersed in sea water, and especially in saturated brine, for some time together, preserve the lustre of the eye and the ruddiness of the cheek, longer than those in fresh water, of an equal temperature, and such persons exhibit the the vital re-action stronger when removed from it.

I preferred

^{*} It may be supposed that the degree of impregnation of the river water with falt, must depend on the time of the tide when it is taken up. Thus, that it must be salter at the height of the flood, when the tide has run fix hours from the fea, than at the period of low water, when it has run nearly the fame time from the land .- I expected to find this the cafe. and endeavoured to ascertain the difference; but in a trial which I made by evaporating 40lbs. of water taken up at the height of the flood tide, and the fame quantity taken up at low water, I could perceive no difference in the proportion of falt, a circumstance which I am unable to explain. The small difference that there was in the residuum in favour of the water taken up at the top of the tide, arose evidently from its being mingled with a large portion of dirt—the water at the top of the tide is mixed with dirt and feculencies, that taken at the lowest point of ebb is nearly pure.

I preferred the brine to vinegar, as being cheaper, and more easily procured of the necessary quantity: otherwise, it is well known how grateful vinegar is to patients in sever, and perhaps a mixture of vinegar and water of the proper strength, might be prepreferable even to brine. But though I gave the preference to brine over fresh water, I have very often used the latter, and it is seldom that any danger can result from the want of a saline impregnation, where the cold is employed in so stimulating a form as that which has been described, and for so temporary a duration.

3. In taking the heat of the patient, I have generally used a small mercurial thermometer of great sensibility, with a moveable scale, made for me by Mr. Ramsden, after a form invented by the late Mr. Hunter, and used by him in his experiments on the heat of animals, and I have introduced the bulb under the tongue with the lips close, or under the axilla indifferently; having sound by repeated experiments, that the heat in these two places corresponds exactly, and gives a just indication of the heat of the surface of the body where sheltered by the necessary teguments, and secluded from the contact of the external air.

Finding, however, confiderable rifque in using the straight tubed thermometer in contagious dis-

eases, I got some instruments of this kind made with a small bulb curved at the end. The bulb being introduced under the tongue or the axilla, the observer can stand behind the patient, and mark the rise of the mercury, without coming into the immediate sphere of his respiration. Though no injury was in any case incurred from the use of this thermometer, yet a farther improvement has suggested itself. By introducing a small piece of iron into the tube, after the manner of Mr. Six, a permanent indication of the greatest heat is obtained, and the approach of the observer towards the patient during the experiment, is rendered unnecessary.

fentibility, with a moveshile fetile, made for me by

on the best of animal, and I have introduced the

chap.

CHAP. VII.

Precautions requisite in using the cold affusion, illustrated by Cafes.

was for fome minutes almost toppended; his pulle

1. IT was before remarked, that the cold affusion cannot be used with safety during the cold stage of the febrile paroxyfm: the following case will illustrate this truth. In the fummer of 1792 I was requested by Mr. Hoffman, an ingenious Prustian gentleman, and a furgeon in the army then under the command of the Duke of Brunfwick, to give him an opportunity of feeing the method of ufing this remedy. At that time there was a patient labouring under a tertian intermittent under my care in the infirmary, on whom it could with propriety be exhibited. Accordingly a time for meeting Mr. Hoffman in the fever-ward was appointed, when the hot stage of the paroxysm might be expected to be fairly formed. It happened however that the accession of the fever had occurred an hour later this day than might have been expected, and when we arrived. arrived, the patient was still in the cold stage of the paroxylm; the affiftants however proceeded: he was taken out of his bed shivering, his pulse small and frequent, his extremities shrunk and cold. In this state the brine was dashed over him as usual, but not with the usual happy effects:-his breathing was for fome minutes almost suspended; his pulse at the wrift was not to be felt; the pulfations of the heart were feeble and fluttering; a deadly coldness spread over the surface; and when respiration returned, it was short, irregular, and laborious .-After the use of frictions on the surface, and particularly on the extremities-of a fleady warmth applied for some time to the scrobiculus cordis—and of cordials cautiously administered in small quantitiesthe pulse at the wrist returned; but for some time it was exceffively quick and feeble. He recovered however in the course of an hour, and it was found that the paroxysm of fever had been extinguished: but the circumstances first related, were evidently full of danger, and they produced at the time much apprehension and uneasiness. The same remedy was however used in the hot stage of the ensuing paroxysm, and with the usual happy effects .-Other cases to the same purpose might be adduced if it were necessary.

I have frequently used the cold affusion in the hot stage of the paroxysm of intermittents, and almost always

to be fairly formed. It has beened however that the

always with the immediate folution of the fit; but in general, if no remedy be used in the intermission, the sever returns at the usual period. In some instances, however, the succeeding paroxysm has been prevented by using the cold affusion about an hour previous to the period of its expected return, and the disease been ultimately removed by continuing this practice through sour or sive of the sollowing periods.

The use of the cold affusion in the absence of sever, requires however a constitution in a great measure unbroken; and many of the intermittents which we see in Liverpool, being transmitted to us from the warm climates, adhere to constitutions in which this practice is not perfectly safe. In such cases, it may notwithstanding be adopted in the hot stage of sever with safety and advantage. Indeed it ought never to be forgotten, that an application of cold, which is safe in the violence of sever, is not safe when the sever is removed. Injury has sometimes occurred from continuing the cold affusion in the period of convalescence,

2. Neither is the cold affusion safe after the sweating stage of sever has continued some time, and the body is passing through that cooling process. The following case will illustrate this position. In the summer of 1791, a boy of eight years of age, in D 4 whom

whom I am peculiarly interested, was attacked by fever. On the third day his pulse rose to 130, and 140 in the minute, and his heat to 106° and 107° of Faht. His thirst was very great, and delirium commenced on the fecond day, and continued without intermission. Various methods had been employed to abate the fever, and particularly to excite perspiration, but unsuccessfully. His heat was not leffened by repeated spunging of the surface of the body with cold vinegar and water; and after a copious bleeding, all the fymptoms were as alarming as before. It feemed hazardous to repeat this evacuation, as the blood exhibited no fize, and there was a fuspicion that the disease originated in contagion. The patient had taken antimonials without any apparent effect, and after watching the state of the thermometer, with the bulb at the axilla, upwards of an hour, though the mercury had funk a fingle degree in that interval, it stood at the end of the time as high as 106°. In this state of things we refolved on trying the cold affusion, and every thing being prepared, he was stripped naked and lifted out of bed. As we were about to throw the water upon him, it was observed that a perspiration had broken out all over him, but the heat being fo great, we perfifted in our purpose, and four gallons of fresh water of the temperature of 60°, were dashed upon him; the effects were altogether furprifing. On replacing him in bed, the mercury in the thermometer

tribitw.

only, and the burning heat of the extremities was converted into a coolness that was rather alarming; the pulse had sunk in frequency to 90° but was full and steady. Gentle frictions were applied to the legs and seet, but they were not long continued, for the general warmth speedily returned; the heat in the trunk of the body rose in about an hour to 100° and the pulse to 100. His delirium went entirely off; the sur on his tongue speedily disappeared; and twenty hours afterwards he was found free of every complaint but debility.

Subsequent experience has however convinced me, that though in this case the termination was so happy, the cold affusion was not unattended with hazard. Perspiration had commenced, and the heat was finking. It had perhaps funk more at the moment when the affusion was performed, than was indicated by the thermometer, for the bed clothes often keep the body from cooling under perspiration to the degree that would otherwise be-produced. In this case when the surface is suddenly exposed to the external air, the heat finks rapidly. If the perspiration had continued an hour longer before this remedy was used, the heat would have been still more diminished; a torpor of the vessels of the furface, and of the extremities, would have been produced, followed by a great, and probably a dangerous gerous re-action of the centre. This observation will be illustrated in the sequel.

In recommending the affusion of cold water as a remedy in fever, an express exception is therefore made against its use during the feverish chill, or after the perspiration has begun to flow profusely, and more especially after it has continued to flow profusely for some time. An exception is also made against its being employed in the latter end of fever when the strength is much exhausted, and the heat is fometimes as low or lower than the temperature of health. While, however, the heat rifes one or two degrees above the healthy standard, this remedy may be used even in the latter stages of fever. I have employed it with advantage on the 11th, 12th, and 13th days. In inflances of this kind it will however be prudent to make the degree of cold very moderate, as has been already observed; and as it is scarcely to be expected, that at an advanced period of the disease, the progress of it can be stopped, or its duration much leffened, it may perhaps answer every purpose to employ in such cases the tepid ablution. I have indeed often contented myfelf with fpunging the body all over with tepid vinegar, or vinegar and water, from the 9th or 10th day forwards; but I have frequently in cases where the heat continued high, directed the general affusion of tepid water, by which the heat may always be speedily and

and effectually reduced, when that is the only object in view.

Under these restrictions, the affusion of cold water may be used with perfect safety in the low contagious sever of this country, and the sacts already stated, will show that it is a remedy of great power and essicacy. In the first stages of sever, it appears very generally to cut short the disease almost instantaneously; and even when it sails of this essect, as is usually the case when it is applied in the more advanced stages, it nevertheless moderates the violence of the symptoms, and shortens the duration of the sever.

Cammatory fever, without topical inflammanion, in

contribution a very rate occurrence in this illand;

chap.

CHAP. VIII.

water may be used with period- talety, in the loss

contagious fever of the country, and the facts al-

General remarks on fever. History of a case of sever in which the affusion of cold water was not salutary.

advanced flages, it nevertheless moderates the vio-

SINCE the introduction of scientific arrangements into medicine, diseases have been much reduced in number, and their nature has been more clearly understood. This is especially true of continued fever, which is exhibited by Dr. Cullen under three genera only, Synocha, Typhus, and Synochus. Of these genera, however, the Synocha, or pure inflammatory fever, without topical inflammation, is confessedly a very rare occurrence in this island; the venerable professor used to declare that he had not met with a fingle instance of it in forty years practice.—And the typhus and fynochus feem to be confidered by him as the same disease, modified differently, by the difference of climate, feafon, and conftitution. Both are described as contagious, and as occasionally producing each other. Doubtless the typhus, or low contagious fever, is the prevailing

vailing fever of this island, and of Europe. It is the epidemic of all our great towns, of our jails, hospitals, and manufactories; its origin and progress are clearly ascertained, and its symptoms generally understood. It is to this fever that the preceding observations chiefly apply.

I have my doubts, however, after much reflection and observation, whether we have not simplified too fast in our nosological arrangements of fever. The dreadful difease which prevailed lately at Philadelphia, and which now ravages the West Indies, cannot perhaps be included without some violence within our fystems of Nosology; and its fatality under all the established modes of treatment, whilst it excites our deepest regret, must serve to abate the pride of modern science. Even in our own island, it appears to me that cases of fever occasionally occur, which cannot be referred with advantage to any of the genera of Dr. Cullen*. The following is a description of a fever of this kind; I have not met with it often, but when I have met with it, it has very generally proved fatal, under the eftablished

^{*} I am aware that all questions respecting Nosological arrangement have a tendency to degenerate into verbal disputes, and I willingly avoid them, referring for my accuracy to those who have studied diseases, not in books only, but in the volume of nature.

established modes of treatment; and I am forry to say, that in the only instance of this sever in which I have tried it, the affusion of cold water proved unsuccessful also.

The fever in question does not feem to originate in contagion, or to propagate itself by contagion. I have never been able in a fingle instance to trace it to that fource, nor have I ever found it to be communicated from the patient to any of his attendants. The cases which I have seen have occurred chiefly in the winter feason, in persons in the flower or vigour of life, possessed of considerable sensibility of mind, and in the habits of more than ordinary mental exertion. After some days of indistinct catarrhal complaints, the fever comes on (in general after some accidental exposure to cold) with a very violent and long continued attack of chills and rigor. and to this, as is usual, succeeds a state of heat and re-action. The patient complains of intense headach and of oppression at the præcordia, with occasional but not severe cough, and with some increase in the frequency of respiration. His pulse is not remarkable as to frequency or ftrength; his fleep is not particularly diffurbed; and for fome days the complaint goes on as if produced by catarrhal fever. From the first, however, there appears a great quickness and impatience about the patient : he talks more rapidly than usual; apprehends you quickly,

quickly, and answers you instantly. He cannot, however, command his attention long, and is fatigued with the effort. His heat, which was at first moderate, becomes very great on the 7th and 8th day, reaching 107° or 108° of Faht, ; he becomes delirious and talks inceffantly. Throughout the fever, his fenses of hearing and taste are uniformly acute, and this is true also of his sense of feeling. Great as his heat is, he is much alive to the impressions of cold on the surface of the body, and shrinks from them. At times he appears furprifingly calm and natural, gets out of bed and dreffes himfelf, infifting that he is well. Often he ftarts up fuddenly in bed and opens his curtains, feeming to look round the room for some person he fupposes present; and sometimes he rings the bell violently, if within his reach, without apparent object. Indistinct conceptions rife and vanish in his mind, and the impressions of sense are confounded with the ideas of imagination. As the fever advances, the respiration becomes more hurried and laborious, the pulse more frequent and feeble; and towards the latter end of the difease, but not before, perspirations break out, at first partial, and at length general and profuse, which however, though they reduce the heat, do not otherwise relieve him. The pulse finks; the body is covered with petechiæ; wine, bark, opium, and blifters, afford no relief: the patient dies on the twelfth or thirteenth day of fever_

fever, and after death the body runs rapidly into putrefaction.

I have feen this fever treated by venefection and antimonials in the early ftages, with a strict attention to the antiphlogistic regimen; and by bark and cordials, as the strength began to decline; but without fuccels. I have also seen it treated from the first on the same plan as typhus, but with an equally unfortunate iffue. In a cafe of this fever which occurred lately, I made use of the cold affusion, and as the mercury rose in the thermometer with the bulb under the tongue, to 107°, I employed this remedy with fome degree of confidence. The effects did not correspond with my former experience, or with my hopes. The patient felt the cold most acutely, but was not relieved. His pulse did not diminish in frequency; his heat subsided very little, and that for a few minutes only; neither diaphoresis nor sleep followed. This remedy was not repeated, but the furface of the body was spunged from time to time with vinegar, without however producing fenfible benefit or refreshment.

I have already mentioned that the affusion of coldwater is not to be used after a profuse perspiration has taken place; and that it is not to be used in the

cold stage which begins the paroxysm of fever, nor till the hot stage be fairly formed. In the typhus, however, this last restriction seldom requires us to wait long; the affusion may be used in general in twenty-four hours from the original attack, and often much sooner. The case I have just related is the only instance, out of many hundred trials, in which I found, that even on the fixth day of fever, with the actual heat of the body far above the temperature of health, the affusion of cold water was neither falutary nor refreshing. I have however to observe, that notwithstanding the great heat of the body, producing the utmost restlessness and anxiety, the fensation of heat was interrupted by chilliness, on the flightest application of cold, and that the furface and extremities not only felt chilly, but grew cold even on the accession of the external air. In reality, through the greater part of the fever, the state of the patient had a considerable resemblance to what we see in the paroxysim of an intermittent, when the cold stage is terminating, but the hot stage not fairly formed—when the heat as well as the blood, is accumulated in the centre of the fystem, and the vital power is struggling to give them that propulsion to the superficies, which terminates in profuse perspiration, and carries off the disease. I have little doubt that immersion in the tepid bath continued for some time (of the temperature that feels comfortably warm to the skin) is

the proper remedy in the fever I have described, as it doubtless is in the struggle of the paroxysm of intermittent; and when an opportunity offers, I mean to ascertain the truth on this point. But this opinion will be illustrated when we come to speak of the warm bath more particularly.

If any one should contend that the fever I have described is in reality only a variety of the typhus, or fynochus, I shall not be disposed to contest the matter. The question concerning identity, leads to endless disputes in every branch of science where it occurs, and he must know little of nosology, who supposes it has yet received a confistency, that would render fuch a discussion profitable. It is fufficient for me to observe, that the symptoms of the two diseases are in a considerable degree different, though with that general fimilarity that belongs to all cases of fever; that the state of the nervous system as to impressibility is widely different; and what is of most importance, that the methods of treatment, which according to my experience almost invariably succeed in the one disease, are unfuccessful in the other. Every practitioner knows, that in typhus, the fense of hearing is generally obtuse; and the same may be said of the taste, fmell, and touch; whether the observation is applicable to the fight also, and under what restriction, appears to me doubtful. The acuteness of all these fenfes fenses in the fever which I have described, is very remarkable, and particularly in regard to the sensibility of the surface. I have observed this symptom to be produced by several narcotics, and by some poisons. It is very remarkable in the hydrophobia; and in the last days of a person who died of inanition the senses of touch and vision were extremely acute.*

E 2

when the property of the property of the state of the property of

proper herriescent and loss model a bosburi de la

cool air, feethed to point out the essemal me of

tion, might be more purpositely adapted to the

more analysismic forms of malispox, of he refula

CHAP.

^{*} This case will be found in chap. xvii.

-inel salt of braze CHAP. IX. but salds to the

to the produced by fewnal narcotics, and by forne

posione. It is very remarkable in the hydropho-

fentes in the fever which I have described, is very

Of the use of the affusion of cold and tepid water in small-pox, with cases.

THE fingular degree of fuccess, that on the whole attended the affusion of cold water in typhus, encouraged a trial of this remedy in some other febrile diseases. Of these the small-pox seemed more particularly to invite its use. The great advantage that is experienced in this disease by the admission of cool air, seemed to point out the external use of cold water, which being a more powerful application, might be more particularly adapted to the more malignant forms of small-pox. The result corresponded entirely with my expectation. Of a number of cases in which I witnessed the happy effects of the affusion of cold water in small-pox, I shall give the following only.

CASE. I.

In the autumn of 1794, J. J. an American gentleman in the 24th year of his age, and immediately on his landing in Liverpool, was inoculated under

my care; the prevalence of the small-pox rendering it imprudent to wait till the usual preparations could be gone through, or indeed till the fatigues of the voyage could be recovered. He fickened on the Teventh day, and the eruptive fever was very confiderable. He had a rapid and feeble pulse, a fœtid breath, with pain in the head, back, and loins. His heat rose in a few hours to 107°, and his pulse beat 119 times in the minute. I encouraged him to drink largely of cold water and lemonade, and threw three gallons of cold brine over him. He was in a high degree refreshed by it. The eruptive fever abated in every respect-an incipient delirium fubfided, the pulse became flower, the heat was reduced, and tranquil fleep followed. In the course of twenty-four hours the affusion was repeated three or four different times at his own defire; a general direction having been given him to call for it as often as the fymptoms of fever returned. The eruption, though more numerous than is usual from inoculation, was of a favourable kind. There was little or no secondary fever, and he recovered rapidly.

In fituations where the eruptive fever of smallpox is clearly distinguishable, and where it does not abate sufficiently on the admission of cold air, the affusion of cold water may be resorted to with considence and safety, regulated however in this application, as in every other, by the actual state of the patient's heat, and of his sensation of heat. In the confluent small-pox, however, after the eruption is completely formed, this remedy cannot perhaps be used with advantage. The following case will illustrate this position.

best 130 times in the minute. I encouraged him

breach, with onin in the nead, back, and loins.

H. A. aged 23, an American mariner, fell under my care (Dec. 7,) on the third day of the eruption of the small-pox; that is, on the sixth day of the disease. His pulse 114 and seeble, his heat 100°. His head, back, and loins, ached severely—thirst great—skin livid—small-pox confluent.

He was put on a milk diet—gentle mercurial purgatives were ordered from time to time, and an opiate every night at bed-time. Lemonade was given largely, at first by itself, and afterwards mixed with wine, and the affusion of cold water was directed in the usual way. In ten minutes after the affusion, the pulse was 96, the heat 98°; the livor of the skin was much diminished, but the pains were not relieved.

Dec. 8. Noon—Pulse 96, soft and regular—thirst gone—respiration slow and natural—heat 97°.—

The

The affusion was ordered to be repeated; ten minutes after, pulse 84 and feeble—heat 84°.

Dec. 9. Noon—Pulse 88, heat 93°—the cold affusion was not repeated in this very reduced state of heat; the decoction of bark was ordered, and a pint of wine daily in lemonade.

Dec. 10. Noon—Pulse 116, and full—heat 98°, respiration still easy—expectoration considerable and viscid—thirst less—eyes quite closed—head swelled—a complete union of the pustules on the face.—Bark and wine continued, with the opiate at night.

Dec. 12. Pulse 118—heat 96°. A bucket sull of water of the temperature of 92° was poured over him. He appeared refreshed at the moment; ten minutes after, pulse 112, heat 94°. Complained of being chilly. Respiration still easy—free of pains, and his face less swelled. Complained of his throat. A blister was applied to it all round.

Dec. 13. Noon—Pulse 118—heat 96°—respiration still free, but his throat very fore. Medicines were continued, but the affusion of tepid water was not repeated.

Dec. 14. Noon—Pulse 138—heat 100°—respiration is now become laborious, and the expuition E4 difficult.

difficult. The throat was much swelled. He was frequently spunged with tepid water, and the medicines continued.

Dec. 15. Noon—Unfavourable symptoms increased.

Dec. 16. Noon—Vomiting came on, which was relieved by opium. His fenses and his intellect reremained acute to within an hour of his death, which happened at eight o'clock in the evening of this day.

If this case be more detailed than seems necesfary, let this be excused, as it is the first in which the actual heat in confluent small-pox has been recorded. It is here given accurately from the period when the disease came under my care.

In regard to the effects of the cold affusion, it may be observed, that this remedy was not used during the eruptive sever, nor till three days after the eruption had appeared, and the character of the disease was decided. In the stage in which it was employed, the sever and the heat were abating, as is usual after the eruption; and in all cases in which the heat is sinking, the application of cold must be made with great caution, as has already been mentioned.

oned. After the fecond affusion (on the 8th) the heat funk below its natural standard, and continued below it for some time; so that this remedy became inadmiffible. The difease went through its usual course. The tepid affusion on the eighth day of the eruption (Dec. 12) was used in part to wash off variolous matter, and in part to produce refreshment. The heat which was before 96°, funk two degrees, fo that it could not with fafety be continued, for experience has proved, that the tepid affusion is a powerful means of diminishing heat. The heat rose again with the secondary fever, and the patient died of the affection of the throat, as I believe is general in the confluent fmall-pox.* It will be at once perceived, on the principles already laid down, that in a difease like this, the affusion of cold water could only be effentially useful during the eruptive fever. It is during the eruptive fever that the quantity of the affimilation is determined, as well as its kind. This is invariably found to bear an exact proportion to the eruptive fever, and whether we confider the eruptive fever as the cause or effect of the affimilation, there is every reason to expect, from the laws of the living fystem, that the diminution of this fever will diminish the quantity, and meliorate the quality of the variolous eruption. In

^{*} See Zoonomia, vol. ii. page 237.

In the case just related, the heat during the eruptive fever (judging from trials in fimilar fituations) had rifen to 106° or 107°; but it had funk to 100°, before the cold affusion was employed. It may eafily be conceived that this remedy could have been employed to a much greater extent, and that its effects would have been far more falutary, if it had been used throughout the previous fever. That it would have effentially altered the character of the disease, I presume not to affert. This however, I can declare, that in all the cases in which I have used the affusion of cold water during the eruptive fever, however fevere the fymptoms may have been, these symptoms instantly abated, and the disease assumed a benignant form. The case of Mr. Johnston (Case I.) already given, will illustrate this observation; and fix or seven others I might adduce to the same purpose. As yet my experience extends no farther.

The inoculation of infants is so very rarely sollowed by any serious disease, that as far as respects them, the affusion of cold water may be seldom required. The inoculation of adults is not, however, quite so safe. Inoculation is seldom indeed performed in our island on adults that are natives, but foreigners frequently require it; and in Liverpool, our intercourse with America renders it often necessary to perform it on adults from that continent.

nent. We may also observe, that when the natural fmall-pox is epidemic, the eruptive fever will be generally diffinguished; and wherever it is diffinguifhed with fymptoms of violence, instead of trusting to cool air only, the cold affusion, or cold bath, are strongly recommended. To our brethren across the Atlantic this is more especially addressed .-In America, as well as many parts of the old continent, in consequence of the neglect of early inoculation, the natural fmall-pox at times fpreads alarm and devastation throughout extensive districts. In this island the ravages of the natural small-pox are on the whole very great, yet they occasion little disturbance or alarm. The practice of inoculation among the more opulent classes of fociety, keeps up the contagion in all our populous diffricts, and at the same time by relieving these classes from the apprehension of the natural small-pox in their own families, prevents them from oppoling the calual progress of the disease among the inferior orders, who want the knowledge and the combination neceffary to the use of the means of prevention among themselves. Though therefore it is demonstrable, and has indeed been demonstrated, that the destruction of the natural, or rather the casual small-pox, might be entirely avoided, yet, as it falls almost wholly on the families of the poor, and as it has been an evil that has been long, and that is familiarly known, we fubmit to it through habit, as if it were

were inevitable*. The alarm produced by the cafual contagion is therefore feldom fo great in our large towns, as to give a practitioner frequent opportunities

* The observations on the possible prevention of the casual small-pox, will doubtless suggest to the medical reader, the "Sketch of a Plan" for that purpose, published in 1793, by my respectable neighbour, Dr. Haygarth. That this plan is in itself practicable, and that it would be effectual, I have little doubt. Unfortunately, it requires the assistance of government (as I remarked at the time) and this I fear is a powerful objection. It is possible however that some scheme of this kind (including I hope all contagious diseases) may one day or another be attached to some comprehensive plan for the management of the poor.

The theory that fuggested Dr. Haygarth's plan, but which formed no effential part of it, involved him in a discussion on the length of time that variolous matter may be exposed to the atmosphere, and retain its infectious quality. On this occafion, as there was fome difference of opinion between us, he proposed several experiments to me which would doubtless (See pages 459, 460, of " A have decided the question. Sketch, &c.") It was fully my intention to have undertaken some experiments such as he mentioned, and I even commenced them, but as my attention was forcibly drawn to other fubjects, these experiments, which required extreme accuracy, were not completed, and my engagements have never fince permitted me to recommence them. I have, however, fince that time, inoculated with matter at different periods from its being taken from the patient, and the refult is as follows .-The length of time which variolous matter exposed to the air retains its contagious quality, depends on its superficies. If it fluent small-pox, the only stage of that disease in which medical treatment is likely to be of much avail. Where such an opportunity does occur to the enlightened reader of these pages, it is hoped that the

be fpread very thin on a piece of flat or convex glass, it loses this quality much fooner than when it is collected in a mass, Spread thin upon glass, it sometimes disappoints the inoculator at the end of twenty days, though not generally; and I have known it fucceed in communicating the difeafe, even when diffused over a large surface, at the end of seventy-three days. (See Dr. Haygarth's Sketch, page 447.) But I find from the experience of others, as well as from my own, that this is not common. The instance in which I found variolous matter to retain its infectious quality longest, was the following. On the fecond of February, 1792, I took a confiderable quantity of this matter on a piece of window glass, keeping it as much together as its fluidity would admit. It was exposed immediately to a stream of air, and the surface was speedily dry. On the fecond of March following, after moistening a portion of it with a little water, I inoculated three patients, and all with fuccefs. In the course of the summer I inoculated with another portion of it, previously liquified by the addition of warm water, and with fuccess as before. On the twentieth of July, 1793, I again used a portion of the same matter in the same way, and again with success; but this success was long doubtful, and it was not till the twenty-second day after the operation, that the patient fickened .- I used the same matter in June, 1794, when it entirely failed me. It is now by me and is not mouldy, nor any ways changed in its appearance. Variolous matter kept fome time is certainly flower in producing the disease, even where it does succeed in the end.

the affusion of cold water, or the cold bath, will not be neglected. The Chinese, it is said, have long sollowed this practice with extraordinary success*.

The use of the cold affusion in sever was common among the physicians of the infirmary, (Dr. Brandreth, Dr. Gerard, and myself) while severs were received into that hospital, and its extraordinary success there, has rendered it in Liverpool familiar in private practice, and extended it, as I have already mentioned, to several other sebrile disorders. The

* Sir William Watson, in his tract, entitled "Account of Experiments on the most successful methods of inoculating the small-pox," published in 1761, mentions the case of a young woman, who in the absence of her nurse, got out of bed delirious during the cruptive sever of small-pox, and threw herself into the new river near Islington. She was discovered stoating on her sace; and when taken out of the water, had not the least appearance of life. She was recovered however by the usual methods, and afterwards passed well through the disease.

In some parts of Bengal, according to Mr. Ives, inoculation is practiced among the natives—After the operation is performed, the patient is ordered to bathe in cold water thrice a day, and to live on the most cooling diet. When the fever comes on, the bathing is left off, but it is resumed on the second day after the eruption appears, and continued for the three successive days. It is supposed to fill the pustules.

Ives' Voyage to India, in the years 1755, 6, 7, ch. iv. p. 54.

icarlatina anginosa has for many years prevailed in Liverpool, and though in general mild, has as at particular seasons been malignant and fatal. To this disease, the affusion of cold water has lately been applied by my friend Dr. Gerard, and the particulars that follow are mentioned on his authority.

In the latter end of December 1796, all the children of a family in his neighbourhood, five in number, had been attacked in fuccession with scarlet fever; four of these were recovering, but one was dangeroufly ill, when the father of the family, with whom one of the children had flept, was himself seized with all the symptoms of the disease. He had excessive pain in his head and back, and flying pains all over him. He had frequent rigors, loss of appetite, and fickness, with some flushing of the face, but without any efflorescence on the skin or affection of the throat. This was his fituation when Dr. Gerard was called in, about fixteen hours after the first attack. An emetic, and afterwards a cathartic were ordered, but their operation was flow and imperfect, and on vifiting him, ten or twelve hours afterwards, he was not materially relieved.

Entertaining no fort of doubt of the nature of the attack, these symptoms foreboded that the epidemic would, in this instance, be severe; and Dr. Gerard determined to try the affusion of cold water, from from which in typhus he had seen such happy effects. Accordingly the operation was performed, and with a result that far exceeded his hopes. As he was much debilitated, half a pint of hot wine was given him after it, and on being put to bed, the symptoms of sever were found nearly gone; a genial warmth diffused itself over the extremities of his body, followed by perspiration and sleep. Next day he complained of a slight degree of head-ach and lassitude; Dr. Gerard therefore ordered the affusion to be repeated, as well as the warm wine after it; the symptoms of the disease vanished, and never re-appeared.

A day or two afterwards, a maid who had been hired as an affiftant to attend the fick children, and who had been about a week in the house, was attacked by the precise symptoms already related, and which had uniformly ushered in the epidemic. She took an emetic on the first attack with little benefit, and soon after the cold water was poured over her, the wine being administered after it. In this case the remedies were used earlier than in the former one;—they were used once only: the febrile paroxysm was dissolved, and never returned.

The result of these cases communicated by Dr. Gerard, leads to a variety of important reslections. That the affusion of cold water extinguishes the incipient scarlatina as well as the typhus, can scarce-

ly be doubted; and thus this powerful and simple remedy is extended to another, and a most important class of diseases. That the disease was extinguished without the specific efflorescence of the skin, or affection of the throat, is a circumstance not a little It feems to demonstrate that this efflorescent matter is the product of the eruptive sever; and that the fever itself being destroyed in the first instance, the efflorescent matter is never produced. Thus we are freed from the apprehensions which a false theory might suggest against extinguishing a process by which nature was extricating itself from an acrimony which the fystem had imbibed .- Thus also our conclusion is supported, that the eruptive fever of small-pox is the cause, and not, as some have supposed, the consequence of the progress of assimilation, and that the diminution of this fever by cool air, and still more by the affusion of cold water, actually diminishes the quantity of matter affimilated, and in certain cases might perhaps wholly prevent the affimilation.* This last conclusion is indeed

^{*} Dr. Cullen has assumed, that in all the various degrees of small pox, the quantity of matter assimilated bears an exact proportion to the bulk of the body, and that the difference in the nature and quantity of the eruption depends wholly on the permeability of the skin. This doctrine, which in his lectures he extended to all the other exanthemata, is, in my judgment, one of the weakest parts of his most valuable work.

indeed doubtful, as all analogical inductions must be between different diseases; and particularly where the difference is of this important kind, that the constitution in one case is susceptible of the disease once only, while in the other it may probably receive it indefinitely. We may also observe, that the prevention of the assimilation of small-pox, by wholly extinguishing the eruptive sever, if it were in our power, would not be advisable, since it must leave the patient exposed to the future influence of that contagion.

I have not had an opportunity of repeating Dr. Gerard's practice in the incipient stage of scarlatina, but after the essociation on the surface decides the nature of the attack, I have for the last sisteen months uniformly prescribed immersion in the tepid bath, (from 92° to 96°) and with striking benefit. Whether the assuring of cold water is applicable to the other exanthemata, must be left to surure experience.

In the cases that I have related to illustrate the effects of the cold affusion, the temperature of the water may be judged of from the season of the year. In general it was from 40° to 50° of Faht. In the epidemic which prevailed in the 30th regiment, the water of the river was employed, as has already been mentioned, which, as the season was uncommonly

uncommonly cold, did not, though in the month of June, exceed 58° or 60°. I have, however, very often used the river water in private practice during the fummer months of the last four years, when in general it has been from 65° to 70°, and the effects correspond with those already described. The folution of fever, or to speak in language more precise, but as yet less intelligible, of the morbid febrile catenation, depends on the fudden, general, and powerful impression on the fensations, and this impression is less affected by the difference in the temperature from 40°. to 65°. as far as my observation extends, than might on a first confideration be imagined. Within these limits the efficacy of this remedy, as well perhaps as its fafety, depends on the fuddenness and momentariness of its application. The powerful impression on the fenfations is much weakened when the water is poured flowly on the body, and as the respiration is suspended or convulsed during this application, as well as during the act of immersion in the cold bath, it might in some cases incur hazard to protract it.

F 2 CHAP.

chap. X. a brown and I chap. X. a brown and in

when in general it has been from 65° to 70°,

and the effects correspond with those already

delerabed. The foliation of fever, or to freak in

magan sala-at tiesout god bib thios glacommonau

Of the affusion of tepid water on the surface of the body, in feverish disorders, and of spunging the body with water or vinegar. The affusion of tepid water practised by the ancients.

far as my oblevation extends, than might on a first

I APPLY the term tepid to water heated to that degree which is warm, but not hot to the fenfations, and which in the way of affusion is from 87° to 97° of the scale of Fahr. According to my experience this term, when the body is immerfed, may be applied to water some degrees colder, the reason of which will be easily understood by those who reflect, that under immersion no evaporation from the furface of the body takes place. At first I imagined that the tepid affusion might be beneficial in cases where the heat of the body is below the degree necessary to render the cold affusion safe. I employed it therefore in those stages of fever where the heat did not exceed the temperature of health. A little experience however convinced me that this practice

practice was not without hazard, for I found, that in many cases, at least, the heat of the living body is lowered as speedily by the affusion of tepid water, as by the affusion of water that is cold :- If I mistake not, in some cases the heat is lowered more speedily by the tepid water. To those who reason respecting the heating and cooling of the living body in the fame manner as respecting inanimate matter, this observation will appear paradoxical; I affert it however from actual observation, and a little reflection will explain the phenomenon. The evaporation from the furface is more copious from the tepid affusion, and on this the cooling of the body very much depends. But this is not all; the tepid affufion is little if at all stimulating, and does not, like the cold affusion, rouse the system to those actions by which heat is evolved, and the effects of external cold are refifted. Where the object is to diminish heat, that may be obtained with great certainty by the repeated use of the tepid affusion, suffering the furface of the body to be exposed in the interval to the external air-and if the beams of the fun are excluded, and a ftream of wind blows over it, the heat may thus be reduced where cold water cannot be procured; even in the warmest regions of the earth-on the plains of Bengal, or the fands of Arabia. I have accordingly employed the tepid affusion very generally in those feverish affections where the morbid actions are weakly catenated, de-

pending

pending rather on the stimulus of preternatural heat, than on contagion, miasmata, the morbid contents of the stomach and bowels, or local inflammatory affection. Of this kind are a great part of the feverish affections of children, in which the tepid affusion is a valuable remedy. It very generally produces a confiderable diminution of heat, a diminished frequency of the pulse and respiration, and a tendency to repose and sleep. I have used it also in feverish diforders of various kinds where the lungs are oppreffed, and the respiration laborious, and where of course the oppression might be dangerously augmented, by the fudden stimulus of the cold affusion. is also applicable to every case of fever in which the cold affusion is recommended, and those may receive much benefit from it, whose fears or whose feebleness deter them from that energetic remedy. have not however found its effects fo permanent as those of the cold affusion, and I have never seen it followed by the total ceffation of regular fever, as often occurs after the cold affusion. In the hectic fever, however, where the actions are less strongly catenated than in fynochus or typhus, the paroxyfm is fometimes completely extinguished by the affufion of tepid water* in the commencement of the hot

* See this fact mentioned in the Zoonomia, vol. ii. p. 296, where a relation is inferted by Dr. Darwin, of the author's (Dr. C's.) own case of hereditary consumption.

hot stage. In the hectic paroxysm, the heat seldom rifes more than two degrees above the temperature of health in the trunk of the body, and three or four degrees on the extremities. By moistening the palms of the hands and the foles of the feet with vinegar, its effects may be moderated, for it is from the fensation of heat in the extremities that the stimulus to the fystem is chiefly derived; and this practice ought not to be neglected, if the tepid affusion is not employed generally. In all cases of fever indeed, where the burning heat of the palms of the hands and foles of the feet is present, this method of cooling them should be reforted to; it is uniformly fafe and refreshing. I have not employed the cold affuson in the hectic paroxysm. This disease generally adheres to a debile fystem; the body parts with its heat in it easily; and the lungs being always affected in the pulmonary hectic, the fudden application of cold to the furface might produce unpleasant, and perhaps dangerous effects on the respiration. Neither have I tried it in peripneumony, or meazles.

When the affusion of water cold or tepid is not employed in fever, benefit may be derived, as has already been mentioned, though in an inferior degree, by spunging or wetting the body with cold or warm vinegar or water. This application is however to be regulated like the others, by the actual state of the patient's heat and of his sensations.

F 4

According

According to my experience, it is not only less effectual, but in many cases less safe; for the system will often bear a sudden, a general, and a stimulating application of cold, when it shrinks from its slow and successive application*.

I have

(*) I have purposely avoided entering on the general operation of cold, in hopes of being better prepared for the fubject at some future opportunity. In the mean time I have continued the use of such terms as are most intelligible, and most consonant to the true doctrines as far as I perceive them. To speak of cold in any form acting as a stimulus, seems however to fome learned friends, not merely an error, but an abfolute contradiction in terms. Heat they affirm to be the univerfal stimulus, and cold being merely the privation of heat, must in their opinion, always have a fedative operation. As well, it is faid, may darkness, which is the privation of light, stimulate the eye, as cold, which is the privation of heat, stimulate the general fystem. This observation, which has more importance with me from the quarter whence it comes, than from any intrinsic weight it possesses, leads me into one or two general remarks.

That cold can never act as a stimulus, was a favourite dogma of the late Dr. Brown. It was a necessary consequence of his general doctrines of life. As he admitted only of two classes of diseases, diseases of increased and diminished excitement; so he reduced all remedies into two corresponding classes, such as diminish, and such as increase this excitement. That cold in extreme degrees is a powerful and effectual sedative, is incontrovertible: he therefore held that it is sedative in every degree, without being at the trouble to point out the line of temperature below which the term of cold is applica-

I have also used the affusion of cool water as a remedy in sebrile diseases, but more frequently in paralysis, and in other diseases of debility. By the term

ble. It was the character of Dr. Brown to follow his hypothesis into all its confequences, contemning all facts that rested on the testimony of others, and neglecting more than perhaps any man of his talents ever did, to bring his opinions to the test of experiment. His theory was in his hands the bed of Procrustes. On the operation of cold he was fingularly erroneous. He afferts that cold can never be of fervice in the fevers of the torrid zone, (Elements of Med. vol. i. p. 23.) and the reason he gives is, that in those diseases there is indirect debility, to which cold must not be applied. He declares that it never can be of service but in sthenic (inflammatory) difeases, (vol. i. p. 103.) that for these it is the grand remedy; (p.296) that its operation is the same as in small-pox, in meazles, catarrh, and all other inflammatory difeases; and that " it is " fufficient to cure them all!" These affertions are not merely erroneous, they are dangerous.

Whatever opinion may be entertained of Dr. Brown's fundamental principle—that all the phenomena of life are to be explained, by affuming that the living principle (the exciteability) is accumulated and expended in the inverse ratio of the stimulation, it will not now be disputed, that he was acquainted with only one mode of action of the living principle, that which Dr. Darwin has described under the name of irritation; and that he was wholly regardless of the influence of sensation, volition, and association, since his time so fully illustrated in Zoonomia, and which on every hypothesis must be allowed to have such vast influence on the motions of life. It was to his difregarding the action of cold on the sensation.

term cool, I indicate the temperature from 87° to 75°. It operates as a gentle stimulant, and may be used as a milder form of the cold affusion. Like the

fations, that the error of Dr. Brown is more immediately to be traced.

It is not so easy to explain why Dr. Darwin, who understood the laws of sensation so well, should also have difregarded the influence of cold on this fenforial power. He feems to confider the application of cold as merely fubtracting the stimulus of heat, and thus to be altogether negative; (Zoonomia, vol. ii. p. 757.) without adverting to that most powerful senfation, by which great and fudden changes from a higher to a lower temperature are attended, through which the energies of life are roufed into action, and the fedative influence of cold for a time counteracted. Whatever theory we adopt in regard to the origin of caloric in the living body, it feems to me to be in a great measure owing to the influence of the changes of the external temperature on the fensations, that the uniformity of the heat of the animal is preserved; and this view of the subject appears perfectly consistent with the general doctrines of Dr. Darwin.

The ftimulating action of cold, though short in duration, is powerful in degree. In the torpor of convulsion, when weaker stimuli are unperceived, the affusion of cold water on the naked body will often excite the dormant sensibility, and introduce a new action throughout the nervous system. In the apoplectic state brought on by the sumes of charcoal, this remedy is of all others most efficacious; when dogs are suffocated in the vapour of the Grotto del Cani, it is well known that they are recovered by plunging them in the adjoining

the cold affusion its application should be sudden and momentary, when the object is to increase the tone of the system, or to dissolve a morbid catenation; where it is employed to moderate inordinate heat, it may be used more slowly, provided it does not interrupt the catenation on which respiration depends*.

The practice of giving cold water as a drink in fevers

lake. And in other animals, when the last motions of life are apparently over, the same application will sometimes, as I have observed, renew the contraction of those sibres that feemed before fixed in death.—This observation I hope to illustrate at some future opportunity.

Can darkness stimulate the eye? No—Darkness produces no sensation. Can cold be rendered stimulating? Surely—After what has been observed no one will deny it: unless indeed it should be said, that it is not the cold that stimulates, but the sensation which the cold produces; a point that it would be a waste of time to dispute. It is the stimulant power of cold that renders it so difficult to employ it in instammatory diseases. (See note at the end of chap. xv.)

(*) I borrow this term (catenation) from Dr. Darwin.—
It is employed to express a number of actions linked together, nearly in the sense of association: the sympathies, as they are usually called, are included under this term, and many associated actions to which the word sympathy has not been applied, as the connexion between the heart and the lungs, the stomach and the heart, &c. (See Zoonomia, vol. i. sect. xvii.)

fevers was common among the ancients, and immersion in cold water they occasionally employed, but the affusion of it on the surface of the body seems to have been wholly unknown. Ablution of the surface with cold water was first practised in modern times at Breslaw, in Silesia, as appears from a differtation by J. G. de Hahn, under the title of Epidemia verna quae Wratislaviam, anno 1737, afflixit, to be found in the Appendix to the AEta Germanica, vol. x.* While, however, the laws by which the affusion

^(*) De Hahn fays that he is the better able to describe this difease, as he himself was seized by it; and was cured by " peregrina illa multis vifa medendi methodus." He begins with an account of the causes of the epidemic. That first mentioned is the ftate of the weather. In the Month of May 1736, after a very promifing fpring, the weather became wet and cold, and the fall of rain was so considerable during June, July, and part of August, as to lay a great part of Silesia under water. The wind blew chiefly from the North-West .-The harvest was almost entirely destroyed. In consequence a famine prevailed throughout the province, of which the author gives a dreadful account, and this he mentions as the fecond cause. The third cause was the vitiated air. The stagnation of the waters in the low grounds in the neighbourhood of Breslaw, filled the atmosphere with marsh effluvia. Even the cattle fuffered feverely from this state of the air, and from the want of pure water to drink. The epidemic began in the enfuing fpring. He gives a number of cases. The first is that of a farmer in the neighbourhood of the city, of 40 years of age. "Comites febris erant fubitus virium lapfus, capitis & præcordiorum dolor gravis, fluxio alvi, pervigilium, inquies,

affusion of cold water ought to be regulated were not understood, the use of the remedy must have been

dilirium. Quæ infuperabilia morbi fymptomata excipiebat die undecima his ipsis insuperabilior obitus." The second case which he mentions occurred in the city. It is that of a delicate woman, of 30 years of age, previously afflicted by fevere misfortunes. The fymptoms were as follows. "Subito elanguida febriebat. Dolor capitis illico atrox. Sitis & Secundo die importuni menses. Exin vomitus bilis, alvusque biliofa. Sputa viscida. Syncopticus rigor. Ardor partium internus. Linguæ quafi candenti ferro compressæ, ficca glabrities & restrictio. Vox arentibus faucibus nulla. Angina fine tumore pessima. Repetebant interdum tenacis petuitæ vomitus alvique fluxio. Os fine medela ficcum. Iactatio. Supinus torpor. Mors inter convulsiones gemibunda." The author proceeds to detail four other cases similar to the above, which also terminated fatally. Two females who caught the infection, escaped in confequence of critical eruptions.

At length the alarm became general. The attention of the magistrates was roused, and from a strict inquiry it was found, that more than double the usual number of deaths had occurred at Breslaw in that year. All the usual methods of practice were of no avail. Whether bleeding was employed or abstained from, the disease was equally mortal. In this distress Dr. De Hahn determined to try the effects of the external use of cold water. The first case in which it was employed, occurred in the month of April; it is related as follows. "Mercator xxxii, Annos natus, floridus natura, sed intermoerores suscus in orthostadio, mox febriens, capitisque dolore & nausca affectus. Permittentibus id viribus, altera die mittebamus sanguinem, qui instammatorius. Vespere vomebat aeger, tertia die maculosus. Maculae morbillis aemulae

been extremely hazardous, and the fatal confequences of its improper application, we can eafily believe

to

inconstantes. Mador. Mox præcordiorum intollerabilis dolor & dilirium. Quafi ex temulentia vacillans extra lectum vagabatur aeger. Oculo altero dilatato, altero connivente, utroque gramioso & culigenoso; lingua lævi, arida, candente; fputo refinofo; urina biliofa."-The ufual remedies failing, on the eighth day they had recourse to ablution. "Tum ad externas illas humectationes confugiebamus, indifessa opera spongiis omnem corporis ambitum demulcentes. Id confequamur, ut sensim sputa resinosa fierent magis mobilia; ut flaccida et sufca facies conspiceretur magis turgidula; ut felicius procederet blandus mador; ut refipiscerit, hactenus vel loquax nimium, vel taciturnus ex dilirio æger." The author then relates a cafe of a fimilar nature, in which, the ablutions being rejected, the patient died. Next follows the case of a lady, a relation of his own, 43 years of age, whose symptoms appeared of the most dangerous nature. In this case the attack was sudden. "Inter domesticas occupationes quasi conquassata concidebat. I. Facies primo momento Hippocratica; artubus glacie perfusis & tremulis. Horror repetitus ad quemvis aeris attactum. Vomitus post quævis ingesta. Abdominis turgor. Videbatur mihe advolanti non morbo corripi ægra, fed fiderata mori, vixque amplius cum morte colluctari. Pharmaco, ob vomitus, non erat locus; neque venæ sectioni, ob algida & emarcida membra. II. Jugi abluitione artuum glacies diffluebat inter gemibundos angores. Sitis tandem invitabat potum. Calorem excipiebat æstus. Clysmi solubantur partes abdominis internas. Quarta die magis fibi confcia ægra caput quafi a percussione dolere conquerebatur. * * * Octava demum die certa redituræ fanitatis spes duorum tandem mensium confirmata."

to have prevented its gaining any ground on the continent, or having been adopted in Britain. These laws

The author next relates the case of a man of 64, whom he attended, with two other phyficians, and on whom the affusion of cold water, or ablution of any kind, does not feem to have been tried. This person died on the 10th day, and our author himself was immediately after seized with the difease, having probably caught it of his patient. He was in his 44th year. "Die 1mo, Perfenseram inter vix superabilos labores, infestum aliquamdiu nuchæ dolorem. Hic ifto mane folito ferventior, febrem epidimicam invitabat absque prævio horroris fenfu. Sœvus ille dolor ex nucha ortum ducens-totum tunc ambiebat caput. * * * Fervebat caput, frigentibus pedibus, spasmo ad abdomen rigidis. Angescente de momento in momentum dolore, horror aderat ad quemvis aeris attactum. Languor mox ab initio extremus. Nox inquieta & sudans. 2do, Oculi gravissime dolentes * * * 3tio, Parca mane remissio dolorum, mox intenfiorum; febre, prævio tempore, aucta. Nox clamoso nuchae & capitis cruciatu atrox. 4to, Post hanc dies nocte pejor. Pedum glacies nondum egelata. 5to, Tristibus his fymptomatibus pingebatur pavoris mentifque debilitas fuspecta; 6to, manibus hoc biduo vibicatis, tendinumque subsultu tremulis, milio per dorsum sparso; 7mo, vomitus præterea, aptharum ventriculi testis. Ab hac die abluitio frigida univerfalis in ufum vocata cum antea particularis tantum obtinisset. 8vo, Octavo pulsus tremulus observatur. Gemitus a dolorum ferocia perennis. 9no, vomitus grumum cruoris emoliebatur. 10mo, Obtinebat tunc rerum ab aliquo mentis deliro stupore incuria. 11mo, sudor, armistitium aliquot horarum producens, opportuno corticis usui locum concedebat, cum jam exitialis languor quævis pessima minaretur. Loquela difficilis & balbutiens.

laws are now, I hope, ascertained by ample experience, and practitioners will, I trust, find themselves directed

Stridor dentium. 12mo, Angina apthofa. donicus. Spafmi cynici. * * * Id interim confequabamur corticis ufu, ut quotidiana exacerbatio tardius, & die decima quarta, quæ tota comatosa fuerat, sub noctem demum ingruerit. Sed gravitate tarditatem compensabat, violenti frigore totum corpus quatienti. Mox algidus aderat fudor, deficienti loquela; demiffa citra voluntatis imperium urina." From this apparently desperate state De Hahn however recovered, and chiefly, as he apprehends, by the ablutions of cold water, which were employed even in this advanced stage of the fever. Almost from the beginning of the fever he indulged freely, according to the plan he had followed with others, in liquids, confifting chiefly of water rendered grateful and cooling by the addition of lemon juice, nitre, &c. and every night he took a moderate dose of opium, from which he found the greatest benefit. From all heating alexipharmics he entirely abstained, having the greatest repugnance towards them. On the 7th day, the ablutions with cold water were begun. The method employed feems to have been that of fponging the furface, and this appears to have been continued for fome time together. The expression is neither elegant nor precise. "Jugiter ergo spongiis abluebar." The effects are described as follows. "Sancteque testor, nunquam non refici ad breve temporis momentum languentes marcidae cutis fibras me perfensisse. Succedebat, repurgatis hac ope poris, perspiratio liberior, ac sudor modestus; videbantque cutis obturacula promptius spongia remota, quam pharmaci interni virtute. Recreabat præterea hiantia & purum aerem fitientia cutis ora, liberior aeris accessus." This account of the effects of the practice, corresponds with our experience at present. Dr. De Hahn found the advantage of a plentiful supdirected in fafety in the future use of this powerful remedy.

G

The

ply of fresh air, at that time little understood throughout Germany or Europe, and he fecured it by keeping his windows open. He changed his bed-clothes and linen frequently; and he occupied one bed during the day, and another during the night, a practice which moderated the heat, and often produced great refreshment. This practice deserves indeed to be followed whenever it can be done with convenience. Experience convinces me that it always produces much comfort under feverish heat and irritation, and that it often procures tranquil fleep. Where a change of bed cannot be obtained, the method of Dr. Franklin, of exposing the body to fresh air, and ventilating the bed-clothes, ought to be reforted to. De Hahn took the bark on the 11th day, and found it very grateful to his stomach, and agreeable to his sensations. That day he took five drachms of the powder. On the 12th, he took four drachms; on the 13th, three only; and on the 14th, only one; his relish for it having gone off. The bark feemed to moderate his fymptoms, which returned with violence on leaving it off, and on the 15th day he found himfelf on the point of death. At this very time however he perfifted in the cold ablutions, and to them he afcribes his recovery. Juges illae abluitiones, quibus hujufque recreatus fueram, ea ipsa die, qua mori videbar, non negligebantur; licet enim totus algidus algido fudore perfunderer, non secus ac liquefactae glaciei immerfus; frigida tamen abluebar; maritum, fervatura chariffima conjuge lethales madores ocius diluente. Morbi tandem reliquiæ menstrua equitatione feliciter subactae. De Hahn was attended by his father, also a physician, under whose direction the ablutions were performed, and by whom the greater part of the fymptoms must have been recorded. Infinenced by his example, the same remedy was employed

The affusion of tepid water in febrile diseases was not unknown to the ancients, though seldom employed

with others, and many were faved by it in circumstances apparently the most desperate.

It is evident from this abstract, that De Hahn was not regulated in his use of external ablution with cold water, by rules fimilar to those which I have ventured to lay down from feveral years experience. Inftead of pouring the water over the naked body, he applied fponges foaked in cold water to every part of the furface in fuccession, and feems to have continued the application for fome time together; in my judgment the least efficacious, as well as the most hazardous manner of using the remedy. He does not feem in general to have used the ablutions till the eighth or ninth day of fever, and till the cases were growing desperate from the failure of other means. At this advanced ftage, the ablutions, as might be expected, feem to have been of very inferior efficacy. Yet in the fingle case, in which from the impossibility of the patient's fwallowing medicines, they were used on the fecond day of the fever, the recovery was speedy; it appeared certain on the eighth day; and this might have encouraged an earlier trial of the same practice in other instances. But what appears to me most furprifing is, that he does not feem to have been regulated in the use of this remedy, either by the actual heat of the patient or his fenfations of heat. In his own case he expressly declares, that the cold ablutions were used on the 15th day of the fever, when he was shivering with cold, and covered with cold fweat; circumftances under which I fhould pronounce it to be in the most extreme degree dangerous. Whether my refluctions as to the use of the cold affusion, and the application of cold in general to the body, be too fevere, future observations must decide; but from a general review of the

employed by them*. It is I apprehend new in modern practice, and will be found an important addition to the list of our remedies. The effects differ considerably from immersion in the tepid bath, as will be more fully explained in speaking of that remedy. Though the affusion of tepid water was little in use as a remedy, it was familiarly practised as a part of their daily regimen, by the Greeks and G 2 Assatics

incautious practice of De Hahn, I am not surprised that his boasted remedy, is, so far as I can learn, no longer in use, either in Silesia or in any part of Germany. The means of making this abstract have been surnished me by a friend at Edinburgh, who copied for me the greater part of the memoir of Dr. De Hahn, from the 10th vol. of the Asta Germanica, in the College Library; in my first edition called Asta Nat. Cur. after Dr. Cullen.

^{*} Some traces of its use in sever are to be found in Celsus. "Quidam ex antiquioribus medicis, Cleophantus, in hoc ge"nere morborum, (tertian intermittents) multo ante accessio"nem caput ægri multa calida aqua perfundebat, deinde vinum
dabat. Quod, quamvis pleraque ejus viri præcepta secutus
"est Asclepiades, reste tamen præteriit, est enim anceps.
"Celsus, lib. iii. cap. xiv." This use of the tepid or warm
affusion is altogether different from what I propose, and was
not only, as Celsus supposes, a doubtful, but probably an injurious practice. For by chilling the body it would dispose it
to the accession of the paroxysm. The use of the tepid affusion
as now proposed, may fairly be considered as a new practice.

Asiatics in the earlier periods of their history. That the Greeks in later times, and after them the Romans, immerfed themselves in the baths to which they were fo much devoted, is certain; but in the heroic age of Greece, the affusion of tepid water seems to have been the mode in which this luxury was enjoyed. Abundant proofs of this are to be found in Homer, particularly in the Odyffey, that admirable record of ancient manners. Venus, after her public difgrace, is described as flying to the groves of Paphos, were she is laved by the Graces, and the improvement derived to her beauty is recorded in a strain of poetry worthy of the subject. (Odys. lib. viii. 1. 362, to 367.) Ulysses is represented as laved by attendant nymphs in the court of Alcinoiis, and in the island of Circe the whole process of heating the water, and pouring it over the naked hero, is particularly described.* On this occasion also, nymphs administered to him, who after the ablution anointed him all over with oil; a fervice well calculated in every respect to increase the pleasure, and to diminish the risque of the tepid

The Romans under their Emperors, carried the fystem

^{*} Odys. lib. x. l. 358, &c. compare this with the bathing of Telemachus in lib. xvii. l. 85, &c.

fystem of bathing to a height of luxury and expense which it never reached in Greece or Asia, as the ruins of their magnificient Balnea prove to this day; and the affusion of warm water was one of the methods by which they diversified this favourite gratification.*

fythems at large. Among the ancients the internal

G 3 CHAP.

^{*} See Hieronymi Mercurialis de Arte Gymnast: lib. i. p. 44 and 45.

Table of the CHAP. XI.

and the affition of warm water was one of the

methods by which they diversified this favoure

Often of bathing to a height of luxury and expende

Of the internal use of cold water in fever.

BEFORE I conclude the account of my experience of the use of cold water in fever, it will be necesfary for me to fay fomething of its effects when fwallowed, on the ftomach, and through it on the fystem at large. Among the ancients the internal use of cold water in ardent fevers is recommended by Hippocrates, Galen, Celfus, and most of the celebrated physicians whose works have come down to us: among the moderns, that extraordinary man, Cardanus, wrote a differtation in its favour; and to pass over a multitude of inferior names, Hoffman, though with fome restrictions, recommends it, not in fever only, but in various other difeafes*. In our own country it was proposed as an almost univerfal remedy by Smith, and a treatife has been written on it under the title of Febrifugum Magnum,

^{*} Hoffmanni Opera, vol. i. p. 479.

by Dr. Hancock. In Spain and in Italy the use of cold water in severs, obtained in the beginning of this century, a greater and more general reputation than in any of the other countries in Europe, and at one time, seems to have superseded all other diet, as well as medicine. This treatment was celebrated under the title of Dieta Aquea, and an account of it may be found in the 36th volume of the Philosophical Transactions, by Dr. Cyrillus, a professor at Naples, to which the reader is referred. Besides the internal use of cold water, he mentions the advantage of laying powdered ice or snow on the bodies of the sick.

Nevertheless, the propriety of giving cold water in fevers, has been disputed by men of high character, and particularly by the celebrated Boerhaave. His doctrine, that a lentor in the blood is the cause of fever, led him to insist on the use of warm drink, and the danger of cold; and his commentator Van Sweiten, though he allows cold drink in some instances, yet in general argues against it*. These learned theorists prevailed in their day over the voice of nature, and the precepts of Hippocrates and Hossman.—In the writings of Pringle, Cleghorn, and Lind, we find little or nothing on the subject, G4 though

^{*} See Boerhaavii Aphorism. Sect. 743- with the commentary of Van Sweiten.

though they wrote expressly on fevers; Dr. Cullen mentions cold drink, but gives no opinion on the propriety of its being used, and certainly did not recommend it in practice. He was even doubtful of the extent to which cold air might be admitted*. On the whole it may be afferted, that the use of cold drink in fever is contrary to modern practice, and that where it is occasionally given, it is administered with caution, and rather permitted than enjoined.

It is not however to the doctrine and precepts of Boerhaave alone, that the difuse of cold drink in fevers is to be imputed. The propriety of giving it freely has been at all times controverted, not on the ground of theory only, but from experience of the dangerous, and sometimes suddenly fatal effects of large draughts of cold liquids, various instances of which have been recorded from the earliest periods of medical history. While therefore some physicians have prohibited the use of cold drink in fevers altogether, those who have recommended it from experience of its salutary effects, have introduced various cautions as to its exhibition, sounded on certain theories, generally fallacious, on the manner in which its deleterious influence is produced.

To

^{*} See his First Lines-cure of fever.

To detail the various opinions that have prevailed on this curious and important point, would be to add another chapter to the ample records of human errors. That the danger arising from cold drink depended on the great difference between the temperature of the liquid and of the body, and that it is therefore to be prohibited when the heat of the body is very great, is an opinion very generally received by the moderns; and among the ancients, though their doctrines were less erroneous, yet while the means of ascertaining the real heat of the living body, and the changes it undergoes, were unknown, it cannot be expected that they should have arrived at the truth on this important subject.

The effects of cold water as a drink in fevers I was naturally led to examine by my experience of its effects as an external application. I have made this examination with the thermometer in my hand, and with all the attention in my power; and the following refults, which will fave the reader the fatigue of reading the particulars of various cases and experiments, seem to me to contain all the information necessary to direct our practice.

1. Cold water is not to be used as a drink in the cold stage of the paroxysm of sever, however urgent the thirst. Taken at such times it increases the chilliness and torpor of the surface and extremi-

ties, and produces a fense of coldness in the stomach, augments the oppression on the præcordia, and renders the pulse more frequent and more seeble. Its effects in all these respects are similar to the affusion of cold water on the skin in the same stage of the paroxysm as described in pages 18 and 37, though inferior in degree. If the thirst is gratisted in the cold stage of the paroxysm, it ought to be with warm liquids.

2. When the hot stage is fairly formed, and the furface is dry and burning, cold water may be drank with the utmost freedom. Large draughts of cold liquids at this period, are highly grateful; they generally diminish the heat of the surface several degrees, and they lower the frequency of the pulse. When they are attended with these falutary effects, fensible perspiration and sleep commonly follow.—These effects are similar to those produced by the affusion of cold water on the surface, as already described, but inferior in degree also. Though various cases are on record of the paroxysm of fever being diffolved by cold water, drank in this stage of the disease, my experience does not furnish me with any inftance of this kind*. Indeed fince I became

^{*} See Alpinus. Med. Meth. lib. ii. cap. 3. Where after reciting the effects of cold drink in diminishing heat and thirst and exciting profuse perspirations and large discharges of urine,

came acquainted with the extraordinary efficacy of the affusion of cold water on the surface, I have not trusted the folution of the paroxysm to its internal use. I have however employed cold drink when necessary as an auxiliary. Throughout the hot stage of the paroxysm cold water may be safely drank, and more freely in proportion as the heat is farther advanced above the natural standard. It may even be drank in the beginning of the fweating stage, though more sparingly. Its cautious use at this time will promote the flow of the perspiration, which after it has commenced, feems often to be retarded by a fresh increase of animal heat. A draught of cold water taken under fuch circumstances will often reduce the heat to the standard at which perfpiration flows more freely, and thus bring the paroxyfm to a speedier issue.

3. But after the perspiration has become general and profuse, the use of cold drink is strictly to be forbidden. At this time, I have perceived in more than one instance, an inconsiderate draught of cold water, produce a sudden chilliness both on the surface and at the stomach, with great sense of debility, and much oppression and irregularity of respiration.

he concludes, Mirabile eft, quomodo tale præsidium, has sebres expugnat; nam execretionibus, quas aqua suscitat, hat sebres siniunter. According to the same author this practice was followed by the Egyptians. See lib. ii. cap. 15.

ration. At fuch times, on applying the thermometer to the surface, the heat has been found suddenly and greatly reduced. The proper remedy is to apply a bladder filled with water, heated from 110° to 120°, to the scrobiculus cordis,* and to administer small and frequent doses of tincture of opium, as recommended by Dr. Rush. By these means the heat is speedily restored.

This effect of cold water used as a drink during profuse perspiration is precisely analogous to the affusion of it at such times on the surface of the body, a practice known to be of the utmost danger, and enumerated by Hossman among the causes of sudden death. Inveniuntur in amplissimo regno natura plura qua brevi tempore nocent ac perimunt, ut aqua frigida corpori sudore diffluenti immoderatius superingesta. Vol. i. p. 194.

The pernicious effects of cold water applied internally and externally during profuse perspiration, depend on the same causes, namely, that perspiration itself is a cooling process, under which when profuse, the heat of the body, whatever its actual state may be, is sinking; that under such circumstances,

we

^{*} See the Appendix, No. II. p. 27.

The pernicious effects of cold drink during profuse peripirations was known to the ancient.—See Celfus, lib. i. cap. 3.

we find as a matter of fact, it parts with its remaining heat more easily; and on the sudden application of cold, that this heat sinks to a degree which disturbs, and sometimes wholly interrupts the catenations on which life immediately depends.

Thus then we may fafely adopt the fame general rules for the use of cold water in sever as a drink, that have already been laid down for its external application. It may be used as a drink at any time, when there is no sense of chilliness present, when the heat of the surface is steadily above what is natural, and when there is no general or prosuse perspiration.*

Though we have arrived at these conclusions, the reader is entreated to believe, that the effects of cold water used as a drink have not been inferred from its external application, but made the subject of separate inquiry. Yet, that in the one case, and in the other, they should be similar in kind, though different in degree, will be expected by every one acquainted with the laws of the animal economy, and particularly with the sympathy that subsists between the stomach and the surface.

I have only to add, that in our common contagious

^{*} See p. 16.

gious fever, when I have used the affusion of cold water, I have feldom found it necessary to employ it largely as a drink, and my experience of its effects when drank in large quantities, has been chiefly confined to those cases where the fears or prejudices of the patients, or their friends, have prevented our having recourse to the more powerful method of affusion. For, however burning the thirst may be, it is speedily abated, and even removed, with very little drink, and often without any, by the fuccessful use of the affusion on the surface. Though the affusion in general suffices in our contagious fever, yet where cold water is employed in the dreadful fever of Philadelphia and the West Indies, it is probable that its internal and external use should be combined; a point that must be determined by the actual heat of the patients, measured by the Thermometer, and by their fensation of heat; circumstances of which it is to be regretted that we have as yet no accurate information.

duly to add, that in our common

CHAP.

CHAP. XII.

Of the disease that arises from drinking cold liquids, or using the cold bath, after severe exercise.

IT is here natural to inquire how far the fatal effects proceeding from drinking cold water, not in fever, but in cases where the system has been extremely heated by bodily exertions (of which the records of medicine afford fo many inftances) are to be explained on the principles already laid down. If they are explicable on these principles, we ought to be able to shew, that they have occurred in situations where the fystem, after having been much heated and enfeebled by fevere exertions, is lofing its præternatural heat from profuse perspiration, and in general also from the cessation of the exertions by which this heat was originally produced. Here two powerful causes combine to cool the body, and if under their operation, a sudden application of cold is made either to the stomach or the surface, the living power will, we know, refift it faintly, and the fatal confequences be accounted for.

- 1. In my own experience this fudden death has occurred once only, and that many years ago. It was in the case of a young man who had been engaged a long time in a most severe match at sives. After it was over he sat down on the ground, panting for breath, and covered with profuse perspiration. In this state he called to a servant to bring him a pitcher of cold water just drawn from a pump in sight. He held it in his hand for some minutes, but put it to his head as soon as he had recovered his breath, and drank a large quantity at once. He laid his hand on his stomach, and bent forwards; his countenance became pale, his breath laborious, and in a few minutes he expired. Various methods were employed to restore him, but in vain.
- 2. The following case resembles very exactly that just given. Blasius, Senensis, familiaris noster et condiscipulus, dum longiusculo tempore sub ardentissimo sole pilæ lusu incaluisset, nec sudore adhuc aut satigatione remissis, in subterraneum locum ubi vinaria erat cellula, descendisset, frigidimissimi vini calicem bausit: quo epoto, statim desicit. Benevent. cap. 17. De abditis.
- 3. Elegans & optimæ staturæ juvenis Romanus, eum pilå luderet, et sudore respersus, ac totus madidus, & satigatus ad puteum, pro siti arcenda venisset, exhausta

exhaustâ frigidâ recens per caldarium extractâ, illico in terram cecidit & obiit.

An inflammation in the Hometh fol-

- 4. Alterum novimus ex iis, qui in campo negotiantur, qui quum non minus corporis totius adapertis poris, madidus domum reverteretur, cyathum frigidioris aquæ ebibit, et mortuus fuit. Anat. Lusit. curat. med. cent. 2. curat. 62.
- 5. Forrestus relates, that in the year 1544, Valerius Cordus, a young man of great learning and talents, went during the heat of the dog-days, to collect plants among the Florentine mountains. Exhausted with fatigue and thirst, he incautiously drank of a cold spring which issued out of one of the hills, and was immediately seized with a fever, of which he died; but the symptoms of which he has not recorded. P. Forestus, lib. i. Scholio ad obs. 13.
- 6. Scaliger relates the case of a reaper, who, stooping down to drink at a sountain after severe labour, instantly expired. Scaliger de Subt. ad Cardan. exer. 13.
- 7. In Heister's observations, a case is related of a young man, who, about the Christmas season, had been playing and dancing at a Mill with some young women, and had eaten greedily of some hot H

buttered cake. After this, being extremely thirsty, he took a large draught of some cold water mixed with snow. An inflammation in the stomach followed, terminating in mortification, of which he died. Heister's Med. Sc. Observations, translated by Wiseman, p. 17.

8. Villanum quendam nobis familiarem novimus, qui messis tempore, anno 1597, exhaustus viribus, et totus sitibundas præ nimio solis ardore, domum rediens, cum in magna quantitate, ad sitim explendam, gelidam (aquam) bibisset, exanimatus mox juxta puteum cecidit, ac intra tres horas animam expiravit. Georgius Grasseccius in Theatro Anatomico.

These relations are chiefly taken from the collection of Schenck—they might be greatly enlarged, and I have collected a number of similar cases; which it were a tedious and an useless task to detail. In all of the cases which I have consulted, as well as in those I have related, three circumstances are either expressed or may be clearly inferred—1. The body had been previously heated beyond the temperature of health, by exercise carried to fatigue. 2. To this violent exertion a state of rest had in general succeeded. 3. A profuse perspiration had taken place. So far our reasoning is supported; but as these points are of the utmost consequence

consequence in explaining the operation of cold on the human body, and as direct experiments are attended with extreme hazard, the reader will excuse me, if I attempt to support them farther, by such evidence as history may incidentally afford.

1. In Quintus Curtius, (lib. vii. cap. 5.) an account is given of the march of the army of Alexander the Great in pursuit of Bessus, through the country of the Sogdiani, which is represented as destitute of water, sterile, and covered with scorch-The intolerable heat, fatigue, and ing fands. thirst of the soldiers in their march through this burning defert, are described with all the florid eloquence of the historian. At length fainting under their toils, they reached the banks of the river Oxus, where by indulging in large draughts of the stream, Alexander loft a greater number of his troops than in any of his battles. Sed qui intemperantius hauserant intercluso spiritu extincti sunt; multoque major borum numerus fuit, quam ullo amiserat prælio.*

H₂ = 2. A

^{*} The whole particulars of this march as described by Quintus Curtius, are very interesting. The desert, which contained not a drop of water, was four hundred Stadia across—(per quadringenta stadia ne modicus quidem humor exsistit,) that is, upwards of forty-six English miles. They began their journey in the night, directing their course by the stars, and for some time their march was tolerable, being refreshed by

- 2. A similar story is related by Appian—Appianus Alexandrinus de bellis civilibus, lib. v. tradidit Cornificianos milites a Pompeianis pugna fatigatos & assuantes fontanam aquam avide bibentes, ex iis plurimos emortuos. Marcel. Donat. lib. iv. cap. 6. Hist. med. mirab.
 - 3. A difaster of the same kind is recorded to have

country of the Soguians, which is represented as

the dews of night, and the coolness of the dawn; but when the fun rose, the heat became troublesome, and as the day proceeded, most oppressive; it was equally painful to stand still or proceed. After a day of dreadful fatigue, the van-guard of the army, and Alexander himfelf, reached the Oxus towards evening; and fuch as were themselves refreshed, were employed in carrying water back to the fainting troops behind. arrived in fuccession on the banks of the river, it may easily be supposed that they drank without moderation, and hence the destruction that ensued. It was on this occasion that Alexander displayed his magnanimity in refusing the cup of water brought to him as he advanced, because it was not sufficient both for him and his companions; and that he gave a proof of his genius by ordering fires to be kindled on the high banks of the Oxus, not merely to direct his way-worn foldiers through the darkness, but to animate their fainting exertions by a profpect of the end of their toils. Plutarch alludes to this ftory, but does not relate it fully. I fear it is not to be found in Arrian. A fimilar distress is mentioned by him (lib. vi. p.425) to have occured in Alexander's march through the deferts of the country of the Gedrofi, who inhabited the fouthern part of the Persian empire on the shores of the Indian ocean.

have occurred to the Christian army in the holy wars. Gulielmus Tyrius, lib. iii. c. 16. scribit, Christianum agmen Pisidiam ingressum, regionem arentem & inaquosam, tandem invento sluvio; quod quidem, qui largius aquam frigidam ingurgitaverunt, sitis discrimen evadentes, mortem in aquarum opulentia reperere.

These historical relations support very fully the doctrine I have already laid down. The cautious reasoner may not, on a hasty consideration, be inclined to rest with much confidence on this fort of evidence; but on reflection he will fee that it is entitled to confiderable authority; because the facts are in their nature not liable to be mistaken, and because they are not likely to be misrepresented. It is also entitled to much weight, because it is not given by these historians in support of any particular doctrine; and because the experiment having been made on fuch numbers of persons at once, it acquires an authority hardly to be ascribed to solitary cases, however accurately detailed. These considerations induce me to lay much more stress on evidence of this kind, than on the precepts respecting the effects of cold drink to be found in medical authors, ancient or modern. Nevertheless I have looked into the greater part of the ancients on this point, (for among the moderns there is not much on the subject) and have found nothing, that fully confidered, invalidates the conclusions I have laid down.

Of

Of the ancient physicians, the most copious on the use of water in all its forms, is Galen. He not only used cold drink, but immersion in the cold bath, in burning severs, with extraordinary success. His relations appear to me, in general tedious and obscure, but not destitute of truth; and the weariness of perusing him is occasionally relieved by the pleasure of rescuing a fact that was buried under masses of salse theory. The reader who would consult him on this subject may use the references below.*

clined to reft with much confidence on this fort of

Observations, published by Dr. Rush of Philadelphia, 1789, an account is given of the "Disorder coccasioned by drinking cold water in warm weather," which frequently occurs there, "Three circumstances," he observes, "generally concur to produce disease or death from drinking cold water.

The patient is extremely warm. 2.

The water is extremely cold. And 3. A large quantity of it is suddenly taken into the body.

The danger from drinking cold water is always in proportion to the degrees of combination which coccur in the three circumstances that have been mentioned.

bar * Vol. i. 23. B. no daim non a syaft amabora an

Vol. ii. 78. C.

Vol. vii. 70. A. and forwards throughout the volume. I quote from the Latin edition in folio, published at Venice, 1656.

" mentioned." p. 151. Dr. Rush goes on to state the symptoms of this disease, which are, I apprehend, given with accuracy. His method of cure is in my judgment proper and judicious. "I know " but one certain remedy for this difease, and that " is liquid laudanum. The doses of it, as in other " cases of spasin, should be proportioned to the vio-" lence of the disease. From a tea-spoonful to near " a table-spoonful has been given in some instances " before relief has been obtained. Where the "powers of life appear to be fuddenly fuspended, "the fame remedies should be used which have "been so successfully employed in recovering per-" fons supposed to be dead from drowning." To this I would add the application of a bladder filled with water heated to 110° or 115° of Fah! to the pit of the stomach, from which I have feen powerful effects in restoring the vital heat*. But while I accede to Dr. Rush's practice I cannot subscribe to his notion of the causes of this disease, or to the method of prevention founded on this cation of cold to the body, whether to the fur.noison

Dr. Rush seems to entertain the popular opinion on this subject: the body is extremely warm; the water extremely cold, and a large quantity is introduced H 4 fuddenly

^{*} See Appendix, No. II. p. 31, 32.

fuddenly. He apprehends the danger to arise from the great difference between the temperature of the body and of the water taken in. As a means of prevention he therefore proposes to such as cannot be restrained from drinking cold water when præternaturally heated-1. To grafp the veffel out of which they are about to drink for a minute or longer with both hands; that a portion of heat may be abstracted from the body and imparted to the cold liquor. 2. If they are not furnished with a veffel to drink out of, but obliged to drink at a pump or a spring, always to wash their hands and face previously to drinking, with a little of the cold water. "By receiving," fays he, "the shock " of the water first on those parts of the body, a " portion of its heat is conveyed away, and the "vital parts are defended from the action of "the cold," all gainofler ai are the following and

The fact however is in my mind perfectly established, that there is no situation in which the application of cold to the body, whether to the surface or the stomach, is so safe, or in general so salutary, as when the heat of the body, from whatever cause, is præternaturally great, provided that the body is not already in a state in which it is rapidly parting with this heat, and no disease has taken place in the general sensibility, or in the structure of any of the parts; and where the body is præternaturally heated,

ter ele to Dr. Ruffi's practice I cannot

the degree to which cold water may be drank, may be always decided by the steadiness of the sensation of heat, and the tenacity with which the præternatural heat is actually retained. Thus in continued fevers it may be drank to a greater extent, than in the hot stage of intermittents, because the heat is more firmly retained; the profuse perspiration not being at hand, by which the febrile heat of intermittents is carried off. The ancients who gave cold drink largely in continued fevers, were doubtful of its use in intermittents.* It may however be given (as I have already stated) with great safety in intermittents, provided it be taken in the time that intervenes after the hot stage of the paroxysm is fairly established, and before the perspiration that follows it, has become general and profuse. As however it is only in that interval that it can be given in intermittents with advantage or fafety, we can eafily understand that the ill effects arising from its being accidentally drank in the cold, or the fweating stage of the paroxysm, (in both of which the thirst often demands liquids) may have produced the doubts which fome

^{*} See Sennertus, lib. ii. cap. ix. p. 54. Itaque Græci auctores jubent in statu, cum febris acuta, sitis, inquietudo, cordis & arteriarum pulsatio est vehementissima, ægerque avidissime eam expetit, aquam frigidam copiose exhibere.—

In intermittentibus febribus vero aquæ frigidæ potus nunquam convenit.

have expressed in regard to its use in intermittents, and the interdiction which in such cases, others have pronounced against it. We may explain also from the same considerations, why in the accounts that have been handed down to us of injurious effects from the use of cold drink in severs, the greater part of the cases have been intermittents.

The instances however that are recorded of the fatal effects of large draughts of cold liquids, have more frequently occurred after fevere exercise and fatigue, than even in intermittent fever. The cause of this is obvious—the heat præternaturally accumulated by exercise, is held with less tenacity than even the heat in intermittents. It is diffipated by the perspirations that exercise occasions, and is speedily lost, when to profuse perspiration is added a state of rest. It is then that a large draught of cold liquid is especially dangerous. But while the præternatural heat is sustained by continued exertion, cold liquids may be taken in moderate quantities without producing any injurious effects. They may even I apprehend be drank copiously without producing fuddenly the fatal effects already describedbut in copious draughts they are found oppressive to the stomach during exercise, and they excite languor, nausea, and sometimes vomiting, as I have had occasion to observe. In the narrative already mentioned of the march of Alexander's army the vero some frigide potes and through

through the defert country of the Sogdiani, it is related by the historian, that a few of the foldiers, by the advice of the natives, had provided themfelves with water, of which under their burning thirst they drank immoderately. The consequence was, that they became heavy, feeble, and unable to fupport their arms, and this state of oppression was succeeded by severe vomiting. Graves deinde avide bausto bumore, non sustinere arma; non ingredi poterant; et feliciores videbantur, quos aqua defecerat, quum ipfi fine modo infusam vomitu cogerentur egerere. Q. Curtius, lib. vii. cap. 5. The water thus wasted, or worse than wasted, might have been used to advantage in wetting, from time to time, the garment next the skin. Thus the oppression of its weight on the stomach would have been prevented, and the furface of the body being kept cool by constant evaporation, the heat of the fystem would have been moderated, and the thirst alleviated*.

If

^{*} Postilions understand the difference between giving their horses cold water to drink, during exercise, and after the exercise has ceased. When in their power, they always water their horses two or three miles before the end of their journey. My friend Mr. Charles Aiken assures me, that during his tour on foot through Wales, in company with his brother, of which Mr. Arthur Aiken has given so interesting and useful an account, they drank of the pure streams, as they descended from the mountains, without reserve, during the fervor of the day, taking care however never to rest after drinking.

If this account of the circumstances under which cold drink after severe exercise proves injurious, be just, the precautions of Dr. Rush to those who will drink in such circumstances, are founded on error. By abstracting a part of the præternatural heat of the body before drinking, the danger is not diminished, but greatly increased. This enlightened physician will excuse these observations, drawn from me by a consideration of the importance of the subject, but accompanied by sentiments of sincere esteem and respect.

If the effects of cold water used internally under fevere exercise, are not entirely analogous to the effects produced by its affusion on the skin;-the difference will be easily understood, by those who consider, that where a quantity of water is swallowed, besides the influence of the cold, the stomach suftains a load, from the weight and the bulk of the liquid, particularly oppressive under constant action and agitation of the voluntary muscles, from which the furface, moistened with water, is entirely free; and on the other hand, that the evaporation from the furface promoted by the immediate access of the external air, must operate more directly in cooling the body, and particularly in counteracting the burning rays of the fun, than water taken into the stomach. With these exceptions, the operation of cold liquids on the stomach and on the furface face of the body are analogous in the case of præternatural heat produced by bodily exertion, as in all other cases of præternatural heat. As it is safe to drink cold water in proportion as the heat from exercise is great and steady, so also is it safe according to this ratio, to pour it on the surface, or to immerse the body in the cold bath.

In the earlier stages of exercise, before profuse perspiration has diffipated the heat, and fatigue debilitated the living power, nothing is more fafe, according to my experience, than the cold bath. This is fo true, that I have for some years constantly directed infirm persons to use such a degree of exercise before immersion, as may produce some increased action of the vascular system, with some increase of heat; and thus fecure a force of re-action under the shock. which otherwise might not always take place. The popular opinion that it is fafest to go perfectly cool into the water, is founded on erroneous notions, and fometimes productive of injurious confequences. Thus, persons heated and beginning to perspire, often think it necessary to wait on the edge of the bath until they are perfectly cooled; and then plunging into the water, feel a fudden chilliness that is alarming and dangerous. In fuch cases the injury is generally imputed to going into the water too warm, whereas in truth it arises from going in too cold.*

But though it be perfectly safe to go into the cold bath in the earlier stages of exercise, nothing is more dangerous than this practice after exercise has produced profuse perspiration, and terminated in languor and satigue. Because, as has already been repeated more than once, in such circumstances, the heat is not only sinking rapidly, but the system parts more easily with the portion that remains.

This account of the operation of the cold bath will explain some circumstances very generally mentioned by writers on the effects of cold on the human body, and hitherto not properly accounted for. That the Roman youth in the heat of their exercise in the Campus Martius, frequently plunged into the Tyber,

^{*} Dr. A. Munro Drummond, in his inaugural differtation, "De Febribus Arcendis," the only specimen lest of his admirable talents, speaking of the effects of the cold bath as a preventative against the action of contagion, observes, "Nec "frigida protinus sugienda vel calida temperanda, quamvis "cutis pallida aliquantisper siat vel leviter aliquis inhorruerit. "Si exercitatio antecedat quæ citra lassitudinem & sudorem "est, hæc facile evitari posse experiendo didici: et sere, se "nil aliud obstet, quo ante quisque plus incalueret quam "aquam intraverit, eo calidior emerget.

Tyber, is a fact universally known; they found in this practice a high enjoyment, and they believed it conducive to health, and more especially to sleep. On the other hand various relations may be cited of the injurious effects of an apparently similar practice; the most remarkable of which is the case of Alexander the Great, when covered with dust and sweat, he threw himself into the Cydnus, and was seized with a disease, of which he nearly perished; one of the best authenticated sacts in ancient history.

In the differtation De febribus arcendis, by Dr. A. Munro Drummond, already quoted, these sacts are brought together in the following words.—Alexander, quondam diei fervidissimo tempore, liquore shuminis invitatus, vix dum in Cydnum amnem descenderat, cum gravi inde morbo implicitus est. Romana juventus, post quotidianas in Campo Martio exercitationes, pulvere simul & sudore persusa, Tyberi impune lassitudinem cursus natandi labore deposuit. Mali nimirum adsuetudine duramur in his sicut in cateris rebus omnibus. Thes. Med. vol. iii. p. 154. Doubtless the influence of habit has a considerable share in regulating the estects of cold on the human body; but the circumstances just mentioned seem capable of an explanation on other principles.

On the Campus Martius, the exercises of the Ro-

man youth were carried on with all the vehemence of emulation. Swimming formed a part of those exercises *, and generally terminated the foot-race. The youthful candidates in this exercise directed their course towards the banks of the river, and plunged headlong into the stream. Sometimes the contention did not terminate till the river was once, or perhaps twice swam across. Hence it will easily be feen, that they were accustomed to immerse themselves in the water in the very fervor of their exertions, when the heat was præternaturally great; and not after the body was cooled by profuse perfpirations, or exhausted by long continued fatigue. In this fituation the practice was fafe; without taking into confideration, that the perfons concerned were in the flower of life, fortified by early habits, and partly defended from the shock of immersion by the inunctions which feem to have been generally used among the Romans, before the cold+,

Cur timet flavum Tiberim tangere?
See Vigetius, lib. i. cap. x.

See also Heiron. Mercur. lib. iii. cap. 14.

+ See Horace Satyr, lib. ii. Sat. 1.

Ter unchi

Transnanto Tiberim, sommo quibus est opus alto.

^{*} See Horace, lib i. ode 8. Where the poet, after queftioning Lydia why her lover neglects his accustomed exercises on the Campus Martius, among other particulars asks,

and after the hot bath; and which were particularly employed by the athletæ of Greece and Rome in all their exercises. It was the more safe, because the stream of the yellow Tiber being comparatively scanty and slow, its waters speedily received the influence of the sun, and acquired the temperature of the atmosphere. Where the air and the water are of the same temperature, the rarer element prepares the body for the contact of that which is more dense.

The circumstances under which Alexander plunged into the Cydnus, were different in many essential points. He had marched at the head of his troops to seize a pass in Mount Taurus, necessary to facilitate his passage into Cilicia. After having secured his

The subject of the ancient unquents is treated of by Hieronymus Mercurialis, de arte Gymnastica, lib. i. cap. x.

The same author, (lib. iii. cap. 14) speaking of swimming, so much in estimation among the ancients, observes, quo pacto vero similis exercitatio perageretur, tanquam res vulgatissma silentio praeterita suit sere ab omnibus, uno excepto Antyllo, qui nataturos prius moderate unxisse, et frictione corpora praecalsecisse deinde e vestigio in aquam se projicere debere voluit, quasi omnes sanitatis gratia natantes eas regulas observarent.

This method of previously anointing the body, might be recommended to our modern swimmers, not only as defending them from the shock of immersion, but as enabling them to glide through the water with less effort. his object, he descended from his elevated station, through a road, difficult and full of defiles, to the city of Tarfus, which was fituated at the bottom of the mountains. His whole march, which probably continued feveral days, was attended by extraordinary exertion, not from the nature of the country only, but also from the pressure of circumstances. In ascending the heights, he had to haften forwards, left the enemy should pre-occupy the pass by which Cilicia must be entered. In descending from the immense elevation he had reached, he was stimulated by the defire of faving the city of Tarfus, the capital of the province, which the Persians threatened to consume by fire. In both of these enterprizes Alexander was successful. It was at the close of this extraordinary march, according to the testimony of all the historians, that the conqueror, advancing into Tarfus at the head of his troops, covered with dust and sweat, and exhausted with long continued toil, stripped himself in the fight of his army, and plunged into the pure and cold waters of the Cydnus, which ran through the city. The lymptoms that followed, are described with sufficient minuteness and precision. Vixque ingressi subito borrore artus rigere caperunt : pallor deinde suffusus est, et totum propemodum corpus vitalis calor reliquit. Exspiranti similem ministri manu excipiunt, nec satis compotem mentis in tabernaculum deferunt. Q. Curtii, lib. iii. cap. v. From the length and difficulty of the moles del dres return set e guo march,

march, it is natural to suppose that Alexander must have been cooled as well as debilitated by extraordinary perspiration and fatigue, and under such circumstances, immersion in the cold and rapid Cydnus, was followed by the consequences which we should expect from the principles already laid down *.

In Prudence which one I 2 family fonce knew an

* If an objection be urged against our laying much weight on a relation taken from a period of history so remote, I reply, that I am far from resting any part of my conclusion upon it, having the sirmer foundation of actual observation. But as this adventure of Alexander is mentioned by almost every writer on the effects of cold on the living system, to exemplify the danger of its being applied when the body is præternaturally heated, it seemed proper to shew, that it is capable of a more just and natural explanation.

In regard however to the fact itself, of Alexander having suffered from bathing in the Cydnus, it might be easily shewn, that we cannot reject it without denying entirely the weight of historical evidence. 1. This exploit was performed before thousands of witnesses, by some of whom it is recorded, and it is mentioned by all the historians. 2. The circumstances that preceded it, and the consequences that followed it, are also recorded with great minuteness, and form a natural chain of events. 3. The act itself of stripping before his army, and springing into the Cydnus, may have an air of improbability; but is not so inconsistent with Greek, as with modern manners; and however unlike the dignity of great commanders in our own days, is perfectly characteristic of Alexander—of that ardent and daring temper, and of that buoyant imagination, which were equally regardless of danger and decorum;

In his effay on swimming, Franklin makes the following observation: "During the great heats of "fummer there is no danger in bathing, however "warm we may be, in rivers which have been thoroughly warmed by the sun. But to throw ourselves into cold spring water, when the body has been heated by exercise in the sun, is an imprudence which may prove fatal. I once knew an "instance"

a temperament, that joined to extraordinary talents, attached his foldiers to him in fo fingular a degree, and which in the æra of the world in which he lived, was calculated to carry him to the aftonishing elevation which he attained. 4. The exploit itself was of a nature not likely to be feigned; and the issue of it had nothing marvellous—there was nothing in any part of it, by which the son of Ammon, or his followers, could be in any way gratified.

The effects of Alexander's bathing in the Cydnus, gave a celebrity to this river which it otherwise would not have acquired. I have called it cold and rapid, meaning to speak of it comparatively with the Tiber. The Cydnus was one of the principal streams that descended from Mount Taurus, (Strab. xiv. p. 462.) at the bottom of which stood the city of Tarsus. From the great elevation of the mountain, the Cydnus must in this situation have been rapid and cold; and colder because rapid. From the same cause it would be easy to shew (if it were at all to the purpose) that the transparency ascribed to its waters was probably ascribed justly. Quintus Curtius imputes its coldness to the rays of the sun being excluded by the shadyness of its banks: the more accurate and philosophic Arrian, mentions as the cause, the lostiness of the mountains

" instance of four young men, who having worked at harvest in the heat of the day, with a view of refreshing themselves, plunged into a spring of cold water; two died on the spot, a third the next morning, and the sourth recovered with great difficulty. A copious draught of cold water in the same circumstances is often attended with the same effect in North America." The

whence its fountains descended. Quintus Curtius speaks of it as gliding through a gentle descent, (leni tractus e fontibus labens,) and Tibullus of its silent and placid stream.

At te, Cydne, canam, tacitis qui leniter undis, Cæruleus placidis per vada serpis aquis.

Strabo, on the other hand, (lib. xiv. 392.) mentions its current as most rapid, and this to some of the scholiasts seems a contradiction. It is not so in reality. If the declivity was uniform, the current might be at once fwift and filent. The river Moffat (the fource of the Annan) defcends from the mountain of Hartfell, by a declination, which in three thoufand yards, amounts to two hundred and thirty feet, with fuch uniformity, that in the course of some miles there is hardly a fingle ripple. To finish a digression by far too long, but which may relieve the dryness of a professional subject, it may be observed, that our account of the illness of Alexander receives confirmation from a difference among the hiftorians respecting its causes; some imputing it to his having bathed in the Cydnus, and others to his previous fatigue: at the distance of two thousand two hundred years, it may be clearly determined that both circumstances combined in producing his difease.

authority of the American Bacon is of great weight in medicine, as in every branch of science; and particularly in what respects immersion in water; for doubtless he spent more time in this element than any philosopher of modern days. It may however be eafily supposed that he adopted the commonly received opinion, that the injury arose from the perfons in question going in when bot, instead of from going in when cooling, after having been heated; to which last circumstance, it can hardly be doubted, that the fatal accident he relates, was to be imputed. It is not however to be denied, that the difference between the temperature of spring and river water, at certain feafons of the year, will make a great difference (in any circumstances of the body) in the consequences of immersion. Springs which issue copioufly from the earth, are, as we know, both from reasoning and observation, nearly of the temperature of the earth at some distance under the surface; and this temperature, when unaffected by chemical changes, is probably (for actual experiment is not yet fufficiently extensive to affirm the proposition univerfally) of the mean heat of the atmosphere, taking the year round, under which it lies*. On the other hand, rivers even of the largest fize, acquire

at

^{*} See Dr. John Hunter's valuable paper on this subject, in the Philosophical Transactions for 1788, p. 53.

at a certain distance from their source, the temperature of the earth, and rife towards the end of fummer, to nearly the highest degree of the summer heat*. In countries where the temperature of the seasons has little variation, the difference between the heat of the rivers and springs will be very inconfiderable. That this is the cafe in Jamaica, we know from the experiments of Dr. Hunter; but in Penfylvania, to which it is probable that Dr. Franklin's observations more particularly refer, this difference at certain feafons of the year must be very great, the difference of the temperature of the feafons being fo very confiderable. The rivers of Penfylvania in the latter end of fummer, are probably nearly as warm as the Buxton bath, or from 78? to 82°; and in the Buxton bath there is reason to believe, that even after confiderable fatigue, the lial chill choic my remarks on the circumstances

which render the cold bach title or hazardous aff

^{*} Thus the Mersey at Liverpool, in the middle of the tide, is in the month of August from 65° to 69°. On the 20th of August 1795, it rose in the middle of the tide to 70°, and under the rocks at High Park, the mercury stood on the 21st, at high water, at 72°.—In the air, at the same time, it was at 77°.—The weather had been uncommonly warm for sourteen days before, and 72° was probably the average heat of that fortnight. It seems to be this average that the tide of the Mersey acquires. Observations made in our different æstuaries every week, would give the actual and the comparative heat of the seasons (on the coast at least) with much more accuracy than observations in the atmosphere.

chill of immersion is too inconsiderable to produce much hazard. The fprings of Pensylvania, and more especially the deep funk wells, if they retain the fubterranean temperature (the medium heat of the feafons) will probably be in the heat of fummer and autumn, twenty or twenty-five degrees cooler than the rivers or the atmosphere*; a difference capable of producing, in the fituation of the body which renders the application of cold dangerous, the most serious effects. It is doubtless owing to the great comparative coldness of their springs and wells. that fatal accidents from drinking cold water, and bathing in cold springs, are so frequent in Pensylvania. In the West Indies, where water is seldom to be procured below 77°, fuch accidents, if they ever occur, are very rare.

I shall close my remarks on the circumstances which render the cold bath safe or hazardous after exercise, with the following narrative:—

On the first of September, 1778, two students of medicine at Edinburgh, set out on foot on a journey, a considerable part of which lay along one of the rivers of Scotland. They started by surrise.

^{*} The wells at New York are from 54° to 56°, according to Dr. Routh. See Philosophical Transactions for 1788, p. 61.

rife, and proceeded with alacrity in the cool of the morning. At the end of eight miles, they breakfasted, rested for an hour, and then resumed their journey. The day grew warm as it advanced, and after a march of eight miles more, they arrived heated, but no fatigued, on the banks of the river abovementioned, about eleven in the forenoon.-Urged by the fervor of the day, and tempted by the beauty of the stream, they stripped instantly, and threw themselves into the river-The utmost refreshment followed, and when they retired to the neighbouring inn, this was succeeded by a disposition to fleep, which they indulged. In the afternoon they proceeded, and travelling fixteen miles farther at a fingle stretch, arrived at the inn where they were to fleep, a little after fun fet .- The afternoon had been warm, and they perspired profusely; but the evening was temperate and rather cool. They had travelled for fome miles flowly, and arrived at the end of their journey, stiffened and wearied with their exercise.

The refreshment which they had experienced in the morning from bathing, induced however one of them to repeat the experiment, and he went perfectly cool into the same river, expecting to relax his limbs in the water, and afterwards to enjoy profound sleep. The consequences were very different. The Tweed, which was so refreshing in the morning,

now felt extremely cold; and he left the water haftily. No genial glow fucceeded, but a feverish chill remained for some time, with small frequent pulse, and flying pains over the body. Warm liquids and frictions brought on at length confiderable heat, and towards morning perspiration and sleep followed. Next day about noon they proceeded on foot, but the traveller who had bathed was extremely feeble; and though they had to perform a journey of a fingle stage only, as some part of it was difficult and mountainous, he was obliged to take the affiftance of a carriage which overtook them on the road. It was feveral days before he recovered his usual vigour. This relation will not I hope be deemed of the less authority, because it is given by the person who fuffered by his imprudence*. It is unnecesfary to point out the application of these incidents to the doctrines already laid down.

It

I may here observe, that illuesses similiar to the above occur frequently, from the imprudence of the country people, (the Boltoners, as they are called) who make an annual visit to the river below Liverpool, for the purpose of bathing.—In such cases, I have found the disease that followed, was not instammatory, as might be supposed, but rather a sever of debility, as in the instance detailed.

^{*} Those who know the road from Edinburgh to Mosfat will be at no loss to mark the different stages of this adventure.

It feems to be a general truth, that from whatever cause the heat of the body is increased, in proportion to this increase (provided no local disease has occurred) is the fafety with which cold may be applied. In the celebrated experiments of Dr. Fordyce, Sir Charles Blagden, and others, of which an account is given in the Transactions of the Royal Society, vol. lxv. p. 111, and 484, it is repeatedly mentioned. that these gentlemen passed from a room heated to 200° and upwards, into the cold air with perfect fafety. "During the whole day," fays Sir Charles Blagden, "we paffed out of the heated room," (where the temperature of the air feems to have been 240°, and sometimes 260°,) " after every ex-" periment, immediately into the cold air, without " any precaution; after exposing our naked bodies " to the heat, and fweating most violently, we in-"flantly went into a cold room, and flaid there, " even fome minutes before we began to drefs; yet " no one received the least injury." Ibid. p. 494.

The freedom from injury, and even from inconvenience, which these gentlemen experienced, depended on the increased heat of the body, and the increased action of the arterial system.—Had they continued exposed naked to the cold air till the heat sunk as low as its natural standard, and the heart and arteries subsided into their usual state of action, their situation would have been very hazardous.

In fimilar experiments repeated at Liverpool, of which a detail is given by Dr. Dobson, in the same volume of the Philosophical Transactions, p. 463, the gentlemen concerned, passed from the heated room into the cold air, with equal impunity. My friend Mr. Park assures me, that after remaining some time in the stove, where the heat was as high as 202°, he went into the external air without a great coat, or any other than his usual clothing, during a hard frost, and perceived neither injury nor inconvenience.

In many manufactures, the persons employed are exposed to extraordinary degrees of heat, particularly in the manufacture of glass. Such persons, guided by nature, often endeavour to obtain relief, by exposing themselves to degrees of cold, which on the commonly received opinions, would in their situation appear extremely dangerous. Thus at Glasgow in the spring of the year 1780, I learnt, that it was common for the workmen in the glass manufactory, after enduring for some time the consuming heat of their surnaces, to plunge into the Clyde; a practice which they found in no respect injurious.

A great variety of the phenomena respecting the influence of cold on the living body, receive an easy explanation by attending to the principle already mentioned.

mentioned. Thus after the heat of the body is increased two or three degrees in the hot bath, it is not only fafe, but refreshing, to plunge into the cold bath, as I have repeatedly experienced. A practice of this kind prevails, as is well known, in Ruffia, where it is common, after remaining fome time in the hoth bath, to roll naked in the fnow, and return to the warm bath as before. The Ruffian hot baths vary from 10610 to 1160 of Faht.* and I find by experiment that the actual heat of the living body is increased in the hot bath (unless under particular circumstances afterwards to be explained) when the temperature is no higher than 100°. We may therefore fafely conclude, that in making thefe fingular transitions, the heat of the Russian is in the first instance increased beyond the natural standard; and it is to this increase that the safety, as well as the agreeableness of the practice is to be attributed.

It is in this way that we must account for the safety of a practice which prevailed at Rome under

the

^{*} See Dissert. physica experimentalis de calore Animalium, by J. A. Braun, in the Now. Comment. Academiæ Scientiarum Imperialis Petropolitanæ, vol. xiii. The Abbé Chappe D'Auteroche says, the Russian baths are as high as 60° of Reaumer, (160° of Fahrenheit) Philosophical Transactions, vol. lxv. p. 112; but this is incredible, unless he speaks of vapour baths.

the first of the emperors. After the overthrow of the republic, the Romans confoled themselves for the lofs of their freedom, by a more unbounded indulgence than ever in those fensual gratifications which had led to their fall. Of these, the pleasures of the bath formed a diftinguished part, they fought every means of heightening and divertifying them, and connected them with other epicurean enjoyments. The mild and tepid immersion of the Greeks no longer satisfied them; they heated their baths to the utmost pitch of endurance; and as they rose reeking from their furface, veffels full of cold water were dashed over their naked bodies, as a high gratification in itself, and a means of stimulating the senses to gratifications still higher. Such practices could not however be continued without injury: In the bagnios of Imperial Rome, the last of the Romans, the followers of Zeno and of Cato, fought to footh their forrows, and were melted down into flaves; the powers of their minds became enfeebled, the vigour of their frames decayed, and they loft for ever the bold impressions of freedom and of virtue*. It

^{*} This subject might lead to digressions, which my limited plan does not admit. I shall content myself with a few observations. Though private baths were in use in Rome from the earliest periods of their history, yet the system of public hot baths did not, it is understood, commence till the days of Augustus. According to Dion, Mæcenas introduced it.—Agrippa followed his example—It was soon carried to an asso-

It was supposed by Dr. Cullen, who paid much attention to the operation of temperature on the living

nishing height, and the construction of baths where the people might be accommodated gratis, was an established and a successful method of gaining their affections. The extraordinary expense and magnificence of those structures are well known—the remains of the baths of Caracalla and Dioclesian testify their former grandeur in our own days. According to Fabricius, there were 856 public baths at Rome, and some of these were large enough to contain at once 1800 persons. The rage for hot bathing in Rome, under the first Emperors, exceeded all bounds.

In regard to the exact temperature of their baths, as the thermometer was not known to the ancients, we cannot speak precisely. We have however the authority of Seneca and Plutarch for saying, that the bath of the Greeks was of a moderate temperature; and as Alexander sound refreshment in the bath in the burning sever of which he died, it probably did not exceed 93°, or at most 96°, of Fahrenheit. But at Rome, in the days of Seneca, the hottest baths were most in estimation, and those of Nero seem to have exceeded all others in heat—We may infer this from one of the best of Martial's epigrams.

Si temperari balneum cupis fervens
Faustine, quod vix Julianus intraret,
Roga, lavetur, Rhetorem Sabineum,
Neronianas is refrigerat thermas.

Lib. iii. 25.

A person was employed at one time to regulate the heat of the baths, but in Seneca's days this had fallen into disuse. (Epist. 37.) This rage for bathing was checked by Adrian, and regulated by Severus. The fashion of heating the baths to this extraor-

living system, that in all sudden changes from a higher to a lower temperature, a fensation of cold takes place, even though the lowest point of change be such as when permanent; becomes oppressive by its heat. Thus that a change of the external air from 90° to 85°, is acompained by a fensation of cold, though this speedily goes off, and is succeeded by a fensation of heat, if the temperature of 85° be continued*. This observation ought, I think, to be restricted to degrees of heat inferior to that of the human body, for it does not feem true when applied to superior degrees of heat. If for instance, a person is immersed in the bath heated to 104°, and a quantity of cold water be speedily admitted, so as to lower it to 98°, a sensation of a very agreeable nature takes place, not however to be denominated cold-Where the actual heat of the body has been raised, and the circulation accelerated, a much greater reduction of the temperature of the bath must take place before it becomes cold to the railed lin tobarray avad or mand oraid to stone bar fenfations.

in heat-We may inter this is one one of the wift of Warti

dinary degree did not continue—We have the authority of Galen, who flourished soon after Seneca and Plutarch, that in his days very hot baths were no longer in use. See Hier. Mercur. lib. i. cap. 10.

^{*} See Differt. Med. Inaug. De frigore, by A. Cullen, published at Edinburgh 1780, (p. 8 and 9) which contains the best general view of the doctrines of the author's celebrated father on the operations of cold.

fensations. In the same manner a person under the steady heat of sever, (102° to 106°) suddenly throwing off his bed-clothes, and exposing himself naked to the external air, does not perceive it cold, but merely cool, a sensation too which goes speedily off. It is true there are some exceptions to this in sever, of which I have given one instance in detail. (See page 46.)

The manner in which our fensations are affected by changes of temperature, is a subject of importance, as well as of difficulty, for without a more precise knowledge of this, the action of temperature on life cannot be understood. In the state of health, I believe, however, that our sensations of heat and cold may be reduced to general principles.—But in certain of the Neuroses, as well as of the Pyrexiæ, these sensations are much perverted, and even in health they are much influenced by constitution, habit, and situation. To enlarge on this point would however lead into the general subject of the operation of cold on the living system, for which I am not sufficiently prepared, and which I have studiously avoided.

the ficady bear of fever, (102° to 106°) inddeply throwing off his b.IIIX . TAHO expenses himself

but merely cool, a fenfation ton which goes freedily

nie sido or andicomo chis in

Use of the cold Bath in convulsive Diseases—In Insanity—General Remarks.

IT might now be expected that fome observations should be offered on the operation of cold water applied to the surface, or received into the stomach, in the various cases that have been related; but besides that it is advisable the reader should form his conclusions in a great measure for himself, we shall be better prepared for such general reasonings as may suggest themselves, when we have observed the operation of the cold bath on convulsive affections, and on other diseases, classed by Dr. Darwin under the same general title, the diseases of volition*.—To save repetition, the reader is here supposed to have perused the paper in the Appendix, N°. I, of which he will consider the following observations as forming a sequel.

Since

[#] The third class of Dr. Darwin .- See Zoonomia, vol. ii.

Since the original publication of this paper, I have feen three different cases of tetanus arising from wounds, in which the affusion of cold water was employed alone, and in all of these the disease terminated fatally. I have seen a sourth case in which the affusion of cold water was conjoined with the liberal use of wine and opium, where the patient recovered. And I have seen two other cases in which the cure was trusted to wine and opium alone, in one of which the issue was fortunate, in the other unfortunate.

It is however proper to observe, that in the three first cases, the disease was far advanced, having refifted other powerful remedies before the patients fell under my care; and as the power of deglutition was nearly loft, the affusion of cold water was employed under desperate circumstances, as a dernier refort. It was befides impossible to use this remedy to any extent, for the powers of life were too much exhausted to sustain the continued application, or the frequent repetition, of so violent a stimulus.-There is also in the advanced stage of tetanus, an objection to the use of a remedy that requires much change of posture. In such cases, every exertion of the will on the voluntary muscles has a tendency to induce a general convultion.—From a confideration of all these circumstances, and from more extensive experience, I should now be difinclined to the use

of the cold bath in any of its forms, in the symptomatic tetanus, unless in the earlier stages of the disease, when the vigour is less impaired, and the convulsive actions less firmly catenated.

The practice, first introduced by Dr. Rush, of giving wine in large quantities in tetanus, of which a successful case is detailed in the paper already mentioned*, seems to me to be fully confirmed, as the most efficacious treatment of this very dangerous disease. The wine ought however to be combined with opium in considerably larger doses, according to my later experience, than in the case already referred to.

In the Appendix, N°. I, several cases are given in which the cold bath was used in other convulsive disorders. This remedy I have now employed in various diseases of this class for sourteen years, and on the whole, it has been attended with considerable success.—One conclusion, of rather a singular nature, was drawn from those cases, which my subsequent experience has uniformly corroborated:—" That the efficacy of the cold bath in convulsive disorders, is much promoted by its being employed during the presence of convulsion*;" or,

as

^{*} See Appendix, No. I. p. 9.

as I would now express it, that the chief benefit derived from the cold bath in convulfive diseases, depends on its being used in the paroxysm of convulsion; that its efficacy consists in resolving or abating the paroxysm; and that when this effect is produced, the return of the paroxysm is greatly retarded, if not entirely prevented. To this singular fact, for which I am unable to account, I invite the attention of physiologists: it may throw some light on the laws of association; a subject which Dr. Darwin has shewn to be of such vast importance in physiology, and which hitherto has been so little understood. The following case, taken from the records of our Insirmary practice, is in many respects curious, and will serve to corroborate the position just laid down.

John Westmore, aged 22, was admitted into the Infirmary on the 11th of October 1792. In consequence of a fright, he had been seized with sits two months before, which now recurred several times every day, of various duration, from two minutes to an hour. During these his consciousness was wholly abolished. These sits occurred without warning, and were peculiar in their appearance. At first the muscles of one side were strongly retracted, then those of the other, alternately; and then the muscles of both sides acting together, the whole trunk of the body was drawn upwards to the head; this action resembling very exactly that of K3

the victim of the law, suspended in the agonies of death. His faculties of mind did not feem as yet impaired, nor the animal functions much diffurbed. No medicines were ordered for this patient, but a bucket-full of cold water was directed to be thrown over him, the instant of the accession of the fit. Some circumstances prevented this being done for a few days, during which he went into the cold bath daily, when the fits were off him, and with feeming advantage; the number of paroxysms being reduced from eight or ten in the twenty-four hours, to two or three. At length he was thrown into the cold bath on the inftant of the accession of one of his fits, which was speedily terminated, and from this time forth he had no return for fourteen days, when he was discharged as cured. Westmore continued free of complaint till the beginning of February following, when his fits returned. He was re-admitted on the 24th of that month, and at that time his fits recurred fix or feven times in the day. He was again directed to use the cold bath dailyand to have a bucket of water thrown over him on the accession of every fit. By this practice he speedily recovered, and fince that time the disease has never returned. He is now, (March 1798) in perfect health.

I have used the cold bath during the paroxysm of epilepsy in various instances, but in general without

without the patient being roused to consciousness or fenfation, and without advantage. In one case of this difease, which occurred in the hospital practice, the paroxyfm returned periodically every afternoon. In this instance a cure was effected, by applying a cataplasm, formed chiefly of tobacco, to the scrobiculus cordis, about half an hour before the expected return, by which a powerful impression on the fystem was produced, and the paroxysm of epilepfy prevented. This practice repeated feveral days at the expected periods, destroyed the diseased catenation, and the cure was permanent.-In this mode of application, the peculiar effects of tobacco on the nervous system is speedily produced, and it is perhaps the fafest way of exciting them; because the application can be discontinued at pleasure; an advantage of great importance in the use of so deleterious a medicine, and which cannot be obtained when it is thrown into the flomach or rectum. I was induced to use tobacco on this occasion, from having observed it to succeed in the cure of obstinate intermittents, when applied in the same manner, previous to the expected accession of the paroxysm. In two recent cases I have employed this remedy in a different manner. Each of the patients laboured under general convulsion, and the paroxysm had returned fo frequently as to produce continued coma, and in the bye-standers an absolute despair of recovery. I ordered a decoction of half a drachm of tobacco K 4

tobacco in four ounces of water, to be thrown up as an enema. This powerful agent penetrated the fyftem to its very centre, roused the sensibility which the affusion of cold water, and other external impressions could not awake; excited sickness, vomiting, and profuse perspiration, and interrupted the convulfive actions, which have never fince returned. In each of these cases the recovery was altogether unexpected. The use of tobacco smoke thrown up the rectum in obstinate constructions of the alimentary canal, has been long established. I believe the effects may be obtained with more ease and greater certainty by employing the decoction. The reasonings used in explaining the operation of the affusion of cold water, are perfectly applicable in accounting for the effects of tobacco; and there is reason to hope that convulsions stopped in their paroxysm, by the one or the other remedy, are not in many cases liable to return.

In spasmodic diseases which rise into general convulsions, I have seen these convulsions rendered less frequent, and for a time seemingly diminished in violence, by medicines used during the intervals; but seldom effectually cured, except by means used during the act of convulsion, at the instant of its accession, or at the period of its expected return. Of the medicines employed during the intervals of epilepsy, the oxyd of zinc seems to me to have some

fome efficacy; but the digitalis purpurea still more. -In consequence of its powerful influence, I have used this last medicine in many cases of epilepsy, both in private and hospital practice; of twelve of which I have preferved particular registers. In every one of these, it diminished the number of paroxyfms, and the intervals were fo much lengthened, that in several instances we entertained hopes of a cure. It did not however finally fucceed in any one of them; and when, as the constitution became accustomed to its influence, the fits began to return, and we increased the dose of the medicine, its deleterious qualities, fo well described by Dr. Withering, prefented a danger greater than that of the difeafe, and obliged us to abandon it. On one occasion, we had an opportunity of observing, that pushing this medicine to the utmost extremity did not effect a cure. A female patient took it fo rashly as to produce the utmost danger of her life. Her pulse funk to thirty-two in the minute, her heat at the axilla to 89°; -her fight failed, her respiration became laborious, and inceffant vomitings threatened immediate diffolution. She was however recovered by the warm bath and opiate clysters, but as her ftrength was reftored the epileptic paroxyfms returned.

If it were my object to give a general view of the effects of the cold bath, or the affusion of cold water, water, in spasmodic affections, I might quote largely from the systematic writers of the last century, and I might detail various cases which have been furnished me by respectable living practitioners.-My friend Dr. Ford has mentioned to me the cafe of Mr. C. of Briftol, who was inftantly relieved of an obstinate stricture of the neck of the bladder of thirty hours duration, (during all which time not a drop of water had paffed) by placing his feet on a marble flab, and dashing cold water over the thighs and legs. The effect was instantaneous; the urine burst from him in a full stream, and the stricture was permanently removed. The common remedies, particularly opium and bleeding, and each of these very largely, had previously been used in vain. I record this case with the more pleasure, as it will recal fimilar cases to the learned reader's memory, and indicate a numerous class of diseases of the fame nature, in which this powerful remedy may be employed.

Cold water cannot be used as a drink during the paroxysm of convulsions, and of course we cannot shew the analogy between its external and internal use in these, as in other diseases. That its effects taken internally, are most falutary in a numerous class of chronic diseases, is however well known, though perhaps not acknowledged to the full extent of the truth. A considerable part of the virtue of mineral

mineral waters is doubtless to be attributed either to the diluting quality of the pure element itself, or to the invigorating effect of cold on the stomach, and through it on the system at large. The subject is however sufficiently extensive, as well as important, to require and to deserve a distinct examination. In hypochondriacal, hysterical, and dyspeptic affections, cold water taken internally has produced the most salutary effects. Hossman praises it in head-ach, whether arising from indigestion or some primary affection of the nerves of the head. The following case, extracted from his sirst volume, p. 475, will shew the use of cold drink in certain convulsive affections.

frigore, incidit in ingentes præcordiorum anxietates, cum tussi, virium lapsu, extremorum tremore, intercurrente borripilatione & interno æstu. Tertia die post, manus è pedes convulsivis motibus & subsultibus agitati; per horæ dimidium insistentibus.. Inde magis magisque increverunt convulsivi motus, ut dorsum & collum aspectu borrendis incurvationibus quassarentur, idque aliquoties quotidie. Sub paroxysmo pedes frigebant, respiratio erat creber, pulsus celer & durus, sensus tamen constabant. Post paroxysmum languebat corpus universum, appetitus vero cum somno erat integer. Adhibiti in consilio medici multi; exterorum etiam consilia conquissita Plurimi verminosam progeniem subesse causam rati, aliqui

aliqui per terrorem invectum esse causam putarunt. Sed data, tam ex illorum scientia anthelmentica, quam ex horum præscripto antepeleptica, adhibitis simul balneis, suerunt prorsus frustranea. Malum potius perstitit idem quin subinde factum acerbius. Tandem miser aquam sontanam frigidam, ad unam vel dimidiam mensuram, quotidie per vices bibere jussus est. Quo intra 14 dies motus isti convulsivi paullatim compositi, nec in hoc usque tempus iterum infestarunt." This simple and successul practice deserves to be imitated.

In the Appendix, N°. I, p. 17, I have mentioned the connexion between convultive difeases and infanity, and conjectured that the same principles are in many cases applicable to the treatment of both. This connexion has been since explained, and the subject very happily illustrated by Dr. Darwin, (Zoonomia, vol. ii. c. 3) who has arranged the whole of these diseases under one class, and thrown a new light on one of the most curious and obscure parts of pathology. The following case taken from the register of our Asylum, may serve as a new proof of this connexion.

June 2, 1796. L. I. aged 32, a man of very irregular habits of life, was admitted into the Afylum, in a state of surious infanity. His disease was supposed to have been brought on by excessive drinking.

drinking. As cases of infantity, including the whole method of cure, are seldom recorded, I will give this in detail.

Very powerful methods of coercion were required in the first instance, and the bowels were afterwards cleared by a saline purgative. At night he took eighty drops of laudanum, which produced imperfect sleep, and in the morning sollowing he was more wild than ever.

Having in some cases of infanity, found very extraordinary effects from such doses of opium as induced prosound sleep, I directed that five grains of this medicine should be given every four hours, till sleep succeeded. Next day (4th June) it was reported to me, that he had taken twenty-seven grains of opium in the twenty-sour hours. From the first five grains there was no apparent effect; after the second, he slept an hour and a half; after the third, three hours. Though the medicine had been twice repeated in doses of six grains each, he had no return of sleep—He appeared wild, agitated, and feeble, rather in a state of alarm than of sury. His pulse 100 and soft, with gentle and general perspiration.

It feemed not prudent to push the medicine farther ther; but having often in cases of this nature obferved the soporific effects of opium to come on, many hours after the medicine had been discontinued, he was directed to have milk as his food, and to abstain from medicines of every kind for twenty-four hours.

The report of the 5th is as follows. "He remained in a perpetual state of alarm till eight o'clock last night; after that hour he slept, but not foundly. He has taken nine pints of milk in all, having been very thirsty. It agreed with him till ten this forenoon, when he vomited confiderably. Since that time he has been more tranquil and feeble." In this state, I directed all medicines to be discontinued, and ordered him a pint of wine, with two pints of gruel daily, and fuch folid food as his ftomach could receive and digeft. This plan was continued till the 17th, when his strength was considerably restored, but his infanity as violent as ever. It was now directed that he should go daily into the bath heated to 95°, and that he should take half an ounce of the infusion of the digitalis, as recommended by my friend and kinfman, Dr. Currie of Chefter. (Mem. of the Medical Society of London, vol. iv.)

He continued on this plan with little variation till the 29th, the report of that day is as follows:

"Till the 25th the digitalis produced no obvious effect. On that day sickness came on and his pulse was hardly perceptible. The digitalis was therefore omitted. He has been regularly in the bath, and while in it, was once seized with diliquium animi, but the mental derangement did not afterwards abate. His pulse is now about 94, seeble and irregular.— He is a good deal emaciated, but more sierce than ever, he talks incessantly, with the utmost incoherence, and the association of his ideas seems wholly perverted."

In this state I ordered the digitalis and the tepid bath to be discontinued, as well as all other medicine. I directed a nutritious diet, but without wine or opium, and placed him in a situation to be as little as possible exposed to external impressions. For three days there was little change in his situation—his appetite improved, and he perspired much. At the end of this period, he had suddenly a lucid interval of several hours duration; but before I could witness it, he was again become surious.

At this time (the 16th July) I directed that he should take three drachms of peruvian bark, half a drachm of vitriolated iron, and the same quantity of aromatic powder, in the form of an electuary, daily; and that twice every day, he should be subjected to the tepid affusion for a minute or two at a time.

time, the water being poured gently over him of the heat of 88° more and a second reb tell no

perceptibles The diminish was theres

The following is the report of the 16th July: "The tepid affusion was continued till the 14th, but without any tranquillizing effect; the other medicines have been given regularly. On the 9th he became calm and rational: and flept the following night for eleven hours profoundly; but started fuddenly from this fleep, as wild and furious as ever. This state of mind continued twenty-four hours, when he again became tranquil, and continued so till he went to bed. He slept a few hours quietly, but awoke as before, under extreme agitation, and continued fierce and raving for twelve hours and upwards. Since this time, he has had a fuccession of paroxysms of infanity, alternating with periods of tranquillity of mind, of nearly the fame duration. In general, he goes to bed tranquil, and fleeps quietly till about three in the morning, when he awakes under the horror of some frightful dream; and this introduces his paroxysm of infanity and fury, which continues from ten to fourteen hours."

In this fituation I ordered an emetic to be administered to him late in the evening, in hopes that it might interrupt this beginning catenation. Being disappointed, I had again recourse to an opiate at bed-time; and this also proving inefficacious, I directed

rected that he should be awoke before the expected accession, and kept awake beyond its period of attack. By these means we disturbed the regularity of the alternations, but without any advantage; on the contrary, his lucid intervals became shorter in duration, and less complete.

Perplexed with these extremes, and keeping in mind the success of the cold bath in convulsive diseases, I ordered it to be tried on the present occasion. The infanity returning with great violence on the 21st, he was thrown headlong into the cold bath—He came out calm, and nearly rational, and this interval of reason continued for twenty-four hours. The same practice was directed to be repeated, as often as the state of infanity recurred.

The following is the report of the 30th: "The direction has been followed, and on the morning of the 23d, he was again thrown into the cold bath in the height of his fury, as before. As he came out, he was thrown in again, and this was repeated five different times, till he could not leave the bath without affiftance. He became perfectly calm and rational in the bath, and has remained fo ever fince."

This patient continued with us for fome time afterwards, bathing every other day, and taking the L oxyd

oxyd of zinc in small quantities. He never relapsed, and was discharged some time afterwards, in persect health of body and mind.

It has been contended by Dr. Darwin, that as convulfive diseases arise from inordinate action in the muscles, so maniacal diseases arise from the fame inordinate action in the organs of sense:-and as the muscles and the organs of sense are the proper instruments of the will, the diseased actions of the one and the other are by him arranged as one class. under the title of Diseases of Volition. But in the fystem of Dr. Darwin, the actions of the organs of fense constitute our ideas; a deranged action of these organs is therefore no other than what is usually denominated a derangement of mind; in other words infanity*. As therefore convulsion and madness, may be considered as the same disease differently fituated; and as the feat of the difease may be changed from various causes, and extended through various sympathies and catenations, we have thus an explanation of that connexion between convultive affections and infanity, and particu-

^{*} If, as is generally believed, the motions of the organs of fense rather excite ideas than constitute them, the reasoning of Dr. Darwin is not affected by this change of terms—That madness is a disease of the body, cannot be disputed, whether we admit motions in the organ of sense or not.

larly of their alternations with each other, which have been observed in every period of medicine.-That feveral of the modes of treatment applicable to the one, are also applicable to the other, has long been known. The arnica montana, the oxyd of zinc, the digitalis purpurea, and opium, are medicines used equally in convulsions and in infanity. The case which has just been related extends this analogy, and is the more striking in this point of view, because the success of the cold bath was almost instantaneous, and occurred in the paroxysmi of infanity, in the same manner as has been before observed of its operation in convulsion.

the cold bath in the parexylm of madness, or of con-

It is supposed by Dr. Darwin, that the motions excited by volition and by fenfation, proceed in different directions; the first passing from the central parts of the fystem towards the superficies, the fecond from the superficies towards the central parts. Hence he concludes that they are opposed to, and mutually destructive of each other. To increase the force of voluntary actions must therefore be to leffen positively, as well as relatively, the actions that arise from sensation, and the converse of this proposition is equally true. Whatever faith we may give to the hypothesis of Dr. Darwin, his conclusion feems to be supported by a great variety of phenomena. That the higher degrees of volition and fensation are incompatible with each other, is a L 2 truth

truth of great importance in every part of the science of life, whether moral or phyfical*. It was the doctrine of another great physiologist (whose praise will now be heard with approbation, fince he is beyond the reach of praise) "that no two actions can take place in the same constitution, nor in the fame part at the fame time;" + and though he applied this doctrine chiefly to show the incompatibility of different diseased actions with each other, yet it feems applicable also to explain the operation of a great part of the more powerful remedies. The position of Dr. Darwin resolves itself into this more general principle of Mr. Hunter, and the operation of the cold bath in the paroxyfm of madness, or of convulfion, receives an eafy explanation under either doctrine. We may fay with Dr. Darwin, that the powerful stimulus given to sensation, destroys the previous diseased volition, or in the more general chas from the Superficies towards the central parts.

Hence he concludes that they are opposed to, and

^{*}The reader who is not already acquainted with this part of Zoonomia, will not be able to understand how convulsions, which in common language are considered as involuntary, are classed as diseases of volition. He is therefore referred to the work in question for a full explanation. In the mean time we may observe, that Dr. Darwin considers all motions as voluntary which have originated in volition, though from their violence no longer subject to the will. Volition is used by him in a sense somewhat different from its common acceptation.

⁺ John Hunter. Treatife on the Blood: Introd. p. 3.

and more fimple language of Mr. Hunter, that the fudden application of cold water to the furface, produces an action of the fystem with which the previous difeafed action is incompatible. In order however to the fuccess of the cold bath in either of these species of disease, it will be readily perceived, that it is necessary that no considerable læsion of structure shall have taken place, and that the morbid actions be not fo strong as to produce infenfibility to impression, and shall not have been fo often repeated as to form indiffoluble catenations. It feems also necessary that the diseases in question shall be such as are susceptible of general paroxysms; a circumstance not so easily explained. Finally, it is necessary that the digestion shall not have been much impaired, or the vigour of the circulation much debilitated, lest the action of cold be too strong for the living powers. Under these restrictions, the cold bath, according to my experience, may often be applied with advantage, and always with fafety, in convulfive difeases, and in infanity; for though in these diseases, even in their more violent forms, the temperature of the body is little, if at all, increased, yet the actual heat is retained with great tenacity, as I have found by many obfervations.

The refiftance which the maniac opposes to cold, he opposes also to contagion, and to other noxious powers, powers, as has often been observed with surprise and admiration. This refistance feems to belong to every species of increased voluntary exertion, where the functions of digeftion and nutrition are not impared. The illustration of this position would not be difficult, and the conclusions to be deduced from it are in a high degree interesting and important; but a discussion of the subject would lead us too deep into the general doctrines of life.*

^{*} Dr. Marcard, of Pyrmont, in his work on the effects of warm and cold baths, entitled, Ueber die Natur, und Gebrauch der Bader, lately published, has quoted my paper on the effects of the cold bath in convulfive difeafes (Appendix No. I.) and it appears, that he does not affent to my conclufions. Dr. Marcard supposes indeed, that the cold bath may be useful in convulfive diseases, where there is no local affection stimulating the system, but he contends that it must be hurtful in cases where convulsions originate in stimuli, as worms, or crudities in the alimentary canal, teething, &c. It does not appear to me that Dr. Marcard has understood my paper perfectly, and I do not feel myfelf disposed to yield my experience to his speculations. The doctrine which Dr. Marcard feems to maintain, that where one stimulus is already acting, the application of another must be injurious, is contrary to experience, and as I apprehend, to the laws of the animal occonomy. The work of Dr. Marcard is a valuable one, and deserves to be translated; but it seems to be too full of those general and sweeping observations, which though nearly banished from other sciences, continue in the writings of phyficians, and ferve at once as a proof and a cause of the impersection of our art. I fear this volume will not be found free of them.

near paylicing in every period of the world, impa-

CHAP. XIV.

General view of doctrines respecting sever.—Hippocrates—Galen, &c.—The ideas of the author respecting the nature of this disease.

THE great difficulty which men have in all ages experienced in the acquisition of knowledge, has arisen from the promptitude of the human mind to decide in regard to causes. To the weak and ignorant, prefumption is as natural, as doubt is intolerable, and with fuch, belief is almost always a creature of the imagination. Nor is this peculiar to weakness and ignorance: to retain the mind unprejudiced and undecided in the investigation of striking and interesting phenomena, till by the painful steps of induction, the hidden cause is revealed, is an effort of the most difficult kind, and requires the highest and rarest powers of the understanding. The records of every part of science, bear ample testimony to this truth, particularly the records of medicine, and in a still more especial manner that part of medicine which treats of the nature of fever. The most eminent physicians in every period of the world, impatient of observing and delineating, have been eager to explain and even to systematize; and the science of life owes its corruptions more to the misapplication of learning, than even to the dreams of superstition. Passing over the theories derived from the splendid sictions of the Greek philosophy, various are the salse doctrines introduced into medicine in modern times, and from the more certain branches of science. The principles of mechanics, of chemistry, of magnetism, and of electricity, have in succession been employed to explain the motions of life, and have served only to mislead and to deceive.

Hippocrates who lived at too early a period to be acquainted with the collateral branches of science, studied life and disease in the book of nature, and had the merit of an original observer. Perceiving the increase of heat to be the most remarkable symptom in sever, he assumed this for the cause, and sounded his distinctions of severs, on the disferent degrees of the intensenss of this heat. He had not an instrument that could measure this exactly, and necessarily trusted to his sensations. In forming his diagnosis, he placed his hands on the breasts of his patients, depending more on the degree of heat than on the state of the pulse, the nature and connexions of which he did not understand.

fland*. His practice appears to have been natural and judicious, and founded on his theory. He directed linen dipped in cold water to be applied to the hottest parts; drew blood away both by cupping-glasses and the lancet; and administered cold water and cooling drinks, particularly barley water and honey. This simple practice, will acquit the Coan sage with the candid and reslecting, of many of the idle theories which have been imputed to him, and which doubtless were the offspring of after times.

In the days of Galen, philosophy and science had made some advances, and the corruptions which he introduced into medicine from those sources are well known. Preserving however the doctrine of his master, that heat was the cause of sever, his practice sounded on this notion, was in some respects bold and successful, as was before observed. To him followed the Arabians, who still retained the original doctrine of Hippocrates, but with new modifications and corruptions, by which the real nature of sever was still farther obscured.

In our own country, and in the last century, arose Sydenham. He also was an original observer, and to him medicine is doubtless much indebted.

ndy affuded) as well as other chemi-

That

^{*} Riollay's Introduction, &c. p. 9.

That he recorded symptoms with great accuracy; that he established many important facts as to the treatment of disease, and that he was a more cautious reasoner than his predecessors or contemporaries, are points that are indisputable. But, though he affected not to theorize, he was a theorist in every page of his works, and though he founded his reasonings on what appeared to him self-evident principles, yet time has proved them to be defective and fallacious. It was the postulate of Sydenham, that every difease is nothing else but an endeavour of nature to expel morbific matter of one kind or another, by which her healthy operations are impeded. In this endeavour she is not to be obstructed but affifted, and the process carefully watched and promoted by which she accomplishes her purpose. By one or other of the emunctories, this is finally effected, and till it be effected, health cannot be restored. Under this general notion, the inordinate actions of fever are perpetually compared to the motions of fermentation, by which nature feparates the vitiated particles from the blood previous to their expulsion. The doctrine of Sydenham may be traced to remote ages; under his fanction it spread over Europe, and has descended in one form or other to our own times. Sydenham's doctrine introduced the fystem of Boerhaave, (to which we have already alluded) as well as other chemical theories of leffer fame, and the erroneous modes

modes of treatment to which they gave birth. The inordinate heat which Hippocrates confidered as the cause of sever, in the eye of the chemists appeared naturally enough, a necessary, and within certain limits, a salutary consequence of the process stirred up by nature to expel the disease!

The dawn of a juster pathology of fever is to be found in the works of Hoffman. Though he also undertook to be the interpreter of nature's intentions, he contemplated her process in fever with more fagacity, and rejecting chemical and mechanical analogies, endeavoured to discover the cause of fever in the peculiar nature of the vital motions. He supposed the noxious cause producing sever (in the language of the schools the remote cause) to operate first on the living folids, occasioning a general spasm of the nervous and fibrous system, beginning in the external parts and proceeding towards the centre. In consequence of this a contraction of the veffels of the extremities must of course take place, impelling the circulating fluids in an increafed ratio on the heart and lungs; which stimulating these organs to increased action, the fluids are thereby repelled towards the extremities, and thus the phenomena of fever are produced. There are therefore two distinct sets of motions in fever, the first from the extremities towards the centre, arifing immediately from the spasm, and accompanied by a fmall small pulse, anxiety and oppression; the second, from the centre towards the surface, which is the effort of nature to resolve this spasm, and marked by a sull strong pulse, and increased heat. The first of these sets of motions are baneful, and sometimes satal; the second are medicinal and salutary. By these views the physician is to be directed in counteracting the morbid actions, and assisting the sanative process of nature*.

Since to theorize feems to be the irrefiftable propensity of men of genius, this theory of the celebrated Hossiman, which accords with appearances, and has a considerable connexion with practice, may be allowed the tribute of applause.

The system of Hossiman, produced that of Cullen, which may perhaps be considered as the prevailing doctrine at present, when however theory of every kind has lost much of its weight on mature understandings.—According to Hossiman, the first effect of the remote cause of sever is the spasm, producing a re-action, as has already been mentioned. Dr. Cullen introduced a previous link into the chain—He contended that the first effect of the noxious essential that the sirst effect of the debility,

^{*} See Hoffman's Works, vol. i. tom. ii. page 10;-Gene-

debility, affecting the fenforium commune. To this debility he attributed the spasm, and to the spasm the re-action of the heart and arteries; which re-action continuing till the spasm is resolved, removes the debility and the disease. According to Hoffman the spasm belongs to the class of motions that he denominates baneful; but Dr. Cullen presumes it to be falutary, and therefore ascribes it, in the language of the schools, to the vis medicatrix natura. It is not my purpose to appreciate these changes introduced into the theory of Hoffman, or to attempt to estimate with precision the Cullenian doctrine of fever, as it arranges phenomena, or applies to practice. Debility of a peculiar kind, spasm, and reaction of the heart and arteries, feem all of them links in the chain of fever, and in the first lines of Dr. Cullen, the hiftory of the difease is recorded with extraordinary minuteness and accuracy. I bow with respect and sensibility to the genius of this illustrious man; but his doctrine seems erroneous, in as far as it enters into the supposed intentions of nature, and defective, in passing over the morbid heat, and the morbid affociation, which form the fuccessive links of the chain. Cullen, formed according to Dr. Brown the effence

This doctrine of fever, as well as all the other doctrines of its celebrated author, was affailed by Dr. Brown. Affuming the existence of an unknown principle, as inherent in the living fibres, to which

which he gave the name of excitability, he explained all the phenomena of life and difease, as dependent on this principle, acted on by stimuli; which stimuli, applied in the due proportion, produce the just degree of excitement, that is, the state of If the stimuli are diminished below the health. healthy proportion, he supposed the excitability to accumulate; if encreased beyond this proportion, he supposed it to be expended; and on these different conditions he attempted to found a theory of disease. Diseases he divided into two classes, Sthenic and Asthenic, or diseases of increased and of diminished excitement, in the last of which classes he placed typhus, the fever of which we treat. In the excitability, Dr. Brown admitted of no change, except in regard to quantity; in the excitement, no variation, except in regard to strength; and in all universal diseases he supposed the whole system to be equally affected.* Having therefore affigned to fever its place in his feries of descending excitement, he refused to inquire into its symptoms, or to enlarge on its treatment. It is evident that fuch a theory could embrace none of the characteristics of the disease. Debility, the first link in the chain of Dr. Cullen, formed according to Dr. Brown the effence of fever. The existenc of spasm he denied, re-action of its celebrated author, was pikifled by

^{*} This was his general position; it is true he relaxed a little from it in his explanations.

he derided, and the morbid heat and morbid affociation, he wholly overlooked. It is not however to be disputed that his general views of disease had a salutary influence on the practice in typhus, by encouraging a more liberal use of wine and opium, remedies of the utmost importance, and now universally adopted.

The various modes of action of the living principle, or to use his peculiar language, of the sensorial power, which were unknown to Brown, have been observed and illustrated by the author of Zoonomia.

In the theory of fever delivered by Dr. Darwin, the influence of morbid affociation is very fully explained. I am not however disposed to adopt this theory, which the learned and candid author himself considers as incomplete, because it assumes the truth of his peculiar doctrines, to which it cannot be expected that a hasty affent should be given—considering their vast extent and importance, and considering also that many other fabrics of genius of the same kind, have crumbled down into the sand of which they were formed.*

Dismissing

v, to be applied to it, and the

^{* &}quot;What I have thus delivered, I beg to be confidered "rather as observations and conjectures, than as things ex"plained and demonstrated; to be considered as a foundation

Difmissing therefore as far as it is possible, all theories from the mind, let us briefly confider the process of nature in fever, and examine what inferences may be fafely drawn from the fuccession and catenation of the leading fymptoms, and the remedies which experience has pointed out as most fuccessful. Our observations must of course be confidered as applying to the typhus, or low contagious fever of Britain, more particularly, but generally to all idiopathic fevers; and if they should be found as ower, which were unknown to Brown, have

[&]quot; and a fcaffolding, which may enable future industry to erect " a folid and a beautiful edifice, eminent both for its fimpli-" city and utility, as well as for the permanency of its mate-" rials-which may not moulder, like the structures already " erected, into the fand out of which they were composed; " but which may frand unimpaired like the Newtonian philo-" fophy, a rock amid the waste of ages !" Zoonomia, vol. ii. p. 625. Much as I admire the profound reflection, and fingular ingenuity of Dr. Darwin, the position common to him and Dr. Brown, that the fenforial power, or excitability, is a fubstance which accumulates and diminishes in the inverse ratio of the stimulation, on which his doctrines of fever are founded, has not my affent. On the contrary, it appears to me, that whatever the nature of the vital energy (the fenforial power or excitability) may be, it does not act in a way that admits the relations of quantity to be applied to it, and the contrary doctrine, which introduces into the fystem of Dr. Darwin fo many epicycles, feems to me the mortal part of his immortal work. I am aware that Dr. Darwin holds the doctrine mentioned with certain modifications; which however do not appear to me fatisfactory.

little fatisfactory in their refult as the conclusions of others, the learned and candid reader will not, it is hoped, suffer his opinion of the accuracy of the author's facts to be affected by the fallacy of his reasonings, but extend his forgiveness to one other abortive attempt to illustrate the nature of sever, an important, but difficult and long-contested subject.

Medical science has not ascertained the various remote causes which may produce fever, and I avoid this inquiry, on which I have nothing new, that is fatisfactory, to offer*. The first symptom of the disease is a sense of debility or languor, very distinctly expressed on the countenance, followed by paleness, cold, and tremor; the functions of the mind are weakened in a correspondent degree with those of the body, and all the actions of the system feem enfeebled. This general debility is peculiar in its kind. The fymptoms refemble those produced by inhaling certain gases whose properties are known; and this circumstance serves to strengthen those views of the nature of contagion which the M new

^{*} In confining the remote causes of sever to contagious human essential, and to marsh miasmata, Dr. Cullen probably simplified too far. Cold, under certain circumstances, seems to produce sever, and the constitution seems at times to take on sever from unknown and internal causes.

new chemistry has suggested. They seem also to have a refemblance, though a more remote one, to the effects of certain poisons operating on the stomach, or introduced into the circulation by the abforbents. The remote cause of fever itself, may perhaps be confidered as a poison, acting directly on the fenforium commune. Where this poison is peculiarly concentrated and malignant, or where the fystem is much debilitated, the powers of life are fometimes oppressed and extinguished in the first stage of the disease. In general however, a re-action or resistance commences; the heart and lungs are roused into increased exertion by the pressure of the accumulating fluids, and strive as it were, to repel them back on the furface and extremities; while a spasmodic, or morbid stricture of the extreme veffels opposes the reflux of the fluids, and thus maintaining the inordinate pressure on the centre, excites the heart, arteries, and lungs, to still more violent exertions. In this contest the stomach is usually brought into sympathy, and nausea and vomiting are often induced.

This constitutes what appears to be a struggle between the living energy, and the morbid cause—between the power of the centre, and the resistance of the extreme vessels—in the course of which a degree of preternatural heat is generated, and the phenomena of the hot stage are produced. When

the powers of life prevail, the stricture on the extreme veffels, and on the exhalants of the skin, at length gives way, and a profuse perspiration being poured out on the furface, the heat is carried off, and the febrile fymptoms subside. Without entering into detail, this may be confidered as a general view of the fingle paroxysm. It often however occurs that the folution of the spasm on the extreme vessels does not take place, or is incomplete, and that the morbid heat is not carried off. In this case the morbid actions go on, and the difease runs into continued fever. For some time the spasm on the extreme vessels remains, and the heat is preternaturally great; but these symptoms do not continue through the whole course of the disease. contrary, the heat towards the latter stages, will fometimes be found as low as the natural standard, fometimes lower, and the capillaries of the skin be completely relaxed; while the inordinate action of the heart and arteries continues, being carried on by the morbid affociation produced in the course of the disease, which retains its influence in this as in other cases, after the cause that produced it ceases to operate.

To this general view of fever, it is probable that little objection will be offered, by those who are familiar with the disease. But exceptions may be taken to the language as not sufficiently clear of theory,

M 2 against

against which the author had declared. It may be faid, for instance, that in applying the active forms of speech to the re-action or resistance of the constitution, we feem to run into the Sthaalian doctrine, or at least to assume the existence of the vis medicatrix naturæ of Dr. Cullen; and that to enumerate spasm as a link in the chain of fever, is in fact to admit the leading peculiarity of his fystem. To this it may be replied, that there is no subject on which the imperfection of language is fo much perceived, as in our attempts to describe the phenomena of life. In the strictness of speech we properly employ the passive mood of our verbs, in recording the phenomena of inanimate matter, and confine the active mood to those vital motions, which are accompanied with consciousness or design; but there are various actions peculiar to life which are not accompanied by consciousness, or subject to the will, and which depend on laws wholly diftinet from those which regulate inanimate matter. If we borrow the phraseology and the mood in which we speak of inanimate matter in recording these actions, we are apt not only to form indiffinct conceptions of their nature, but to apply the principles which regulate inanimate motions to the motions of life. This is an obvious, and as experience teaches us, an ample fource of error. If, on the other hand, we employ the active forms of speech, these immediately fuggest consciousness and design, and the

the imagination forms to itself an ideal being, as directing these actions, in whom consciousness and defign may refide. This also is a natural and fruitful fource of error.-Both these extremes we would avoid. But the penury of language obliges us to use one or the other mood in treating of the phenomena in question; and on reflection it appears, that in our present ignorance of the first principles of life, a clearer notion will be formed of the nature of those motions in which the whole system sympathizes, destitute though they be of consciousness and defign, by the occasional use of the active forms of speech, since they have a much stricter analogy with those vital actions, to which these forms are justly applicable, than with the motions of inanimate matter. Confiderations of this kind may have influenced the late John Hunter in the use of his terms, which are chiefly active, and which, though far from perfect in themselves, are more happy than those of former physiologists, as his doctrines of life were more original, more unprejudiced, and in general more accurate and profound. blood flows imprediardy back into

Debility of a peculiar kind, is then the first operation of the remote cause producing sever—the necessary consequence, or as some contend, the concomitant esset, is a spasm, or contraction of the arteries; but more especially of the extreme vessels, and the capillaries of the surface—hence M 2 follows

follows an accumulation of blood on the heart and lungs—the re-action of those organs—the generation of morbid heat—and of morbid association. On each of these links in the chain of fever, let us offer a few remarks.

- r. It has been afferted by some inveterate theorists, that the debility in fever has nothing in it peculiar. It is enough to reply, that simple debility often exists even in extreme degrees, without producing the phenomena that constitute sever.
- 2. It has been afferted, that a spasm on the extreme veffels does not exist in fever, or that if it does exist, it may fafely be difregarded. Those who hold the first opinion, suppose the paleness and shrinking of the surface to arise merely from the enfeebled circulation, by which the blood is not propelled into the extreme parts; and doubtless this has its effect. In fyncope, however, where this paleness and shrinking arise from pure debility, as soon as the heart and arteries recover their energy, the blood flows immediately back into the extreme veffels as before. That this is not the case in fever, the most cursory observation will convince us. After the re-action of the heart and arteries has been some time established, the blood is indeed propelled through the veffels of the furface that circulate the undivided fluid; but the capillary arteries that **feparate**

separate the aqueous part remain constricted, as well as the orifices of the skin that pour it out on the surface.

3. It is however contended, that this stricture may be difregarded, fince it is now very generally admitted, that the perspirable matter is not excrementitious, and fince we find that its obstruction in other circumstances, and from other causes, produces little or no injury to the constitution. A little reflection will serve to refute this position. Spasm on the extreme veffels must necessarily affect the condition of the heart and lungs, from the increafed pressure of the circulating fluids on these organs: it must also affect the stomach not only from the connexion of that organ with the heart and lungs, but from its direct sympathy with the surface. - Dr. Cullen has put these points beyond controversy. But this is not all; the obstruction of perspiration-of profuse perspiration—in the paroxysm of fever, obstructs the process by which the constitution expels the morbid heat, and thus leaves the fystem under the influence of a general stimulus of the most powerful nature. It was a position of the celebrated Boerhaave, that the morbid heat in fever, being a fymptom only, might therefore be difregarded .-But can we suppose, that a heat fix or seven degrees greater than that of the blood in health, however generated, will not have the most important effects on the fystem, and if it stands in the relation of effect to the preceding symptoms, that it will not operate as a cause on those which succeed?—Doubtless this morbid heat re-acts on the heart and arteries; stimulates the inordinate exertions by which it was excited; irritates the spasm of the extreme vessels which prevents its being expelled; and prolonging the increased action of the heart and arteries, establishes a morbid association, which carries on this action after the heat itself has subsided, and the spasm relaxed.

4 What then, it may be inquired, is the actual state of the animal heat in the different stages of the paroxysm of intermittent, as measured by the thermometer? In different trials it feemed liable to confiderable variety, but the following is the general refult, which however the reader will receive as subject to the corrections that a more extensive experience may fuggeft. In the cold ftage, the heat is diminished, not on the surface only, as some have imagined, but very probably over the whole fystem. I have found it under the tongue, and at the axilla, as low as 94°, 93°, and 92°; and on the extremities many degrees lower. At this time the ftomach feels cold, and univerfal tremor takes place. Warm, and even hot liquids, are highly grateful, and the warm bath, heated to 100°, more grateful still. The drinking of warm liquids should be freely freely indulged in, and there is reason to believe, that immersion in the warm bath (as practifed by Galen) would be found as falutary as it is grateful. By degrees the pulse increases in frequency and force, and the heat begins to rife, but with much irregularity. It doubtless accumulates first in the centre of the fystem, and from the centre is pushed along with the blood, towards the furface. This process does not however go on in any regular progression, but like almost all the other animal processes, with alternate increase and relaxation. At times it appears retrograde; and even when it is so far advanced that the heat taken at the axilla, and under the tongue, is greater than the standard of health, a slight acceffion of external cold will produce a general chillness, and bring back the oppression on the heart and lungs. On the furface itself the restoration of heat takes place with no regularity.-Certain parts are heated first. In certain parts the heat is above what is natural, while in others, it remains below this flandard; and hence arises that mixed sensation of heat and cold, which every one acquainted with fever must have experienced in the transition from the cold to the hot stage of the paroxysm.

The irregularity with which the heat of the furface is restored, probably arises from this circumstance, that two causes conjoin in producing this effect—the reslux of warm blood from the centre,



and the sympathy between the stomach and skin. To this last is attributed the circumscribed heat of certain parts of the furface, while the reft remains cool, this local heat being excited by the increased action of the veffels of these parts, awakened by the restored heat and tone of the stomach, which in other circumstances produce similar effects. In general the fense of cold predominates, even after a morbid heat has taken place at the axilla, under the tongue, and in different parts of the abdomen and thorax: because a considerable part of the surface, and the whole of the extremities, still remain below the natural temperature, and the fenfation which this produces, arifing from parts of great fenfibility, overcomes the fenfation of heat from the deeper feated parts, whose fensibility is so much inferior. At length however the heat of the furface becomes general and uniform, rifing to 102°, 103°, 104°, and fometimes 105° of Fahrenheit. I have not seen it higher than this last degree in the paroxysm of intermittent, and feldom in continued fever; though different authors speak of febrile heat four, or even five degrees higher. The fensation of heat is now strong and steady, and the accession of external air does not produce a return of chillness as before.-This is the time for the affusion of cold water, as has already been mentioned. The fensation of heat is most powerful on the extremities, particularly on the palms of the hands and the foles of the feet .-This

This arises in some degree from the great sensibility of these parts, but in some degree also from this circumstance, that in their natural state they are two or three degrees cooler than the trunk of the body, and cannot be raifed to the general temperature of health without an uneafy fense of heat, which is frequently relieved by a partial perspiration. When raifed to the general fever-heat, their temperature is therefore increased in more than a proportional degree beyond its usual standard.

After remaining some time in this hot stage, the fkin begins to relax, and to become fofter and fmoother to the touch; infenfibly a moisture takes place, terminating generally in profuse sweat .-Neither however does this process go on regularly; after the moisture commences, it sometimes goes off, and a dry and burning state of the skin returns: again the fweat commences, and when it becomes general and continues steady, it is critical, that is, it terminates the paroxysm. As it commences, the heat declines, though at first very slowly: as it proceeds, the heat subsides gradually; and when it terminates, the temperature of health is restored. The finking of the morbid heat is however a good deal regulated by the quantity of the bed-clothes, and the closeness with which the body is enveloped .-Where the whole covering that was laid on during the cold stage, is kept on during the whole paroxysm Augitavriide tooromus suom saud das (as (as often happens through a mistaken wish to force the perspiration) the heat is carried off with dissiculty, and it requires a very profuse perspiration to reduce it to its natural standard. On the contrary, if the clothes be lessened after the hot stage is fairly established, the heat passes off with less difficulty, and the perspiration requisite to remove it is much less profuse.

These details will not be thought superstuous or tedious by those who consider, that after all that has been written on the subject of sever, thermometrical observations on the rise and progress of sebrile heat, are in a great measure new*.

elsee recommunity account to profite (west,-

5. Whether it be owing to the nature or force of the contagion, to the state of the constitution, or to some other unknown cause, the re-action of the system does not always resolve the spasm or morbid stricture on the extreme vessels. When this stricture is not resolved, the sweat is not essued, the morbid heat is not carried off, and the diseased actions run into continued sever.

emperature of health is restored. The

If

^{*} I intended to have introduced here one or two registers of the heat and pulse, taken every half hour during the paroxysm of intermittent, but this is delayed till I am enabled to speak from more numerous observations.

If a person is confined in a hot room, or in the hot bath, till his heat rifes four or five degrees above the natural standard, his pulse will be found of a feverish rapidity; wandering pains will foon be felt over the body; languor, lassitude, and at length great debility will take place, with most of the fymptoms of regular fever. It is evident that these fymptoms cannot be expected to go off till the inordinate heat is removed, and if the person remain fome time in the heated medium, he will find that the inordinate action of the heart and arteries continues after leaving it, and even after his own heat has fubfided to its natural standard. This depends on a principle peculiar to life, to which the name of habit or affociation has been given and which extends its influence to all the vital phenomena, whether intellectual or corporeal.

In fever this morbid heat does not arise from the surrounding medium, but from certain motions in the system itself. But however generated, a heat sive or six degrees above the natural standard must be a powerful agent, and it cannot be expected that the diseased actions should subside under so strong a stimulus. The rapidity of the circulation, and the labour of respiration, are consequences of a heat of this degree, from whatever cause arising, and must continue till the heat is reduced.

We may here observe, that it is not by any means easy to make an accurate comparison between the state of the fystem under the heat of fever, and under an equal degree of heat introduced from the furrounding medium. Because it is in fact very difficult to raise the heat of the body in a state of health, to the highest degree of fever heat, either by heated air, or heated water. The experiments recorded by Sir Charles Blagden, prove this in respect to a heated atmosphere; the heat as it is accumulated in the fystem, being diffipated by profuse perfpiration, or by fome other living process, of which perspiration is a concomitant effect. I have found the fame difficulty in heating the body in the hot bath, for there also the heat is with difficulty increased beyond 100° or 101°, when fweat, as is usual, breaks out over the furface. Even in these degrees the greater part of the fymptoms of fever will be felt. The heat however is moderated, and, when the temperature of the external medium is reduced, it is carried off by the discharge from the surface. But in fever this is prevented; for the peculiar debility induced by the remote cause, occasions, or is attended by, a spasin, or morbid stricture of the capillaries of the furface and of the skin itself, by which the insensible perspiration is prvented from increasing in proportion to the heat, and the fensible perspiration obstructed. Hence the difficulty of reducing the heat in fever; and thus it appears, that the spasm

on the extreme veffels, which some of our latest physiologists have wholly neglected, is perhaps the most important peculiarity of the disease.

It may be observed, and it furnishes a subject of interesting inquiry, that this stricture, which often gives way under a moderate re-action, maintains itself with rigidity where the re-action is great. Thus where the heat rises to 99°, 100°, or 101°, the orifices of the skin often relax, and carry it off by increased perspiration; while at 104° and 105°, with a proportional force of circulation, they are in general obstinately constricted. This is very striking in fever, but not peculiar to it. Even in the state of health, while a moderate, gradual, and general stimulus from heat opens the pores of the skin, a fudden and confiderable increase of heat constricts them; the violence of the stimulus, in this, as in other inftances, occasioning a morbid refistance of the living fibre on which it acts*.

But, it may be justly remarked, that after fever is established, the reduction of heat to the natural standard,

^{*} This refistance to inordinate stimuli extends to every part of the living system, and belongs to every species of vital action; it characterizes the mental as well as the corporeal functions, and a full consideration of its influence might afford the most important instruction, not to physicians only, but to moralists and legislators.

standard, however desirable, is attended with difficulty and hazard, fince the patient cannot in general bear the continued exposure to external cold necefsary for this purpose.

Even in the hot stage of fever, if the patient be exposed naked to a cold atmosphere, though he supports the application for some time with comfort, and even with advantage, yet as the heat of the furface approaches the standard of health, a sudden sense of chilliness comes on, with a return of oppresfion on the heart and lungs, and all the fymptoms of the first stage of the paroxysm. The length of application, and the degree of external cold producing this effect, are indeed various in various constitutions, and in different stages of the disease: the degree of reduction of the animal heat at which the chilliness commences, is also uncertain; but the general truth of the position laid down is established on ample experience, and will not, I believe, be controverted. In cooling the furrounding atmofphere in fever, or in any other continued application of cold, care must be taken to keep it within the limits in which it is grateful to the sensations; and within these limits the cool treatment, as far as my observation extends, is uniformly advantageous. But though it moderates re-action, it does not remove the spalm on the capillaries, or break the catenation or habit by which fever is prolonged.

This

This is effected by the fudden affusion of cold water over the naked body; and the mode of its operation will now be eafily understood. The fudden, general, and powerful stimulus given to the syftem, dissolves the spasm on the extreme vessels of the furface, and of the various cavities of the body: the fudden and general evaporation carries off a large portion of the morbid heat accumulated under the fkin; and the healthy action of the capillaries and exhalents being restored, the remaining superfluous heat passes off by sensible and insensible perspiration. The stimulus of morbid heat and of morbid stricture being removed, the inordinate action of the heart and arteries fubfides, and the haraffed, and toil-worn patient finks into that peaceful fleep which nature has provided as the folace of our pains and forrows, and the restorer of our strength.

If fuch be the explanation of the diminution of the heat, and of the frequency of the pulse, which follow the affusion of cold water on the surface, it may here be inquired how the diminution of the thirst is accounted for. Are we to suppose, that a portion of the liquid is taken up by the absorbents of the surface, as is generally believed to be the case, during immersion in the bath? I apprehend not. The absorption in the bath, if it occur at all, (of which there is no sufficient evidence) has been greatly exaggerated, and the diminution of thirst

after the cold affusion is too speedy to be ascribed to fuch a cause. It admits of a more satisfactory explanation. The morbid stricture on the extreme vessels, to which we impute so much, is not confined to the capillaries of the furface, but extends, as far as we have an opportunity of observing, to the capillaries of all the cavities of the body. The fense of thirst is produced by the heat and dryness arifing from the morbid stricture of the exhalants in the inner furface of the mouth, fauces, and stomach; and this stricture being removed by the affusion on the surface of the body, from the sympathy between the furface and these cavities, so often alluded to, the healthy action of the exhalants of these parts, as well as those of the skin, is restored; and their moisture being poured out, the heat and dryness are removed, and the thirst diminished.

This indeed feems to be the way in which thirst is relieved in the first instance, by liquids taken into the stomach itself, where the effect is too sudden to arise from the absorption of any portion of the liquid into the mass of circulating sluids. We cannot indeed perceive this effect on the exhalants of the stomach, but we may reasonably suppose it to take place where we observe the swallowing of drink to be sollowed by a relaxation of the exhalants on the skin, which must proceed from their sympathy with those of the stomach previously relaxed; and it may illustrate

trate and support this position, that when liquids, swallowed even in large quantities in the hot stage, do not produce a fenfible increase of perspiration on the Ikin, the relief they afford to the thirst is momentary only. In fuch cases we presume the stricture on the exhalants of the stomach has not been resolved. The relief from thirst afforded by liquids taken into the stomach is in general however more effectual and complete, than from affusion of water on the surface, because when the stricture on the capillaries is removed, a rapid abforption takes place from the stomach, and the circulating fluids are diluted. But on the furface of the body this absorption is doubtless flow and imperfect, and it will be afterwards shown, that there is reason to doubt whether it takes place at all. It was remarked that the affusion of tepid water on the furface, though effectual in diminishing heat, (page 70) is not followed by the entire ceffation of fever, as is often the case after the cold affusion. The cause of this is obvious-the coolness produced by the tepid affusion, arises from the increased evaporation, and continues only while this continues. The stimulus given by the tepid affusion is comparatively slight and transient-it does not refolve the spasm on the surface, or on the other cavities of the body, and of course it does not destroy the morbid catenation. The heat therefore speedily returns, when the remedy is withdrawn.

N 2

This explanation of the operation of the affusion of cold water seems to me satisfactory. The cases which have been related show, however, that it is only in the early stages of sever that it produces a solution of the disease. In the after stages, though uniformly advantageous while the morbid heat continues, its effects are not so decisive, the morbid actions having acquired the force of habit, or in other words being carried on by the associations or catenations that have been produced.

What then ought to be the indications of practice in fever? To diminish the cold in the cold stage; to moderate the heat in the hot stage; and to resolve the stricture on the extreme vessels by which the heat is retained, and the re-action prolonged; and where the inordinate action of the vafcular system continues after these objects have been attained, to support the powers of life till the morbid affociations, or habits of action, gradually die away, from the removal of the causes by which they were introduced. In addition to these general indications, it will be effential to fecure the proper action of the bowels; and in every case to unload the alimentary canal of its morbid contents; whether these contents have become diseased through the action of general fever, or, as there is reason to believe in some of the fevers of the warm climates, be the remote cause by which sever is produced.

If these general observations should excite attention, they will doubtless excite objections also, and perhaps animadversions; but whatever be the fate of his reasonings, the author rests with some confidence on the stability of the facts by which they have been fuggefted. In the view which he has taken of fever, the fupposed intentions of nature are not inquired into; and of course no attempt is made to arrange the fymptoms as they arise from the direct agency of the febrile poison, or the exertion of her medicating power. Such attempts are as unnecessary as they are difficult. It is not indeed to be denied, that the living fystem, after being excited into morbid action, paffes frequently unaffifted, through a fuccession of symptoms into a state of health; differing in this as in every other respect, from any of the arrangements of inanimate matter. But it is equally true that it often finks in the course of this process, sometimes destroyed in fever by the excess of its own re-action, and fometimes by the habits of action, or affociations produced in the course of the disease.

It is a serious error to suppose that the sebrile poison, being received into the system, is the principal cause of the symptoms, and that they consist of a struggle of nature to expel it, without which health cannot be restored. It is safer to consider the sebrile poison as an agent that excites the system

N 3

into

into fever, which however is carried on, not by the continued prefence and agency of this poison, but by the principles which regulate the actions of life. We are not therefore to wait for the fanative process by which nature is supposed to separate this virus, and to throw it off, watching her motions, and affifting her purposes; but to oppose the fever in every stage of its progress with all our skill, and to bring it to as speedy a termination as is in our power. When we dispel the morbid heat, and reduce morbid re-action in the hot stage of the original paroxysm, by the powerful means of the cold affusion, the whole of the febrile symptoms vanish; a fufficient proof that in this stage of the disease these symptoms arose from inordinate heat and inordinate action, and not from a poison circulating with the blood. Hence the fafety and the wisdom of decifive measures in the earlier stages, before the ftrength is materially impaired, or the diseased habits established. Those who practice within the tropics, where fever runs its dreadful course with fuch rapidity, ought especially to be aware of this truth, with which indeed the most enlightened of their number feem to be sufficiently impressed. The writings of Dr. Jackson and Dr. McLean inform us, that the affusion of cold water begins now to be employed in the destructive disease (popularly named the yellow fever) of Jamaica and St. Domingo, mingo, and with falutary effects*. In those climates, however, the difease ought to be combated, not merely by affusion of cold water on the surface, but by immersion of the patient in the cold bath. In this fever, hours, nay minutes are precious, and as the cold stage is short, the application of this remedy admits little delay. As foon as the fecond, or hot stage is formed, a judicious and resolute application of cold might supersede all other remedies. But the temparature of the fea, and even of the fprings within the tropics, rifing generally as high as 77° or 78° of Fahrenheit, affords a stimulus to the fenfations from fimple affusion, of a weaker kind, and hardly permits us to hope, that in the form of affusion, the cold bath can be followed by those effects in the ardent fever of the torrid zone, which in the milder fever of this climate it so happily pro-

^{*} See Dr. Jackson on the "Fevers of Jamaica." London, printed by Murray, 1791, and Dr. McLean's inquiry into the nature and causes of the great mortality among the troops in St. Domingo.—London; Cadell and Davies, 1797. Had I seen Dr. Jackson's work earlier, I should have mentioned him sooner; but I was referred to him for the first time by the work of Dr. McLean just published, and which reached me after this chapter was nearly printed off. The works of these experienced physicians deserve great praise; but their accounts of the severs of the West Indies—like all the other accounts, are without information in regard to the actual state of the heat of the patients. The thermometer has not yet been employed in the diseases of the tropics. (October, 1797.)

mingo, and wish falutary effects, to those cliduces. Artificial modes of cooling the water may indeed be reforted to, and when this can be effected, the repeated use of the affusion may in all probability extinguish the disease. But on military duty, as well as in many other fituations, artificial methods of cooling a fufficient quantity of water it may often be impossible to employ. In fuch cases, when the physician has ascertained that the heat of his patient has rifen to the degree which justifies the practice, let him plunge him fearlefsly into the cold bath, or the sea. The duration of immersion must depend on the effects on the pulse, on the fensations, and on the heat measured by the thermometer; and a greater degree of coolness will be produced by alternately raifing him into the air, where the wind blows over his naked body, and finking him into the water, than by continued immersion. (See Appendix, No. II.) The utmost care is necessary in a process of this kind, to guard against the effects of fatigue. Various instances might be adduced of the success of this practice in persons who have in the delirium of fever on shipboard, plunged into the sea-The case of Sir John Chardin, who, when at Gambroon in the year 1673, was cured by this means of a burning fever, attended by delirium, after every other remedy had proved ineffectual, is a celebrated inftance of this kind, and long ago fuggested to Dr. Glass the propriety " of making a farther trial of this un-" common and furprifing remedy." I have received a variety of authentic relations of the same nature. These have happened on board our ships at sea, during the delirium of sever, and chiefly in the warm climates; and in every instance where the patient has been fortunately taken up, recovery has followed.

An accident of this kind occurred about three years ago to Captain S-, of this port, in the Irish channel. He sprung out of his cabin window in the height of delirium, and was upwards of twenty minutes in the water. He was taken up perfectly calm, and speedily recovered .- If it be confidered, that in the Irish channel, even in summer, the temperature of the water is seventeen or eighteen degrees colder than in the West Indies, the importance of this fact will be clearly perceived. while the different modes of applying cold water to the furface are employed, it ought also to be poured into the stomach in large quantities, when the patient's heat will permit it; and the presence of nausea and vomiting is no objection to this practice, if a chilliness of the stomach is not produced.

When we consider the general adoption of cold drink in the ardent sever of the ancients, its success in the Hungarian sever, to which the yellow sever bears such analogy, and the pointed recommendation it has received from Hossman in bilious vomitings and dysentery

dysentery, it appears rather extraordinary that it has not hitherto been resorted to in the diseases of the West Indies.

The falutary effects of the cold bath, and of cold drink in fever, strongly recommend the adoption of these remedies in the plague. On the northern and eastern shores of the Mediterranean, where the liberal use of cold drink in fevers prevailed throughout antiquity, and on fome parts of which it feems to have been carried to a great extent in the beginning of the prefent century, there is reason to believe that it has now fallen into disuse. If we give credit to the answers made to the queries of Mr. Howard, in regard to the treatment of the plague, the administration of cold drink forms no part of the regular practice in that terrible disease. The Jew physician of Smyrna indeed fays, " that " the Turks in the violence of the fever, take " handsfull of fnow and apply it over their bodies, " and also eat it; and likewise sometimes throw " cold water on their feet." But the learned Hebrew cannot determine whether it is of fervice, " as these people in other respects pay no regard to " the rules of diet." Howard on Lazarettos, page 39. Morandi, physician at Venice, observes, " that " fome failors at Constantinople, in the phrenzy " of the plague, have thrown themselves into the " fea; and it is faid, that on being taken out, they "have recovered." But this happy temerity the regular practitioners do not appear to have imitated, a fact that is further established by Dr. Russel, in his account of the treatment of the plague at Aleppo*.

The administration of cold drink in the plague, as well as the external application, must however be regulated by the actual state of the patient's heat, and of his sensations of heat, of which also we have to lament that we have no accurate information. If it should be found that in this malignant

* When Mr. Howard was printing his work on Lazarettos, at Warrington, in 1788, I communicated to him an account of the fuccesses of our first trials of the affusion of cold
water in fever, in the Liverpool Infirmary, and he promised
to make more particular inquiries in regard to the empirical
use of this remedy, in his ensuing journey to the East, in the
earlier part of which, it is well known, his career of benevolence terminated.

Though the use of cold water in sever seems to be abandoned in the countries visited by Mr. Howard, there is evidence in the writings of modern travellers that it obtains on some parts of the southern shores of the Mediterranean, as well as in the interior of Asia and Africa, whither the influence of a false theory has not extended.

In his Letters on Egypt Savary observes, that "if heat were "the source of their disorders, the Said would be uninhabitable. The burning fever (the Causus of the Greeks) is the

nant disease the heat is little or not at all above the natural standard, (as is the case in the advanced stages of the confluent small-pox) our hopes from the

"only one it seems to give rise to, and to which the inhabitants are subject. They soon get rid of it by regimen,
drinking a great deal of water, and bathing themselves in
the river." page 242.

of a ship (a man of credit) as related by himself, who having taken some sailors on board affected by the plague at Constantinople, caught the insection. "I selt (says the captain) an excessive heat which made my blood boil, my head was very soon attacked, and I perceived I had but a few moments to live. I employed the little judgement I had lest to make an experiment. I stripped myself quite naked and laid myself for the remainder of the night on the deck: the copious dew that sell peirced me to the very bones; in a few hours it rendered my respiration free, and my head more composed. The agitation of my blood was calmed, and after bathing myself in sea-water, I recovered.

"Masuah," says Bruce "is very unwholesome, as indeed, is the whole coast of the red sea from Suez to Babelmandeb, but more especially between the tropics. Violent severs, called there Nedad, are very prevalent, and generally terminate on the third day in death. If the patient survives till the fifth day, he very often recovers by drinking water only and throwing a quantity of cold water over him even in his bed, where he is permitted to lie without attempting to make him dry, till another deluge adds to the first." Bruce's travels vol. iii. p. 33.

the exhibition of this powerful remedy would probably be disappointed. But the trial is recommended by every consideration.

He describes the same sever as prevailing in Abyssinia, especially in all low marshy grounds. " It is really," he says, " a malignant tertian. It always begins with a shivering and " headach, a heavy eye, and inclination to vomit." " The " face assumes a remarkable yellow appearance." This is doubtless the yellow fever of the West Indies and America. The practice of using cold water internally and externally in this fever in the lower Egypt, Nubia, and Abyffinia, has not indeed been interrupted by the influence of a false theory: but we may afcribe its continuance in part to another circumstance. The water in the springs of those uniformly fultry climates is never much colder than the air, and hence the fatal accidents from the improper use of cold drink or the cold bath, have feldom if ever occurred. (See p. 119, and 120, of this volume.) At Masuah, Bruce found the thermometer to range from 83° to 92° which gives an aftonishingly high medium temperature, and he advises the traveller in Nubia never to feruple throwing himself into the coldest spring he can find, whatever may be his heat. Vol. iii. p. 47.

mi and fire or emed a coldinery of period and

to the mode recieft univale, to am

CHAP. XV.

He deferibes the force fewer as prevailing in Abyffinia.

Animal heat—its origin.—Perspiration—its cooling influence.—Mode of operation of other remedies in Fever—Antimonials—Opium, &c.

I HAVE elsewhere observed, that if a definition of life were required, it might be most clearly established on that capacity, by which the animal preferves its proper heat under the various degrees of temperature of the medium in which it lives. The more perfect animals poffess this power in a superior degree, and to the exercise of their vital functions this is necessary. The inferior animals have it in a lower degree, in a degree however fuited to their functions. In vegetables it seems to exist, but in a degree still lower, according to their more limited powers, and humbler destination. As the capacity of preferving nearly an uniform temperature in all the varieties of climate and feafon, is a criterion of life in the more perfect animals, fo among individuals of the fame species, the degree of this capacity capacity may be confidered as a criterion of the strength of the living principle.

There is reason to believe, that while the actual temperature of the human body remains unchanged, its health is not permanently interrupted by the variation in the temperature of the medium that surrounds it; but that a few degrees of increase or diminution of the heat of the system, produces diseases, and death. A knowledge therefore of the laws that regulate the vital heat, seems to be the most important branch of physiology.

Modern chemistry assumes to itself the discovery of the origin of animal heat, which is supposed to depend on those changes in the ingesta, which the functions of life are always producing. Among the breathing animals, the principal of these changes is occasioned by the lungs on the atmospheric air which they inhale, and respiration is considered as the principal fource of the heat of this class of ani-That the oxygen contained in the atmofphere furnishes the largest portion of the pabulum of life, is now scarcely doubted, respiration being a process in which it passes from its gaseous to its concrete state, giving out a portion of the heat it held in vapour, in the same manner as steam in its conversion into water, gives out a portion of its heat. The heat thus extricated, being conveyed by the circulation

culation to all the parts of the body, is the principal means by which the whole is warmed and animated*.

Some physiologists of our own country, and on the continent, have of late supposed, that a similar process is carried on by the surface of the body, through which also, though in an inferior degree, the oxygen of the atmosphere is inhaled. Experiments are wanting to establish this position, and there is reason to believe that the principal function

of

^{*} Respiration is a case of compound elective attraction, in which the oxygen gas is decomposed, and enters into new compounds; a part feems to be absorbed by the general mass of blood circulating through the lungs; a part unites with the hydrogen of the venous blood, and forms water; another part with the carbon of this blood, and forms carbonic acid gas. In the two first cases the language in the text applies strictly, but in the last case the heat extricated does not arise from the oxygen becoming concrete, but from its entering into the composition of a gas which has a less capacity for caloric than itself. The heat disengaged in these different processes, unites with the blood, which in lofing its hydrogen and carbon (hydro-carbonat) is converted from venous to arterial, and has its capacity for caloric increased. Thus inflammation in the lungs is prevented. But the arterial blood, in the course of circulation, again abforbs by little and little a portion of hydro-carbonat, parting with caloric in the fame proportion. Hence the uniformity of temperature throughout the body. This feems the modern theory of respiration, which it must be confessed is not without its difficulties.

of the skin, in the breathing animals at least, is of a different and opposite nature. That an animal possesses to a certain extent, the faculty of rendering fenfible heat latent, or to speak more philofophically, of reducing caloric, from a free to a combined state, in cases in which the stimulus of heat might otherwise overpower the living energy, there is reason to believe, from a variety of experiments and observations. And that this is in part performed by the perspiration from the surface, can scarcely admit of a doubt. The process of perspiration which is continually going on from every part of the body, is in this point of view the converse of that of respiration: as in respiration a gas is constantly converted into a solid or fluid, and thus heat evolved, so in perspiration a fluid is continually converted into a vapour, and thus heat is absorbed. If then we suppose, that while the proportion of oxygen received into the fystem continues the same, the temperature of the atmosphere is increafed, we can understand why our heat is not increased, by supposing an increase of perspiration. And if the temperature of the atmosphere remains unchanged, while the oxygen received by the lungs is increased, we can still explain the stability of our heat, by fuppofing an increase of perspiration. The first of these suppositions is nearly realized, when a warm day comes on after cold weather; the fecond is realized when an increased respiration

pears to have a principal share in regulating the animal heat, and the chain of life seems connected with the physical world by two links, which the recent discoveries in chemistry enable us to unveil*.

A veffel

* I have faid nearly realized, in the beginning of the fentence. Supposing the same bulk of air breathed in warm as in cold weather, the proportion of oxygen will be rather less in the first case, from the diminished density of the air. There is reason to believe also, that an animal contaminates the air more slowly in warm weather, that is, receives a less proportion of the oxygen it inhales, into the system. But these particulars, in the rapid and general views which I offer, are necessarily overlooked.

Dr. Mitchel, of New York, in his ingenious paper on the nature of contagion, has unguardedly afferted, that the actual heat of the animal is always proportioned to the quantity of oxygen inhaled; but a very curfory examination of facts will shew that this is erroneous. By the experiments in No. II. of the Appendix (and various others might be adduced) it will be feen, that the living body has the power of preferving its heat nearly unchanged for a confiderable length of time in fo dense a medium as water 500 colder than the blood. This could not arife from an increased inhalation of oxygen. The phenomenon is partly to be explained from the fudden contraction of the perspiratory organs; but principally, from the powerful impression on the sensations rousing the vital principle to increased exertion, and to increased extrication of heat. It is to be observed too, that it was the trunk, and more folid parts of the body, that preferved their heat, in the experiments referred to; the extremities became cold.

A veffel filled with water, and exposed to the atmosphere cannot be raised above 212° of Faht. by any quantity of fuel, because, as heat is applied from below, evaporation carries it off from the furtace. Hot springs are of the same heat at their fountains throughout the year, probably because the evaporation from the furface of their waters increases and diminishes with the heat of the atmosphere. In like manner we may suppose the heat of the living body to be kept uniform by the evaporation from its furface increasing or diminishing according to the quantity of heat extricated in the fystem, or received from the furrounding medium. But the cautious reasoner, aware of the deceptions arising from such analogies, will here very properly inquire-Does the perspiration by the skin, the body remaining at the temperature of health, actually go on more rapidly in warm than in cold weather? Is it greater under exercise than when at rest? Is it more plentiful when oxygen is received in abundance into the fystem, than when it is imbibed more sparingly? We run little hazard in answering the two first of these questions in the affirmative, but in regard to the last, farther experiments are requisite to enable us to decide.

The infensible perspiration attracted the notice of physicians very early, and it was a common dogma among the ancients, which has descended with

0 2

too

too little examination to our own times, that the whole furface of the body inhales and exhales. The fubject has been confidered more accurately fince the experiments of Sanctorius in the middle of the last century; and the great quantity of matter supposed by him to pass off by this process. pointing out its importance in the animal œconomy, theories of disease, founded on the diminution or obstruction of this discharge, have since his time prevailed univerfally. The quantity of the matter of perspiration is not easily ascertained, and this uncertainty must continue while the absorption by the lungs, (and perhaps by the skin) from the atmosphere, remains uncertain. It was for some time believed, that the quantity of perspiration from the furface, had been exaggerated by Sanctorius, but fince a notion has been entertained of the great absorption by the skin, his calculations have been fupposed to fall short of the truth. If however we calculate that fifty ounces of water are converted into vapour on the furface of the body in twentyfour hours, this must produce a very considerable absorption of heat, and have a powerful influence on the temperature of the living system.* And however

^{*} Very various calculations have been made of the quantity of matter perspired. The only experiments on this point that deserve much notice are those of Mr. Cruikshanks. He appears to have lost upwards of seven pounds weight in

however we may diffrust the experiments hitherto made, on the quantity of the matter of perspiration, there is little reason to question the points in which they all agree—that this discharge is greater in warm weather than in cold—under exercise, than when at rest.

Important as the evaporation from the furface of the body must be in regulating the heat, it is remarkable how little it has hitherto been considered in this point of view. In the year 1755, Dr. Cullen published his discovery of the cold produced by the evaporation of sluids, (Essays and Observations Physical and Literary, vol. ii.) a phenomenon long known in Asia, but till then unobserved in Europe, and which has paved the way to so many other discoveries of the modern chemistry. It does not however appear that Dr. Cullen applied his discovery in explaining the function of perspiration. The effects of evaporation from the surface in cooling the body, was however soon after observed by

this way in twenty four hours, when at rest, in an atmosphere of 71°; but under exercise, a much larger quantity. This exceeds the calculation of Sanctorius, and the experiments of Mr. C. point out what Sanctorius did not know, the great absorption by the lungs; for there is reason to believe, that the absorption of air, vapour, or water, by the skin, if it occur at all, has been much exaggerated.

Dr. Franklin. In a letter, dated June 1758, is the following observation, "During the hot Sunday at " Philadelphia, in June 1750. when the thermome-" ter was up at 100° in the shade, I fat in my cham-" ber without exercise, only reading or writing, " with no other cloaths on than a shirt and a pair of " long linen drawers, the windows all open, and a " brisk wind blowing through the house; the fweat " ran off the backs of my hands, and my shirt was " often fo wet as to induce me to call for dry ones to " put on. My body however never grew fo hot as " the air, or the inanimate bodies immerfed in the " air*" It does not indeed appear that Dr. Franklin actually measured his heat, but he makes the inference of his comparative coolness from remembering, that all the bodies about him, even the shirt out of his drawer, felt warm to the touch, and he concludes that he was kept cool "by the continual fweating, " and by the evaporation of that fweat." Doubtless this reasoning was suggested to Dr. Franklin, and perhaps the whole circumstances recalled to his mind, by the paper of Dr. Cullen, then recently published. When, by the observations of Mess. Du Hamel and Tellet, in France, and the experiments of Dr. Fordyce and Sir Charles Blagden in England, a discovery was made of the very extra-

^{*} See Franklin's Letter and Papers, p. 365.

ordinary degrees of heat which the living body can fupport, philosophical minds naturally endeavoured to account for fo fingular a phenomenon. That the animal possesses a power of generating cold as well as heat, was the more general opinion: but the conclusiveness of the experiments of our English philosophers to establish this opinion was questioned by many, and among others by my late ingenious friend Dr. Bell, in the paper that forms the first number in the Memoirs of the Society of Manchester. In this paper the effects of the evaporation from the furface in reducing the heat during the experiments in question, is particulary infifted on. In discussing this subject with Dr. Bell, it naturally fuggested itself, that the principal office of the infenfible perspiration might be to regulate the animal heat, and this opinion, which feems fo reasonable, has been confirmed by reflection and observation.

But whatever may be the influence of the infensible perspiration in cooling the body, it does not seem to be the only process by which this effect is produced. From some experiments in the hot bath, it appears to me that the temperature of the body is with difficulty increased after the sweat begins to slow profusely, and as there can be no evaporation from the skin when the body is immersed in water, should my observation be consisted by

04

future

future experience, it will shew, that in the production of sweat itself, a degree of heat is absorbed, and thus explain in some degree the reduction of heat that follows profuse perspiration in other cases. It was indeed supposed by Albinus, Haller, and William Hunter, that the fweat, as well as the infensible perspiration, is an exudation of the watery part of the blood through the cuticle; but this opinion is contrary to all analogy, and depending on experiments made on the dead body, may be fafely rejected, notwithstanding the illustrious names that countenance it. The opinion of Dr. Fordyce and Mr. Cruikshanks seems to be the true one. The matter of perspiration is separated from the blood by the capillary arteries, and thrown out on the furface by organic pores in the cuticle (however difficult to be discovered) connected with the extremities of these arteries; and in this process there is not a separation merely, but a new combination, as in fimiliar instances. In this combination there may be an absorption of heat, and thus the coolness produced by sweating be in part accounted for. The absorption of heat may however take place in various other processes, besides the production and evaporation of perspirable matter; as doubtless its extrication takes place in various other processes besides respiration; and after all, the sudden changes of temperature that take place in certain circumstances, render it probable that the animal possesses powers

powers over its heat not yet understood, and which no chemical theories can explain*.

Leaving this point for future inquiries, the importance of perspiration to the animal economy we may safely affirm. By this process the stimulus on the extreme vessels arising from morbid distension, is diminished or removed, and whatever other methods the system may possels of diminishing its heat, this discharge seems in certain circumstances essential to that object. In situations where the organs of perspiration are spalmodically constricted, while by the increased momentum of the circulation, heat is preternaturally evolved (as in the commencement of sever) we can easily understand the disorder of the system that ensues.

It is observed, that persons who engage in ex-

^{*} Various ingenious arguments have been used by different friends, and correspondents, to show the improbability of an absorption of heat in the formation of the matter of perspiration; this being a supposition which infers, that the matter of perspiration, has a greater capacity for heat than the blood from which it is secreted, which is contrary to analogy. I have thrown out the suggestion; but would not be thought to rest any weight upon it. The phenomena which occur in the heating and cooling of the living body, I am far from thinking that we can account by for changes of chemical combination.

ceffive hard labour, speedily fink under it, unless they perspire freely, and support their perspiration by drinking fome thin and moderately stimulating fluid. This is the case with the reapers in Penfylvania, where the harvest occurs in the hottest feafon of the year, and who by means of profuse perspiration, are enabled to work in the funshine, in which the thermometer rifes very often above the heat of the blood.* This is the case with our coal-heavers, who probably lofe a fifth or fixth part of their whole weight by this discharge daily, and who supply this waste chiefly by large draughts of porter. In inftances of fuch extreme labour, a fudden stoppage of perspiration from the cessation of exertion, and the accession of cold, is frequently fatal, as has already been explained.

Europeans who go to the West Indies are more healthy in proportion as they perspire freely, especially if they support the discharge by a moderate use of gently stimulating liquids, stopping short of intoxication, and guard against the essects of too excessive an evaporation when the vigor is impaired by fatigue. In the extreme heats of the torrid zone, this indeed is not easy. The highly excitable system of the youthful European is acted on powerfully by the climate. If the orifices of the skin do not pour

^{*} See Franklin's Letters and Papers, p. 366.

pour out a proportional quantity of perspiration, difease must ensue from the direct stimulus of heat; and if the necessary quantity of perspiration takes place, the fystem is enfeebled by the evacuation, and the extreme veffels lofing their tone, continue to transmit the perspirable matter after the heat is reduced as low as its natural flandard, or perhaps lower-as occurs in the cold colliquative fweats in the latter stages of fever. In this fituation we can eafily suppose, that even a flight degree of external cold may become dangerous. The damps and chilliness of night, joined to the increased evaporation produced by the night breeze, must often prove pernicious, especially after exercise and profuse sweating. They rob they furface of its necessary heat, produce a torpor and contraction of the extreme veffels, and the dangerous re-action of the centre, which has already been explained. The proper treatment of the European on his landing in the West Indies, may probably consist in his being placed for a time in some inland and elevated station, where the land being cleared, the air is pure, and less fultry than on the coast; on his avoiding as much as possible any violent exercise; on his guarding against all intoxicating liquids, and whatever increases the heat and diminishes the strength of the body; taking care however to supply the circulation with liquids proportional to the waste. fuch means his fystem will gradually accustom itself to the climate, the organs of perspiration adapt themselves to the increased heat, and the dangers of sudden change be averted.

The perspirable matter of the European, does not appear to be well fitted to the torrid zone. Confifting nearly of pure lymph, it is speedily diffipated by evaporation, and without a profuse discharge, the furface cannot be kept moist and cool. On the other hand, the profusion of the discharge debilitates the fystem, and renders even the cold of evaporation dangerous, especially when this evaporation is increased by an accidental breeze, or the effect of the cold promoted by rest after fatigue. The negro, fitted by nature to the climate, is less accessible to the stimulus of heat, and bis unctuous fweat less easily diffipated, keeping his skin uniformly moift, fustains a more uniform perspiration as well as evaporation, and guards the system against the waste and danger of profuse sweating under fatigue, as well as the other consequences already explained. The pungent and stimulating quality of his perspirable matter will promote these salutary effects. this view of the subject, it may be worthy of investigation, whether the practice of the ancients of using unguents on the skin, still general among the eastern nations, and common among all favages, whose defective cloathing obliges them to guard by this means against the viciffitudes of temperature, Mould

should not be revived by our countrymen in the West Indies. It has been imagined, that those unguents effectually obstructed the perspiration; but this notion has been adopted without inquiry, and against probability. These unguents, like the perspirable matter of the negro, may be formed of a confiftence, that while it retards excessive sweating, does not obstruct moderate and necessary perspiration, and being themselves evaporable, they may keep up a coolness that shall diminish the necessity of the natural discharge. In the warm climates the tepid bath feems a necessary adjunct to inunction. It has been for fome time used among the French in the West Indies, and is now, (as we are informed) beginning to be adopted by the English; and with proper restrictions, it may produce the happiest effects. After exercise, under a burning fun, when the strength is wasted by fatigue and perspiration, immersion in the bath heated from go" to 95° of Fahr. will be found most grateful and falutary. It restains the profuse and colliquative fweating, keeps up the heat of the furface and extremities, and prevents the dangerous re-action of the centre, which a torpor of the extremities fo generally produces. It allays the violent action of the heart and arteries, and fooths the fystem of senfation. On leaving the bath, friction, and inunction of the furface prevent the chill of evaporation, and while they enfure the fafety of the practice, they

prolong this classic gratification. The refinement of modern manners has introduced linen teguments next the body, and abolished inunctions of the surface. Hence the warm bath has become less safe, as opening the pores of the skin, and exposing the naked body to the chills of evaporation; and thus through the greater part of Europe it no longer forms a part of the regimen of health. In the east it still prevails. If it should be revived among our countrymen in the west, they must conjoin with it the inunctions of the furface, and the fine flannel teguments next the skin, which accompanied its use among the Greeks and Romans. In recommending a trial of unquents to the practitioners of the torrid zone, I may shelter myself under the authority of the celebrated Lord Verulam, unqueftionably the first physician, as well as the first philofopher and statesman of the century in which he lived. " Inunctio ex oleo, et byeme confert ad sa-" nitatem, per exclusionem frigoris, et æstate, ad deti-" nendos spiritus, et probibendam exolutionem eorum, et " arcendam vim aeris, quæ tunc maxime est præda-" toria. Ante omnia igitur usum olei vel olivarum vel " amygdali dulcis, ad cutem ab extra unguendam, ad " long ævitatem conducere existimamus."

^{*} See Fran. Bacon. Opera fol. Francofurti ad Moenum, 1665 p. 536, 537. 538, where the practice of various nations refpecting unquents, is confidered at some length, and the cautions requisite in adopting this practice, least the body should be overheated, &c. are very ingeniously detailed.

The operation of unquents on the furface of the body prefents indeed a subject for important and original observation. The effects of the warm and tepid bath, though more investigated, are scarcely better understood; for perhaps there is no part of medicine on which fo much has been written, and so little been decided. These subjects are connected together, and might form, with great advantage, part of an experimental inquiry into the laws of animal heat. The commonly received opinion, that the warm bath relaxes and enfeebles the fystem, must, I apprehend, be admitted with many restrictions. Immersed in water, or in air, heated to the degree that quickens the circulation, we are doubtless speedily enfeebled. But by a heat short of this effect, it may be disputed whether debility is ever produced. The degree to which the bath must be heated in order to quicken the circulation, approaches nearly to that of the blood. In my own case, I have found the pulse became more frequent at 96° when the ftomach was empty, but at 94° after dinner; and the practice, faid to have been adopted by the ancients, of going into the bath to relieve themselves from the oppression of an overloaded stomach, would appear to me to have been attended with fome hazard, and with great inconvenience. It does not indeed appear how the defired effect was obtained, except through the means of profuse sweating. In future experiments refpecting. specting the effects of the warm bath, it will be of importance to observe the heat at which the sensible perspiration begins to flow; which probably varies a good deal in different constitutions, and which as it varies, will materially affect the results. It will be important also to notice the state of the stomach as to sullness; the condition of the pulse; the previous degree of exercise; and the actual heat of the surface; all of which, according to my observation, will influence the experiments.

In all inflammatory diseases, it is of importance not to use the bath heated to the degree that materially quickens the circulation; where this is not attended to, the fymptoms are heightened, unless indeed a speedy and profuse perspiration ensues. In the degrees in which it does not quicken the circulation, the warm bath is foothing and fedative, efpecially when the immersion is prolonged; and it is the temperature from 95°. to 90°. that is fo fingularly restorative after fatigue, though a still lower heat is fafe and refreshing, as those who have used the baths of Buxton can teftify.—The warm bath is frequently employed to excite a fensible perspiration, which may be prolonged after leaving it; and in this way it may be used with great advantage. Where this object is in view, it is advisable to immerfe the patient in the water heated to 94°. or 95°. and very gradually and flowly to increase the heat

to 97°, or perhaps 98°, watching its effects. When the fweat begins to appear on the forehead, if the pulse remains calm, and the patient feels at ease, an increase of heat to 99°, or even 100°, may be ventured on with fafety, should profuse perspiration be required. But when the bath is heated in the first instance to 100° the stimulus of heat produces a feverish circulation, which the subsequent defective perspiration cannot allay. The injurious effects are still greater where the bath is heated at first to 105°. or upwards, and fwear, instead of flowing more freely in confequence of the increased temperature, is discharged with greater difficulty, the stimulus of the heat, and the fuddenness of its application, inducing a contraction of the orifices of the extreme veffels of the skin. In this respect, as in most others, the analogy is perfect between the stomach and the furface; for it will uniformly be found, that fweating is excited more eafily by draughts of tepid liquor, than an equal quantity of the same sluid fwallowed as hoteas it can be borne.

The fensible perspiration excited in the warm bath does not lower the temperature of the body while immersed, but being prolonged afterwards, it becomes powerfully refrigerant, and is a remedy of great efficacy. These observations however shew the importance of regulating the temperature of the warm bath by the thermometer, and demonstrate

P

65

how little dependance can be placed on the observations of those writers who speak of its effects without noting its temperature. A variation of two or three degrees, often impossible to be ascertained by the sensations, will not merely vary the degree, but alter the nature of its effects.

The view given of the nature of fever, and of the functions of the perspiration, seems to afford an obvious explanation of the operation of those antimonial emetics, and fudorificis, by the early exhibition of which, fevers are often stopped in their first stages. The direct action of these medicines on the stomach resolves the spasmodic stricture of the capillaries of that organ and of the furface, by which the morbid heat is confined; while the concustion given to the whole fystem, diffolves the morbid catenation, and terminates the difeafe. These remedies are however uncertain in their beneficial effects, and always highly debilitating. They cannot enter into competition with the affusion of cold water on the furface, but may be employed when the presence of local inflammation precludes the use of that powerful remedy.

These general views cast some light also on that most curious subject, the operation of opium on the living principle. In sormer times it was disputed whether opium was in its nature cold or hot; in our own days it has been disputed whether it is stimulant or sedative. Such abstractions afford exercise for the ingenuity, but in the way in which they have been conducted, they foar above utility and truth. Opium relieves pain, and produces fleep, and these effects being in general concomitant, may be treated of together. But though opium generally produces fleep, yet we are fometimes disappointed in this effect, and especially in those feverish disorders where sleep is of such high importance. About nine years ago, I attended a boy of ten years of age, the fon of a friend, dangeroufly ill of fever. His nights had been fleepless, and the general irritation was very great. Opium had been administered in vain. I determined to give it in larger quantities, and to watch its effects. At ten at night he took forty drops of laudanum; at twelve the same dose was repeated. As I stood by his bed two hours afterwards, he appeared in an imperfect and agitated fleep: his eyes were half open, his face was livid; his lips and skin parched; he was evidently much diffurbed, and moaned in his breathing. At this time I perceived that he had toffed the cloaths off his feet, which lay naked, and that he grasped the cold bed-post with one of his hands. On examining his hands and feet they felt dry and burning. There was tepid vinegar in the room, and I moistened first his hands, and then his feet with it, without awaking him, repeat-P 2

ing

ing this practice from time to time: he became more quiet. Thus encouraged, I gradually moiftened his legs and thighs in the fame manner, and at length his breaft and neck, and removing the rest of the bed-cloaths, lest him covered with a sheet only. In a little while the agitation subsided, and he sunk into a sound, and seemingly peaceful sleep, which continued several hours, the surface of his body being bedewed with a gentle perspiration.

This interesting case pointed out to me clearly, that the sedative effects of opium are often counteracted by the stimulus of heat on the surface and extremities, and that the actual state of the heat of the patient is a circumstance requiring particular attention in the administration of this powerful remedy. The following observations are the result of the inquiry which these resections suggested.

Opium given in health generally produces sleep, if the mind be vacant, the stomach empty, and external impressions be excluded. In this case, its first effect is to increase the sensibility, to give a gaiety and livelines to the imagination, and to diffuse a genial glow over the surface and extremities. The actual heat of the trunk of the body is little if all increased, but the surface and extremities are brought up to the general temperature, (97° or 98°) and a gentle

gentle, and as it has been called, a breathing perspiration is diffused over the skin. In this state we sink into those happy slumbers which are ill exchanged for the realities of life. On the approach of sleep, the pulse is generally increased in frequency by four or five pulsations in the minute, and the respiration is slightly irregular, as in the gentler degrees of pleasurable emotion; but as the sleep becomes fixed and profound, the pulse subsides to its slowest and most tranquil state, and the respiration becomes regular, deep, and rather slower than natural. The various circumstances that prevent this savourable operation of opium, even in the state of health, it is not our purpose to consider at present.

When opium is given in fever, if the heat be ewo or three degrees or upwards above the natural standard, and the skin dry, it seems very generally to increase the heat and restlessness. There are exceptions. If the heat, though præternaturally great, is subsiding, and the skin beginning to soften, though not yet moist, opium very often accelerates the perspiration, and by this means diminishes the heat. In such cases its salutary effects generally sollow—tranquillity and sleep. Thus it will happen, that an anodyne draught given early in the evening shall occasion increased heat and agitation, which is deferred till two or three in the morning would have

have produced perspiration and repose. In the evening the exacerbation of fever is on the increase, or at its height, which towards morning is subfiding, the difference in the actual heat of the furface being often not less than two degrees or upwards. A dose of opium given in the hot stage of intermittent fever, will often accelerate the fweating stage and shorten the paroxysm, which under similar circumstances in continued fever, has a contrary effect; because in the former case the disposition to perspire is more eafily excited. But even in intermittent fever, opium when given in the hot stage, will be much promoted in its diaphoretic and falutary effects, by moderate draughts of tepid, or if the heat be great, of cold liquids. In continued fever where the heat is great and the skin dry, it is proper to lower the temperature of the furface, and if poffible to excite fenfible perspiration before opium is administered, if we wish to ensure its diaphoretic and foporific effects. But even after opium has been exhibited, when the inordinate heat prevents its fedative operation, it will be found fafe and falutary to use the tepid or cold affusion; and when the heat is by this means reduced, repose and sleep will follow. Tepid or cold drink will produce, though in a weaker degree, fimilar benefit. These methods of promoting the diaphoretic effects of opium, seem more certain and advantageous in fever, than the practice of combining it with ipecacuanha cuanha, or the preparations of antimony; but where opium is to be used in inflammatory diseases, or in dysentery, doubtless this last method is to be preferred. These remarks must be considered as applying to opium in its ordinary doses, that is, from half a grain to two or three grains of the extract, or from ten to sixty drops of the tincture. Perhaps it has been too much the practice of late to give this medicine in large doses, and to overlook its effects in smaller quantities. Experience has convinced me that considerable effects are produced on the system by a very sew drops of the tincture properly administered, and that it is always unwise to employ it in doses larger than necessary to produce the desired effect.

The effects of alcohol (under which term I comprehend the spirituous and vinous liquors) on the system, have a striking resemblance to those of opium. Alcohol is more heating, less diaphoretic, and less soporific. When however the mind is vacant, the stomach empty, and external impressions excluded, alcohol like opium has a tendency to induce sleep. As sleep however approaches, the heat of the body rises, especially on the surface and extremities, and stimulating the heart and arteries to increased action, produces a state of agitation with which sleep is often incompatible. Or if the dose has been such as to benumb the senses, with the

first return of sensibility the drunkard is roused by intolerable heat from his apoplectic flumbers. The degree of heat in the fever of intoxication is doubtless various in various constitutions. I have observed it to rife to 105°, nearly the highest degree of heat in the typhus of this island; and Dr. Alexander in his own case found it as high as ro70.-Under these degrees of heat, the thirst, agitation and confequent debility, are very confiderable, the fkin being in general obstinately constricted. proper treatment in this, as perhaps in all other cases of morbid heat and dry fkin, when local inflammation is absent, is to pour large draughts of cold water into the stomach, or to affuse it on the furface. By this practice the heat is reduced, the orifices of the skin opened, sensible perspiration is produced, and one half the mischief of intoxication is prevented. Similar effects may be produced, though in a weaker degree, by tepid drinks or the tepid affusion.

The administration of alcohol in fever requires the same precautions as that of opium. It should not be given when the heat is great, and the skin dry and burning: but it may in general be given with safety and advantage, when the heat is nearly of the natural standard; and even when it is somewhat above it, if the orifices of the skin are open, and the surface is most. Opium, as was before remarked, is more sudorific than alcohol, and given in that stage of the sever of intoxication in which the skin begins to relax, it often accelerates sensible perspiration, and sometimes occasions it, when it might not otherwise have taken place. In this way I explain a phenomenon which has occurred to others as well as to me, that a moderate dose of opium sometimes gives extraordinary relief to the drunkard, under the sufferings through which he must pass to sanity and strength.

These observations are not offered to the reader as including a complete view of the subject of which they treat. My principal object has been to establish the use of a new and powerful remedy in sever, and the general remarks and reasonings which are added, are chiefly with a view to explain its operation and enforce its fafety. As however these additional observations are on subjects of great importance, it is my intention at some future period to refume them. In the mean time, it may be objected, that in this rapid sketch affertions are sometimes given instead of experiments and proofs. In fuch cases however, it will be found, that the points at iffue may be readily brought to the test of experiment; and as general and obscure expressions are every where avoided, it will be easy to resort to this trial; a circumstance which will procure from candid minds a patient hearing, as well as forgiveness for those errors into which I may have fallen.

I have only to add, that the application of cold under any form, where fever is combined with local inflammation, I have purposely avoided to discuss. It is a subject of much difficulty, and my observations upon it must wait for the elucidations of suture experience*. For the same reason I have avoided a particular

That the action of cold on the living body is under certain circumstances stimulating, is maintained in note, p. 72, introduced into this edition, to which the reader is referred; and this stimulation depends on the sensation by which the application of cold is attended. In sever with local inflammation, there is a morbid degree of heat, for which the application of cold would seem the natural practice. But in such diseases the sensibility to cold (as well as to other powers that stimulate) is also much increased, and hence the difficulty of resorting to it as a remedy. Cold acting powerfully on the sensations will in the first instance stimulate the system; and it appears to me probable, that in proportion to this stimulation is the difficulty of reducing the actual temperature. Hence it may happen, that the injudicious application of cold in instam-

^{*} Finding that some of my correspondents are disposed to extend the affusion of cold water to symptomatic and inflammatory severs, and that they are even inclined to think, that in such affections it must act with peculiar advantage, I subjoin to this second edition, a few observations on this point, but rather with the view of inculcating caution than the hope of directing practice.

a particular examination of the action of cold on the body in health, whether as a strengthening or a debilitating

matory diseases shall increase the action of the heart and arteries by the stimulus of sensation, without subtracting pertuanently the stimulus of inordinate heat. This observation extends to meazles, catarrh, &c. in which a sense of cold is so easily excited.

To apply cold with fuccess in inflammatory diseases, it feems to me probable, that the fensation which attends it fhould be as much as possible moderated, and that the application should be so regulated as to ensure the effectual reduction of the temperature of the patient. If this be the cafe, the fudden and temporary affusion of cold water, so advantageous in typhus, will not be falutary, or indeed fafe, in inflammatory fever, in which it appears to me that the patient, if cold is employed, should fink himself flowly in the cold bath, continuing the immersion for such a duration as the state of his heat, and of the pulfation of his heart and arteries will admit. By the experiments in Appendix No. II, it appears, that under continued immersion in cold water, the sensation of cold speedily abates, and that the frequency of the pulse is in a most striking and uniform manner reduced. And though the heat in the trunk of the body is wonderfully fustained, yet in the extremities, and over the furface, it is speedily and permanently lessened. It is true these effects may be modified by difeafe in fuch a way as to contradict experiments made on a person in health; and though in pure synocha the cold bath would probably be advantageous, yet ferious objections may present themselves against employing it in pleuritis or peripneumonia; objections which I do not think infurmountable, but which experience neither enables me to invalidate or to strengthen. In phrenitis, gastritis, enteritis, &c. especially a debilitating power, an inquiry that ought to be first entered into, in a regular investigation of the effects of temperature, and its changes upon life.

CHAP.

in their desperate forms, I think immersion in the cold bath ought to be hazarded.

It may ferve to support this reasoning, that the local application of cold to parts inflamed, is regulated by the principles I have laid down. It must not be sudden and temporary, but great in degree, and permanent in duration. Hence the success with which ice, and snow, and the clay-cap, are applied to different parts, for the purpose of preventing or reducing inflammation. In all such cases the sensation of cold speedily subsides,—and even though ice be lying on the part affected.

It is well known that the general action of cold may be extended over the fystem by its application to a part. The use of cold in hemorrhagies is often regulated by this maxim. I have found that hemorrhagy from the lungs may be stopped by immerting the feet in cold water, and perhaps this may be done ftill more certainly by a permanent application of cold to the penis and fcrotum, which part with their heat more eafily than any other portion of the furface of the body. have found that a ftill more powerful effect was produced in hemoptoe, by immerfing the body up to the pubes in cold water, a practice that I can speak of from experience as often fafe and efficacious in this difeafe. But in all these instances a permanent, as well as powerful application of cold was requifite, and from the analogy between hemorrhagies and inflammations, this may support our reasonings respecting the proper method of employing cold as a remedy in these last difeafes.

CHAP. XVI.

Population of Liverpool.—Prevalence of fever among the poor, with documents.—Benefit derived from the Manchester House of Recovery in contagious diseases.

—Propriety of the general adoption of similar establishments.

THE Typhus, or low contagious fever, prevails in all large cities and towns to a degree that those are not aware of, who have not turned their attention to the subject, or whose occupations do not lead them to mix with the labouring poor. In Liverpool it has been supposed that this disease is seldom to be met with, and it is certainly true, that the upper classes of the inhabitants are not often subjected to its ravages. It has never, in the last sixteen years, spread among the higher classes so as to occasion any general alarm; and when the extent to which it is constantly present among the poor, shall be proved by authentic documents, this circumstance will serve to demonstrate the narrow sphere

fphere of the contagion, and to shew how much it is within the limits of human power to lessen the frequency of the disease. Before we produce these documents, it will be proper to offer a few general observations.

The population of Liverpool, as taken by actual enumeration, amounted on the first day of January 1790, to nearly 56,000 persons, and admitting the increase since that time to be in the ratio of preceding years, it would now (October 1797) exceed 63,000. But as the progress of the town has doubtless been checked by the war; something may be deducted from this last number; though we shall probably be under the truth in fixing it as low as 60,000*. Of the inhabitants of Liverpool, it is ascertained, that about 7,000 live in cellars under ground, and nearly 9,000 in back houses, which in general have an impersect ventilation, especially in the new streets on the south side of the town, where a pernicious practice has lately been introduced

^{*} The enumeration mentioned, was made by Mr. Makin Simmons, and published in Mr. Gore's directory in the year 1790, but omitted in the last edition. It occupied him three months, viz. from October 13th, 1789, to January 13th, 1790, and was executed with great care and accuracy. Mr. Simmons gave the population of each street separately, enumerating also the front houses, back houses, or cottages, and the

duced of building houses to be let to labourers, in small confined courts, which have a communication with the street by a narrow aperture, but no passage for the air through them. Among the inhabitants of the cellars, and of these back houses, the typhus is constantly present, and the number of persons under this disease that apply for medical assistance to the

inhabited cellars, with the number of persons in each of these descriptions of habitation. The general result was as follows:

Front houses 6540, containing 39,118 Inhabitants.

Back houses, or Cottages, 1608, containing 7,955 Inhabitants.

Inhabited Cellars 1728, containing 6,780 Inhabitants.

The Workhouse and other Charities, 1,879

In all, 55,732.

There were besides \\
Empty houses \\
\frac{717}{2}

Making in all 10,593 Houses, or if the Cellars be Deducted 1,728, 8865 will be the number.

There is no town of an equal fize in the kingdom that possesses (as far as I have heard) so authentic a document of its population. The numeration of Mr. Simmons fell short of what many supposed to be the number of our inhabitants; but popular calculations on this subject always exceed the truth—perhaps not less in general than by a fourth part.—The popular calculation of the inhabitants of Liverpool, compared with other towns of which there is no accurate enumeration, may be taken at 75,000.

the charitable inftitutions, the public will be aftonished to hear, exceeds 3,000 annually*.

The public charitable institutions in Liverpool for the relief of the fick poor, are the Infirmary and the Dispensary. Fevers, by one of the original rules of the Infirmary, are excluded from that hofpital, though this rule, as may be seen from the narrative with which this volume commences, has been occasionally departed from. But with a few exceptions, the care of the poor in fever falls on the Difpensary. Of this institution I was for several years one of the physicians, and such of my observations respecting it as are not supported by authentic documents, are founded on personal observation. Unless in cases where they are removed into the Infirmary, the care of almost all the inhabitants of cellars when fick, and of a great part of the inhabitants of the back houses or cottages, devolves on the physicians, furgeons,

^{*} From the first of January 1787, to the last of December 1796 inclusive, a space of ten years, it will appear by the table which follows, that 31,243 cases of sever were admitted on the books of the Dispensary alone, which on an average is 3124 yearly. If it be supposed that some cases may be denominated typhus by mistake, let it be considered, how many cases of this disease do not appear on the books of the Dispensary, though occurring among the poor, being attended by the surgeons and apothecaries of the benefit clubs to which they belong, &c.

furgeons, and apothecaries of the Dispensary. This description of persons, amounting in all to about 16,000, do not appear to be healthy, the number of cases admitted annually on the books of the Dispensary, on the average of the last ten years, exceeding 13,000. In the commencement of the year 1780, a register of these cases was begun by Mr. Avison, late apothecary to this charity, and continued by him to the summer of the present year, when he terminated a life of spotless integrity, singular usefulness, and incessant exertion*. From this valuable record I have formed the following table.

Q

GENERAL.

* I infert the diseases of September 1790, extracted from the Register, which at the same time will convey a specimen of our most prevailing diseases, and shew that inflammatory and symptomatic severs are not included under the general title of sever.

September 1790.

Febris	310	Catarrhus	24	Debilita	s 19
Ophthalmia	59	Dyfenteria	2	Hydrop	s 8
Cynanche	13	Cephalagia	10	Scorphu	ila 11
Pneumonia		Apoplexia		Rachitis	THE .
Hepatitis		Paralyfis	4	Syphilis	24
Gaftritis		Dyspepsia	21	Icterus	1
Phthifis	10	Epilepfia	1	Dyfuria	5
Rheumatifmus		Afthina	16	Vermes	5
Eryfipelas		Pertuffis		Pfora	142
Variola	-	Colica	24	Partus !	Difficilis r
Rubeola	29	Cholera	24	Uftio	14
Scarlatina	-	Diarhœa	24	Ulcus	61
Aphtha	13	Hysteria	4	Vulnus	89
Hæmorrhois		Convulfio	2		
Hœmorrhagia		Infania	1	Ala	1093

			,
,		,	H
и	6	3	•
L		•	٠
			١
В	=	3	8
۲	3	7	١
Į.	۰	•	۱
и	3	s	,
	5	ā	6
	ä	8	Ē
V	2		۰
К	Ξ		2
4	3		
		ŧ	•
			۱
Б	=	=	2
ø	•	•	•
E	9	g	٠
			8
,		,	ŗ,
Ľ	•	9	•
H			c
ľ	5	ı	,
u	۰	ı	1
P	ī	ř	7
,	٠,	4	e
ú	ı	ı	i
ľ	B	i	5
ı	f	í	i
H	H	þ	
B	ı		
ď	Ħ	í	í

the	00 0	00 00	000	000	-	13	0 4	+ 01	1	2	1 º
o, till the y year. Tatal.	1174	1118	11119	11100	1177	13237	14200	1383	1366	1300	213,305
fan. 1780, of every J Dec. 7	855	760	1000	959	1140	1095	1093	1159	1273	1236	17603
fir A of 3	730	704	1055	858	951	982	1274	1215	1300	1300	17 660
from the first of Jan. and of each month of e	825	736	934	8000	927	908	1257	1335	1022	1259	17022
benfary f b year, e Sept.	557	722	929	930	1000	10017	1241	1128	1160	1004	16672
Liverpool Difpenfary, numbers of each year, aly. August Sept.	672	619	870	1010	CITIS	930	1112	1065	6801	945	15088
the Liver the number July.	787	686	895	816	916	1039	1601	1077	9901	1200	16507
Marie and Printers	805	1605	934	791	903	1070	1166	1005	1105	1571	17804
on the books of s, distinguishing	827	9601	968	986	944	1113	1176	1247	1112	1020	17803
	995	1243	969	696	1002	1153	8111	1228	1138	1171	IOI72
f Patients ad a space of 17	1147	1190	976	1111	882	1361	1323	1188	1170	2003	20475
ther of 1 796, a 1 Feb.	888	1041	916	760	096	1053	1126	1037	1175	1309	17388
Shewing the number of Patients admitted tast of Dec. 1796, a space of 17 year Year Jan. Feb. Mrcb. April.	893	1106	738	982	1022	1415	1223	1088	1901	1298	18220
Shewing last of Year	1780	1782	1784	1786	1788	1789	1971	1793	1794	1795	-

By the foregoing table it appears, that the number of patients admitted under the care of the Difpenfary annually, taking the average of the last seventeen years, is 12,548. But if the last ten years only be taken, the average number will be found to be 13,355. It appears that the greatest number of patients admitted in any one year, was in 1795, amounting to 15,235, exceeding the average of the last ten years by 1880; and that the smallest number admitted for the last seven years, was in 1796, falling short of that average by 354, and of the number admitted in 1795, by 2234. When it is confidered that the difference between these two years falls chiefly in the winter and fpring months; that the winter and fpring of 1795 were the coldest, and those of 1796 the mildest, experienced in our climate for a long feries of years; thefe ftriking facts will corroborate the statements of Dr. Heberden, in proving that the common opinion, which supposes a mild winter into our island to be unhealthy, and a fevere winter to be favourable to health is in both particulars the reverse of the truth. It appears also, that though the number of patients in the winter months of 1794-5, greatly exceeded the general average, yet it was in February, and more especially in March, that the excess rose so very high; on this last month the astonishing number of 2003 patients having been admitted. In the three preceding months the cold had been intenfe,

intense, but in March the weather became warmer, uncertain, and stormy. Hence it appears, that though intense cold is directly unfavourable to the health of our poor, yet that it is in the transition from this intense cold to warmer weather, that their constitutions sustain the severest shock; a position that is still more fully supported by the decisive evidence that the month of March is the most unhealthy of the year, and next to it the month of April, taking the average of seventeen years together. On the same evidence we may state the fuccession of the months in respect to their unhealthiness in the following order-March-April-January—October— May — December — November — February - June - September - July - and August.* But though on the whole, fummer and autumn are are more healthy than winter and spring, the difference is perhaps less than might have been expected; and the difference even between the two months that forms the extremes, is not very great. In the feventeen years recorded in the table, the average number of patients for each month of the whole is 1045; the average number in the month of March

is

^{*} It appears that June stands in the table higher than May or December, but this arises from the vast number admitted in June 1782, when the influenza was prevalent, and was confined nearly to that month. Allowance being made for this, June will stand as inserted.

is 1204; in August 940. These observations however are strictly applicable only to a particular description of persons in a particular situation—to the poor of Liverpool—and without the corroboration of other documents, must not be extended to the general mass of society, or to the kingdom at large.

I proceed to give, from the same authentic register, the table of typhus, the more immediate object of our inquiries.

A TABLE

Q3

0	80,						1	23	10											
	every year	(Your	ירעם	iki				700						S.P.		19	1	00	21	
	Januar th of ev	Total.	1917	2113	2873	1000	2662	\$17	2205	3177	3107	2930	3470	3344	3151	2925	3405	2970	2002	48367
A TABLE,	each mon	Dec	133	223	220	1/3	167	102	161	320	295	208	5602	371	174	228	320	180	247	4047
	om the	Nov.	150	208	150	25/	309	244	107	283	315	204	355	335	212	224	258	317	153	14201
	d on the books of the Liverpool Diffeenfary fr	100		234	0	310	247	250	211	234	341	214	340	344	330	305	290	239	329	14592
		Sept.	129	101	143	205	200	180	691	218	258	212	310	300	211	338	303	961	254	4025
		Aug.	150	127	140	227	230	180	188	192	311	162	270	348	223	161	245	159	182	3431
	the Lir	July	161	157	120	280	270	173	159	255	271	184	343	266	236	199	162	248	203	3852
	books of Bing th	Fune	183	154	159	212	225	187	155	356	245	205	247	240	237	334	305	230	176	3850
	on the l	May	168	187	148		232	219	202	313	253	391	281		261		337	234	288	4206
	2	Apr.	173	200	0	207	247	294	244	234	255	323	204	230	278	250	280	204	242	4250
	vers ae	Mch.	179	180	231	184	245	296	216	301	213	338	337	277	269	221	383	546	266	14682
	r of Fe	Feb.	125	146	265	158	194	268	991	200	174	176	248	247	261	200	230	265	191	13602
	of De	Jan.	150	130	268	210	170	285	101	256	226	210	176	252	250	174	157	152	107	3683
	to the last of Dec. 1706, inclusive:	Year.	1780	1781	1782	1783	1784	1785	1786	1787	1788	1780	1700	1041	1702	1702	1704	1705	1796	

Thus it appears, that of 213,305 patients admitted in seventeen years at the Dispensary, 48,367 have laboured under typhus; nearly a fourth of the whole; an immense proportion! It is curious to observe, that March takes the lead in respect to the prevalence of fever, as well as of difeases in general, and that August, as before, brings up the rear. It appears also that the month of March 1795, which exceeded every other month in the production of diseases in general, exceeds in a still greater proportion in the production of fever; no less than 546 patients having been admitted in that month under this fingle disease. The year 1795 has not however the same unfortunate pre-eminence here as in the table of general disease, being surpassed in the number of fevers by feveral others. But the year 1796 enjoys a fimiliar distinction as formerly, having produced fewer fevers than any other of the last ten, and 545 fewer than the average of the whole. Taking the feventeen years in the table, the average number of fevers annually is 2845, and the average number monthly is 237. The average number of March rifes to 275, and of August, finks to 202. On the whole the prevalence of fever is greater, and the influence of feafons upon it less, than might have been expected.

For the last ten years there have been, on an average, 119 patients ill of fever constantly on the books

books of the Dispensary; or if the last seventeen years be taken, the constant average number will be 109. Of convalescents, unfit for labour, the average number will be nearly as great. Thus, in Liverpool, 240 of the poor may be considered as constantly rendered incapable of earning their sub-sistence by this single disease; and as the poor seldom lay up any part of their earnings for a season of sickness, the expense of their maintenance must, in one form or other, fall on the public. If we take this as low as 101 for each, it will amount to 24001 annually.*

It would afford me particular pleasure to introduce authentic documents of the success attending the treatment of the typhus by my brethren of the Dispensary; but the valuable register which furnished

^{*} These calculations will not be expected to be minutely correct, but I apprehend they will in general be found under the truth. The average number of severs annually for the last ten years, is 3124, and the mean duration of each sever is supposed to be fourteen days; $\frac{3124 \times 14}{365} = 119.8$ daily. If the average of the last seventeen years be taken, this will give 2845 annually, which produces on the same principles 109 daily. I calculate that the period of convalescence after each fever is sourteen days, or upwards, which gives the same number of convalescents as of patients in fever, deducting those who may die, a much smaller proportion than could well be supposed.

nished materials for the preceding tables does not mark the iffue of the particular cases. In general however it may be afferted, that where the affiftance of the Dispensary is resorted to in time, even under the disadvantage of being attended in confined and dirty habitations, a great proportion of cases terminate favourably. The general treatment of fever at the Dispensary, consists in giving antimonial emetics in the first instance, and afterwards bark, opium, and wine. Nourishing food is occasionally adminiftered. Ablution with cold water has feldom been used, and indeed it is in general difficult, if not impossible, to employ it in the obscure cellars in which this disease is usually found. The fever that prevails among our poor is remarkably uniform; it is the pure typhus, to which the cordial treatment can be applied with fafety, and to a great extent. Whereas among the higher classes, fever is often attended with inflammatory symptoms in the first instance, sometimes with pneumonic symptoms through a confiderable part of the difease; and in fuch cases the indications of practice being contradictory, fuccess is much less certain.

Though the cure of this disease is a principal object of our charitable institutions in Liverpool, it is to be lamented that little or nothing has been done for its prevention; although it may be confidently afferted, that this object, great as it is, is not beyond

beyond the reach of human power.—This infection arises from a want of cleanlines, and ventilation; and its influence is promoted by damp, fatigue, forrow, and hunger. When the subject shall be seriously taken up by the public, the means of securing cleanliness and ventilation, even in the greater part of our subterraneous dwellings, may be clearly pointed out; and though the secondary causes, it I may so call them, seem less capable of being removed, yet much may be done to alleviate them, when a comprehensive inquiry into the condition and wants of the poor (the most pressing of all objects in the eye of enlightened patriotism) shall engage the attention of the legislature.*

Important

^{*} The want of a diet fufficiently nutritious, is doubtless one of the causes that promote the typhus, and other diseases, among our poor. This does not feem to arife in general from the price of their labour being inadequate to furnish such a diet, but from their ignorance in the most advantageous modes of cookery, and still more from their indulging in articles that confume their means without adding to their fuftenance. In the eighteen hundred cellars in Liverpool, there are many in which animal food is not tafted more than once a week; but there are very few in which tea is not drunk daily; it is often indeed drunk twice a day. The money fpent on tea is worse than wasted. It is not only diverted to an article that furnishes no nutrition, but to one that debilitates the empty fromach, and incapacitates for labour. Hence the vaft number of dyspeptic complaints among our patients at the public charities, which are almost all to be traced to

Important information in regard to the means of preventing the generation and progress of contagion, may be found in the writings of physicians of our own county and neighbourhood; of Dr. Percival and Dr. Ferriar of Manchester, of Dr. Campbell of Lancaster, and of Dr. Haygarth of Chefter. The persevering exertions of Dr. Percival, Dr. Ferriar, and the other gentlemen of the faculty attending the Manchester Infirmary, have produced an inftitution in that town, which deferves to be imitated in every large city and town in the kingdom; a house of reception, or as it is justly called, a House of recovery for the poor, when labouring under fever. Into this afylum, on the first notice of the disease, the patient is immediately removed, and proper methods taken to purify his habitation. By these means the contagion is extinguished in its birth; and the patient himself being removed to a fituation where all the refources of art can be employed with advantage for his benefit, and removed at a period of the difease when these resources are of great avail, is very generally restored in health to his family and to the community.

The

the use of tea or spirits, often indeed assisted by depression of mind. At the Infirmary and Dispensary together, this class of patients exceeds five hundred annually. The great majority are semales.

The benefits of this admirable inftitution in Manchester, no longer rest on the basis of theory; they are established on the immoveable soundation of fact and experience. The prevalence of sever in the town is diminished to a degree that has exceeded all rational expectation; and the sears entertained, that the House of Recovery might spread the contagion through the streets in its vicinity, are sound to be entirely groundless.*

The

^{*} The following facts are extracted from the Report of the Board of Health, dated Manchester, May 26th, 1797. Since the establishment of the House of Recovery, the whole number of home patients (i. e. patients requiring attendance at their own habitations) is reduced one half; but the number of fever cases, in a far greater proportion. In January 1796, 226 cases of fever were admitted at the Infirmary of Manchester, but in January 1797, only 57. Fears had been entertained that the fever might spread from the House of Recovery into the neighbouring buildings, and fo through the town, and this had given rife to a confiderable opposition. The streets in the immediate vicinity of the House of Recovery, are Silver-street, Portland-street, &c. In the first eight months after the establishment of the House of Recovery, the number of cases of fever admitted on the books of the Infirmary from these streets was 25. In eight months of the preceding year, viz. from the 20th of September 1795, to the 20th of May 1796, the same streets furnished 267 cases of sever. From the 20th of Sept. 1794 to the 20th of May 1795, they furnished 389 cases; and from the 20th of Sept. 1793 to the 20th of May 1794, 400 cases. Facts such as these, require no comment; they have made a forcible impression in Manchester, 25

The utility of the House of Recovery in Manchester has been still more completely ascertained in the course of the present summer.—" Within "this last four months (I copy a paragraph in a letter from Dr. Ferriar) "we have been threatened "with the return of an epidemic sever, and danare gerous cases of typhus have appeared in every quarter of the town; yet by selecting those patients who were, from their symptoms and situatients who were, from their symptoms and situatients who were, from their symptoms and situa-

might be expected, and the Report I quote contains public testimonies in favour of this excellent Institution from the Board of the Instrmary, the Stranger's Friend Society, and The Special Board of Overseers of the Poor. The total number of sever-cases received into the House of Recovery from its being opened, the 19th of May 1796, to the 31st of May 1797, is

Of these are discharged cured,

Dead

Remaining,

7

371

The benefit of this institution is not to be calculated by the number of persons cured. Every single removal into the House of Recovery, probably prevents on an average two or three cases of the disease. As the institution goes on, it may be considently predicted, that the proportion of cures will be still greater than here stated, because experience will induce the poor to apply for relief in the early stages of sever, when the power of medicine may be exerted with the greatest advantage. I have the authority of Dr. Ferriar for saying, that since the 31st of May to the 25th of September, two persons only have died in the House of Recovery.

" tion, most likely to communicate infection, and " by removing them into the fever ward, the week-" ly lift of home-patients has been prevented from " increasing much beyond their usual number. " Our House of Recovery, which contains accom-" modations for twenty-eight patients, is fufficient " for Manchester at present. In the event of a " peace, and the influx of new inhabitants which " it must occasion, it may be necessary to provide " accommodation for a greater number. But I " think fever-wards capable of containing fifty pa-" tients, with separate rooms for the reception of " cases of Scarlatina Anginosa, Measles, or Small Pox, " would answer every purpose here, in times of the " greatest emergency, even supposing a very con-" fiderable increase of population."

'Till the latter end of the year 1787, there was no place for the reception of fever, or any other contagious difease, in Liverpool. At that time the fever broke out in the Infirmary, described in page the fixth of this volume, and two wards being fitted up for the patients insected, one for each sex, these were afterwards occupied by such cases of sever as occasionally presented themselves. The reception of sever into these wards continued for sive years and one month, during which time 232 cases of sever passed through them. The wards were small, and under ground; in every respect inconveniently fituated.

fituated. They held at most eight beds each, but were properly adapted to six beds only. But they admitted of cleanliness and ventilation, and it was there that an opportunity was afforded of employing the affusion of cold water; of remarking its effects; and of establishing the rules that ought to regulate its application.

From the first however, the admission of fever into the Infirmary, (which was contrary to one of the original rules of the institution) was objected to by fome of the friends of that charity. They allowed the benefit arifing from a receptacle for fever, but they contended that the Infirmary was not adapted to that purpose; and that the attempt to render it a hospital for fever, the wards already mentioned being all that could be allotted to that difease, did not afford the relief required, while it obstructed the original purposes of the institution-The want of an afylum for this disease elsewhere, overcame these objections. But about the beginning of 1793, a contagious fever having prevailed in the work-house, two spacious wards were fitted up in that building by the Parish Committee, for contagious diseases: and an arrangement having been made for the reception of fuch cases of fever as presented themselves at the Infirmary, into the wards of the work-house, from that time the admission of fever into the Infirmary has ceased.

The wards for fever at the work-house, have fecured that immense hospital, often containing 1200 persons, from the spreading of fever, to which, previous to their establishment, it was perpetually liable; and though they are in the very centre of the building, and cannot be entered but through the common stair-case, yet no single instance has occurred of the contagion extending from them into the other parts of the house. They have also relieved the Infirmary from the necessity of admitting those miserable wretches, who, under the influence of fever (perhaps caught on ship-board) and refused admittance into private lodging, were brought up to that hospital, where, if not received, they might have perished in the streets. But they have not been of fufficient fize to answer the purpose of a general House of Reception for the poor labouring under fever throughout the town, nor has their power of accommodation been employed to its full extent with this view.* The physicians of the Dispensary, who attend the work-house, have authority

^{*} These sever wards contain eighteen beds very conveniently, and from the 20th of August 1793 to the 31st of August 1797, a space of sour years, only 530 cases have passed through them. They have not therefore on an average contained more than six sever-cases at a time. Of these 530 cases, 51 have died, a small proportion, considering that 16 died in 24 hours after admission, and were in articulo mortis when received.

authority to remove patients on the first appearance of fever, from any part of the workhouse itself, into these wards; but they have not the same authority to remove them from the town. The delay occasioned by the forms to be passed through in obtaining the admission of these patients has very generally prevented the attempt being made, and sometimes where it has been made and succeeded, has occasioned the relief to come too late.

In consequence of considerations of this kind, a motion was made at the annual vestry in March 1796, that a fet of fever-wards should be constructed at the expense of the parish, for the general use of the poor, on the plan of the Manchester House of Recovery. Two wards, each capable of containing twenty beds, were stated to be sufficient for the town; it was proposed that they should be erected on the elevated and open ground contiguous to the workhouse, and be supported out of the parish rate. And it was afferted that this would be a plan, not merely of charity, but of œconomy. The measure of establishing fever-wards was unanimously agreed to, and a committee appointed to carry it into effect; but differences of opinion having arisen as to the execution of the plan, all proceedings on the subject have been suspended. It is earnestly to be hoped that they may speedily be refumed.

Contagious diseases, and more particularly fevers, have in general been excluded from the hofpitals of England, those of London perhaps excepted; but the evidence of our own Infirmary and work-house, of the Chester Infirmary, and of various fimilar inflitutions, proves, that under proper regulations they may be admitted under the fame roof with other diseases, without any danger of the infection spreading through the building. It is not however to be denied, that the attendance and the arrangement requisite for patients under fever, render it a matter of convenience and propriety that they should be received into a separate building; and it is a point capable of demonstration, that hofpitals for such diseases stand pre-eminent in point of utility over all other hospitals, those for accidents that require the immediate affiftance of furgery alone perhaps excepted. The benefit derived from hospitals in other cases consists in removing disease, and is confined to the patient himself; but in cases of contagion, the evils prevented are much greater than those remedied, and the benefit is by this means extended from the patient himself to the circle by which he is furrounded. The eftablishment of fuch hospitals was first suggested by tracing the infection which propagates fever, to its origin, and afcertaining the power of ventilation and cleanliness in preventing and in alleviating the disease. The arguments for fuch hospitals are strengthened by the discovery of the chemical methods of destroying this contagion; and if I do not greatly deceive myself, they are still farther strengthened by the success of that mode of practice in sever, which it is the chief object of this publication to explain and to establish. A vigilant exercise of all the means of prevention and of cure, might indeed in a short period, supersede the use of hospitals for fever, by extinguishing the disease; a prospect in which the philanthropist might indulge with more safety, if he could calculate with the same considence on the wisdom, as on the power of his species.

ha affahalfhat been intificeent induction of facts, and

if it o cur at all, it was loggeofed that the quantities

CHAP

by the different to the chimical methods of the

CHAP. XVII.

Is there an Inhalation by the Skin?—Case of P. M. Esq. who died of Inanition.—General Restections.—Conclusion.

IN speaking of the origin of animal heat, and of the nature and office of perspiration (p. 192, 196) it was questioned whether the inhalation of vapour or water through the skin into the circulating sluids be established by a sufficient induction of sacts, and if it occur at all, it was supposed that the quantity had been exaggerated. Among the ancients however, the doctrine in question was universally received, and it has been admitted by the most celebrated physiologists, not merely of the past, but of the present times. The sacts and arguments in support of it are detailed by Haller in the 5th volume of the Elementa Physiologia; but on examination they will be found to afford no direct proof of its

truth.* Of the great variety of authors quoted on this occasion, no one affirms the actual increase of the weight of the body in the warm bath, Dessault excepted, and this is a point that Haller himself seems never to have subjected to experiment.

The inhalations through the furface in vegetables, in infects, and in certain of the amphybia, are the analogies that give the most direct support to this doctrine, applied to the human species. It is however controverted by many facts, and if it should ultimately be established, it must be reared on a new foundation. When I was at Buxton in 1788, I made an experiment on the effects of bathing on the weight of my body, and after half an hour's immersion, I found it rather diminished than increased. This experiment had I believe been made

* The arguments drawn from the appearances that occur in the dead body, may be wholly set aside; as may the instances of certain narcotic vegetables, applied to the skin, (e. g. tobacco and white hellebore) occasioning sickness at the stomach, which are cases of sympathy, or of absorption after the cuticle had been eroded. The arguments drawn from the increase of weight in moist air, and from the vast discharge of urine in diabetic cases, are also wholly fallacious. In diabetes we now know there is no absorption by the skin, and the phenomenon in question, which has been much exaggerated, must be accounted for by the action, not of the skin, but of the lungs.

before by Dr. Pearson, with the same result; it has fince been repeated frequently with great care,* and it is afcertained as a fact, that no increase in the weight of the body is produced by immersion in water of the heat of 82°. In the year 1790, I had a patient in diabetes, whose cuticle, as is usual in that disease, was in a morbid state; and being defirous of trying how far the inordinate action of the kidneys might be affected by a gentle stimulus applied to the skin, I immersed him in a bath of the temperature of 96°, weighing him before and after immersion. There was no variation in the weight. This experiment has lately been made by Dr. Gerard, in another case of Diabetes, an account of which is given in the publication of Dr. Rollo;+ and as it was repeated a great number of times with the utmost care and accuracy, it may be confidered as established, that immersion in the warm bath in Diabetes produces no increase of weight. I have made five different experiments of the fame kind on myself, varying the heat of the bath from 87°. to 95°. but never in any instance found my weight augmented. It may be faid however, (and it has been faid) that though in Diabetes, where

^{*} In one case, a sum of money depended on the issue, and a friend of mine submitted to be the subject of the experiment.

⁺ Rollo on Diabetes Mellitus, v. ii. p. 72.

the epidermis is diseased, no liquid is inhaled; and though in health, where the vessels are full, no absorption may take place; yet that where the body is wasting from a want of the proper food through the stomach, the plastic powers of nature may be exerted to supply the defect, and to excite an inhalation by the absorbents under the cuticle, or through those pores on the surface, by which exhalation is usually performed.

The following case will throw some light on this point. I give it however not merely with this view, but as containing other particulars that appear to me curious and interesting.

In August 1795, a gentleman of ——shire, aged 66, applied for my assistance in a dysphagia, (obstructed deglutition) with which he had been for ten or twelve months assisted. At first the complaint was slight, and occurred only when he attempted to swallow dry and hard substances; it afterwards extended to solids of every kind; and at the time I first saw him, though still able to pass down liquids, the quantity he could swallow was not sufficient for his nutrition, and he was considerably reduced. On the introduction of a bougee into the gullet, it passed about two inches easily, but then met with an obstruction, which by a moderate pressure was overcome. It then passed easily

feven or eight inches more, but experienced a firm refistance in the lower part of the tube, towards its termination in the cardia, which the skill and patience of Mr. Park could not furmount, and which finally baffled every resource of his art. Though we had not an opportunity of examining the Oefophagus after death, yet the nature of this obstruction was evident. It doubtless proceeded from a schirrous tumour, gradually increasing, which at first diminished the passage, and in the end obstructed it wholly; and it existed in that part of the tube, which, from its fuddenly narrowing, is, as Dr. Hunter has remarked, most exposed to the causes, and most frequently the seat of this disease. Mr. M. never had any confiderable pain from the preffure of the bougee on the tumour, and though he expectorated (if the expression may be used) much mucus, there never was any purulence discharged, or any reason to believe that the tumour had ulcerated.

The last time Mr. M. swallowed in any quantity, was on the 17th of October. On the evening of that day a sudden increase of the obstruction came on, and from this time forth he was able to swallow only a table spoonful of liquid at a time, and with long intervals. It was with difficulty that he got down seven or eight spoonfuls of strong soup in the day, and this quantity gradually diminished.

On the first of November, the passage seemed wholly obstructed. As he wasted rapidly, we were obliged, from the 18th of October to employ the subsidiary modes of nutrition, and he went into the tepid milk, and water bath, and had a nutritive clyster administered once a day.

As Mr. M. himself was far from despairing of his recovery to the last, the affectionate friends around him did not wish him to be informed of his real situation. He continued therefore to swallow two or three table spoonfuls of milk daily, till about the 15th of November, on the supposition that some might pass; but though it rested in the Oesophagus some time, it was constantly ejected at last; and being satigued by this expectoration, which disturbed his rest, and by which he perceived that the whole of the milk was thrown up, for the last twenty days of his life he made no attempt to swallow, but contented himself with washing his mouth and sauces with wine and water.

Mr. M.'s friends, though they could not but foresee the issue of his case after all attempts to remove the obstruction had failed, yet desired that his life might be prolonged to the uttermost. The following plan was therefore adopted with this view, on the last days of October, and it was persevered in with the variations afterwards to be mentioned.

tioned, till the commencement of the delirium that preceded his death.

Every morning at eight o'clock, he had a clyfter, confifting of eight ounces of strong broth, made chiefly of membranous parts of beef (being the most nutritious, according to Dr. Fordyce) into which were rubbed two yolks of an egg, and to which were added forty drops of laudanum. This was repeated at three in the afternoon, and again at nine in the evening; but in the evening, instead of forty, one hundred and twenty drops of laudanum were added, to compensate the want of an anodyne draught, to which he had been accustomed at the hour of rest. Previously to this however, he was placed up to the neck in a tepid bath, of which one fourth was milk, and the rest water; the whole quantity amounting to twenty-four gallons. The temperature was fixed at 96° to accommodate his fensations, and the time of immerfion was gradually prolonged from forty-five minutes to an hour.

The fize and repetition of the clyfters were determined by our experience of what the rectum would retain; the laudanum being added to increase its retention, as well as to allay irritation, and to stimulate the living energy. In a few days the retention of the rectum improved, and enabled us to enlarge the clysters to 10 oz. of broth and three yolks of eggs each; to which, from the 12th of November, 8 oz. of white wine were added; the laudanum being increased to 60 drops for each of the clyfters in the day, and to 150, 180, 200, and 250 drops for the evening clyfter. Thus the whole of his nutriment for twenty four hours confifted of 30 oz. of broth, 24 oz. of wine, 9 yolks of eggs, and from 250 to 380 drops of laudanum, by clyfter; with what liquid might be supposed to be taken up by the absorbents of the surface in the bath. Mr. M. was very tall, and naturally corpulent. In health he had weighed 24olbs. But imperfect nutrition had gradually reduced him, and at the time of commencing this process, he only weighed 179lbs. On the 20th of November he was reduced to 154lbs.; on the 24th to 149lbs. This was the last time we had an opportunity of weighing him. He feems to have wasted about 5lbs. in four days, and on the fecond of December, when his delirium commenced, he probably weighed 138 or 139lbs. having loft upwards of 100lbs. of his original weight. He lived till the 6th of December, but from the 2d, the rectum no longer retaining the clysters, they were of course omitted, as well as the bath.

About the 25th and 26th of October, there was reason to suppose that Mr. M. would have had a shorter

a shorter date. He had swallowed a very small portion of nutriment for the preceding eight days, and the plan of nutrition just mentioned, though adopted in part, was not fully established. His tongue and lips were at this time become dry; a blifter applied to the fternum, discharged a thick coagulable lymph only; and his urine, reduced to a few ounces in the twenty-four hours, was become extremely high coloured, and almost intolerably pungent. In a few days however this state of the tongue went off; the skin became perfectly foft, and was often covered with a very gentle perspiration; the urine flowed without pain, and of the natural appearance, the quantity being from 24 oz. to 36 oz. in the twenty-four hours-more than the whole of his loss of weight in the same space of time.

Mr. M. had in general a stool after every fifth or fixth clyster, that is, three or four times a week. The substance of these stools was solid, and nearly of the natural scetor, and the mass perfectly homogenous. The colour was lighter than natural, as is the case in obstructions of the bile duct; their whole appearance resembling what we find in cases of dissection in the termination of the lleum or beginning of the Colon, with the exception of being much less liquid. The concoction of the Fœces, if I may so speak, was much more perfect

perfect towards the end, than at the beginning of this period; at first they looked curdy, but at last they approached to the natural appearance.

The heat, examined by the thermometer, was natural, and nearly uniform from first to last. After the clysters there was generally some slushing of the cheek, and drowsiness, with some increase in the warmth of the extremities; but under the axilla there was seldom any variation, and when it rose there a single degree, it was speedily lowered by a gentle diaphoresis.

The pulse, during the period of the scarcity and pungency of the urine, was diffurbed, and more frequent. It afterwards subsided to the natural state, and during the whole of the month of November, was not to be diftinguished from that of a person in health. On the morning of the first of December it became small and frequent; and still more frequent, though stronger, after the delirium commenced; the state of the pulse, as well as almost all the other fymptoms, from that time forth, very nearly resembling the symptoms of the last stage of fever, when it terminates fatally. During this delirium, a perpetual and indiffinct muttering occurred, with great reftlessness and agitation; the furface and extremities were fometimes of a burning heat, fometimes clammy and cold. The eyes loft their

their common direction, the axis of each being turned towards the nofe. In this state however the fensibility of the retina was not impaired, but rather encreased, for he screamed out on the light of the window being admitted, to which before he had been accustomed. At this time also the sense of touch feemed more than usually acute, for he appeared diffurbed with every accidental breath of air. The delirium, and the derangement of vision commenced nearly together, but we observed the derangement of vision first. On the first of December he complained that he faw objects fometimes double; but it was not till the fucceeding day that any confiderable incoherence of mind was observed. The pulse became feeble and irregular on the fourth; the respiration, which had been fingularly undiffurbed, became laborious; the extremities grew cold; and in ninety-fix hours after all means of nutrition, as well as all medicine had been abandoned, he ceased to breathe.

Mr. M. complained very little of hunger. Occasionally he expressed a wish that he could swallow, but not often, nor anxiously; and when questioned on the subject of his appetite, always declared that he had no hunger to occasion any uneafiness. The clysters evidently relieved the sense of hunger, and the opium they contained, feemed to have a principal share in producing this relief. It occasioned

quiet and rest after each clyster, and allayed every kind of desire or appetite.

Neither was Mr. M. much disturbed with thirst. This sensation was indeed troublesome during the first days of his abstinence; but it abated, and, as he declared was always removed by the tepid bath, in which he had the most grateful sensations.

Mr. M.'s spirits were uncommonly even, and his intellect perfectly found. He occupied himself a good deal in his private concerns, and as usual interested himself in public affairs. As we discouraged all unnecessary exertion, he spent a great part of his time in bed; but till the last few days of his life, he dreffed and undreffed himfelf daily, and walked, not only about his room, but through the house. His nights were quiet, and his sleep found, and apparently refreshing. Towards the end of this period, he faid he had very lively dreams, but they were all of a pleafant nature. The last conversation I had with him was on the morning of the fecond of December. He told me that he had had a very gay evening with two Yorkshire Baronets, whom he named; they had pushed the bottle about freely; many jokes had paffed, at the recollection of which he laughed heartily, a thing uncommon with him. When I observed that this was a very lively

lively dream, he feemed to endeavour to recollect himself, and for a moment appeared sensible that the scene had passed in sleep. But recurring to the circumstances, he spoke of them again as realities, and I made no further attempt to undeceive him. He faid he had been walking out that morning, and was pleafed to fee the spring so far advancedhe had never felt the air so refreshing. At this time, though his pulse was become rapid, his refpiration was natural, and his speech distinct, and without effort. He was extremely weak, and scarcely able to turn himself in bed, to which he had for feveral days been entirely confined. No man had ever perhaps approached death by fteps more easy; and it was earnestly to be wished that his remaining moments might be few. This wish was not gratified. In about an hour afterwards his delirium became complete, during which his strength evidently increased, and the struggle commenced which has already been described, and which lasted ninety-fix hours. He died in his fixty feventh year.

I have been the more particular in noting this case, because I have not found the methods of supporting life under impaired or obstructed deglutition, sufficiently detailed, nor the death from inanition described with sufficient accuracy. A knowledge of the first may on many occasions be impor-

tant; and an acquaintance with the last may possibly throw light on some difficult questions that respect not merely the vital, but the intellectual phenomena.

I have not found any case deserving of credit, in which life was fo long preferved without the aid of the stomach, two excepted, one recorded by Ramazzini,* and the other mentioned by Mr. Cruikshanks. The case described by Ramazzini was that of a nun; and the obstruction was not mechanical, but spasmodic; or as he himself supposes, arising from a paralysis of the Oesophagus. She is faid to have lived fixty-fix days without fwallowing, the whole of her nutriment confifting of a fingle clyfter of broth, and two yolks of eggs, administered daily. The dysphagia, which came on fuddenly, feems also to have gone fuddenly off, and the finally recovered. The case is not given with fuch minuteness by Ramazzini as to inform us whether any attempts at fwallowing were made during the dysphagia. As the obstruction was not mechanical, it is probable that fuch attempts were made daily; and if this be supposed, a suspicion may be entertained from the nature of the affection,

S that

^{*} Ramazzini, Opera Omnia Genevæ, 1717, p. 173. Many hundreds of fabulous relations of long continued abstinence may be found, some of which are noticed by Haller.

⁺ The Anatomy of the Absorbing Vessels, London, 1786, p. 101.

It is observed by Ramazzini, that the sufferings of his patient were so great on the 30th day, that she was anxious for death, and refused to admit the clyster, which however was administered during Syncope, and with immediate and happy effects. In this respect her situation was perfectly contrasted by that of Mr. M. who enjoyed ease and serenity of mind till the last final struggle, as has already been mentioned.

The case of Mr. Cruikshanks is not given at length, but alluded to in the following words. "That the furface of the skin absorbs other fluids " which come in contact with it, I have not the " least doubt. A patient of mine, with a stricture " of the Oesophagus, received nothing, either solid " or liquid, into the stomach for two months; he " was exceedingly thirsty, and complained of mak-" ing no water. I ordered him the warm bath for " an hour evening and morning for a month; his " thirst vanished, and he made water in the same " manner as when he used to drink by the mouth, " and the fluid descended readily into the stomach." It is to be regretted that a case so curious is not given in detail. It does not appear what other means of fustenance Mr. Cruikshanks employed befides the tepid bath, especially during the month of abstinence, in which the bath itself does not feem

to have been used. On some suture occasion it is to be wished that he may supply these deficiencies; especially as the information I am about to give, will shew that the conclusion he draws from this case, of the absorption of the skin, is fallacious.

Notwithstanding what I had observed formerly, I fully expected in the case of Mr. M. an increase of weight from the immersion, because he always expressed great comfort from the bath, with abatement of thirst, (which however was never great) and because subsequent to the daily use of it, the urine had flowed more plentifully and become less pungent. I expected it also from the authority of general opinion, and was curious to know the extent of the absorption. I weighed him three different times. The first time he was weighed with his cloathes on, before he went into the bath; and naked when he returned from it. Our process was therefore complicated. He appeared to have gained 30 oz. in weight, which gave him much comfort; and this circumstance prevented me from undeceiving him when I afterwards discovered the error.* The fecond time he stepped perfectly naked upon Merlin's balance immediately before immersion, and again immediately after it, his body being previoufly

^{*} I discovered that we had been incorrect before the fecond experiment.

oufly dried. The weights were never moved. The refult furprifed me: I could not diftinguish the flightest variation in the weight of his body, though my beam would have detected a single drachm. I repeated this experiment four days afterwards, and with a similar result—On both of these occasions the immersion continued an hour, and a constant friction had been kept up nearly the whole of this time on the inner surface of the thighs, with the view of increasing the action of the absorbents.

It is worthy of observation that there was neither increase nor decrease of weight in the bath; yet Mr. M. was at this time wasting twenty ounces in twenty-four hours, and confequently if there was no absorption in the bath, it might be supposed that he would have loft 5-6ths of an ounce during the hour of immersion. It is the more remarkable that there was no decrease, because in the bath the skin always foftened, and the forehead became covered with a gentle moisture, which probably arose from the condensation of the vapour from the bath. At the time however I confidered this as the matter of perspiration, because after leaving the bath, and going into bed previously warmed, be usually fell into a gentle and general diaphoresis, the soother of every irritation, and the harbinger of refreshing sleep. So far the case is tolerably complete. I lament that I cannot

I cannot give the appearances on diffection, as we were not permitted to examine the body after death.

The reflections suggested by this case are numerous: I shall present one or two that seem important.

The natural state of the pulse during the first month of Mr. M.'s abstinence, when his strength was daily decaying, demonstrates, that the motion of the heart and arteries gives no certain indication of the strength of the system. Pure debility, within certain limits, does not seem to produce a frequent pulse, nor in all cases, a feeble one; but when carried to a certain degree, it excites that commotion of the system, in which—first the contractions of the heart, and afterwards of the lungs, become more rapid; which in popular language may be denominated the last struggle of nature, and which may be considered as indicating the approach of death.* To endeavour to reconcile these phenome-

S₃ na

^{*} In a recent case (April 1798) where deglutition was obstructed from a different cause, the same methods of supporting life as in the case of Mr. M. were resorted to. But the tepid bath was soon abandoned, the patient conceiving that it irritated a cough to which he was subject. In this instance,

na to any of the prevailing theories of vital motion, would be a fruitless attempt. Death has not an uniform aspect, but its phases do not appear to me numerous. The concluding scene in the life of Mr. M. may illustrate the form of its approach, as well as the nature of the symptoms in the last stage of sever, and of various other diseases, in which similar symptoms occur.

The natural state of the heat, during the inaction of the stomach, might prove, if it were necessary, how little the vital heat is connected with the function of digestion, and renders it probably, that the general increase of heat which follows the ap-

plication

as in that of Mr. M. the approach of death was gentle and eafy. The patient, who was aware of his real fituation, met his fate with perfect equanimity, and possessed his understanding to the last. Though the pulse became more frequent two or three days preceding his death, yet there was little or nothing of that struggle described in the case of Mr. M. This gentleman had long been wasting under an incurable disease, and had taken little nutrition for some time previous to the commencement of his abstinence.-He lived twentythree days after he defifted from all attempts at deglutition. He never complained of hunger, and the thirst which he felt occurred at intervals only, and was not very fevere. The case was unfavourable to more particular observations; the patient having been long fubjed to a difease of the system by which the epiglottis and the furrounding parts were finally destroyed, his symptoms were of a mixed nature.

plication of many stimulating bodies to this organ, is not the direct effect of the action of the stomach itself, but of its influence on the heart, or diaphragm, or both. It is afferted by Haller, that in cases similar to that of Mr. M. the heat has been found diminished. The observations made by the thermometer in this case, throw a doubt on the affertion; while the great facility with which Mr. M. parted with his heat, concurs with a variety of corresponding sacts to support the position, that in those situations, where the generation of heat is natural, the power of retaining it in the system is in proportion to the force of the living principle.

But what shall we conclude respecting the abforption of water by the surface of the body in the
warm bath?—Admitting that it did not take place
in the case of Mr. M. it may be supposed this did
not arise from the impermeability of the cuticle,
but from the heat of the bath being too high. It
may be added, that the analogy of the vegetable
kingdom supports this supposition, since vegetables
are found to exhale in warm and dry air during the
day, but to absorb largely under the moisture and
coolness of night. Considerations of this kind
might probably influence some learned physicians in
London who were consulted in this case, to propose
that the heat of the bath should be as low as 80°.—
It was however raised to 96° for two reasons: the

first.

S 4

first, that I had found no increase of weight in the Buxton bath at 82°; the second, that as Mr. M. parted with his heat easily, he could not support immersion with comfort in a lower temperature.

It is however more likely to be supposed, that though there was no increase of weight in the case of Mr. M., there might be an absorption in the bath which was countervailed by the increased exhalation.-The experiments I have detailed do not enable us to decide this point absolutely; but from them, and from various other confiderations, I entertain little doubt, that though the exhalants of the skin pierce the epidermis, and come into contact with the external air, the mouths of the absorbents terminate under it, and are covered by it; and that while it remains found and entire, no abforption of folid, liquid, or aeriform elaftic fluid, takes place on the furface. In the inflances that are supposed to favour the contrary opinion, it will be found, that the article absorbed is forced through the epidermis by mechanical pressure; or that the epidermis has been previously destroyed by injury or difease; or if found, that the article applied to it is of an acrid nature, which first erodes this tegument, and then coming into contact with the mouths of the lymphatics under it, is of course absorbed.*

At

^{*} The abforption of mercury, fulphur, oil of turpentine, and fimilar fubflances, rubbed on the skin in the form of un-

At the instant that these observations are committed to the press, my attention has been pointed to the valuable memoir on this subject read before the Academy of Sciences by M. Séguin, of which some account is given by Fourcroy in the third volume of "La Médicine eclairée par les Sciences physiques, p. 232." The conclusions of M. Séguin correspond in every particular with those which I have already laid down, and being sounded on a great variety of experiments made for the express purpose of determining on the cutaneous absorption, they appear to me to give his inferences all the solidity of truth. The objection, that though there

guents, comes under the first of these descriptions; the absorption of the fame fubstances, and in some rare instances of lead from the furface of wounds, and of herpetic, or other eruptions, under the fecond; the absorption of cantharides, aloes, &c. under the third. I am not ignorant of the experiments of Dr. Falconer and Mr. Albernethy, nor wanting in due refpect for those gentlemen; but my own observations, supported by the experiments of M. Séguin, induce me to reject their conclusions. A probable fource of error in the experiments fupposed to afford a direct proof of cutaneous absorption, is, that they were made on a fmall part of the body, e.g. the arm or leg, and not on the whole; though if the mouth and noftrils were left free for respiration, it would be as easy, and much more correct, to make experiments for this purpose, as well as for determining the quantity of exhalation, on the whole of the furface, as on a part. Dr. Marcard, who wrote expressly on the effects of the bath, has adopted the theory of absorption through the skin, without making any experiments.

there is no increase of weight in the bath, there may be an absorption, which is countervailed by the exhalation, M. Séguin found a fatisfactory mode of invalidating. He dissolved in the water of the bath, in which he made his experiments, fubstances which produce a specific effect when received into the fystem, by which their inhalation might be afcertained. He employed the oxygenated muriat of mercury in folution, on a number of venereal patients, and while the epidermis was entire, he never perceived a fingle instance of falivation, or even of amendment of their complaints. But in cases where a considerable portion of epidermis was injured or destroyed, as in the itch, the specific effects of mercury on the system were produced. The experiments of M. Séguin are very numerous, and appear to be devifed fo as to meet every objection. He concludes from the whole, that while the epidermis is entire, the absorbents of the furface take up neither water nor air, nor any fubstance diffused or diffolved in either.

Admitting that the inferences of the friend and coadjutor of Lavoisier are just, how are we to explain the refreshment and abatement of thirst, experienced by Mr. M. in the bath. Referring to what is said in pages 177, &c. on the nature of thirst, we may here shortly observe, that as liquids thrown into the stomach relax the exhalants of the surface.

furface, so doubtless this action is reciprocal; and that the exhalants of the surface being relaxed in the warm bath, a relaxation in those of the stomach ensues, by which the sense of thirst is abated. In the cold, as well as in the hot state of intermittent sever, the thirst is great; but it is always lessened, and generally removed, on the accession of the sweating stage, though no liquid should have been swallowed.

Will this explanation, it may be asked, account for the benefit derived in accidents at fea, when no drink can be produced, from immerfing the body in water, or applying wet fubftances to its furface? It will explain it in part perhaps, but not wholly .-In the torrid zone, where the greater part of these inflances have occurred, the immersion of the body in water, or the application of wet clothes to the skin, will keep the surface cool, by defending it against the direct rays of the fun, and thus prevent the debility produced by an accelerated circulation; it will also defend it, in whatever climate; from the wasting effect of evaporation, and thus be a means of prolonging the strength. This reasoning is supported by a fact which may be inferred from the circumstances I have mentioned, and which is established by the experiments of M. Séguin, that the body wastes much less in water than in air.

In the Buxton bath I found my weight very flightly diminished; but in the case of Mr. M. and in all our experiments on diabetic patients, there was neither diminution nor increase in the bath. In the experiments of M. Séguin, there was always fome diminution in the bath of whatever temperature, which he ascribes to the pulmonary perspiration. How is this difference accounted for? Though the cutaneous exhalation may have been stopped in Mr. M. during immersion, yet allowing the perspiration by the lungs to have gone on, this alone, fupposing no absorption, ought to have produced a diminution of weight in the bath. To this it may be replied, that though no absorption through the skin takes place in the bath, that by the lungs goes on, while in the heat of 96°. the evaporation from the furface of the water, by loading with moifture the air inhaled, prevents its taking up the moifture in the lungs, and thus diminishes the pulmonary perspiration.-Against this source of error it is probable that M. Séguin guarded.* It is in this way that the increase of the weight of the body in moist air.

^{*} I fay probable, because I have not seen the memoir of Séguin himself, but only the abstract of it given by Fourcroy, who does not enter into the details.—An account of this abstract is given very accurately, and pretty much in detail, in the Monthly Review Enlarged, vol. xii. p. 514. I have subjoined an extract of the article of Mr. Fourcroy to this second edition. See Appendix, No. III.

air, which is faid fometimes to occur, and which has been supposed to prove a cutaneous absorption, may be accounted for. In moist air the absorption of the lungs from the atmosphere goes on unchanged, while the pulmonary, as well as the cutaneous perspiration, is diminished, and thus the increased weight of the body may be explained. On this point however more accurate experiments are required.

It was observed, that the discharge by urine alone, in the case of Mr. M. exceeded much in weight the waste of his whole body; and it cannot be doubted that the discharge by stool and perspiration exceeded the weight of the clysters. Thus it appears that the egesta exceeded the ingesta, in a proportion much greater than the waste of his body will explain. How, it may be said, is this accounted for, unless by cutaneous absorption? In the same way, it may be answered, by the action of the lungs.

That the oxygen inhaled in respiration from the atmosphere unites in part with the carbon of the blood, to form carbonic acid gas, and in part with the hydrogen, to form water, is generally admitted. In instances of inanition we may suppose the proportion in which it unites to these different substances to be varied by the powers of life; that the formation

formation of water is greater than usual, and that of carbonic acid gas less. Thus a portion of the oxygen, which, in the ordinary course of things is exhaled in the form of an elastic vapour, being absorbed in the form of water, will serve to retard the waste of the body, and to surnish the blood with the proportion of lymph, which affords a proper supply to the secretions of urine and perspiration. In the case of Mr. M. there was no appearance of that acrimony of the sluids which might naturally have been expected from the cessation of the supply of chyle, and which Haller describes as the constant effect of inanition.

This theory of the increased formation of water in the lungs has been employed by Dr. Rutherford of Edinburgh, to account for the vast discharge by the bladder in diabetes. It seems also to account for the phenomenon of the egesta exceeding the ingesta, so often observed in that disease. It illustrates the doctrine of Séguin, in the memoir already quoted, of which Fourcroy gives an account in the following words—Le diabète ne provient point de l'eau absorbée dans l'air par la peau, mais du ressur de celle qui ne peut pas être enlevée aux poumons par l'air trop chargé d'humidité—a doctrine which appears to be desective, from not pointing out the source of the increased quantity of water in the lungs. Beautiful as the theory of Dr. Rutherford

is, it cannot be confidered as established in diabetes. or in inanition (to which I have extended it) till the air expired by patients under these diseases shall be examined. In the mean time I may observe, that the application of it to the case of Mr. M. accords with the usual economy of the living principle, which in remedying the defect of particular organs, does not transfer their powers entire to other parts of the system, but modifies or increases the natural action of some other organ, by which that deficiency may be to a certain degree countervailed. Such an exertion of the living principle was evident in the case of Mr. M. in the increased powers of the rectum, which have already been described. These vicarious actions of the fystem are however defective, whether arifing spontaneously, or by the affistance of art. No applications of food to the inner furface of the rectum can long fupply the absence of food in the stomach, nor can the want of liquid in that organ be long countervailed by applications of moisture to the surface, aided by the modified action of the lungs. After a certain time, though immerfed in water to the chin, the thirst will prove intolerable without drink, as the shipwreck described in the Appendix will testify, (Appendix, p. 24) and the fufferings of Tantalus,* which the theories of

Dr.

^{*} Dr. Marcard, in the work quoted, p. 135, after afferting the great inhalation of water by the skin in the bath, declares, that no naturalist will estimate the sufferings of Tan-

Dr. Marcard led him to deride, will be rated as highly by the naturalist as by the poet.

If the non-absorption of the surface of the body be established; it will ascertain, that contagion is received into the system, in the ordinary course of things, by the lungs only; and perhaps justify a practice common, as I am informed, among our more experienced seamen on the coast of Guinea, and other warm climates, who when exposed during the night to a breeze from the marshes, wrap their heads in a sea-cloak, or other covering, and sleep fearlessly

talus highly. It feems probable, that immersion in tepid water is more effectual in abating thirst, than immersion in cold water.

The account I have given of Dr. Rutherford's theory of diabetes, I derive from the valuable inaugural differtation of Dr. Marcet, of Geneva, just published, presented to me by the ingenious author. To Dr. Marcet I am also indebted for pointing out the memoir of M. Séguin, whose doctrine he himself has controverted, from giving, I apprehend, more weight to the authority of Haller, than on inquiry, it will be found to deserve on this subject.

The explanation offered by Dr. Marcet of the increased heat in diabetes will be found I believe irrelevant.—In three cases of this disease which I have examined, the heat previous to the medical treatment was less, not greater than natural, notwithstanding the almost unanimous concurrence of authors in the contrary statement; a proof among many others of the inaccuracy of the accounts we have of temperature in disease.

fearlessly on the deck with the rest of their bodies nearly naked, in consequence of the heat.

It is with pleasure I lay down my pen, thought other observations crowd upon me. If what I have already offered seem tedious, the interest of the subject will plead my apology. It is no otherwise connected with the rest of the work, than as it regards the most important functions of the living principle.—But I have not been very studious of rigid method in this volume, which is enlarged beyond my original design.—Should the facts which it contains deserve and obtain attention, the fate of the reasonings by which they are accompanied will be a matter of little anxiety. In the present state of our science the enlightened reader will thank me for avoiding systematic arrangement; and recommending myself to his candour, I bid him adieu!

fearlefely on the deel with the reft of their bodiers nearly naised, in suplequence of the bear.


It is with piculine it has doors enough to thought or her observed of the following the arrest of the following the arrest of the following the arrest of the following the training particle of the following that will plead my appellage. It is no concretife door nather meth integers of the fiving principle. The meth integers of the fiving principle.

The meth integers of here very finalizes of rigid method?

Journal Religiouses which is endinged involved my original Religious of the realized matter. There and obtain accompanied that hate and the realized of leader which the realized of leaders and the realized of the contains of leaders and the arrest of the method of the standard of the method of the standard of the method of the standard o

APPENDIX.

Nº. I



Of Tetanus, and of Convulfive Diforders.

Read before the Medical Society of London, May 10, 1790, and extracted from the 3d volume of their Memoirs.—Page 147.

THE various objects which engage your attention will not admit of much of your time being employed on a fingle paper, and I have undertaken to write on a subject of some importance. It will be proper for me therefore to come directly to the point, and to avoid as much as possible every thing that is superfluous. What I have to offer shall be confined to the treatment of convulsive disorders, and especially of tetanus.

For this difease, so terrible in its progress, and so generally fatal in its issue, several new remedies have been proposed within the last thirty years, and each in its turn has had some share of public considence. But opium, mercury, the cold bath, and wine, are those only which seem entitled to any reputation, and of each, it has been my fortune to have had some experience. This experience I proceed to give you with all the clearness and faithfulness in my power.

T₂ In

but

In the course of nine years practice I have seen seven cases of this disease; of which two were idiopathic, and five the consequence of wounds. Of these in order.

1. George Gardner, a foldier in the Staffordshire militia, was put under my care by his officers, on the 20th of February, 1781. About a fortnight before, after fevere dancing and hard drinking at a country wedding, in which he had been employed two days and nights, he fell fuddenly into a fit, which lasted an hour and a half, during which his confciousness was abolished. On recovery he was affected with flight twitchings, which gradually encreased, and were followed by fixed spafmodic contractions in different parts of the body, but more affecting the left fide than the right. He had, when I saw him, all the symptoms of tetanus. The head was pulled towards the left shoulder, the left corner of the mouth was thrown upwards, the eyes were hollow, the countenance pale and ghaftly, the face and neck bedewed with a cold fweat; but his most distressing fymptom was a violent pain under the enfiform cartilage, with a fudden interruption of his breathing every fourth or fifth inspiration, by a convulsive hiccup, accompanied by a violent contraction of the muscles of the abdomen and lower extremities. He felt on this occasion as if he had received an unexpected blow on the scrobiculus cordis. Before I faw him he had been bled and vomited repeatedly, and had used the warm bath, not only without alleviation but with aggravation of his complaints. The three first remedies mentioned were used here in succession, viz.opium, mercury, and the cold bath.

He first took a grain of opium every other hour, afterwards a grain every hour, and at last two grains every hour; but he grew worse and worse during the two days on which this course was continued. The spasms extended to the back and shoulders, the head was at times retracted, and the muscles of the abdomen partook of the general affection. Being no longer able to fwallow the pills, he took no medicine of any kind on the night of the 22d, in the course of which general convulsions came on, and returned once or twice in every hour. The tincture of opium (liquid laudanum) was now directed to be given, and an ounce of the quick-filver ointment to be rubbed in on each thigh. In twenty-four hours he took two ounces and a half of the tincture without fleep or alleviation of pain. The dose being increased, in the next twenty-fix hours he swallowed five ounces and a half of the laudanum, a quantity which, at that time, was I believe unexampled. He lay now in a state of torpor. The rigidity of the spasms was indeed much leffened, and the general convultions nearly gone; but the debility was extreme; a complete hemiplegia had fupervened; the patient's eyes were fixed, and his speech faultering and unintelligible.

As this young foldier appeared on the utmost verge of life, it seemed no longer safe to continue the laudanum, which had relieved spasm only in so far as it had brought on general paralysis. Intermitting this medicine therefore, we gave small doses of camphor from time to time in a liquid form, but the chief attention was directed to supporting the strength by such nourishment as could be swallowed. Gruel, with a small quantity of wine, was ordered for him, though with much caution; for at that time I was not instructed with what safety and efficacy this last article might have been administered. For the next six days he seemed to revive: the general convulsions kept off, though the

twitchings and convultive hiccup continued. But on the night of the first of March he was seized, during sleep, with a convulsion as severe as ever, and this was followed by a return of all his symptoms with their former violence. The jaws were indeed more completely locked than before, deglutition was become impossible, and the pain under the ensistent cartilege was so extreme as to force from the patient the most piercing cries. At this time the effects of the quicksilver ointment were apparent in the sector of the breath, and in a considerable salivation.

Had poor Gardner been a man of any rank, or indeed had he been furrounded by his family, it is most probable that we must now have abandoned him to his fate. But our proceedings being obstructed neither by the prejudices of ignorance, nor the weakness of affection, another, and a last effort for his life was refolved on. Having heard that the cold bath had been employed with fuccess in tetanus in the West Indies, particularly by Dr. Wright of Jamaica, and Mr. Cochrane of Nevis, and this practice corresponding with certain speculations of my own, I had recourse to it on this occasion with some little confidence. With the confent of his officers. Gardner was carried to the public faltwater baths of this town, then of the temperature of 36° Fahr, and thrown headlong into it. The good effects were instantaneous. As he rose from the first plunge, and lay struggling on the furface of the water, supported by two of his fellow-foldiers, we observed that he stretched out his left leg, which had been for some time retracted to the ham. But his head did not immediately recover the fame freedom of motion, and therefore he was plunged down and raifed to the furface fucceffively for upwards of a minute longer, the muscles of the neck relaxing more and more after every plunge.

plunge. When taken out, we felt fome alarm: a general tremor was the only indication of life, the pulse and the respiration being nearly, if not entirely, suspended. Warm blankets had however been prepared, and a general friction was diligently employed. The respiration and pulse became regular, the vital heat returned, the muscles continued free of constriction, and the patient fell into a quiet and profound fleep. In this he continued upwards of two hours, and when he awaked, to the aftonishment of every one, he got up and walked acrofs the room, complaining of nothing but hunger and debility. The convultive hiccup indeed returned, but in a flight degree, and gave way to the use of the cold bath; which he continued daily a fortnight longer; and in less than a month we had the fatisfaction of feeing our patient under arms, able for the fervice of his country.

That the opium, though it failed in effecting a cure, had confiderable influence in mitigating the difeafe, and prolonging life, is, I think, apparent.

That the mercury had little effect is clear from the fecond convultions coming on foon after the falivation appeared.

The fuccefs of the cold bath in circumstances so apparently desperate was not lessened by bad effects of any kind. Though the patient was in a falivation when thrown into it, yet this was not stopped suddenly; it lessened indeed immediately, and foon disappeared, but without any of those bad consequences so well described by Sir John Sylvester and Dr. Dobson. Subsequent experience has taught me to attribute some part of the suddenness of the

benefit obtained in this instance to a circumstance that distressed me much at the moment. The very instant that we were about to immerge poor Gardner, he was seized with a general convulsion. We hesitated, but kept our purpose, and happily plunged him into the water with the convulsion upon him. I am also inclined to think that our success is in part to be ascribed to the powerful, general, and sudden application of the remedy, and under this opinion I should prefer immersion in water, to its affusion, the method employed by Dr. Wright.

2. Soon after this I was sent for by a poor woman, who, in consequence of difficult labour, and, as she imagined, of local injury in some part of the uterus, was seized with the spasmus cynicus, locked jaw, and other symptoms of tetanus. She was immediately taken to the cold bath, and thrown into it in the same manner as the former patient, and with similar good effects. The spasms disappeared, and though they afterwards returned to a slight degree, they gave way entirely to a second immersion.

In the first of these cases I was affisted by Mr. Walker, surgeon's mate of the Staffordshire militia; and in the second case the effects of the cold bath were witnessed by him and my friend Mr. Park.

So far of the idiopathic tetanus; my experience of the effects of the cold bath in the disease originating in wounds is neither so satisfactory nor so complete.

3. The first case that occurred to me was that of a master of a vessel returning from sea in the year 1784. The injury had been received on the leg, but the wound had little

little inflammation on it, though the patient was fo far advanced in the disease as to go off in a general convulsion, before any remedy could be tried.

- 4. The second case seemed rather an instance of locked jaw, properly so called than of tetanus. The injury having been received on the jaw itself, the affection was confined to the muscles of that part, and the disease seemed to be the trismus inflammatorius of Sauvages. It gradually yielded to bleeding and the topical application of warm vapour, the cold bath producing no good effect.
- 5. In the third case we contented ourselves with pouring cold water in successive buckets on the patient's head, partly because this was the most convenient method, and partly because it is that employed by Dr. Wright, whose paper on the subject in the fixth volume of the Medical Observations and Inquiries, had by this time appeared. While we were bufy in this way, and flattering ourselves with some appearance of fuccefs, the unhappy patient fuddenly threw himfelf on his back, and a quantity of water entered his mouth. The effects were highly diffreffing. The muscles of deglutition were unable to carry down or to expel the fluid, and fuch convulfions took place as led us to expect inftant death. He recovered, however, fo far as to regain his former fituation, but the effects of his fufferings on the bye-standers prevented our continuing this practice, and he was carried off a few hours after.
- 6. A fourth instance of this disease I saw about two years and a half ago. There was no room for new practice, for the patient was too far advanced. He had been treated with opium and mercury as usual, and as usual died.

These cases afford little inference either in favour of or against the use of the cold bath in tetanus arising from wounds, but they are stated shortly, that the account of my experience in this disease may be fair and complete.

It does not appear from Dr. Blane's Hiftory of the Difeafes of the Fleet, that this remedy was used in any of the cases of tetanus that occurred among the seamen in the late war. In fpeaking of this difease Dr. Blane indeed mentions, that though Dr. Wright and Dr. Cochrane had found the cold bath fuccessful in cases without local injury, they had acknowledged the practice to be ineffectual in the difeafe when arifing from wounds. Dr. Blane may perhaps be accurate respecting Dr. Cochrane, whose publication on this fubject I have not been able to procure, but he is miftaken respecting Dr. Wright. This gentleman, in the paper already quoted, informs us, that fince he employed the cold bath, he has not failed in effecting a cure in a fingle instance of tetanus, without distinction or reservation; and of the fix cases which he has detailed, in three the difease appears to have arisen from local injury. This point is of importance, and a mistake respecting it in a work like that of Dr. Blane's, which will no doubt be much confulted in the future practice of the fleet, may, if paffed unnoticed, have ferious confequences.*

It is worthy of remark, that the use of the cold bath in tetanus is a practice of very ancient date. Dr. Wright, whose precision and candour are exemplary, mentions that he received the first hint of it from Dr. Lind of Haslar. Hippocrates, however, was acquainted with it, as appears by Aphor.

^{*} This is cerrected by Dr. Blane in his fecond edition,

Aphor. 21. lib. 5. It is also noticed by Avicenna, lib. 3. cap. 7. and in the collection of Schenkius, p. 120, an account of the success of this practice in two cases of tetanus is quoted from Jason, cap. 21, De morbis cerebri, ex Valesco. Hippocrates indeed, and after him Avicenna, restrict the use of this remedy to the young and corpulent, and to the summer season. They also confine it to cases in which the disease has not originated in local injury, for it appears, by another of his aphorisms, that the father of physic thought convulsions supervening to a wound, always fatal*. The experience of Dr. Wright encourages us to extend this remedy to tetanus originating in wounds, but if it should fail we are still not to despair, as the following case will testify.

7. In July 1787, a labouring man was brought into our Infirmary with a wound in his hand. He was under my friend Mr. Park's care, and the fore was nearly healed, when an alarming rigidity appeared in the motion of the jaw. Opium and mercury were immediately prescribed, with which were conjoined first the cold, and afterwards the warm bath. Mr. Park, however, finding the disease to proceed with the most unfavourable omens, called a confultation, at which, in the absence of the rest of our colleagues, we agreed to make a trial of bark and wine on the authority of Dr. Rush of Philadelphia, whose excellent paper on this fubject, in the fecond volume of the American Philosophical Transactions, had just appeared. We had, however, little or no hope, for the disease was far advanced, and its progress had been very rapid. At this time the jaw was not only rigidly contracted, but the spasms had extended to the neck and back, the pain under the enfiform cartilage was most

acute

^{*} Hippoc. Aph. lib. iv. fect. v. aph. 2.

acute, and twice or thrice in every hour he was feized with general convulsions, each of which lasted about half a minute.

The patient was, however, a man of a vigorous mind, and in his perfect fenses; and his danger was not concealed from him. Death, he was told, must be the inevitable confequence, unless he swallowed wine in large quantity, but this we verily believed would fave him. At first bark was infused in the wine, but he could not swallow the mixture, and therefore we trusted to wine alone. It was wonderful to fee the exertions which this poor fellow made. If the liquid was offered to him at an improper time, the effort of deglutition brought on a general convultion; nay a general convulsion was the consequence of advancing it at such a time towards his head. But watching the remission of the fpasms, he was able to swallow a table spoonful or more at once, he himself giving the fignal when the wine should be administered.* In this way, through the opening made by the lapping of the upper jaw over the under one, he drew up and swallowed a quart of port wine in the course of two hours; at the end of which time he thought himfelf refreshed, and was encouraged to proceed. At the end of twentyfour hours he had finished his third bottle, and at this time it was evident that the down-hill progress of the disease was checked. Though the pain under the enliform cartilage (the most excruciating of all the symptoms of tetanus) was little abated, yet he felt himself more able to bear it; and the general

^{*} This circumstance suggested to me the similarity between this case and hydrophobia. I noticed it to my respected friend Dr. Percival, who has mentioned it in his valuable essay on that disease. v. Percival's Essays, vol. II. p. 366. 4th edition. I differ however from this experienced physician in his supposition that the two diseases are the same, or of the same origin.

general convultions were certainly less frequent and less severe. We continued our plan with patience and vigour, but for a long time life and death seemed to hang in equal scales. On the sisteenth day of this course he was affected with a slight nausea and vomiting, which soon went off, and thenceforward his recovery seemed more apparent. It was not however till he had been forty-two days under this treatment that his safety could be ascertained, and during this time he swallowed a hundred and ten bottles of Port wine.

In this case bark was administered along with the wine towards the latter period of the complaint; and during the whole of it, gruel or milk in the quantity of a quart a day, or upwards, was given by way of nourishment. The milk, however, he preferred, as agreeing better with the wine, and tending, as he conceived, to allay a burning sensation at his stomach, the effect of an acid serment there.

Though the wine was given in fuch quantity, yet it never produced any fymptom of ebriety—it foothed the irritation of his nerves, and comforted his mind, and, without increasing the frequency of his pulse, it augmented his strength.

Every night he took from fixty to a hundred and twenty drops of the tincture of opium, and, with the wine, this small dose seemed to have a more composing effect than thrice the quantity taken before he began it.

To allay the pain under the entiform cartilage, clothes dipped in æther were applied externally with good effects. As I flood by him one day, preffing the doffils with my open hand to the scrobiculus cordis, he expressed himself relieved

by the force which I applied. The same relief seemed to follow from a gentle and uniform pressure on the parts suffering from spasm. In consequence of this, rollers were applied round each thigh where spasmodic twitchings had been very distressing, and afterwards, at his own request, a general pressure of the same kind was applied to the back and abdomen. These bandages he would not suffer to be removed, but we poured æther on them occasionally, over the parts most affected by spasm, guarding in the usual way against the cold produced by too speedy an evaporation.

At length our patient recovered. He is now a watchman, and calls on me occasionally. He complains of a stiffness at times in the muscles of his back, and though naturally a strong and healthy looking man, his features retain the indelible impressions of his disease. His eyes appear hollow, his face sharp and pale, his cheeks and lips are skinny, and the masseter muscles hard and shrivelled.

I have lately had an account of a case of this kind which occurred in a sailor on the coast of Guinea, who recovered under the care of Mr. Robson, an ingenious surgeon in the African trade, from a state apparently desperate, by a very liberal use of bark and ardent spirits.

To conclude this part of my subject, I have known two instances of this disease in horses, both originating in wounds; the first of these was treated in the method first described. The cold bath was used repeatedly, and large doses of opium employed in the form of clysters; but the disease terminated satally. The second of these occurred in a horse of my friend Dr. Camplin (whose affistance in the case of the watchman I ought to have acknowledged), about

about a twelve-month ago in his journey to Bath. The doctor informs me that he gave him wine and opium, mixing fixty drops of laudanum with every pint of wine, and by this practice he had the pleafure of recovering him, though not till he had drunk as much wine as he was worth. Brandy or gin may perhaps be as fuccessful in such cases.

It may be supposed, that these instances of success in the use of the cold bath and wine in tetanus, would lead me to the employment of these remedies in other convulsive disorders. This has accordingly been the case; but of the use of wine I am not yet prepared to speak. Of the use of the cold bath in such disorders I can speak with some considence, as my experience of it is now of eight years' duration.

In the convultions of children I have found the cold bath a most useful remedy, whether the disorder originated in worms or other causes. I have seldom known it to fail in stopping the paroxysins, at least for some time, and thereby giving an opportunity of employing the means fitted to remove the particular irritation. I must however observe, that in early infancy I have used it with caution; fometimes tempering the water when the weather was cold, and fometimes pouring it on the patient, rather than immerfing the patient in it. I have in general made the application of cold in this way fudden and transient, have employed means to fecure re-action, and have avoided the remedy entirely in all cases where the vital energy seemed much exhausted. With these precautions I have seen great fuccess attend this remedy in a variety of instances. Minutes of nine of these I have preserved, but I mean to give one only in detail.

John

John Slater, aged eight years, came under my care in the month of January, 1782. About two years before, while at play, he was feized fuddenly with a convulfion which continued for half an hour, and had returned ever fince at short intervals. Various means had been employed for his relief but without fuccess; the fits were become more and more frequent, a hemiplegia had fupervened, and the intellect was apparently loft. For twenty-four hours he had lain in a state of insensibility, motionless on the left fide, and the mufcles of the other fide only agitated by the convulfive tremors. We put him immediately into a tub of cold water, which inftantly stopped the paroxysm and threw him into a deep fleep, out of which he awaked after two hours with a shriek, and fell into convulsions as before. The cold bath was repeated, and afterwards continued daily, present relief being always obtained by it. The interval was employed in administering the tin powder, which was worked off with calomel, but no worms were observed to be discharged. After several days the convulsions returning, though with a confiderable abatement, I became diffatisfied with the mode in which the bathing had been performed, the fize of the tub employed never having admitted of fudden immersion. On this account we had the child conveyed to the public bath, into which he was thrown headlong, his father being stripped in the water to receive him. The temperature of the bath was 43° Fahrenheit. He was repeatedly plunged down and taken up for half a minute; was taken out of the water free of convulsion, fell immediately into a profound fleep, and awaked clear of complaint. In these respects this case resembles that of Gardner, first described. It resembles it also in this respect, that convulsion had taken place at the moment of immersi-The paralytic weakness of the fide remained, however for some time, but by the continued use of the bath it was at length entirely removed; and the powers of the mind, which had been totally suspended, gradually returning, at the end of six months were perfectly restored.

He continued well upwards of twenty months, was healthy and vigorous in body, and in the acquirement of knowledge remarkably acute. But on the 30th of December, 1783, his mother having struck him and frightened him much, he was seized in the night during sleep with general convulsion, in which he continued several hours without intermission. Being again called to him, I employed the same remedy, but at the time of immersion the convulsion was not on him. Sleep and refreshment followed as before, but the paroxysm returned in the evening, though in a slighter degree. However, by perseverance in the daily use of the bath, and throwing him twice into the water with the convulsions upon bim, in six days his health was restored.

I lament over the fate of this amiable boy. He continued from this time three years and a half in perfect health; grew strong and tall, and shewed great affection of temper and sensibility of mind. But being violently agitated with fear and grief, he fell again into convulsions in the month of July, 1787. I was sent for as usual, but was absent from town at a considerable distance. Having on sormer occasions launched him myself into the bath, his mother, a superstitious woman in a low walk of life, would not permit this remedy to be used till my return. He lay for thirty hours convulsed and senseless, and the first intelligence I had of his illness was accompanied by the account of his death.

U This

This case differs materially from the convulsions of early infancy. I give it therefore not as an instance of thefe, but as a specimen of a disease affecting children from four to twelve years of age, and which, though not very uncommon, as far as I can judge, has not yet found its place in any systematic work that I know of. The nosology of convultive diseases is indeed very imperfect, and the terms we employ in describing them are by no means precife. Those who have seen tetanus and the disease last mentioned, will know, that what is called convulsion in both instances, is in the one case and the other a very different affection. The convulsion of tetanus is a sudden and violent contraction of all the voluntary muscles, throwing the frame into firong contortions, but lafting only a minute at most, and feldom so long. The patient, where the case terminates fatally, probably dies at last, from the spasmodic affection extending to the heart, or fometimes perhaps, from the sufpension of respiration. The convulsion in the other case, though it begins with violence, soon subsides, leaving the body in a state approaching to coma: and though the separate fibres of many of the muscles keep in constant vibration, and the turbulence of general convulsion returns occasionally, yet in the intervals the respiration is natural, and the pulse little disturbed; voluntary motion and consciousness, are, however, completely suspended.* If this state continues long, the violent commotions return more frequently, the breathing becomes laborious, the strength is at length exhausted, and the patient dies with the fymptoms of apoplexy.

I conclude

^{*} This species of convulsion has been denominated elonic, in opposition to tonic, but with little propriety. All convulsions may be said to be clonic.

I conclude this paper, already too long, with a few general remarks.

- 1. It appears to me that the efficacy of the cold bath in convulsive disorders is much promoted by being employed during the presence of convulsion. How I came to be led into this opinion may be seen in the case of Gardner. Much experience is required to establish this as a general truth, and I give it as an opinion only. As madness is, I think, best combated in the height of the phrenzy, so I believe convulsive diseases are most capable of remedy in the agitation of convulsion. I seem to have a glimpse of a general principle in nature that connects these facts so apparently different; but the ground is not yet firm enough for the foundation of a theory.
- 2. It may however feem to strengthen the above remark, that in spasmodic diseases, which rise not to general convulsion, the cold bath seems to be of inferior efficacy. In Chorea Sancti Viti I have tried it frequently, but never found it of any service. This is one of the sew diseases in which electricity is of decided advantage.
- 3. In the hysteric paroxysm, the cold bath, or indeed the plentiful affusion of cold water, is an infallible remedy. Those who suppose that the terror it occasions ought, in this case, to prevent our having recourse to it, are, in my opinion mistaken. Though the hysteric paroxysm be the offspring of passion, it is never occasioned, I will venture to assert, by the passion of sear. A sense of danger will always, I believe, prevent it; or indeed a powerful dread of any kind. I have known a tub of cold water kept in readiness, with the certainty of being plunged into it on

the recurrence of the paroxysm, cure this disease, without the remedy being ever actually tried. I know the hysteric paroxysm often takes place when danger is over, but that is another case.


but my experience of its effects is as yet too recent and too imperfect to be detailed. I am not so fanguine as to expect that experience will accumulate rapidly on this subject. Many there are who will think my practice too hazardous to be copied; even after the account of its success. This relates especially to the use of the cold bath in convulsive disorders in private practice; but in hospital practice, and in the practice of the fleet and army, the same objections do not present themselves; and in these departments of the profession it may be expected that its effects will be ascertained.

must, that in topicsodic different which alle not to general

Dr. Girdlestone, in his account of the diseases among the troops in India, mentions that hot Madeira wine is given with success in the tetanus which occurs in that climate; but he tells us, though not from his own experience, that the cold bath has been very unsuccessful. To what circumstance it may be owing that a remedy, which has been so efficacious in the West Indies, should have failed in the East, does not appear; though if the circumstances under which it was employed were fully related, the difficulty might perhaps be explained. It may, however, be observed, that in the sultry climate of India, where the human frame is greatly relaxed, it will not be safe to use the cold bath with the same freedom as in more northern regions. And it will undoubtedly be more safe, in any future trials of this remedy in that country, to use the method of affusion employed.

ployed by Dr. Wright, and so successful in his practice in a similar climate, instead of immersion, which it appears was employed in the cases mentioned by Dr. Girdlestone.

I present these facts and observations to you, Gentlemen, with the confidence of a man whose intentions are good. I have not spoken under the warmth of new conceptions, for my experience is, you see, of several years' duration; and I anxiously hope that time will stamp my narration with the seal of truth.



the river Markey into the United the algert; and bear a part of the wreck, where they paid the night; and

a figural which they made being differential next day home. Hillberry filted, a bear want off, though at a great rife,

mained twenty-three hours on the wreek; and of fourtons,

in the end recovered. Of the times that perished one was the mafter of the velled a another was a passenger who had

been a mafter, but had loft or fold his thip in America :

played by Dr. Wright, and to fuccefeld in his practice in a finitar climate, inflead of immertion, which it appears was cusployed in the cafe mentioned by Dr. Girdleftone.

Nº. II.

I prefent these facts and observations to you, Gentlemen, with the confidence of a man whose intentions are good.

An Account of the remarkable Effects of a Shipwreck on the Mariners; with Experiments and Observations on the Influence of Immersion in fresh and salt Water, hot and cold, on the Powers of the living Body.

Read before the Royal Society, April 19, 1792, and extracted from Part I. of their Transactions for that year. — Page 199.

THE following narrative is submitted to the Royal Society, as containing in itself some curious circumstances, and as having suggested the experiments afterwards to be recited.

On the 13th of December, 1790, an American ship was cast away on a sand-bank that lies in the opening of the river Mersey into the Irish Channel. The crew got on a part of the wreck, where they passed the night; and a signal which they made being discovered next day from Hillberry Island, a boat went off, though at a great risk, and took up the survivors. The unfortunate men had remained twenty-three hours on the wreck; and of sourteen, the original number, eleven were still alive, all of whom in the end recovered. Of the three that perished one was the master of the vessel; another was a passenger who had been a master, but had lost or fold his ship in America;

the third was the cook. The bodies of these unfortunate persons were also brought off by the men from Hillberry Island, and were afterwards interred in St. Nicholas's church-yard, amidst a great crowd of spectators. The cook, who was a weakly man, died a few hours before the boat reached the wreck, but the two masters had been long dead, and this united the sympathy for their loss with a curiofity to inquire into its circumftances and causes. When the following particulars came to be known, this curiofity was increased. Both the masters were strong and healthy men, and one of them a native of Scotland, in the flower of life, early inured to cold and hardships, and very vigorous both in body and mind. On the other hand, feveral of the furvivors were by no means ftrong men, most of them had been inured to the warm climate of Carolina, and what was fingular enough, the person among the whole who seemed to have fuffered leaft was a negro.

What is extraordinary is feldom long unaccounted for in one way or other, and the death of the two masters was said to have been owing to their having taken possession of a keg which had contained cherry brandy, and which still contained the cherries;—these, it was reported, they had kept to themselves, and eaten in large quantities after the shipwreck; and this having produced intoxication, was supposed to have hastened their death. Some experienced seamen were satisfied with this account, which indeed seemed very rational; for though spirituous liquors may fortisty the body against the effects of heat combined with moisture, and may perhaps support it for a short time under great satigue, they are, I believe, uniformly hurtful when taken under severe and continued cold. Pleased to see a doctrine becoming popular which has been so ably supported by

Dr. Aikin, and others, I believed it might receive a striking confirmation from this catastrophe, into the particulars of which I determined to examine accurately. I therefore obtained access to the survivors of the crew, and from them, but more especially from Mr. Amyat, the mate, I received the information which I required.

22

In repeated conversations with this intelligent young man, I learnt that Capt. Scott, the master of the vessel, died in about four hours after the ship struck; and that Capt. Davison, the passenger, died in about seven : but that the flory of their having eaten cherries infused in brandy was entirely without foundation: of this he was certain, for he flaw the keg, which contained the cherries, staved, while Capt. Davison was endeavouring to fill it with water to make grog for the crew; the cherries fell on the wreck and were immediately washed into the sea. Mr. Amyat expressed his surprise at the early death of the two masters, but could not affign any cause for it. He said there was no liquor of any kind faved, nor any fort of food; that the whole crew were on an equality in all points, except that some were deeper in the water than others, but that the two mafters had the advantage in this respect, for they sat on the only part of the wreck that was out of the fea, whereas the poor negro, who escaped almost unhurt, was perhaps deepest in the sea of any. He explained this in the following manner. When the ship struck they cut away her masts to prevent her from oversetting, and after this she drifted over the fand bank, into what he called a "fwash" on the other fide. Here she floated, and they let go their best bower anchor; but it dragged, and the vessel struck again in a few minutes on another bank. In this fituation fhe espedar which has been to phly topported by

^{*} See Transactions of the Philosophical and Literary Society of Manchester, v. I.

the lay fome time, beating against the fand, and the sea breaking over her. In a little while Mr. Amyat faw the tar barrels, which formed her cargo, floating towards the land, and foon after the bottom parted entirely, and was carried in the same direction. Happily for the men, the part of the wreck on which they were lashed was held by the anchor, and floated in the water, a finall portion of the after part of the quarter deck being above the furface. On this fat the two mafters, generally out of the sea, but frequently overwhelmed by the furge, and at other times expoied to heavy fhowers of fleet and fnow, and to a high and piercing wind. The temperature of the air, as nearly as can be gueffed, was from 30° to 33° of Faht, and that of the sea, from trials in similar circumstances, from 38° to 40°. * Immediately before the two masters was Mr. Amyat himself. As he was fitting, and the deck sloped pretty rapidly, he was generally up to the middle in the water. The fituation of the rest may be supposed; some of them were up to the shoulders. They were not at any time able to change their position, but kept their legs in pretty constant motion to counteract the cold, their arms being employed in holding by the wreck.

The mafter of the ship, Captain Scott, a native of North-Carolina, and about forty years of age, died first. As they were in the dark, Mr. Amyat could not see his countenance; but he was first alarmed by hearing him talk incoherently, like one in the delirium of sever. By degrees his voice dwindled into a mutter, and his hearing seemed to fail. At length he raised himself up in a fort of convulsive motion, in which he continued a few seconds, and then

there feemed no doubt, that the death of the two mailers

fell

^{*} From some recent experiments I have reason to believe that the temperature of the sea was three or four degrees lower than it is here stated.

fell back dead on the deck. This happened about eight in the evening; four hours after the ship went aground. Soon after this, Captain Davison, who was about twentyeight, began to talk incoherently, in the same manner as the other. He struggled longer, but died in the same way, at about eleven at night. The cook died in the forenoon of the succeeding day. He was a low-spirited man, and desponded from the beginning. All the rest held out, as has been already mentioned, though forely pinched with cold and hunger, till they were taken up about three in the afternoon. Mr. Amyat faid that his hands and feet were fwelled and numb, though not absolutely senseles; he felt a tightness at the pit of his stomach, and his mouth and lips were parched; but what diffressed him most were cramps in the muscles of his sides and hips, which were drawn into knots. Though immerfed in the fea, they were all of them very thirsty; and though exposed to such severe cold, Mr. Amyat himself was not drowly, nor were any of the men drowfy, nor did fleep precede death in those that perished. These facts are curious.

Reflecting on the particulars of this melancholy story. there seemed no doubt that the death of the two masters was to be imputed to their peculiar position on the wreck. Exposed to heavy showers of sleet and snow, they might suffer from being wet with fresh, rather than falt water: they might also suffer from being exposed to the cold of the atmosphere, probably seven or eight degrees greater than that of the sea.* The chilling effects of evaporation might operate

^{*} This is I now believe a wrong supposition. Two or three degrees was more probably the difference, if indeed there was any difference. The water spreads over a vast expansion of sand, and as the cold had been some time great, it had probably brought both elements to the same temperature. (1798.)

operate against them, promoted as these must have been by the high wind; or they might receive injury from their frequent immersions in the sea, producing an alternation in the media surrounding. This last supposition did not, indeed, strike me at this time; the others dwelt on my mind.

Of the powers attending animation, that which feems fundamental, is the capacity of the living body of preferving the fame heat in various degrees of temperature of the fame medium, and, indeed, in media of very different denfity and pressure. If a definition of life were required, it is on this faculty that it might best be founded. It is known that some fluids, applied to the skin, vary in their effects according to their impregnation. In the fame degree of temperature, pure water on the furface of the body is much more hurtful than water in which falt is diffolved. Seafaring men are univerfally acquainted with this, and a striking proof of the truth, as well as of the importance of the obfervation, may be found in the Narrative of Lieut. Bligh. Probably the faline impregnation may stimulate the vessels of the skin in some way that counteracts the sedative or debilitating action of the cold. At any rate, it feemed not unlikely that some light might be thrown on this curious subject, by observing the effects of immersion in fresh and falt water, of equal temperature, on the animal heat. And this might also assist in accounting for the death of the unfortunate men already mentioned.

-loggid sammagnot onal selections of med vilegge som

I placed a large vessel, containing one hundred and seventy gallons of salt water, in the open air. The atmosphere

phere was damp, and what is called raw. The thermometer stood at 44° in the air, and this also was the temperature of the water. The subject of my experiment was Richard Edwards, a healthy man, twenty-eight years of age, with black hair, and a ruddy complexion. The hour chofen for his immersion was four in the afternoon, about two hours after his dinner; a time appointed rather for my own convenience, than as being most proper for the purpose. His heat was 98° before undreffing, his pulse 100 in the minute. He was undressed in a room where the mercury was at 56°; and afterwards stood naked before the fire till his heat and pulse were examined again, and found as before. He then walked pretty brifkly through a paffage paved with ftone into an open court, where the north-east wind blew fharply upon him: he was exposed to it for a minute, and then plunged fuddenly into the water up to the shoulders. The thermometer which had been kept in a jug of warm water, at the heat of 100°, was introduced into his mouth, with the bulb under his tongue, as foon as the convultive fobbings occasioned by the shock were over. The mercury fell rapidly, and a minute and a half after immersion it stood at 87°. He remained motionless in the water, and the mercury rose gradually; at the end of twelve minutes it stood at 93°1. While he fat in the water it occurred to me to examine his heat when he rose out of it into the air: I had reflected on the power that must be employed to keep up his heat in a medium fo denfe as water, and where an inanimate body, of the fame bulk, would have cooled fo much more speedily than in air of the same temperature. Supposing that this heat-producing process, whatever it may be, might continue its operations fome time after the extraordinary stimulus (the pressure of the water) was removed, I expected to fee the mercury rife by the accumulation of his

S CONTROLL

heat, on changing the medium of water for air, and therefore kept him exposed, naked, to the wind two minutes after taking him out of the bath. To my furprise, although the attendants were rubbing him dry with towels during this time, the mercury fell rapidly. He was put into a warm bed, and his heat when examined under the tongue was 87°, at the axilla, 89°. Frictions were used, and brandy mixed with water administered; but I found on this, as on all future occasions, the best mode of counteracting the cold, was to apply a bladder, with hot water, to the pit of the stomach (the scrobiculus cordis), a fact which I think important: this being done, his shiverings, which before were fevere, foon ceafed, and he became more comfortable. Three hours afterwards, however, he had not entirely recovered his former heat; but by eight at night he was in all respects as ufual. If even a oven distancement out of to the beat not

I have been very minute in detailing the circumstances under which this experiment was made; some of the particulars which, at the time, I thought of little consequence, I found afterwards of importance. The experiment itself I determined to repeat as exactly as possible.

to say thoughts, and described me to inquire again into

EXPERIMENT II.

On the next day, at the same hour, the same person was again immersed, as before. His pulse previously was 85, his heat 100°. He had been put to bed an hour before to save the time spent in undressing. The heat of the water and of the atmosphere 44°. The wind north-east, and strong. On this occasion, as before, there was a rapid fall of the mercury; the following table will save words:

2 min. after immersion 89° ½	7 Ther. 9 min. after immersion 95°3
3 - 90 1	10 94 ½
5 100 _ 1500 _ 10 94 ½	12 95
7 - 95 4	14 and 15 - 95
8 95 \$	The Sale of the sale on Sale

At the end of fifteen minutes he was taken out, and stood three minutes, naked, exposed to the north-east wind, at the end of which time the mercury had sunk to 88°. A draught of ale was given him, and he was put into a warm bed; in three minutes afterwards the mercury rose to 93°. An hour after his heat was 95°.

The effects produced by this alternate exposure to water and air of the same temperature, gave a new direction to my thoughts, and determined me to inquire again into this singular phenomenon. The most obvious method would have been to have prolonged the process of alternation, and replunged the person cooled by the external air into the bath; but this was running too great a risque, unless some more sudden and certain method could be found of restoring the heat that might be lost. It was prudent, therefore, to proceed more cautiously. In the next experiment I resolved to try the methods of heating, as well as of cooling the body.

EXPERIMENT III.

On the following day, at the same hour, the same person was again immersed in the salt-water bath. His heat previously was 98°, his pulse 100. The temperature of the air and the atmosphere, as before 44°. The mercury sunk rapidly to 90°.

a minutes

2 minutes after		10 minutes after - 94° 1
3 my per mor effethe and	88	11 - 94 4
4	88 1	The state of the s
5	90 1	13 96
6	92	14 96
7	92	96
8	94	16 96
9 - moins and oth ave	94	At the fame hour of ano

He was now taken out, and stood in the wind three minutes, shivering violently. This circumstance rendered it dissicult to ascertain exactly the fall of the mercury, which was, however, considerable. When examined in the room in which he undressed, it stood at 90°. He was now plunged into a fresh-water warm bath, heated to 97°½. What is very surprising, the mercury fell two degrees.—The following table will shew the progress of the return of his heat.

min. after immersion in the warm bath, mer-	5 minutes after, - 94°
cury, 88° 2 minutes, 92 3 92 4 94	7 — 96 8 — 96 9, 10, 11, 12, to 16, 96

If the rife of heat in the cold bath at 44°, and the warm bath at 97°½, be compared, the first will be found more slow; but that after being sixteen minutes in the one and in the other, the heat was the same in both cases, when taken at the mouth. It must, however, be acknowledged, that in the cold bath, the extremities were chilled and cold, while in the hot bath, the heat was equally diffused.—When Edwards got out of the hot bath, he put on his clothes, and was remarkably alert and cheerful the whole evening. Encouraged by the safety of these experiments, I resolved

resolved to increase the time of immersion in the cold bath, and to inquire more generally into its effects on the sensations, as well as heat.

EXPERIMENT IV.

At the same hour of another day, the same person was again immersed as before, his heat being previously 97° 1, and that of the water 42°. Wind north-east, and brisk.

It will be observed, that in the above table there are blanks left in the report. At fuch times the thermometer was taken out of Edwards's mouth, to admit of his answering the questions put to him. He faid, that on plunging into the water he felt an extreme cold, which he could not but think was partly owing to his being exposed, naked, to the wind before; that this cold diminished, and in a little while he felt comfortable, but that after a while the fense of coldness returned, though less than at first; diminishing again, but in a less degree. At length his sensations became pretty fixed. In this state, when the water was at rest, he should not even have known, by his feelings from the upper part of his cheft to the pubes, that he was in the water at all. His feet and legs were very cold; fo were his hands and arms; and so also the penis and scrotum. He mentioned, likewise, that he felt a cold circle round the upper

part of his body, though not constantly. On examining into this, I found it was greatest at first, and that it extended over the space which, from the undulations left in the bath by the plunge of immersion, was alternately above and under the furface of the water: when the bath fettled, it was little felt; but by agitating the fluid, I could reproduce it at any time when the cold in the extremities was not so great as to prevent its being felt. This curious particular ferves to explain a circumstance much dwelt on by Mr. Amyat, in giving an account of his fufferings on the wreck; that what he felt most severely was the cramps in the muscles of his hips and sides, parts which, from his fituation on the wreck, already described, must have been alternately under and above the furge. Here I must obferve, that the fea did not break over the fufferers all the time they were on the wreck. The wind moderated, as well as the waves, and for the last fifteen hours, they were not at any time overwhelmed, or at least Mr. Amyat himfelf was not. The cold never abated. Being all lashed to the wreck, they never changed their positions; the bodies of those who died occupied the space where they were originally placed. Mr. Amyat, therefore, during the whole time fat nearly up to the middle in water, but subject to the variations occasioned by the motion of the sea.

To return—When exposed naked to the wind, the mercury, in this case, sunk as usual five or six degrees, and his shiverings were great. Desirous of restoring his heat as speedily as possible, we incautiously heated the hot bath to 104°: but after being half a minute in it, he screamed out with pain, especially in his extremities, and about his scrotum. When taken out, his shiverings almost amounted to convulsion. The bath was lowered to 88°,

and he was replaced in it, and its temperature progressively, but pretty rapidly increased to 100°. He continued, however, to shiver much, his heat remaining about 90°; but a bladder, with very hot water, being introduced under the surface of the bath, and applied close to his stomach, the good effects were instantaneous, his shiverings ceased, and his heat mounted rapidly to 98°.

All these experiments having been made on one person,
I determined to repeat this last on another.

EXPERIMENT V.

R. Sutton, aged 19, of a pale complexion, and a feebles frame, was immerfed in the bath, under the circumstances of the preceding experiment. His heat was previously 96°½.

I a minute after,	heat 92	1 18 minutes 93°4
I minute -	- 90	19 931
2 minutes -	- 883	20, 21 94
3	- 89	22 921
4	- 90	23 924
5	- 92	24 924
6	- 924	25 94
7 to 10	- 92	26 94
11	e la no	27 92 2
12 to 15	- 92	28 924
16	- 922	29 94
17	- 93	30 94

Though this person seemed to bear the cold bath well, having lost in thirty minutes only $2\frac{1}{2}$ degrees of heat, yet when exposed afterwards to the wind, he shivered violently, and lost his heat very fast. He was put into a warm bath, heated to 96°, but recovered his heat very slowly, as the following table will shew.

1 minute

İ	minute after,	heat 88°	
2	minutes -	- 90	
3	AND DESCRIPTIONS	- 90₹	
4		- 90°	great shivering.
5	a ao thomsiqu	- 90	here the bath was heated to
			100°.
6	San town of the D	- 90	shiverings still.
7	The state of the state of	- 90	ditto.
8,	9	- 90½	ditto.
10	CONTROL STATE OF	- 92	ditto.
11	AND STATE	92	bath heated to 104°.
12	THE RESERVE TO SERVE	- 94	distribution to over hands
13	The state of the s	- 93	heated to 108°. Shi-
	Sall and the sall	to the case of	verings.
14	Table 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	93	a bladder with very hot water
			applied to the flomach.
15	Talana Talan	- 94	Charles and the same of after a
16	-	96	very comfortable.

EXPERIMENT VI.

Richard Edwards, the original subject of experiment, was again immersed in the cold bath, of the temperature of 40°, and remained in it three quarters of an hour. His heat previously was 97° his pulse 90 in the minute. The mercury sell at first to 92°, was stationary for a few minutes, and then mounted, though as usual, with no regularity. In twenty-two minutes it stood at 96°; it then began to decline, and in twenty-three minutes more had sunk to 94°. Being exposed as usual to the wind, the mercury sunk as usual, and he shivered violently. In the warm bath at 96° his shiverings continued several minutes, his heat remaining at 90 and 91°. In seven minutes the mercury began to rise saft, and sive minutes after was at 96°.

EXPERIMENT VII.

The effects of forty-five minutes immersion in the cold falt-water bath, at 40°, were proposed to be tried on Richard Sutton. He was much under the impressions of fear, and his heat previously raised the mercury only to 94°. The mercury funk, as usual, on his immersion, but to an unufual degree. It did not stop in its fall till it got to 83°, which perhaps might in part be accounted for by the extraordinary chattering of his teeth, admitting fome contact of the air. It then mounted in the usual irregular way, and at the end of thirteen minutes had got to 92°. Here it. stood for nineteen minutes longer with little variation; at the end of this time it began to fall rapidly, though irregularly, and in three minutes was down at 85°. He had now been thirty-five minutes in the water, and I did not think it fafe to detain him longer; we therefore hurried him into a warm bath, heated to 96°, where he shivered much. The bath was heated gradually to 109°, and in this heat he recovered his proper temperature in about twenty-eight minutes. Being then put into a warm bed, he fell into a profuse perspiration, which left him in his usual health.

One general remark will ferve for the pulse in all these experiments. It was not possible to keep the subjects of them from some degree of previous agitation, and this always quickened the pulse. The natural pulse of Edwards was about 70 in the minute; but it may be observed, that it was never under 85 before immersion, and generally above it. However this might be, it invariably sunk to 65, or from that to 68, in the water; became firm, regular, and small. After being long in the bath, it could hardly be felt at the wrist, but the heart pulsated with great steadiness and due

due force. In the last experiment, when the heat funk rapidly, Sutton faid that he felt a coldness and faintnes, at his stomach, which he had not perceived before, and when I examined the motion of his heart, it was feeble and languid. In some future trials of the effects of immersion in fresh water (one of which I shall detail) the same coldness at the stomach preceded a rapid fall of the mercury; and these facts, together with the effects I found from applying a confiderable heat to this part when the body was chilled with cold, convince me that there is some peculiar connection of the stomach, or of the diaphragm, or both, with the process of animal heat. Whoever will consider the rapidity with which a dead body would have cooled immerfed in water of the temperature of 40°, may form fome estimate of the force with which the process of animal heat must have acted in the experiments already recited. These experiments, however, do not coincide with our generally received theories of animal heat. The increase of heat, in fever, has led fome perfons to believe that animal heat is produced by, or immediately connected with, the action of the heart and arteries; here, however, it may be observed, that while heat must have been generated in the bath with more than four-fold its usual rapidity, the vibrations of the arterial fystem were unufually slow. Another, and a very beautiful theory of animal heat, supposes it immediately to depend on respiration; but in the bath, after the first irregular action of the diaphragm from the shock of immersion was over, the breathing became regular, and unufually flow. Laftly, the curious phænomenon of the heat rifing, and falling, and rifing again, in the bath, with the body at rest, and the temperature of the surrounding medium unchanged, is, I think, fatal to those theories of animation which confider the living body as a mere machine, acted

Cold of the benefit X 3 were as and of assertion on

on by external powers, but not itself originating action, and differing from other machines only in the peculiarity of the powers which are fitted to set it in motion.* I have said that the temperature of the medium continued unchanged, but it may be supposed that the bath was heated a little during the experiments; it was so; but being exposed, with a large surface, to the open air, the wind blowing briskly over it, its heat was little altered; in twelve minutes immersion it had gained nearly one degree, and in forty-five minutes, the longest duration of any of the experiments, it had gained three degrees. As this accession was regular, if it had been greater it would not have invalidated the foregoing observations.

Many other trials were made on the effects of immerfion in water on the human heat, which I shall speak of generally, under the general conclusions which they suggested.

The experiments already recited, suggested to me the notion, that in all changes from one medium to another of different density, though of the same temperature, there is a loss of animal heat. I found, however, that this conclusion requires many restrictions.

or extensionaming of school of the lamine to vector for My

^{*} The author does not mean to object to the theories of Dr. Crawford and Dr. Goodwin. The decomposition of the atmosphere in respiration, is in all probability the principal source by which caloric is received into the system, but the actual state of heat in the body cannot safely be inferred (as some have supposed) from the quantity of air respired. Caloric is received into the system by other processes also, particularly from the food and drink in the state of digession, and the evolution of it into active heat appears to be modified by the living power on its own peculiar principles, and cannot be explained by chemical or mechanical principles or analogies, so far as we are yet acquainted with the subject.

- 1. My experiments being made on bodies of such very different density as air and water, do not admit an universal inference of this fort.
- 2. Being all made in a temparature fifty degrees under the human heat, no certain conclusion can be drawn as to what might happen in degrees of heat much higher, where it is probable, the effects of the change, if it appeared at all, might be less striking. It would seem, however, that after a person is long chilled in cold water, the first effect of passing through the external air into the warm bath, is first a fall of heat in the air, and after this a still greater fall in the warm bath, followed, however, by a speedy rise.

The air and the water being equally cold, and both 45° or under, I found the lofs of heat in passing from the one to the other to be regulated in the following way.

- I. If, instead of being exposed naked to the wind previous to immersion in the water, the body was kept warm by a slannel covering, the mercury fell much less on the plunge.
- 2. If, after plunging in the water, the person continued in it only a minute or two, a subsequent fall of the mercury did not always take place, on his emerging into the air. On the contrary there was sometimes a rise on such occasions in the mercury, especially if the atmosphere was at rest.
- 3. In one instance, after continuing in the water fifteen minutes, on rising into the air in a perfect calm, though during a frost, there was little or no seeming diminution of

the heat; while exposure under similar circumstances, with a north-east wind blowing sharply, though the air was many degrees warmer, produced a rapid diminution. The effects of the wind in diminishing the human heat are indeed striking, and are not in my opinion explained by the common suppositions.

4. The loss of heat by a change of media, depends much on the rapidity of the change, for the plastic power of life in varying the process of animal heat, so as to accommodate it to the external changes, acts for a time with great celerity, though this celerity seems to diminish with the strength.

EXPERIMENT VIII.

I placed in a large room, where the mercury stood at, 36° two flipper baths at the distance of fix yards from each other. One was filled with cold falt-water of the temperature of 36°, the other with water heated to 96°, which was my own heat. Undreffing myself in an adjoining room by a fire, I afterwards flipped on a loose flannel dress, and defcended flowly into the cold bath, where I remained two minutes; I ascended flowly into the air, and then funk myself in the warm bath, where I remained two minutes also: I returned to the cold bath, where I staid two minutes as before, and removed from it again to the warm bath. But during all these changes of media and temperature, the thermometer with its bulb under my tongue never varied from 96°. I attribute this partly to the heat of my body being in some degree defended by the flannel dress, partly to the calm of the air, but chiefly to the flowness of motion in these changes. It may be said that the time of staying in the

the different baths was not long enough to produce any sensible change in the heat of circulating sluids of such a mass, but this is not consistent with many of the other facts.

5. The influence of the application of cold water to the furface of the body on the heat, is in some respects regulated by the animal vigour, as the following experiment will show.

EXPERIMENT. IX.

In the fame room I placed a large empty veffel, in which two young men fat down in succession, each with the bulb of a thermometer under his tongue. A man standing on a bench with a bucket of cold falt-water containing four gallons, poured the whole on the head and shoulders, suffering it to run down on the rest of the body. This process took up nearly a minute, during which I examined the mercury, and found it unchanged. They were both directed to continue fitting without motion for a minute after, during which, in both inftances, the mercury rofe two degrees. A third, much inferior in vigour, fubmitted to the same experiment, and the mercury continued during the affusion of the water unchanged, but in a minute after funk half a degree. In fevers where the heat is generally increased from two to fix degrees above the ftandard of health, pouring a bucket of cold water on the head always reduces the pulse in frequency, and commonly lowers the heat from two to four or five degrees. Of this falutary practice I hope foon to fpeak at large to the public.

6. The power of the body in preserving its heat under the impressions of cold, and the changes of temperature, and

and of media, feems in some measure regulated by the condition of the mind. That fear increases the influence of cold, and of many other noxious powers, will not be doubted; but the state of the mind to which I allude, is that of vigorous attention to other objects. This, it is well known, will to a certain degree deaden, or, indeed, prevent the fenfation of cold; and what does this, I apprehend, prevents, or at least weakens, its physical action. The astronomer, intent on the objects of his sublime science, it is faid, neither feels, nor is injured by, the damps or the chillness of the night; and in some species of madness, where the ideas of imagination are too vivid to admit the impressions of sense, cold is resisted to an extraordinary degree: I have feen a young woman, once of the greatest delicacy of frame, struck with madness, lie all night on a cold floor, with hardly the covering that decency requires, when the water was frozen on the table by her, and the milk that she was to feed on was a mass of ice.

7th. There are particular conditions of the atmosphere, not perfectly understood, that seem to have an instruence in depriving us more speedily of our animal heat, than others where the cold is greater.

It may feem that by this time I had renounced my intention of trying the effects of immersion in fresh water on the animal powers, and particularly on the heat. Some trials, I have, however, made, of which I shall only relate the following.

EXPERIMENT X.

In the same vessel containing an equal bulk of fresh water, Richard Edwards, the subject of my first experi-

ments, was immersed, at the same hour of the day. His heat previously was 98°, his pulse beat 92 in the minute: the heat in the air was 41°½, that of the water 40°. The wind was now in the west, so that in the court where the bath stood there was a persect calm. As I had some sears of the issue of this experiment, instead of exposing him for a minute naked to the wind before immersion, he was covered with a stannel dress from the air, till the instant he descended into the water, into which he was suffered to sink himself slowly, with the bulb of the thermometer under his tongue. These are important circumstances. The sollowing table exhibits the result.

Immediately on immersion,	14 minutes after - 96° 1/2
heat - 98°	15 96
1 minute after, 97½	16, 17, 18, 19, 20, 96
2 minutes 97	21, 22, 23 24, -
3 98	25 95
4 97½	26 94
5	27 931
6	28, 29 94
7, 8 96	30 93
9 97	31, 32 94
10 - 97	33, 34 921
11, 12, 13,	of I was about in imposits and

He now got out into the air very flowly, and flood in it three minutes, the wind not blowing on him. He lost one degree of heat at first, which he recovered. He was then put into a warm bath at 90°, which at first he felt warm, and his feet and hands were pained, but in two minutes he fell into a very violent shiver, and his heat fell two degrees. The bath was then heated to 95 and 96°, but still

still he felt cold. It was heated to 99°: he continued in it five minutes and his heat was 91°. The heat was gradually raised to 106°, when the sense of coldness of which he had complained at the pit of the stomach gradually went off. Before this I had usually kept him in the warm bath till his natural heat was nearly recovered, but after being half an hour in the heat of 106°, his own heat was still 93°. He now became sick and very languid, a cold sweat covering his face, his pulse very quick and seeble. He was removed into bed, but passed a feverish night, and next day had wandering pains over his body, with great debility, resembling the beginning stage of a fever. By cordials and rest this went off.

This experiment clearly enough confirms the greater danger of being wet with fresh than salt-water; but in itself points out nothing certain besides, except that it is not to be rashly repeated. I mean to try some of these experiments to a greater extent on the brute creation, when I have procured thermometers better suited to my views. The thermometers I employed had not a sufficient mobility for very nice experiments, and I am well aware, that in particular instances this may have misled me, though the general results, which is all that is of importance in such experiments as these, will I hope be found just and true.

Before I conclude, I must offer a few observations on the subject that led to these experiments.

that where there is only the choice of being wet with falt or fresh water, it is always safest to prefer the first. In the beavy showers of rain, hail, or snow, by which gales of wind

wind are generally accompained, the men that must be exposed to them, ought, like Lieutenant Bligh and his crew, to wring their clothes out of falt-water.

- 2. In all cases where men are reduced to such distress by shipwreck or otherwise, that they have it only in their power to chuse between keeping the limbs constantly immerged in the sea, or of exposing them to the air while it rains or snows, or of being exposed to it, where the sea is at times washing over them, it is safest to prefer a constant immersion; because, in the northern regions, where the cold becomes dangerous to life, the sea is almost always warmer than the air, as the experiments of Sir Charles Douglas shew; and because there is not only a danger from the increased cold produced by evaporation, but also from the loss of heat by the rapid changes of the surrounding medium, as the foregoing experiments point out.
- 3. Whether, in high and cold winds without rain or fnow, and where a fituation may be chosen beyond the reach of the waves, it is fafer to continue in the air, or to feek refuge in the sea, must depend upon several circumstances, and cannot perhaps be certainly determined. The motives for choosing the sea will be stronger in proportion as the wind is high and cold, and in proportion as the shore is bold.

The foregoing narrative shows that men may survive twenty-three hours immersion in the sea, of the temperature of 38° or 40° (as great a cold as it almost ever possesses) without food or water, and almost without hope of relief; but that any man ever survived an equally long exposure to the

^{*} These numbers ought I think to have been 33° or 35°. See note, p. 24-

the higher degrees of cold of the atmosphere, in the same circumstances, does not appear. Though in the case related, immersion in water did not prevent thirst, yet there is no doubt that it alleviated it, a circumstance of high importance towards the preservation of life.

Liverpool,
December 25, 1791.

P. S. I have purposely avoided any reasoning on the causes of the loss of vital heat on the change of media in the experiments recited. It may be supposed that during immersion, the water immediately in contact with the skin, having become heated to a certain degree, the naked body, on rifing from it into the air, was in fact exposed to a colder medium, and thus the lofs of heat, in this instance, produced. My examination of the heat of the water during immersion, not having been made in contact with the body, I will not deny that there is some foundation for the remark; and the cases, it must be allowed, are by no means exactly parallel between immersion in an open vessel, however large, and immersion in the sea, where the constant undulation may be prefumed to occasion a continual change in the furrounding fluid. But whatever allowance may be made for the circumstance mentioned, I am persuaded that the difference between the denfity of air and water being confidered, it is not fufficient to explain the loss of heat in the instance alluded to. The changes of temperature in the living body are governed by laws peculiar to itself. I have found, in certain diseases, greater and more sudden variations than any mentioned, from applications of cold very gentle in degree, and momentary in duration.

In his mafterly " Experiments and Observations on Animal

Animal Heat," Mr. Hunter has objected to taking the heat of the human body by introducing the bulb of the thermometer into the mouth, because it may be affected by the cold air in breathing. The objection is well founded if the bulb be placed on the upper furface of the tongue, but if it be under it and the lips shut, the effects of respiration may be disregarded, as I have found from many hundred experiments. The heat may be observed in this way with eafe and certainty, by employing thermometers curved at that end to which the bulb is affixed (the bulb being introduced at the corner of the mouth) fome of which have been made for me by Mr. Ramiden, accordto a form given, as well as others on Mr. Hunter's plan. From repeated trials it appears to me, that when the usual clothing is on, the heat of the living body may be taken with nearly the fame refult and equal certainty, under the tongue with the lips shut, at the axilla with the arm close to the fide, and in the hollow between the fcrotum and the thigh; every other part of the furface is liable to variation and uncertainty. It is evident, that of these three methods, the first only can be employed (as far as I can discover) when the trunk of the body is immerfed in water; and even when the naked body is exposed to the cold air, the first method feems the best, the heat remaining most steady under the tongue; the axilla is the next best in order, and the worst the lower part of the groin; for the scrotum and the parts of generation lofe their hear on the application of cold more speedily perhaps than any other part of the body, the extremities not excepted.

N. B. The water employed in the experiments related, contained falt in the proportion of one to twenty-four.

APPENDIX

No. III. I all bles and to

I AM happy to add to this fecond edition, extracts from two communications of Dr. Wright, with whose narrative this volume commences, in which he gives some farther information respecting the application of cold water to the body in fever. The first of these is from a letter of that gentlemen to Dr. Garthshore published in the 7th volume of the medical facts and observations 1797, on the treatment of acute diseases and particularly those of the West Indies. The letter is dated Edinburgh, December 10th 1794.

"In a former letter I remarked to you, that the application of cold water externally had been, for fome time,
practifed by Dr. Gregory, professor of physic in the university, in cases of typhus, with remarkably good essect;
but he has never carried it to the extent I did in my own
case, and in that of others, several years ago*. Instead
of dashing cold water on the naked body, Dr. Gregory
orders the bodies of his patients to be washed with a
fponge, dipped in cold water and vinegar, at least twice
a day. This operation I shall call lavatio frigida. The
earlier this mode is practised the better; because, in
typhus the patient grows daily worse, for in the second
week there is a great increase of sever, and a proportional

^{*} See London Medical Journal, Vol. vii. p. 109, (this reference of Dr. Wright is to the narrative given at the beginning of this Volume.)

" loss of strength: but even then Dr. Gregory has found " the application of the wet sponge act as a charm; nor have " delirium or petechiæ been confidered by him as any bar " to the adoption of this remedy; on the contrary, where " these have been present, and the pulse much quickened, " he has, by the lavatio frigida speedily reduced the pul-" fations from 110 to 90 in the minute, and the delirium " and other threatening fymptoms have foon after difap-" peared. About a fortnight ago, a student of physic, who " had been ill for some days before Dr. Gregory was ap-" plied to, had, besides a great degree of sever and deliritum, " numerous fpots, or petechiæ, on his breaft, belly, and " extremities. The lavatio frigida was used on the day the Doctor first visited him, and by next morning the " delirium had ceafed, and the petechiæ difappeared. The " pulse, which on the preceding day had been 110, was " now at 80; and by continuing the application of the " wet sponge now and then, the pulse became natural on "the fourth day after the Doctor first faw him. Many " fimilar cases might be adduced from the books of the cli-. " nical ward of the Royal Infirmary."

"Successful as this method has been in the hands of "Dr. Gregory, and some others, besides mine, I am well "aware that much judgment and caution are necessary in putting it in practice. In all cases where there are visce- ral obstructions, cold bathing does much mischief; and in fevers of this sort, with inflammatory diathesis, there is reason to suspect topical inflammation of the viscera; in this last case, if cold bathing were made use of, the patient would run the risque of his life, and the physician justly lose his character."

Medical Facts and Observations, vol. vii. p. 2.
Y

The second extract is from a report drawn up by Dr. Wright (since his return to the West Indies) as physician to the army, and director of the military hospitals in Barbadoes, respecting the diseases most common among the troops in that climate, and the best mode of treating them. This valuable paper is inserted in the second volume of the Annals of Medicine, for 1797.

" TYPHUS.

"Under this head we class the nervous fever, the ship fever, the hospital sever, the jail sever, the pestilential fever, the yellow sever, &c. All these are different names for the same disease, and differ only from each other in malignity or violence, from local circumstances, the state of the atmosphere, or season of the year.

" TYPHUS NAVIUM, SHIP FEVER.

"In transporting troops from England to the West " Indies, this fever often breaks out, and rages with great " violence. The patient is fometimes indisposed a day of " two before the difeafe be marked; but frequently is taken "ill at once, with laffitude, proftration of strength, irre-" gular chills and heats, nausea, and sometimes vomiting. " He has a flight head-ach, is reftless in bed, has confused " ideas, and troublefome dreams; there is no great heat of " the fkin, it is rather moift, cold, and fweating. The " tongue at first is white, moist, and slimy; afterwards " dry and parched. The countenance is pale and funk, 44 the eyes dull and languid, the belly irregular, the urine pale and fecreted in too great a quantity." The doctor goes on to enumerate the remaining symptoms, and to point out the causes, which differ in no respect from the symptoms

toms and causes of Typhus in general. Of the cure he speaks as follows:

"The fick on being landed were washed, either with cold sea-water, or water made milk warm; they were completely shifted, and placed in clean well-aired wards, with dry bedding. If the fever was recent, an emetic was given, and then the primæ viæ cleansed by a solution of natron vitriolatum, and manna, or cream of tartar. After this the bark was given without loss of time, and with such success, that in less than a week the men were in general discharged, quite recovered.

"In the beginning of the ship sever, the cold bath had the best effects; and through the day, when the sick were hot, washing the hands and face suddenly in cold water and vinegar was exceedingly refreshing. Light covering in bed was directed, especially where there was any præternatural heat. In the advanced stages where there were symptoms of inflammatory diathesis, we had recourse to small doses of antimonial powder alone, or mixed with a few grains of calomel. Where the body was costive, sive grains of calomel proved to be the best laxative, or purge."

Dr. Wright goes on to mention the use of cold acidulated watery drinks, fresh air, blisters in obstinate cases, and in the latter stages, bark, wine, and nourishment. He observes that what is said of the ship-sever, will equally apply to the hospital or jail-sever.

[&]quot; TYPHUS ICTEROIDES. YELLOW FEVER.

[&]quot;The yellow-fever," fays Dr. Wright, "appeared Y 2

degree of malignity." * * * " The diforder began with alternate rigors and hot fits, giddiness and dimness of fight; the patient could not bear the light, the eyes looked dull, and half-closed when asseep; the white of the eye was tinged with yellow, the eyes themselves funk, and the countenance fallen. The pulse seemed varied; sometimes it was natural, but in general it was small and tremulous. The breathing was difficult, attended with sighing, anxiety, and restlessness.

"The skin in general was hot, and had that biting feel so common in all malignant severs; at other times the skin was cold and clammy. There was uniformly a great prostration of strength, a loss of appetite, and a constant inclination to vomit; at first the contents of the stomach were thrown up, afterwards an abundance of bile. The tongue was dry and furred, the thirst unquenchable, the body costive, the urine scanty, high-coloured, and burning or scalding the urethra. Universal pains, especially in the joints, calves of the legs, and tip of the shoulders, often took place. The sleep was disturbed, and the patient inclined to delirium. The fever was without any sensible remission at first, but afterwards there were evident remissions and exacerbations.

"Inter in different cases; and instances have occurred where it finished its course in twenty-four, or forty-eight hours. Every symptom rapidly increased, the senses were more disturbed, delirium and coma were constant; at times the delirium was low, but sometimes surious. The vomiting of bile was incessant; at last, whatever

"was brought up had the colour of coffee; and this was denominated the Black Vomit. Hæmorrhages from the nose, the mouth, and even from the pores of the skin, were frequent and fatal appearances. The body now became yellow or livid, with cold or clammy sweats; the countenance hypocratic, the pulse sunk, and death put an end to the patient's misery.

"Cure. The first intention was directed to a speedy evacuation of morbid matter; the second intention, to prevent the secretion and accumulation of more; the third to relieve the most urgent symptoms; and the fourth, to obviate the mischief already done to the system.

" the black yomir, had taken place, the captionen has " First, early and brisk purging was put in practice. " After some trials of cathartics, we had recourse to large " doses of calomel, repeated at short intervals, until a plen-" tiful discharge by stool was obtained. By this means, " the vomiting instead of being increased, was gradually " abated, and at last subdued. By calomel, the pores of " the /kin were opened, a resolution of the fever was brought " about, and the patient happily recovered. Where pa-" tients were received in the advanced stages of yellow fever, " we had ftill recourse to calomel, and at the same time " when it was needful, to mercurial frictions, and the " warm bath; and we recollect of no instance where mer-" cury had been freely given, and perfevered in till it shew-" ed itself in the mouth, which was not attended with the " happieft confequences.

In the beginning of the yellow fever, the cold bath fuceeeded admirably, as in other species of typhus, but in
Y 3

" the advanced stage much caution was necessary. | Some "Lucky expedients, however, have been practifed, which "fuccess alone could justify: thus, when the most urgent " fymptoms had been fubdued, the patients were wrapt up in a wet blanket, a profuse sweat was brought on, and an " immediate recovery the consequence. In cases of exces-" five vomiting, effervescing saline draughts have at times " been fuccessful; but calomel, prudently administered, will have the defired effect. In all fevers where the sto-" mach was irritable, and bile was pumped up, our de-" pendence was on calomel; where it failed, danger was " apprehended. We did not, however, despair; we gave " capficum pills (Cayenne pepper made into pills) with " the most marked success; and even where melaena, or " the black vomit, had taken place, the capficum has " fnatched the patient from the most imminent danger.

"In all the cases of the yellow sever which we have feen, we never found those enormous quantities of calomel mel necessary, which are mentioned by many late writters. Ten-grain doses, indeed, were given, for two or three turns, until it operated by stool If, after this, the fever was obstinate, the dose of calomel was reduced to two grains every three hours, till symptoms of resolution had appeared. It was then discontinued.

"Respecting antimonials, the stomach was in two irritable a state to bear them in any form, and the lancet
(blood-letting) was not only unnecessary, but dangerous
in the extreme."

The report of Dr. Wright extends to the Remitting and Intermitting Fever, Cholera Morbus, Diarrhœa and Dysentery;

Dysentery; it is signed by all the gentlemen of the medical staff of Barbadoes; and is in every point of view deserving of great attention. It is indeed to be wished that the information conveyed to us by this experienced and able practitioner, respecting the severs of the West Indies, had been more precise. We have still to lament our ignorance in regard to the actual heat of the patient, and the successive changes that it undergoes in the progress of the disease; and we are equally uninformed as to his sensations of heat. It affords me however particular satisfaction, that the experience of Dr. Wright (whom I have not the pleasure of knowing) so far as it goes, corresponds so much with the result of my own observations.

It was in the beginning both of the typhus and of the yellow fever that he found the cold bath produced the happiest effects; and doubtless these effects, though not particularly defcribed, corresponded with his former experience, and appeared in the diminished frequency of the pulse, the diminished heat of the body, and the flow of fensible perspiration. It is worthy of remark that the calomel, to the efficacy of which in this difease Dr. Wright has given a decided testimony. though its first action was that of a cathartic, yet operated in the end as a sudorific. " By calomel, the pores of the " fkin were opened, and a refolution of the fever brought " about." The history indeed of all fevers, of every country, fo far as I have been able to trace them, uniformly points out the falutary confequence of early and profuse fweating, of which even the epidemic, named from this fymptom, which prevailed in England in the end of the 15th and beginning of the 16th century, under the name of the fweating fickness (fudor anglicus) is not an exception.-It is only however when it is early, general, and profuse, that Y 4 fweating fweating is so strikingly falutary; when scanty, it is insignificant, and when it occurs in the latter stages of the disease, it is often injurious. Early and copious sweating seems no less salutary when excited by applications which act immediately and in some measure mechanically on the pores of the skin, than when excited in any other way. Mr. Thomas Graham, an ingenious and experienced furgeon in the Royal Navy, informs me, that when ferving on board Sir John Jervis's fleet in the Mediterranean, in the present war, he was often enabled to produce profuse perspiration with all its happy consequences, by a method which he employed of covering the whole surface of the body, in the early stages of fever, with warm vapour. In the directions for the cure and prevention of the plague, published by Count Berchtold, at Vienna in 1797, of which an account is given in the fecond volume of the Annals of Medicine, p. 327, it is faid, " Immediately after a person is perceived to be infected with " the plague, he must be taken into a close room, and over a brazier of hot coals, with a clean sponge dipped in " olive oil, his body must be very briskly rubbed all over, " for the purpose of producing a profuse sweat." * * " The friction ought not to be continued more than four " minutes, and a pint of oil is enough to be used at each " time. In general the first rubbing is followed by a very " copious perspiration; but should it fail of this effect, " the operation may be repeated, first wiping the body with " a warm dry cloth; and in order still farther to promote " perspiration, the patient may take any warm sudorific " drink" * * * The operation ought to be begun " on the first appearance of the symptoms of the disease; " if neglected till the nerves and the mass of blood are af-" fected, or a diarrhœa has commenced, little hopes can be " entertained of a cure."

This

This practice has, it is faid, been followed with great success at the hospital of St. Anthony in Smyrna, and the benefit seems to depend on the early profuse sweating that is produced. It is evident that this mode of treatment, as well as that mentioned by Mr. Graham, may be adopted in the cold stage of the paroxysm, which it probably brings to a speedier issue. It corresponds with the warm bath employed in this stage of the disease by Galen, and being applicable to sever at an earlier period than the affusion of cold water or the cold bath, it does not superfede the use of these remedies if it should afterwards be found requisite to employ them. It deserves a trial in sever in general, especially in those severs of the warm climates which run their course with such rapidity.

Before I quit the subject of fever, I must shortly obferve, that some of my friends have expressed their surprise at the vast number of cases which appear on the records of our Difpenfary, and their wish that this point were enquired into more narrowly. I have discussed this subject with the late Mr. Avison, who admitted that some deduction ought to be made for cases classed under this title, which might not turn out regular fevers, amounting perhaps to a fifth or a fourth part of the whole. This point is particularly attended to at prefent, and some account of the refult will be given in a future edition of this volume, should another edition be called for. In the mean time, it may be observed, that though the numbers are likely to be fomewhat reduced, the refult of the most accurate investigation ferves to establish more clearly the great prevalence of typhus among our poor, and to enforce the establishment of a receptacle for fever, (fee p. 241) and the adoption of some regulation respecting inhabited cellars, and other

other confined habitations, where the contagion is conflantly kept up.

THERE is no part of the preceding volume which has met with more opposition than the opinion I have delivered respecting the non-absorption of the surface of the body, and the experiments and observations in support of this opinion in chap. xvii. I retain, however, this opinion, though I do not consider the question as decided; but if the contrary doctrine should be established in the end, it is evident that it must be fixed on a new foundation. Leaving the matter to future inquiry, I give, in the mean time, an extract from the report of Fourcroy on the memoir of M. Séguin, quoted p. 234, as the volume which contains it is scarce in England, hoping that we may soon be furnished with a more full account of the experiments of this distinguished philosopher.

M. Fourcrov begins with mentioning the circumstances which led M. Séguin to doubt the commonly received opinion, respecting cutaneous absorption, and to make new experiments on the subject. The memoir read by him contains the result. "Nous n'entrerons point" says M. Fourcroy, dans le détail des nombreuses expériences qui y sont consignées, et que l'on affoibliroit par la rapidité nécessaire dans un rapport; nous nous contenterons de rappeller à l'Académie les principaux résultats qu'elles ont donnés. Le premier résultat tiré de trente-trois expériences saites fur lui-même, c'est que le corps n'augmente pas de poids dans le bain, qu'il perd moins dans l'eau que dans l'air, et que cette perte suit sur-tout la raison de la tempéra-

" ture de l'eau du bain; que la perte de poids dans l'eau à dix ou douze degrés (baromètre à vingt-huit pouces), est " à celle dans l'air comme 6 5 est à 17; qu'à quinze à dix-" huit degrés de température, cette perte dans l'eau est à " celle dans l'air comme 7,5 est à 21,7; que dans l'eau " chaude à vingt fix ou vingt-huit degrés, elle est à celle " dans l'air comme 13 est à 23. Deux autres personnes ont offert des réfultats différens dans ces pertes relatives, mais elles ont toujours moins perdu dans l'eau que dans l'air. " M. Séguin attribue cette perte moindre à ce que la ma-" tière de la transpiration insensible n'est point exposée au " contact de l'air qui doit la dissoudre dans l'état ordinaire. " Il rend raison de la différence de ces pertes à diverses températures de l'air de la manière suivante. La perte de " poids qu'on éprouve dans l'eau à dix ou douze degrés est 66 beaucoup plus foible que celle qui a lieu dans l'air, parce " qu'il n'y a point de transpiration cutanée; il n'existe " alors que la transpiration pulmonaire : celle qui se fait " dans l'eau à dix-huit degrés est un peu plus soible que la " feule transpiration pulmonaire, parce qu'outre qu'il n'y " a point de transpiration cutanée dans ce cas comme dans le " précedent, l'air qui entre dans le poumon est chargé d'hu-" midité, et ne dissout pas toute celle qui se dégage de ce " viscère ; enfin, la perte de poids qu'on fait dans un bain " d'eau à vingt-huit degrés est plus considérable que celle " qui est produite par la seule transpiration pulmonaire dans " l'air à cette même température, parce qu'alors le corps " perd, et par cette dernière transpiration, et par la sueur " qui fort des vaisseaux exhalés, en raison de l'augmentati-" on de mouvement du cœur et des artères, qui, comme M. Séguin l'a prouvé dans son mémoire sur la transpira-" tion, est la seule cause de la transpiration sensible, ou de 4 la sueur. Mais, malgré la différence de ces trois résultats " qui

" qui dépendent de la température de l'eau du bain, il n'est " pas moins certain qu'il n'y a point augmentation du poids " du corps par le bain, et qu'il y a seulement une perte " moins forte que dans l'air, dépendante de l'absence de " celui-ci et de la privation de sa qualité dissolvante par " rapport à la matière de la transpiration.

"Les premières expériences dont nous venons d'expofer " les réfultats généraux les plus importans, prouvoient " bien que le corps n'augmente pas de poids dans le bain, " mais elles ne décidoient point encore l'absorption ou la " non absorption par la peau, car on pouvoit objecter à leur " auteur que la perte moindre que dans l'air dépendoit de la " portion d'eau absorbée par les vaisseaux inhalans. M. " Séguin a pensé que, pour répondre à cette objection, il " falloit faire baigner des individus dans des dissolutions de " fubstances dont les effets sur l'économie animale fussent " bien tranchans. Il a employé la dissolution de muriate " oxigéné de mercure, à des doses connues, en pédiluves, " sur plusieurs malades attaqués de symptômes vénériens, et " il a constamment observé que lorsque la peau étoit bien " faine et l'épiderme bien entier, il ne paffoit pas de fublimé " corrosif dans leurs humeurs, ils n'éprouvoient aucun des " accidens dus à ce sel, et aucune amélioration dans leurs " maladies, tandis que dans le cas où l'épiderme étoit affec-" té et entamé, comme dans la gale, &c. ce sel pénétroit " le corps et produisoit alors les effets qui en font recon-" noître l'existence dans l'économie animale.

"Non content de ces expériences faites dans un hôpital fur plusieurs individus malades, M. Séguin a cru devoir les recommencer sur un sujet sain, les suivre avec une forupuleuse exactitude, jusqu'à ce qu'il eût levé tous les doutes

" doutes, et il s'est choisi lui-même pour sujet de ces nou-« velles tentatives. En tenant à un grand nombre de reprises différentes, pendant long-temps à chaque fois, une es partie de son bras plongée dans une dissolution connue de " sublimé corrosif à différentes températures, et disposée " dans un manchon de verre recouvert de taffetas gommé, " de manière à ce qu'il n'y eût point d'évaporation fenfible, " le reste de son corps, excepté sa bouche, étant d'ailleurs se enfermé dans l'enveloppe imperméable de taffetas ciré, " afin de pouvoir apprécier la transpiration pulmonaire " comme il avoit apprécié la perte dans l'air de la partie du bras plongée dans la dissolution, M. Séguin est parvenu " à obtenir des réfultats aussi singuliers que nouveaux : " nous ne rapporterons ici que ceux qui ont trait à l'absorp-" tion par la peau. Quand la dissolution de deux gros de " fublimé dans dix livres d'eau est à dix et à vingt-huit de-" grés de température, la quantité de ce sel dans le bain est très sensiblement la même après l'expérience qu'aupara-" vant, et conséquemment il n'y en a pas d'absorbé; mais " dans la même diffolution à dix-huit degrés, si la presque total té du bras tres-sain, et dont l'épiderme est bien en-" tier, y reste plongée, il y a par heure 1, 2 grains de sub-" limé absorbé, quoique l'eau du bain qui tenoit cette por-" tion de sublimé en dissolution ne soit point elle-même 46 absorbée. M. Séguin tire de ce singulier résultat l'in-" duction que ce n'est pas par les vaisseaux lymphatiques " que se fait cette absorption du sublimé, car ces vaisseaux " absorberoient bien plus facilement l'eau, qui cependant. " n'éprouve pas de diminution, mais par les vaisseaux ex-46 halans. L'auteur explique ce phénomène d'une manière " très ingénieuse : il pense que ces vaisseaux exhalans ref-" ferrés par des températures basses telles que douze, en " évatuant des gouttelettes de fueur continuelles par un " effort

" effort plus considérable du cœur et des artères produit par " la température de vingt-huit degrés, et n'absorbant con-" féquemment aucune parcelle de fublimé dans les deux " circonstances, se trouvent tellement disposés à la tempé-" rature de dix-huit degrés, que suffisamment dilatés pour " que l'eau de la dissolution soit en simple contact à leur " extrémité avec l'humeur transpiratoire, sans être repouf-" sée par l'écoulement de cette humeur, comme cela a lieu " à vingt-huit degrés; alors ce contact fans mouvement " permet à l'humeur de la transpiration de dissoudre de " proche en proche une partie du muriate oxigéné de mer-" cure dissout dans le bain, de le partager avec sa première " eau de dissolution, de se mettre avec elle dans un vérita-" ble état d'équilibre, comme cela a lieu dans toutes les diffolutions falines mêlées avec de l'eau pure. Ainfi, " fuivant lui, une diffolution faline à dix ou à vingt-huit " degrés étant mise en contact avec la peau humaine bien " faine, il n'y a point d'absorption ni de la part de l'eau ni " de la part du fel ; la même diffolution à dix-huit degrés, " mise en contact avec la peau qui ne transpire point dans " l'eau à cette température, permet à l'humeur de la tranf-" piration de partager le fel de la dissolution jusqu'à l'équi-" libre de faturation et de le porter dans la circulation : cet " effet n'a pas lieu lorsque les pores des vaisseaux exhalans " sont resserrés par une température basse ou traversés par " les courans de sueur qui repoussent la dissolution de sel. " Les vaisseaux lymphatiques n'enlèvent ni solide, ni li-. " quide, ni fluide élastique à la surface du corps ; les vaif-" seaux exhalans n'absorbent jamais ni solide insoluble, ni " liquide. ni gaz, parce qu'ils sont toujours pleins de la li-" queur transpirable. L'absorption qui n'a lieu que dans " une certaine température, et qui tient à la tendance à nu req salloumitation mount sidemental rance and "ril'équilibre

" l'équilibre entre un liquide non faturé et un liquide sa-" turo, est par cela même très-bérnée.

M. Séguin, en poursuivant la description de ses ex-" périences, confirme les affertions précédentes, par l'hif-" toire de plusieurs malades vénériens, chez lesquels la dif-" folution de sublimé employée en lotion n'a rien fait tant " que l'épiderme n'a point été entamé; d'autres à qui le " fublimé ainsi que le muriate d'ammoniaque et de mercure " ou le fel alembroth, appliqués à sec, n'ont produit d'ef-" fets que lorsque la peau s'entamoit par l'âcreté de ces sels ; " de quelques-uns qui n'ont éprouvé ni érofion, ni action " consequemment du muriate de mercure ou mercure doux " appliqué fur la peau. Le tartrite d'antimoine et de po-" taffe ou le tartre stibié, appliqué ainsi sur le ventre, a " purgé après avoir produit des boutons dans le lieu de l'application; la gomme gutte, la scammonée, appliquées fue le ventre à sec, et recouvertes comme les matières " précédentes d'un emplâtre agglutinatif à fa circonférence, " n'ont fait naître aucun effet sensible. L'onguent mercuriel n'agit que par une friction qui fait pénétrer l'ox-" ide de mercure sous l'épiderme, et le met dans le cas " d'être absorbé; un onguent fait avec le sublimé corrosif " entame la peau, et devient un des moyens les plus prompts et les plus actifs de faire pénétrer du mercure très-oxidé " dans le torrent de la circulation.

"Il n'est pas nécessaire d'entrer ici dans de plus grands détail sur les expériences très-nombreuses de M. Séguin, pour faire concevoir les résultats généraux qu'elles lui ont sournis et qu'on peut réduire aux suivans; 1°. Les vaisseaux absorbans n'absorbent dans aucun cas ni l'eau, ni l'air, ni les matières qui y sont mêlées ou dissoutes; "2°. l'épi-

" 2°. l'épiderme qui les recouvre exactement dans l'état " fain, les empêche absolument de faire cette fonction, et " ils ne l'exercent qu'au-dessous de cette croûte; 3º. Les " matières dissolubles sont peu-à-peu enlevées à l'eau qui " les dissout par l'humeur de la transpiration placée à l'ex-" trémité des vaisseaux exhalans, lorsque cette humeur ne " coule point en torrent comme dans la fueur, ou lorfque " les vaisseaux ne sont point resserrés comme par une tem-" pérature trop basse ; 4°, les matières liquides ou fluides " élastiques ne sont point admises dans les vaisseaux exha-" lans, toujours pleins de l'humeur transpiratoire qui v " féjourne ou qui y est dans un mouvement inverse à celui " de l'absorption ; 5°. les matières caustiques séches ne sont " absorbées qu'après avoir détruit et corrodé l'épiderme; " 6° les matières fèches non folubles ne peuvent paffer " dans le système lymphatique, que lorsque par une fric-" tion plus ou moins forte on les a fait pénétrer à travers " les mailles et les pores de l'épiderme jusqu'à l'espace ou " s'ouvrent les bouches des vaisseaux absorbans.

"A ces énoncés, qui réfultent immédiatement des ex"periences indiquées, M. Séguin en ajoute d'également
"importans, qui ne sont que des conséquences nécessaires
des premieres et dont nous exposerons ici les principaux:
"1°. les maladies épidémiques se contractent par la voie
de la respiration, et les miasmes dissous dans l'air déposé
dans les poumons sont absorbés par les vaisseaux absorbans
de ces viscères qui, dépourvus d'épiderme, jouissent d'une
force absorbante très-entière; 2°. le diabète ne provient
point de l'eau absorbée dans l'air par la peau, mais du restructe qui ne peut pas être enlevée aux poumons
par l'air trop chargé d'humidité; 3°. les amas d'eau ou
les diverses espèces d'hydropisses ne dépendent que de la
"différence

différence d'action entre les vaisseaux absorbans et les vais-" feaux exhalans ; 4°. l'abforption commune dans les abfor-" bans par le vuide qui v est produit, soit par la diminution " de pression, soit par celle des stimulus, leur structure " valvulaire interne, détermine le mouvement des fluides " de leur extrémité vers le système des vaisseaux sanguins; " 5°. les matiéres acres et stimulantes, en faisant contracter " les vaisseaux, arrêtent l'absorption; l'affinité des sub-" flances à absorber avec les vaisseaux absorbans détermine " également cette fonction; telle est la différence d'action " des matiéres nourrissantes et des purgatives, par rapport " au système des vaisseaux absorbans abdominaux; 6°. " enfin les virus contraires pénètrent par les poumons; " l'épiderme est un rempart qu'ils ne peuvent franchir dans " l'état sain et dans l'intégrité parfaite de ce tissu, et ils ne " peuvent pas être absorbés par la peau."

> La Médicine eclairée par les Sciences Physiques, vol. iii. p. 234.

I think it unnecessary to add a single observation to this extract, which will convince every candid mind that our conclusions respecting cutaneous absorption have been hasty, if not wholly erroneous.



of difference of allians sentre has validations ablantante extlet visits of haves extralamental. Tables, even communication has the reliable part of grodule, for part is diminusion of both part period extra even grodule, for part is diminusion of other prefixed, for part evine, des fermine, lear drochlute of the part extre which the following of mousement des finishes of destrock farguines of the reliable with the reliable to the reliable to the reliable to the reliable of the reliable to the reliable to the reliable to the reliable to the reliable to

La M. Joine celas vie de Science

I think it more refers to all a fingle obforcation to give, some objects which which will control of the control that and that control of the free beauty.

Set of

INDEX.

produced delication to continue to be the second

Alexander the Great latherings of his army from the im-

A

CO CONTRACTOR OF THE PERSON OF		P.
Ablution of the body with cold water, its use in feve	er,	71
employed by	De	34
Hahn, in the fever of Breslaw, 1737,	76	83
Affusion of cold water on the surface of the body use	d by	
Dr. Wright,	1	5
employed in a fever in the L	iver-	
pool Infirmary,	6	8
employed in a fever in the 30th regiment,	9	13
manner of using it in fever, with cases,	17	32
used with different impregnations, .	33	35
precautions required in using it, illustrated	l by	
cafes,		37
case of fever in which it was not salutary,	46	48
ufed in finall pox,	52	53
fcarlatina anginofa, .	63	65
temperature of the water employed,		66
- used in the burning fever on the coast of	the	
Red Sea, and in Abyssinia,	son J	188
Affusion of tepid water on the surface of the body	, its	
cooling effects,		69
how and when to be employed, .	69	71
- not so effectual in fever as the affusion of	cold	
water, .		70
this explained,		79
Aikin, Mr. Charles, referred to on the use of cold d		moul
during exercise,	1	107
Z2 A	lexand	der

INDEX.

Alexander the Great, fufferings of his army from th	e im	. P.
prudent drinking of cold water, .	99	100
his own illness from bathing i		
Cydnus, explained,	113	117
Alexander, Dr. his account of the increased heat		
intoxication,		216
Alkohol, observations on its effects in health and	dic.	
eafe,	215	217
Alpinus quoted,		
В	Alla	30
Bacon, Lord Verulam, quotation from him in favo	ur of	tota:
the use of unguents,		206
Bath, cold, fome account of its use in small-pox,	il one	62
- ufed by Galen in fever,	to sto	102
when fafe, in health and difeafe,		108
cases and histories illustrating this,	110	122
use of, in convultive diseases,	ant a	130
oase of infanity in which it was successful	113	140
precautions necessary in employing it in	infa-	140
nity or convulfions,	****	149
proposed to be employed in the fever of	f the	
West Indies	1	184
Bath, warm, used in infanity,		142
- use of, proposed to be revived in the	West	77-
Indies,		205
its restorative effects after fatigue,	101	ibid.
general observations on it,		207
Baths, ancient, observations on,		125
Ruffian, their temperature,		ibid.
Bell, Dr. his paper in the Manchester Transactions	re-	100
ferred to,		199
Blagden, Sir Charles, his experiments in a he	eated	. 22
room referred to	123	198
Boerhaave, opposed the use of cold drink in fever,		87
Brandreth, Dr. his employment of cold water extern	nally	The state of
in fever,	2 200	62
mountain.	Brow	

INDEX.	iii
and the second of the second o	P.
Brown, Dr. John, his doctrines on the use of cold as a re	1
medy controverted, 72	74
Bruce, Mr. his account of the fevers in Nubia, &c.	188
services, account alcold of along the published by	
TOI SET UNSINGEN	
Cardanus, wrote in favour of cold drink in fever,	86
Celfus, quoted and referred to, . 83	86
Chardin, Sir John, cured of fever by plunging into the fea	, 184
Cold, its stimulant powers afferted and explained, 72	75
- external, dangerous after profuse sweating,	203
- its operation in inflammatory diseases, 218	220
Convulsive diseases, cases of, with various remedies, 130	138
Cullen, Dr. his definition of typhus,	16
his arrangement of fever incomplete,	44
an opinion of his controverted,	65
- his discovery of the cold produced by evaporation	1, 87
his account of the effects of change of tempera-	12/25
ture on fenfation,	126
Currie, Dr. Wm. his use of the digitalis in infanity,	142
Cruikshanks, Mr. his calculation of the quantity of pers-	None of the last
pirable matter, &c.	196
case of inanition mentioned by him,	258
Cyrillus, Dr. his account of the dieta aquea referred to,	87
D	
Danis Da samakan kisanawa afaka asawi	
Darwin, Dr. remarks on his account of the operation,	
of cold,	74
his theory of mania and convultions,	
quoted, reflections on his fystem,	
De Hahn, Dr. his account of the fever of Breslaw,	The state of the s
Digitalia was din infanitu	2134
Digitalis used in infanity,	142
Dispensary, Liverpool, table of diseases admitted there,	-
table of fevers,	230

THE DESCRIPTION OF THE PERSON		D
Drummond, Dr. A. M. quoted,		P.
nat E to necessarily		
Emetics, their operation in fever explained,		210
Evaporation, account of cold produced by, publishe	d by	
Dr. Cullen, 1755,	10	197
fupposed to regulate the animal heat,		198
parent and referred to,		100
All probabilitions gargainty of Fact to know and land,		
Ferriar, Dr. his account of the house of reception	for	100
fever at Manchester,		235
Fever, histories of .	I	17
cases of, detailed,	21	32
- remarks on the nofological arrangements of,		44
- case of, not included in these,		46
fcarlet, cured by affusion of cold of water,		63
hectic, remarks on,		71
at Breflaw, in 1737, account of,	76	83
- theories of, ancient and modern,	0 428	151
- table of, admitted at the Liverpool Difpensary,	1912	230
- house of reception for, at Manchester,		235
Ford, Dr. cafe by him,		138
Fordyce, Dr. George, his experiments in a heated r	oom	
referred to,	123	198
Franklin, Dr. his air-bath recommended in fever,		81
his account of the circumstances that rea	nder	
the cold bath dangerous,		116
a letter of his respecting the effects of evap	ora-	
tion on the furface of the body,		198
quated, rededingson bit fyftem it co		
a, Dr. his account O the fever of Breflaw,		
Galen, gave cold drink in fever,		86
- ufed the cold bath alfo,		102
his account of the temperature of the Roman b	A work	3 1.575
his doctrine of fever,	139	153
Z Drumport	Ger	ard.

INDEX.

4	P.
Gerard, Dr. his use of the affusion of cold water in fever	
in fcarlet fever,	63
Graham, Mr. Thomas, royal navy, his account of the	A
use of the vapour-bath in fever, append	
Gregory, Dr. James, his use of ablution with cold water	The Chi
in fever, appendi	× 46
rmary, Liverpool, account of a lever that broke out	
H	
Haller, has collected the facts and arguments for the in-	
halation of the skin,	244
Haygarth, Dr. his plan for the prevention of small-pox	
referred to, .	60
experiments on variolous matter fuggested	
by him, . 60	62
Heat, animal, state of in fever, cases which shew this, 21	32
method of examining, 35 36, appendix	45
history of the changes of in the pa-	
roxysm of fever,	172
its origin,	191
how regulated,	192
Heironymus Mercurialis, his account of the unguents	
used by the ancients before swimming,	112
referred to respecting the an-	
cient baths,	128
Hoffman, quoted on the danger of pouring cold water	
over the body when fweating,	92
Howard, Mr. his account of the treatment of fever in	
Italy, &c. referred to,	186
communication to him by the author,	187
Hunter, John, his physiology referred to,	
his peculiar language defended,	165
Hunter, William, his opinion on the formation of the	
perspirable matter opposed,	200
Hunter, Dr. John, his account of the temperature of the	
fprings in Jamaica,	118
Supply Su	100

red for wearly of the calculation of calculation for the	1743
Jackson, Dr. his account of the fevers of Jamaica re-	
ferred to, amount with aver answord T ald amed	182
Immersion in water salt and fresh, cold and warm, ex-	
periments on, appendix	, 25.
Inanition, case of,	247
Infirmary, Liverpool, account of a fever that broke out	
there, . II	6
Inhalation by the skin, generally believed by the ancients,	195
afferted by Haller, and the mo-	1
derns in general,	244
- experiments which render it doubtful,	246
denied by Séguin,	265
extract of an account of his expe-	
riments respecting it, appendi	x 56
Inunction of the furface of the body mentioned by Homer,	102 - 1
use of by the Romans,	112
general among the ancients, and	
still common in the east,	204
recommended by Bacon,	206
Liverpool, population of, with remarks,	221
prevalence of fever among the poor,	223
charitable establishments, account of,	224
comparative fickliness or healthiness of the	
different months among the poor,	227
	PARTY.
M M and mode wind add to	
cord, ble his sociant of the treatment of fiver in	
Maclean, Dr. his account of the diseases of St. Domingo	
referred to, alt al . Il as malas in man	182
Mania, remarkable case of, cured,	140
Maniacs, their reliftence to cold and contagion,	149
Marcard, Dr. objects to the cold bath in convultions,	150
affumes the inhalation of the skin,	271
Mersey, river, at Liverpool, not salter at high than low	
water,	34
Men	fley,

INDEX.	vii,
The topperation Tilly to world alderingers from	P.
Mersey, river, experiments on its temperature,	119
Morandi, Dr. quoted,	186
C CONTANT METERS IN	
and the standards of the southwester	
Opium, used in a case of infanity,	141
operation of in fever and in health, remarks on,	211
P	185
Percival, Dr. his support of the house of recovery at	mol T
Manchester,	225
Perspiration, insensible, its effects in cooling the body,	235
more plentiful in warm than cold weather,	
The state of the s	195
quantity of, fenfible, remarks on,	
The state of the s	199
matter of, in the European, not fitted to the	204
Torrid Zone,	187
Plague, cured by immersion in the sea, - case of, cured by exposure naked to the dews of	
	188
night,	
cold drink and the cold bath recommended in, inunctions of oil employed in, append	iv EA
inunctions of oil employed in, append	24
R	9790
Ramazzini, case of inanition related by him,	257
Respiration, modern theory of,	192
Rush, Dr. his account of the diseases arising from the	1000
incautious use of cold drink,	92
Rutherford, Dr. his theory of the peculiar action of the	20 E 1990 YE
lungs in diabetes, "	270
lungs in diabetes,	
S	
Savary, his account of the treatment of fever in Egypt,	188
Séguin, M. denies the inhalation of the skin,	
extract of an account of his experiments or	2000
this subject, append	
Shipw	ALL DO

at the title of with an annual	P.
Shipwreck, remarkable history of, with an account	
	ndix 20
Small-pox, cases of, treated by the affusion of cold	
and warm water,	52
observations and experiments respecting,	59
Sydenham, Dr. his doctrine of disease,	153
T.	
	pendix 1
Thermometers, account of,	35
how to be applied in taking the hum	an
heat, the aft and come after appe	ndix 45
Thirst, nature of,	177
- observations on the relief obtained in, by the :	if-
fusion of water, or immersion in the bath, 17	8 267
Tobacco, its use in convulsive diseases,	135
d.Zone, and	
cured by framerican is We feet, 187	Spirita
Water, cold, used as a drink in fevers by the ancients,	, 76
authorities for this practice,	86
against it, and bed death blos	88
rules for using it in fever,	89
in health,	95
cases illustrating the circumstances und	
which cold drink is dangerous,	96
case of convulsions cured by it,	139
how to be used in the sever of the West I	
dies, and in the plague,	185
how used by the Egyptians in their burning	NAME OF STREET
fever,	188
in Nubia and Abyffinia,	189
Wright, Dr. his narrative,	1' 5
his account of the use of ablution with co	
water in fever at Edinburgh, appen	
extract from his report on the treatment	70 = 0
acute diseases in the West Indies, appen	
append	100 40

