

A clinical introduction to the practice of auscultation, and other modes of physical diagnosis : intended to simplify the study of the diseases of the lungs and heart / By H.M. Hughes.

Contributors

Hughes, Henry Marshall, 1805-1858.

Publication/Creation

London : Printed for Longman, Brown, Green, and Longmans, 1845.

Persistent URL

<https://wellcomecollection.org/works/sbmaybyk>

License and attribution

This work has been identified as being free of known restrictions under copyright law, including all related and neighbouring rights and is being made available under the Creative Commons, Public Domain Mark.

You can copy, modify, distribute and perform the work, even for commercial purposes, without asking permission.



Wellcome Collection
183 Euston Road
London NW1 2BE UK
T +44 (0)20 7611 8722
E library@wellcomecollection.org
<https://wellcomecollection.org>



E. XII. 2.


William Roth

29,630/B

P. e 39.

18247

H 20/3/07



ON

AUSCULTATION.

Wilson and Ogilvy,
57, Skinner Street, Snowhill, London.





A
CLINICAL INTRODUCTION
TO THE
PRACTICE OF AUSCULTATION,
AND OTHER MODES OF
Physical Diagnosis :
INTENDED TO SIMPLIFY THE STUDY OF THE DISEASES
OF THE
LUNGS AND HEART.


BY
H. M. HUGHES, M.D.
FELLOW OF THE ROYAL COLLEGE OF PHYSICIANS,
ASSISTANT PHYSICIAN TO GUY'S HOSPITAL, ETC.

LONDON :
PRINTED FOR
LONGMAN, BROWN, GREEN, AND LONGMANS,
PATERNOSTER ROW.

1845.



TO THE
PUPILS OF GUY'S HOSPITAL,
FOR WHOSE USE IT IS MORE ESPECIALLY WRITTEN,
THIS LITTLE WORK,
PRODUCED FROM MATERIALS PRINCIPALLY
THERE COLLECTED,
IS AFFECTIONATELY INSCRIBED,
BY THEIR SINCERE FRIEND,
THE AUTHOR.



P R E F A C E.

THE object of the present work,—for while so many books on Auscultation already exist, it may be necessary to explain the object of a fresh candidate for public favour,—will be perhaps best explained by a sketch of conversations, in which I have on several occasions taken part, and from which the notion of writing it may be stated to have in great measure originated.

“ I want, Sir, to learn something about Auscultation,” a pupil has said to me ; “ what book would you advise me to read ? ” “ The book of nature, Sir ; go to the bed-side, and to the inspection room.” “ Yes ! but I know nothing about it, and I want some little information beforehand.” “ Certainly ! Read Andral, Bouillaud, Laennec with Forbes’ notes, Hope, Stokes, Williams.” “ Oh ! but I have not time to read half of them, or

perhaps really to study one of them. I only just want something to give me a little insight into the matter, and then I mean to work away in the wards." "Then look over Martinet's Manual, or get Sharpe's little compilation, or Spittall, or Cowan's Bedside Manual, or Dr. Newbigging's translation of the exquisite little work by Barthe and Roger." "Well! I have got one of those you mention, but I can make nothing out of it. There is such a profusion of terms, such lots of 'bruits and râles,' that I cannot get on at all. I wish you would write a book that beginners could make use of; it is really much wanted for pupils, and for other persons who have been out of the way of clinical instruction for some years."

In conversations to this effect I have on several occasions taken part. Whatever, therefore, may be thought of the judgment of the individuals in applying to myself to relieve them of their difficulties, and howsoever such an application may be regarded, as a mere incense-offering to my self-complacency, the difficulties complained of I have presumed to be real. I know indeed that, after the perusal of works especially treating of diseases of the chest, pupils, and practitioners too, are often quite incapable of instituting an ordinary

examination, and of distinguishing the more common physical signs. With terms they may be sufficiently familiar; the theory of morbid sounds may be not unknown to them; but how they are to elicit some of these sounds, and how they are to distinguish others, they know not. They cannot, in fact, *practise* auscultation and percussion; and they are unacquainted with the mode of setting about it. How, then, is this knowledge to be attained? Clearly by clinical instruction, most efficiently; by the precepts delivered and the illustrations offered at the bedside. But it unfortunately happens, that comparatively few students attend to these instructions, whether of precept or example, and that numbers find out their deficiencies only when it is too late for them to devote their attention to this infinitely most desirable mode of supplying them.

It has, then, appeared to myself as well as to others, that a work was wanted, in which should be simply explained to the student of auscultation, not merely the origin, character, and diagnostic value of certain physical signs, but also the manner in which he should proceed to elicit them; and which should direct him how to

percuss, as well as state the indications afforded by percussion.

It may, indeed, be asserted that all this, and much more than this, has been already done by several preceding writers. It may be so. Still the tyro in auscultation is at a loss. Preceding writers do not appear to have afforded the required information in an available form. The present attempt may be equally unsuccessful. But though I have certainly perused nearly all, and I believe, with one or two exceptions, all the works which have been recently published on diseases of the chest in the English language, and many that have not been translated, I know of no writing which approximately answers the purpose I desire to effect; unless it be the articles "Thorax, exploration of," and "Auscultation," by Dr. Forbes, in the *Cyclopædia of Practical Medicine*. Had these articles been published in a separate form, and been somewhat extended, it is possible that the present attempt would not have been made, and this work would have never appeared; though from my recollection of those articles, perused a long time ago, I trust some useful additions may be found herein.

I cannot forbear here to recommend, and to

express the delight I derived from the perusal of, the admirable little book of Barthe and Roger. It contains a great amount of information, conveyed in a very agreeable way; but it treats of auscultation alone, and is in some other important particulars unsuited for the object I have in hand.

That object is *not* to attempt to teach the practice of auscultation and percussion;—that I feel assured can be attained only at the bed-side of the patient—but it is to point out to the student the way in which he may learn it by himself. It is *not* to treat of the diagnosis of thoracic diseases; upon that I profess not here to enter; but it is to point out the physical signs of those diseases, and, as far as I am able, simply and intelligibly to explain the causes of those signs; it is to instruct the beginner in the mode, by which he is to obtain a knowledge of them, as well as to direct him how to interpret them.

Consistently with this object, I have desired that my statements should be plain, my directions distinct, and my explanations simple, and devoid of anything approaching to erudite mystery. Consistently with this object, also, I have thought it desirable to avoid all quotations, as well as any

Unable to display this page

rived from an experience of more than fifteen years, especially devoted to the practical study of diseases of the chest; during the whole of which I have neglected scarcely a single opportunity of examining the signs of disease in the living, or the products of disease in the dead subject; and during nearly the whole of which I have possessed the advantages afforded by one of the largest hospitals in the British dominions.

With the exception of the article by Dr. Forbes, referred to in the body of the work, I have not made a single extract from any author; to that article I have referred simply for the purpose of examining, and partially copying, the author's regional divisions of the chest. Though, therefore, the statements herein contained *may* have been repeatedly made, and the opinions herein delivered *may* have been again and again expressed by others; and though I may have seen them recorded by one or a dozen writers; yet, unless it be so acknowledged in reference to any particular fact or opinion, I believe I may with confidence assert, that I have given no directions, of which I have not by long practice proved the utility, and that I have made no statements, which, from whomsoever they

may have been originally derived, are not "*the result of my own observation.*" Facts and signs are matters of experience and of observation ; of these I have not hesitated to express myself with decision. Explanations of facts and signs are often merely matters of opinion ; of them I trust I have ever spoken with caution, and with becoming deference to the opinions of others. At the same time, when I have entertained settled and decided opinions upon any debatable question, with whatever difficulty or doubt that question may be surrounded, I have thought it right to give expression to those opinions ; while upon some matters, the explanation of which appears to me to be still involved in obscurity, I have not hesitated to acknowledge my ignorance.

I have thought it preferable to make use of the terms ordinarily employed, as, though open to some objection as stated by some previous authors, they have an acknowledged meaning among auscultators, and are so generally received, that I believe the changes proposed by those writers would not be attended with any advantage, commensurate with the confusion which would arise from their introduction. The language

of auscultation is already difficult enough to the uninitiated ; it appears therefore unadvised, particularly in a work intended to simplify, to increase the difficulty by a " confusion of tongues."

As to the style in which the work is written, I may observe that my chief desire has been to write plainly, so as to be intelligible by all. I have studied utility rather than elegance, and, if capable of attaining the "*apte*" and "*distincte*," I have not been very solicitous about the "*ornate*" of the Roman orator. I have written for those especially who are engaged, or who are likely to become engaged, in the practical study of thoracic disease at the bed-side ; and I have written for them as I would address them, in a familiar and almost colloquial manner. At the expense of frequent repetitions, of homely phrases, and, it may be, of some literary coarseness, I have attempted strongly to impress upon the mind of the reader an important fact, rather than to polish an expression or to round a period. Of this just complaints may perhaps be made ; but it must be recollected that what may appear grating to the ear of the literary critic, may be smooth and easy to the less refined, and that—to borrow the simile

of a philosophical writer of the last century—
“the luxurious receive no greater pleasure from
their dainties than the peasant does from his
bread and cheese; but the peasant, whenever he
goes abroad, finds a feast, whereas the epicure
must be well entertained to escape disgust.”

H. M. H.

14, ST. THOMAS'S STREET,

Oct. 1845.

TABLE OF CONTENTS.

	Page
Dedication	v
Preface	vii
Table of Contents	xvii
Explanation of Plate	xxiii

CHAPTER I.

PRELIMINARY OBSERVATIONS AND DIRECTIONS.	1
1. Cautions as to the conduct and demeanor of Examiner	3
2. Necessary qualifications of the Student of Auscultation	5
3. Position of the Patient	10
4. Divisions of the chest into regions	13
5. Different modes of exploration in diseases of the chest	16

CHAPTER II.

INSPECTION, OR OCULAR EXAMINATION.	17
1. Method to be pursued, and circumstances to be noticed on inspection	17
2. Indications afforded by inspection	21

CHAPTER III.

PALPATION, OR MANUAL EXAMINATION.	28
1. Mode of employing palpation	28
2. Indications afforded by palpation	29

CHAPTER IV.

	Page
PERCUSSION.	34
1. Different modes of percussion	36
1. Immediate percussion	36
2. Mediate percussion	37
2. Percussion in the normal state of the chest . . .	45
3. Natural resonance of different regions on percussion	46
4. Indications afforded by percussion	51
1. In diseases of the lungs and bronchial tubes .	51
Bronchitis	51
Pneumonia	52
Phthisis	54
Venous congestion	58
Malignant disease, &c.	59
Emphysema of the lung	60
2. In diseases of the pleura	61
Pleuritic, or other effusion	61
Malignant disease, &c.	63
Pneumothorax	63
3. In diseases of the heart	65
Disease of the heart	65
Pericardial effusion	67
4. In disease of the anterior and posterior Medi-	
astinum	69
Aneurism	69
Abscess, &c.	70

CHAPTER V.

AUSCULTATION, OR EXAMINATION BY THE EAR.	72
1. Immediate auscultation	73
Cautions respecting it	75
2. Mediate auscultation	78
Stethoscopes	78
3. Mode of using them	84

	Page
4. Cautions respecting mediate auscultation	86
5. Combined use of immediate and mediate auscultation	90
6. Auscultation of the organs of respiration in health	91
1. Of the natural respiration	92
2. Of the voice	102
Pectoriloquy	103
Bronchophony, &c.	103
7. Auscultation of the organs of respiration in disease	106
1. Of the respiration in disease	107
2. Dry sounds in disease of the air passages and lungs	108
<i>a.</i> Larynx and trachea	108
<i>b.</i> Bronchial tubes	111
Sonorous rattle	112
Sibilating rattle	113
Emphysema	114
<i>c.</i> Phthisis	118
Pneumonic consolidation	118
Tubular breathing	120
Amphoric and fistulous breathing	121
Metallic respiration and tinkling	122
Malignant disease	124
3. Dry sounds in diseases of the pleura	125
Pleuritic rubbing	126
Crackling	128
Jerking respiration	129
Tubular breathing	130
Absence of respiration	131
Exceptive cases	132
Amphoric respiration	135
Metallic breathing	135
Metallic tinkling	137
4. Moist sounds in diseases of the air passages and lungs	137
Comparison of moist rattles with each other	139

	Page
Tracheal rattle	141
Mucous rattle	141
Gurgling	142
Muco-crepitating rattle	144
Crepitating rattle	145
Sounds resembling it existing in health	147
“ <i>Râle crépitant réduit</i> ”	150
Moist crackling of disorganized lung	151
Phthisis	152
Muco-crepitation	153
5. Moist sounds in diseases of the pleura	154
Gurgling, &c.	155
Metallic tinckling	156
8. Auscultation of the voice and cough	158
1. In diseases of the air-passages and lungs	158
Larynx	159
Dilatation of bronchial tubes	160
Emphysema	161
Pneumonia	161
Venous congestion, &c.	162
Malignant disease	165
Phthisis	166
Pectoriloquy	167
2. In diseases of the pleura	170
Ægophony	170
Ægophonic bronchophony	171
Exceptive cases	173
Absence of vocal resonance	174
Metallic resonance	174
6. Auscultation of the heart	175
1. Auscultation of the heart in health	176
The Impulse	177
The Rhythm	180
The Sounds	184
The Second sound	185
The First sound	186

CONTENTS.

xxi

	Page
The natural range of the sounds of the heart .	188
2. Auscultation of the heart in disease . . .	191
1. The Impulse	191
2. The Rhythm	197
3. The Sounds	200
<i>a.</i> Increase of the sounds	200
<i>b.</i> Decrease of the sounds	202
<i>c.</i> Abnormal sounds or murmurs	204
Direct	205
Regurgitative	213
Illustrative examples	215
Anæmic murmurs	221
<i>d.</i> Pericardial sounds	228
<i>e.</i> Mixed sounds	232

CHAPTER VI.

MENSURATION, OR EXAMINATION BY ADMEASUREMENT. 236

Mode of measuring the chest 240

CHAPTER VII.

SUCCUSSION, OR EXAMINATION BY SHAKING. 243

Mode of effecting succussion 244

100	The history of the...
101	...
102	...
103	...
104	...
105	...
106	...
107	...
108	...
109	...
110	...
111	...
112	...
113	...
114	...
115	...
116	...
117	...
118	...
119	...
120	...

CHAPTER II

121	...
122	...
123	...
124	...
125	...
126	...
127	...
128	...
129	...
130	...

CHAPTER III

131	...
132	...
133	...
134	...
135	...
136	...
137	...
138	...
139	...
140	...

EXPLANATION OF PLATE.

THORACIC REGIONS.

1. The Acromial regions.
2. The Superior Sternal region.
3. The Infra-Clavicular regions.
4. The Inferior Sternal region.
5. The Mammary regions.
6. The Axillary region (left.)
7. The Lateral region (left.)
8. The Scapular region (left.)
9. The Infra-Scapular region.

ABDOMINAL REGIONS.

10. The Epigastric region.
11. The Hypochondriac region (left.)
12. The Dorsal region (left.)
13. The Umbilical region.
14. The Iliac region (left.)
15. The Lumbar region (left.)

EXPLANATION OF PLATE

TO GREAT BRITAIN

1. The London region
2. The English Channel
3. The Irish Channel
4. The Irish Sea
5. The Atlantic Ocean
6. The English Channel
7. The Irish Channel
8. The Irish Sea
9. The Atlantic Ocean

ADDITIONAL NOTES

1. The English Channel
2. The Irish Channel
3. The Irish Sea
4. The Atlantic Ocean
5. The English Channel
6. The Irish Channel
7. The Irish Sea
8. The Atlantic Ocean

A

CLINICAL INTRODUCTION

TO THE PRACTICE OF

AUSCULTATION,

&c.

CHAPTER I.

PRELIMINARY OBSERVATIONS AND DIRECTIONS.

IT cannot be necessary at the present day to insist upon the advantages of auscultation and percussion to the practical physician. The remarkably rapid improvement which has recently taken place in the diagnosis of diseases of the chest, which, not more than thirty years ago, were the most obscure, but which may be now considered as among the most easily recognized affections to which the human frame is liable,

B

is almost entirely attributable to their employment. Their importance may be fairly supposed now to be generally, if not universally, conceded. It is, however, as I conceive, a duty incumbent upon all who venture to write, however briefly, upon these modes of investigating disease, to endeavour to impress upon the mind of the student, what has been frequently and in various ways forced upon my own mind; viz. that it is, to say the least, highly desirable, if not absolutely essential to his future success in combating disease, that he should become practically familiar with auscultation and percussion while at the Hospitals. They require much time—much practice—much training; in other words, a prolonged education is necessary to their advantageous employment. When, indeed, facility has been previously attained, and confidence already acquired by repeated trials, and by frequent reference to the dead body, in confirmation of opinions formed during life, they may be used with benefit to the patient and satisfaction to the attendant, in private practice. But I believe that skill and aptitude in their use are rarely acquired in private, and that, from the comparatively limited opportunities of studying morbid anatomy at the homes of patients, dexterity and confidence are not often attained except in public institutions. I would therefore earnestly recommend every medical pupil at the Hospitals to become acquainted with the

practice of auscultation, and familiar with the products of disease *now*, while he is able to do so; as I feel assured that he will never regret the time expended in the attainment of these objects. If, on the contrary, the present opportunity be allowed to pass unimproved, it is not impossible that he may never enjoy another, and it is highly probable that he never will have so wide a field, or so suitable a time, for the study of that which he will find of vast service to him in his future course.

1. *Cautions as to the conduct and demeanor of the examiner.*

Some care is occasionally necessary lest persons of an excitable and irritable temperament, whose chests are about to be explored, should be alarmed by the anticipation of the examination, the nature of which they do not understand, and the trouble and fatigue of which they consequently exaggerate. It is therefore in such cases desirable, that, if it have been previously made known that an examination of the chest is to be made, the process should be represented to be, as it really is, free from pain and attended with little inconvenience, though, if thoroughly effected, necessarily occupying considerable time. With the same view of not exciting alarm in the mind of such nervous individuals, all unnecessary display of stethoscope and

pleximeter, and parade of every kind, should be avoided; otherwise some undefined notion of an operation may be produced, and the patient may become unwilling, or unable, to submit to the ordeal. If everything be quietly done, and all treated as a matter of course, and managed with delicacy and gentleness, I believe very few persons, unless exhausted by disease, will be either unwilling without hesitation to submit to, or unable without injury to bear, a tolerably minute exploration of the chest.

The examiner should also endeavour at the commencement to wear a cheerful aspect, and be careful not to allow the results of his investigation to escape him in the hearing of the patient, or even to permit them in any way to affect the expression of his countenance, each shade of which is sometimes watched by patients with the most intense anxiety, and even slight changes in which occasionally have a very depressing effect upon susceptible individuals.

In all examinations the benefit of the person examined should be the primary consideration.

—To this the mere gratification of the curiosity or interest of the examiner should be ever made to yield. The determination of a nice point of diagnosis should never be allowed to interfere with the real good of the invalid. When, therefore, patients are likely to become seriously exhausted by

being examined, the exploration should not be attempted, or, if already commenced, should be at least temporarily suspended: unless, indeed, the immediate determination of a particular question should itself, and at that time, necessarily involve the benefit of the person examined. This caution is more particularly necessary in hospital and other public practice, in which it sometimes becomes necessary, for the sake of the patient, and the interests of humanity itself, to restrain the ardour of pupils and other examiners.

2. *Necessary qualifications of the student of Auscultation.*

1. A certain amount of knowledge of anatomy and physiology is necessary to the practical auscultator. It is absolutely essential that he should be acquainted not merely with the situation and boundaries of the heart, lungs, liver, spleen, &c., the course of the large vessels, the attachments of the diaphragm, &c., but that he should be also well informed in the physiology of the circulation and respiration.

2. Though a thorough familiarity with the history and constitutional symptoms of disease, its morbid anatomy and pathology, is not in the first instance necessary for the successful prosecution of the study of the physical signs of thoracic diseases, as both may be assuredly studied

conjointly with advantage at the bed-side of the patient by even the tyro in medicine; yet some general knowledge of the ordinary symptoms and results of morbid processes is requisite for the student of auscultation at even the very commencement of his course. If he possess not such information, if his conclusions are derived alone from what he hears, he will find himself frequently mistaken. He may in consequence be induced to regard physical diagnosis as of little practical value, and to desist from the pursuit of that which may become to him a most important acquisition in his future discrimination of disease.

3. It is also clear that the student should have some general acquaintance, derived from reading, lectures, or clinical instruction, with the connection existing between certain sounds and certain forms of disease. He should know, for example, that dulness on percussion, a crepitating rattle and increased resonance of the voice, usually accompany particular stages of pneumonia, and that a rubbing noise is attendant upon some forms of pleuritis and pericarditis. That he may escape the danger of forming too hasty an opinion, and avoid the error, so common with young and inexperienced auscultators, of incorrectly designating a particular sign, and consequently of misinterpreting its import, it is highly desirable that he should in the first place have pointed out to him, either by his clinical instructor, or by some more

advanced pupil, a good characteristic example of each sign as it occurs in practice. Of this he should endeavour to obtain a distinct impression. If he obtain such an impression, he will not be likely afterwards to forget it, and to it therefore he will be able in his future investigations to refer; so that he may himself afterwards say to others, as he has already been told—"Here is a good example of pleuritic rubbing," and "there a fine specimen of crepitating rattle."

4. It is necessary that the student should have some knowledge of the more common doctrines of acoustics; such as the origin and nature of sound, the variation in the conducting power of different media, and the facility or difficulty with which the vibrations of sounding bodies are transmitted to others of different density and elasticity. He should recollect, what appears to have been forgotten by some writers upon auscultation, that, as regards this art, in diseases of the chest, the vibrations of sound, however originating, however conducted, and however modified before reaching the solid parietes of the chest, are transmitted either mediately by the stethoscope, or immediately to the ear by the vibrations of these solid parietes themselves; and therefore that the sound, whether originating in air, or transmitted through water, when it has reached these parietes, is in all cases better conducted by a solid than by any other medium; and that it is so, not

merely because solids are in themselves better conductors than either fluids or gases, but because the medium of conduction is identical with that by which the sound necessarily reaches the surface of the chest.

5. It is scarcely necessary to state, that the auscultator's sense of hearing should not be defective. But to become a proficient in the art of auscultation, something more is necessary than the mere absence of defect. For this purpose what musicians call "a good ear," or a delicate appreciation of minute differences of sound, is an important if not an essential qualification.

6. There are few attainments really valuable which can be mastered without the expenditure of much time and trouble. Auscultation and percussion are certainly not exceptions to this general rule. Patience and perseverance, then, are certainly requisite for the successful prosecution of auscultation. A long education, multiplied opportunities of examining disease, and comparing the sounds heard during life with the revelations of the inspection room, are assuredly necessary. A caution may, perhaps, be here introduced with advantage. The student should ever follow his patient to the *post-mortem* room, and should carefully mark the localities in which certain signs have been previously observed, and compare those with the condition of the subjacent parts; but he should also bear in mind the period

of time which has elapsed between his examination and the death of the patient, and recollect the changes that may have occurred in the state of those parts in the interim. From a neglect of these considerations in the mind of the necroscopists, I have not unfrequently seen students abashed and disconcerted, and the diagnosis of the physician improperly and unjustly repudiated.

7. It is essential to the successful prosecution of auscultation and percussion that their value should be fully, but at the same time fairly, estimated. If the individual employing them be influenced merely by fashion, and labour under the misconception that they are in reality of little importance to the formation of a correct diagnosis, it is highly probable that, in his hands at least, they will prove so. If, on the other hand, the student conceive that by their use he will be enabled in every case to determine the exact condition of a diseased organ, he will be most assuredly liable to much disappointment. They should never be, and there is no valid reason why they ever should be, employed to the exclusion of the other modes of investigation. When the history, and the general or constitutional symptoms of disease, have been inquired into with the same minuteness and interest as in other cases, then, but usually not till then, should percussion and auscultation be employed, or then, at any rate, and not till then, should any deductions be made

from the indications afforded by the physical signs. To a neglect, or comparative disregard of the history and ordinary symptoms of disease, and a too implicit confidence in their important art, is it probable that some of the errors of even accomplished auscultators are to be attributed. There is, however, I repeat, no valid reason why the auscultator should not use with equal diligence, and be equally skilled in the employment of, all other modes of investigation, as the individuals who discard the stethoscope, and make no use of the ear in forming their diagnosis, and why he should not enjoy the advantage of his art *in addition* to all those possessed by others. Let, then, auscultation and percussion be regarded as ancillary only to other, or rather as two among many, modes of investigating disease; as two stout and strong additional strings to our bow; albeit the information they afford is in some cases equal, if not superior, to that derived from all other modes combined.

3. *Position of the patient.*

The object to be desired in reference to the position, is so to place the individual to be examined that the parietes of the chest may be put upon the stretch, without contraction of the superficial muscles, or any exertion on his part likely either to induce fatigue, or in any way to affect his perfect ease. For this purpose, when the fore part of the chest is to be ex-

amined, if the patient be able to sit up, I have found sitting upon a chair with a high, rounded, and reclining back, against which the head and shoulders can be supported without any muscular effort, to be the most suitable position. He should be placed opposite to a good light, with the shoulders thrown back, and the arms hanging down by the side of the chair. If a chair with a high reclining back be not at hand, the patient may be seated on an ordinary chair, with his arms passed over its back, and his head and shoulders as far retracted as his ease will admit. When the side, or axillary space, is to be examined, the patient's hand should be placed upon the occiput or back of the neck, and the body inclined to the opposite side. To explore satisfactorily the posterior regions, a "good back" should be made by inclining the head and bending the body forwards, and folding the arms across the breast. At the same time the scapulæ must be firmly pressed down by the examiner, as, by their elevation, they otherwise remove the supra- and infra-spinal fossæ from those parts of the lung which they ordinarily overlie, the careful examination of which is particularly important, particularly in phthisis.

Exactly the same plan may be adopted in exploring the posterior and lateral parts of the chest, if the patient keep his bed, but is able to sit thereon. If he be incapable of enduring this fatigue, he must be turned, while lying, to either side, as far

as his strength and circumstances will admit. The examination of these parts, in such cases, must almost necessarily be any thing but satisfactory. In the exploration of the front of the chest, on the contrary, an arrangement can generally be easily made so as fairly to examine every part without materially interfering with the supine position of the patient. This may be advantageously effected, without much trouble, by placing two pillows below the shoulders, and one only below the head, so that the former may be higher than the latter, and the chest be thereby somewhat rounded. If the position of the body be favourable, and we have the benefit of a good light thrown directly upon the front of the chest, it is not necessary that the surrounding curtains and furniture be particularly regarded, as they have really very little or no effect in modifying the sounds proceeding from that cavity in the process of exploration. It is, however, desirable that as little noise as possible should be made in the chamber, or ward, in which the examination is being made, and it is especially important that no part of the patient's or the examiner's dress, or any thing in contact with either, should itself, by friction, give rise to a rubbing or rustling noise, which might interfere with the purity of, or be actually confounded with, those proceeding from the chest.

4. *Division of the Chest into regions or compartments.*

Opinions of the nature of diseases are materially influenced by their locality. The most important diagnostic signs are frequently elicited by a comparison of the indications afforded by one part of the chest with those derived from another. It is often important accurately to define the limits of disease affecting certain organs; to be able to mark their progress or their secession, or to be assured of their remaining stationary; to determine if the organs themselves are in their natural situation, or to ascertain to what extent they have diverged therefrom. It has consequently been found convenient to divide the chest, as well as the abdomen, into various regions or compartments. For this purpose vertical lines passing between certain fixed points, and transverse lines passing round the body on a level with other fixed points, and intersecting the vertical lines, have been employed.

The following divisions are those usually adopted, with the exception of some slight modification of the useful additions and alterations advocated by Dr. Forbes, in his valuable articles, "Abdomen," and "Chest, exploration of," in the "Cyclopædia of Practical Medicine." By three vertical lines, one passing on each side of the sternum, and the other running the whole length

of the dorsal and lumbar spine, the chest is first divided anteriorly into a central position, and a right and left side, and both the chest and abdomen are posteriorly divided into equilateral compartments. A vertical line is then drawn from the scapular extremity of the clavicle, on each side, to the spinous process of the pubis; another from the posterior boundary of each axilla perpendicularly to the crest of the ileum, and another along the inner margin of each scapula, between two transverse lines to be immediately mentioned. The transverse lines encircle the body—1st, on a level with the clavicles; 2dly, on a level with the junction of the cartilage of the fourth rib with the sternum; 3dly, on a level with the extremity of the xiphoid cartilage; 4thly, on a level with the end of the last rib; and, 5thly, on a level with the spinous processes of the ossa ilii. All these points are well defined, can be almost always distinctly felt, are not liable to vary (excepting from disease), and may be, without difficulty, remembered. By these vertical and transverse lines the chest is divided into a superior, two central, and seven other regions upon each side. The superior, or *acromial* region, embraces all the space between the lower part of the neck and the level of the clavicles, but is superiorly bounded behind by the seventh cervical vertebra, and before gradually tapers off or merges into another region at the upper end of

the sternum. The two central regions are, the *Superior sternal* and the *Inferior sternal*. Of the other regions, two upon each side are anterior; viz. the *Subclavian*, or *Infra-clavicular*, and the *Mammary*; two are *lateral*, the *Axillary* and the *Sub-axillary*, or *Lateral*; and three upon each side are posterior, the *Scapular*, the *Inter-scapular*, and the *Sub-scapular*.

The abdomen by these lines is divided into thirteen regions. Of these, nine are more or less anterior; namely, the *Epigastric*, the *Umbilical*, and the *Hypogastric*, which are all central; and the *Hypochondriac*, the *Iliac*, and the *Inguinal*, situated upon each side of them. Four of the abdominal regions only are posterior; viz. the right and left *Dorsal*, and the right and left *Lumbar*.

Some of these regions, which are commonly and properly described as belonging to the abdomen, are very often implicated in diseases of the chest, as the lungs descend laterally and posteriorly much below the level of the xiphoid cartilage, and the heart sometimes materially encroaches upon the epigastrium. On this account it is desirable that the regional division of the abdomen should be introduced together with those of the chest. The student is presumed to be acquainted with anatomy. It is therefore not necessary here, at least, to introduce any observations respecting the particular relation of individual

organs to the various regions which have been thus succinctly marked out, and the boundaries of which, so far as regards the anterior and lateral surfaces of the chest, will be sufficiently evident by a glance at the plate at the commencement of this volume. The posterior boundaries are so clearly defined by prominent parts, as to require no illustration.

5. Different modes of exploration in diseases of the Chest.

The means adopted for investigating diseases of the chest by physical signs, in contradistinction to the rational symptoms, are principally included under the terms Inspection, Palpation, Percussion, Auscultation, Mensuration, and Succussion. Upon each of these it is proposed to offer a few brief directions and observations.

CHAPTER II.

INSPECTION, OR OCULAR EXAMINATION.

By Inspection is intended the simple viewing of the form and working of the parietes of the chest. This mode of examination, though too often little regarded, and much too lightly performed, is certainly only secondary in importance, as a means of diagnosis, to auscultation and percussion; if indeed it may not in some cases be properly esteemed of higher value than either of them.

1. Method to be pursued, and circumstances to be noticed, on inspecting the Chest.

In order that the contour and play of the thoracic parietes may be observed, it is obvious that the surface must be either entirely bare, or clothed with a tight elastic vest. For the efficient performance of Inspection, therefore, a certain amount of exposure, even in the case of females, can be scarcely avoided. This exposure, however, need be but very trifling. If the patient be in bed, the lower part of the thorax may be concealed by the bed-clothes while the superior regions are sub-

mitted to inspection, and these in their turn may be covered with a light shawl or handkerchief thrown loosely across the shoulders while the inferior regions are submitted to the same process. When the patient is seated in a chair her own garments can be easily arranged so as to perform the same office as the bed-covering. If this be delicately managed, and the vestments be studiously arranged by the examiner, so as to conceal whatever is not required to be exposed, I believe that little or no difficulty will be generally experienced in acquiring, without offence to the most fastidious, all the information which Inspection is calculated to afford. It is rarely, very rarely indeed, necessary that the *mammæ* themselves should be exposed. The glands, if naturally developed, form so dense a covering to the portion of the *parietes* over which they lie, as generally to conceal the irregularities and deficiencies immediately beneath them.

Upon inspecting the posterior portion of the chest it is merely necessary to place a pillow behind, or to gather the bed-clothes around, the loins, and to retain the light shawl or handkerchief across the shoulders, to obviate any appearance of indelicacy. When these arrangements have been made, the capacity and the general form of the chest should be first noticed. The shape of the sternum and the connected cartilages of the ribs should be observed, and any unnatural

prominence, shortness, or deformity in either of them should be noted. It should be ascertained if the sides are equal in size and in form; if any flattening, contraction, or prominence, exist in one or other, or any part of either of them, and if any fulness of the præcordial region, any unusual separation and bulging of the intercostal spaces, or prominence of the hypochondria, be present.

After the form of the anterior and lateral regions has been accurately observed, which may be effected almost at a glance, the movements of the chest during respiration should be examined. In the ordinary respiration of health the motions of individual parts of the chest, upon inspection, appear to be comparatively slight, in consequence of the cavity being enlarged in every direction nearly simultaneously. They resemble the easy ebb and flow of a soft wave. When, however, a deep inspiration is taken, it is observed that the sternum is slightly but steadily projected forward; that the abdominal parietes are gently dilated; that the lower ribs are first and most considerably raised, and that the elevation of the separate ribs takes place,—(the gentle cost wave moves)—gradually, evenly, and regularly upwards, notwithstanding which each and all appear to move at the same time. Every part acts separately, but each in perfect harmony with the other. At the same time, if the patient be sitting up, or lying upon his back, there may be generally

noticed a gentle rising or undulation below the left breast, dependent on the impulse of the heart against the parietes. This undulatory motion of the general parietes usually occurs, in health, about eighteen times in a minute, whilst that dependent on the impulse of the heart is commonly four times as frequent.

The examiner, then, should first observe the movements of the parietes in ordinary respiration; he should subsequently direct the patient to take two or three deep but slow inspirations, and should take notice if every portion of the chest be equably raised, if the ribs move freely and (if I may use the expression) independently of each other, and should be especially careful to compare the action of one side with that of the other. Posteriorly, the general configuration of the chest, and the mobility of the ribs, should be examined with similar attention. But in addition to the observations made upon the anterior surface, the shape of the spine should be particularly noticed, as upon the irregularities and curvatures of this bony column very often depend contractions and other alterations in the form of the chest, which, without such notice, might not improbably be attributed to diseases of different origin and of another kind. The discovery of a simple curvature of the spine not unfrequently leads to the satisfactory explanation of many symptoms, and the elucidation of various physical signs, which had

been previously unwittingly ascribed to other causes.

2. *The indications afforded by Inspection.*

When a portion of the pleura lining the ribs is affected with inflammation, the parietes connected with or overlying the part, in consequence of the pain induced by motion, are less raised during inspiration than other parts of the thoracic walls. When, after the subsidence of the inflammation, the lung has, through the medium of the effused inflammatory product, become adherent to the parietes, the part affected, though it may be raised to an equal extent with the corresponding part upon the opposite side, is moved less freely or more *stiffly* than previously to the attack. It is raised as an unyielding case, rather than with the suppleness and pliancy observed between the ribs in other parts. When, again, considerable effusion has taken place into one of the pleuræ from inflammation of the membrane, the lung is compressed, and a portion of the air squeezed out; less air than in the healthy condition gains admission into its cells in consequence of that compression, and the ribs therefore covering the effusion are less raised during the act of inspiration. When the fluid has been present a considerable time, and in large quantity, not only is the side or part of the side in which it exists immoveable during respiration, but it becomes

enlarged or misshapen, bulging at one part and contracted at another. When, again, the fluid has been at length absorbed, the side not only remains less pliable than before the attack, but, partly in consequence of the imperfect expansibility of the formerly compressed lung, already bound down by adhesions, which prevent its expansion commensurately with the absorption of the fluid, and partly in consequence of the contraction of the newly deposited fibrinous matter, it becomes flattened, or "falls in," to an extent varying in different cases. Hence it follows that imperfect elevation, defective pliancy, or perfect immobility of the ribs, and that enlargement or contraction of one side, or of a portion of the thoracic parietes, become physical indications of an attack of pleurisy, or of its recent or remote consequences.

Now as pleurisy almost always coexists with a considerable deposition of tubercles, if near the surface of the lung; as these bodies are almost uniformly first deposited in the upper lobes, and most commonly on one side sooner, or in greater number, than on the other; as the adhesion resulting from this pleurisy interferes with the free expansion of the portion of the lung which it has invested; and as moreover the presence of tubercles in the lung must exclude some portion of the air, which would otherwise have reached the air cells, it follows that imperfect elevation, or

flattening, of one or both infra-clavicular regions, may become, and not unfrequently is, one of the first physical indications of tubercular phthisis.

In the more advanced stages of phthisis the indications afforded by Inspection are more striking and definite. When, for example, the upper part of the lung has become invested with a thick and dense pleuritic coating, which is firmly united to the parietes, and when cavities have formed in the lung, and have partially collapsed, the flattening and defective mobility of the regions in which the disease exists are necessarily much greater. It should, however, be recollected, that phthisis may exist in its advanced, as well as in its incipient, stage without this flattening and imperfect elevation of the parietes. This fact is sometimes observed in cases of very acute disease, or in those examples of the complaint in which the deposit and its consequences are almost entirely confined to the upper and posterior parts of the lung.

In pneumonia accompanied, as it usually is, with pleurisy, the indications afforded by Inspection are similar to those existing in phthisis, and dependent in a great measure upon similar causes. An important distinction, however, it must be recollected, exists in the fact that phthisis almost always first affects the upper lobes; while simple pneumonia, in a great majority of instances, attacks the lower lobes of the lungs.

When malignant disease in a massive form invades a large portion of a lung or pleura, the parietes covering it are imperfectly raised during inspiration, in consequence of a diminished quantity of air reaching the cells, and sometimes, also, in consequence of pleuritic adhesions. As the disease advances, the parietes become misshapen, either from bulging outwards, or from contraction.

In emphysema of the lungs the volume of the organ is increased, in consequence either of the simple dilatation of the air-cells, or of the escape of the air into the space between the lobules, or beneath the pleura. The pulmonary tissue partially loses its natural contractility. The air is not efficiently expelled during expiration. The lungs are, in fact, more full of air than in the healthy state, and are incapable of ridding themselves of their abnormal burden. The process of the production of emphysema (independently of accidents and a few very rare cases) is in general a slow and gradual one; each successive attack of bronchitis or bronchial obstruction contributes its small proportion. Every act of coughing tends to increase the general effect. In process of time the continued pressure of the lung upon the interior of the parietes effects their dilatation, and, if the disease be general, causes them to assume a rounded form. The patient is capable of taking but little air into the lung, in

consequence of the organ being already distended, and the parietes of the chest being already considerably dilated. Hence it follows, that a full rounded form of the parietes, together with imperfect elevation of the ribs upon inspiration, is one of the physical signs of emphysema of the lung.

In very long-continued cases of general emphysema, a notable exception to what has been stated may be observed in the lower portion of the chest. The ribs of the lateral regions are then bent inwards, and are depressed instead of being elevated, or approach to, rather than diverge from, the centre of the body, during inspiration. This appears to arise from the ribs having been already elevated, and the bony parietes permanently dilated to nearly the utmost extent, so that they at length become almost motionless during the inspiration. The comparatively little air which is taken into the chest gains admission, therefore, by means of the contraction of the diaphragm, the connections of which muscle to the inferior ribs cause them to be drawn inwards and downwards during its action. After long continuance of this inward motion during inspiration, the ribs at the parts indicated become permanently depressed, as in slighter and less persistent cases of emphysema they become permanently elevated from a somewhat similar cause. This inward inclination

of the lower ribs becomes very obvious upon inspection, when emphysema has existed for a long time, and the patient is already advanced in life.

When fluid of any kind is present in the pleura in considerable quantity; if, that is to say, the complaint called hydrothorax or pneumothorax exist, the air-cells cannot be filled, the lung expanded, or the chest dilated upon inspiration, as in health. Imperfect elevation of the ribs, or total immobility of one side of the chest, therefore, become characteristic signs of these complaints.

When the heart has been enlarged for a considerable time, or when fluid effusion has long existed in the pericardium, a general fulness or roundness of the præcordial region may be sometimes observed. When aneurisms of the large vessels of the heart have caused absorption of the internal parietes, or have merely become adherent thereto, a slight or considerable elevation, according to circumstances, may be frequently noticed, which sometimes increases synchronously with the pulsation of the artery. The consequences of fractures, penetrating wounds, or of abscesses affecting the internal organs, are also commonly indicated by Inspection of the parietes.

The diseases, then, in which diagnosis may be rendered more clear by inspection of the chest, and in which, therefore, it may be advantageously employed, are pleuritis, and its consequences,

whether in the form of simple effusion, of empyema, or of contraction, pneumonia, phthisis, emphysema, hydrothorax, pneumothorax, enlargement of the heart, hydropericardium, or other pericardial effusions, aneurism of the large vessels, fractures, abscesses, &c.

In every primary examination, and in all in which an opinion is expected to be delivered as to the condition of the contained organs, Inspection should never be omitted. It should, however, be constantly borne in mind that curvatures of the spine, and congenital or acquired malformation of the skeleton, may so entirely change the configuration of the parietes as to cause them to present, upon mere Inspection, the appearances which ordinarily coexist with almost every one of the diseases which have been mentioned.

CHAPTER III.

PALPATION, OR MANUAL EXAMINATION.

THIS mode of examination, by which the sense of touch affords its aid to the other senses in the diagnosis of disease, notwithstanding what has been said concerning the "*tactus eruditus*," requires but little skill or experience for its exercise, so far at least as relates to the complaints of the chest.

1. *Mode of employing Palpation.*

Palpation is chiefly used in two ways. The first of these is by pressure of the sentient extremities of the fingers, as in the ordinary exercise of the sense of touch, and requires no comment or explanation. The other mode is by placing the palmar surface of the appressed fingers, or of the whole hand, upon the part to be examined, and using such an amount of pressure as to enable it to participate in the vibrations, or to appreciate the defective motion, of the parietes of the chest, and then comparing the impressions communicated to it with those communicated to the other hand, similarly placed upon the corresponding part of the opposite side.

2. *Indications afforded by Palpation.*

When it is inconvenient to expose the chest of a patient—when, for example, the *appearance* of delicacy might be considered to be thereby sacrificed—or when there may be danger, or a liability to an increase of symptoms by the surface of the body being bared, the appressed fingers of one hand placed flatly, and pressed firmly upon the infraclavicular region of one side, while the other is similarly placed and pressed upon the corresponding region of the other side, are often capable of distinctly appreciating a flatness of one side, or a difference in the pliability or expansibility of the two sides, in the early stage of phthisis.

When it is doubtful whether the eye has correctly measured a supposed slight difference in the action or mobility of the two sides, Palpation, exercised in the mode just mentioned, may be usefully employed to remove or to confirm that doubt. Any considerable deficiency of elevation of the ribs, or mobility of the parietes, or any great disparity between the two sides, is immediately detected, even without exposing the patient, by the hands placed upon the lateral regions—whether that deficiency depend upon pleuritic effusion, hydrothorax, pneumonia, pneumothorax, or any other cause. The exact nature of the complaint must be decided by farther examination of a different kind.

When disease obviously exists in one side of the chest, and the lung is clearly consolidated from some cause, it is sometimes difficult to decide whether that consolidation arise from structural disease of the lung itself or from pressure upon its exterior, as from effusion in the pleura, &c. In such cases assistance may be occasionally derived from Palpation; for though, when fluid exists in the pleura to a small amount, the vibration of the voice is communicated to the parietes and may be appreciated by the ear, it is not distinguished by the hand.

The explanation of the fact appears not to be clear—but the fact itself is now acknowledged to be certain—that the vibration of the voice is communicated through the parietes to both the ear and hand when the lung is consolidated, as from pneumonia, and that it is communicated to the ear and not to the hand in some cases, at least, of effusion into the pleura. The most probable explanation appears to be that, though the sound may be more shrill, it is not so loud in effusion as in consolidation, and though the vibrations are more frequent, they are less strong in the former than in the latter.

Whatever may be the explanation of this curious circumstance, Palpation may thereby sometimes become a means of distinguishing pleuritic effusion from consolidation of the lung, and thus afford aid in a point of diagnosis which is not always sufficiently distinct.

When the surface of the pleura or pericardium, which naturally glide smoothly, is roughened by solid effusion between its layers, a rubbing or grating noise is heard upon applying the ear or stethoscope to the neighbouring parts of the chest. The vibration thereby communicated to the parietes can often be perceived by the hand placed, as previously directed, upon the part affected.

The same, with slight modifications, may be said in reference to loud noises arising from obstruction to the passage of air in the bronchial tubes, whether that obstruction depend upon thickening of the lining membrane, external pressure, or superabundant secretion.

By the sense of touch the position of the heart may be generally ascertained, and, in some of its diseases, important information may be derived from manual examination. While the body is erect, the heart, when in a natural condition, is commonly felt to strike the parietes about an inch below and to the inner side of the nipple. While lying upon the back its impulse is greatly decreased, and is usually felt somewhat nearer the sternum. When the body is turned to the left side, the impulse is felt in a direct line with, or often nearly an inch to the outer side of a line passing vertically over, the nipple; while on the contrary, when the body is turned to the right side, it is felt between the cartilages of the ribs, close to the sternum, or sometimes cannot even

be felt at all. These changes are stated not to take place when the pericardium is adherent, and the want of these, therefore, is assumed to be diagnostic of that condition. But this fact I have myself never had an opportunity of verifying in simple adherent pericardium without enlargement of the heart, though, when the organ is enlarged, it is well known that the exact position of the impulse is sometimes not clearly defined.

When the parietes of the heart are thickened, and the force of its impulse is consequently increased, the hand placed over the præcordial region becomes at once sensible of its abnormal force, though the pulse at the wrist may at the very same time be small and feeble.

When the cavities of the heart are dilated, with or without any increase of the thickness of their walls, the impulse is often perceptibly extended over a larger space than natural, and may be felt not only above, below, and around its ordinary site, but also in the scrobiculus cordis, and sometimes even on the right of the sternum.

When the heart is removed from its natural situation by gaseous or fluid effusions into the pleura, by tumors, abscesses, &c., it is by manual examination that the fact can generally be best determined.

When obstruction exists in the valves, a trembling motion, or "purring tremor" ("*frémisse-*

Unable to display this page

CHAPTER IV.

PERCUSSION.

PERCUSSION is the art of striking the parietes of the cavities of the body in such a manner as to enable the examiner to judge of the amount of resonance of the parts beneath, and is one of the most important means of physical diagnosis in diseases of the chest. When the bony parietes of the chest are smartly struck by the knuckles, or by the tips of the appressed fingers, the resulting sound varies according to the nature of the organ, or the part of the organ which lies under the portion of the parietes which is struck. The healthy lung is a light spongy tissue, the cells of which, and the tubes leading to which, are occupied by air. If the ribs, then, be struck over a portion of this tissue, when in a healthy condition, the contained air, like that within a cask or drum, resonates with the parietes; or, in other words, by its elasticity and compressibility, allows such an amount of free vibration of the parietes as to produce a deep hollow sound. If, on the other hand, the chest be struck over a portion of the lung from which the air has been excluded,

either by pressure from without, as in pleuritic effusion, or by deposit within, as in pneumonia or pulmonary apoplexy, then, in consequence of the parts below resonating only very imperfectly with the parietes, from defective compressibility and elasticity preventing their free vibration, a dull, heavy, or dead sound is produced—as in a cask filled with water, or a drum with dough. When the portion of the parietes which covers the heart is similarly struck, the resulting sound is also dull, in consequence of the slight amount of compressibility and elasticity possessed by that organ and its contained fluid. When, therefore, the heart is considerably enlarged, when its investing membrane is distended with effused pus or serum, or when its principal vessels are greatly enlarged by aneurism, the extent of dulness in the præcordial region is increased. These few examples will suffice to illustrate the great importance of percussion in the diagnosis of diseases of the chest. The practice of this mode of exploration, apparently so simple, requires great nicety, tact, and delicacy, for its efficient application. Great care is also required, in its exercise, that the examiner be not misled and deceived by trifling circumstances connected with the manner in which it is performed or with the parietes of the patient.

1. *Different Modes of Percussion.*

Percussion may be either mediate or immediate. It is immediate, when the striking body, whether it be the hand of the examiner, or any other instrument, falls directly upon the parietes of the chest, or when some article of clothing only intervenes between them and the percussing body. It is mediate, when some solid material, as a disc of wood or ivory, a piece of leather, or the finger of the left hand, is interposed between the parietes and the striking body for the purpose of increasing or modifying the sound.

1. *Immediate Percussion* may be performed with any suitable instrument, as the end of a stethoscope, or with the knuckle of the contracted fore-finger of the right hand, but is most efficiently practised with the points of the appressed fingers. The extremities of the fore, the middle, and the ring finger, should be pressed close together, and brought as nearly as possible to a level surface, beyond which the nails should not extend, and to which they should scarcely reach. The patient being in the position pointed out in a preceding chapter, and the parietes being rendered tense, the knuckle of the fore-finger, or the points of the appressed fingers arranged as just explained, should be smartly struck, in a direction perpendicular to the surface, successively upon the

sternum, clavicle, the ribs, and scapulæ. The corresponding regions of the two sides should be percussed immediately after each other, so that the resulting sounds may be accurately compared, as dulness, not absolute only, but comparative, often affords most important indications. It may be well to state that the tips, the actual bony extremities, of the fingers should be the parts coming in contact with the chest, and not the soft bulbous sentient palmar surfaces of the last phalanges, otherwise the sound will not be so clear, as the percussing instrument will not be so firm, and therefore not so well calculated to elicit it. The examiner should also be especially careful that the fingers, or other percussor employed, strike the chest at a right angle with the surface struck; as, if they fall upon the parietes in a direction only slightly oblique, the effect will be greatly diminished.

2. *Mediate Percussion*.—Direct or immediate percussion may be sometimes employed with advantage, particularly when great nicety of discrimination is not required, and when it is not necessary that the examination should be prolonged. It is, however, when extensively employed, open to some and not slight objections. When, for example, the patient is an irritable or nervous person, each stroke of the fingers or percussor causes excitement, and the frequent repetition of

the slight tappings upon the unprotected chest induces a state of the circulation and respiration entirely opposed to the calm and quiet condition, which it is so desirable should be preserved during the future stages of the examination. When the patient is thin, or the skin delicate, the direct percussion of the slightly covered osseous structures causes inconvenience, sometimes amounting to pain. When the individual, on the other hand, is fat, or very muscular, when the integuments are œdematous, or when in the female the mammary glands are large and full, or loose and pendulous, direct percussion cannot be employed with advantage, in consequence of the dense and non-elastic soft tissues preventing the stroke of the finger reaching the bony parietes with sufficient force and sharpness to induce effectual vibrations. There are also some parts, in which, from the absence of bony covering, immediate percussion is entirely useless.

On these accounts, and perhaps some others, mediate percussion has been employed, and a variety of instruments have been proposed for its performance. Thin discs of wood, ivory, cork, and India rubber, with and without different kinds of tongues or lips to be used as handles, have been introduced and advocated under the name of *Pleximeters*, for the purpose of receiving and conveying the stroke of the finger, or of a variety of little hammers, under the title of

Plessors. Each of these may possess some advantages, and each has had something and some one to recommend it for general employment, but each participates more or less in the objection of itself giving rise when struck to a sound, which interferes with that dependent on the vibration of the thoracic parietes.

If any substance intervene between the striking instrument and the chest, if any *pleximeter* be employed, it is clear that its vibratory power should be as similar as possible to that of the vibrating parietes ; that it should be composed of materials identical with them,—should in fact be made of bone covered with skin. Such materials we have already prepared. A *pleximeter* of bone covered, with skin admirably adapted for the purpose, and always ready for use, we possess in the fingers of the left hand. This pleximeter possesses many advantages in addition to those already mentioned.

It is not open to the objection of sometimes causing pain by firm pressure upon fat and thin persons, as are the wood or ivory discs which are occasionally employed. It is capable of being much more nicely adjusted to uneven surfaces than either these rigid substances, or stiff leather, and can be much more easily maintained in the required situation than flexible but elastic India rubber. In thin persons the intercostal spaces form hollows which cannot be filled by unyielding pleximeters; air consequently

intervenes between them and the parietes, and by its resonance, when the pleximeter is struck, increases the sound dependent on the vibrations of the instrument itself, and thereby adds to the sources of fallacy thence arising. The stiff pleximeters, moreover, cannot be conveniently applied in the axillæ, or, in very thin persons, above or immediately below the clavicles. The firm pressure of the fingers on the contrary causes no pain; they are capable of being perfectly adapted to the surface of the chest, however uneven; they are therefore equally efficient as pleximeters in fat and thin persons; they can be applied in the hollow of the intercostal spaces; they can be employed in every part of the chest, and, when struck, give rise to very little independent sound. On these several accounts it is believed that the very best *pleximeter* is produced by the fingers of the left hand, and that the very best *percussor* is produced by the appressed fingers of the right hand.

Having, then, decided upon the instruments to be employed in Percussion, the next question to be considered is the mode of employing them. The fingers of the right hand being arranged as before mentioned, the palmar surface of the fore, the middle, and the ring fingers (one or all) is to equably placed and firmly pressed upon the part to be examined. If the whole chest is to be submitted to the operation, it is convenient to begin

above and to pass regularly downwards. The fingers should be so firmly pressed as to make them as much as possible one with the parietes, so that the impulse given to the fingers may be directly communicated to the parietes, and that the vibrations of the latter may be correctly appreciated by the former. It is essential that the whole extent of the finger or fingers should be equally and firmly pressed upon the surface of the chest. If the pressure be unequal, and particularly if one portion of the finger struck be in contact, and another portion not in contact with the skin, either a very imperfect sound is elicited upon percussion, or a noise similar to that produced by striking a cracked earthen vessel (*"bruit de pôt fêlé"*) is brought forth. The points of the three appressed fingers of the right hand are now to be several times smartly struck upon the middle phalanx of one of the fingers of the left hand thus carefully placed, and, after the impulse is communicated, to be immediately withdrawn from contact with it.

After the sound elicited by the Percussion of one region has been thus cautiously elicited, and carefully observed, the fingers of the left hand should be removed to, and similarly placed upon, the corresponding region of the opposite side of the chest, and the sounds caused by Percussion upon each region, and every part of each region, should be compared with those proceeding from the other. When the præcordial region is per-

cussed, this comparison of the two sides of the chest is of course not available. I have said that the stroke should be directed to, and concentrated upon, the middle phalanx of *one* finger. Which finger this may be is perhaps a matter of indifference, but it is better, for the sake of the skill and aptitude acquired by habit, always, excepting in particular circumstances, to employ the same. The middle finger, closely flanked and supported by that on each side of it, is that which I, from habit, always employ.

In mediate, as in direct, percussion the direction of the stroke should be perpendicular to, or at right angles with, the surface to which it is applied; otherwise much of the effect will be lost, and incorrect conclusions may be the result. If any space or spot be found unusually dull, or particularly resonant, upon either side, the examiner should first bring to mind, as far as he is able, the natural amount of resonance of that particular part. If in the normal condition it be similarly dull, or resonant, as the space over the heart and liver in the one case, and that covering the stomach in the other, he must attempt to define the extent, and, as far as possible, the degree of dulness or unusual resonance, and then mentally compare it with that existing in the healthy well-formed chest, to which he should be able to refer as to a standard, and with the sounds of which he should therefore be perfectly familiar.

It must, however, be acknowledged, that the

assumed standard itself is exceedingly variable, and that it is quite impossible to define exactly the amount and extent of dulness in the healthy condition. In two individuals equally free from any indication of disease, and equally well formed, the extent of the præcordial dulness, for example, may vary considerably. An approximation to a healthy standard, however, may be attained by practice, and with this the comparison must be made.

If, on the contrary, the part found to be dull should, in the natural state, be moderately resonant, it should be first ascertained if there exist any cause of dulness exterior to the chest. If none be discovered, and it be determined that the cause exists within, the examiner should next attempt to decide whether it arise from solid, or from fluid material, and whether it be deep, or superficial, or both. This latter particular is effected by varying and regulating the force of the impulse communicated by the stroke of the fingers. Thus, if between a solid organ and the parietes a thin layer of healthy lung intervene, a slight and delicate stroke, or tapping, will elicit the resonance afforded by that layer, while a firm heavy stroke will detect the dulness caused by the deeply seated solid body. Conversely to this, if the source of the dulness be only superficial, a light stroke will afford little or no sound, and a firm and strong one will afford proof of the presence of a resonant substance below.

These opposite conditions may be fairly illustrated by first percussing the lower part of the right mammary and lateral regions, which are naturally resonant when only light, and dull when strong, Percussion is employed ; and then similarly examining the scapular regions of a thin person, in which an exactly contrary state of things naturally exists.

Beginners, and other inexperienced persons, are generally in the habit of giving much too strong and firm a stroke while exercising Percussion ; forgetting, or not being aware, that to distinguish minute differences, or delicate shades in the quality, as well as the amount, of resonance—as, for example, in the comparison of that elicited from the subclavicular regions in the early stage of phthisis—light Percussion is far more efficient than the heavy strokes which are frequently employed. A gentle tap will often display a marked difference in the two sides, when a strong blow will be utterly inoperative for the purpose.

When the source of the dulness has been proved to be superficial only, or superficial as well as deeply seated, the examiner should endeavour to ascertain if the material causing the dulness be fluid or solid. This may be sometimes, but certainly not always, determined by combining Palpation with Percussion. In other words, it is effected by estimating the amount and character of the resistance communicated by

the parietes of the chest to the finger of the left hand at the time the stroke is inflicted by those of the right hand. Thus, if the cause of the dulness be a solid, the resistance afforded by the parietes is greater, and the impressibility, if I may so term it, or the resiliency, less than if it have a fluid origin. If, on the contrary, the source of the dulness be fluid, the resistance is less, and the resiliency greater, than when it proceeds from a solid material. The capability, however, of appreciating this difference, which is merely one of degree, is not likely to be attained but by long habit and experience, and cannot always be appropriated by the most skilful of auscultators. Fortunately there are other means of determining this sometimes important, but often difficult question.

2. Percussion in the normal state of the Chest.

To be capable of judging accurately of the sounds which are morbid, it is clearly necessary that the examiner should be familiar with those which are natural, and therefore that he should be conversant with the amount of resonance on Percussion in the different regions of the chest in persons free from disease. This varies considerably in different individuals, quite independently of shape or congenital malformation. This difference of resonance on Percussion in healthy persons depends upon a variety of circumstances;

among which may be enumerated the size of the chest, the thickness and looseness of the skin, the presence, absence, and amount of fat below the integuments, the development and firmness of the muscles, and the general elasticity or tonicity of the tissues. Still, when the internal organs are healthy, and when the external parts are natural in their congenital formation, and have not been materially affected by belts, stays, or other articles of dress, the various regions bear, in regard to resonance on Percussion, a definite relation to each other, with which it is absolutely essential that the student of auscultation should be acquainted.

3. *Natural resonance of the different regions of the Chest upon Percussion.*

It has been stated that even in health the regions of the chest vary in their amount of resonance on Percussion in different individuals. The remark is applicable to the thorax generally, but it is more especially true in reference to the *acromial* and some other regions. In stout persons with short necks, indeed, percussion of the acromial region affords no information, as the entire space which it embraces is uniformly dull. Even in thin persons it requires some tact and management to make the part resound. It is, however, so important carefully to examine this region, and to

compare one side with the other, particularly in reference to phthisis, which sometimes almost solely affects the very apices of the lungs, situated from an inch to an inch and a half above the lower edge of the clavicles, that the information derived from the careful investigation of this region will amply repay the examiner for the expenditure of a little extra time and trouble. It is the neglect or superficial observation of this region which undoubtedly sometimes causes the disease to be overlooked.

To be enabled to judge of the comparative resonance of the two sides—and it is to this that the chief value of the examination is attached—the shoulders of the patient must be firmly depressed, and the head and neck slightly inclined to the side opposite to that which is being examined. The finger of the left hand is then to be strongly pressed downwards, inwards, and backwards, above the clavicle, close to the bottom of the neck, and firmly percussed by those of the right hand. When the amount of resonance has been tested upon one side, the other should be similarly explored. The fingers should then be removed to the top of the shoulder and similarly percussed, and should, alternately upon one side and the other, be successively passed round to each portion of the space between the scapular end of the clavicle and the spine. By careful management, firm pressure, and strong percussion, a considerable amount

of resonance, though inferior to that of some other regions, may be brought out, and a difference in the corresponding portions of the two sides detected. It must be, however, repeated, that even in the most favourable circumstances the resonance of this region is defective compared with that of some other regions.

The infra-clavicular or subclavian regions are fairly resonant, excepting in stout muscular subjects, when the large pectoral muscles materially interfere with the sound on Percussion. The superior sternal region, though occupied in part by the large vessels of the heart, contains also the trachea and its primary division, as well as the approximating upper lobes of the two lungs, which cause it to be very resonant on percussion. The inferior sternal region is almost always dull, from embracing a great portion of the space occupied by the heart. Superiorly, the lungs, in various degrees, overlap that organ, and on this account the upper part of this region is often resonant upon light, and comparatively dull on firm Percussion. The mammary regions vary in their amount of resonance in different parts. On the left side, over the cartilages of the fifth, the sixth, and sometimes the seventh rib, it is rather dull, particularly on firm percussion, in consequence of the left ventricle and apex of the heart being there covered with only a thin layer of pulmonary tissue. It is often on the outer side very resonant, nearly as high as

the fifth rib, from the stomach being distended with gas, and sometimes dull to a less extent, from that organ being replete with food. On the right side the Mammary region is superiorly resonant; but below the sixth rib, or the sixth intercostal space, it is dull on a forcible stroke being inflicted, in consequence of the upward projection of the convex surface of the liver, though a gentle tap will easily detect the resonance arising from the intervening portion of the lung. The Axillary regions, where uncovered by the converging fibres of the large pectoral muscle, are very resonant, as the ribs are therein covered only by the flat serratus muscle, the skin, and loose cellular membrane. The Lateral regions vary like the Mammary, being resonant superiorly, but on the right side dull on forcible percussion, in consequence of the underlying convex surface of the liver, and on the left side variable, according to the state of repletion or emptiness of the stomach, and the nature of its contents. In its inferior and posterior part it is sometimes also rendered dull to a variable extent by the greater or smaller size of the spleen. When this organ and the liver are enlarged, or pushed upwards by large abdominal effusion, they, independently of any thoracic disease, materially interfere with, or entirely destroy, the natural resonance of both lateral regions. The Scapular regions are dull, from thick layers of muscles which intervene between the integuments

Unable to display this page

disease in the chest, but influenced by the constantly varying condition of the hollow organs of the abdomen, may be arranged as follows, according to their amount of resonance.

Tympanitic, (occasionally,) from distension of the stomach, &c., the lower part of the left Mammary, left Lateral, and left Infra-scapular regions.

Very resonant.—The Superior Sternal, the Axillary, and the upper part of the Infra-scapular regions.

Resonant.—The Subclavian or Infra-clavicular, the upper part of the Mammary and Lateral, and the Inter-scapular regions, excepting the inner edge of the left Mammary.

Imperfectly resonant.—The Acromial and the lower part of the right Mammary, Lateral, and Infra-scapular regions.

Dull.—The Inferior Sternal, the inner edge of the Left Mammary, and the Scapular regions.

4. *The indications afforded by Percussion.*

1. *In diseases of the Lungs and Bronchial Tubes.*

As in simple *Bronchitis* there exists merely an increased supply of blood in, together with some thickening of, the lining membrane of the tubes, there is no appreciable alteration of the resonance of the chest. Natural resonance of the chest, therefore, is one of the signs of simple bronchitis. The consequences of the complaint, however, often modify the sound elicited by percussion. Thus, in the debilitated patient, pulmonary congestion,

particularly of the more dependent parts of the lung, may be a direct result; and this may exist to such an extent as to be accompanied by dulness on percussion. From the long continuance, or frequent repetition of bronchitis, emphysema arises, which is attended with increased resonance of the chest.

In the early stage of *Pneumonia*, the pulmonary tissue contains a larger supply of blood than in the natural condition, and the air-cells and intervening spaces are, at least, partially filled with serous fluid. The ordinary quantity of air, therefore, cannot gain admission to the interior of the cells, the elasticity of the tissue is decreased, and the resonance on percussion is consequently diminished. When, instead of being partially occupied by fluid, the air-cells are filled with solid or semi-solid matter, or are closed by the thickening of their walls, and the air is entirely excluded from their interior, as in the advanced stages of the complaint,—when, in fact, *hepatization* is established,—the dulness on percussion becomes much more marked, and the resistance afforded to the finger more decided. As, in the progress of cure, the engorgement of the pulmonary tissue decreases, and the cells become partially free, the amount of dulness, together with the resistance communicated to the finger, gradually diminishes.

When, on the contrary, pneumonia passes on to the third stage, and the effused matter softens down, though air may gain admission to the

disorganized tissue, the resonance on percussion does not increase in proportion : excepting in the comparatively rare case of large pneumonic abscess, variation in the amount of dulness is often scarcely perceptible. The comparatively slight change of the dulness in such cases probably arises from the contraction of the surrounding tissue, and simultaneous depression of the ribs, as well as from the commingling of purulent or softened albuminous matter with the inspired air.

Percussion, then, forms an important element in the physical diagnosis of pneumonia. It not merely assists the physician in ascertaining the existence of the disorder, but, when the case is carefully watched, it enables him to determine its stage and progress.

Is dulness on Percussion, then, always attendant upon pneumonia ?—It certainly is not. The disease may certainly be present without any appreciable dulness. Thus, when inflammation affects separate lobules of the lung, disseminated through one or both organs, and these lobules are surrounded by a comparatively large amount of healthy crepitant pulmonary tissue, no dulness can be discovered on percussion. It is from this form of the disorder being more frequent among children, as well as from physical examination being in them effected with less facility than in adults, and from the patients frequently dying before the disease has arrived at the stage

of consolidation, that the pneumonia of childhood is less commonly attended with dulness on percussion, than that of grown persons. There are other cases in which pneumonia cannot be recognized by dulness: as when a defined portion of hepatized lung occupies the centre of the organ, and is on every side surrounded by healthy structures, or when it exists in situations normally dull, as in the thin edge overlapping the heart, or in that passing down before the liver.*

In *Phthisis* the dulness on percussion is very variable. In some few cases it cannot be discovered during their entire course. Its importance, like that of almost every other physical sign in this complaint, is associated with the comparison of one part with another, or of corresponding regions in the two sides of the chest. Phthisis almost always primarily affects the apex of the lungs, and most commonly that of one before, or to a greater extent, than the other. Upon this circumstance do the distinctive physical signs of the complaint mainly depend. When, therefore, tubercles are largely and equally distributed throughout the whole of one or both

* It has been stated, upon high authority, that in some cases of consolidation of the lung, the side has been more resonant than natural, and even tympanitic, upon percussion. This curious circumstance has been differently accounted for by different writers, but as the author has never witnessed such a case, the exception has not been introduced in the text.

lungs, the characteristic physical signs of the complaint are absent. It has been previously stated, that when tubercles are deposited in considerable numbers in any part of the lung, the pleura covering that part almost always becomes inflamed; plastic matter is effused, which connects the pulmonary and costal layers of the membrane. The expansibility of the pulmonary tissue is thereby diminished, from its free motion being impeded, and from the contraction which sooner or later ensues in the newly formed membranous covering. The tubercles also encroach upon the space to which, in the healthy state, air is admitted.

From these causes less air enters the cells of that part of the lung. Hence, even in the early stages of phthisis, arises a difference of resonance in the corresponding regions of the two sides, or between different regions of the same side. The regions thus affected are almost universally the infra-clavicular, the acromial, or the scapular. There may be no actual dulness either absolute or comparative. Both sides may sound well. But when they are accurately compared by gentle tapping, as well as strong percussion, in a compressed as well as in an expanded state of the lungs, a difference of *tone* or *quality* of the sound elicited by percussion may be often observed. This difference of tone or quality of sound, especially when existing in combination

with other physical signs, should be carefully noted, as an important particular in the diagnosis of incipient phthisis.

When tubercles become more numerous, when the pleuritic coating has increased in thickness and density, and especially when the lung surrounding the tubercles is rendered hard and firm by pneumonia, the sound on percussion not only varies in tone, but differs in degree. Dulness then becomes decided, and obvious upon comparison with that afforded by healthy parts. It may exist upon both sides, but at this period of the complaint it is most commonly present on one side only, or is much more marked upon one than upon the other.

During the stage of softening, the dulness is scarcely altered, unless it be so from an increase and extension of the surrounding pneumonic consolidation,

In the latter stages of the complaint, even when cavities have been formed, unless the vomicæ be large and superficial, the sound upon percussion is far from good; it is nearly, if it be not actually, dull. This arises partly from the thick and dense pleuritic covering, and the layer of consolidated lung, which intervene between the cavities and the parietes, partly from the irregular form of most vomicæ being unfavourable to the resonance of the contained air, and sometimes, in part, from the presence of fluid secretion within them.

The sound upon forcible percussion is not, however, absolutely dull, as in some of the former stages of the complaint, or in the consolidation of pneumonia, but rather resembles that of a sounding body to which a damper has been applied.

When a cavity is very large, occupying the greater part of an entire lobe of the lung, the sound may be not only more than ordinarily resonant, but absolutely tympanitic. Such cases are, however, rare, and as a general rule tubercular cavities are not, as is supposed by many, accompanied with increased resonance on percussion.

In a few cases a peculiar sound is heard, which has not been inaptly compared to that produced by striking a cracked pot (*bruit de pôt fêlé*). It appears to be connected with the imperfect resonance of the air contained within a cavity of moderate size, in consequence either of its being partially occupied by fluid, or of the vibration of its parietes being interrupted by the entrance of one or more large tubes.

In that form of phthisis, then, in which tubercles are generally disseminated throughout the lungs, the resonance of the chest upon percussion is not appreciably altered. In the earlier stages even of the more common form of the complaint, there may exist a variation in the resonance, or rather in the tone of sound elicited by percussion, on comparing the infra-clavicular or acromial with the mammary or axil-

lary regions of the same sides, or corresponding regions of different sides with each other. In the second and third stages of the disorder,—those, namely, of consolidation and softening,—the dulness on percussion is obvious, when compared with the sound proceeding from a healthy lung. In the latter stages of the complaint the sound varies according to the size and arrangement of the cavities: it may be almost dull, it may resemble a muffled resonance, it may be tympanitic, and lastly, it may approach to that arising from striking a cracked pot.

In simple *Venous congestion of the lung*, arising from the difficult transmission of blood through the heart, dulness on percussion exists in a degree and extent proportioned to the amount and duration of the disease, and the debility or exhaustion of the patient. In such cases the dulness is generally observed in the most dependent parts of the lungs, or in those most remote from the centre of circulation. It is not, however, of so marked and decided a character as that arising either from pneumonic consolidation or pleuritic effusion. But when the congestion reaches to such a height as to cause pulmonary apoplexy, in which the part affected is rendered perfectly destitute of air, and reduced to a state resembling damson cheese, the dulness on percussion, and the resistance afforded to the finger, is complete. This disease, it may be observed, often affects

detached masses of the lung, or the narrow edges of the organ, and is consequently not always detected even upon minute examination.

In *Œdema of the lung* arising from diseases of the heart, or from other causes, serum partially occupies the space ordinarily filled by air. This affection, therefore, which almost always occurs in the most dependent parts of the lung, is also accompanied with a moderate amount of dulness on percussion.

In *Malignant disease of the lung*, the pulmonary tissue is pushed aside, and the air is entirely excluded from the parts affected by the cancerous or fungous growth. When, therefore, the disease occupies a large space, or invades a considerable amount of pulmonary tissue in the massive form, there exists perfect dulness on percussion, and great resistance to the finger.

Malignant disease, however, like scrofulous tubercles, lobular pneumonia, and pulmonary apoplexy, may exist in small rounded disseminated masses, surrounded by healthy pulmonary tissue, to a comparatively large extent. Under such circumstances, the healthy portion of the lung surrounding the malignant deposit, as in the other diseases specified, conceals the dulness which would be otherwise produced by it. When, therefore, malignant disease attacks the lung in a massive form, as it usually does, when originally developed therein, it is accompanied

by dulness on percussion. When, on the contrary, it is distributed through the lung in the form of variously sized tubercles, as it ordinarily is when arising secondarily to its appearance in other parts, it is often not accompanied by any alteration of the resonance of the chest.

Emphysema of the lung.—The diseases hitherto noticed have been accompanied with a decrease of the natural resonance of the chest: Emphysema of the lung produces exactly the opposite condition. This affection, as has been previously explained, is dependent upon the dilatation, or rupture of the air-cells, or of atrophy, and consequent yielding of the partitions between them. In any case, and in all cases, whether the affection be merely vesicular, interlobular, or sub-pleural, the proportion of air to the solid and fluid constituents of the pulmonary organs is increased. The lung, either locally or generally, is increased in volume. The dimensions of the parietes are enlarged, and the resonance upon percussion is augmented. Not only are the parts ordinarily resonant rendered more so by this affection, but those which, in the natural condition, are dull, become well-sounding. Thus, in consequence of the enlarged lung extending over the heart, the præcordial region often becomes perfectly resonant on percussion, excepting towards the xiphoid cartilage, where, from the increased

size and distension of the right side of the heart, the dulness is even greater and more extensive than natural. Thus, also, in the parts which are ordinarily dull from the underlying liver, the sound on percussion becomes comparatively resonant.

2. *Indications afforded by Percussion in diseases of the Pleura.*

When fluid of any kind is effused into the pleura, the lung, to a corresponding extent, is compressed, and the air, in nearly an equal proportion, is squeezed out of, and prevented regaining admission to, the pulmonary cells. Whether, therefore, the disease arise from inflammation or from obstruction, whether the fluid be blood, pure serum, turbid serum, or pus, dulness on percussion exists to an extent, and, up to certain limits, to a degree also, commensurate with the amount of fluid. The dulness in each case may primarily affect only the lower part of the serous cavity, and gradually extend upwards as the fluid increases, and, by its increment, displace the lung. It may, on the contrary, in each case only extend over a defined space, to which it is confined by previously existing pleuritic adhesions.

When the pleura is free from such adhesions, the fluid, from whatever part of the membrane it proceeds, may in each case gravitate to the lowest part of the cavity, and its site may be

changed according to the varying position of the patient's body; in each case, therefore, the part in which the dulness is observed may also vary together with the changes of position. This change in the situation of the fluid, and of the consequent dulness, according to the position of the body, is, however, far more common in hydrothorax, than in either simple pleuritic effusion or empyema, in which diseases the fluid is much more frequently confined to a defined space by surrounding adhesions. Though, therefore, the change of the situation of the dulness from position may serve to distinguish that arising from hydrothorax, from that produced by pneumonia, it cannot be trusted as necessarily, or even commonly, diagnostic between pneumonia and empyema.

Even when pleuritic fluid has been removed by the use of medicine, or by the employment of paracentesis, dulness on percussion, though in a less degree, often remains for a considerable time. This probably arises from two causes, viz. the imperfect expansion of the layer of lung formerly in contact with the inflammatory effusion; and the presence of a coating of solid lymph left upon the surface of the serous membrane. When the fluid of empyema is absorbed, or is evacuated either by the trochar or by a spontaneous opening, the lung is generally so thickly covered with solid matter, so firmly bound and plastered down by bands and adhesions, as to be almost incapable of

expansion. The lung of the opposite side becomes enlarged, and partially fills up the space. But it does so partially only ; its supplementary action is not complete. Hence the ribs of the diseased side fall in upon the space formerly occupied by the fluid, the chest becomes misshapen, and the spine more or less distorted. Hence, also, though not a single drop of fluid may be present in the pleura, though the serous cavity may be entirely closed by adhesions, dulness on percussion may remain as a consequence of empyema long after the disease itself is cured. Natural resonance may never return.

In *Malignant disease* of the pleura, as in that of the lung, the pulmonary tissue is pushed aside, and dulness exists on percussion commensurately with the extent of the solid deposit.

Tubercles, when confined to the pleura, rarely exist to so great an extent as to give rise to dulness on percussion.

In *Pneumothorax*, the air in the pleura pushes aside and compresses the lung to an extent varying according to the amount of the disease existing in the lung itself, and of the adhesions connecting it with the costal pleura. The part of the chest into which the air has escaped is generally more resonant than usual, and not unfrequently tympanitic upon percussion. When air and fluid co-exist in the same serous sac, dulness is present at one part, and tympanitic

resonance at another. The two conditions vary in extent according to the respective amount of the gas and the fluid. Under these circumstances the fluid almost always occupies the lowest part of the chest, and gravitates, according to the position of the patient, to the space which is the most dependent. The posterior will therefore be dull, and the anterior part of the chest resonant on percussion, while the patient lies upon his back, while exactly the converse state of things will be observed, when, if able to do so, he turns upon his face.

Is, then, tympanitic resonance upon percussion always present in pneumothorax?—It certainly is not. The air may be in quantity so small, the pleura so thickly lined with solid matter, or the pneumothorax may be so centrally situated at the base of the lung, in contact with the diaphragm, and therefore so far removed from the vibrating parietes, and the intervening substances may be such indifferent conductors of the vibrations produced therein, that no increase of resonance may exist upon percussion. The side may, indeed, be even more dull than natural; though this is of rare occurrence. The absence of tympanitic resonance in pneumothorax is more frequent in those cases in which air enters the pleura in consequence of the partial evacuation of an empyema through a bronchial tube, than in those which result from the communication of a

tubercular cavity with the serous membrane. In the former cases the lung has usually been already compressed to the utmost, and air only enters the pleura as the purulent fluid is expectorated. A portion of the space formerly occupied by the purulent effusion is also filled up by the return of the distended ribs and detrudded diaphragm to their accustomed boundaries. The air in the pleura, therefore, is sometimes in quantity so small as not to give rise to resonance on percussion. The same relative state of parts sometimes exists in pneumothorax arising from other causes, when there is a gradual accumulation of pleuritic effusion, and the comparatively heavy liquid excludes the light air from the cavity of the serous membrane. This condition is far from unfrequent in the last stages of pneumothorax connected with tubercular phthisis.

3. Indications afforded by Percussion in diseases of the Heart.

The extent of dulness of the præcordial region varies considerably in different individuals, quite independently of disease. This difference arises, perhaps, in part from congenital formation, and in part from the effects of dress. The normal præcordial dulness cannot, therefore, be defined by any precise boundaries. It may be stated, as an approach to the average of its ordinary extent, that it occupies the entire inferior sternal region, and that portion of the left mammary region

which is covered by the cartilages of the fifth, sixth, and seventh ribs. If the præcordial dulness much exceed these limits, the excess may usually be correctly attributed to disease; though it by no means necessarily follows that that disease is connected with the heart, or with its investing membrane. It may arise from local pleurisy, or consolidation of the lung. The præcordial dulness may, on the other hand, not extend to its ordinary boundaries without disease being present in any organ. It may be almost entirely cancelled by an overlying emphysematous lung, without any cause for the diminished dulness existing in the heart. Still further, though great enlargement of the heart may be present, it may happen that, excepting in the scrobiculus cordis, no dulness on percussion may be perceptible, in consequence of the emphysematous lung intervening between the hypertrophied, or dilated organ and the parietes.

These circumstances being duly considered, and the exceptions borne in mind, it may be stated generally that all enlargements of the heart, and all effusions into the pericardium, are accompanied with increased dulness on percussion of the præcordial region. The cause of this must, from what has been previously stated, be sufficiently obvious, and need not be here explained. When, therefore, from any obstruction to the course of the blood existing in the aortic valves, or in the course of the thoracic

aorta itself, or from any other cause, the left ventricle is greatly dilated and thickened, an increase in the extent of the præcordial dulness is observed, especially over the costal cartilages, and below the left nipple.

When obstruction exists in the mitral valve, when the auriculo-ventricular opening is contracted, or admits of regurgitation, the left auricle is imperfectly emptied during the diastole, and is more than ordinarily distended during the systole, of the ventricle. The auricle at length becomes permanently dilated and thickened. This may be often ascertained by an extension of the præcordial dulness to the cartilages, and adjoining bone of the left fourth rib, the third intercostal space, and even the third rib. When the lungs are extensively affected with emphysema, when, in consequence of disease in the mitral valve, the circulation of the blood through the lungs is obstructed, or when, from congenital or acquired smallness of the calibre of, or from any other disease in, the pulmonary artery, there exists an impediment to the free transmission of the blood from the right ventricle, repletion and distension of that cavity, and of its adjoining auricle, necessarily ensues. The evidence of this exists in the extension of the ordinary præcordial dulness to the right side of the sternum, or the inner edge of right mammary region, and particularly to the scrobiculus cordis.

In Pericardial effusion.—From the difficulty,

or rather the impossibility, of accurately defining the natural limits of the præcordial dulness in individual cases, the small increase which may be presumed to be dependent on the plastic effusion of recent pericarditis is not discoverable by percussion. The presence even of fluid effusion cannot always be ascertained by percussion, unless it exist in considerable quantity.

When, however, the quantity of fluid is large, the dulness on percussion is perceptibly increased. If the serous membrane be free from adhesions, the increased dulness commences below, extends on either side of the bottom of the sternum, and gradually passes upwards.

When the membrane is largely distended with fluid, the increased dulness is sometimes very remarkable and extensive. Its boundaries are enlarged in every direction, but especially transversely.

When partial adhesions exist in the membrane, and prevent its distension in certain directions, the fluid effused in the free portions of the serous sac causes the outline of the dulness to be irregular in form.

Hence it happens that pericardial effusions may exist in considerable amount without being distinguishable by any increased dulness; or, in other words, the dulness may so slightly extend beyond the limits naturally existing in some healthy individuals, as not to be certainly appreciated as the result of disease. In other cases the dulness may

be general and extensive, and in a third set of cases it may be limited and local, though at the same time definite and decided.

4. Indications afforded by Percussion in diseases of the Anterior and Posterior Mediastinum.

When aneurism proceeds from that portion of the aorta which is just above the valves, the dulness to which it gives rise may be undistinguishable from that proceeding from disease of the heart itself. It may extend upon either side of the heart, though it most commonly extends to the right side. Independently of the dulness arising from the tumor itself, which varies of course according to the size of the sac, the disease often displaces the heart so as considerably to modify the normal dulness of the præcordial region.

When aneurism exists in the ascending aorta, and is of large size, its presence may generally be recognized by an alteration of the resonance on percussion over the upper sternal, and the inner edge of the right subclavian region. Unless, however, the dimensions of the aneurism are considerable, no absolute dulness is perceptible, in consequence of the overlying resonant lung concealing that which the presence of the tumour would otherwise induce.

The same may be said in reference to aneurism of the arteria innominata. But even when no dulness, strictly so called, can be discovered, there

may be often distinguished a variation of the tone or quality of the sound when corresponding regions of the two sides are percussed, and accurately compared. This variation is not to be disregarded; in doubtful cases it may be of very great importance.

When aneurism exists in the arch of the aorta, it is more likely to be recognized by dulness on percussion than when the ascending portion of the vessel is the part affected. This arises from the lung towards the central line being thinner, and more easily displaced, or entirely pushed aside, than in the parts previously referred to.

Aneurism of the descending aorta, if of large size, may be attended with dulness on percussion on either side of the spine, though it more commonly extends to the left side. It may project directly forward, and, by pushing aside the lungs, be the cause of increased dulness in the præcordial region.

Abscesses of the anterior mediastinum are accompanied with dulness on percussion of the particular part which they affect; and enlargement of the bronchial glands *are said* to be recognizable by dulness of the interscapular regions. This latter fact I have not yet been able to verify, and I cannot but think that the enlargement must be very considerable, to produce appreciable dulness on the interscapular region.

The diseases, then, in which Percussion may

afford useful information, are Bronchitis, Pneumonia, Phthisis, Emphysema, and Malignant Disease of the Lung; Hydrothorax, Pleuritic effusion, and Pneumothorax; Dilatation of the heart, Pericardial effusion, Aneurism of the large vessels, and Abscesses or other diseases of the Mediastina.

CHAPTER V.

AUSCULTATION, OR EXAMINATION BY THE EAR.

GREAT as have been the advantages which have accrued from the modes of physical investigation which have been already, and which are still to be, noticed, it is to Auscultation more especially, combined with the study of morbid anatomy, that is attributable the vast progress which has been made during the present century in discriminating diseases of the chest. It is to Auscultation that the profession is mainly indebted for that measure of light which has supervened upon comparative obscurity.

By Auscultation, as regards the pathology of the chest, is meant the practice of listening to the sounds arising from the exercise of the functions of respiration and circulation, and of comparing those emanating from diseased organs with those ascertained, by previous experience, to exist in the healthy condition. To this may be properly added the explanation of the difference of the sounds occurring in disease, by the physical condition of the organs produced thereby.

Auscultation, like percussion, may be immediate or mediate. It is immediate, when the unguarded ear is placed upon the chest, either bare or protected with only a slight covering; and when, therefore, the vibrations of the elastic parietes are communicated directly to the organ of hearing. It is mediate, when between the ear and parietes is interposed a body, which conducts the vibrations of one to the other. The opinions originally entertained of the great advantages of the stethoscope are now acknowledged to have been erroneous. Some auscultators prefer mediate, others immediate auscultation. Each mode possesses some advantages, and each is liable, in certain circumstances, to some objections.

1. *Immediate Auscultation.*

The ear, when this mode is employed, is successively applied to, and firmly pressed upon, the different regions of the chest, either quite bare or only slightly covered; in such a way that the organ may, as much as possible, become one with, and therefore participate in the vibrations of, the solid parietes upon which it is placed. This is in many respects a desirable mode of auscultating. It requires, or appears to require, less abstraction; it is more easily acquired; it causes no pain to the patient by the pressure employed, which, in some instances, almost neces-

sarily arises from the use of the stethoscope; and it is, usually at least, as effective as mediate auscultation, when large surfaces are implicated by disease. Not unfrequently, indeed, sounds which are indistinct, and appear indefinite through the cylinder, become sufficiently obvious and determinate upon the application of the ear.

But immediate auscultation is open to several objections. The rustling of the clothes of the patient, in consequence of a larger surface being covered by the side of the head than by the end of the stethoscope, is more likely to interfere with the sound proceeding from the chest. The hair of the auscultator himself, by the motion of the patient's body, frequently gives rise to sounds nearly resembling those arising from disease. These may be incorrectly supposed to proceed from the chest, and may thus become a source of error.

It is sometimes important, as an aid to diagnosis, to be able to define the exact situation in which a particular sound is heard with the greatest intensity or distinctness; as in affections of the valves of the heart, the aorta, &c. By the unguarded ear, we cannot effect this so perfectly, we certainly do not possess the power of localizing particular sounds so accurately, as by the aid of the stethoscope. By some persons it may be considered indelicate to apply the naked ear, not merely to the bare chest, but even to the unrobed,

though slightly covered, parietes of female patients. In other persons, motives of a different nature will incline the auscultator himself to avoid contact, or a too near approach to the integuments of his patient. In the one case, modesty, or prudery, may forbid the employment of immediate auscultation; in the other, uncleanness may render it undesirable. Independently, however, of the real or presumed indelicacy of applying the ear in the case of female patients, it is often quite impracticable to do so, with good effect, in persons with largely developed mammary glands.

There are also, it must be recollected, some situations in which the ear cannot be fairly and efficiently employed; as in the space above the clavicle, in the upper part of the axilla, and in very thin persons, immediately below the clavicle, and in the supra-spinous fossa of the scapula.

Immediate auscultation, however, notwithstanding what has been said, may be employed with the best success in many cases, and, in most cases, may be usefully applied for the confirmation of opinions derived from other sources, or the verification of sounds indistinctly heard by means of the stethoscope.

Cautions respecting immediate Auscultation.

These have been, for the most part, already referred to in the preceding observations; but it

may be desirable that they should be more distinctly and separately stated.

First, then, it is necessary that, in the case of female patients, the appearance of the slightest approach to indelicacy should be avoided. If, therefore, for the purpose of confirming a doubtful opinion, or for any other reason, it be thought desirable to apply the ear itself to the anterior part of the chest, the skin should be covered with some soft yielding material, either appertaining to the patient's dress, or obtained from the attendants. A soft towel, an old dinner napkin, or even a silk handkerchief, may answer the purpose perfectly well; though cotton should generally be preferred to articles of linen, wool, or silk, as these not unfrequently give rise to some rustling or crackling sound, when tumbled or pressed, which may interfere, or be confounded with, those emanating from within the chest.

In this mode of auscultation, it is most assuredly the solid parts of the ear, and the bones of the head, which transmit the vibrations of the solid parietes of the chest to the auditory nerve. The protection referred to as a covering is, therefore, the more necessary; because, as has been before stated, the entire ear should be evenly and firmly pressed upon the parietes of the patient, so as to cause them to become, as much as possible, one vibrating body. The napkin or towel, employed as a covering to the chest, should be

smoothly spread, and tightly drawn, over the surface, and all other garments should be so far removed, as not to come in contact with the auscultator's ear or head; extraneous sounds will otherwise assuredly arise, which may not improbably mislead him.

The examiner should carefully put aside his hair from around his ear, and, as far as may be, from the side of the head, in contact with the chest; since if any portion intervene between the ear, and the chest, or come in loose contact with the patient's dress, a noise will be produced, which may be easily mistaken for sounds proceeding from within.

It can be scarcely necessary to state, that in the case of uncleanly individuals, male or female, some protection should be used, lest the examiner should derive from his patient more than he would desire to retain.

From what has been already mentioned, it may be scarcely requisite to remind the auscultator, that when the chest of male patients is thickly clothed with hair, the hair must be removed before immediate auscultation can be satisfactorily performed. In females with large pendulous mammæ, the gland must be drawn aside, and the skin rendered as tight and firm as it can be, without serious inconvenience to the patient; her own hand, where it can be employed, is the best instrument for the purpose.

2. *Mediate Auscultation.*

The principles of mediate auscultation appear not to be always understood, even by many who constantly employ it. The stethoscope, of whatever form, or however composed, is simply a conductor between the parietes of the chest and the ear. The vibrations of those parietes are caused, or are transmitted, by certain sonorous or resonating substances within the cavity. Whether these vibrations originate in air, or in blood,—whether they are conveyed by solid or semi-solid bodies, by fluid or by gas, they all ultimately reach the solid walls of the chest. It is by the vibrations of those solid walls that they ultimately reach the surface; and by the vibrations of those solid walls that they are all conducted to the ear. The stethoscope is merely a conductor. It may, indeed, as in the flexible stethoscope, conduct by the air contained in the hollow of the tube alone; it may, as in the ordinary stethoscope supplied with a nipple-like appendage for the external meatus, conduct both by air within the bore of the cylinder, and by its solid walls,—or it may conduct by the solid walls alone.

What form of stethoscope, then, is the best for general use? To this it may be generally replied,—that, by which each individual can hear best. In this respect fancy and prejudice may effect something; habit much more. But to answer the

question satisfactorily, it should be, in the first place, recollected that solids are better conductors of sounds than either fluids or gases; and secondly, that the vibrations of sound are considerably modified, and decreased in intensity, by transmission from one medium to another; that however originating, and by whatever elastic substance they may be transmitted, the waves constituting sound are greatly altered, and, by partial interruption, are decreased in their extent, upon their entrance into a medium of a different kind. The waves of sound, therefore, become less powerful by being transmitted from solids to gases, or to fluids, as well as from either these to solids. As, therefore, solids are in themselves better conductors than either fluids or gases; and as the solid parietes of the chest (confining attention to that cavity) are the medium, by the vibrations of which the sounds of the thoracic organs reach the surface; it appears sufficiently evident that, upon theoretical grounds at least, the solid stethoscope is the best that can be employed.

I believe that, generally speaking, what is rendered probable upon theoretical grounds will be verified by experience; and that in the vast majority of cases the solid stethoscope will be found the most useful instrument in practice. I call the ordinary stethoscope solid, because, from long experience, and from numerous and

various experiments, it appears to me quite evident that it is by the solid walls of the stethoscope that sound is, in all cases principally, and in many cases entirely, conducted.

It is not necessary, nor do I think it at all incumbent upon me, to point out the advantages, or what I conceive to be the disadvantages, of any particular form of stethoscope; but it may be desirable that I should mention that particular form which long practice, as well as theory, has induced me to regard as the best.

Though, then, each person had better make use of that stethoscope which, by experience, he finds best suited to his ear, or by which, from habit, he can hear best; yet I may observe, that as from repeated trials I have long convinced myself that, in the wooden instrument, the sound is principally conveyed through the wood itself, that, in fact, the solid is the chief, and, in a great majority of instances, the sole conductor, the more simply the wooden instrument is constructed, the better will be its conducting power. I do not intend hereby to deny that with some persons, particularly those who employ a nipple to the ear-piece of their stethoscope, some vibrations of sound are conducted by the air in the bore of the cylinder, directly to the *membrana tympani*; but I believe that in all cases the greatest effect, by far, is communicated by the wood itself to the solid parts of the ear, and from them to the bones of the head; and

I am convinced that this is the sole mode by which many, and myself in particular, appreciate sounds in auscultation.

If the external meatus be closed with lint or chewed paper, the sounds are almost equally audible. If a perfectly solid cylinder be employed, or the bore of a hollow instrument be stuffed with any soft substance, the effect is scarcely perceptibly decreased. That it is at all decreased, in however trifling a degree, is dependent, I imagine, in the one instance, upon the mass of solid being greater, and therefore requiring more force to produce an effect between its molecules, equivalent to that created in a mass of less extent; and in the other instance, upon the fact that the substance introduced into the hollow of the cylinder, or of the meatus, interferes, in some slight degree, with the vibrations of the solids themselves.

What, then, it may be asked, is the professed object, and in what consists the real utility, of the bore of the solid stethoscope? The professed object is, that the vibrations of sound may be transmitted through the air in the tube. Its real utility depends upon its making the instrument lighter, upon its probably facilitating the vibrations of the solid, and, in some cases, particularly in those instruments in which a nipple is appended to the ear-piece, upon its contributing to the effect of such vibrations by the addition of

those which are communicated by the air occupying the hollow space.

The distinctions sometimes made between the sounds of the breathing, of the voice, and of the heart, may, I think, be disregarded in deciding upon the form of the stethoscope to be employed. Each and all of these sounds are communicated to the surface through the solid parietes, and by a solid they are all best conducted to the ear. At such conclusions we should arrive from purely theoretical considerations, and though, as regards auscultation, experiment and theory do not always appear exactly to coincide; though some rare circumstances it seems to be difficult to explain upon common acoustic principles, yet in this case, as regards myself at least, practical experience exactly tallies with the theory.

The stethoscope, then, which appears to be most convenient, and advantageous for general use, is made of one material, without joints, or additions of any kind. The material, (cedar) of which it is usually made, is probably as good as any other. It should be from four to six inches long, with a clean, polished quarter-inch bore. The shape may be according to the student's fancy; but it is desirable, for the sake of lightness, to have it somewhat hollowed in the stem. The rim of the bell-shaped or thoracic extremity should be broad, and well rounded; so that it may be firmly pressed upon the parietes without causing pain, or much

inconvenience, to the person examined. The ear-piece should be broad, so as to embrace the entire ear, and nearly flat, or so little hollowed as not to admit the presence of air between the wood and the ear. The wood itself should be rather dense, and the instrument stout in every part, that it may not be chipped or broken by every accidental fall.

Such is the stethoscope which, after trial of several kinds, appears to me to be the best suited for practical purposes, and which I have for many years employed. A sketch of an exceedingly good instrument, now in my possession, may perhaps be here introduced with advantage, lest the directions above given should not be suffi-



ciently explicit. It has been reduced one-third.

It may, however, be acknowledged that an auscultator can often hear better with a bad instrument, to which he has been long accustomed, than with a stethoscope of superior form and construction, with the use of which habit has not rendered him familiar. It may be indeed repeated, that the form of the instrument is of very trivial importance, and that the best stethoscope for any individual is clearly that with which he can hear best.

There are, indeed, some cases in which the flexible stethoscope may be employed with advantage, even by those who are accustomed to use the wooden cylinder. But they are cases only in which it is inconvenient or impracticable to apply the solid tube; those, namely, in which it is important to examine the posterior parts of the chest, but in which it would be unsafe or imprudent to raise the patient, and he is himself unable to turn for the purpose. In such cases, the infundibular end of the instrument may be passed under the body, without seriously disturbing the invalid, and important information may be occasionally acquired thereby.

3. Mode of using the Stethoscope.

The instrument, grasped in the centre by the two fingers and thumb of the right hand, as a pen,

should be first flatly placed, and firmly pressed, upon the parietes of the chest. The cylinder must always be at right angles with the surface to which it is applied, and the head of the examiner must be always bent to the extremity of the instrument, and the instrument not made to incline to his head. Thus, if the patient be upon his back, and the front of the chest be the part under examination, the stethoscope must be perpendicular, or nearly so; if the space between the front and the sides of the chest be submitted to investigation, it must be placed obliquely; and if the side of the chest be the region explored, it must project horizontally therefrom, and the head of the examiner must be depressed, or elevated accordingly. The ear should be evenly adjusted upon the broad, nearly flat end of the instrument, and the pressure of the head be so firm as to make the parietes of the chest, the conducting medium, the stethoscope, and the ear, one connected vibrating body. This cannot be effected without considerable pressure, which, however, should not be so great as to cause pain to the patient. To prevent this, not unfrequent, result of the application of the stethoscope, the edge of the funnel-shaped extremity of the instrument should, as before stated, be broad, and smoothly rounded off. The contact of the fingers with the instrument is then to be removed, and observations made; first, as regards

the breathing, and then voice, and the cough, as will be further explained in a future page.

The undivided attention of the examiner must be directed to the sounds passing through the tube; his senses should be dead to all besides. He must hear nothing with the other ear; the abstraction must be perfect. Generally speaking this capability of abstraction is to be attained only by habit; but if noises in the chamber, the ward, or the adjoining room, or street, interfere with the examiner's power of discriminating the sounds proceeding from the chest, he may place a finger of the disengaged hand over the meatus of the unemployed ear, and thus exclude all other sounds. He may by the very act contribute to the abstracted attention, which, in many cases, is absolutely requisite to arrive at satisfactory conclusions. It has been said that the stethoscope should be grasped by the fingers of the right hand, when, of course, the right ear would be employed; but it will conduce greatly to the student's aptitude in exploration to be able to employ both hands and both ears with equal facility.

4. *Cautions respecting mediate Auscultation.*

It may be well to repeat, in the form of cautions, some particulars which have been already mentioned in the preceding directions. Let, then,

the examiner be especially careful that the entire circle of the thoracic end of the stethoscope come in close contact with the parietes. If, from unequal pressure, improper position of the patient, or oblique direction of the instrument, the contact be not entire, the vibrations of the walls of the chest will be very imperfectly transmitted.

In some cases, as in persons naturally very thin, or greatly reduced by exhausting diseases, the depressions existing between the ribs in some regions of the chest, and those frequently the most necessary to be minutely examined, are so great as to render it impossible that the whole circle of the stethoscope can come in contact with the skin. The intercostal spaces are left untouched, while the skin over the ribs is submitted to severe pressure. Under such, and all similar circumstances, little, or nothing, can be heard through the stethoscope.

Two expedients may be then adopted: the examiner may either make a pad of soft linen, or lint, by which the depressions between the ribs may be filled up, and the skin over the ribs protected; or he may adopt the other, and usually the preferable expedient, of discarding the stethoscope altogether, and using the unprotected ear. The latter plan is the more desirable of the two, as the pad necessarily in some measure interferes with the facility with which the sounds are transmitted to the ear. The naked ear should

therefore be in such cases adopted, excepting in those parts of the chest to which it cannot be fairly applied.

The stethoscope must be carefully prevented coming in contact with the patient's garments, with the bed-clothes, with his extremities, or with any other part of his body than the chest itself. This caution is even more necessary in mediate than in immediate auscultation. The slightest friction of the garments, or the hand, is so perfectly conducted by the instrument, as most materially to interfere with the distinctness of, if it does not overpower, the sounds proceeding from the chest.

In very stout persons the pressure of the instrument must be increased, so that the adipose tissue may be compressed, and, as far as possible, rendered identical, as regards conducting power, with the bony parietes which it covers, and the free vibrations of which it in some measure prevents. In persons whose integuments are œdematous, the pressure of the instrument must be equally great, or even greater, than in fat persons; so that the serum may be entirely squeezed out of the cells immediately below the stethoscope. Loose fat and diffused serum are bad conductors of sound, when compared with solids. Hence the importance of the caution just delivered.

In reference to œdema, however, another cau-

tion is required. This arises from the frequently observed fact, that the escape of serum from the cellular membrane, by the pressure of the instrument itself, gives rise to a sound very similar to one frequently proceeding from within the chest. The sound referred to is *pleuritic rubbing*, one form of which, possessing a jerking, or interrupted character, the noise produced by the escape of the serum from the cells under the pressure of the stethoscope, so exactly resembles, that if the patient continued to breathe during the whole of the examination, the two sounds would be distinguished with the greatest difficulty, if, indeed, they could be at all distinguished from each other. An obvious distinction of course exists in the fact that the rubbing of the pleura is only present during respiration; while the œdematous creak, so long as any serum remains to be squeezed out of the cells, persists though the breathing be temporarily suspended. The sound so precisely represents certain forms of the rubbing existing in disease of the pleura, that I have, in the presence of pupils, occasionally illustrated the latter sound by pressing the stethoscope upon an œdematous thigh.

If, indeed, the œdema be soft, and the integuments flabby, the peculiar sound to which I have referred is not always produced by pressure, or it is of a much less sharp and decided character than when the swelling is firm and tense. On

these two accounts, then, first, because the infiltration of the integuments obstructs the sounds proceeding from the chest ; and, secondly, because on pressure it gives rise to another sound, which may be easily confounded with one of intrathoracic origin ; the serum of œdema should be carefully pressed out of the cellular membrane before auscultation is attempted.

5. Combined use of mediate, and immediate Auscultation.

From what has been already said, it will be evident that, on some occasions, and in some parts the ear, and on others the stethoscope, may be most advantageously employed. As, therefore, it will conduce to the efficiency of his investigations, it is desirable that the student should be practically conversant with both modes of auscultating. Let him on no account neglect the use of the stethoscope, because he finds that, under ordinary circumstances, he can hear better with the unguarded ear ; as in practice cases will constantly occur to him in which he will be compelled to employ the cylinder, or to leave certain parts of the chest unexamined.

The plan of proceeding which will perhaps be found on the whole most useful, is in each case to practise both mediate and immediate auscultation. Thus, on the anterior part of the chest,

and particularly in the acromial, the axillary, and the infra-clavicular regions, the stethoscope may be used; while in the lateral regions, and all the posterior surface of the chest, excepting the space above the spine of the scapula, the ear will be found at least as effective, when unguarded, as when supplied with the solid conductor. If the patient be very fat, or the integuments be infiltrated with serum, mediate is preferable to immediate auscultation; as with the stethoscope more direct and local pressure can be made than with the ear, which, though not so far removed, in such cases comes not into such near acoustic relation with the solid parietes, as when the solid conductor intervenes between them.

In all examinations of the larynx and trachea, it is almost always necessary to employ the stethoscope. In all affections of the heart, excepting when great tenderness exists upon pressure, the stethoscope is to be preferred to the ear; while in the auscultation supplementary to succussion, to be afterwards noticed, and in the superficial examinations made through stays and petticoats, the ear, if it be evenly placed, and the patient's body be firmly pressed against it by the the hand placed upon the opposite side, will be ordinarily a more effective auscultatory instrument than the cylinder.

6. *Auscultation of the Organs of Respiration.*

It is absolutely necessary, as has been before

stated, that the student be perfectly familiar with the sounds naturally existing in the chest, previously to his becoming acquainted with, or being capable of judging of, those which are abnormal; just as he must be conversant with the healthy appearance of organs, before he can fairly expect to become a morbid anatomist. The natural sounds of the respiration and the voice, therefore, will be first noticed, and subsequently those which exist in disease.

1. *Of the Natural Respiration.*

To enable the examiner to judge correctly of the healthy, or morbid condition of the respiration, ordinary inspiration is scarcely sufficient. It is desirable that the air should be more generally diffused through, and should more perfectly inflate, the pulmonary cells, than it does in common inspiration. In nice examinations, therefore, the patient should be desired to fill his chest by a succession of forced inspirations. The examiner will be thereby enabled to make his observations upon the sound, and duration of the expiration, as well as upon those of inspiration; the former of which, in ordinary breathing, is sometimes scarcely audible, and differs from the latter in some other important particulars, which will be afterwards referred to. To prevent con-

fusion it may be well to confine attention, in the first instance, to the inspiration*.

If the ear be applied to the stethoscope, evenly placed upon the larynx, or trachea, and gentle pressure be made, the sound, caused by the ingress and egress of air during the act of respiration, will be observed to be hoarse and hollow; like that produced by blowing, with the lips considerably compressed, through a pipe of large calibre, as through the bore of the stethoscope;

* Simple as the process may appear, it is by no means an easy matter to make some persons take a really deep inspiration, in such a way as to enable the auscultator to judge of the condition of the lung. Their awkwardness is sometimes so great as really to have the appearance of design. First they raise the shoulders, without inspiring at all; then they inspire deeply; then they draw in the air through the compressed lips, or relaxed fauces, in such a manner as to give rise to noises, which overpower the sounds originating within the chest; afterwards they protrude the abdominal muscles, and then perhaps go through the entire series of awkward efforts again. To tell them to take a deep breath, to heave a sigh, to fill the chest, to draw in the air, to raise the ribs while they breathe, and even to shew them how to do so, are of no avail. The more you talk, and the more they appear to try to effect the object you desire, the farther do they remove from that object; the more stupid in this respect do they seem to be. But they often involuntarily do exactly that which, after many and diversified attempts, they have been unable to effect. Thus, if told to cough, and to continue coughing for a short time, they will, from the mere exhaustion caused by the repeated short expirations, which constitute that act, spontaneously take such a soft, quiet, yet deep inspiration, as all previous efforts had failed in causing them to accomplish.

or that which is sometimes produced by a draught of air through the key-hole of a door. It is called *tracheal* respiration. When a similar sound is heard, as the result of disease, in other parts of the chest, the breathing is said to be *tracheal*.

If, instead of being placed upon the neck, the stethoscope be now put upon the upper bone of the sternum, below the sterno-clavicular articulation, or between the scapulæ upon either side, and the individual be desired to breathe deeply; a sound very much like, but not exactly similar to, the tracheal respiration will be heard by the auscultator. The sound may be generally, but, it must be acknowledged, by no means universally, observed in the state of health in the situations indicated. It is sometimes absent in all the localities, but sometimes, when inaudible on the front of the chest, may still be heard close to the spine, at a level with the centre of the scapulæ. It is not so loud, or so hollow, as the tracheal sound. It may be represented as its diminutive. It resembles the noise occasioned by blowing through a small reed, or a quill of moderate size. It arises from the rush of air through the larger bronchial tubes, and, wherever heard, is therefore called *bronchial respiration*, or *tubular breathing*. If discovered in other situations than the upper part of the sternum, the inner side of the infraclavicular, or the interscapular regions, it may

with tolerable certainty be regarded as morbid ; though it may be occasionally heard less distinctly than in those regions in the axillary spaces ; particularly that of the right side.

It may be here observed, that the natural breathing not only presents varieties, as to tone and distinctness, in different individuals ; but also that similar regions, upon opposite sides of the body, may vary in these respects, as in some others, quite independently of disease.

When the stethoscope is placed upon most other parts of the chest, than those previously specified, a sound is heard by the auscultator during inspiration very different from those which have been mentioned before. It is perhaps difficult to define its character correctly. It is soft, scarcely sonorous, equable, breezy, and, during inspiration, continuous. It may be pretty accurately copied by gently drawing the air through the partially compressed lips ; it may be compared to the song of a gentle gale in the thick summer foliage ; or to the whisper of the retiring wave upon a sandy coast. It is called the *respiratory* or *vesicular murmur*, on account of its being the characteristic of healthy respiration, and of its being clearly connected with the vesicular tissue of the lung. It probably depends upon the admission of the air into the irregular cavity, and its passage over the uneven surface, of the pulmonary vesicles.

It varies considerably in intensity in different regions of the chest. It is most distinct in the acromial, the central and lower part of the superior sternal, the infra-clavicular, the axillary, and the subscapular regions. It is less distinct in the lateral, the right mammary, the scapular, and the interscapular regions; still less in the hypochondriac; and least of all in the inferior sternal and the inner part of the left mammary region.

Independently of the variation of the intensity of the sound in the different regions of the chest, whether the variation arise from the position of the organs, the amount of pulmonary tissue beneath the ear, or the facility or difficulty with which the inspired air reaches the pulmonary cells, the two sides of the chest frequently vary a little in respect to the loudness of the respiratory murmur. Thus it is, usually, rather louder in the acromial, scapular, and infra-clavicular regions of the right side, than in the corresponding regions of the left side; but in so slight a degree, as to be scarcely worthy of consideration, in a practical point of view.

The respiratory murmur may, both locally and generally, be more loud, or less distinct than natural, in persons who are quite free from any appreciable disease. It may also be harsh, or rough, scarcely audible, or altogether absent.

Thus, in childhood, and in youth, the respiratory murmur is more loud than in adult life, and especially than in old age. This probably arises

from the more dense tissue, and the greater functional activity of the lungs in the young than in persons advanced in life. From this circumstance, it is called, whether existing in youth or in age, *puerile respiration*.

But as it is frequently observed to exist in the healthy portion of a lung, or in an entire lung, when the remaining portion in the one case, or the opposite lung in the other, is greatly embarrassed, and prevented performing its accustomed function; and as this healthy portion, or this sound lung, seems under such circumstances to have its functional activity increased, and thus to supply the deficiency of the diseased parts, it is sometimes called, not inaptly, *supplementary respiration*. It is always heard when the healthy respiration is more than ordinarily active, as in persons "out of breath," as it is called, from strong exertion, as running, dancing, &c., or after the respiration has been voluntarily suspended for a time, and the individual breathes quickly to restore the normal balance of the circulation through the pulmonary organs.

The respiratory murmur of some persons appears naturally to be harsh or rough, or to be so at least without any other evidence of pulmonary disease than is afforded by such harshness or roughness. The cause is not evident, though, from what is observed in disease, it seems not improbable that it is dependent upon some

congenital or acquired peculiarities in the smaller divisions of the bronchial tubes, rather than upon the vesicular tissue of the lung itself. It may be either local, or general. Its more common seat, when local, is the axillary space, in which it is far from unfrequent. If it be observed in other situations, and, at the same time, not heard over the whole of those regions in which the pure respiratory murmur is commonly heard most distinctly, it must be regarded with some suspicion.

Harsh or rough breathing is, to the educated ear, sufficiently distinct from puerile respiration, though, to the inexperienced in auscultation, it may be difficult to convey a clear notion of what constitutes the distinction, and still more difficult to make any but those possessed of a delicate sense of hearing appreciate the difference. It is, however, necessary to make the attempt. In harsh respiration the character of the sound is changed or modified; in puerile respiration its power is simply increased. Though the former may exist in health, it is ordinarily indicative of disease, however slight. The latter is merely an evidence of increased activity in the function of respiration in the lungs, or in that portion of lung in which it is observed; whether that increase be dependent upon the tender age, the accidental rapidity of the circulation and respiration of the individual, or the defective action of some other portion of the pulmonary tissue. Puerile respi-

ration is in fact a loud vesicular murmur, an exaggeration of the natural sound of respiration. It is as the "*forte*" to the "*piano*" of the same note on the same soft-toned instrument; while the harsh or rough breathing is the "*forte*" of the same note, but upon a loose and jarring string; or, to carry on the former illustration, in the one the same soft breeze passes through a greater number of trees, in the other the breeze is increased to a moderate gale.

The power of the respiratory murmur, instead of being increased, may be diminished, though no disease exist in the chest. This imperfection of the respiratory murmur is usually observed either in parts of the lungs which have been little used, as in the lower regions of the chest of females accustomed to tight lacing; or in persons suffering from deformity, whether congenital or acquired; or in the chests of persons considerably advanced in life. In each case it probably arises from defective respiratory function. In the two first mentioned cases the defect results from want of use, and consequent imperfect expansion, of the lung; in the last case it proceeds from atrophy of the pulmonary tissue. It may, indeed, be regarded as the natural character of the respiratory murmur in old people, and may therefore be called "*senile*," as that existing in childhood is termed "*puerile*," respiration. Occasionally the respiratory murmur is entirely absent from one, or a part of one lung, though no disease be present in

the organ itself. This condition, however, probably never exists without either some mechanical obstruction to the ingress of air, either in the air tubes, or upon the exterior of the organ.

The more common causes of the absence of respiration in a portion of the lung, independently of disease in the pulmonary tissue, are the presence of foreign bodies in the bronchial tubes, or the compression of their calibre by aneurisms or other morbid growths. It may also arise from penetrating wounds of the chest, by which the lung itself remains uninjured, but in consequence of which it collapses from the admission of air into the pleura.

None of these conditions, however, nor any of those arising from malformation, or compression of the ribs by stays, can, properly speaking, be regarded as states of health, though in none is it necessary that the pulmonary tissue, or any of the respiratory organs, should be themselves actually diseased.

The preceding observations, as previously hinted, have especial reference to the murmur of *inspiration*, from which that of *expiration* differs in some important particulars. The difference consists in the decreased loudness and duration of the expiratory, as compared with the inspiratory sound. The tone and general character are the same in both,—one is merely louder and longer than the other.

An attempt has been made accurately to define

the difference, by figures. But as probably no two persons would exactly coincide, even in a given case, as to the purely relative proportions between the two, which they possessed no means of determining accurately by admeasurement, the attempt was not likely to be successful, or generally applicable.

The difference has been much overstated. The proportion pronounced to exist in the healthy condition, by the proposer of the plan for marking the difference by figures, was, in regard to intensity as well as to duration, as 10 to 2; that is to say, the sound of inspiration was considered to be naturally five times as long, and five times as loud, as that of expiration.

Now, though a marked difference does exist in the state of health,—and it is of great importance that this should be particularly noticed, in consequence of its bearing on disease, in which the proportions are changed, and sometimes even reversed,—yet it may be safely asserted that the disparity is not so great as that which has been stated by the proposer of the employment of figures for its admeasurement. Perhaps the proportion of 5 to 2 may be regarded as nearer the truth; though each individual examiner will probably form a standard for himself on this, and on many other subjects which cannot be accurately measured, or strictly defined.

As to tone and softness, as has been before

stated, the murmur of expiration naturally resembles that of inspiration. If, therefore, its character be different, or if its intensity, and its duration, or either of them, be materially increased,—if, that is to say, they nearly approach, equal, or surpass, those of inspiration,—the condition may be regarded as an abnormal one; and the cause may be almost universally attributed to structural alteration, and, if not occurring in a person in advanced life, to disease properly so called.

2. *Of the Voice.*

Some facts may, in disease, be occasionally elicited by auscultation of the Cough, in contradistinction to that of the Voice. But as the sound of each is connected with the same portion of the respiratory organ, as they are similarly conducted by the same parts, and as in health they produce very much the same effect upon the ear, when heard through the parietes of the chest,—as, moreover, cough can scarcely be properly regarded as existing in health,—the Voice and the Cough may, in the following observations, be spoken of together.

If the end of the stethoscope be placed upon the larynx of a healthy person, and, while one ear of the examiner is applied to the free end of the instrument, and the other ear is closed by a finger of the disengaged hand, the individual

examined be directed to talk, his voice will pass through the stethoscope directly into the ear, and will appear as distinct as, and perhaps even more loud than, if his lips were close to the ear. This is *perfect Pectoriloquy* (chest-talking), and the best possible illustration of that important sign, as it exists in some forms of disease.

If now the end of the cylinder be removed from the larynx, and placed upon the lower part of the trachea, and the person be directed to talk as before, it will be noticed that, though the voice appear in some measure to permeate the tube, and though the individual still seem to speak into the ear, yet the voice is not so clear and distinct as in the former case. Such is *imperfect Pectoriloquy*.

Let the examiner now move the stethoscope to the upper bone of the sternum, to the cartilages of the second and third ribs, or to the inter-scapular regions, and again direct the person to talk. A fresh modification of the voice is now observed. It does not appear to permeate the instrument at all, and though loud is comparatively distant and indistinct. The sound appears to be obstructed by the intervention of a bad conductor between the mouth of the speaker and the ear of the listener. This is *Bronchophony* (bronchial sound).

If, finally, the end of the stethoscope be placed upon other parts of the chest than those specified,

the sound of the voice is still heard, and differences of intonation can be recognized, but under ordinary circumstances, articulation can be scarcely said to exist. The distinctness, as well as the loudness of the sound, diminishes the farther the stethoscope, or the ear, (if that be now employed) is removed from the larynx, and the larger bronchial tubes: the former is the source of the sound, and the latter are its best conductors. Thus it will be found that, at the most distant parts, as over the thin edge of the lungs, the voice, if heard at all, is reduced to an indistinct buzzing.

Such, then, are perfect pectoriloquy, imperfect pectoriloquy, bronchophony, and the distant voice in the natural condition; such are they also if well developed when they occur in disease. It must not, however, be supposed that there exists any defined line of demarcation between each, so that it can be said here ends pectoriloquy, and here begins bronchophony. They all gradually merge into each other; they are not distinct species, but merely varieties of sound. So that what one auscultator may call imperfect pectoriloquy, another may probably denominate bronchophony.

Independently, however, of the differences of the resonance of the voice resulting from the distance of the source of sound, and of a good or bad conductor, and irrespectively of the acknowledged fact,

that perfect pectoriloquy gradually becomes imperfect, that this imperfect pectoriloquy gradually merges into bronchophony, and bronchophony into the ordinary state of the voice, as heard generally over the parietes of the chest, there are some varieties, in respect to the thoracic voice, which are worthy of attentive consideration. Thus it is very generally rather more distinct upon the right side, and particularly below the right clavicle and over the right scapula, than in similar situations upon the left side. This may, I believe, be regarded as the rule, though some exceptions occasionally occur. The voice is also much more resonant in some persons than in others. Thus a shrill or acute voice is generally more distinctly audible than one which is hoarse or grave. It is usually also louder in thin persons than in those whose integuments are loaded with fat, or infiltrated with serum, and, the power of the voice being equal, and both free from disease, it is more resonant in persons with contracted chests than in those in whom the thoracic cavity is largely developed. It is generally more distinct, *cæteris paribus*, in females than in males, probably from the greater shrillness of the voice in the former than in the latter.

In some few persons indeed—most commonly females—the voice is, in certain situations, as below the clavicles, or over the scapulæ, so clear, distinct, and loud, as to approach, if not to

attain, the character of imperfect pectoriloquy. Hence arises the necessity of the caution in all examinations of the voice, always to compare the corresponding parts of opposite sides with each other. If after such comparison both sides be found equal, or nearly equal, in resonance, and if after minute examination no other physical signs of disease can be discovered, the unusual resonance may be ascribed with tolerable safety to natural peculiarity.

In consequence of some tones or syllables being better conducted than others (and it is remarkable that it is not always the same tones which in different persons possess this superiority), it is desirable that the same words or syllables should be pronounced during the examination of every part and of each side. It is on this account that it is usual to desire the person under examination to repeat the numerals, one, two, three, four, &c., as in them little or no variety is likely to occur.

7. Auscultation of the Organs of Respiration in Disease.

It will be found convenient in examining the phenomena afforded by auscultation in disease, as in the healthy condition of the respiratory organs, first to confine attention to the sounds, which coincide with the breathing itself, and after-

wards to consider the alterations which take place in the voice and cough.

1. *Of the Respiration in Disease.*

It cannot be too forcibly impressed upon the mind of the student, and it may therefore be here again brought to his recollection, that there are few, very few single signs, if indeed there be any, which are purely characteristic or pathognomonic of disease. The more he observes for himself, and the more attentively he studies the products of disease, the more convinced will he become of this truth. Auscultation and its hand-maid arts often enable us, with almost perfect certainty, to predict various morbid changes, but of the nature of those changes they tell us little or nothing. Physical signs merely indicate certain physical states, which may be produced by two, or by a variety of causes. The physical condition of the organs may be very generally detected by auscultation, and the other different modes of exploration treated of in this work; but information in respect to the morbid process by which it has been produced, must be sought for in other channels, and be determined by other means; as, by the general symptoms of disease, or by the history of the respective cases.

The morbid sounds of the respiration may, to avoid confusion and perplexity, be conveniently first divided into dry and moist sounds. By dry

sounds are intended those which arise from alterations of the calibre of the larynx, trachea, or bronchial tubes, from changes in the capacity, or from obstruction, of the pulmonary cells, or from roughening of the pleural surfaces. By moist sounds are meant those in which, either without, or in addition to, the alteration, obstruction, or roughening giving rise to the dry sounds, fluid secretion or effusion exists in the passages or cells, or air is mixed with fluid in the pleura.

2. Dry Sounds in Diseases of the Air-passages and the Lungs.

(a.) *Larynx and Trachea.*—If, from inflammatory engorgement of the lining membrane, from solid effusion upon the internal surface, or from fluid effusion beneath it, or if, in consequence of preceding ulceration, any contraction exist in the cavity of the larynx, or in the course of the trachea, which gives rise to constriction of the tube, and thereby impedes the free ingress and egress of air, a wheezing, whistling, or crowing sound is produced by the act of respiration. It is loud or slight, according to the amount of the obstruction, and to the power of the patient, and is generally louder during the inspiration than during the expiration. Of this nature is the harsh or crowing respiration existing in croup, and the wheezing of œdema of the glottis. A foreign body

fixed in the œsophagus, and pressing upon the trachea, or situated in the trunk of the air-passages itself, gives rise to similar sounds.

If a tumor,—as an abscess, an aneurism, or a morbid growth,—exist in the vicinity, and exert such pressure upon the cartilaginous box, or tube as to produce obstruction to the breathing, the mechanical results are nearly the same; and the sounds which result from it, though they may vary a little in different cases, are also nearly the same. To the crowing convulsions or false croup of children, in which, though the cause is different, the mechanical effect is the same, so far as regards the sounds produced by respiration, as in laryngitis of the adult, similar observations may be applied

In each of these cases the noise is usually sufficiently obvious, and the evidence of obstruction is sufficiently clear, independently of auscultation. By the stethoscope we are enabled merely to say that obstruction exists. But the nature of that obstruction it telleth not. We may, indeed, be able to state that, in the case of a foreign body, the cause of obstruction is fixed or moveable—that it is now in one part and now in another—but the feelings of the patient, if he be conscious, are usually amply sufficient to assure us of this.

Is, then, auscultation useless in diseases of, or in accidents occurring to, the larynx and trachea? It certainly is not. It is true that it merely indi-

cates that obstruction exists, and does not acquaint us with the nature of that obstruction; but by it we are enabled to effect that which is often of the utmost importance—viz. to point out the exact seat of the obstruction. Without the stethoscope it may be doubtful whether the impediment to the breathing is situated in the larynx, or in the trachea, or whether it is high up or low down in this tube, as the sounds heard by the ear are very similar, if not identical, in either case. But by gently placing the end of the stethoscope upon the larynx, and carefully tracing along the respiratory tube, and attentively listening to the sound of the breathing as we pass it onwards, we are usually enabled to say, pointing to a particular spot, “There is the seat of the obstruction.”

This capability of localizing the source of obstruction is the only benefit afforded by auscultation in diseases of the larynx and trachea, properly so called, with which I am practically acquainted. But though single, it is of immense importance, and may be the means of restoring life when death is impending. It need scarcely be said, that when obvious tumors are pressing upon the trachea or larynx, we are often enabled by means of auscultation to state that such tumors are, or are not, aneurismal.

When, instead of being constricted, the narrow opening of the larynx is widened; if, instead of being thickened or contracted, the vocal cords be considerably ulcerated or destroyed, the natural

reed-like sound which exists in the healthy respiration disappears, and a mere blowing noise supplies its place, similar to that produced by blowing through a reed of large calibre, without more compression of the lips than is necessary to support the tube, or through the hollow of the stethoscope, with the mouth opened sufficiently to present no obstruction to the expired air.

(b.) *Bronchial tubes*.—It has been already stated that the sound of the air passing through the bronchial tubes in the natural condition is, in most parts of the chest, inaudible, and that, under ordinary circumstances, it can only be heard over and near to the upper part of the sternum, and especially at the root of the lung in the interscapular space, where the larger tubes are near to the surface, and little or no pulmonary tissue intervenes between them. Sometimes it is scarcely audible even there. This probably arises from the respiratory murmur contributing to obscure the bronchial sound, which is but feebly conducted by the soft spongy tissue of the lung itself. It will be seen farther on that the case is altogether altered when the vesicular murmur is absent, and when the lung, from consolidation, or other causes, becomes a good conductor of sound.

When the larger bronchial tubes are in any way contracted, whether by pressure upon their exterior, from deposit upon their interior, or from thickening of the structures of which they are composed, the transit of the air through them

is of course impeded. The obstruction thus produced gives rise to sounds which vary considerably, according to the nature and extent of the obstructing cause, and the quality of the conducting medium existing between the tubes and the surface of the chest.

Thus, when the mucous membrane of the large tubes is merely thickened, and they are free from morbid secretion, and not dilated, as in the very early stage of bronchitis ; or in the quiescent state of the membrane of those persons who habitually suffer, after slight exposure, from a chronic form of the malady, or when the walls of the tubes are themselves thickened from long-standing inflammatory change, or "hyper-nutrition," a loud, rough, deep, snoring noise will generally be heard upon the application of the ear to the chest, which clearly depends upon the vibration of the tubes resulting from their narrowing, and the consequent impediment they afford to the free passage of the air. This is the ordinary form of *sonorous rattle* or *rhoncus* ("râle sonore").

When, together with this general constriction, there exists enlargement of the tubes in certain parts,—when, that is to say, partial dilatation is combined with general thickening and contraction, then in addition to the loud snoring noise, there is often a deeper sound, a sort of puffing or blowing, which sometimes approaches the character of the sound produced by blowing into a wide mouthed bottle. If the dilatation be very great,

so as to produce, as it sometimes does, a cavity of considerable size, the same sound may be heard even though no constriction exist at other parts. The physical conditions necessary for its production are the passage of air through a tube, or through a moderately-sized opening into a free space much larger than the opening itself.

When, instead of the larger, the smaller tubes are the parts principally constricted, the sound is of a higher pitch ; hissing or squeaking noises are then heard, which constitute the *sibilating rattle* (“*râle sibilant*”) of authors.

The *sonorous* rattle then depends upon some obstruction, either by compression or thickening, absolute or comparative, of the larger bronchial tubes, while the *sibilating* or hissing rattle, or rhoncus generally arises from a similar affection of the smaller tubes. It is to the sonorous rattle or rhoncus what the fife or the kit is to the flute or the violoncello. These rattles may exist either singly or together, independently of the presence of fluid secretion in the bronchial tubes, and consequently without any moist rattles or rhonci, though they are most commonly conjoined with the latter sounds.

Together with this obstruction of the larger or smaller bronchial tubes, and very commonly as its direct mechanical effect, there exists a dilated condition of the pulmonary vesicles, and consequent thinning, or it may be even rupture, of the membrane which separates them from each other.

This condition of the pulmonary tissue constitutes emphysema of the lungs. As the affections of the tubes and of air-cells are so commonly connected together, as cause and effect, it may be well to consider them in immediate succession.

When the air-cells are considerably dilated, and their partitions, or the tissue intervening between them, are consequently reduced in thickness, though the bulk of the lung, or of the part of the lung affected, is increased, the amount of pulmonary tissue in any given portion of the diseased organ is decreased. The natural elasticity of the tissue is diminished, and in extreme cases is destroyed. The area of the cells is not only absolutely greater than in health, but, compared with the calibre of the tubes, it is also enlarged. The lungs, when emphysema is general and extensive, are in a state of forced inspiration. They are too full of air, and are incapable of expelling it. The necessity of fresh air begets a constant effort to draw still more into the chest by the extraordinary efforts, and forced contractions of the muscles of inspiration, while expiration is but feebly assisted by the muscles, which under certain circumstances contribute to that act. Expiration, in fact, unaided by the elasticity of the pulmonary tissue, and assisted in only a trifling degree by the abdominal muscles, is principally, or almost solely, effected by the elasticity of the ribs. The necessary results, as regards auscultation, of this disease of the lung,

will then be readily understood to be — 1st. That the act of inspiration is shortened, in consequence of the quick and strong effort to effect it. 2nd. That the natural inspiratory murmur is greatly decreased, both as regards intensity and duration, in consequence of diminished density of the vesicular tissue of the lung on the one hand, and of the small amount of air inhaled on the other. 3rd. That the duration of the expiration is greatly prolonged by reason of the deficiency, or absence of elasticity in the pulmonary tissue; and 4th, that a loud wheezing noise accompanies the expiration, vastly exaggerating the natural sound, the result of the obstruction which exists to the exit of the air through bronchial tubes comparatively or absolutely contracted.

The genuine auscultatory signs,* then, of emphysema of the lungs, are a short inspiration

* With *crepitation*, as the direct result of pure emphysema, (the "*râle crepitant sec à grosses bulles*" of Laennec, and other auscultators, who had followed him,) I have no practical acquaintance; and though I am not prepared, and am far from being disposed, to say, that it never does nor can exist, I believe I may truly state, that I never saw an unexceptionable case, in which it could be proved to have been present; and from multiplied observations I have no hesitation in expressing my assured conviction that the crepitant rattles, so constantly heard in emphysema, arise very generally, if not universally, from fluid secretions in the dilated cells, and smaller bronchial tubes.

with a feeble inspiratory murmur, and a loud, protracted, and wheezing expiration.

The loud and prolonged expiratory wheezing of emphysema is very remarkable, and may be satisfactorily explained as above. But not only is the expiratory wheezing extended vastly beyond the ordinary duration of the time employed in expelling the air from the chest, and greatly exceeds that occupied by inspiration; but in extreme cases it may sometimes be observed that this wheezing obviously continues after the act of inspiration has commenced. However apparently difficult may be the explanation, the fact is quite certain. It may be perhaps presumed to arise as follows.

The great effort of persons affected with extensive emphysema, is to gain a sufficient supply of fresh air. To accomplish this, they must exclude from the lungs as much as possible of the air which distends them. The act of expiration, as has been already explained, is chiefly effected by the elasticity of the ribs, which have already become considerably dilated, by the long-continued pressure exerted upon them by the enlarged lung from within. The elevation of the ribs upon inspiration, however small may be its extent, is quickly effected, and when their elasticity comes into play by the cessation of the inspiratory effort, their tendency to return to the situation from which they were raised is immediate. So

far then as the ribs are concerned, the expiration in emphysema may be as short, or even shorter than inspiration. Not so as regards the included air, which cannot escape from the multitudinous cells in which it is contained through its only outlets, the absolutely or comparatively contracted bronchial tubes, with a rapidity equal to that with which the ribs fall. The air, therefore, in the chest is temporarily compressed, and its tendency to escape does not cease when the ribs have ceased to fall. More than this, the elastic fluid within the chest does not regain its equilibrium, as regards density with the external air, until the elevation of the ribs actually occurs. This elevation does not, as in the state of health, take place immediately upon the contraction of the muscles of inspiration; nor, in consequence of the already largely dilated state of the chest, and the consequently unfavourable position in which those muscles are placed for efficient action, is it effected till the muscles have already contracted to some extent.

Such appears to be the most probable explanation, as well of the sounds heard in emphysema, as of the curious fact to which reference has been made. The theory of the matter may be defective, or physiologically incorrect. The facts remain the same.

Supposing then that, together with general emphysema, there exists thickening, or obstruction of the larger bronchial tubes in one part,

and dilatation in another,—contraction of the smaller canals here, and enlargement of them there,—the loud sonorous rattle arising from the contraction may be combined with the puffing or blowing of dilatation, the hissing of capillary narrowness, and the wheezing of emphysema. Under such circumstances, which are far from uncommon, the sounds, even independently of moist rattles, are of course exceedingly various. The note of a base viol, the cooing of pigeons, the squeaking of young mice, the hissing of the escape of steam, snuffling and snoring, whistles and wheezes, puffing and piping, in various degrees and combinations, may be occasionally heard, and may be employed to characterise the *strange and multifarious noises* which are then audible.

Phthisis.—In the earlier stages of Phthisis, when but few tubercles are deposited, and no surrounding consolidation exists in the lung, a slight alteration of the respiration is often the only recognizable physical indication of the presence of the abnormal deposit. The tubercular bodies, whatever their origin, whether from inflammation, or from depraved nutrition, and whatever their exact seat, whether in the air-cells themselves, or in the vascular tissue which constitutes their boundaries, necessarily encroach upon the pulmonary vesicles. They consequently impede the free expansibility of the lung, and must, in some degree at least, detract from the elasticity of the organ. They must also exert some amount of pressure upon

the capillary branches of the bronchial tubes. The consequence of this condition is, not merely that the soft breeze-like inspiratory murmur is perceptibly, however slightly, diminished, but that it is replaced by comparatively a harsh and hoarse sound. This latter sound, it is presumed, is dependent upon the pressure exerted upon the small and flexible bronchial tubes by the abnormal tubercular deposit, as the defective inspiratory murmur is upon the imperfect expansion of the pulmonary vesicles. From the same cause the intensity of the expiratory murmur is increased, and its duration prolonged. As was stated on a former occasion, whatever may be thought of the explanation here offered, or whether it be really correct or incorrect, it is certain that among the earliest physical indications of the presence of tubercles in the apex of the lung, are deficient respiratory murmur, hoarse respiration, and prolongation of the expiration.

When, in a later state of phthisis, the portion of the lung affected becomes in some measure consolidated, partly from the increase of tubercles, but principally in consequence of surrounding pneumonia, or when pure pneumonia attacks a lung, and proceeds without ulceration to firm dry consolidation, the air-cells are entirely obstructed, and the vesicular murmur in the part is consequently absent. But in both these cases, in the tubercular combined with the pneumonic, and in

the purely pneumonic consolidation, auscultation affords other signs than the mere absence of the respiratory murmur. Among the most important of these is bronchial respiration, or tubular breathing. The spongy tissue of the lung being converted into a firm solid mass, is changed from a very bad to a good conductor of sound, and the sound produced by the passage of air through the larger bronchial tubes is thus better conveyed to the surface of the chest; it also becomes more distinct from the absence of the natural sound of the respiration, which ordinarily interferes with, or conceals it.

Does, then, tubular breathing necessarily indicate tubercular, or pneumonic consolidation? It will be seen in a future page that it certainly does not. It merely indicates this: that the bronchial tubes are either actually nearer the surface, or, acoustically speaking, are made to approach it by the intervention of an improved conductor; and, if the respiratory murmur be at the same time absent, that the cells of the lungs are either compressed or obstructed. To arrive at the cause of this physical condition, other means must be adopted than the mere auscultation of the breathing.

The character of tubular respiration varies according to the extent and situation of the part affected; that is to say, according to the size and amount of the tubes passing through, or near to,

the consolidated tissue. It is also, of course, modified in its purity and intensity, by the presence or absence of the respiratory murmur, and of morbid rattles. Thus it may be of a shrill tone if the tubes are small; or it may be deep if they are large; but in either case, it resembles the sound produced by blowing through a reed, a quill, or a straw, with little, or without any, compression of the lips, according to the perfection with which it exists. The more complete the consolidation, or the more pure the sound, the less obstruction does there appear to be to the transmission of air through the reed; the less compression of the lips does it require accurately to represent it; the more perfect, in fact, is the tubular character of the sound.

If, as it sometimes though rarely happens, a bronchial tube be dilated into a large cavity; if, as it more frequently occurs, a moderately dilated tube be surrounded by a mass of consolidated lung, or be very close to the surface of the chest; or if, whether from tubercular degeneration or pneumonic softening, a large cavern be formed in the substance of the lung itself, and be not very deeply seated, or have only diseased pulmonary tissue intervening between it and the parietes, then either a deeply-toned buzzing sound, or a sonorous and hollow kind of whistling, or hissing noise, takes the place of bronchial respiration. The former is named *amphoric breathing*,

from its frequently exact similarity to the sound caused by blowing into a wide-mouthed bottle. The latter is termed *fistulous breathing*. It may be so denominated either from its resemblance to the sound caused by air permeating a reed which *is split*, or from its presenting to the ear of the examiner the impression of elastic fluid passing through a fistulous opening. Both of these sounds are also occasionally heard when a communication exists between a bronchial tube and the pleura.

Nearly allied to these sounds is the *metallic respiration*. It sometimes accompanies one or other of them, and very generally exists in very nearly similar physical conditions of the lung or pleura. It gives an impression to the ear of a large cavity, equally with the amphoric breathing; but it possesses a ringing or brazen twang, which may be perhaps correctly represented by blowing through a tube into a moderately-sized bell or glass vase, the sides of which vibrate slightly from the blast.

It is often, but not necessarily, associated with another sound, (*metallic tinkling*,) and appears to depend upon the free vibration of air against the resonant parietes of a large cavity. The whole of these three sounds are generally, if, indeed, they be not uniformly, confined to the act of inspiration.

In what then consists the difference of the cause

of these three sounds? What are the physical conditions necessary for the production of each? They are all similar, but not identical, in character. They all give to the ear an impression of air passing into a cavity of considerable size, they all likewise exist under circumstances nearly similar, and they sometimes replace each other. Though, therefore, the physical conditions necessary for the production of each must be very similar in all of them, it is clear they are not identical. In what, then, does the difference consist? To this I, at least, am not able to reply with confidence. I may, however, state that I believe the difference of the fistulous, and the amphoric respiration, to depend upon the size of the opening leading into the cavity of the lung or pleura; that the opening is small, or obstructed when the former, and comparatively large, when the latter sound is heard; and therefore that amphoric respiration may become fistulous, by the tube, through which the air gains admission into the free space, becoming obstructed with secretion, or being diminished in its calibre from any other cause, and that fistulous, may be converted into amphoric, breathing by the removal of such obstruction, or by the tube becoming otherwise enlarged. I conceive also that the metallic ringing of the respiration is dependent rather upon the nature of the walls of the cavity itself, than upon the form or size of the opening leading to it, and that it is from the vibration of these walls, being communi-

cated to the air contained within them that the peculiarity of the tone arises. Hence I believe it happens that metallic respiration is very rarely heard, unless some portion at least of the walls of the cavity are formed by the solid parietes of the chest, or by a thin layer of firmly adherent pleura lining them. The determination of this question, however, is fortunately not one of importance in a practical point of view, whatever interest it may possess in other respects.

The auscultatory signs produced by *Malignant disease* of the lung are exceedingly variable, according to the situation, amount, and distribution of the morbid deposit. Sometimes malignant tubercles are irregularly scattered through the lung; but as they give rise to no general symptoms, the chest is frequently not particularly examined, or, if it be examined, it may afford little or no indication of their presence. At other times, disease of much less amount may, from its position near to, and its pressure upon, a considerable branch of a bronchial tube, produce a loud, hoarse, and harsh tubular breathing, or wheezing, and from its merely mechanical effect, may at the same time give rise to great general distress.

In other forms of the complaint, when the malignant deposit exists in a large isolated portion of the organ, and the bronchial tubes are not materially compressed or pushed aside, the physical signs, as might indeed be expected, are much the same as those arising from consolidation of

the lung from other causes, and, as regards the breathing, are almost confined to the absence of the respiratory murmur and the presence of bronchial respiration. When, again, the malignant mass is so large, or is so situated, as to push aside the lung altogether, or to compress and actually to close the bronchial tubes, all sound of respiration, whether vesicular or tubular, ceases in the part affected; excepting so far as it is conducted by the consolidated mass from the lung of the opposite side of the chest, or a comparatively healthy portion of the lung on the same side. The sounds, under such circumstances, appear not to be close to the ear, but to come, as in truth they do come, from a distance.

Such, then, are the chief and the most important of the *dry* sounds connected with the respiration in disease. For the sake of distinctness, they have been mentioned separately. Separately they often exist; but the student must recollect that they are more frequently variously combined with each other, and with the different moist sounds or rattles, which will be spoken of presently, and which often detract from the clearness of their character, and consequently render them more difficult of discrimination.

3. *Dry sounds in diseases of the Pleura.*

The smooth, moistened surfaces of the pleura, during the respiratory movements of health, softly

glide over each other without giving rise to any sound. They act like a beautiful piece of machinery in good trim and well oiled.—That they do so glide over each other in health, appears sufficiently evident from the effects of disease, when the machinery is out of order, the surfaces lose their polish, or, it may be, are not supplied with the amount of oil necessary to make them play easily upon each other.

The first physical effect of inflammation in a serous membrane is supposed to be to render it abnormally dry. It is presumed that this inflammatory dryness may itself, independently of any effusion, give rise to a morbid sound, just as the absence of oil on a piston, or in a joint, may be the cause of an unusual one in machinery. This is possible; but it may be questioned whether the mere dryness of a serous membrane coexisting with, and the pure result of, the first stage of inflammation, has been proved ever to have produced such an effect.

When, however, effusion has taken place, and the serous membranes are coated with an albuminous layer however thin, when the smooth polished surface of the pleura is replaced by a rough and plastic one, it is not merely probable, but in numerous cases it has been proved, that the folds of the membrane, in place of gliding smoothly and noiselessly over each other, stick to each other on the way, and by their motion produce a sound. Two rough surfaces, in fact, are

now in contact, and the friction of one against the other, like that of other rough surfaces, makes a noise, or in other words, produces a vibration of the substances of which they are composed. This sound is appropriately called *pleuritic rubbing*, *friction sound of the pleura*, or, "*frottement ascendant et descendant*." This last term has been employed in consequence of the sound usually accompanying both the elevation and the falling of the ribs.

The sound varies much in character. This variation probably depends upon the quality, and the quantity of the solid matter intervening between the folds of the pleura. When the effused lymph is recent and soft, it appears as if the opposed surfaces stuck together by a slightly adhesive material. The sound which results from the motion of the parts, under such circumstances, is interrupted and jerking. It closely resembles that which may be produced by pressing firmly together, and moving slowly over each other, the palms of the hands, moistened in the slightest degree possible with perspiration. It may be also likened to the stiff or crisp noise caused by squeezing a packet of arrow-root.

When the plastic effusion, in consequence of the separation, or the absorption of some of its liquid constituents, has become less moist and adhesive; when a solid, dry, rough coating is left upon corresponding portions of the pul-

monic and costal pleura, a simple rubbing noise is heard upon inspiration, and expiration, which may assimilate to that arising from the attrition of two pieces of cloth. At other times the sound is very loud, harsh, and grating (*"raclement"*), and can often be heard and felt by the patient himself, as well as by the physician and the attendants, if the hand or ear be placed upon the chest. This harsh, rough, grating sound ordinarily, but not universally coexists with the inspiration alone. It may be compared to that arising from the attrition of two pieces of rough wood.

There is yet another peculiarity in reference to the sound arising from morbid deposit in the pleura. It sometimes assumes a curiously creaking, or squeaking character, and really most exactly simulates the noise caused by bending a piece of stiff leather (*"craquement de cuir"*). In this case it is ordinarily heard only towards the end of the inspiration; a circumstance that certainly renders probable the opinion, that it depends upon the stretching and tension of old adhesions. Sometimes, indeed, the different varieties of pleuritic rubbing, excepting perhaps the first, are heard only at the end of a forced inspiration, when the lung is most distended with air, and when, if the pressure be not greater, the extent of contact between the two folds of the pleura is increased.

The first and second variety of pleuritic rubbing above referred to, may certainly exist either previously to the effusion of fluid, or subsequently to its absorption; but it is at least probable, that the harsher kinds of sound are connected with chronic diseases, and therefore ordinarily follow the absorption of fluid. Immediately, however, that fluid intervenes between the two surfaces in the case of recent disease, or so long as any remains behind in that of longer date; while, in fact, the presence of fluid prevents the contact, and consequently the friction of the two opposing surfaces of the pleura, no rubbing sound can be heard.

In the earlier stages of phthisis, the murmur of inspiration below the clavicles and in the acromial region often presents an irregularly interrupted, jerking, or catching sort of sound ("*saccadé*"). It rather accompanies, than supplies the place of, the respiratory murmur. Its origin is not certain, but it appears most probable that it is connected with a roughened state of the pleura, the result of effusion, and that it is, in fact, a modification of pleuritic rubbing.

At other times, though much more rarely, there is heard during the inspiration, in the same situations, a soft rustling, or crumpling noise, approaching to that produced by squeezing together soft gauze paper—*pulmonary rustling* ("*froissement pulmonaire*"). It has been so named, on

account of its supposed connection with an alteration of the pulmonary tissue; but, like the former sound, it appears to myself to be more probably connected with some change in the serous membrane, which prevents the two layers gliding over each other with perfect smoothness, and without noise. Neither of these sounds are of much value in a diagnostic point of view, especially as their origin is uncertain. But as they indicate some change in that part of the lung, or of its covering, in which tubercles are, in a vast majority of instances, first deposited, they should not, in doubtful cases, and in connection with other physical signs, be altogether neglected. I believe that they result from changes in the pleura, and that these changes depend upon the presence of tubercles, either in the lungs, or in the membrane itself. It is, however, right to say, that I have no solid grounds for this opinion, as I have never had an opportunity of investigating their origin at the dissecting table; nor, indeed, is it likely that such should be the case, as they are, comparatively, so rare, and of so little importance (the second more especially), and occur only in so early a period of phthisis, that the opportunity of investigating the physical condition upon which they depend, must be exceedingly rare indeed.

Where either fluid or solid matter exists in the pleura, to an extent sufficient materially to com-

press the spongy tissue of the adjoining lung, the respiratory murmur, of course, ceases in that part, in consequence of the air-cells not being expanded during inspiration. The respiratory murmur being absent, and the spongy tissue being compressed, and, by its compression, being converted into a body of superior conducting power, the sound of the air passing through the bronchial tubes is conveyed to the surface. Thus tubular breathing, or bronchial respiration, together with the absence or the distance of the respiratory murmur, becomes a valuable sign of effusion into the pleura.

It is, however, a sign of only a limited amount of fluid, excepting under peculiar circumstances, and in particular parts. For, if the lung be not adherent to the parietes of the chest, and if it be otherwise in a healthy condition, as the fluid accumulates from below, it presses the lung more and more upwards. It is consequently farther and farther removed from the general surface of the chest; so that at length, when the fluid is in large quantity, it is compressed into a very small compass close to the spine. Under such circumstances it is clear that the bronchial tubes themselves, in which the sound, in part at least, originates, and by which it is primarily and best conducted, must be far removed, together with the lung, from the greater portion of the parietes. The source of the sound, therefore, being far removed, though

the fluid *may be* a better conductor of sound than the natural spongy structure of the lung, not only is the respiratory murmur lost; but the bronchial respiration itself, excepting at the root of the lung between the shoulder-blades, is greatly diminished, and in many cases quite inaudible.

Some exceptions, as has been already hinted, exist to the uniform correctness of these general statements. The first of these depends upon the gravitation of the fluid. When the fluid is not in large quantity, is not confined by adhesions, and is not combined with a large proportion of plastic or albuminous matter, it may change its situation, according to the varying position of the patient's body. The part therefore in which no respiratory murmur is audible, and in which tubular breathing is distinct, while the patient is supine, may be destitute of tubular breathing, and afford a very tolerable respiratory murmur, when he turns upon his face. This fact is much more commonly observed in hydrothorax, than when the effusion has resulted from pleurisy, in which it is certainly not so common as it has been represented to be.

The next exception arises from adhesions. Though the fluid may be in large quantity, and though the lung may, in the main, be greatly compressed, and, as regards its chief expanse, be far removed from the general parietes, it may be bound down at one or more parts to the costal, or phrenic pleura, by old adhesions, too firm and

strong to yield under the pressure of the fluid. At this part, then, the lung is not removed from the surface, and though no trace of pure respiratory murmur may be distinguishable, it is yet quite possible, and indeed it often occurs, that very well-marked bronchial respiration is heard under such circumstances in parts far removed from the root of the lungs.

It may be here observed, though the observation can be scarcely regarded as an exception to the general statements previously made, that, in all these cases, bronchial respiration can be heard, even more distinctly than naturally, in the interscapular region. It is more loud and pure upon the diseased than upon the healthy side, in consequence of the fluid pressing the large bronchial tubes against the solid parietes of the chest, and excluding the air from the surrounding, and intervening lung.

The last exception that will be mentioned is the most curious one, and by far the most difficult of explanation. In a few rare cases it has happened, that though the pleura has been supposed to be full of fluid, and in one case was a few days afterwards found to be so upon inspection after death, not only has tubular breathing been general, but a shrill resonance of the voice has been distinctly audible over the whole side. Though the lung was believed to be compressed into the smallest space, and was actually found to be so a short time subsequently, and though, therefore,

the bronchial tubes were far removed from the solid parietes, in all parts, excepting in the neighbourhood of the spinal column, still their vibrations were communicated to those parietes, and from them to the ear. How did this happen? It is, indeed, difficult to say; and because difficult to explain, the fact itself may be doubted. The same thing, however, in regard to the voice, though not in respect to the tubular breathing, occurred in a man, from whose chest, immediately after the examination, seventy-two ounces of fluid were drawn off by paracentesis. In reference to this case, then, there could be no doubt; and what occurred in one case in regard to the voice, might surely occur in another in regard to the breathing, or to both. How, then, is it to be explained? It is one of those cases in which ordinary acoustic principles appear, most probably from our not being aware of the whole circumstances, to be insufficient for the satisfactory explanation of what is heard in the practice of physical diagnosis. If a conjecture might be ventured upon the subject, it would be this:—So long as there exists any portion of the lungs which is capable of farther compression, so long, therefore, as the parietes are slack, and the fluid contained within them, though in large quantity, is moveable, so long is bronchial respiration inaudible over the general surface of the side. But when the compression is great enough to render the parietes and the fluid contained within them tense, but not so great

as to flatten and close the largest bronchial tubes, then the vibration of the tubes is communicated to the spine, or to the bony parietes, against which they rest, and thence, as in a well-braced drum, to the walls of the entire cavity. The fluid is so tightly squeezed, that it moves as a solid with the solid walls. The circumstance appears, indeed, to bear some analogy to the equally curious one previously mentioned, and stated to have been observed on several occasions by Irish physicians, viz. the tympanitic resonance upon percussion, which has been noticed in some cases of entire consolidation of the lung. As to the breathing being sometimes not heard, when the voice is audible and shrill, this probably depends upon the greater or less extent of the vibrations, from which the two proceed.

When a communication exists between one or more bronchial tubes, and the cavity of the pleura; whether the communication be the result of a tubercular abscess bursting into the serous membrane; of a collection of purulent fluid in the pleura emptying itself through the lung; or of gangrene, which are the three most common causes of pneumothorax; a considerable quantity of air usually finds its way into, and continues within, the serous sac. Here then, if at any time, might be probably presumed to exist, the *fistulous breathing*, the *amphoric respiration* or *bottle blowing*, and the *metallic resonance*, which have been previously mentioned, while speaking of the

diseases of the lung; and here it is that they do, in fact, exist most frequently, and in their greatest perfection. It would be useless to attempt to describe them again; they should be heard to be perfectly known, and thoroughly appreciated. To hear them once will more conduce to their future recognition than whole pages of descriptions and similitudes. It cannot either be necessary to explain their origin. It is sufficient to say, that they indicate with tolerable certainty a large cavity with resonating walls, and a free ingress of air.

A caution is here requisite, to guard the student against a very important, and in some respects, not a very unfrequent source of fallacy. The whole of the sounds last described, and particularly that mentioned in the next paragraph, may be distinctly heard in examination of the chest, and appear to be close under the ear; while they in reality exist in the abdomen, and arise from the motion of air and fluid in the distended stomach. The sounds may, in truth, be under the ear, in consequence of the largely distended stomach, by pushing up the diaphragm and displacing the lung, coming into near contact with the ribs, as high even as the fourth or fifth; or, they may be so in appearance only, in consequence of their being conveyed through the lung, increased in its conducting power by consolidation, or by the compression resulting from fluid effusion.

There is another sound connected with disease of the pleura, as well as of the lung, the tone of which is assuredly often dry, and which may possibly sometimes arise independently of the presence of fluid. But as "*metallic tinkling*," the sound to which I refer, is certainly most frequently connected with the vibration of fluid, and as it is not perhaps proved that it ever exists independently of the presence of fluid, it may be better here barely to refer to it, and reserve its fuller consideration for the next section.

4. *Moist sounds in diseases of the Air-passages, and Lungs.*

With the view of relieving the student of some portion of the perplexity, with which he may possibly have hitherto contemplated the names of the various rattles, or moist sounds which occur in disease, a few simple statements will be made in reference to the whole of them, as a class, previously to either mentioning them individually, or particularizing the complaints during the progress of which they arise.

All the moist sounds heard in diseases of the respiratory organs, it may be then first remarked, arise from the passage of air through fluid, or the admixture of air with fluid. They may occur during the act of inspiration, or of expiration; may be situated in the larger, or in the smaller

bronchial tubes, which may contain mucus, blood, serum, or pus; they may originate in a portion of disorganized lung, or in a cavity, either pulmonary or pleural; they may be produced independently of any respiratory act by the simple motion of body; they may vary in tone and in character, in site and in pathological origin, but their origin, physically speaking, is the same in all. It is the mechanical admixture of fluid and air. Separately, either of these is incapable of giving rise to moist sounds. Air will certainly not do so, and fluid cannot without the presence of air. In a tube replete with air, no moist sound will be produced by agitation, neither will a cask quite full of fluid, however shaken, give rise to any rattling or squashing noise.

The moist sounds similar, and for the most part identical, in their mode of production, have been differently named, according to the size of the bubbles producing them, or, what in a great majority of cases, amounts to the same thing, according to the calibre of the tube through which, or the size of the cavity into which, the air passes. Thus the bubbles of air arising from its admixture with fluid either are actually of greater dimensions in the trachea, than in the bronchial tubes, and in the larger, than in the smaller tubes; or the different extent of the space, in which the bubbles are formed and burst, pro-

duces a corresponding difference in the resulting resonance, which causes them to appear to be so.

When the bubbles are very large, or their resonance is very full, as they occur in the trachea, or in a tubercular cavity, they are called *tracheal rattles*, or *gurgling*. When they are somewhat smaller, as in the bronchial tubes of first or second size, they are called *mucous*, or *bronchial rattles*. When they are still smaller, as in the finer divisions of the bronchial tubes, they are termed *muco-crepitating rattles*; and when in the air-cells themselves, or in the capillary branches in immediate connection with them, they are named, *crepitating rattles*.

Let the student imagine the bubbles of the tracheal rattles or gurgling to be the size of a horse-bean; those of the mucous rattle the size of a pea, or a tare; those of the muco-crepitating rattle the size of a millet-seed; and those of the crepitating rattle the size of a poppy-seed, and he will form no very incorrect notion of the relation of these sounds to each other; and, bearing in mind the manner in which they are produced, he will, in most cases, not be far wrong as to their seat and their import.

The moist sounds vary as to their distinctness and quality, according to the facility or difficulty with which the vibrations producing them reach the surface of the chest; or, in other words, according to the goodness or badness of the conductor, intervening between their source, and the

parietes. They are, for example sake, more shrill and loud when the lung is consolidated, than when the pulmonary tissue is in a healthy condition. They also vary, in a slight degree, according to the nature of the fluid in which they originate. They are somewhat more shrill, or sharp, if produced by blood in the tubes, than by serum, and in stiff tenacious mucus, than in the ordinary bronchial secretion.

The preceding statements are assuredly sufficiently simple, and must, it is believed, be intelligible to all; yet it is hoped that, by bearing them constantly in mind, the student will be enabled, not merely easily to follow, and to understand the succeeding remarks, which can be regarded as simply an extension of them, and an application of them to particular cases; but that he will be himself competent to recognize, and in some degree to appreciate the value of many of, the fluid sounds, which he hears in his examinations of pulmonary disease.

When, either in addition to, or independently of ulceration, constriction, or compression, fluid is present in the larynx or trachea, the passage of the air through the fluid, in the acts of inspiration and expiration, and the consequent formation and bursting of bubbles, produces a loud and loose bubbling, or gurgling noise. It may be, and generally is, audible without approximating the ear, or applying the cylinder to the throat; but it is most distinctly heard, and its situation

is most accurately defined by means of the stethoscope, gently and evenly placed upon the side of the neck. The sound is a little, but only a little, modified by the quality of the fluid in which it originates. It is rather sharper in tone when the fluid is thick and viscid, than when it is thin and bland. Thick purulent, or muco-purulent matter, and tough viscid mucus, give rise therefore to a sound rather more shrill than common bronchial mucus, watery phlegm, or blood.

The difference arising from the quality of the fluid, may be fairly illustrated by blowing through a straw into simple water, and into a somewhat denser fluid, as soap and water, or syrup. The moist tracheal sound is the same as that which, in the last few hours of life, results from the failure of the nervous energy, is familiarly called "the rattles," and by uninformed persons is very generally, but incorrectly, considered to be the certain precursor of death. By auscultators it is called "*tracheal rattle*."

When the bronchial tubes of first or second-rate size contain fluid, the bubbles which arise from the transmission of air through them, appear to be smaller than those existing in the trachea and larynx. The resulting sound is called *mucous rattle* ("*râle muqueux*."). It is present whenever fluid exists in the tubes, and the free passage of air admits of the production of bubbles. The sound is most commonly produced

by the pressure of bronchial mucus in the tubes. It is hence called mucous rattle. It may, however, arise from other fluids. Its designation is not changed on that account, but whether originating in mucus or pus, in blood or serum, it is still called mucous rattle, as the name is generic, and has reference rather to the character of the sound, than to the nature of the fluid in which it originates.

It must not, however, be supposed, that in the mucous rattle, the bubbles are, or appear to be, all exactly of the same, and that a medium, size; as, independently of the difference of the calibre of the tubes themselves, it is clear that small bubbles may arise in a large tube, though large bubbles cannot exist in a small one. The name and the description are derived from the general, and most prominent character of the sound, which gives an impression to the ear of the bursting of bubbles of a moderate size.

When general dilatation of the tubes is present, the bubbles appear to be larger than usual; especially in those parts of the lung in which the tubes are naturally small.

When, instead of general dilatation of the tubes to a moderate extent, there exists such an extension of the walls of one, or of several of them, as to form considerable, though it may be irregular, cavities, as when caverns are present in the lungs, whether those caverns result from

tubercular disease, or from pneumonia, a large tracheal rattle, or more properly, *gurgling* (*"gourgouillement"*), is heard in the part affected upon each act of inspiration. Are, then, it may be asked, tracheal rattles and gurgling precisely the same thing? Not exactly. They are very similar, but not identical. Independently of tracheal rattles being, strictly speaking, confined to the larynx and trachea, though both the tracheal and gurgling rattles appear to result from the bursting of large bubbles of air, and both, therefore, also appear to proceed from a space of considerable size, there is still a slight difference between them. Gurgling gives an impression to the ear of a thicker or more dense fluid than the ordinary tracheal rattle. The sound, too, generally corresponds with the actual condition of the fluid in which it originates. For as the rattles in the trachea usually proceed from common bronchial mucus, or from blood, gurgling is ordinarily produced by pus, thick muco-purulent or softened tubercular matter. Does true gurgling, then, necessarily indicate the presence of either a largely dilated tube, or a cavity in the lung? Certainly not. It generally does so, but not necessarily. For if a bronchial tube of moderate size, or a tube dilated in only a slight degree, contain thick purulent mucus, and be surrounded by a dense mass of consolidated lung, the sounds which result from the

passage of air through the thick fluid in the tube, and which, under ordinary circumstances, would give rise only to a mucous rattle, will be so increased in loudness and fulness, by the improved conducting power of the solid mass, as exactly to resemble the gurgling of a cavity. Let the student, therefore, be careful not to pronounce too confidently upon the existence of a cavity, from the presence of gurgling alone, or he will assuredly be liable to error, and will sometimes find himself mistaken. *Gurgling*, then, is a large and resonant mucous rattle not existing in the trachea. It commonly arises from a cavity in the lung, or from a largely dilated bronchial tube; but it may proceed from a tube of normal size, which is filled with thick fluid, and is also surrounded by a mass of consolidated lung.

When the smaller tubes are the parts solely or principally diseased, or rather when morbid secretion, or other fluid, is present in the minute branches of the bronchial tree, the moist sound, which is produced by the passage of air, appears to arise from the bursting of bubbles smaller than those constituting the mucous rattle, and larger than those, which are next to be mentioned, and which give rise to *crepitation*. The sound is therefore appropriately called the *muco-crepitating rattle*. Excepting as regards the apparent size of the bubbles producing it, and the calibre of tubes in which it exists, it resembles the

mucous rattle, to which it bears much the same relation as the dry *sibilating* does to the *sonorous* rattle or rhonchus. Like the mucous rattle, it is rendered more shrill and sharp by the presence of consolidation of the lung, and from causes similar to those which have been already mentioned, when speaking of the mucous rattle and gurgling. When fluid is present in the air-cells of the lungs themselves, the sound which results from the admixture of air with the fluid appears to arise from the bursting of bubbles still smaller than those previously referred to; and, if the fluid be strictly confined to the cells, the bubbles, like the cells themselves, appear all to be nearly equal in size. It matters not what may be the nature of the fluid,—supposing it to be situated in, and confined to, the pulmonary vesicles,—whether it be blood, serum, or viscid mucus, the sound which is heard upon inspiration is like that produced by the bursting of bubbles of very small and of equal size.

Whether, therefore, the cells are partially filled with serum, as in œdema of the lung,—with blood, as in apoplexy of the lung,—or with thick viscid mucus, as in inflammation of the lung, a very fine sound, similar to that which is heard when salt is thrown upon the fire, and is called *crepitation*, arises from the intermingling of the inspired air with the fluid so situated. It is hence called the *crepitating rattle* (“*râle crepitant*.”) When

existing in the greatest purity, it is most correctly represented by rubbing a lock of hair between the finger and thumb close to the ear.

Is there, then, no difference between the crepitating rattle of œdema, pulmonary apoplexy, and pneumonia? It would be, perhaps, incorrect, or at least questionable, to state that a difference does always, and must necessarily, exist between them; but in practice it is found that a difference, in a great majority of cases, actually does exist. The truth is, that the *pure* crepitating rattle is very rarely heard, either in pulmonary apoplexy or in œdema. It is generally combined with the muco-crepitating rattle of the small tubes, into which, as well as into the air-cells, the serum or blood is almost always likewise effused.

As the air intermingles with the fluid in the tubes upon its exit from, as well as upon its entrance to, the lungs, the mucous and muco-crepitating rattles are heard during the expiration as well as during the inspiration. The same occurs, and for the same reasons, when serum or blood is present in the capillary branches of the tubes. Now, although it be quite true that capillary bronchitis very frequently exists in combination with pneumonia, and though, when such is the case, it may be difficult, or even impossible, to state with certainty, and from the sound alone, that the crepitating and muco-crepitating rattle may not arise from pulmonary œdema or apoplexy; yet it is

also true that pneumonia is often present without any affection, or without any physical indication of any affection, of the small tubes.

It is in such, and in such cases only, that the crepitating rattle is heard in purity and perfection. It is then very small, very crisp, very equal, and heard only at the end of the inspiration. The only sound which exactly resembles it is one which is sometimes heard for a short time, under circumstances which render it probable that no disease at all exists, either in the lung or pleura. When, for example, a person who has been breathing quietly, is submitted to examination, and directed to take a deep inspiration, it not very unfrequently happens that at the very end of the inspiratory effort, there is heard, at the base of the lung, a very pure and perfect specimen of crepitation. But it is heard only three or four times. The examiner feels assured, if fever be present, that he has a case of latent pneumonia. If he have used his unguarded ear, to avoid fallacy, he puts his hand aside; if the stethoscope have been employed, he may apply the naked ear to the chest, to make "surety doubly sure;" or he may direct a pupil's attention to the sound, as a good example of what he may have been long searching for. But the pupil does not hear it. The examiner himself tries again, but he can hear it no more. It is gone, and pure respiratory murmur has supplied its place. I have not unfrequently been thus deceived myself; and my

pupils have also been disappointed of an expected prize.

What is the source of this sound? Present usually at the base of the lung, heard only at the end of a deep inspiration, and disappearing altogether after a few forced expansions of the chest and of the pulmonary cells, it appears at least probable, that it may arise from the unfolding, or the separation of the delicate walls of some pulmonary vesicles, which, from not being used in ordinary respiration, may have become somewhat dry, and may thence, like a dried bladder, give rise to the sound upon their insufflation.

The crepitation of emphysema is larger and looser, and more unequal, than that of pneumonia; as before stated, it most probably arises from the presence of serum, or of mucus in the dilated air-cells and the capillary tubes.

The attempt, however, to describe these minute differences of sound is almost useless. The sounds must be heard to be appreciated; no description, however exact, and however minute, can render them practically available to the uninitiated. Auscultation cannot be learned by book—except for the purpose of passing an examination. The student must go to the bed-side,—must go frequently, and stay long. Without this he cannot become a practical auscultator—without this he cannot become a skilful practitioner in diseases of the chest.

In pneumonia, then,—in œdema, in apoplexy,

and in emphysema of the lung,—crepitating rattles may be heard. In all, the sound probably arises from causes nearly similar, viz. the effusion of fluid into the cells of the lung, and the intermingling of the inspired air with that fluid. But in all of them, excepting in pneumonia, the crepitating rattle is almost always combined with some amount of muco-crepitation, arising from the presence of fluid in the smaller branches of the bronchial tubes. In pneumonia the sound may be likewise so combined, but it frequently occurs alone. It is the especial characteristic of inflammation of the pulmonary tissue; and if perfectly pure, equal, persistent, and existing only at the end of the inspiration, it may perhaps be still regarded as nearly pathognomonic of that complaint*.

Crepitating rattle then in its purest form, is

* I say this, though I am acquainted with the discussions which have recently taken place concerning the crepitating rattle, which is present in some cases of pleurisy after the fluid has been absorbed; and though I am perfectly familiar with the sound to which those discussions refer. That sound I have been accustomed to regard as the "*râle crepitant redux*," that is to say, as the genuine crepitation of pneumonia, which had been either previously obscured by the presence of effusion, or, after its temporary disappearance from consolidation or compression, had been reproduced, in the thin layer of pulmonary tissue, which is more or less inflamed in almost all cases of so-called "simple pleurisy." I am certainly not acquainted with any sound, liable to be mistaken for the pure crepitating rattle of pneumonia, which has in any case been proved to proceed from the pleura.

present in, and is especially characteristic of, the first stage of pneumonia, when the air-cells are still unclosed, but contain fluid secretion. When the cells are quite obstructed, either from the tumefaction of their vascular parietes, from the deposit of solid matter, or from both causes combined, all vesicular sound, all sound, that is to say, connected with the air-cells, ceases in the part affected; and tubular breathing, or bronchial respiration, supplies its place.

When the disease is on the wane, and the cells become capable of admitting air, the physical condition of the part is very nearly similar to that existing previously to consolidation; the physical signs are consequently similar. The crepitating rattle reappears, in the first instance in a very pure form; but, as the complaint gradually declines, the rattle becomes more and more loose and large, and therefore more and more resembles the muco-crepitating rattles of capillary bronchitis, which it also resembles in accompanying the act of expiration, as well as that of inspiration. This is the "*râle crepitant redux*" of the French stethoscopists.

When, instead of retrograding, the disease goes on to induration and contraction of the pulmonary tissue, tubular breathing becomes permanently established in the part, and if, in the progress of time, the bronchial tubes entering it become dilated, amphoric breathing, or, when fluid is present

in the tubes, gurgling, may be constantly heard therein.

When, instead of becoming indurated and contracted, the lung, in consequence of the unhealthy, unorganizable nature of the deposit, begins to break up from the softening of the inflammatory product, air is again admitted to the interior of the lung, and, by mixing with softened and semi-fluid matter, induces a sound which is very peculiar, and which, as far as my experience extends, exists only in that condition of the lung, and in a similar condition occurring in the progress of phthisis. This sound is neither crepitating, nor muco-crepitating rattle, nor gurgling, but it is a combination of the two latter sounds; or rather, it is the crepitating rattle altered in character by the thick viscid quality of the fluid into which the air passes, and by the increased conducting power of the surrounding consolidated mass. To the ear, indeed, it appears to be connected with precisely those physical conditions, and to arise from exactly that state of the respiratory organs, which actually exist; viz. the passage of air into a soft mass, such as the rotten lung, reduced nearly to the consistence of hasty-pudding or thick treacle. It is in fact a sharp, shrill, and loud muco-crepitating rattle, and may be not inaptly represented by the sound produced by passing the saliva to and fro, through the teeth with the mouth closed. It is, indeed, in

other words, a diminutive gurgling, or large moist crackling.

As the lung continues to break down, the bubbles become larger and larger, and the rattle, though still unusually shrill, assumes more and more of a mucous character; till at length, in consequence of the removal, by expectoration, of the semi-fluid mass constituting the disorganized lung, a cavity is formed, and amphoric breathing, and gurgling, supply the place of other sounds. It must, however, be recollected, that though a cavity be formed, and the physical signs of such cavity be sufficiently evident, it may be still surrounded by pulmonary tissue in various stages of degeneration; and that this disorganized tissue may give rise to many of the fluid sounds, previously mentioned, in addition to those of gurgling and amphoric respiration.

In the early stages of *phthisis*, the first *moist* sound that is heard is most commonly a fine and delicate muco-crepitating rattle, audible only at the end of the inspiration, and at the commencement of the expiration; it appears to be situated in the smallest branches of the bronchial tubes, and to be most probably dependent upon slight bronchitis, or increased bronchial secretion, the result of the irritation produced by the presence of tubercles.

This is commonly the earliest moist sound which accompanies the deposition of tubercles.

But it sometimes happens, that even previously to the appearance of this sound, there is heard in those parts of the lung in which the tubercles are well known to be almost always first deposited in the ordinary form of phthisis, a sort of click, which, from the nature of the sound, may be presumed to arise from the obstruction afforded to the passage of the air, by a portion of adhesive mucus, or other sticky substance, in a large tube. Even when this sound is present, which is not very frequently, it is only heard occasionally; that is, it is not heard at each inspiration, and perhaps not oftener than once in four or six inspirations. Its origin is doubtful, and as it is somewhat unusual, and therefore of comparatively little diagnostic importance, it is unnecessary farther to dilate upon it; excepting for the purpose of stating that it may be represented by the sticky noise produced by the removal of the tongue from the roof of the mouth.

As the number of tubercles increases, or more particularly as they become surrounded by pneumonia, the bronchitis and its accompanying physical signs increase also. The muco-crepitating rattle becomes more general, and also more shrill; as it is now better conducted by the greater or less consolidation of the surrounding tissues. It is also sometimes mixed with the genuine crepitation of pneumonia, and the mucous rattle of the larger tubes. From this stage of

the disease, the moist sounds of phthisis so generally resemble those which occur in the unhealthy form of pneumonia, accompanied with serofulous, or unorganizable deposit, which sounds have been just described, that it is scarcely necessary that they should be again mentioned. It may be, however, stated, that the shrill muco-crepitation, the loud mucous rattle, and the gurgling, succeed each other with varying celerity in different cases, and are variously combined accordingly as smaller communications with the bronchial tubes, or larger cavities, are produced in the destructive progress of the disease. It may also be not irrelevant to remind the student, that when little, or scarcely any, indication of disease can be discovered below the clavicle of either side, a cavity of moderate size may be present in the very apex of one or both lungs, the *physical* evidences of which can be detected only by applying the stethoscope at the side of the neck above the clavicles, or more especially at the posterior and highest part of the acromial region.

5. *Moist Sounds in diseases of the Pleura.*

It has been already stated, that when fluid exists in the pleura, whether the quantity be large or small, unless air be likewise present, no moist sounds are produced, either by the act of

inspiration, or by the motions of the body. It may, however, perhaps be questioned whether the sticky noise which results from the friction of recently effused plastic matter, or the somewhat similar sound which arises after the fluid portion of a pleuritic effusion has been in a great measure absorbed, may not be fairly considered to constitute an exception to this general statement; as the impression given to the ear is certainly not that of the friction of perfectly dry substances. It appears at least doubtful, if anything more nearly approaching to a moist sound than this ever arises from simple pleurisy, and it is exceedingly questionable, if sounds simulating the mucous, and even the crepitating rattles, have been ever demonstratively shown to originate from fluid, or from any other, effusion into the pleura, without the presence of air.

When, however, a communication exists between one or more bronchial tubes, and a defined cavity in the pleura, and when this cavity is of moderate size, and is occupied partly by purulent, or sero-purulent fluid, and partly by air, the *physical conditions* are very nearly similar to those, in which a cavern, resulting from phthisis or pneumonia, is present in the substance of the lung, and is situated close to the surface. The *physical signs* are therefore similar also. In each case fistulous, or amphoric respiration may be observed; in each case metallic ringing or reso-

nance of the breathing may exist ; and in each case the examiner may be able to detect, either separately or in combination with the other sounds, the large, thick, sonorous mucous rattle, called gurgling.

One other fluid sound remains to be considered. It is most commonly present in conjunction with disease of the pleura ; but it is also, though rather rarely, found in connection with cavities in the lungs, without any especial affection of the serous membrane. When, in consequence of a communication with a bronchial tube, a large space in the pleura contains a large proportion of air, and also a small quantity of fluid ; or when a phthisical cavity of very large size is similarly circumstanced, there is every now and then heard a very peculiar sort of tinkling noise upon examining the chest. It more nearly resembles the sound caused by shaking a pin in a decanter, than any other with which I am acquainted. This is *metallic tinkling* (“*tintement metallique.*”) It is most commonly heard only at intervals—that is, it may occur once in three, four, or forty inspirations. It rarely, if ever, attends the expiration. It may cease altogether, and reappear after a considerable time. In this respect it seems not unfrequently to be influenced by the position of the patient's body.

It is most probably produced by the continued and rapid reverberation of a delicate sound against

the firm and vibrating walls of a large cavity—it is an an echo in a small space. The original sound from which the echo proceeds appears most commonly to arise from the bursting of a bubble of air, or from a drop of liquid falling, upon the surface of fluid in the bottom of the cavity. But it sometimes seems to be likewise produced by the passage of air over a loose portion of *membrane*, or *thick secretion*, situated in a tube at the entrance of the cavity. The physical conditions necessary for its production appear to be a large cavity with resonating walls, and containing a large portion of air, with a small quantity of fluid. When, as occurs in many, perhaps most, protracted cases of pneumothorax, the cavity in the pleura previously to death becomes in a great measure filled with fluid effusion, and the air is consequently displaced or absorbed, the metallic tinkling disappears, and, unless the fluid so accumulated be evacuated, permanently ceases.

Let the student, however, beware that he is not in this respect, as well as in those before mentioned, deceived by the stomach, which, containing a mixture of fluid and gas, and passing high up beyond its ordinary boundaries, sometimes so far encroaches upon the chest as to give rise to physical signs, many of which exactly resemble those of pneumothorax, and among which a sound very nearly approaching to, if not precisely identical with, *metallic tinkling*, is occasionally heard.

7. Auscultation of the Voice and Cough.

1. In diseases of Air-passages and the Lungs.

The changes produced by disease in the auscultation of the voice in many respects resemble those which have been noticed in reference to the breathing. Thus, it may be generally stated, that whatever increases the power of conducting sound in the thoracic organs, which are connected directly or indirectly with the larynx, also increases the resonance of the voice, heard through the parietes; just as it increases the loudness and shrillness of the morbid sounds which accompany the respiration. But as the voice in disease, as well as in health, is formed by the passage of air from the lungs, or by the expiration; and as many, if not the majority, of the morbid sounds which accompany the breathing are produced by the passage of air to the lungs, or by the inspiration, many and important alterations, affect the one, which are in no respect applicable to the other. Thus, mucous rattles may be heard throughout the whole chest, and the voice may remain perfectly natural; and the voice may be quite inaudible, and yet the respiratory murmur may continue pure in every part. Sounds also which accompany the respiration may, and often do, originate in the pulmonary tissue itself, and

exist independently of any sound arising in, or conveyed by, the bronchial tubes; but these tubes, as regards the voice and the cough, are, both in health and in disease, the principal and primary channels by which the sound is conveyed to the parietes.

In the *diseases of the larynx* the alterations of the voice which exist in ordinary conversation are also observable upon the application of the ear to the chest, or of the stethoscope to the side of the neck. The only assistance afforded by auscultation, in disease of the larynx, with which I am practically conversant, is, that it enables us to ascertain the exact seat of the disease, or of the obstruction which gives rise to the alteration, or to the loss, of the voice. Of the nature of that disease, or of the character of that obstruction, it telleth not. The same acknowledgment was made while treating of the respiration.

As in ordinary *bronchitis* the texture and consistence of the lung is unchanged, the voice, for the most part, remains unaltered. But when, in consequence of long-continued bronchial obstruction, the lung becomes considerably congested with blood, or infiltrated with serum, the organ is rendered comparatively solid. Commensurately with its degree of solidity it is also rendered a better conductor of sound. The voice, therefore, and the cough are, in the parts so affected, louder than in other parts of the chest. Those parts are

ordinarily at the base of the lung posteriorly, as they are generally the most dependent parts, and the fluids, consequently, have a tendency to accumulate therein from gravitation.

When the bronchial tubes are generally, but at the same time slightly and simply dilated, as a result of old bronchial obstructions, the resonance of the voice is often only very slightly, or not perceptibly, increased; as not only are the parietes of the tubes themselves less thick and dense than in the normal condition, but the surrounding pulmonary tissue is often more or less affected with emphysema; the necessary consequence of which is, that the voice is less perfectly conducted to the parietes than in the natural state of the organ. But when dilatation is partial, and at the same time extensive; when, that is to say, the dilatation affects one, or only a few tubes, in a defined portion of its, or of their course; when, in fact, one or more circumscribed cavities are formed by the dilated tubes; the resonance of the voice may be considerably increased over the part in which such dilatation exists; and if it be near to the surface of the lungs, and particularly if it be surrounded by consolidated tissue, the voice may be not merely increased in resonance, but the increase may amount to *bronchophony*, or even to perfect *pectoriloquy*. These terms have already been explained and illustrated, while speaking of the natural voice (see p. 103). It

is therefore not necessary here to repeat the account of them. But if the student have any doubt of his capability of discriminating between them, let him turn to that page, and repeat the experiments there recommended to his notice.

In simple emphysema of the lung, the density of the organ is diminished, and its conducting power consequently decreased; the resonance of the voice, therefore, as heard through the parietes of the chest, is proportionally lessened.

When the lung is consolidated from pneumonia, its conducting power is increased; and if the tubes passing through, or in the vicinity of, the consolidated mass, be of tolerable magnitude, as in the upper part, or central portions of the lung, the voice is more easily transmitted, and *bronchophony* is consequently heard. But if the consolidation exist only at the edges of the lung, removed a considerable distance from any tube of large calibre; or if, though in the neighbourhood of large tubes, those tubes are closed by compression from without, or by secretion from within, the voice is not only not increased in resonance, but it may be less loud than natural; as the tubes, which are the principal conductors of the voice from the larynx to the parietes, do not, under such circumstances, in consequence of the absence of air, vibrate with sufficient power to affect the solid walls of the chest. Thus, at the angle between the diaphragm and the ribs, perfect con-

solidation of the lung may exist without any increase of the natural resonance of the voice. Thus also when, as sometimes, though rarely, happens, the tubes are plugged up with plastic secretion, while the lung itself is converted into an airless mass, the voice may be scarcely audible through the parietes.

In persons who, being either constitutionally feeble, or having been debilitated by disease, have been confined for a long time to one position in bed; as in the case of fracture of the old, and in typhoid fevers of all ages; blood accumulates in the more dependent parts of the lung, the vessels of which become dilated, and the tissue of which is rendered less elastic, and at the same time soft and feeble, from the increase of fluid, and the comparative deficiency of air. From long continuance of this state of passive congestion, it is not improbable that structural change may sometimes occur, and that a low form of pneumonia may be induced, similar to that, which, occurring in the last few days, or hours, of life, is called by the French pathologists "*pneumonie des agonisans*," or the pneumonia of the dying. But whether the changes induced in the lung amount in any, or in most, cases to pneumonia, it is certain that the density of the lung is increased, that a certain degree of consolidation is produced, that its power of conducting sound is improved, and that, as a direct consequence, the resonance of

the voice over the part becomes so much greater than natural as not unfrequently to amount to perfect bronchophony.

The preceding observations are, in the majority of instances, applicable to that condition of the lung which ordinarily exists in pulmonary apoplexy. They are, indeed, perfectly relevant, as regards the state of the lung when blood has not escaped into the pulmonary tissue. But while the passive congestion, arising from debility, always exists in the most dependent parts of the lung, that resulting from obstruction, as from diseased heart, or other similar causes, may exist in any part; though it is most frequently found in those portions of the lung which are the farthest removed from the central organ of the circulation.

When the tissue of the lung is broken down, or its cells are filled with effused blood, when the bronchial tubes are not compressed or plugged up, and when the part of the lung affected is favourably situated for the conduction of the voice, the bronchophony, which is present in consolidation from other causes, is also present in that resulting from pulmonary apoplexy. But, as in both these respects, the lung is often unfavourably situated; as the part affected is frequently the most distant from the larynx, as well as from the heart; and as the tubes are often compressed or obstructed by the effused blood; increased resonance of the voice is not so common

in this affection as in the consolidation of the lung from other causes.

When, in consequence of the destructive progress of unorganizable pneumonia, a cavity is formed in the lung, the state of the voice varies according to the seat of the cavity, and the condition of the surrounding textures. If it be at the thin edge of the lung, if the tubes leading to it be small, and if the pulmonary tissue, intervening between it and some one or more large tubes, be not consolidated, the voice may be scarcely increased in resonance. If, on the contrary, it be in the upper part of the lung, or near large tubes; or if it be close to the surface of the chest, or surrounded by consolidated pulmonary tissue, then either bronchophony, or imperfect, or even perfect pectoriloquy will be present, according as the situation, and the condition of the diseased parts are more or less favourable to the transmission of the voice. Pectoriloquy by no means necessarily exists under such circumstances. There may be a cavity in the lung without pectoriloquy; as pectoriloquy may be present without a cavity.

When, instead of being softened, and broken down, and by expectoration reduced to a cavity, a portion of lung affected with pneumonia becomes dense, firm, and contracted, and by adhesions is rendered almost one with the parietes; and when into this portion of the lung there enter large bronchial tubes, whether the large size of the tubes be natural or the result of disease, the

voice is often rendered exceedingly loud indeed. When, in addition to this state of parts, the disease is situated near to the larynx, the whole of the circumstances contribute to increase the vocal vibration. The resonance not merely equals that existing when the patient speaks into the tube, or when the tube is placed upon the larynx; it is not simply pectoriloquy; but the voice becomes so shrill and jarring as to be actually painful to the examiner.

In *malignant disease of the lung*, the abnormal deposit is sometimes, though comparatively rarely, diffused through a portion of the pulmonary tissue, without materially compressing the bronchial tubes. Under such circumstances (of which I have only seen one example), the voice is rendered more loud than natural, commensurately with the amount of consolidation of the part, its permeation by large bronchial tubes, and its vicinity to the larynx. At other times the malignant mass encroaches upon the lung, and, instead of invading its texture, pushes it aside, and with it the bronchial tubes by which it is traversed. Under such circumstances there exists a solid mass, containing no conducting pipe from the larynx, or, in other words, no prime conductor; the resonance of the voice, therefore, is not only not increased, but that naturally existing in the part affected, is diminished, and sometimes even lost. According to its position and arrange-

ment, then, malignant disease of the lung may be accompanied with increased, or with decreased, resonance of the voice.

In the earlier stages of *phthisis* the degree of resonance of the voice depends upon the amount of tubercular deposit, the consequent density of the lung, and its resulting conducting power. Even in very early stages there may be sometimes heard a slight increase of vocal resonance in the acromial or infra-clavicular regions of one side over those of the other side. The student, however, must be here especially vigilant against error. He must recollect, that in many, and especially in thin persons, the voice is naturally very resonant below the clavicles, and that it is sometimes slightly more resonant upon the right than upon the left side. He should, therefore, never form an opinion in consequence of the resonance of the voice in the superior parts of the chest on one side, without comparing it with that of the other. He should also never forget that greater resonance may exist upon one side than upon the other without disease; and that this increase, which is independent of disease, *usually* exists upon the right side. Unless, therefore, the increased resonance of the voice be confined to one side of the chest; unless it be combined with other physical signs, or with obvious constitutional symptoms; or unless it be of so decisive a character as not to be satisfactorily explained by natural conforma-

tion, it should not be regarded as an evidence of disease.

As the consolidation of the lung increases, whether by the greater number of tubercles, or by attendant pneumonia, the voice becomes more and more resonant, and bronchophony, or imperfect pectoriloquy, is observed in the part affected. The farther it is from the larynx, the less likely is it to be the result of natural conformation; the more distant it is from the larger divisions of the trachea, the more probably does it result from disease; and therefore the more important is it, as an indication of disease at the top of the scapula, than on its inner edge, and near to the acromial process, than close to the sternum.

When, in the progress of the complaint, cavities are formed in the lung; as those cavities are usually near to the apex of the lungs, and therefore not far removed from the larynx; as they are almost universally surrounded by consolidated pulmonary tissue, which is adherent to the costal pleura, and which therefore becomes one firm mass connected with the parietes; and as, moreover, these cavities are generally near to the surface of the lung, the whole arrangement of the parts is most favourably circumstanced for the conduction of the voice. Pectoriloquy, perfect or imperfect, is accordingly a very frequent sign of a tubercular cavity.

But though a frequent, is pectoriloquy a certain,

sign of a cavity? It has been already stated that it is not. But in a tubercular subject, one evidently suffering from advanced phthisis, may it not be so regarded? No! Well, but if tubular breathing, and contraction, or flattening of the part, exist at the same time, is not pectoriloquy a proof of a cavity? No! not even then. All this, and much more, may exist, and yet the presence of a cavity in the lung cannot be with certainty predicted, even though the part affected be the apex of the organ. Even then it is merely a question of probability, as all these signs may arise from a consolidated and contracted lung, combined with dilated tubes. This condition of the organ, as has been already stated in reference to pneumonia, may not merely give rise to perfect pectoriloquy, but may render the voice more loud, shrill, and ringing, than when the patient speaks into the trumpet-shaped end of the cylinder. The truth is, that pectoriloquy, that sign which was formerly regarded, and is still esteemed by some, who are not practically conversant with auscultation and with morbid anatomy, as the especial and sure sign of a cavity, is now well known, by those who are familiarly acquainted with the revelations of the inspection-room, to be among the most uncertain of the physical signs afforded by auscultation.

When a cavity in the lung is of very large size, the voice, like the breathing, sometimes possesses a peculiar ringing or metallic resonance,

in consequence of the reverberation of the sound against its solid walls.

The observations made in reference to the voice are also almost literally applicable to cough. The only remark, indeed, that it appears necessary to make, in reference to the resonance of the cough, in contradistinction to that of the voice, is, that the former can sometimes be heard when the latter is inaudible, and that the act of coughing sometimes causes the voice to be heard in a part in which it had been previously inaudible. Thus, in the signs afforded by the breathing, as well as by the voice, a tube may be obstructed by thick mucus, or other secretion, which, by impeding the passage of air into the diseased part, may prevent those signs appearing which depend upon the respiration, and greatly modify those which are connected with the voice. Under such circumstances, if the patient be desired to cough frequently and forcibly, the obstruction is often removed, and the signs are developed, if they had not been previously observed, or reappear, if they had temporarily ceased. It occasionally happens, also, that the cough is resonant when the voice remains unaffected, or is affected only in a slight degree. Upon the cause of this peculiarity it is not always easy to decide; but as it is not removed by the repeated effort of coughing, it appears not to be dependent upon the cause previously referred to.

2. *Auscultation of the Voice in diseases of the Pleura.*

When a small amount of effusion is present in the pleura, whether that effusion be solid or fluid, or whether it be the result of inflammation, or of obstruction, and when that effusion is thinly spread over the surface of the lung, the spongy tissue is compressed commensurately with the amount of that effusion. The compressed lung, and the effusion being singly, as well as in combination, better conductors of sound than the pulmonary tissue in its normal condition, the sound of the voice is then conveyed more perfectly and distinctly than in the natural condition to the parietes of the chest.

But it, moreover, frequently happens, under such circumstances, that the voice is not only rendered increasingly loud, but that it is curiously altered in tone and quality. It often becomes remarkably shrill, trembling, or bleating. The tone is supposed to resemble that of the bleating of a goat; it is therefore called *egophony*, and the voice is said to be *egophonic*. At other times the voice so conducted has a peculiar nasal twang, very similar to that of the *fantoccini*. It is hence appropriately termed Punch's voice ("*voix de polichinelle*.") On other occasions, the voice resembles that of a person speaking through a trumpet.

The causes of these peculiarities of the voice are not very evident. They are, however, supposed to be produced by the trembling vibration of the fluid modifying the sound passing through it. Upon what the *differences* of the ægophonic sound depend, seems at present to be quite inexplicable; nor, unless the actual condition of the parts at the time could be clearly ascertained, does it appear probable that they will be ever clearly and demonstratively explained. The whole of these sounds are by some called *ægophony*, or considered as modifications of *ægophony*. If the whole be considered as coming under that designation, ægophony is of very frequent occurrence; but pure ægophony, the true goat-bleating noise, is exceedingly rare. It may be perhaps regarded as among the most uncommon of auscultatory signs.

Sometimes the voice possesses a tone and quality not precisely similar to, or rather not exactly amounting to, either of the characters which have been just mentioned. It is increased in power; it is louder than natural; but it is not simple bronchophony; it has a different tone from the natural voice; it possesses greater shrillness, but the shrillness does not possess the qualities of either of the varieties of ægophony. It partakes of the character of both, but does not actually resemble either of the two. The tone of the voice under such circumstances, has been

not inappropriately termed *ægophonic bronchophony*. The condition of the organs of respiration, with which this character of the voice is associated, is uncertain. It is not improbable that it may be connected with slight consolidation, together with effusion.

Is, then, ægophony a certain sign of effusion? Far from it. So far, indeed, is this from being true, that as ægophony of the pure bleating kind is among the rarest, so ægophony of every sort or kind is certainly among the most uncertain of the physical signs, considered as a diagnostic of effusion. Ægophony, ordinarily so called, may, it is at least probable, exist without effusion into the pleura, and effusion into the pleura it is certain may be present without ægophony. On no occasion, therefore, should ægophony be regarded as positively indicative of effusion.

The situations in which it is most commonly heard, are the infra-scapular, and the inter-scapular regions. But in these regions, ægophony is in some persons said to exist independently of disease. Hence arises the very important practical rule, that ægophony should never be regarded as of any importance, and particularly of any value as diagnostic of effusion upon one side of the chest, unless it have been previously ascertained whether it is also present upon the other side. It is, indeed, true that fluid may be present, and give rise to ægophony upon both sides. This

in active disease is not very frequent. But when it occurs, its presence must be determined by the employment of other means, than by the auscultation of the voice.

As the quantity of fluid in the pleura increases, the ægophonic character of the voice gradually disappears. Bronchophony for a time supplies its place; but as the fluid accumulates, the lung, and with it the bronchial tubes, are removed farther and farther from the parietes, so that the resonance of the voice gradually decreases, till it becomes almost perfectly inaudible.

An exception of course exists, in this respect, in reference to the inter-scapular region, in which are situated the large bronchial tubes at the root of the lung; as the greater the amount of fluid, the more closely are they commonly pressed against the parietes. In this region, therefore, the voice is ordinarily even more distinct and loud than in the natural condition, at the time that it is scarcely audible in other parts of the same side of the chest. The exceptions, which were referred to while speaking of bronchial respiration as the result of effusion, and of its disappearance under ordinary circumstances, when the fluid accumulated beyond a certain extent, are likewise applicable to the voice.

The very curious and rare exception to the general rule of tubular breathing and vocal resonance disappearing, except in the inter-scapular

region, when fluid exists in large quantity in the pleura; the fact, I mean, of both these sounds being sometimes heard all over the side of the chest, when a large amount of fluid has been proved to be present in the pleura, was once observed in reference to the voice and not to the breathing. In one case, that is to say, the voice was resonant all over the side of the chest, from which seventy-two ounces of fluid were drawn off by paracentesis, though the tubular breathing was only heard in the inter-scapular region.

Another curious fact in reference to effusion may be here referred to. It sometimes happens that the voice appears more shrill to the ear than in the natural condition, while the vibration of the parietes, as appreciated by the hand, is less than natural, or even entirely lost. *Tactile vibration*, in other words, is absent, though the *vocal resonance*, as appreciated by the ear, is, or at least appears to be, increased; and yet both are communicated by the vibration of the same parietes. The cause of this, which I have on several occasions verified, I must confess myself totally at a loss to explain. I have not even a remote conception of the manner in which it is to be explained, though of the fact, I believe, there can be no doubt.

When, in consequence of a communication existing between the pleura, and a bronchial tube, a tubercular cavity, or a portion of gangrenous

lung, air is present in the serous membrane in considerable quantity, the voice and the cough possess the same ringing or metallic resonance which has been already noticed, while speaking of the breathing in pneumothorax. It usually accompanies both the voice and the cough; but it is sometimes observed to attend the one, and not the other. Under such circumstances, it is generally, I believe, the cough which possesses, and the voice which is destitute of, the metallic character. This probably arises from the greater muscular effort exerted, and the consequently greater vibration excited in the air contained in the cavity, by the act of coughing, than by that of speaking. The character of the sound itself it is unnecessary to attempt to explain, or to illustrate farther than it has been already done. It should be heard to be properly appreciated. When it has been once heard, it is scarcely possible that it can ever after be mistaken.

6. *Auscultation of the Heart.*

While speaking of the auscultation of the respiratory organs, it was stated that it was necessary for the student to be acquainted with the signs existing in the healthy state, previously to commencing the investigation of the physical effects of disease. This previous knowledge is more especially necessary in reference to the heart; as without clear and definite notions of

the action, and familiarity with the sounds, of the organ in health, it will prove utterly futile for him to attempt fairly to appreciate the physical signs of its diseases. A brief, and, as far as practicable, a simple account, will therefore first be given of the auscultatory signs afforded by the heart in the healthy condition, and afterwards will be made a short exposition of those which occur when the organ is diseased.

1. *Auscultation of the Heart in health.*

The physical circumstances which are to be observed in reference to the auscultation of the heart are included under its *Impulse*, its *Rhythm*, and its *Sounds* *.

* It may be right to state that I have never either myself made, or have been present while others have made, any experiments upon the lower animals, in reference to the physiological action of the heart; and that my information, so far as regards observations upon the uncovered heart, is derived entirely from the statements and experiments of others. But while acknowledging my obligations to the experimenters for the *facts* which they have established—(if, indeed, they may be now considered to be established), I must with all deference crave permission entirely to differ from some of them in their *reasonings* concerning the cause of the heart's sounds.

The opinions, in reference to the sounds of the heart, adopted in the text, are those which have been ably advocated, and were, I believe, first promulgated in this country by Dr. Billing, and in France by M. Rouanet; though, considerably after their promulgation by these gentlemen, they were adopted by myself quite independently. They have ever appeared to

The Impulse of the Heart.—When the expanded hand is placed upon the left side of the chest below the nipple, or more particularly when the points of the fingers are placed upon the space between the fifth and sixth ribs, a little to the inner side of a line running vertically over the nipple, a gentle and regular pulsation is felt, and when the ear or the stethoscope is applied to the same part, a slight shock is perceived by the examiner. This is the *Impulse*, or the stroke of the heart against the parietes. It is nearly or quite synchronous with the pulse at the wrist, and with the systole or contraction of the ventricles. In

me to be capable of explaining, so far as it is at all explicable, all that is observed in reference to the sounds of the heart either in health or disease. They have never been shaken, in the slightest degree, by the conclusions of the experimentalists; as I could never bring myself to believe that by taking out the heart of dogs and donkeys, cutting out the valves, skewering them up with bradawls, and suspending them with fish-hooks, &c. &c., they were at all likely to arrive at a correct explanation of the *natural* sounds of the human heart. Though, therefore, I am quite aware that, without any experiments of my own, I should hesitate before I take the liberty of differing from the wiser and more learned gentlemen who have made many experiments to illustrate the subject, I feel at the same time that I should neither fulfil the object of this work, nor my duty to those who favour me with a perusal of it, if I did not upon this question, as upon all others, simply state the result of my own observations and reflections, without feeling myself called upon to state, upon every occasion, the why and the wherefore, cavilling at the opinions of others, or answering probable objections to my own.

the healthy adult it occurs from sixty-eight to eighty times in the minute. It has been variously explained at different times, but is now generally believed to arise from the apex of the heart striking the parietes. A tilting forwards is produced by the contraction of the ventricle, which causes the apex of the heart to move in the section of a circle, of which the base of the heart is the axis or centre, and to strike against the ribs: hence arises the Impulse.

The part in which the Impulse is felt, in the healthy condition of the organ, varies considerably, according to the position of the body. While it is erect the impulse is felt about an inch to the inner side of a line running vertically over the nipple. When the individual examined lies upon his back it is decreased in force, and, at the same time, felt about half an inch nearer to the median line of the body. When the trunk is turned directly upon the left side, the impulse is perceived either immediately below, or sometimes considerably to the outer side of, the nipple; while, when the trunk is turned directly upon the right side, it is either not perceptible at all, or only just perceptible between the cartilages of the ribs close to the sternum.

The Impulse of the heart, like the pulse at the wrist, is naturally much more frequent in some individuals than in others. It also, quite irrespectively of disease, varies very considerably in

character. It is full and powerful in the strong and robust; while in the weak and delicate it sometimes amounts to scarcely more than an undulation, the precise situation of which it is occasionally difficult to determine. In the vigorous but calm individual, it is strong, heaving, and diffused; in the weak, nervous, and excitable, it is smart, smacking, and very defined, or as it were concentrated. When, indeed, the organ itself, and the arteries leading from it, are healthy, the Impulse of the heart resembles in character and in power, as well as in frequency, the pulse at the wrist. The precise *situation* of the Impulse may be best defined by the fingers, or by manual examination; though by those who have already become accustomed to the use of the stethoscope, its *character* and *power* may generally be best appreciated by the sensation communicated thereby to the side of the head. In one case, for example, it may shake the head scarcely at all; in others it may effect a sharp knocking stroke; while in a third class of cases it may, without knocking, lift up the head, or communicate to it a heaving sensation.

Here it may be desirable to caution the student against confounding together a sharp or smacking Impulse, and a forcible or heaving one. This is particularly necessary, as the former, by those who are little conversant with the matter in hand, is not unfrequently, or is, perhaps, generally,

denominated a strong impulse, and is most incorrectly supposed to indicate a thick powerful heart; whereas it almost always characterises the very opposite condition of the organ. A sharp, quick, smacking Impulse, merely indicates an irritable heart, and *usually* a weak one—like a pulse of the same character;—whereas it is a deep, heaving Impulse, like a large, full, and incompressible pulse, which accompanies a powerful or hypertrophied heart.

The Rhythm of the Heart.—If the ear, or the stethoscope, be placed upon the præcordial region, and the heart be acting naturally, a regular succession of sounds, and cessation of the sounds or of intervals, will be noticed by the most casual observer. In the first place will be noticed a sound comparatively long, then a sound comparatively short, and then an interval without any sound; after which, again, will occur the long, then the short sound, and then the interval. The long and short sound, together with the interval, make up the time occupied by one complete circuit of the heart's function; and as these sounds and this interval have been observed constantly to bear a definite relation to each other, the period occupied by them, or the complete circuit of the heart's action, has been divided into corresponding portions of time.

Supposing, then, that the whole period of this

time be divided into fifths,—the first sound has been calculated to occupy two-fifths, the second sound to occupy one-fifth, and the interval between the sounds to occupy the remaining two-fifths. The period occupied by the whole corresponds to the time between one stroke of the pulse and another. So that if a person's pulse beat eighty times in a minute, there would be eighty long sounds, eighty short sounds, and eighty intervals. Similar relations would exist, supposing that they could be correctly estimated, and that the pulsations were at the same time regular, if there were one hundred, or a hundred and sixty strokes of the pulse in a minute. This regular division of the time occupied by one complete circuit of the heart's function constitutes the *Rhythm* of the organ.

This may be perhaps termed the external, or obvious rhythm ; as there exists, also, an internal rhythm, upon which the external one depends, and of which, therefore, it is the indication.

It has been observed, while watching the uncovered heart of the lower animals, that after the interval, or repose of the heart, the contraction of the appendix of the auricles first occurs, and is immediately followed by the systole of the two ventricles, which contract simultaneously. At the same time occur the first sound, the distension of the aorta, and of the pulmonary artery with blood, comparative fulness of the auricles, and the tilting forward of the apex of the heart. Immediately after the systole or contraction of

the ventricles, follows the sudden diastole or dilatation of the ventricles. At the same time occur the reaction of the distended aorta and pulmonary artery upon their contained blood, the second sound, and the comparative flaccidity of the auricles. Then occurs the pause or interval between the sounds. The ventricles are meanwhile becoming full, but not distended with blood; the appendices of the auricles again contract; a small additional quantity of blood is sent into the ventricles, which are supposed to become thereby distended, and by such distension to be excited again to contract, when the series of actions takes place as before.

Such are, very briefly, the actions, as related by experimentalists, upon which the *Rhythm* of the heart depends, and such, therefore, is what may be denominated its *internal Rhythm*.

When the sounds of the heart, and the interval between them, bear the relation to each other which has been mentioned above, the rhythm is said to be *natural*: when they do not bear that relation to each other, the rhythm is said to be *unnatural*, and the action of the heart is said to be *irregular*. Irregularity of action, then, is the same thing as an unnatural rhythm.

When the pause, or interval between the sounds, is longer after some contractions than after others; and when this increased duration of the pause or interval occurs at regular periods, whether it be every third, sixth, or sixteenth beat of the heart,

the action of the organ is said to be *intermittent*. But when, though the sounds and intervals bear the same relative proportion to each other as in the natural condition, one stroke of the heart is observed to be more powerful than the stroke, or than the four, six, or any other number of strokes, which precede and succeed it, the action is said to be *unequal*. *Irregularity* of the action of the heart, then, depends upon an alteration in the natural succession of sounds and intervals, and is connected with the Rhythm. *Intermittence* is the regular recurrence at stated intervals of that irregularity, and is also connected with the Rhythm. *Inequality* exists when one stroke of the heart is stronger than another, or than several others, and is connected with the *Impulse*. It is, in fact, an *irregularity of the impulse*.

Is, then, a natural rhythm indicative of a sound heart, and an unnatural rhythm necessarily characteristic of an unsound one? Both these questions must be answered in the negative.

A heart greatly diseased, so long as the circulation is undisturbed, and the patient remains quiet, mentally as well as bodily, may pulsate with perfect regularity. The rhythm of a merely irritable heart, as that of an hysterical girl, a dyspeptic, a valetudinarian, a hypochondriac, or of a person greatly alarmed or excited, may exhibit all sorts of irregularities. The Rhythm, therefore, considered alone, is of little diagnostic im-

portance, as the mere fact of making an examination for the purpose of ascertaining its character may, in a nervous subject, alter it altogether.

The Sounds of the Heart.—The natural sounds of the heart, the regular recurrence of which has been already referred to as constituting the Rhythm, may be fairly represented by the following syllables, *too-to—too-to*.—The origin of these sounds has been the source of much discussion, and the object of a great variety of experiments. The student may see an account of them in most systematic treatises on diseases of the heart.

It does not consist with the design of this work to enter into a critical examination of this or of similar questions. It may be simply observed, that the opinions hereafter expressed, are those which appear, and ever have appeared, to the author to be the most probable. It is entirely opposed to the object I have in hand to engage in any arguments in support of those opinions, or conjecturally to answer any objections which may be possibly raised against them.

I proceed, then, at once to inquire what is the efficient cause of the two sounds of the heart? In answering this question it will be convenient, first to consider the origin of the second sound, as being the more simple, and *now* admitting of less dispute; and subsequently to consider the

cause of that which precedes it, concerning which there still exists much diversity of opinion.

The origin of the second sound.—If the cock of a leaden pipe, which is not firmly fixed to its solid support, be turned, and a full stream of water be allowed to pass through the aperture which that cock controls, little agitation of the pipe is induced by the mere passage of the fluid. If, however, while the water is passing with rapidity and force, the cock be suddenly turned, and the passage of the fluid through the aperture be in consequence suddenly arrested, a very considerable agitation of the pipe will be immediately perceived, and a corresponding noise will be heard. This agitation, and this noise, are clearly the consequence of the vibrations excited in the fluid by the sudden arrest of its onward progress; which vibrations are communicated to the solid walls of the pipe, and thus become obvious to the eye, as well as to the ear. Another illustration may be taken, from what is seen to occur at the closure of a lock on a river. So long as the water continues to flow through the constantly narrowing opening between the two arms of the lock, there is no agitation in the back water. But immediately the two arms meet, a violent shaking is observed in the wood of the lock itself, and in the water which is behind it. This is evidently produced by the sudden stoppage of the onward progress of the stream.

Now the circumstances attending the second sound of the heart are precisely similar to those which are observed in the pipe and in the lock; excepting that the former are connected with living and elastic tissues, and the latter occur in connection with comparatively inelastic and dead matter. The blood is, by the contraction of the ventricles, propelled into the aorta and pulmonary artery, and these vessels become thereby distended. When the pressure from behind is removed by the sudden diastole of the ventricles, the large vessels react (whether by actual contraction, or by mere elasticity, does not affect the present question), powerfully react, upon the fluid which distends them. The blood is forced back towards the ventricles. Some of the fluid, it is almost certain, must re-enter the cavities. But its retrograde motion is suddenly arrested by the flapping back of the sigmoid valves. These flood-gates suddenly stop the backward progress of the blood; and this sudden stoppage of the forcible current, as in the case of the pipe and cock, or in that of the lock, produces a violent agitation in the particles of the blood, which is communicated to the coats of the vessel, and thence to the parietes of the chest. *Hence arises the second sound of the heart.*

Origin of the first sound.—The first sound of the heart appears to arise from causes which are, *mutatis mutandis*, almost precisely similar to

those which produce the second sound. If the ventricles be put for the aorta and pulmonary artery, the auriculo-ventricular for the sigmoid valves, and positive for probable contraction, or elasticity, then one description might be almost literally applied to both. When the contraction of the ventricles commences, the passage to the auricle, it must be recollected, is at least partially open. There is a space open for the blood to pass into the auricles as well as into the large arteries. Some blood, it may be pretty confidently asserted, at first does actually pass back into the auricles; though the question, whether a small quantity does or does not pass, does not affect the theory of the sound. The blood, strongly compressed by the forcible contraction of the ventricles, has at least a tendency to pass into the auricles as well as into the aorta and the pulmonary artery; but the flood-gates at the entrance shut the opening; its backward progress is suddenly stopped by the auriculo-ventricular valves. This sudden stoppage of the blood, as in the case of the lock upon a stream, produces a violent agitation in the fluid, which is communicated to the body of the ventricles, and thence to the parietes of the chest. *Hence arises the first sound of the heart.*

The difference in the tone, and in the duration of the two sounds, appears to be sufficiently explained by the different constitutions of the parts

engaged in producing and conducting them. The size of the auriculo-ventricular valves, and the thickness of the ventricles, compared with that of the valves, and of the coats of the vessels; and the irregularity and great inequality of the surface, the muscoli pectinati, the carneæ columnæ, and tendinous cords of the ventricles, compared with the smooth lining of the two great arteries, appear amply sufficient to account for the difference.

Natural range of the Heart's sounds.—It is exceedingly difficult indeed, to define the natural limits of the sounds of the heart; as in scarcely any two individuals are they precisely similar. There are so many circumstances, entirely extraneous to the heart itself, which tend to increase or diminish the area over which they are heard, that scarcely any defined line can be strictly said to include it. The sounds, for example, are heard more extensively in a thin, than in a fat person; in an individual of diminutive size, than in one of large bulk; and in a person of a naturally irritable disposition, than in one of calm temperament. In some excitable females, indeed, and in males who approach them in character and configuration, the sounds of the heart may be heard over almost the whole thorax; while in others of large frame, with a deep chest, and much adipose tissue below the integuments, of a quiet and unexcitable disposition, they are scarcely audible

in the præcordial region itself. From the mere extent, then, over which the sounds can be heard, considered *per se*, no conclusion can be properly derived, as to the heart being healthy or diseased.

Irrespectively also of any disease in the organ itself, there are many complaints which cause the sounds of the heart to be heard in some directions more distinctly than in others. Thus, for example, a consolidated lung may cause them to be heard in certain situations, in which they would not be otherwise audible. Hence it happens, that the fact of the sounds of the heart being heard more distinctly than usual in one, or both infra-clavicular regions, while they are not more loud than natural in other parts of the chest, becomes one of the physical signs of phthisis.

An effusion of fluid, or of gas into the pleura, a morbid growth, and even a distended stomach, may so displace the heart as to prevent its sounds being heard in the normal situation, while they become audible in situations, which are abnormal, as on the right of the sternum, &c. When the lungs are extensively affected with emphysema, the heart may be so far separated from the parietes, by those bad conductors of sound, as to be almost or entirely inaudible in every part, excepting at the scrobiculus cordis.

Perhaps it would be on the whole most correct, as it would certainly be most intelligible, to say,

that the heart, when free from disease, in the perfectly natural condition of the organ, and of the surrounding parts, is heard only over the *præcordial region*. By the *præcordial region* is here, and throughout this work, intended, not a part included in any imaginary or arbitrary line, but the space on the anterior surface of the chest which covers the heart, and the origin of the great vessels. It may in addition be stated, that while the first sound is heard most distinctly over the junction of the fifth rib with its cartilage, the second sound is most clearly audible on the sternum, near the third intercostal space of the left side.

But these sounds, it has been already mentioned, very often indeed extend beyond these limits, without any appreciable disease in the heart, or in other thoracic organs. What, then, is the usual order of such extension? What are the regions in which the sounds are, under ordinary circumstances, successively audible? To answer these questions with exactitude, the sounds should first be considered separately, and afterwards in connection with each other. The first sound, then, will generally be found, in the first place, to extend over the left mammary, then over the inferior sternal region, to the *scrobiculus cordis*, and then over the superior sternal region; while the second sound, if separately regarded, may generally first be heard over the superior sternal,

then in the right, and afterwards in the left infra-clavicular region. After extension to the regions just named, the two sounds will commonly be heard with varying distinctness in the other regions, nearly in the following order: the right mammary, the left lateral, and left axillary,—the inter-scapular, the left, and right scapular, the right lateral, and right axillary,—the left and right acromial, the left subscapular, and hypochondriac, and the right subscapular, and hypochondriac regions.

The sounds of the heart are somewhat modified, though considerably less than the impulse, by the position of the body. Thus they are heard more distinctly in the erect position, when the organ more nearly approaches the surface, than in the recumbent position, in which the heart is farther removed from the surface. They are also heard less distinctly than usual upon the left side of the chest, when the body is turned to the right side, and *vice versâ*.

2. *Auscultation of the Heart in disease.*

1. *The Impulse.*

When the pulse at the wrist is materially affected by disease, of whatever kind, a corresponding alteration will be observed in the Impulse of the heart, unless some mechanical impediment exist to the course of the blood. If, in fevers and phlegmasiæ, the pulse be strong and vigorous,

the Impulse is so likewise ; if the pulse be feeble and depressed, so is the Impulse ; when the pulse is contracted and vibrating, as after hæmorrhage, or in anæmia from other causes, as chlorosis, abstinence, &c. the Impulse of the heart partakes of the same character.

When hypertrophy exists,—when, in other words, the parietes of the ventricles, or of the auricles, but of the left ventricle more especially, are thickened—whether that hypertrophy arises from obstruction existing in the valves, or in the aorta, or from other causes,—the Impulse is powerful and heaving. When, together with thickening of the parietes, there also exists dilatation of the cavities, the Impulse is more extensively diffused, as well as more powerful and heaving than natural, and may often be felt over nearly the whole of the præcordial region.

When the parietes of the ventricle, instead of being thick and powerful, are thin and weak ; or when the muscular substance is pale, flabby, or loaded with fat, the Impulse is feeble, but is sometimes also smart, or, as it were, concentrated. When, together with this weakness, thinness, flabbiness, or accumulation of fat, the cavities of the ventricles are also dilated, the Impulse, though sometimes so feeble as to amount to scarcely more than a gentle tremor, is observed over an extent of surface larger than that in which it is ordinarily felt.

When the individual examined is nervous, and excitable, is labouring under chlorosis, or is anæmic from other causes, the impulse is often exceedingly sharp and smart. This kind of beat is constantly mistaken, by the uninitiated, for a powerful Impulse. The patient is said to have a strong Impulse, and is supposed on that account to labour under hypertrophy of the heart. This is far from being true. It is often, perhaps indeed generally, the very reverse of the truth. The Impulse of hypertrophy is full, strong, heaving, and sometimes diffused; the Impulse of chlorosis, anæmia, and irritability, is sharp, smart, and concentrated. The latter appears to proceed from a quick stroke of the apex of the heart against the parietes, like a smart smack of the points of the fingers; while the former seems to be a forcible elevation of the parietes of the chest, by the body of the ventricle, as if the hand itself were applied within, and pushed them forwards.

When obstruction exists in the pulmonary artery, in the mitral valve, or, indeed, in any portion of the heart, or of the large vessels, which is anterior, in the course of the circulation, to the tricuspid valve, the Impulse is generally more or less perceptible at the scrobiculus cordis. This arises from the continued distension of the right ventricle, which results from such obstruction, inducing increased energy of contraction, and

thus leading to hypertrophy of the parietes, and to dilatation of the cavity of the ventricle.

When, in consequence of great and long continued obstruction of the mitral valve, or of such enlargement of the left auriculo-ventricular opening as renders the valve incapable of preventing regurgitation, the blood returns in part through the opening into the auricle, the whole heart, with the occasional exception of the left ventricle itself, becomes enlarged. Under these circumstances, in addition to the Impulse felt at the *scrobiculus cordis*, dependent upon enlargement of the right ventricle, there is frequently a perceptible Impulse between the cartilages of the third and fourth, or even of the second and third ribs of the left side. This superior Impulse is not always exactly synchronous with the Impulse below the nipple, and at the *scrobiculus cordis*. When it is synchronous with that Impulse, it probably arises from regurgitation through the mitral valve. The blood is then propelled back again into the auricle, and by such propulsion may produce the Impulse in the situation specified. When it is not exactly synchronous, but appears immediately to precede the *infra-mammillary* Impulse, it appears most probably to proceed from the muscular action of the auricle itself, hypertrophied by its constant hard work.

When fluid is effused into the pericardium, as

a result either of inflammation, or of obstruction, the Impulse varies according to the quantity of the fluid effused. If the fluid be moderate in quantity, the Impulse below the nipple is only slightly decreased in power, while a gentle rising is sometimes felt between the cartilages of the second and the fourth ribs. If the fluid be in large quantity, the regularity of the Impulse is lost, and its position is often altogether changed. A confused sort of tumbling motion only, at once irregular and unequal, is felt here and there over the præcordial region, instead of the natural Impulse.

But does—the student may inquire—a powerful Impulse necessarily indicate hypertrophy? No! Is, then, hypertrophy always accompanied with a strong impulse? Assuredly it is not!

In explanation of the negative to the former question, it may be stated that an aneurism of the descending, or even of the left side of the ascending aorta, may push the heart aside, and by dilating and producing partial absorption of the parietes, may communicate a powerful heaving Impulse to the part in which the natural Impulse is felt.

In explanation of the negative to the second question, it may be observed, that, though the heart be powerful, and the parietes of the ventricles very thick and strong indeed, the action of the heart may be so hampered by the consequences of obstruction in the valves, and its con-

tractions may be so impeded, and so overpowered by fluid accumulated either within or without its cavities, as to be almost entirely mastered, and to be only just enabled to flutter a little, and thus to rid itself of a small portion of its load. Such, indeed, is very commonly the condition of the organ, in long-standing disease of the valves, and particularly in that of the mitral valve, when the nervous energy of the individual is considerably reduced; even though the muscular power of the heart, abstractedly considered, be much greater than in health. Such is commonly the condition of the Impulse, in the last few days of the life of persons affected with disease of the valves, accompanied with hypertrophy. If they do not die suddenly, as persons so affected frequently do, it may indeed be considered to be the natural termination of such cases.

A very large and powerful heart, therefore, may be accompanied by a very feeble, irregular, and fluttering Impulse, and a very small and feeble heart, may be, and usually is, accompanied with a very smart and "smacking" one.

It has been stated, (but whether it has been proved by inspection after death, or is only surmised, does not appear), that when adhesion of the pericardium exists, without any disease of the heart itself, the situation of the Impulse is not changed, as it is in the healthy condition of the organ, and of its investing membrane, according to the varying position of the body; and that this

immobility of the Impulse arises from the heart being by such adhesion tied down to the diaphragm. Now, when the pericardium is adherent, there generally also exists disease within the heart itself, the consequence of which is the enlargement of the organ. Under such circumstances it is sufficiently obvious that the situation of the Impulse is not so likely, as in the state of health, to be altered by any change in the position of the body, (by reason of the increased bulk of the heart); but whether, in *simple* adhesion of the pericardium, this presumed fixity of the organ really and necessarily occurs, my own experience does not enable me to decide.

2. *The Rhythm.*

Any moral impression, or any physical impediment, which unnaturally excites the action of the heart, may at the same time disturb its Rhythm. Any suddenly supervening emotion of the mind, therefore, whether of joy, fear, or anger, may, in susceptible individuals, almost instantaneously alter its character. Persons whose temperament is naturally excitable, or those in whom it has become so from disease or anxiety, dyspeptics and valetudinarians, are especially liable to, what is called, nervous palpitation; which is merely an excited action of the heart, accompanied with an unnatural Rhythm, and an unequal Impulse. The remote and exciting causes, therefore, of an

unnatural Rhythm are very numerous. Its proximate cause is an interruption of the regular successions of contractions and dilatations of the auricles and ventricles. One cavity contracts before its time, or another is dilated too quickly; the natural series of actions is interrupted; the regular succession is broken.

Such is an unnatural Rhythm. What, then, are the organic changes affecting the heart; what are the physical conditions of the organ which principally tend to induce it? In answer to this question, it may be replied, that the physical condition which interferes most with the equable flow of blood through the chambers of the heart; that the organic changes which in the greatest degree contribute to the interruption of the circulation through the organ, are most likely to produce an unnatural Rhythm.

Among the most frequent causes, therefore, of an unnatural Rhythm, are diseases of the valves; in consequence of which, the cavities behind the seat of obstruction become so distended with blood, and the ventricles thereby so hampered and fettered in their action, that they make various irregular efforts and ineffectual struggles to get rid of the load which oppresses them. This irregularity of the action, or unnatural condition of the Rhythm of the heart, occurs much more commonly in disease of the mitral valve, than of the other valves. This fact is most probably explained by

disease of the mitral valve being more frequently accompanied with regurgitation than that of the other valves. Yet, even when the aortic valves do admit of regurgitation, an unnatural Rhythm does not so frequently accompany the complaint, in proportion to their number, as it does the imperfections of the mitral orifice. Upon what this difference depends it is not easy to determine.

Another very frequent cause of an unnatural Rhythm is thinness, great dilatation, or weakness of the parietes of the ventricles; from which arises a similar incapacity of emptying their cavities as exists in diseases of the valves. From this cause also probably arises the irregular and defective action, with unnatural Rhythm, which often coexists with ossification of the coronary arteries; in consequence of which disease, the muscular structure of the heart is insufficiently supplied with blood, and imperfectly nourished.

An unnatural Rhythm often arises also from large effusions into the pericardium, which probably induce the irregular action of the heart by preventing the cavities being equably and sufficiently filled with blood.

In all these cases the unnatural Rhythm may vary from a slight occasional irregularity to a perfect jumble of sounds, among which one can be scarcely, if at all, distinguished from another.

It is unnecessary, as it would be futile, to attempt to describe the different alterations of the Rhythm which are occasionally met. Their

individual peculiarities are for the most part inexplicable; and a detailed description of their varieties would be attended with no advantage to the student.

One, however, is so peculiar as to require notice. This is the *treble beat*. The heart, on some rare occasions, is found to give rise to three sounds instead of two, as in the healthy condition. It is, indeed, said that the sounds are sometimes quadruple, but this fact I have myself never had an opportunity of witnessing. The treble beat, or sound, I have heard on many occasions. It has been composed of two short and one long sound, and may be represented thus—to-to-too, to-to-too, to-to-too. The origin of this peculiarity is uncertain; but it appears to be most probably connected with the pericardium; a slight and short rubbing noise arising from which either overpowers a portion of the first natural sound, and causes it to appear double, or occupies a portion of the interval.

3. *The Sounds.*

The sounds of the heart may be unnaturally increased, unnaturally decreased, or altogether abnormal in character; in other words, the natural sounds may be louder or less distinct than in health, or they may be overpowered and concealed, or supplanted by morbid sounds, called murmurs.

a. Increase of the Sounds.—It has been

already mentioned that the natural sounds of the heart may appear to be increased in resonance from the surrounding lung being consolidated, and becoming thereby a better conductor of the sounds. With these causes of the increased resonance of these sounds we have here nothing to do. Our present observations apply to the increase of the sounds dependent solely upon the heart itself.

This condition is chiefly observed when, the valves being healthy, the chambers of the heart are dilated, or the parietes of the ventricles are thinner than natural. The ordinary result of this change appears to be, that, while the flapping action of the valves is free, the contraction and dilatation of the ventricles is more rapid and smart, and that the agitation produced in the fluid which they contain is greater than in the natural condition of the organ. The vibrations, therefore, communicated to the surrounding solids, is increased, and, from the thinness of the intervening barrier, is more readily communicated to the surface of the chest than in the state of perfect health.

It is in such cases that what has been called the "back-stroke" is more particularly observed. This is, in truth, nothing more than a loud, clear, and flapping second sound, dependent upon the rapid dilatation of a large ventricle, and accompanied with the sensation of a body retiring from the ear.

When the sounds are loud, as well as clear, enlargement of the cavities, as well as thinness of their walls, will generally be found to exist; when the sounds are clear or shrill, without being particularly loud, the parietes are commonly thin, but the cavities are not dilated.

The particular regions, to which the sounds of the heart are most commonly successively extended, have been previously mentioned, and need not, therefore, be again referred to in this place.

b. Decrease of the Sounds.—A diminution of the natural sounds of the heart may arise from a great variety of causes. Anything which interferes with the free action of the heart itself, or impedes the free motion of its valves; anything which interrupts the full and free contraction of the organ, and consequently decreases the easy and rapid backward flapping of the membranous curtains, tends, in an equivalent degree, to diminish the heart's natural sounds. Overdistension of the organ, therefore, or effusion into the pericardium, from whatever cause arising, may produce this effect: stiffness or want of pliancy in the valves themselves may, without giving rise to a murmur, be attended with a similar result. Morbid sounds, or murmurs, arising from one valve, may obscure the natural sounds arising from the others; and pericardial

murmurs, and bronchial rattles, may drown them altogether.

It is also observed that an unnatural thickness of the walls, or simple hypertrophy, of the heart, causes the sounds to be less clear and loud than when the organ is perfectly healthy. The sounds, under such circumstances, may be more deep and full, but they are not so sharp and loud. The cause of this appears to be two-fold. In the first place, when the organ is thickened, the contraction of the ventricle, though more strong, is less quick and sharp; it produces, therefore, a less rapid, though it may be a more tight, closure of the valves, and consequently excites less agitation in the particles of contained fluid, and less vibration in the surrounding solids. In the second place, the thicker the parietes of the heart, the greater mass of solid matter is there to be moved by the vibrations, from which the sounds originate; the less perfectly, therefore, are they communicated to the surface of the chest, from which they are actually more distant, in proportion to the thickness of fleshy substance through which they have to pass. Supposing then, that they were, as originally produced, of equal intensity or resonance to those existing in health, they might still reach the exterior of the chest, by which they are communicated to the ear, with diminished power, in consequence of the greater thickness of muscle through which they have to

be conveyed ; for nearly the same reason that they are not so distinct and loud in fat, as in thin, persons.

It may be stated to be consistent with general experience, that a thick heart is accompanied with a powerful, heaving impulse, and with feeble or dull sounds ; that a thin heart is accompanied with a smart, and smacking, but not strong, impulse, and with loud, or sharp sounds ; and that a dilated heart is accompanied with a weak but diffused impulse, and with loud and extensively audible sounds. These observations have reference, of course, to the several specified conditions of the heart, independently of valvular obstruction, or inordinate distension of the cavities.

c. Abnormal Sounds or Murmurs. — The gentle current of a quiet stream, with a level bottom and even banks, passes on with but little ripple, and without noise. If, however, a large stone, or any other resisting body, be thrown into the stream, and, by projecting upwards from the bottom, cause considerable obstruction to the free passage of the water, various little eddies and undulations are observed upon the surface ; and if the fall be sufficiently great, or if, in other words, the current be sufficiently strong and rapid, a noise is produced by the agitation of the water thence arising : the quiet “meandering

stream" becomes a "babbling brook." If, again, instead of a large stone being thrown into the centre of the stream, a portion of its bank fall in, and be sufficiently resistant to constitute a decided obstruction to the free passage of the water, a similar ruffling and eddying of the surface will ensue, and a similar noise will, under similar circumstances, arise from the agitation of the water. But if the mass be so large, and the obstruction so great, as to dam up the water towards its source, and thus to impede the force, and diminish the rapidity, of the current, little or no noise will arise; or it will not arise until the back water has become gradually accommodated to the impediment to its downward progress, and the current has resumed its former strength and celerity.

These observations in reference to the stream may not be in every respect, strictly speaking, illustrative of obstructions to the flow of blood through the heart and great vessels; but they are, I believe, in the main perfectly applicable, and they will, I trust, contribute to render the succeeding remarks both intelligible, and easily available at the bed-side.

When the healthy blood passes through the healthy heart in a quiet stream, no noise results from its direct transmission. The valves of the large arteries are appressed to the sides of the vessels, and those between the auricles and ven-

tricles leave a space sufficient for the free passage of the blood. No obstruction exists; no inordinate agitation or eddying of the fluid is produced; and no noise results from the transit*.

When, however, the valves become thickened by inflammation, as in endocarditis, they lose their pliancy and flexibility; they do not set close to the walls of the vessels, and do not leave a free passage for the blood: when their surfaces are covered, partially or entirely, with plastic lymph, or their free edges are studded with minute granules or vegetations; when they are puckered, wrinkled, stiffened, and corrugated, from the remains of old disease imperfectly repaired; or when they are rendered unyielding and uneven by the deposition of atheromatous or ossific matter below the lining membrane; when, indeed, any mechanical obstruction exists to the free passage of the blood through the apertures which are guarded by the valves, an unusual agitation is excited among the particles of the blood, which, if the rapidity of the current be sufficiently great, gives rise to a morbid sound, or "murmur." Of this character, and of similar origin, is the vast majority, at least, of morbid sounds or murmurs arising within the heart and large

* The production of the natural sounds of the heart is not opposed to the correctness of this statement, as they, it must be recollected, arise not from the direct transmission of the blood, but from its being forced backwards.

vessels.—Let it, then, be recollected, that the natural sounds of the heart are produced by the flapping back of the valves, preventing the retrograde movement of the blood, and that the morbid sounds, or “murmurs,” of the heart are produced, at least generally, by the passage of the blood through the valves.

The morbid character of the sounds may amount merely to an unusual harshness, and increased duration of the natural sounds, as sometimes occurs in the very early stages of rheumatic endocarditis; when, simple as it may appear, this harshness becomes a most important indication of future mischief, and of more obvious signs of the malady too soon to make their appearance.

It may be only a slight puff or whizzing noise, very accurately resembling that produced by the use of a pair of bellows—hence called “*bellows murmur*,” (“*bruit de soufflet*”); or it may be like that caused by rasping or filing wood—“*rasping murmur*,” (“*bruit de râpe*”); or like that arising from sawing wood—“*sawing murmur*,” (“*bruit de scie*”); or, finally, it may approach to the tone of certain musical instruments, as of a bass viol, a bassoon, or oboe—“*musical sound*”*.

Are, then, these different sounds, it may be

* Of the direct cause of the musical tone in certain diseases of the heart my own experience does not enable me to speak with certainty; as in the four or five cases of the kind which I have

asked, characteristic of different states of disease, or are they all merely indicative of obstruction to the progress of the blood? Is the harshness of one sound dependent upon a rigid obstructing medium, and the softness of another sound dependent upon a pliable obstructing medium? Does the nature of the deposit materially affect the character of the murmur? Is it harsh and saw-like, with a bony deposit, and soft and bellows-like, with an albuminous deposit? These questions have been answered (it is presumed conjecturally) in the affirmative. My own individual experience necessitates a direct negative. It may, indeed, be pretty confidently asserted that the truth is the direct reverse, and that no such necessary connection, as, *à priori*, might be deemed at least probable, is found in nature; but that, on the contrary, soft murmurs coexist with hard deposits, and that harsh murmurs arise from soft deposits; that nothing can be determined as to the quality of the obstructing cause, from the character of the sound; as weekly

had an opportunity of examining after death, it has rather curiously happened that two conditions have existed, either of which might by possibility have given rise to it, or both of which might have contributed or have been necessary to its production, These two conditions have been, disease of the mitral valve and of the tendinous cords, and cellular adhesion of the pericardium to the pleura costalis. I believe, however, that the sound really arises from the vibration of the tendinous cords, or other parts of some of the valves, rendered tense by disease.

experience demonstrates that a very soft murmur may depend upon a rough bony valve, and that a harsh grating murmur may arise from soft vegetations, or have its origin in even a simple ulcerated opening through a valve.

But is a murmur indicative of valvular disease capable of being referred to any particular valve, or does it merely imply obstruction somewhere? Are there any signs by which the disease of one valve may be distinguished from the disease of another valve? If such signs really exist, what are they? In reply to the first question, it may be stated that certain peculiarities do, for the most part, accompany the disease of particular valves; a brief sketch of the anatomical relations of the parts concerned will be requisite, previously to giving a satisfactory answer to the second and third questions.

The valves of the pulmonary artery are situated almost exactly opposite the junction of the cartilage of the third rib with the left of the sternum. The vessel passes forwards and to the left, but is soon directed backwards to be divided into its right and left branches.

The aorta is at its origin partly covered by the pulmonary artery. One perpendicular section would divide part of the valves of both. But the aortic valves, taken as a whole, are about half an inch to the right of, and about the same distance lower down than, those of the pulmonary artery. The aorta, emerging from behind, passes in a

different direction from the pulmonary artery. It is directed upwards, forwards, and to the right, till it arrives near to the junction of the cartilage of the second rib with the right side of the sternum; when it crosses behind that bone to form the arch, and then dips downwards and backwards, and becomes the descending aorta.

The mitral valve is situated nearly an inch below those of the aorta, and extends considerably more to the left side. It is situated about opposite the space between the cartilages of the fourth and fifth ribs of the left side. The body of the left ventricle, to which it is attached, rests posteriorly upon the spine, the contents of the posterior mediastinum alone intervening between them.

The tricuspid valve is considerably to the right of, and anterior to, the mitral, and it is for the most part covered by the sternum.

Now when these anatomical relations of the different valves are remembered, and when it is also recollected that the morbid sounds or murmurs originate in the circulating blood, and that the vibrations therein excited are conveyed along its current, and are communicated to the solids by which it is surrounded, it will be easily understood, that the murmurs resulting from disease of the different valves, usually occur, or are most distinctly heard, in different parts of the chest.

It may be then stated, as consistent with

theory, as well as with experience, that the murmurs resulting from disease of, or from obstruction in, the pulmonary valve, are heard most distinctly opposite to the cartilage of the third, or to the interspace between the second and third ribs. It will be also understood, why they appear to be very superficial, to pass a short distance to the left, and then abruptly to terminate, or to diminish very considerably in their intensity.

Diseases of the aortic valves are most commonly accompanied with a sound, which appears to be less superficial than that arising from the pulmonary valves. It is most distinctly heard over the sternum, near to its junction with the third rib of the left side, or the space between the third and fourth ribs. It is heard, though with gradually decreasing intensity, in the course of the aorta, upwards and to the right; it often extends as high, or nearly as high, as the clavicle, and sometimes along the top of the sternum.

The murmurs dependent upon disease of the mitral valve are ordinarily most distinctly heard opposite the junction of the cartilages with the fourth and fifth ribs; or immediately below the nipple. They are not so plainly conveyed in the course of the aorta as those of the aortic valves; they are commonly audible in the axilla, and are very frequently heard on the left of the spine opposite to the sixth, seventh, and eighth dorsal vertebræ, or thereabouts; for it is scarcely pos-

sible, excepting in very thin persons, exactly to mark the particular vertebræ in the dorsal region.

Murmurs, resulting from disease or imperfection of the tricuspid valve, are usually loudest over the central and lower part of the sternum. They are frequently audible at the *scrobiculus cordis*, but are not conducted in the course of the aorta, nor (I believe) heard in the back.*

Such are the situations in which, and the circumstances under which, the "Murmurs" arising from disease of the different valves are generally heard most distinctly. Such may be said to be the ordinary rule. But, whether the fact arise from an alteration in the relative situation of the different parts of the heart, produced by enlargement and distension, or from other causes, it is assuredly a rule liable to occasional exceptions, and to exceptions moreover which are not always easily to be explained by examination of the body after death. There are, however, many circumstances in connection with other physical signs, and with the general symptoms, which tend very materially to facilitate the diagnosis of the diseases of the different valves: but upon these it is not now my province to dilate.

Hitherto those murmurs, which arise from the

* Disease of the tricuspid valve, independent of more extensive mischief in the Mitral or the Aortic valves, is so rare, that it cannot perhaps be positively stated, what are the especial physical signs of that particular affection.

direct, or forward progress of the blood, have been alone referred to. But it often happens that the valves are so stiffened and crumpled, so contracted and shortened, or that the openings, which it is their peculiar province to guard, are so enlarged, that the valves are incompetent to close them; or that, as the result either of rent or of ulceration, an aperture exists in one or other of the valves, which admits of the backward passage of the blood. Under any of these circumstances a portion of the blood, upon the occurrence of the diastole, returns into the cavity from which it had been propelled by the systole. If then the imperfection exist in the tricuspid or mitral valve, part of the blood returns into the auricles on the contraction of the ventricles; if the imperfection exist in the sigmoid valves, part of the blood returns into the ventricles on the contraction of the large vessels. In its return, the blood passes over a rough surface, or through a contracted space. The space may be absolutely contracted, or it may be contracted only comparatively with the size of the cavity to which it is the mode of access. The backward passage of the blood, through this space, may, and frequently does, give rise to such agitation among the particles of the fluid, as to produce sounds, similar to those which have been referred to as resulting from the direct passage of the blood through an obstructed valvular opening. There

are, therefore, "Murmurs of regurgitation," as well as "Murmurs of obstruction."

Murmurs, then, may arise from the direct passage of the blood through a contracted valve; they may arise from the retrograde motion of the blood, through an imperfect valve; or they may arise from both causes combined. When a murmur arises from the direct passage of the blood through one valve, and the retrograde motion of the blood through another valve, though the cause be double, the murmur appears almost universally to be single, because they occur at the same time. When, on the contrary, a murmur arises from the direct passage, and also from between the retrograde motion, of the blood through the same valve, or the same pair of valves, the murmur is double because it occurs at different times.

If the student bear these observations in mind, and reflect upon the circumstances upon which they are founded; if he recollect the origin of murmurs, obstructive and regurgitative, and at the same time remember the action of the heart, and of its valves, he will easily understand the following illustrations, and will I believe be capable of unravelling most valvular murmurs. Exceptions of course will occur to him, as exceptions occur to the most experienced; but to the one as to the other they will be only exceptions to a very general rule.

When regurgitation takes place through the aortic valves, the disease giving rise to which does not in any material degree produce obstruction to the direct current of the blood, a murmur is heard most distinctly over the sternum, opposite to the third rib; it occupies the time or place of, and often entirely conceals, the natural second sound of the heart; though the second sound may be sometimes indistinctly heard together with it between the cartilages of the second and third rib, upon the left side.

When both obstruction exists in, and regurgitation takes place through, the aortic valves, a double murmur, or "see-saw" sound is heard over the sternum, opposite the third rib, and passes upwards in the course of the aorta, but gradually decreases in power, as the stethoscope is farther and farther removed from the situation of the valves. Should the same double murmur, or "see-saw" sound, occur opposite the second intercostal space,—should the murmur be pretty much confined to that situation, and not follow in any degree the course of the aorta,—it may be presumed, rare as, comparatively speaking, is the affection, that obstruction exists in, and regurgitation takes place through, the pulmonary valves.

When a murmur, which accompanies, or occupies the time and place of the first sound of the heart, is most distinctly heard below the nipple, is audible in the axilla, and in the left inter-

scapular region close to the spine, or over the bony column itself; when it gradually diminishes in power as the stethoscope is removed from these localities; and when it is not at all, or only very imperfectly, conducted along the course of the aorta, it may be presumed, that regurgitation takes place through the mitral valve. This regurgitation, however, it must be ever borne in mind, may occur through the mitral valve, and through other valves, in consequence either of the valves themselves being diseased and contracted, and thereby unfitted to perform their natural function in closing their respective apertures, or of the apertures being dilated, and the valves, though healthy, being consequently incapable of doing so.

Supposing, then, that a loud murmur occupies the time of the first natural sound, is equally loud behind the sternum, opposite the third rib, and below the nipple; that it follows the course of blood in the aorta, and is also distinctly heard near the spine, opposite to, or rather below, the angle of the scapula—what, according to the preceding statements, may be presumed to be the probable nature of the affection? If the student reflect, he will reply, “Probably direct obstruction of the aortic valves, and regurgitation through the mitral valve.”

It must surely be unnecessary to illustrate this subject farther. Let the student reflect upon

these examples—let him, as it were, follow them home—let him see, and fully appreciate, “the why and the wherefore” of each case, and I believe he will be capable of making out *most* of the instances of murmurs, or morbid valvular sounds, which occur under his notice.

He will be greatly facilitated in doing this, by the recollection, that valvular disease on the right side of the heart is exceedingly rare, comparatively with that of the left side, and that a direct murmur of the mitral valve is by no means common, and therefore that murmurs connected with the mitral valve arise, in the great majority of cases, from regurgitation.

These statements may be perhaps considered of sufficient importance, in reference to diagnosis, to be repeated in a somewhat different form.—Valvular disease exists much more frequently upon the left than upon the right side of the heart, and, even when present upon the right side, almost always exists to a greater extent upon the left side.—Murmurs connected with the mitral valve arise much more frequently from regurgitation, than from obstruction to the direct passage of the blood. As a rule, therefore, though like most other rules it is liable to many exceptions, it may be generally considered, that murmurs, arising from valvular disease, proceed from the left side of the heart, and that, if synchronous with the diastole of the ventricle, they arise

from regurgitation of the blood through the aortic valves.

But do murmurs necessarily indicate either obstruction from valvular disease, or regurgitation from valvular imperfection? Is valvular disease always accompanied by murmurs? Both these questions must be answered in the negative.

Morbid sounds, resembling murmurs, may in the first place arise from diseases of the pericardium. These will be separately considered hereafter.

- But as, in the case of the stream previously referred to for the purpose of illustrating morbid sounds, any solid substance of sufficient density, which encroaches upon the channel by projecting from its banks, may, like a solid body placed in the centre of the current, produce a strong ripple or eddy, and give rise to a liquid rustling sound; so any substance projecting inwards from the parietes of the heart, or from the coats of the large vessels, or so compressing them as materially to encroach upon the channel through which the blood passes, necessarily gives rise to an agitation of the fluid, similar to that resulting from obstruction in the valves, and may, therefore, like it, give rise to a variety of murmurs.

Aneurism, independently of the local pressure which it, as a tumour, exerts upon the heart, and other diseases of the aorta, may also give rise to murmurs.

When the stethoscope is pretty firmly pressed upon a large artery, as the abdominal aorta, the axillary, the femoral, or the carotid, so as partially and locally to obstruct, but not to stop, the current of the blood, a whizzing noise or bellows murmur is constantly heard, which is synchronous with the systole of the ventricle. This evidently arises from the increased agitation, caused by the obstruction among the particles of the fluid, being communicated to the walls of the vessel, and thence through the stethoscope to the ear. Such a sound, created by the examiner himself, has often given rise, and may, with the incautious, still give rise, to the incorrect supposition of the presence of aneurism. It may indeed prove a source of fallacy to the most careful and circumspect. Nor can it be regarded as surprising that such should be the case, when it is recollected that the sound of an aneurism is often very similar, if not absolutely identical, and that it arises from a precisely similar cause, viz. obstruction to the current of the blood.

Like the pressure of the stethoscope, the pressure of a tumor, as of an abscess, or of malignant growth, upon the course of a large vessel, may produce a murmur, and may thus very exactly simulate an aneurism.

The mere dilatation of a vessel does, I believe, not give rise to a murmur; nor does the entrance of blood into an aneurismal pouch necessarily produce a whizzing noise. But if the dilated

vessel have its lining membrane roughened ; if it be studded with atheromatous or ossific matter ; and if this roughening, or this deposit, exist to such an extent as materially to interfere with the free passage of the blood, then a turmoil, agitation, or eddying, is excited in the current, like that of a rapid stream with a rough bottom ; and such vibrations of the coats of the vessel are produced, as constitute a bellows murmur, or sometimes a harsher sound.

If an aneurism press not upon the calibre of the vessel from which it arises, or upon that of any other vessel of large size ; if the opening, by which the blood enters the pouch, be as large, and as direct, as that by which it passes along the parent trunk ; if, in fact, there be no obstruction, direct or indirect, to the passage of the blood, no morbid sound will be produced. But if the converse of this exist ; if, that is to say, the aneurism do press upon the vessel from which it arises, or upon some other vessel ; if the opening by which it communicates with the artery be not large, or be oblique ; so that obstruction does exist either to the direct passage of the blood, or to its admission into the aneurismal sac, then will be produced the indications of such obstructions ; a whizzing noise will then arise, or some other morbid sound will be excited.

Aneurism, or other disease of either the ascending aorta, or of the arch of the aorta, may, therefore, give rise to sounds very similar to, if not

identical with, those arising from obstruction in the valves. They may, however, be very generally distinguished by their position, by their superficial character, and by the general symptoms or local signs, which result from pressure.

Aneurism of the descending, and even of the ascending aorta, may give rise to sounds below the nipple, or in the back, or in both situations, so as very exactly to resemble the morbid sounds arising from regurgitation through the mitral valve. The diagnosis in such cases is often a matter of extreme difficulty. It can, indeed, be rarely determined with positive certainty, unless the aneurism project externally.

Anæmic Murmurs.—In certain states of the system, or, it may be, with certain conditions of the circulating fluid, as in chlorosis, or in anæmia from hæmorrhage, or from other causes, murmurs frequently arise from the passage of the blood, independently of absolute disease of the heart or great vessels. These are termed *anæmic murmurs*, or "*chlorotic bruits*."

They are ordinarily of the softer kind, and resemble the blowing of a pair of bellows ("*bruit de soufflet*"), but they are sometimes quite harsh, and resemble the rougher morbid sounds, as that of filing or sawing ("*bruit de râpe* and *bruit de scie*")

They are very generally supposed to be confined to the aortic openings. This is certainly a

mistake. They are most assuredly very frequently connected with the pulmonary artery, in which murmurs, quite independent of any disease of the vessel, or of its valves, are far from uncommon.

Murmurs often arise from some body pressing upon this vessel ; as a solid mass, the result of pleurisy, of pneumonia, or of phthisis, or enlarged bronchial glands, abscesses of the anterior mediastinum, &c. &c. The murmurs frequently also coexist with chlorosis, or with other forms of anæmia. Are these latter murmurs, then, whether in the pulmonary artery, or in any other part of the circulating system, to be distinguished with tolerable certainty from morbid sounds, the result of organic obstruction within, or without the heart or large vessels ?

Generally speaking, they may, I believe, be distinguished from each other : but they certainly cannot always be so ; and never with absolute certainty by the mere character of the murmur alone. There are, I feel assured, some examples of these anæmic murmurs, which can be proved to be simply functional, and not to arise from organic disease of the heart or its vessels, or from pressure upon them, only by the results of treatment.

Let, then, the student be careful not to assert too confidently that a patient, on the one hand, has organic disease of the heart, or great vessels, merely because he has a harsh murmur over the

aorta, an occasionally irregular rhythm, and a vibrating pulse, which usually coexist with an anæmic condition of the body; or he may cause unnecessary alarm and anxiety; nor let him, upon the other hand, too hastily determine, that, because a murmur is soft, and his patient is an hysterical girl, with a pale face, and is subject to leucorrhœa, and to amenorrhœa, that she has no organic disease; or some day, to his great surprise, grief, and mortification, and possibly also to his disgrace, he may find she has died suddenly with diseased heart.

Anæmic murmurs, however, it may be stated, are very local, and are generally pretty much confined to the situation of the sigmoid valves, either aortic or pulmonary, or both; they do not follow the course of the large vessels so fully, or so frequently, as do the murmurs arising from disease of the valves, or of the arteries; they occur only during the systole of the ventricles; and, as they cannot arise from regurgitation through the mitral valve, they are not heard very distinctly below the left nipple; they are always, so far as I know, accompanied with a smart smacking impulse; they generally disappear for a time while the individual is quiet, mentally as well as bodily, if by that quiet the heart assume a natural impulse; and they are always diminished, and generally disappear entirely, under suitable treatment.

The origin of these anæmic murmurs has latterly been very generally attributed to a watery condition, or a diminution of ordinary viscosity, of the blood; in consequence of which it is believed that the particles of the fluid move more easily over each other, are therefore more freely agitated, and thus give rise to the vibrations which produce the murmur. This may have some, and perhaps an important, influence in producing them.

But there are other circumstances which also appear to play an important part in their causation. The principal of these is the remarkably quick and sudden contraction of the ventricles; in consequence of which the fluid contents of the cavities are propelled through the comparatively small area of the mouths of the large arteries in a shorter time than during the leisurely contractions of health, or the frequent, but not sudden, contractions existing in some other forms of disease. Though, therefore, no actual contraction exists, an obstruction is practically produced by the increased velocity with which the blood is propelled, through the aortic and pulmonary openings. The increased agitation in the fluid thence arising, it is at least probable, has a principal part in the production of anæmic murmurs.

If the heart beat quietly, and the impulse be natural, however decided the pallor of the face,

and whatever the watery condition of the blood, no murmur, I believe, exists when no mechanical obstruction is present.

It is also possible that the *quantity* of the circulating fluid is decreased in such cases, in addition to its *quality* being altered, and that while, by the elasticity of their coats, the arteries are capable of accommodating themselves to the diminished quantity of the fluid, the cavities of the ventricles retain their normal capacity, and that on this account an absolute, as well as a comparative obstruction, may exist to the transit of the blood.

Concurrently with these anæmic murmurs at the origin of the large arteries, there is often heard, upon the application of the stethoscope to the side of the neck, a curious sort of humming noise, which ceases when firm pressure is exerted upon the jugular vein at a point above that on which the end of the stethoscope is placed. It is continuous, not intermittent like the arterial murmur, and is therefore sometimes called the "continuous humming," as well as the "venous murmur"—"*bruit de diable*," &c.

It most probably depends upon partial obstruction to the quickened flow of blood through the veins. Strong pressure causes it to cease; but without pressure, *directly or indirectly applied*, it is, I believe, never heard. Like the anæmic murmur of the arteries, it is supposed to be asso-

ciated with a watery condition of the blood, and it is, we are told, a frequent, if not a constant, attendant upon that state of the system with which such a watery condition of the blood is a concomitant.

This statement is not made from my own observation, but if true, the venous hum may be perhaps considered a useful assistant indication of the anæmic state.

But great obstruction to the blood may, as has been previously hinted, exist; extensive disease may be present in the valves of the heart, or in the large arteries, and yet no murmur may be heard. This arises from circumstances which may be, as they have already been partially, illustrated by the stream, in which a certain rapidity of the current is necessary to produce such an agitation of the water as will give rise to sound. Though the bottom of a rivulet be very uneven, and its banks exceedingly irregular, yet if the current be not tolerably strong, little or no ripple will be produced, and no sound will be generated. It is just so with the blood; rapidity of the current of the blood, as well as obstruction thereto, is necessary to produce such an agitation among the particles of the fluid as will give rise to sound.

Hence it often happens that a heart with extensive disease of the valves may be without murmur while the patient is quiet, and the circula-

tion is slow ; though immediately the circulation is accelerated, either by physical exertion or by mental emotion, a murmur becomes distinct. Hence, also, it happens, that when the cavities of the heart become greatly distended, in consequence either of the magnitude of the obstruction, or of defective nervous power, the ventricles are frequently incapable of acting upon and propelling their contents with sufficient force to produce a murmur. The channel is irregular enough, but the rapidity of the current, and of the resulting vibrations, are not equal to the generation of sound. Hence, likewise, it arises, that when fluid is present to a large amount in the pericardium, the heart may be so oppressed with the accumulation upon its exterior, that, though great obstruction exist within, no murmur is produced. Thus it will be often observed that when the obstruction is greatest, the murmur, if even it be heard at all, is very feeble ; and that when the obstruction is small, the murmur is very loud ; thus also, in persons who, for weeks and months, and even years, have presented notably morbid cardiac sounds, these sounds, if the individuals are not carried off suddenly, very frequently, or perhaps even generally, cease altogether some days before death.

The cause of this, as before stated, is either that the heart does not contract with sufficient power, or, if it act forcibly, that it cannot act

upon, and propel through the contracted orifices, the large quantity of blood which distends its cavities with a rapidity sufficient to give rise to sound.

Let, then, the student ever bear in mind the truth, that mere obstruction is not in itself sufficient, but that a certain force or rapidity of the circulation must be necessarily combined with that obstruction, to give rise to morbid endocardial sounds. Murmurs may exist without any obstruction of an organic kind; but without a certain degree of force in the circulating current they cannot exist.

Pericardial Murmurs. — The movements of the heart in the pericardium, when it is healthy, are not accompanied with any sound. The folds of the glistening and slippery membrane, like those of the pleura, glide over each other without noise. But when the surface of the membrane is rendered rough by inflammation; when solid or semi-solid plastic lymph is effused upon one or other of its folds, the smooth gliding motion ceases, attrition occurs, and a superficial rubbing noise is heard over the pericardial region upon each motion of the heart. This is “pericardial rubbing,” “exocardial murmur,” or “*frottement*.”

It may exist in, and be confined to, a small space; or, commencing locally, it may gradually

extend over the entire surface, or it may simultaneously appear over the whole, of the membrane at once.

As the fluid of pleurisy removes pleuritic rubbing, so the intervention of fluid between the roughened surfaces of the pericardium prevents their attrition, and most commonly removes the rubbing sound.

Under such circumstances the sound may gradually disappear from below upwards, like that of pleurisy, as the fluid accumulates; and, after its entire cessation, it may gradually reappear from above downwards, as the fluid becomes absorbed, and become again audible over the entire surface occupied by the pericardium.

When only a thin layer of fluid exists in the serous sac, the rubbing noise may, after its temporary cessation, be sometimes made to reappear in certain situations by altering the position of the body, and thus causing the effused fluid to gravitate to the lowest part of the bag, and to leave the uppermost surfaces comparatively dry. It may also, under similar circumstances, be occasionally reproduced by strong local pressure of the ear or stethoscope; by which the two surfaces of the membrane are brought into contact by the fluid being squeezed away from the part to which the pressure is applied.

After entire absorption of the fluid, the two layers of the membrane commonly adhere, when the rubbing noise of course ceases. But some-

times the absorption is incomplete, or other causes prevent the adherence of the two layers of the membrane, the surfaces of which continue permanently roughened. The pericardial rubbing consequently remains behind, and, unless a fresh attack of inflammation occur by means of which adhesion is produced, or a fresh effusion of fluid takes place which separates one layer from the other, it becomes permanent.

The character of the deposit, in the process of time, becomes altered. It may consist of non-plastic lymph; it may be converted into a hard tough substance, approaching the consistence of fibro-cartilage, like the lining of a calf's mouth; or it may consist partially of fluid, and partially of cellular membrane. The sound, which results from the attrition of the two surfaces, varies accordingly. It may be a simple rubbing, "*frottement*;" it may resemble a harsh grating or scraping noise, "*raclement*;" it may simulate precisely the creaking of new leather, "*cracquement de cuir*;" or, finally, its character may very nearly approach the bellows murmur of the valves, "*bruit de soufflet*."

Is then a pericardial sound always to be distinguished from an endocardial or valvular murmur? Can it always be positively stated—this is a pericardial rubbing, and that is an endocardial murmur? This is inside the heart, and that is outside the heart? By experienced auscultators it may indeed be generally so stated,

but I believe not always. For myself, at least, I must confess that I am sometimes in doubt, and moreover sometimes wrong, in my conjectures upon this question. The pericardial rubbing occasionally so closely approximates, in character and situation, the valvular murmur; it appears indeed, as regards the sound, so perfectly identical with it, that I hesitate before I give an opinion, or I feel compelled to acknowledge my inability to form one. These are, however, exceptive cases, and they constitute very rare exceptions to a very general rule.

What, then, it may be inquired, are the especial characteristics of pericardial sounds? They are, or commonly appear to be, more superficial than even pulmonary or aortic murmurs; they approximate more to a rubbing or grating noise than the endocardial murmurs, which have more of a blowing or whiffing character; they ordinarily occupy the period of both the diastole and the systole of the ventricles, and very commonly also a portion of the interval or pause; they are, in fact, less decidedly *intermittent* than the valvular murmurs, and often afford the impression of a circular rubbing motion, like that of a mullar used in grinding paint, rather than of the puff or whizz of a valve; they are more localized—*i. e.* they extend beyond the surface, by the attrition of which they are produced less than,

and do not follow the course of the blood so much as, valvular sounds.

Though then, it is acknowledged, that it is not possible to state with absolute certainty, in all cases of old standing disease, "This is a pericardial rubbing," and "that is an endocardial murmur;" yet the cases in which this difficulty occurs are on the whole very rare; the sounds are ordinarily sufficiently distinct to a person with a moderately good ear. So distinct, indeed, are they on some occasions, that in cases of acute endocarditis accompanied with pericarditis, the experienced auscultator can often in the first place detect the endocardial murmur, and then the gradual supervention of pericardial rubbing, when the two sounds may be heard at the same time:—he can then observe, that, as the pericardial effusion increases, the rubbing becomes so loud as to drown the feebler sound of the valves,—that it then gradually decreases as the plastic matter becomes partially absorbed, and that the two sounds are again heard in combination; till at length adhesion of the pericardium takes place, when the endocardial murmur becomes again, as it was in the first instance, pure and unalloyed.

Mixed Murmurs.—Independently of the combination of sounds, which has been just referred

to, there are some other irregular sounds of the heart, which require a brief notice, and which have been deferred to this place, in consequence of their being dependent upon affections of other organs conjointly with the heart.

It occasionally happens, that, though the pericardium is adherent, or quite free from disease, the pleura which is in contact with its exterior becomes inflamed, and roughened. Under such circumstances, though the motions of the heart, in its proper serous membrane, give rise to no noise, the attrition of the two surfaces of the roughened pleura does.

This sound cannot be always with certainty distinguished from the proper rubbing of pericarditis. It may, however, be sometimes distinguished by the following circumstances. When the lungs are inflated, they press with more force against the heart bag, than when they contain a less amount of air. A fresh surface also of the pulmonary pleura is probably brought in contact with that covering the pericardium during inspiration. The consequence of this is, that the rubbing noise dependent upon the action of the heart against the roughened pleura, is sometimes only heard, and is generally most distinctly heard, during the inspiration. It possesses, therefore, a double synchronism. It keeps time with the respiration, and with the heart also; it disappears or decreases during the expiration, and reappears

or increases with each act of inspiration; yet, when occurring, it coincides with each pulsation of the heart.

Similar to the preceding, in many respects, is another sound. This, however, differs in being endocardial, while that, though not pericardial, is strictly speaking exocardial. It is a valvular sound, a pure bellows murmur, "*bruit de soufflet*," which, though audible during the inspiration, and during the inspiration only, is perfectly synchronous with each systole of the ventricle. It is, as far as I have observed, almost always audible over the valves of the pulmonary artery. It may be supposed to arise from the small additional pressure, caused by the inflation of the lungs, being sufficient under peculiar circumstances to create such an obstruction to the progress of the blood, as to give rise to a sound. How, as it exists in some few cases, and in very few cases only, the simple inflation of the lung is capable of producing this effect, I must confess I do not understand.

It occasionally happens, also, that a gangrenous, a phthisical, or a pneumothoracic cavity is situated in the immediate vicinity of the heart, and that by each impulse of the organ an agitation is produced in the gaseous and fluid contents of that cavity.

Two such cases only have occurred in my experience. In both, the sound was very peculiar, and

in both the precise physical conditions, which after death were found to have given rise to the sound, were predicted during life. In both the noise was very similar. It occurred during the dilatation as well as during the contraction of the ventricles, and evidently arose from the impression made by the stroke of the heart upon the sides of the cavity, and the agitation produced by that impression in its mixed fluid and gaseous contents. The noise may be perhaps best compared to that produced by dashing a spoon into a basin of porridge, or of hasty pudding, and may be pretty accurately represented by the syllables "blob blob—blob blob." I never heard a similar sound produced by the heart. When once heard, I imagine it could not be mistaken; nor indeed do I believe, that it could be well mistaken though it had never been heard before

CHAPTER VI.

MENSURATION, OR EXAMINATION BY ADMEASUREMENT.

MENSURATION is assuredly at once the least important, and the least satisfactory mode of physically examining the chest. In most diseases it is of no use whatever, and even when it is apparently most applicable, its employment rarely affords results which are perfectly satisfactory. It very frequently happens, that a side, which to the eye is evidently bulging and misshapen from the presence of fluid, or other consequences of disease, is found upon admeasurement to be scarcely, if at all, larger than the opposite and healthy side.

The various modes formerly employed of determining the capacity of the chest, by measuring the quantity of air a person takes in, or exhales, by a natural, or forced inspiration, or expiration, are now rarely had recourse to. The simple capacity of the cavity can be as well ascertained by *Inspection* of its form and proportions as by any other method. It is indeed true, that, together with a large and round chest, the quantity of air which can be inspired may be compara-

tively, or absolutely, small; yet this fact also may be estimated, with accuracy sufficient for all practical purposes, by watching the amount of elevation of the ribs, and the general expansion of the whole cavity, upon a deep inspiration.

Of the mode of examining the comparative depth of the two sides of the upper part of the chest, in suspected, or in the early stage of, phthisis, by means of instruments, I must acknowledge that I have no experience. I have never employed them, because I have never felt need of them. I cannot but believe that, with a quick and correct eye, and a sensitive touch, they are unnecessary. From my own experience of other modes of mensuration, indeed, I am quite satisfied, that a slight flattening, or sinking, of one or other of the infra-clavicular regions, might be sufficiently obvious to the eye, and that the imperfect elevation of one side might be sufficiently appreciable by both the eye and hand, though neither might be indicated by the employment of instruments. Still, I must repeat, that of the use of callipers as instruments for measuring the comparative size of the two sides of the chest, I have no experience. I may therefore be, perhaps, considered unqualified to form an opinion of their supposed advantages.

Mensuration is principally available, as a means of diagnosis in diseases of the chest, in the case of inflammatory, or of other effusions into the pleura.

It has been already stated more than once, that the first mechanical result of fluid effused into the pleura is to compress the spongy tissue of the lung, and that, supposing the lung to be free from adhesions to the costal pleura or the diaphragm, and to be otherwise healthy, as the quantity of fluid increases, a corresponding portion of air is usually squeezed out of the organ; till it is at length reduced to a small dense mass closely pressed against the spine. The mediastinum is at the same time pushed over to the opposite side of the chest. It is then, but ordinarily not till then, that the less yielding parietes of the chest begin to be enlarged,—it is then, therefore, but commonly not till then, that mensuration becomes effective as a means of diagnosis of effusion into the pleura. One side of the chest, therefore, may be quite full of fluid, without any appreciable enlargement. It is not until it becomes, what may be said to be, more than full, that the ribs begin to be expanded, or the intercostal spaces begin to bulge.

But it must be ever borne in mind, that one side of the chest may be enlarged without any fluid existing in the pleura,—as from aneurism, from malignant disease of the lung, of the liver, or spleen, and from other morbid growths, as well as from local emphysema the product of local obstruction. It must also be remembered, that the pleura may, on the other hand, be quite full of fluid, and

may yet be not only not larger, but even smaller, than the opposite or healthy side. After one or more attacks of pleurisy, a portion of the fluid may have been absorbed, and, from the lung being firmly bound down by adhesion, and therefore incapable of expansion, the ribs may have fallen in, and the side have consequently become decreased in size in proportion to the amount of fluid removed.

Mensuration, then, should not only never be considered as a determinate mode of deciding upon the presence of fluid in the pleura, when taken alone; but it should ever be regarded as of comparatively little value in confirming the diagnosis of its presence, derived from other sources, or dependent upon other modes of exploration.

This is more especially necessary to be considered where the operation of paracentesis is contemplated. For if, in a case of urgent dyspnoea threatening immediate suffocation, the cause of which is supposed to be connected with effusion into the pleura, the trochar were plunged into the larger side, considered, on account of its size alone, to be the diseased side, it would not unfrequently happen that the healthy side would be punctured; the not improbable result of which might be, that the healthy lung would be collapsed by the admission of air into the pleura, and the patient fall dead at the operator's feet.

This has already happened more than once

from a mistake in regard to percussion, similar to that which is now contemplated in reference to mensuration. The more dull was incorrectly supposed to be the diseased side, and to be filled with fluid, while, in truth, it possessed its natural degree of resonance, and only *appeared* to be dull in consequence of the tympanitic resonance of the opposite side, which resulted from pneumothorax which was the real cause of distress. The trochar was introduced, the air rushed in, the healthy lung was collapsed, and, that of the other side being already collapsed, the patients immediately dropped down dead. Hence arises the very important practical rule, *never* to introduce the trochar into one side of the chest supposed to be diseased, unless it has been previously ascertained that the lung of the opposite side, if it be not absolutely healthy, is, at least, capable of carrying on the respiration.

It has been frequently stated, as if it were of constant occurrence, that the side is enlarged in pneumothorax. It is true that it may be enlarged; but it may be also smaller than natural, and it may be of the same dimensions as the opposite one. There is no rule which can be in this respect depended on.

Mode of measuring the chest.—The simplest and best method of measuring the comparative size of the two sides of the chest appears to be

the following:—A piece of tape is passed round that portion of the chest especially selected for admeasurement, and, while the two extremities of the tape are brought forward, it is marked with ink at the part at which it crosses the spine. The finger of an assistant keeps it fixed at that part, and the ends are then pulled tight, and carefully marked upon the exact spot at which they meet in the centre of the sternum. The space between the marks at the ends, and in the centre of the tape, of course indicates the difference, if any, in the bulk of the two sides.

The examiner will be often much surprised to find how very little this difference really is, though to the eye the enlargement of one side appears to be very decided. He must be especially careful that the tape is even and straight; as a slight twist, or a very little obliquity in its direction, may be sufficient to reduce to nothing the very trifling difference which may really exist.

If now the examiner be desirous to measure the expansibility, or the capability of expansion, of the two sides, he has only to request the assistant to keep his finger firmly pressed upon the mark of the tape upon the spine; to draw the ends moderately tight up to the marks as before; to hold them loosely in his hand, so that they may slip easily; and direct the patient to take a deep inspiration or two. The portion of tape which

slips beyond the mark on each side will be a measure of the absolute expansibility of each.

When, as it indeed often happens, it is thought desirable to attempt to ascertain whether the process of absorption is going on, or whether the fluid is upon the increase,—whether, in fact, the side is progressively getting larger or smaller,—the tape should be carefully preserved for future use and comparison.

This, indeed, really appears to be one of the most important applications of mensuration, as regards effusion. Upon all future admeasurements, the tape will be an index, or a sort of tide-gauge, of the amount of fluid in the pleura, or, at any rate, will correctly indicate the increase or diminution which has taken place in the comparative size of the two sides.

CHAPTER VII.

SUCCUSSION, OR EXAMINATION BY SHAKING.

SUCCUSSION is by far the simplest as it is one of the oldest modes of examining the chest in disease.

When a cask containing beer, or other fluid, is really "full to the bung," or when a glass vial is filled with air alone, no sound is excited by their being shaken. But when the cask contains air as well as beer, or when there is a little water in the vial, a distinct noise is heard when they are respectively agitated, in consequence of the free motion of the different fluids over each other; the vibrations produced in which are communicated to the wood of the cask, or the glass of which the vial is made, and thence to the surrounding air. Similar facts are observed in disease.

When the pleura contains fluid alone, whether it be in large or in small quantity, no sound results from shaking the body. When it contains only air, as in cases of simple pneumothorax, even violent agitation of the trunk is not accompanied with sound. But when, from any cause, air and fluid exist in the pleura at the same time; when, whether it arise from an empyema giving way, and partially emptying itself through, a bronchial

tube, from a phthisical vomica bursting into the pleura, or from an opening with the external air being effected by gangrene of the lung, gas gains admission into the serous membrane, and pleuritis with effusion results; then a very well-marked sound, resembling the rattling, or squashing of fluid in a cask, may very generally be heard upon any violent agitation of the patient's body. This is *Succussion*. It is unnecessary to describe it farther. It is like the noise produced by shaking fluid in a cask. It can scarcely be mistaken.

Mode of effecting Succussion.—Sometimes the mere jerking of the body, or a sudden twist made by the patient himself, is sufficient to produce Succussion, and it may be generally made to appear, by an assistant shaking the body, while the ear of the examiner is applied to the chest. The most common, though somewhat clumsy mode of eliciting the sound, is for the person examining to give the body a good shaking, and then to put the ear down to the chest. But as individuals suffering from the complaints with which pneumothorax is associated, are ordinarily feeble, and often greatly exhausted, the least distressing, and, upon the whole, the most satisfactory mode of eliciting Succussion, is by pressing the ear firmly upon the parietes of the chest, and while grasping the patient's body with one arm, swinging it gently to and fro, and then suddenly stopping the motion. When the sound is

distinctly audible, this simple mode of proceeding will almost certainly produce it.

The sound of Succussion may differ in tone, as in distinctness, according to the comparative amount of the air and fluid present in the cavity. It is loud, deep, and resonant, when the cavity is large, and when the quantity of air is comparatively great. It is more feeble and treble when the cavity is small, or when in a cavity of large size there exists only a small quantity of air.

As it not unfrequently happens in pneumothorax with effusion, particularly in the latter stages of the complaint, that the gas is either absorbed, or driven out of the pleura, in consequence of the gradual increase of the fluid effused, the succussion may gradually decrease in distinctness and resonance, and at length entirely disappear. This usually arises, not, as has been sometimes incorrectly supposed, in consequence of the aperture into the pleura being closed, but in consequence of the serous membrane being quite full of fluid, and therefore incapable of admitting the presence of air.

But is Succussion pathognomonic, or only a characteristic, of pneumothorax with effusion? It is certainly a characteristic, but, as certainly, it is not a pathognomonic sign of the disorder. It has been stated, that a very large vomica containing some fluid, together with the air with which such cavities are ordinarily filled, has on

some very rare occasions been known to have given rise to Succussion, as well as to metallic tinkling. I have never myself known it to arise from such a cause, nor have I any recollection of meeting any individual who had himself heard it under such circumstances. But it is clear that such a case might occur, and it is probable that some such cases have already occurred.

But I have frequently heard that which, without great caution, is exceedingly apt to lead to error,—I mean the exactly similar sound which often arises from shaking the body, when the stomach contains a mixture of air and fluid. Both sounds are often heard and felt by the patient himself, and both are sometimes heard by persons standing near to him, without applying the ear to the chest. The best mode, and in some cases the only mode, of discriminating between the two, is to examine the patient frequently, when the stomach is empty, as well as when it is full, and before as well as after his meals. Other physical signs, and other general symptoms, and the history of the complaint, will of course materially contribute towards the diagnosis; though, without repeated and careful examinations, it is sometimes exceedingly difficult to arrive at certain conclusions upon the matter.

THE END.

THE
LONDON
MEDICAL GAZETTE,
OR
JOURNAL OF PRACTICAL MEDICINE.

LECTURES in the course of publication in the New
Series, commenced May 2.

*** The whole of these Lectures are published under the sanction,
and with the advantage of the revision, of the several Lecturers.*

MR. ARNOTT'S, at Middlesex Hospital.

DR. GEORGE BURROWS', at St. Bartholomew's.

MR. BRANSBY COOPER'S, at Guy's.

DR. FARADAY'S, at the Royal Institution.

DR. PEREIRA'S, at the London Hospital.

MR. ASTON KEY'S, at Guy's.

MR. LAWRENCE'S, at St. Bartholomew's.

MR. B. PHILLIPS', at Westminster Hospital.

DR. JULIUS VOGEL'S (on Microscopical Anatomy,
with numerous engravings) translated by
Mr. W. S. KIRKES, of St. Bartho-
lomew's.

DR. C. J. B. WILLIAMS', at University-College
Hospital.

*** The publication of Mr. ALFRED TAYLOR'S Lec-
tures on Medical Jurisprudence will commence very shortly.*

[The

The *greatly increased size* of the MEDICAL GAZETTE allows of the insertion of a still greater number of

ORIGINAL ARTICLES

than heretofore ; and the Journal continues to be with the Profession a most extensive medium for the communication of Medical and Scientific information. Since the commencement of the New Series, on May 2,

Upwards of One Hundred of the most distinguished Metropolitan and Provincial Physicians and Surgeons have contributed in this department.

The other departments of the Journal consist of—

Hospital and Infirmary Reports.	Medical Intelligence.
Reviews of every new Medical Publication.	Selections from British and Foreign Journals.
Leading Articles.	Tables of the Mortality of the Metropolis.
Proceedings of Societies.	Lists of Gentlemen who pass
Medical Trials and Inquests.	Examinations at the College
Correspondence.	and Hall, &c. &c. &c.

Subscription to the Stamped Edition, 1*l.* 10*s.* per annum, or 7*s.* 6*d.* per quarter, payable in advance.

Orders for the Stamped Edition received by all Newsvenders.

PUBLISHED EVERY FRIDAY MORNING, BY
LONGMAN, BROWN, GREEN, AND LONGMANS;
ALSO IN MONTHLY PARTS.

October, 1845.

A CATALOGUE OF NEW WORKS AND NEW EDITIONS

PRINTED FOR
LONGMAN, BROWN, GREEN, AND LONGMANS,
LONDON.

ANALYTICAL INDEX.

AGRICULTURE & RURAL AFFAIRS.

	Pages
Bayldon on Valuing Rents, etc.	5
Crocker's Land Surveying	9
Davy's Agricultural Chemistry	9
Greenwood's (Col.) Tree-Lifter	12
Hannam On Waste Manures	13
Johnson's Farmer's Encyclopædia	16
London's Encyclopædia of Agriculture	18
" Self-Instruction for Young Farmers, etc.	18
" (Mrs.) Lady's Country Companion	18
Low's Breeds of the Domesticated Animals of Great Britain	19
" Elements of Agriculture	19
" On Landed Property	19
" On the Domesticated Animals	19

ARTS, MANUFACTURES, AND ARCHITECTURE.

Brande's Dictionary of Science, Literature, and Art	7
Budge's Miner's Guide	7
De Burtin on the Knowledge of Pictures	9
Gwilt's Encyclopædia of Architecture	12
Haydon's Lectures on Painting & Design	13
Holland's Manufactures in Metal	13
Loudon's Encyclopædia of Cottage, Farm, and Villa Architecture and Furniture	18
Porter's Manufacture of Silk	24
" Porcelain & Glass	24
Reid (Dr.) on Warming and Ventilating	25
Steam Engine (The), by the Artisan Club	28
Ure's Dictionary of Arts, Manufactures, and Mines	31
" Recent Improvements in Arts, Manufactures, and Mines	31

BIOGRAPHY.

Aikin's Life of Addison	5
Bell's Lives of the most Eminent British Poets	6
Dover's Life of the King of Prussia	10
Dunham's Lives of the Early Writers of Great Britain	10
" Lives of the British Dramatists	10
Forster's Statesmen of the Commonwealth of England	11
Gleig's Lives of the most Eminent British Military Commanders	11
Grant (Mrs.) Memoir and Correspondence	11
James's Life of the Black Prince	16
" Lives of the most Eminent Foreign Statesmen	16
Leslie's Life of Constable	17
Mackintosh's Life of Sir T. More	20
Maunder's Biographical Treasury	22
Roberts's Life of the Duke of Monmouth	25
Roscoe's Lives of Eminent British Lawyers	26
Russell's Correspondence of the Duke of Bedford	26
Shelley's Lives of the most Eminent Literary Men of Italy, Spain, and Portugal	27
" Lives of the most Eminent French Writers	27
Southey's Lives of the British Admirals	27
Waterton's Autobiography and Essays	31

BOOKS OF GENERAL UTILITY.

	Pages
Acton's (Eliza) Cookery Book	5
Black's Treatise on Brewing	6
Collegian's Guide	8
Donovan's Domestic Economy	10
Hand-Book of Taste	13
Hints on Etiquette	13
Hudson's Parent's Hand-Book	15
" Executor's Guide	15
" On Making Wills	15
Loudon's Self Instruction	18
Maunder's Treasury of Knowledge	21
" Scientific and Literary Treasury	22
" Treasury of History	22
" Biographical Treasury	21
" Universal Class-Book	22
Parkes's Domestic Duties	23
Pycroft's Course of English Reading	24
Riddle's English-Latin and Latin-English Dictionaries	25
Short Whist	27
Thomson's Domestic Management of the Sick Room	29
" Interest Tables	30
Tomlins' Law Dictionary	30
Webster's Ency. of Domestic Economy	31

BOTANY AND GARDENING.

Abercrombie's Practical Gardener	5
" and Main's Gardener's Companion	5
Calcott's Scripture Herbal	7
Conversations on Botany	8
Drummond's First Steps to Botany	10
Glendinning On the Culture of the Pine Apple	11
Greenwood's (Col.) Tree-Lifter	12
Henslow's Botany	13
Hoare On Cultivation of the Grape Vine on Open Walls	13
" On the Management of the Roots of Vines	13
Hooker's British Flora	14
" and Taylor's Muscologia Britannica	14
Jackson's Pictorial Flora	15
Knapp's Gramina Britannica	16
Lindley's Theory of Horticulture	18
" Guide to the Orchard and Kitchen Garden	18
" Introduction to Botany	18
" Flora Medica	18
" Synopsis of British Flora	18
Loudon's Hortus Britannicus	19
" Lignosus Londinensis	19
" Encyclopædia of Trees & Shrubs	18
" Gardening	18
" Plants	19
Lindley's Suburban Garden and Villa Companion	19
" Self-Instruction for Young Gardeners, etc.	18
Repton's Landscape Gardening and Landscape Architecture	25
Rivers's Rose Amateur's Guide	25
Roberts on the Vine	25

	Pages
Rogers's Vegetable Cultivator -	26
Schleiden's Scientific Botany -	26
Smith's Introduction to Botany -	27
„ English Flora -	27
„ Compendium of English Flora -	27

CHRONOLOGY.

Blair's Chronological Tables -	6
Calendar (Illuminated) and Diary, 1846 -	15
Nicolas's Chronology of History -	23
Riddle's Ecclesiastical Chronology -	25
Tate's Horatius Restitutus -	29

COMMERCE AND MERCANTILE AFFAIRS

Kane's (Dr.) Industrial Resources of Ireland -	16
Lorimer's Letters to a Young Master Mariner -	18
M'Culloch's Dictionary of Commerce and Commercial Navigation -	20
Steel's Shipmaster's Assistant -	23
Thomson's Tables of Interest -	

GEOGRAPHY AND ATLASES.

Butler's Sketch of Ancient and Modern Geography -	7
„ Atlas of Modern Geography -	7
„ „ Ancient Geography -	7
Cooley's World Surveyed -	8
De Strzelecki's New South Wales -	10
Forster's Historical Geography of Arabia -	11
Hall's New General Atlas -	12
M'Culloch's Geographical Dictionary -	20
Malte-Brun's Geography -	20
Murray's Encyclopedia of Geography -	23
Parrot's Ascent of Mount Ararat -	8

HISTORY AND CRITICISM.

Adair's (Sir R.), Memoir of a Mission to Vienna -	5
„ Negotiations for the Peace of the Dardanelles -	5
Addison's History of the Knights Templars -	5
Bell's History of Russia -	6
Blair's Chron. and Historical Tables -	6
Bloomfield's Translation of Thucydides -	6
„ Edition of Thucydides -	6
Bunsen's Egypt -	7
Cooley's History of Maritime and Inland Discovery -	8
Crew's History of France -	9
Dahlmann's English Revolution -	9
Dunham's History of Spain and Portugal -	10
„ History of Europe during the Middle Ages -	10
„ History of the German Empire -	10
„ History of Denmark, Sweden, and Norway -	10
„ History of Poland -	10
Dunlop's History of Fiction -	10
Fergus's History of United States of America -	11
Grant (Mrs.) Memoir and Correspondence -	11
Grattan's History of Netherlands -	11
Guicciardini's Hist. Maxims -	12
Halsted's Life of Richard III. -	12
Haydon's Lectures on Painting and Design -	13
Horsley's (Bp.) Biblical Criticism -	14
Jeffrey's (Lord) Contributions to the Edinburgh Review -	16
Keightley's Outlines of History -	16
Laing's Kings of Norway -	16
Lemprière's Classical Dictionary -	17
Macaulay's Essays contributed to the Edinburgh Review -	20
Mackinnon's History of Civilisation -	20
Mackintosh's History of England -	20
„ Miscellaneous Works -	20

	Pages
M'Culloch's Dictionary, Historical, Geographical, and Statistical -	20
Maunder's Treasury of History -	22
Milner's Church History -	22
Moore's History of Ireland -	22
Müller's Mythology -	23
Nicolas's Chronology of History -	23
Ranke's History of the Reformation -	25
Roberts's Rebellion, etc. of the Duke of Monmouth -	25
Rome, History of -	26
Russell's Correspondence of the Duke of Bedford -	6
Scott's History of Scotland -	26
Sismondi's History of the Fall of the Roman Empire -	27
„ History of the Italian Republics -	27
Stebbing's History of the Christian Church -	28
„ History of the Reformation -	28
„ Church History -	28
Switzerland, History of -	29
Sydney Smith's Works -	27
Thirlwall's History of Greece -	30
Tooke's History of Prices -	30
Turner's History of England -	31
Zumpt's Latin Grammar -	32

JUVENILE BOOKS.*Including Mrs. Marcet's Works.*

Boy's (the) Own Book -	6
Hawes's Tales of the N. American Indians -	13
Howitt's (M.) Child's Picture and Verse Book -	14
„ (W.) Jack of the Mill -	14
„ Boy's Country Book -	15
Mackintosh's (Sir Jas.) Life of Sir T. More -	20
Marcet's Conversations—	
On the History of England -	20
On Chemistry -	21
On Natural Philosophy -	21
On Political Economy -	21
On Vegetable Physiology -	21
On Land and Water -	21
On Language -	21
Marcet's Game of Grammar -	21
„ Willy's Grammar -	21
„ Lessons on Animals, etc. -	21
Marryat's Masterman Ready -	21
„ Settlers in Canada -	21
„ Mission; or, Scenes in Africa -	21
Maunder's Universal Class Book -	22
Pycroft's (the Rev. J.), English Reading -	24
Summerly's (Mrs. Felix) Mother's Primer -	28
Uncle Peter's Fairy Tales -	31

MEDICINE.

Bull's Hints to Mothers -	7
„ Management of Children -	7
Copland's Dictionary of Medicine -	8
Elliotson's Human Physiology -	10
Holland's Medical Notes -	13
Lefevre (Sir Geo.) on the Nerves -	17
Marx and Willis (Drs.) On Decrease of Disease -	21
Percira On Food and Diet -	24
Reece's Medical Guide -	25
Sandby On Mesmerism -	26
Wigan (Dr.) On Insanity -	32

MISCELLANEOUS

Black's Treatise on Brewing -	6
Bray's Philosophy of Necessity -	7
Clavers's Forest Life -	8
Collegian's Guide -	8
Colton's Lacon -	8
De Burtin On the Knowledge of Pictures -	9
De Morgan On Probabilities -	9
De Strzelecki's New South Wales -	10
Dunlop's History of Fiction -	10

	Pages
Good's Book of Nature - - -	11
Graham's English - - -	11
Grant's Letters from the Mountains -	11
Guest's Mabinogion - - -	12
Hand-Book of Taste - - -	13
Hobbes (Thos.), English Works of -	13
Holland's Progressive Education -	13
Howitt's Rural Life of England -	14
" Visits to Remarkable Places -	14
" Student-Life of Germany -	15
" Rural and Social Life of Germany -	14
" Colonization and Christianity -	15
" German Experiences -	14
Humphreys' Illuminated Books -	15
Illuminated Calendar and Diary for 1845 -	15
Jeffrey's (Lord) Contributions to the Edinburgh Review - - -	15
Lefevre (Sir Geo.) On the Nerves -	17
Life of a Travelling Physician - -	17
Loudon's (Mrs.) Lady's Country Companion -	18
Macaulay's Critical and Historical Essays -	19
Mackintosh's (Sir J.) Miscellaneous Works -	19
Marx and Willis (Drs.) On Decrease of Disease - - -	21
Michelet's Priests, Women, and Families -	22
Müller's Mythology - - -	23
Pycroft's English Reading - - -	24
Sandby On Mesmerism - - -	26
Sandford's Parochialia - - -	26
Seaward's (Sir E.) Narrative of his Ship- wreck - - -	26
Smith's (Rev. Sydney) Works - - -	27
Summerly's (Mrs. Felix) Mother's Primer -	28
Taylor's Statesman - - -	29
Walker's Chess Studies - - -	31
Welsford On the English Language -	32
Wigan (Dr.) On Insanity - - -	32
Willoughby's (Lady) Diary - - -	32
Zumpt's Latin Grammar - - -	32

NATURAL HISTORY IN GENERAL.

Catlow's Popular Conchology - - -	8
Doubleday's Butterflies and Moths -	10
Gray's Figures of Molluscous Animals -	12
" Mammalia - - -	12
" and Mitchell's Ornithology -	12
Kirby and Spence's Entomology - -	16
Lee's Taxidermy - - -	17
" Elements of Natural History -	17
Marcet's Conversations on Animals, etc. -	21
Newell's Zoology of the English Poets -	23
Proceedings of the Zoological Society -	24
Stephens's British Coleoptera - - -	28
Swainson on the Study of Natural History -	29
" Animals - - -	29
" Quadrupeds - - -	29
" Birds - - -	29
" Animals in Menageries -	29
" Fish, Amphibians, & Reptiles -	29
" Insects - - -	29
" Malacology - - -	29
" the Habits and Instincts of Animals - - -	29
" Taxidermy - - -	29
Transactions of the Zoological Society -	30
Turton's Shells of the British Islands -	31
Waterton's Essays on Natural History -	31

NOVELS AND WORKS OF FICTION.

Bray's (Mrs.) Novels - - -	7
Doctor (the) - - -	10
Dunlop's History of Fiction - - -	10
Howitt's (Mary) Diary - - -	14
" " Home - - -	14
" " Neighbours - - -	14
" " President's Daughters -	14
" " The H—— Family, etc. -	14
Marryat's Masterman Ready - - -	21
" Settlers in Canada - - -	21

Marryat's Mission; or, Scenes in Africa -	21
Willis's (N. P.) Dashes at Life - -	32

ONE VOLUME ENCYCLOPÆDIAS AND DICTIONARIES.

Blaine's Encyclopædia of Rural Sports -	6
Brande's Dictionary of Science, Litera- ture, and Art - - -	6
Copland's Dictionary of Medicine - -	8
Gwilt's Encyclopædia of Architecture -	12
Johnson's Farmer's Encyclopædia -	16
Loudon's Encyclopædia of Trees & Shrubs -	18
" Encyclopædia of Gardening -	19
" Encyclopædia of Agriculture -	19
" Encyclopædia of Plants - -	19
" Rural Architecture - - -	19
M'Culloch's Dictionary, Geographical, Statistical, and Historical -	20
" Dictionary, Practical, Theo- retical, etc. of Commerce -	20
Murray's Encyclopædia of Geography -	23
Ure's Dictionary of Arts, Manufactures, and Mines - - -	31
Webster's Encyclopædia of Dom. Economy -	31

POETRY AND THE DRAMA.

Alkin's (Dr.) British Poets - - -	27
Bowdler's Family Shakspeare - - -	27
Chaloner's Walter Gray - - -	8
" Poetical Remains - - -	8
Costello's Persian Rose Garden - - -	9
Goldsmith's Poems - - -	11
Horace, by Tate - - -	29
L. E. L.'s Poetical Works - - -	17
Macaulay's Lays of Ancient Rome - -	20
Montgomery's Poetical Works - - -	22
Moore's Poetical Works - - -	22
" Lalla Rookh - - -	22
" Irish Melodies - - -	22
Moral of Flowers - - -	23
Nisbet's (Jas.) French in Rheinstadt, etc. -	23
Reynard the Fox - - -	25
Southey's Poetical Works - - -	28
" British Poets - - -	27
Spirit of the Woods - - -	28
Thomson's Seasons - - -	30
Turner's Richard III. - - -	31
Watts's (A. A.) Lyrics of the Heart -	31

POLITICAL ECONOMY AND STATISTICS.

Kane's (Dr.) Industrial Resources of Ireland - - -	16
M'Culloch's Geographical, Statistical, and Historical Dictionary - - -	20
M'Culloch's Literature of Polit. Economy -	20
" On Taxation and Funding -	20
Strong's Greece as a Kingdom - - -	28
Tooke's History of Prices - - -	30

RELIGIOUS AND MORAL WORKS, ETC.

Amy Herbert, edited by Prof. Sewell -	5
Bailey's Essays on the Pursuit of Truth -	5
Bloomfield's Greek Testament - - -	6
" College and School ditto -	6
" Greek and English Lexicon to New Testament - - -	6
Burder's Oriental Customs - - -	7
Burus's Christian Philosophy - - -	7
" Christian Fragments - - -	7
Calcott's Scripture Herbal - - -	7
Cooper's Sermons - - -	8
Dibdin's Sunday Library - - -	29
Doddridge's Family Expositor - - -	10
Englishman's Hebrew and Chaldee Con- cordance to the Bible - - -	10
" Greek Concordance to the New Testament - - -	11
Fitzroy's (Lady) Scripture Conversations -	11
Forster's Historical Geography of Arabia -	11

	Pages
Forster's Life of Bishop Jebb - - -	11
Gertrude - - -	11
Hook's (Dr.) Lectures on Passion Week	13
Horne's Introduction to the Study of the	
Scriptures - - -	14
Abridgment of ditto - - -	14
Horsley's (Bp.) Biblical Criticism - -	14
Psalms - - -	14
Jebb's (Bp.) Practical Theology - - -	16
Pastoral Instructions - - -	16
Correspondence with Knox - - -	16
Knox's (Alexander) Remains - - -	16
Keon's History of the Jesuits - - -	16
Laing's Notes on the German Catholic	
Schism - - -	16
Marriage Gift - - -	21
Micheler's Priests, Women, and Families	22
Milner's Church History - - -	22
Parables (The) - - -	23
Parkes's Domestic Duties - - -	23
Riddle's Letters from a Godfather - -	25
Robinson's Greek and English Lexicon	
to the New Testament - - -	26
Sandford On Female Improvement - - -	26
On Woman - - -	26
's Parochialia - - -	26
Sermon on the Mount (The) - - -	27
Smith's Female Disciple - - -	27
Spalding's Philosophy of Christian Morals	28
Stebbing's Church History - - -	28
Tate's History of St. Paul - - -	29
Taylor's (Rev. C.B.) Margaret; or, the Pearl	29
" Sermons - - -	29
" Dora Melder - - -	29
" Lady Mary - - -	29
Tomline's Christian Theology - - -	30
Turner's Sacred History - - -	31
Wardlaw On Socinian Controversy - - -	32
Weil's Bible, Koran, and Talmud - - -	32
Wilberforce's View of Christianity - -	32
Willoughby's (Lady) Diary - - -	32

RURAL SPORTS.

Blaine's Dictionary of Sports - - -	6
Hansard's Fishing in Wales - - -	13
Hawker's Instructions to Sportsmen - -	13
London's (Mrs.) Lady's Country Companion	18
Stable Talk and Table Talk - - -	28
Thacker's Coursing Rules - - -	29
" Coursing's Remembrancer - - -	29

THE SCIENCES IN GENERAL, AND MATHEMATICS.

Bakewell's Introduction to Geology - -	5
Balmain's Lessons on Chemistry - - -	5
Brande's Dictionary of Science, Litera-	
ture, and Art - - -	6
Brewster's Optics - - -	7
Conversations on Mineralogy - - -	8
De la Beche on the Geology of Cornwall, etc.	9
Donovan's Chemistry - - -	10
Farey on the Steam Engine - - -	11
Fosbroke on the Arts of the Greeks and	
Romans, etc. - - -	11
Greener on the Gun - - -	12
Herschel's Natural Philosophy - - -	13
Astronomy - - -	13
Holland's Manufactures in Metal - - -	13
Hunt's Researches on Light - - -	15
Kane's Elements of Chemistry - - -	16
Kater and Lardner's Mechanics - - -	16
Lardner's Cabinet Cyclopædia - - -	17
Hydrostatics and Pneumatics - - -	17
" and Walker's Electricity - - -	17
Lardner's Arithmetic - - -	17
Geometry - - -	17
Treatise on Heat - - -	17
Lectures On Polarised Light - - -	17
Lloyd On Light and Vision - - -	18

	Pages
Mackenzie's Physiology of Vision - -	20
Marcel's (Mrs.) Conversations on the	
Sciences, etc. - - -	21
Moseley's Practical Mechanics - - -	23
Engineering and Architecture - - -	23
Narrien's Elements of Geometry - - -	26
Astronomy and Geodesy - - -	26
Owen's Lectures On Comparative Anatomy	23
Parnell On Roads - - -	23
Pearson's Practical Astronomy - - -	24
Peschel's Physics - - -	24
Phillips's Palæozoic Fossils of Cornwall, etc.	24
Guide to Geology - - -	24
Treatise on Geology - - -	24
Introduction to Mineralogy - - -	24
Portlock's Report on the Geology of	
Londonderry - - -	24
Powell's Natural Philosophy - - -	24
Quarterly Journal of the Geological Society	25
Ritchie (Robert) on Railways - - -	25
Roberts's Dictionary of Geology - - -	26
Sandhurst Mathematical Course - - -	26
Scoresby's Magnetical Investigations - -	26
Scott's Arithmetic and Algebra - - -	26
Trigonometry - - -	26
Thomson's Algebra - - -	30
Wilkinson's Engines of War - - -	32

TOPOGRAPHY & GUIDE BOOKS

Addison's History of the Temple Church	5
Guide to ditto - - -	5
Costello's (Miss) North Wales - - -	9
Howitt's German Experiences - - -	15
(R.) Australia Felix - - -	15

TRANSACTIONS OF SOCIETIES.

Transactions of the Entomological Society	30
Zoological Society - - -	30
Linnæan Society - - -	30
Geological Society of London - - -	30
Institution of Civil Engineers - - -	30
Royal Institute of British Architects	30
Proceedings of the Zoological Society -	24
Quarterly Journal of the Geological Society	
of London - - -	25

TRAVELS.

Cooley's World Surveyed - - -	8
Costello's (Miss) North Wales - - -	9
De Cistine's Russia - - -	9
De Strzelecki's New South Wales - - -	10
Erman's Travels through Siberia - - -	8
Harris's Highlands of Ethiopia - - -	13
Howitt's Wanderings of a Journeyman	
Tailor - - -	15
German Experiences - - -	14
(R.) Australia Felix - - -	15
Laing's Notes of a Traveller - - -	17
Residence in Norway - - -	17
Tour in Sweden - - -	17
Life of a Travelling Physician - - -	18
Parrot's Ascent of Mount Ararat - - -	8
Paton's (A.A.) Servia - - -	23
Modern Syrians - - -	23
Postans's Sindh - - -	24
Seaward's Narrative of his Shipwreck - -	27
Strong's Greece as a Kingdom - - -	28
Von Orlich's Travels in India - - -	31

VETERINARY MEDICINE

Field's Veterinary Records - - -	11
Morton's Veterinary Toxicological Chart	23
Medicine - - -	23
Percivall's Hippopathology - - -	24
Anatomy of the Horse - - -	24
Spooner on the Foot and Leg of the Horse	28
Stable Talk and Table Talk - - -	28
Turner On the Foot of the Horse - - -	31
White's Veterinary Art - - -	32
Cattle Medicine - - -	32

CATALOGUE.

ABERCROMBIE.—ABERCROMBIE'S PRACTICAL GARDENER, AND IMPROVED SYSTEM OF MODERN HORTICULTURE, alphabetically arranged. 4to. Edition, with an Introductory Treatise on Vegetable Physiology, and Plates by W. Salisbury. 12mo. 6s. boards.

ABERCROMBIE AND MAIN.—THE PRACTICAL GARDENER'S COMPANION; Or, Horticultural Calendar: to which is added, the Garden-Seed and Plant Estimate. Edited from a MS. of J. Abercrombie, by J. Main. 8th Edition. 32mo. 3s. 6d. sewed.

ACTON (MISS).—MODERN COOKERY, In all its Branches, reduced to a System of Easy Practice. For the use of Private Families. In a Series of Practical Receipts, all of which have been strictly tested, and are given with the most minute exactness. Dedicated to the Young Housekeepers of England. By Eliza Acton. 2d. Edition, improved. Foolscap 8vo. with Woodcuts, 7s. 6d. cloth.

"Miss Eliza Acton may congratulate herself on having composed a work of great utility, and one that is speedily finding its way to every 'dresser' in the kingdom. Her Cookery-book is unquestionably the most valuable compendium of the art that has yet been published."
Morning Post.

ADAIR (SIR ROBERT).—AN HISTORICAL MEMOIR OF A MISSION TO THE COURT OF VIENNA IN 1806. By the Right Honorable Sir Robert Adair, G.C.B. With a Selection from his Despatches, published by permission of the proper Authorities. 8vo. 18s. cloth.

ADAIR (SIR ROBERT).—THE NEGOTIATIONS FOR THE PEACE OF THE DARDANELLES, in 1808-9; with Despatches and Official Documents. By the Right Honorable Sir Robert Adair, G.C.B. Being a Sequel to the Memoir of his Mission to Vienna in 1806. 2 vols. 8vo. 28s. cloth.

ADDISON.—THE KNIGHTS TEMPLARS.

By C. G. Addison, of the Inner Temple. 2d Edition, enlarged. Square crown 8vo. with Illustrations, 18s. cloth.

ADDISON.—THE TEMPLE CHURCH IN LONDON:

Its History and Antiquities. By C. G. Addison, Esq., of the Inner Temple, author of "The History of the Knights Templars." Square crown 8vo. with 6 Plates, 5s. cloth.

Also,

A FULL AND COMPLETE GUIDE, HISTORICAL AND DESCRIPTIVE, TO THE TEMPLE CHURCH. (From Mr. Addison's "History of the Temple Church.") Square crown 8vo. 1s. sewed.

AIKIN.—THE LIFE OF JOSEPH ADDISON.

Illustrated by many of his Letters and Private Papers never before published. By Lucy Aikin. 2 vols. post 8vo. with Portrait from Sir Godfrey Kneller's Picture, 18s. cloth.

AMY HERBERT.

By a Lady. Edited by the Rev. William Sewell, B.D. of Exeter College, Oxford. A New Edition. 2 vols. foolscap 8vo. 9s. cloth.

BAILEY.—ESSAYS ON THE PURSUIT OF TRUTH,

And on the Progress of Knowledge. By Samuel Bailey, author of "Essays on the Formation and Publication of Opinions," "Berkeley's Theory of Vision," etc. 2d Edition, revised and enlarged. 8vo. 9s. 6d. cloth.

BAKEWELL.—AN INTRODUCTION TO GEOLOGY.

Intended to convey Practical Knowledge of the Science, and comprising the most important recent Discoveries; with Explanations of the Facts and Phenomena which serve to confirm or invalidate various Geological Theories. By Robert Bakewell. Fifth Edition, considerably enlarged. 8vo. with numerous Plates and Woodcuts, 21s. cloth.

BALMAIN.—LESSONS ON CHEMISTRY,

For the Use of Pupils in Schools, Junior Students in Universities, and Readers who wish to learn the fundamental Principles and leading Facts: with Questions for Examination, Glossaries of Chemical Terms and Chemical Symbols, and an Index. By William H. Balmain. With numerous Woodcuts, illustrative of the Decompositions. Foolscap 8vo. 6s. cloth.

BAYLDON.—ART OF VALUING RENTS AND TILLAGES,

And the Tenant's Right of Entering and Quitting Farms, explained by several Specimens of Valuations; and Remarks on the Cultivation pursued on Soils in different Situations. Adapted to the Use of Landlords, Land-Agents, Appraisers, Farmers, and Tenants. By J. S. Bayldon. 6th Edition, corrected and revised by John Donaldson, Land-Steward, author of a "Treatise on Manures and Grasses." 8vo. 10s. 6d. cloth.

BEDFORD CORRESPONDENCE.—CORRESPONDENCE OF JOHN, FOURTH DUKE OF BEDFORD, selected from the Originals at Woburn Abbey: with Introductions by Lord John Russell. Svo. vol. 1 (1742-48), 18s. cloth; vol. 2 (1749-60), 15s. cl.

"The second volume includes a correspondence having relation to the period from the Peace of Aix-la-Chapelle to the death of George II. Its most remarkable portion bears upon an important question, on which there still exist some differences of opinion, viz. the intrigues which led to the junction of the Duke of Newcastle and Pitt, in 1757. The letters respecting the state of Ireland under the Viceroyalty of the Duke of Bedford also, are not a little interesting."—Morning Herald.

* * Vol. III. to complete the work, is in preparation.

BELL.—LIVES OF THE MOST EMINENT ENGLISH POETS.

By Robert Bell, Esq. 2 vols. foolscap Svo. with Vignette Titles, 12s. cloth.

BELL.—THE HISTORY OF RUSSIA,

From the Earliest Period to the Treaty of Tilsit. By Robert Bell, Esq. 3 vols. foolscap Svo. with Vignette Titles, 18s. cloth.

BLACK.—A PRACTICAL TREATISE ON BREWING.

Based on Chemical and Economical Principles: with Formulæ for Public Brewers, and Instructions for Private Families. By William Black. Third Edition, revised and corrected, with considerable Additions. The Additions revised by Professor Graham, of the London University. Svo. 10s. 6d. cloth.

"I take occasion, in concluding this article, to refer my readers to the 'Practical Treatise on Brewing' by Mr. William Black, a gentleman of much experience in the business. This little work contains a great deal of useful information."

Dr. Ure's Supplement to his "Dictionary."

BLAINE.—AN ENCYCLOPÆDIA OF RURAL SPORTS;

Or, a complete Account, Historical, Practical, and Descriptive, of Hunting, Shooting, Fishing, Racing, and other Field Sports and Athletic Amusements of the present day. By Delabere P. Blaine, Esq., author of "Outlines of the Veterinary Art," "Canine Pathology," etc. etc. With nearly 600 Engravings on Wood, by R. Branston, from Drawings by Alken, T. Landseer, Dickes, etc. 1 thick vol. Svo. 2l. 10s. cloth.

BLAIR'S CHRONOLOGICAL AND HISTORICAL TABLES,

From the Creation to the present Time: with Additions and Corrections from the most authentic Writers; including the Computation of St. Paul, as connecting the Period from the Exode to the Temple. Under the revision of Sir Henry Ellis, K.H., Principal Librarian of the British Museum. Imperial Svo. 3ls. 6d. half-bound morocco.

"The student of history, long accustomed to the Doctor's ponderous and unmanageable folio, will rejoice over this handsome and handy volume. It is the revival and enlargement, in a far more compact and available form than the original, of the celebrated 'Chronological Tables' of Dr. Blair. It comprises additions to our own time, and corrections from the most recent authorities. The outline of the plan is faithfully preserved and carried out, with every improvement of which it was susceptible."—Examiner.

BLOOMFIELD.—THE HISTORY OF THE PELOPONNESIAN WAR.

By Thucydides. Newly Translated into English, and accompanied with very copious Notes, Philological and Explanatory, Historical and Geographical. By the Rev. S. T. Bloomfield, D.D. F.S.A. 3 vols. Svo. with Maps and Plates, 2l. 5s. boards.

BLOOMFIELD.—THE HISTORY OF THE PELOPONNESIAN WAR.

By Thucydides. A New Recension of the Text, with a carefully amended Punctuation; and copious Notes, Critical, Philological, and Explanatory, almost entirely original, but partly selected and arranged from the best Expositors: accompanied with full Indexes, both of Greek Words and Phrases explained, and matters discussed in the Notes. The whole illustrated by Maps and Plans, mostly taken from actual Surveys. By the Rev. S. T. Bloomfield, D.D. F.S.A. 2 vols. Svo. 38s. cloth.

BLOOMFIELD.—THE GREEK TESTAMENT:

With copious English Notes, Critical, Philological, and Explanatory. By the Rev. S. T. Bloomfield, D.D. F.S.A. 5th Edit. improved. 2 vols. Svo. with a Map of Palestine, 40s. cloth.

BLOOMFIELD.—COLLEGE AND SCHOOL GREEK TESTAMENT;

With English Notes. By the Rev. S. T. Bloomfield, D.D. Fourth Edition, enlarged and improved, accompanied with a New Map of Syria and Palestine, adapted to the New Testament and Josephus, and an Index of Greek Words and Phrases explained in the Notes. 12mo. 10s. 6d. cloth.

BLOOMFIELD.—GREEK AND ENGLISH LEXICON TO THE NEW

TESTAMENT: especially adapted to the use of Colleges, and the Higher Classes in Public Schools; but also intended as a convenient Manual for Biblical Students in general. By Dr. Bloomfield. 2d Edition, greatly enlarged, and very considerably improved. 12mo. on wider paper, 10s. 6d. cloth.

BOY'S OWN BOOK (THE):

A Complete Encyclopædia of all the Diversions, Athletic, Scientific, and Recreative, of Boyhood and Youth. 20th Edition. Square 12mo., with many Engravings on Wood, 6s. boards.

BRANDE.—A DICTIONARY OF SCIENCE, LITERATURE, AND ART;

Comprising the History, Description, and Scientific Principles of every Branch of Human Knowledge; with the Derivation and Definition of all the Terms in general use. Edited by W. T. Brande, F.R.S.L. and E.; assisted by J. Cauvin. The various departments are by Gentlemen of eminence in each. 1 very thick vol. Svo. illustrated by Wood-engravings, 3l. cloth.

(BRAY MRS.)—MRS. BRAY'S NOVELS AND ROMANCES,

Revised and corrected by Mrs. Bray. In 10 vols. fcap 8vo., with Frontispieces and Vignettes from Designs and Sketches by the late Thomas Stothard, R.A.; C. A. Stothard, F.S.A.; Henry Warren, Esq.; etc.

Vol. I. "*The White Hoods*," with portrait of the Author, a view of her residence, and General Preface to the Series;—Vol. II. "*De Foix*";—Vol. III. "*The Protestant*";—Vol. IV. "*Fitz of Fitz-Ford*";—Vol. V. "*The Talba*";—Vol. VI. "*Warleigh*."

. To be continued monthly, and completed in 10 volumes; each containing an entire Work, printed and embellished uniformly with the "*Standard Novels*."

Vols. 7, 8, 9 & 10 will contain—

Nov. 1.—TRELAWNY.

Dec. 1.—TRIALS OF THE HEART.

Jan. 1.—HENRY DE POMEROY.

Feb. 1.—COURTENAY OF WALREDDON.

BRAY.—THE PHILOSOPHY OF NECESSITY;

Or, the Law of Consequences as applicable to Mental, Moral, and Social Science. By Charles Bray. 2 vols. 8vo. 15s. cloth.

BREWSTER.—TREATISE ON OPTICS.

By Sir David Brewster, LL.D. F.R.S. etc. New Edition. Foolscep 8vo. with vignette title, and 176 Woodcuts, 6s. cloth.

BUDGE (J.)—THE PRACTICAL MINER'S GUIDE:

Comprising a Set of Trigonometrical Tables adapted to all the purposes of Oblique or Diagonal, Vertical, Horizontal, and Traverse Dialling; with their application to the Dial, Exercise of Drifts, Lodes, Slides, Levelling, Inaccessible Distances, Heights, etc. By J. Budge. New Edition, considerably enlarged, 8vo. with Portrait of the Author, 12s. cloth.

BULL.—THE MATERNAL MANAGEMENT OF CHILDREN,

IN HEALTH AND DISEASE. By Thomas Bull, M.D. 2d Edition, revised and enlarged. Foolscep 8vo. 7s. cloth.

BULL.—HINTS TO MOTHERS,

For the Management of Health during the Period of Pregnancy and in the Lying-in Room; with an Exposure of Popular Errors in connexion with those subjects. By Thomas Bull, M.D. Physician Accoucheur to the Finsbury Midwifery Institution, etc. etc. 4th Edition, revised and considerably enlarged. Foolscep 8vo. 7s. cloth.

"*Excellent guides, and deserve to be generally known.*"

Johnson's Medico-Chirurgical Review.

BUNSEN.—AN INQUIRY INTO THE HISTORY, ARTS AND SCIENCES,

LANGUAGE, WRITING, MYTHOLOGY, and CHRONOLOGY of ANCIENT EGYPT: with the peculiar Position of that Nation in reference to the Universal History of Mankind. By the Chevalier C. C. J. Bunsen. Translated from the German, under the Author's superintendence, by C. H. Cottrell, Esq.; with additional matter, furnished by the Author. 2 vols. 8vo. with numerous Plates.—*Preparing for publication.*

BURDER.—ORIENTAL CUSTOMS,

Applied to the Illustration of the Sacred Scriptures. By Samuel Burder, A.M. 3d Edition, with additions. Foolscep 8vo. 8s. 6d. cloth.

BURNS.—THE PRINCIPLES OF CHRISTIAN PHILOSOPHY;

Containing the Doctrines, Duties, Admonitions, and Consolations of the Christian Religion. By John Burns, M.D. F.R.S. 5th Edition. 12mo. 7s. boards.

BURNS.—CHRISTIAN FRAGMENTS;

Or, Remarks on the Nature, Precepts, and Comforts of Religion. By John Burns, M.D. F.R.S. Professor of Surgery in the University of Glasgow, author of "*The Principles of Christian Philosophy*." Foolscep 8vo. 5s. cloth.

"*The author manifests throughout a sound judgment, a cultivated literary taste, and, best of all, a heart deeply impressed with the solemn realities of religion. His sentiments are evangelical, and his spirit devout.*"—Watchman.

BUTLER.—SKETCH OF ANCIENT AND MODERN GEOGRAPHY.

By Samuel Butler, D.D., late Lord Bishop of Lichfield and Coventry; and formerly Head Master of Shrewsbury School. New Edition, revised by his Son, 8vo. 9s. boards.

The present edition has been carefully revised by the author's son, and such alterations introduced as continually progressive discoveries and the latest information rendered necessary. Recent Travels have been constantly consulted where any doubt or difficulty seemed to require it; and some additional matter has been added, both in the ancient and modern part.

BUTLER.—ATLAS OF MODERN GEOGRAPHY.

By the late Dr. Butler. New Edition; consisting of Twenty-three coloured Maps, from a New Set of Plates; with an Index of all the Names of Places, referring to the Latitudes and Longitudes. 8vo. 12s. half-bound.

BUTLER.—ATLAS OF ANCIENT GEOGRAPHY.

By the late Dr. Butler. Consisting of Twenty-three coloured Maps: with an Index of all the Names of Places, referring to the Latitudes and Longitudes. New Edition. 8vo. 12s. half-bound.

. The above two Atlases may be had, in One Volume, 4to. 24s. half-bound.

CALLCOTT.—A SCRIPTURE HERBAL:

With upwards of 120 Wood Engravings. By Lady Callcott. Square crown 8vo. 11. 5s. cloth.

CATLOW.—POPULAR CONCHOLOGY;

Or, the Shell Cabinet Arranged: being an Introduction to the modern System of Conchology; with a sketch of the Natural History of the Animals, an account of the Formation of the Shells, and a complete Descriptive List of the Families and Genera. By Agnes Catlow. Foolscap. 8vo. with 312 Woodcuts, 10s. 6d. cloth.

CHALENOR.—WALTER GRAY,

A Ballad, and other Poems; including the Poetical Remains of Mary Chalenor. 2d Edition, with Additions, fcp. 8vo. 6s. cloth.

CHALENOR.—POETICAL REMAINS OF MARY CHALENOR.

Fcp. 8vo. 4s. cloth.

CLAVERS.—FOREST LIFE.

By Mary Clavers, an Actual Settler; author of "A New Home, Who'll Follow?" 2 vols. fcp. 8vo. 12s. cloth.

COLLEGIAN'S GUIDE (THE);

Or, Recollections of College Days; setting forth the Advantages and Temptations of a University Education. By *****, M.A., — College, Oxford. Post 8vo. 10s. 6d. cloth.

COLTON.—LACON; OR, MANY THINGS IN FEW WORDS.

By the Rev. C. C. Colton. New Edition, 8vo. 12s. cloth.

CONVERSATIONS ON BOTANY.

9th Edition, improved. Foolscap 8vo. with 22 Plates, 7s. 6d. cloth; with the Plates coloured, 12s. cloth.

CONVERSATIONS ON MINERALOGY.

With Plates, engraved by Mr. and Mrs. Lowry, from Original Drawings. 3d Edition, enlarged. 2 vols. 12mo. 14s. cloth.

COOLEY.—THE WORLD SURVEYED IN THE NINETEENTH CENTURY;

Or, Recent Narratives of Scientific and Exploring Expeditions (chiefly undertaken by command of Foreign Governments). Collected, translated, and, where necessary, abridged, by W. D. Cooley, Esq., author of "History of Maritime and Inland Discovery" in the Cabinet Cyclopædia, etc. 8vo.

The First Volume of the Series contains "The Ascent of Mount Ararat." By Dr. Friedrich Parrot, Professor of Natural Philosophy in the University of Dorpat, Russian Imperial Councillor of State, etc. 8vo. with a Map by Arrowsmith, and Woodcuts, 14s. cloth.

*** Each volume will form, for the most part, a Work complete in itself, and the whole Series will present an accurate and luminous picture of all the known portions of the earth.

The Second Work of the Series will be "Erman's Travels through Siberia." 8vo.—In the press.

*** On this traveller, the President of the Royal Geographical Society, in his anniversary address last year, bestowed the following encomium: "If we regard M. Adolph Erman as an astronomical geographer and explorer of distant lands, we must all admit that he stands in the very highest rank." And in his address delivered in May last, the President again made honourable mention of this traveller in the following terms: "In announcing to you with pleasure that the excellent work of your distinguished foreign member and medallist, Adolph Erman, is about to appear in English, I must not lose the opportunity of stating, that the last communication sent to us by M. Erman is one of very great importance."

"We hasten to apprise English readers that they have now an opportunity of making personal acquaintance with a book that has for years been an European celebrity; and we cordially welcome the first of a series to which the able editorship of Mr. Cooley must ensure an extensive and distinguished reputation."—Morning Post.

"The commencement of what promises to be a most admirable series of books, conducted by one of the most able and competent of living geographers."—Examiner.

COOLEY.—THE HISTORY OF MARITIME AND INLAND DISCOVERY.

By W. D. Cooley, Esq. 3 vols. foolscap 8vo. with Vignette Titles, 18s. cloth.

COOPER (REV. E.)—SERMONS,

Chiefly designed to elucidate some of the leading Doctrines of the Gospel. To which is added, an Appendix, containing Sermons preached on several Public Occasions, and printed by desire. By the Rev. Edward Cooper, Rector of Hamstall-Ridware, and of Yoxall, in the County of Stafford; and late Fellow of All-Souls' College, Oxford. 7th Edition. 2 vols. 12mo. 10s. boards.

By the same Author.

PRACTICAL AND FAMILIAR SERMONS, designed for Parochial and Domestic Instruction. New Editions. 7 vols. 12mo. 1l. 18s. boards.

*** Vols. 1 to 4, 5s. each; Vols. 5 to 7, 6s. each.

COPLAND.—A DICTIONARY OF PRACTICAL MEDICINE;

Comprising General Pathology, the Nature and Treatment of Diseases, Morbid Structures, and the Disorders especially incidental to Climates, to Sex, and to the different Epochs of Life, with numerous approved Formulæ of the Medicines recommended. By James Copland, M.D., Consulting Physician to Queen Charlotte's Lying-in Hospital; Senior Physician to the Royal Infirmary for Children; Member of the Royal College of Physicians, London; of the Medical and Chirurgical Societies of London and Berlin, etc. Vols. 1 and 2, 8vo. 3l. cloth; and Part 10, 4s. 6d. sewed.

*** To be completed in one more Volume.

COSTELLO (MISS).—FALLS, LAKES, AND MOUNTAINS OF NORTH WALES; being a Pictorial Tour through the most interesting parts of the Country. By Louisa Stuart Costello, author of "A Summer among the Bocages and Vines," "A Pilgrimage to Auvergne," "Bearn and the Pyrenees," etc. Profusely illustrated with Views, from Original Sketches by D. H. McKewan, engraved on wood, and lithographed, by T. and E. Gilks. Square 8vo. with Map, 14s. cloth, gilt edges.

"We have visited many of the places here written of by Miss Costello, and can bear good testimony to the general excellence of her work; to its usefulness as a 'Hand-Book for Travellers in Wales;' and to the truthful sketches it contains, literary and pictorial. No one who intends visiting the falls, lakes, and mountains of North Wales, should depart unprovided with this most admirable 'Pictorial Guide.'"—Atlas.

COSTELLO (MISS).—THE ROSE GARDEN OF PERSIA.

A Series of Translations from the Persian Poets. By Miss Louisa Stuart Costello, author of "Specimens of the Early Poetry of France," "A Summer amongst the Bocages and the Vines," etc. etc. 8vo. with Borders printed in Gold and Colours. [In October.

CROCKER'S ELEMENTS OF LAND SURVEYING.

Fifth Edition, corrected throughout, and considerably improved and modernized, by T. G. Bunt, Land Surveyor, Bristol. To which are added, TABLES OF SIX-FIGURE LOGARITHMS, etc., superintended by Richard Farley, of the Nautical Almanac Establishment. Post 8vo. 12s. cloth.

"* The work throughout is entirely revised, and much new matter has been added; there are new chapters, containing very full and minute Directions relating to the modern Practice of Surveying, both with and without the aid of angular instruments. The method of Plotting Estates, and casting or computing their Areas, is described, etc. etc. The chapter on Levelling also is new.

CROWE.—THE HISTORY OF FRANCE.

From the Earliest Period to the Abdication of Napoleon. By E. E. Crowe, Esq. 3 vols. foolscap 8vo. with Vignette Titles, 18s. cloth.

DAHLMANN.—HISTORY OF THE ENGLISH REVOLUTION.

By F. C. Dahlmann, late Professor of History at the University of Gottingen. Translated from the German, by H. Evans Lloyd. 8vo. 10s. 6d. cloth.

"Professor Dahlmann's book is, in short, a rapid sketch of the whole of what we call the Modern History of England, from its start at the Coronation of Henry the Seventh, to its intermediate settlement at the Coronation of William the Third. We have no English summary of the history it relates so brief, compendious, and impartial. M. Dahlmann is a very earnest as well as intelligent writer; and the steady advance of the popular principle in England, through an almost uninterrupted march of two centuries, is startlingly reflected in his clear and transparent relation. Mr. Lloyd's translation is very well executed."—Examiner.

DAVY (SIR HUMPHRY).—ELEMENTS OF AGRICULTURAL CHEMISTRY

in a Course of Lectures. By Sir Humphry Davy. With Notes by Dr. John Davy. 6th Edition. 8vo. with 10 Plates, 15s. cloth.

Introduction; The General Powers of Matter which Influence Vegetation: the Organization of Plants; Soils; Nature and Constitution of the Atmosphere, and its Influence on Vegetables; Manures of Vegetable and

Animal Origin; Manures of Mineral Origin, or Fossil Manures; Improvement of Lands by Burning; Experiments on the Nutritive Qualities of different Grasses, etc.

DE BURTIN.—A TREATISE ON THE KNOWLEDGE NECESSARY TO

AMATEURS OF PICTURES. Translated and abridged from the French of M. Francis Xavier de Burtin, First Stipendiary Member of the Royal Academy of Brussels in the Class of Sciences, etc. By Robert White, Esq. 8vo. with 4 Lithographic Engravings, 12s. cloth.

"Mons. De Burtin's whole life has been devoted to the study and acquisition of works of art, and his practical knowledge of every thing connected with the subject is equal to his enthusiasm. He treats of the several qualities that go to make up a good picture, of the characteristics of the different schools and leading masters, and the signatures and prices of their pictures; pointing out the way to judge of their quality, condition, and originality; and describing the best methods of cleaning and preserving them. He also gives a glance at the principal public galleries, and useful hints towards forming and arranging private collections."—Spectator.

DE CUSTINE.—RUSSIA.

By the Marquis De Custine. Translated from the French. 2d Edition. 3 vols. post 8vo. 31s. 6d. cloth.

"A work which those who are desirous to know Russia as it really is, and not as it would fain impose itself on the world to be, would do well to consult. We promise our readers equal surprise and pleasure from the perusal of Mons. De Custine's very clever book."—Gent's Mag.

DE LA BECHE.—REPORT ON THE GEOLOGY OF CORNWALL, DEVON,

AND WEST SOMERSET. By Henry T. De la Beche, F.R.S. etc., Director of the Ordnance Geological Survey. Published by Order of the Lords Commissioners of H. M. Treasury. 8vo. with Maps, Woodcuts, and 12 large Plates, 14s. cloth.

DE MORGAN.—AN ESSAY ON PROBABILITIES,

And on their Application to Life Contingencies and Insurance Offices. By Aug. De Morgan, of Trinity College, Cambridge. Foolscap 8vo. with Vignette Title, 6s. cloth.

DE STRZELECKI (P. E.).—PHYSICAL DESCRIPTION OF NEW SOUTH WALES AND VAN DIEMAN'S LAND. Accompanied by a Geological Map, Sections, and Diagrams, and Figures of the Organic Remains. By P. E. De Strzelecki. 8vo. with coloured Map and numerous Plates, 24s. cloth.

"This is the first systematic and complete natural or physical history of the great southern region which has lately been added to the civilized world. The author has been a traveller and voyager round the globe for twelve years, several of which have been devoted to Australia, in exploring which he seems to have been subjected to considerable peril. In these countries he made a tour of 7000 miles on foot. We recommend his work to all interested in the Australian colonies, or in science for its own sake, as the only existing scientific account of the New World."—Tait's Magazine.

DOCTOR (THE), ETC.

5 vols. post 8vo. 2l. 12s. 6d. cloth.

"Admirably as the mystery of the 'Doctor' has been preserved up to the present moment, there is no longer any reason for affecting secrecy on the subject. The author is Robert Southey; he acknowledged the fact shortly before his last illness to his most confidential friend, an M.P. of high character. In a private letter from Mrs. Southey, dated February 27, 1843, she not only states the fact, but adds that the greater part of a sixth volume had gone through the press, and that Southey looked forward to the pleasure of drawing her into it as a contributor; giving her full authority to affirm that her husband is the author."

Robert Bell, Esq., in *The Story Teller*.

DODDRIDGE.—THE FAMILY EXPOSITOR;

Or, a Paraphrase and Version of the New Testament: with Critical Notes, and a Practical Improvement of each Section. By P. Doddridge, D.D. To which is prefixed, a Life of the Author, by A. Kippis, D.D. F.R.S. and S.A. New Edition. 4 vols. 8vo. 1l. 16s. cloth.

DONOVAN.—TREATISE ON CHEMISTRY.

By Michael Donovan, Esq. M.R.I.A. 4th Edition. Fcap. 8vo. with Vignette Title, 6s. cloth.

DONOVAN.—A TREATISE ON DOMESTIC ECONOMY.

By M. Donovan, Esq. M.R.I.A., Professor of Chemistry to the Company of Apothecaries in Ireland. 2 vols. foolscap 8vo. with Vignette Titles, 12s. cloth.

DOUBLEDAY'S BUTTERFLIES.—THE GENERA OF DIURNAL LEPI-

DOPTERA; comprising their Generic Characters—a Notice of the Habits and Transformations—and a Catalogue of the Species of each Genus. By Edward Doubleday, Esq. F.L.S. etc., Assistant in the Zoological Department of the British Museum. Imperial 4to. uniform with Gray and Mitchell's Ornithology; illustrated with 75 Coloured Plates.

" To be published in Monthly Parts, 5s. each; each Part to consist of 2 Coloured Plates, with accompanying Letter-press. Publication will commence when 150 Subscribers' Names have been received."*

DOVER.—LIFE OF FREDERICK II. KING OF PRUSSIA.

By Lord Dover. 2d Edition. 2 vols. 8vo. with Portrait, 28s. boards.

DRUMMOND.—FIRST STEPS TO BOTANY,

Intended as popular Illustrations of the Science, leading to its study as a branch of general education. By J. L. Drummond, M.D. 4th Edit. 12mo. with numerous Woodcuts, 9s. boards.

DUNHAM.—THE HISTORY OF THE GERMANIC EMPIRE.

By Dr. Dunham. 3 vols. foolscap 8vo. with Vignette Titles, 18s. cloth.

THE HISTORY OF EUROPE DURING

THE MIDDLE AGES. By Dr. Dunham. 4 vols. foolscap 8vo. with Vignette Titles, 1l. 4s. cloth.

THE HISTORY OF SPAIN AND PORTU-

GAL. By Dr. Dunham. 5 vols. foolscap 8vo. with Vignette Titles, 1l. 10s. cloth.

THE HISTORY OF SWEDEN, DENMARK,

AND NORWAY. By Dr. Dunham. 3 vols. foolscap 8vo. with Vignette Titles, 18s. cloth.

THE HISTORY OF POLAND. By Dr.

Dunham. Foolscap 8vo. with Vignette Title, 6s. cloth.

THE LIVES OF THE EARLY WRITERS

OF GREAT BRITAIN. By Dr. Dunham, R. Bell, Esq., etc. Foolscap 8vo. with Vignette Title, 6s. cloth.

THE LIVES OF BRITISH DRAMATISTS.

By Dr. Dunham, R. Bell, Esq., etc. 2 vols. foolscap 8vo. with Vignette Titles, 12s. cloth.

DUNLOP (JOHN).—THE HISTORY OF FICTION:

Being a Critical Account of the most celebrated Prose Works of Fiction, from the earliest Greek Romances to the Novels of the Present Age. By John Dunlop. 3d Edition, complete in One Volume. Medium 8vo. 15s. cloth.

ELLIOTSON.—HUMAN PHYSIOLOGY:

With which is incorporated much of the Elementary Part of the "Institutiones Physiologicæ" of J. F. Blumenbach, Professor in the University of Göttingen. By John Elliotson, M.D. Cantab. F.R.S. Fifth Edition. 8vo., with numerous Woodcuts, 2l. 2s. cloth.

ENGLISHMAN'S HEBREW AND CHALDEE CONCORDANCE OF

THE OLD TESTAMENT; being an attempt at a Verbal Connexion between the Original and the English Translations: with Indexes, a List of the Proper Names and their occurrences, etc. etc. 2 vols. royal 8vo., 3l. 13s. 6d. cloth; large paper, 4l. 14s. 6d.

"The labour bestowed upon this important work has seldom, we should suppose, been equalled; and we have the fullest conviction, from the merely cursory examination we are able to give to such a stupendous task, that the result justifies all the labour, time, and money expended upon it. Indeed, the whole book bears the most palpable evidence of honest carefulness and unwearied diligence—the points of prime worth in a Concordance; and wherever we have dipped into its pages (about 1800), we have, in every case, had our opinion of its neatness, accuracy, and lucid order, confirmed and increased."—Literary Gazette.

THE ENGLISHMAN'S GREEK CONCORDANCE OF THE NEW TESTAMENT; being an attempt at a Verbal Connexion between the Greek and the English Texts; including a Concordance to the Proper Names, with Indexes, Greek-English and English-Greek. 2d Edition, carefully revised, with a new Index, Greek and English. Royal 8vo. 42s. cloth.

FAREY.—A TREATISE ON THE STEAM-ENGINE, Historical, Practical, and Descriptive. By John Farey, Engineer. 4to. Illustrated by numerous Woodcuts, and 25 Copper-plates, 5*l.* 5*s.* in boards.

FERGUS.—THE HISTORY OF THE UNITED STATES OF AMERICA, From the Discovery of America to the Election of General Jackson to the Presidency. By the Rev. H. Fergus. 2 vols. foolscap 8vo. with Vignette Titles, 12*s.* cloth.

FIELD.—POSTHUMOUS EXTRACTS FROM THE VETERINARY RECORDS OF THE LATE JOHN FIELD. Edited by his Brother, William Field, Veterinary Surgeon, London. 8vo. 8*s.* boards.

FITZROY (LADY).—SCRIPTURAL CONVERSATIONS BETWEEN CHARLES AND HIS MOTHER. By Lady Charles Fitzroy. Foolscap 8vo. 4*s.* 6*d.* cloth.

FORSTER.—STATESMEN OF THE COMMONWEALTH OF ENGLAND. With an Introductory Treatise on the Popular Progress in English History. By John Forster, Esq. 5 vols. foolscap 8vo. with Original Portraits of Pym, Eliot, Hampden, Cromwell, and an Historical Scene after a Picture by Cattermole, 1*l.* 10*s.* cloth.

The Introductory Treatise, intended as an Introduction to the Study of the Great Civil War in the Seventeenth Century, separately, price 2*s.* 6*d.* sewed.

The above 5 vols. form Mr. Forster's Portion of the Lives of Eminent British Statesmen, by Sir James Mackintosh, the Right Hon. T. P. Courtenay, and John Forster, Esq. 7 vols. foolscap 8vo. with Vignette Titles, 2*l.* 2*s.* cloth.

FORSTER (REV. C.).—THE HISTORICAL GEOGRAPHY OF ARABIA; Or, the Patriarchal Evidences of Revealed Religion. A Memoir, with illustrative Maps and an Appendix, containing Translations, with an Alphabet and Glossary of the Hamyaritic Inscriptions recently discovered in Hadramaut. By the Rev. Charles Forster, B.D., one of the Six Preachers in the Cathedral of Christ, Canterbury, and Rector of Stisted, Essex; author of "Mahometanism Unveiled." 2 vols. 8vo. 30*s.* cloth.

FORSTER (REV. C.).—THE LIFE OF JOHN JEBB, D.D. F.R.S. Late Bishop of Limerick. With a Selection from his Letters. By the Rev. Charles Forster, B.D., Rector of Stisted, Essex, and one of the Six Preachers in the Cathedral of Christ, Canterbury, formerly Domestic Chaplain to the Bishop. 2d Edition. 8vo. with Portrait, etc. 16*s.* cloth.

FOSBROKE.—A TREATISE ON THE ARTS, MANNERS, MANUFACTURES, and INSTITUTIONS OF THE GREEKS and ROMANS. By the Rev. T. D. Fosbroke, etc. 2 vols. foolscap 8vo. with Vignette Titles, 12*s.* cloth.

GERTRUDE.

A Tale. By the author of "Amy Herbert." Edited by the Rev. William Sewell, B.D., of Exeter College, Oxford. 2 vols. foolscap 8vo. 9*s.* cloth.

"A book, the influences of which cannot fail to be salutary. Practice without profession, brought to bear upon the vicissitudes of every-day life, the power of forbearance and gentleness, and of sacrifice which is to find in itself its recompense; such are the themes here set forth in an agreeable style and an interesting story. Clear and discriminating glimpses of character, and the absence of bitterness and offence, constitute the great charm of this elegant writer, and warrants us in cordially recommending her 'Gertrude' as pleasant and profitable reading."—*Athenæum*.

GLEIG.—LIVES OF THE MOST EMINENT BRITISH MILITARY COMMANDERS. By the Rev. G. R. Gleig. 3 vols. foolscap 8vo. with Vignette Titles, 18*s.* cloth.

GLENDINNING.—PRACTICAL HINTS ON THE CULTURE OF THE PINEAPPLE. By R. Glendinning, Gardener to the Right Hon. Lord Rolle, Bicton. 12mo. with Plan of Pinery, 5*s.* cloth.

GOLDSMITH'S POETICAL WORKS.

Illustrated with Engravings on Wood, from Designs by the Etching Club. Uniform with "Thomson's Seasons." Square crown 8vo.—*In October*.

GOOD.—THE BOOK OF NATURE.

A Popular Illustration of the General Laws and Phenomena of Creation. By John Mason Good, M.D. F.R.S. etc. 3d Edition, corrected. 3 vols. foolscap 8vo. 24*s.* cloth.

GRAHAM.—ENGLISH; OR, THE ART OF COMPOSITION

explained in a Series of Instructions and Examples. By G. F. Graham. 2d Edition, revised and improved. Foolscap 8vo. 7*s.* cloth.

GRANT (MRS., OF LAGGAN).—MEMOIR AND CORRESPONDENCE of the late Mrs. Grant, of Laggan, author of "Letters from the Mountains," etc. Edited by her Son, J. P. Grant, Esq. 2d Edition. 3 vols. post 8vo. Portrait, 1*l.* 11*s.* 6*d.* cloth.

GRANT (MRS.).—LETTERS FROM THE MOUNTAINS.

Being the Correspondence with her Friends, between the years 1773 and 1803. By Mrs. Grant, of Laggan. 6th Edition. Edited, with Notes and Additions, by her son, J. P. Grant, Esq. 2 vols. post 8vo. 21*s.* cloth.

GRATTAN.—THE HISTORY OF THE NETHERLANDS,

From the Invasion by the Romans to the Belgian Revolution in 1830. By T. C. Grattan, Esq. Foolscap 8vo. with Vignette Titles, 6s. cloth.

GRAY.—FIGURES OF MOLLUSCOUS ANIMALS,

Selected from various Authors. Etched for the Use of Students. By Maria Emma Gray. Vol. I. 8vo., with 78 plates of Figures, 12s. cloth.

GRAY AND MITCHELL'S ORNITHOLOGY.—THE GENERA OF BIRDS;

Comprising their Generic Characters, a Notice of the Habits of each Genus, and an extensive List of Species, referred to their several Genera. By George Robert Gray, Acad. Imp. Georg. Florent. Soc. Corresp. Senior Assistant of the Zoological Department, British Museum; and author of the "List of the Genera of Birds," etc. etc. Illustrated with 350 imperial 4to. Plates, by David William Mitchell, B.A.

In course of publication in Monthly Parts, 10s. 6d. each; each Part consisting generally of Four imperial quarto coloured Plates and Three plain, and accompanying Letter-press; giving the Generic Characters, short Remarks on the Habits, and a List of Species of each Genus as complete as possible. The uncoloured Plates will contain the Characters of all the Genera of the various Sub-families, consisting of numerous details of Heads, Wings, and Feet, as the case may require, for pointing out their distinguishing Characters.

** * The Work will not exceed 50 Monthly Parts. No. 18 was published on 1st of October.*

GRAY (J. E.)—THE GENERA OF MAMMALIA;

Comprising their Generic Characters—a Notice of the Habits of each Genus—and a short Character of each of the well-established Species, referred to the several Genera. By John Edward Gray, Esq., Keeper of the Zoological Collection of the British Museum. Imperial 4to. uniform with Gray and Mitchell's Ornithology; illustrated with 175 Plates.

** * To be published in Monthly Parts, 12s. each; each Part to consist of Four coloured and Three plain Plates, with accompanying Letter-press. The Work will not exceed 25 Parts. Publication will commence when 150 Subscribers' Names have been received.*

GREENER.—THE GUN;

Or, a Treatise on the various Descriptions of Small Fire Arms. By W. Greener, Inventor of an Improved Method of Firing Cannon by Percussion, etc. 8vo. with Illustrations, 15s. boards.

GREENWOOD (COL.)—THE TREE-LIFTER;

Or a New Method of Transplanting Trees. By Col. Geo. Greenwood. 8vo. with an Illustrative Plate, 7s. cloth.

GUEST.—THE MABINOCHION,

From the Llyfr Coch o Hergest, or Red Book of Hergest, and other ancient Welsh MSS. with an English Translation and Notes. By Lady Charlotte Guest. Royal 8vo. 8s. each.

Part 1.—The Lady of the Fountain.

Part 2.—Peredur Ab Eirawc; a Tale of Chivalry.

Part 3.—The Arthurian Romance of Geraint, the Son of Erbin.

Part 4.—The Romance of Kilhwch and Olwen.

Part 5.—The Dream of Rhonabwy, and the Tale of Pwyll Prince of Dyved.

Part 6.—Brauwen, the Daughter of Llyr: Manawyddan, the Son of Llyr; and Math, the Son of Mathonwy.

GUICCIARDINI (F.)—THE MAXIMS OF FRANCIS GUICCIARDINI, THE

HISTORIAN. Translated by Emma Martin, author of "A Short History of Ireland." With Notes, and Parallel Passages from the Works of Machiavelli, Lord Bacon, Pascal, Rochefoucault, Montesquieu, Burke, Prince Talleyrand, Guizot, and others. With a Sketch of the Author's Life. Square foolscap 8vo. [*In October.*]

GWILT.—AN ENCYCLOPÆDIA OF ARCHITECTURE;

Historical, Theoretical, and Practical. By Joseph Gwilt, Esq., F.S.A. Illustrated with upwards of 1000 Engravings on Wood, from Designs by J. S. Gwilt. In 1 thick vol. 8vo. containing nearly 1300 closely-printed pages. 2l. 12s. 6d. cloth.

"Gwilt's Encyclopædia ranks high as a work for professional students, containing the mathematics of architecture, with copious details upon all the technicalities of the science. It is a work which no professed architect or builder should be without."—Westminster Review.

HALL.—NEW GENERAL LARGE LIBRARY ATLAS OF FIFTY-THREE

MAPS, on Columbier Paper; with the Divisions and Boundaries carefully coloured. Constructed entirely from New Drawings, and engraved by Sidney Hall. New Edition, thoroughly revised and corrected; including all the Alterations rendered necessary by the recent Official Surveys, the New Roads on the Continent, and a careful Comparison with the authenticated Discoveries published in the latest Voyages and Travels. Folded in half, Nine Guineas, half-bound in Russia; full size of the Maps, Ten Pounds, half-bound in Russia.

The following Maps have been re-engraved, from entirely new designs—Ireland, South Africa, Turkey in Asia; the following have been materially improved—Switzerland, North Italy, South Italy, Egypt, Central Germany, Southern Germany, Greece, Austria, Spain, and Portugal; a new Map of China, corrected from the recent government survey of the coast from Canton to Nankin (to which is appended the Province of Canton, on an enlarged scale, in a separate compartment), has since been added.

HALSTED—LIFE AND TIMES OF RICHARD THE THIRD,

as Duke of Gloucester and King of England: in which all the Charges against him are carefully investigated and compared with the Statements of the Cotemporary Authorities. By Caroline A. Halsted, author of "The Life of Margaret Beaufort." 2 vols. 8vo. with Portrait from an Original Picture in the possession of the Right Honorable Lord Stafford, never before engraved, and other Illustrations, 1l. 10s. cloth.

HANNAM.—THE ECONOMY OF WASTE MANURES:

A Treatise on the Nature and Use of Neglected Fertilizers. By John Hannam. Written for the Yorkshire Agricultural Society, and published by permission of the Council. A new Edition. Fcap. 8vo.—*In the press.*

HAND-BOOK OF TASTE;

Or, How to Observe Works of Art, especially Cartoons, Pictures, and Statues. By Fabius Pictor. 3d Edition. Folscap 8vo. 3s. boards.

HANSARD.—TROUT AND SALMON FISHING IN WALES.

By G. A. Hansard, 12mo. 6s. 6d. cloth.

HARRIS.—THE HIGHLANDS OF ÆTHIOPIA;

Being the Account of Eighteen Months' Residence of a British Embassy to the Christian Court of Shoa. By Major Sir W. C. Harris, author of "Wild Sports in Southern Africa," etc. 2d Edition. 3 vols. 8vo. with Map and Illustrations, 2l. 2s. cloth.

HAWES (BARBARA).—TALES OF THE NORTH AMERICAN INDIANS,

and Adventures of the Early Settlers in America; from the Landing of the Pilgrim Fathers in 1620, to the Time of the Declaration of Independence. By Barbara Hawes. Folscap 8vo. with Frontispiece. 6s. cloth.

HAWKER.—INSTRUCTIONS TO YOUNG SPORTSMEN

In all that relates to Guns and Shooting. By Lieut. Col. P. Hawker. 9th edition, corrected, enlarged, and improved, with Eighty-five Plates and Woodcuts, by Adlard and Brauston, from Drawings by C. Varley, Dicks, etc. 8vo. 21s. cloth.

HAYDON (B. R.).—LECTURES ON PAINTING AND DESIGN,

Delivered at the London Institution, the Royal Institution, Albermarle Street, to the University of Oxford, etc. By B. R. Haydon, Historical Painter. With Designs drawn on Wood by the author, and Engraved by Edward Evans. 8vo. 12s. cloth.

HENSLOW.—THE PRINCIPLES OF DESCRIPTIVE AND PHYSIOLOGICAL BOTANY.

By J. S. Henslow, M.A. F.L.S. etc. Folscap 8vo. with Vignette Title, and nearly 70 Woodcuts, 6s. cloth.

HERSCHEL.—A TREATISE ON ASTRONOMY.

By Sir John Herschel. New Edition. Fcap. 8vo. Vignette Title, 6s. cloth.

HERSCHEL.—A PRELIMINARY DISCOURSE ON THE STUDY OF

NATURAL PHILOSOPHY. By Sir John Herschel. New Edition. Folscap 8vo. with Vignette Title, 6s. cloth.

HINTS ON ETIQUETTE AND THE USAGES OF SOCIETY:

With a Glance at Bad Habits. By *Αγαθός*. "Manners make the Man." 24th Edition, revised (with additions) by a Lady of Rank. Folscap 8vo. 2s. 6d. cloth, gilt edges.

General Observations; Introductions—Letters of Introduction—Marriage—Dinners—Smoking; Snuff—Fashion—Dress—Music—Dancing—Conversation—Advice to Tradespeople—Visiting; Visiting Cards—Cards—Tattling—of General Society.

HOARE.—A DESCRIPTIVE ACCOUNT OF A NEW METHOD OF

PLANTING AND MANAGING THE ROOTS OF GRAPE VINES. By Clement Hoare, author of "A Treatise on the Cultivation of the Grape Vine on Open Walls." 12mo 5s. cl.

HOARE.—A PRACTICAL TREATISE ON THE CULTIVATION OF THE

GRAPE VINE ON OPEN WALLS. By Clement Hoare. 3d Edition, 8vo. 7s. 6d. cloth.

HOBBS.—ENGLISH WORKS OF THOMAS HOBBS,

Of Malmesbury; now first collected by Sir William Molesworth, Bart. The English Works, Vols. 1 to 6, and 8 to 10; and the Latin Works, Vols. 1 to 3, 10s. each.

. Vols. 8 and 9, comprising the Translation of Thucydides, are not sold separately.

HOLLAND.—PROGRESSIVE EDUCATION;

Or, Considerations on the Course of Life. Translated from the French of Madame Necker de Saussure. By Miss Holland. 3 vols. folscap 8vo. 19s. 6d. cloth.

. The Third Volume, being supplementary to the first two, separately, 7s. 6d.

HOLLAND.—A TREATISE ON THE MANUFACTURES IN METAL.

By John Holland, Esq. 3 vols. folscap 8vo. with Vignette Titles, and about 300 Woodcuts, 18s. cloth.

HOLLAND.—MEDICAL NOTES AND REFLECTIONS.

By Henry Holland, M.D. F.R.S. etc. Fellow of the Royal College of Physicians, Physician Extraordinary to the Queen, and Physician in Ordinary to His Royal Highness Prince Albert. 2d Edition. 8vo. 18s. cloth.

HOOK (DR. W. F.).—THE LAST DAYS OF OUR LORD'S MINISTRY;

A Course of Lectures on the principal Events of Passion Week. By Walter Farquhar Hood, D.D., Vicar of Leeds, Prebendary of Lincoln, and Chaplain in Ordinary to the Queen. 4th Edition. Folscap 8vo. 6s. cloth.

HOOKE.—THE BRITISH FLORA.

In 2 vols. Vol. I.; comprising the Phænogamous or Flowering Plants, and the Ferns. By Sir William Jackson Hooker, K.H. LL.D. F.R.A. and L.S. etc. etc. 5th Edition, with Additions and Corrections; and 173 Figures, illustrative of the Umbelliferous Plants, the Composite Plants, the Grasses, and the Ferns. Vol. I. 8vo., with 12 Plates, 14s. plain; with the plates coloured, 24s. cloth.

Vol. II. in Two Parts, comprising the Cryptogamia and the Fungi, completing the British Flora, and forming Vol. V., Parts 1 and 2, of Smith's English Flora, 24s. boards.

HOOKE AND TAYLOR.—MUSCOLOGIA BRITANNICA.

Containing the Mosses of Great Britain and Ireland, systematically arranged and described; with Plates, illustrative of the character of the Genera and Species. By Sir W. J. Hooker and T. Taylor, M.D. F.L.S. etc. 2d Edition, 8vo. enlarged, 31s. 6d. plain; 37. 3s. coloured.

HORNE (THE REV. T. H.).—AN INTRODUCTION TO THE CRITICAL

STUDY AND KNOWLEDGE OF THE HOLY SCRIPTURES. By Thomas Hartwell Horne, B.D. of St. John's College, Cambridge; Rector of the united Parishes of St. Edmund the King and Martyr, and St. Nicholas Acons, Lombard Street; Prebendary of St. Paul's. 8th Edition, corrected and enlarged. Illustrated with numerous Maps and Fac-similes of Biblical Manuscripts. 4 vols. 8vo. (Vol. 2 in 2 Parts), 37. 3s. boards.

HORNE (THE REV. T. H.).—A COMPENDIOUS INTRODUCTION TO THE

STUDY OF THE BIBLE. By Thomas Hartwell Horne, B.D. of St. John's College, Cambridge. Being an Analysis of his "Introduction to the Critical Study and Knowledge of the Holy Scriptures." 7th Edition, corrected and enlarged. 12mo. with Maps and other Engravings, 9s. boards.

HORSLEY (BISHOP).—BIBLICAL CRITICISM ON THE FIRST FOUR-

TEEN HISTORICAL BOOKS OF THE OLD TESTAMENT; AND ON THE FIRST NINE PROPHETICAL BOOKS. By Samuel Horsley, LL.D. F.R.S. F.A.S. Lord Bishop of St. Asaph. Second Edition, containing Translations by the Author, never before published, together with copious Indexes. 2 vols. 8vo. 30s. cloth.

By the same Author,

THE BOOK OF PSALMS; translated from the Hebrew: with Notes, explanatory and critical. 4th Edition. 8vo. 12s. cloth.

HOWITT (MARY).—THE CHILD'S PICTURE AND VERSE BOOK,

Commonly called "Otto Speckter's Fable Book." Translated by Mary Howitt. With French and German on corresponding pages, and illustrated with 100 Engravings on Wood, by G. F. Sargent. 2d Edition. Square 12mo. 7s. 6d. boards.

HOWITT (MARY).—THE H—— FAMILY: TRALINNAN; AXEL AND

ANNA; and other Tales. By Fredrika Bremer. Translated by Mary Howitt. 2 vols. post 8vo. with Portrait of the Author, 21s. boards.

By the same Author and Translator.

THE NEIGHBOURS. A Story of Every-day Life in Sweden. 3d Edition, revised and corrected. 2 vols. post 8vo. 18s.

THE HOME; or, Family Cares and Family Joys. 2d Edition. 2 vols. post 8vo. 21s.

THE PRESIDENT'S DAUGHTERS, including NINA. 3 vols. post 8vo. 31s. 6d.

A NEW SKETCH OF EVERY-DAY LIFE:—A DIARY. Together with STRIFE and PEACE. 2 vols. post 8vo. 21s.

HOWITT.—THE RURAL LIFE OF ENGLAND.

By William Howitt. Third Edition, corrected and revised. Medium 8vo. with Engravings on Wood by Bewick and Williams, uniform with "Visits to Remarkable Places," 21s. cloth.

Life of the Aristocracy.

Life of the Agricultural Population.

Picturesque and Moral Features of the Country.

Strong Attachment of the English to Country Life.

The Forests of England.

Habits, Amusements, and Condition of the People; in which are introduced Two New Chapters, descriptive of the Rural Watering Places, and Education of Rural Population.

HOWITT.—VISITS TO REMARKABLE PLACES;

Old Halls, Battle-Fields, and Scenes illustrative of Striking Passages in English History and Poetry. By William Howitt. New Edition. Medium 8vo. with 40 Illustrations by S. Williams, 21s. cloth.

SECOND SERIES, chiefly in the Counties of DURHAM and NORTHUMBERLAND, with a Stroll along the BORDER. Medium 8vo. with upwards of 40 highly-finished Woodcuts, from Drawings made on the spot for this Work, by Messrs. Carmichael, Richardsons, and Weld Taylor, 21s. cloth.

HOWITT.—THE LIFE AND ADVENTURES OF JACK OF THE MILL,

Commonly called "Lord Othmill;" created, for his eminent services, Baron Waldeck, and Knight of Kitcottie; a Fireside Story. By William Howitt. 2d Edition. 2 vols. foolscap 8vo. with 46 Illustrations on Wood by G. F. Sargent, 12s. cloth.

HOWITT.—THE RURAL AND SOCIAL LIFE OF GERMANY:

With Characteristic Sketches of its chief Cities and Scenery. Collected in a General Tour, and during a Residence in that Country in the Years 1840-42. By William Howitt, author of "The Rural Life of England," etc. Medium 8vo., with above 50 Illustrations, 21s. cloth.

HOWITT.—GERMAN EXPERIENCES:

Addressed to the English, both Goers Abroad and Stayers at Home. By William Howitt. Foolscap 8vo. 7s. 6d. cloth.

HOWITT.—WANDERINGS OF A JOURNEYMAN TAILOR,
through EUROPE and the EAST, during the years 1824 to 1840. By P. D. Holthaus, from
Werdohl in Westphalia. Translated from the Third German Edition, by William Howitt,
author of "The Rural and Social Life of Germany," etc. Foolscep 8vo. with Portrait of
the Tailor, 6s. cloth.

HOWITT.—THE STUDENT-LIFE OF GERMANY.
From the Unpublished MS. of Dr. Cornelius. By William Howitt. 8vo. with 24 Wood-
Engravings, and 7 Steel Plates, 21s. cloth.

HOWITT.—COLONIZATION AND CHRISTIANITY:
A Popular History of the Treatment of the Natives, in all their Colonies, by the Europeans.
By William Howitt. Post 8vo. 10s. 6d. cloth.

HOWITT.—THE BOY'S COUNTRY BOOK:
Being the real Life of a Country Boy, written by Himself; exhibiting all the Amusements,
Pleasures, and Pursuits of Children in the Country. Edited by William Howitt, author of
"The Rural Life of England," etc. 2d Edition. Fcap. 8vo. with 40 Woodcuts, 8s. cloth.

HOWITT (RICHARD).—IMPRESSIONS OF AUSTRALIA FELIX,
During a Four Years' Residence in that Colony: with particular reference to the Prospects
of Emigrants. With Notes of a Voyage round the World, Australian Poems, etc. By
Richard Howitt. Foolscep 8vo. 7s. cloth.

HUDSON.—THE PARENT'S HAND-BOOK;
Or, Guide to the Choice of Professions, Employments, and Situations; containing useful
and practical information on the subject of placing out Young Men, and of obtaining their
Education with a view to particular occupations. By J. C. Hudson, Esq., author of "Plain
Directions for Making Wills." Fcap. 8vo. 5s. cloth.

HUDSON.—PLAIN DIRECTIONS FOR MAKING WILLS
In Conformity with the Law, and particularly with reference to the Act 7 Wm. IV. and 1 Vict.
c. 26. To which is added, a clear Exposition of the Law relating to the Distribution of Per-
sonal Estate in the case of Intestacy; with two Forms of Wills, and much useful Information,
etc. By J. C. Hudson, Esq. 13th Edition, corrected, with Notes of Cases judicially decided
since the above Act came into operation. Fcap. 8vo. 2s. 6d.

HUDSON.—THE EXECUTOR'S GUIDE.
By J. C. Hudson, Esq., of the Legacy Duty Office, London; author of "Plain Directions
for Making Wills," and "The Parent's Hand-Book." 4th Edition. Foolscep 8vo. 5s. cloth.
. *The above two works may be had in One volume, price 7s. cloth.*

HUMPHREYS.—THE ILLUMINATED BOOKS OF THE MIDDLE AGES.
A History of Illuminated Books, from the 14th to the XVIIth Century. By Henry Noel
Humphreys. Illustrated by a Series of Specimens, consisting of an entire Page, of the
exact size of the Original, from the most celebrated and splendid MSS. in the Imperial and
Royal Libraries of Vienna, Moscow, Paris, Naples, Copenhagen, and Madrid;—from the
Vatican, Escorial, Ambrosian, and other great Libraries of the Continent;—and from the
rich Public, Collegiate, and Private Libraries of Great Britain; superbly printed in Gold,
Silver, and Colours.
In course of publication, in Parts. Parts 1 and 2, each containing Three Plates, with Descrip-
tions, Imperial Quarto, splendidly printed, in gold, silver, and colours, in imitation of the
originals, as accurate as can be produced by mechanical means, price 12s.
Large Paper, on Half Imperial (21½ in. by 15), to prevent folding the large Plates, 21s.
Six Parts to form a Volume, Four Volumes completing the work.

HUNT.—RESEARCHES ON LIGHT:
An Examination of all the Phenomena connected with the Chemical and Molecular Changes
produced by the Influence of the Solar Rays; embracing all the known Photographic Pro-
cesses, and new Discoveries in the Art. By Robert Hunt, Secretary of the Royal Cornwall
Polytechnic Society. 8vo. with Plate and Woodcuts, 10s. 6d. cloth.

ILLUMINATED CALENDAR (THE).—THE ILLUMINATED CALENDAR and HOME
DIARY for 1846: containing 12 pages of *fac-simile* from the Calendar of the rich Missal of
the Duke of Anjou, styled King of Sicily and Jerusalem; also 24 pages of Diary, each illu-
minated with an elaborate Border taken from the same MS.; and an Illuminated Title. The
binding designed from the framework of one of the miniature pictures of the same MS.
Imperial 8vo. 42s. bound in an appropriate ornamental cover.—*In the press.*

. *The elaborate Gothic traceries of this MS. form one of the finest monuments of the
art of illuminating. It was executed towards the close of the fourteenth century, more than
a century earlier than the "Hours of Anne of Brittany," which formed the subject of the
Calendar for 1844; and in style and execution it exhibits a totally different style of art from
that work.*

. *The Illuminated Calendar and Home Diary, for 1845; copied from the Manuscript of
the "Hours of Anne of Brittany." Imp. 8vo. in emblazoned printing and binding, 42s.*

JACKSON.—PICTORIAL FLORA;
Or, British Botany Delineated, in 1500 Lithographic Drawings of all the Species of Flowering
Plants indigenous to Great Britain; illustrating the descriptive works on English Botany of
Hooker, Lindley, Smith, etc. By Miss Jackson. 8vo. 15s. cloth.

JAMES.—A HISTORY OF THE LIFE OF EDWARD THE BLACK PRINCE, and of various Events connected therewith, which occurred during the Reign of Edward III. King of England. By G. P. R. James, Esq. 2d Edition. 2 vols. foolscap 8vo. with Map, 15s. cl.

JAMES.—LIVES OF THE MOST EMINENT FOREIGN STATESMEN.

By G. P. R. James, Esq., and E. E. Crowe, Esq. 5 vols. foolscap 8vo. with Vignette Titles. 30s. cloth.

JEBB (BISHOP).—PRACTICAL THEOLOGY;

Comprising Discourses on the Liturgy and Principles of the United Church of England and Ireland; Critical and other Tracts; and a Speech delivered in the House of Peers in 1824. By John Jebb, D.D. F.R.S., Bishop of Limerick, Ardfert, and Aghadoe. 2d Edition. 2 vols. 8vo. 24s. cloth.

By the same Author,

PASTORAL INSTRUCTIONS, on the Character and Principles of the Church of England. Selected from his former Publications. A New Edition. Foolscap 8vo. 6s. cloth.

JEBB (BISHOP) AND KNOX (ALEXANDER).—THIRTY YEARS' COR-

RESPONDENCE between John Jebb, D.D. F.R.S., Bishop of Limerick, Ardfert, Aghadoe, and Alexander Knox, Esq. M.R.I.A. Edited by the Rev. Charles Forster, B.D. Rector of Stisted, Essex, and one of the Six Preachers in the Cathedral of Christ, Canterbury, formerly Domestic Chaplain to Bishop Jebb. 2d Edition. 2 vols. 8vo. 28s. cloth.

LORD JEFFREY.—CONTRIBUTIONS TO THE EDINBURGH REVIEW.

By Francis Jeffrey, now one of the Judges in the Court of Session in Scotland. 4 vols. 8vo. 48s. cloth.

JOHNSON.—THE FARMER'S ENCYCLOPÆDIA,

And **DICTIONARY OF RURAL AFFAIRS:** embracing all the recent Discoveries in Agricultural Chemistry; adapted to the comprehension of unscientific Readers. By Cuthbert W. Johnson, Esq., F.R.S. Barrister-at-Law, Corresponding Member of the Agricultural Society of Königsberg, and of the Maryland Horticultural Society, author of several of the Prize Essays of the Royal Agricultural Society of England, and other Agricultural Works; Editor of the "Farmer's Almanack," etc. 1 thick vol. 8vo. illustrated by Wood Engravings of the best and most improved Agricultural Implements. 2l. 10s. cloth.

KANE.—THE INDUSTRIAL RESOURCES OF IRELAND.

By Robert Kane, M.D. Secretary to the Council of the Royal Irish Academy, Professor of Natural Philosophy to the Royal Dublin Society, and of Chemistry to the Apothecaries' Hall of Ireland. 2d Edition. Post 8vo. 7s. cloth.

KANE.—ELEMENTS OF CHEMISTRY;

Including the most Recent Discoveries and Applications of the Science to Medicine and Pharmacy, and to the Arts. By Robert Kane, M.D. M.R.I.A. Professor of Natural Philosophy to the Royal Dublin Society. 8vo. with 236 Woodcuts, 24s. cloth.

KATER AND LARDNER.—A TREATISE ON MECHANICS.

By Captain Kater and Dr. Lardner. New Edition. Foolscap 8vo. Vignette Title, and 19 Plates, comprising 224 distinct figures, 6s. cloth.

KEIGHTLEY.—OUTLINES OF HISTORY,

From the Earliest Period. By Thomas Keightley, Esq. New Edition, corrected and considerably improved. Foolscap 8vo., 6s. cloth; or 6s. 6d. bound.

KEON (M. G.).—A HISTORY OF THE JESUITS,

Literary, Social, and Political, from the Birth of Ignatius Loyola to the present time. By Miles Gerald Keon. 8vo.—*Preparing for publication.*

KIRBY AND SPENCE.—AN INTRODUCTION TO ENTOMOLOGY;

Or, Elements of the Natural History of Insects: comprising an account of noxious and useful Insects, of their Metamorphoses, Food, Stratagems, Habitations, Societies, Motions, Noises, Hybernation, Instinct, etc. By W. Kirby, M.A. F.R.S. & L.S. Rector of Barham; and W. Spence, Esq., F.R.S. & L.S. 6th Edition, corrected and considerably enlarged. 2 vols. 8vo. 1l. 11s. 6d. cloth.

The first two volumes of the "Introduction to Entomology" are now published as a separate work, distinct from the third and fourth volumes, and, though much enlarged, at a considerable reduction of price, in order that the numerous class of readers who confine their study of insects to that of their manners and economy, need not be burthened with the cost of the technical portion of the work relating to their anatomy, physiology, etc.

KNAPP.—GRAMINA BRITANNICA;

Or, Representations of the British Grasses: with Remarks and occasional Descriptions. By I. L. Knapp, Esq. F.L.S. & A.S. 2d Edition. 4to. with 118 Plates, beautifully coloured, 3l. 16s. boards.

KNOX (ALEXANDER).—REMAINS OF ALEXANDER KNOX, ESQ.

Of Dublin, M.R.I.A.; containing Essays, chiefly explanatory, of Christian Doctrine; and Confidential Letters, with Private Papers, illustrative of the Writer's Character, Sentiments, and Life. 3d Edition. 4 vols. 8vo. 2l. 8s. cloth.

LAING.—NOTES ON THE SCHISM FROM THE CHURCH OF ROME,

called the GERMAN CATHOLIC CHURCH, instituted by J. Ronge and I. Czerzki, in October 1844, on occasion of the Pilgrimage to the Holy Coat at Treves. By S. Laing, Esq., author of "Notes of a Traveller," "The Chronicles of the Kings of Norway," etc. Fcap. 8vo. 5s. cloth.

LAING.—THE CHRONICLE OF THE KINGS OF NORWAY,

From the Earliest Period of the History of the Northern Sea Kings to the Middle of the Twelfth Century, commonly called the *Heimskringla*. Translated from the Icelandic of Snorro Sturleson, with Notes, and a Preliminary Discourse, by Samuel Laing, author of "Notes of a Traveller," etc. 3 vols. 8vo. 36s. cloth.

LAING.—A TOUR IN SWEDEN

In 1838; comprising Observations on the Moral, Political, and Economical State of the Swedish Nation. By Samuel Laing, Esq. 8vo. 12s. cloth.

LAING.—NOTES OF A TRAVELLER

On the Social and Political State of France, Prussia, Switzerland, Italy, and other parts of Europe, during the present Century. By Samuel Laing, Esq. 2d Edition. 8vo. 16s. cloth.

LAING.—JOURNAL OF A RESIDENCE IN NORWAY

During the years 1834, 1835, and 1836; made with a view to inquire into the Rural and Political Economy of that Country, and the Condition of its Inhabitants. By Samuel Laing, Esq. 2d Edition. 8vo. 14s. cloth.

LARDNER'S CABINET CYCLOPÆDIA;

Comprising a Series of Original Works on History, Biography, Literature, the Sciences, Arts, and Manufactures. Conducted and edited by Dr. Lardner.

The Series complete in One Hundred and Thirty-three Volumes, 39l. 18s. (One Volume remains to be published.) The Works separate, 6s. per volume.

LARDNER.—A TREATISE ON ARITHMETIC.

By Dr. Lardner, LL.D. F.R.S. Folscap 8vo. with Vignette Title, 6s. cloth.

LARDNER AND WALKER.—A MANUAL ON ELECTRICITY, MAG-

NETISM, and METEOROLOGY. By Dr. Lardner, LL.D. F.R.S., and C. V. Walker, Secretary of the Electrical Society. 2 vols. folscap 8vo., with Vignette Titles, 12s. cloth.

LARDNER.—A TREATISE ON GEOMETRY,

And its Application to the Arts. By Dr. Lardner. Folscap 8vo. with Vignette Title, and upwards of 200 figures, 6s. cloth.

LARDNER.—A TREATISE ON HEAT.

By Dr. Lardner, LL.D. etc. Fcap. 8vo. with Vignette Title and Woodcuts, 6s. cloth.

LARDNER.—A TREATISE ON HYDROSTATICS AND PNEUMATICS.

By Dr. Lardner. New Edition. Folscap 8vo. 6s. cloth.

LECTURES ON POLARISED LIGHT,

Delivered by Dr. Pereira, before the Pharmaceutical Society, and in the Medical School of the London Hospital. 8vo. illustrated by above 50 Woodcuts, 5s. 6d. cloth.

L. E. L.—THE POETICAL WORKS OF LETITIA ELIZABETH LANDON.

New Edition, 4 vols. folscap 8vo. with Illustrations by Howard, etc. 28s. cloth; or bound in morocco, with gilt edges, 2l. 4s.

The following Works separately:—

The IMPROVISATRICE	- - 10s. 6d.	The GOLDEN VIOLET	- - - 10s. 6d.
The VENETIAN BRACELET	10s. 6d.	The TROUBADOUR	- - - 10s. 6d.

LEE.—TAXIDERMY;

Or, the Art of Collecting, Preparing, and Mounting Objects of Natural History. For the use of Museums and Travellers. By Mrs. R. Lee (formerly Mrs. T. E. Bowdich), author of "Memoirs of Cuvier," etc. 6th Edition, improved, with an account of a Visit to Walton Hall, and Mr. Waterton's method of Preserving Animals. Fcap. 8vo. with Wood Engravings, 7s. cloth.

LEE.—ELEMENTS OF NATURAL HISTORY,

For the Use of Schools and Young Persons; comprising the Principles of Classification, interspersed with amusing and instructive original Accounts of the most remarkable Animals. By Mrs. R. Lee (formerly Mrs. T. E. Bowdich), author of "Taxidermy," "Memoirs of Cuvier," etc. 12mo. with 55 Woodcuts, 7s. 6d. bound.

LEFEVRE (SIR GEORGE).—AN APOLOGY FOR THE NERVES;

Or, their Importance and Influence in Health and Disease. By Sir George Lefevre, M.D. Fellow of the Royal College of Physicians, late Physician to the British Embassy at the Court of St. Petersburg, etc.; author of "The Life of a Travelling Physician," "Thermal Comfort," etc. Post 8vo. 9s. cloth.

LEMPRIERE.—A CLASSICAL DICTIONARY;

Containing a copious Account of all the Proper Names mentioned in Ancient Authors; with the Value of Coins, Weights, and Measures, used amongst the Greeks and Romans; and a Chronological Table. By T. Lemprière, D.D. 20th Edition, corrected. 8vo. 9s. cloth.

LESLIE (C.R.).—MEMOIRS OF THE LIFE OF JOHN CONSTABLE, ESQ.

R.A. Composed chiefly of his Letters. By C.R. Leslie, R.A. Second Edition, with further Extracts from his Correspondence. Small 4to. with two Portraits (one from a new Sketch, by Mr. Leslie, and a plate of "Spring," engraved by Lucas). 2ls. cloth.

"Got up in a peculiar, antique, and handsome manner, congenial to, and worthy of, the subject. The world at large will find much to entertain and instruct in this interesting biography; artists and amateurs in particular will derive great instruction, and every class of readers useful intelligence and agreeable amusement. Mr. Leslie has performed his task with infinite taste and discrimination; giving the opinions of a competent authority upon the productions of his contemporary and friend, and bringing out the remarkable qualities and estimable points of poor Constable's amiable private character in every relation of life with unaffected simplicity and consequent effect."—Literary Gazette.

LIFE OF A TRAVELLING PHYSICIAN,

From his first Introduction to Practice; including Twenty Years' Wanderings throughout the greater part of Europe. 3 vols. post 8vo. with coloured Frontispieces, 31s. 6d. cloth.

LINDLEY.—INTRODUCTION TO BOTANY.

By Prof. J. Lindley, Ph.D. F.R.S. L.S. etc. 3d Edition (1839), with Corrections and considerable Additions, 8vo. with Six Plates and numerous Woodcuts, 18s. cloth.

LINDLEY.—FLORA MEDICA;

A Botanical Account of all the most important Plants used in Medicine, in different Parts of the World. By John Lindley, Ph.D. F.R.S. etc. 8vo. 18s. cloth.

LINDLEY.—A SYNOPSIS OF THE BRITISH FLORA,

Arranged according to the Natural Orders. By Professor John Lindley, Ph.D., F.R.S., etc. Third Edition, with numerous Additions, Corrections, and Improvements. 12mo. 10s. 6d. cloth.

LINDLEY.—THE THEORY OF HORTICULTURE;

Or, an Attempt to Explain the Principal Operations of Gardening upon Physiological Principles. By John Lindley, Ph.D. F.R.S. 8vo. with Illustrations on Wood, 12s. cloth.

This book is written in the hope of providing the intelligent gardener, and the scientific amateur, correctly, with the rationalia of the more important operations of Horticulture; and the author has endeavoured to present to his readers an intelligible explanation, founded upon well ascertained facts, which they can judge of by their own means of observation, of the general nature of vegetable actions, and of the causes which, while they control the powers of life in plants, are capable of being regulated by themselves. The possession of such knowledge will necessarily teach them how to improve their methods of cultivation, and lead them to the discovery of new and better modes.

LINDLEY.—GUIDE TO THE ORCHARD AND KITCHEN GARDEN;

Or, an Account of the most valuable Fruits and Vegetables cultivated in Great Britain: with Kalendars of the Work required in the Orchard and Kitchen Garden during every month in the Year. By George Lindley, C.M.H.S. Edited by Professor Lindley. 8vo. 16s. boards.

LLOYD.—A TREATISE ON LIGHT AND VISION.

By the Rev. H. Lloyd, M.A., Fellow of Trin. Coll. Dublin. 8vo. 5s. boards.

LORIMER.—LETTERS TO A YOUNG MASTER MARINER,

On some Subjects connected with his Calling. By Charles Lorimer. 3d edition. 12mo. with an Appendix, 5s. 6d. cloth.

LOUDON (MRS.)—THE LADY'S COUNTRY COMPANION;

Or, How to Enjoy a Country Life Rationally. By Mrs. Loudon, author of "Gardening for Ladies," etc. Foolscep 8vo., with an Engraving on Steel, and Illustrations on Wood, 7s. 6d. cl.
"A more intelligent and pleasant Country Companion than Mrs. Loudon's lives not even in these days of perpetual instruction and guides to every thing. For a stroll in the fields, for a walk in the garden, for managing the dairy or poultry-yard, for raising the finest flowers and the best fruits, for rural amusements, and for all useful employments to occupy time and produce profitable results, we cordially recommend this excellent work to be taken as a constant companion."—Literary Gazette.

LOUDON (J. C.)—SELF INSTRUCTION

For Young Gardeners, Foresters, Bailiffs, Land Stewards, and Farmers; in Arithmetic, Book-keeping, Geometry, Mensuration, Practical Trigonometry, Mechanics, Land-Surveying, Leveling, Planning and Mapping, Architectural Drawing, and Isometrical Projection and Perspective; with Examples shewing their applications to Horticultural and Agricultural Purposes. By the late J. C. Loudon, F.L.S. H.S. etc. With a Portrait of Mr. Loudon, and a Memoir by Mrs. Loudon. 8vo. *In October.*

LOUDON.—AN ENCYCLOPÆDIA OF TREES AND SHRUBS;

Being the "Arboretum et Fruticetum Britannicum" abridged: containing the Hardy Trees and Shrubs of Great Britain, Native and Foreign, scientifically and popularly described: with their Propagation, Culture, and Uses in the Arts; and with Engravings of nearly all the Species. For the use of Nurserymen, Gardeners, and Foresters. By J. C. Loudon, F.L.S. etc. 1 large vol. 8vo. with upwards of 2000 Engravings on Wood, 21. 10s. cloth.

The Original Work may be had in 8 vols. 8vo. with above 400 8vo. Plates of Trees, and upwards of 2500 Woodcuts, 101. cloth.

LOUDON.—AN ENCYCLOPÆDIA OF GARDENING;

Presenting in one systematic view, the History and Present State of Gardening in all Countries, and its Theory and Practice in Great Britain: with the Management of the Kitchen Garden, the Flower Garden, Laying-out Grounds, etc. By J. C. Loudon, F.L.S. etc. A new Edition, enlarged and much improved. 1 large vol. 8vo. with nearly 1000 Engravings on Wood, 21. 10s. cloth.

LOUDON.—AN ENCYCLOPÆDIA OF AGRICULTURE;

Comprising the Theory and Practice of the Valuation, Transfer, Laying-out, Improvement, and Management of Landed Property, and of the cultivation and economy of the Animal and Vegetable productions of Agriculture, including all the latest improvements; a general History of Agriculture in all countries; a Statistical view of its present state, with suggestions for its future progress in the British Isles; and Supplements, bringing down the work to the year 1844. By J. C. Loudon, F.L.G.Z. and H.S. etc. Fifth Edition. 8vo. illustrated with upwards of 1100 Engravings on Wood, by Branston. 21. 10s. cloth.

The Supplement, bringing down improvements in the art of Field-Culture from 1831 to 1844 inclusive, comprising all the previous Supplements, and illustrated with 65 Engravings on Wood, may be had *separately*, 5s. sewed.

LOUDON.—AN ENCYCLOPÆDIA OF PLANTS;

Including all the Plants which are now found in, or have been introduced into, Great Britain; giving their Natural History, accompanied by such Descriptions, Engraved Figures, and Elementary Details, as may enable a beginner, who is a mere English reader, to discover the name of every Plant which he may find in flower, and acquire all the information respecting it which is useful and interesting. The Specific Characters by an Eminent Botanist; the Drawings by J. D. C. Sowerby, F.L.S. A new Edition (1841), with a new Supplement, comprising every desirable particular respecting all the Plants originated in, or introduced into, Britain between the first publication of the work, in 1829, and January 1840: with a new General Index to the whole work. Edited by J. C. Loudon, prepared by W. H. Baxter, Jun., and revised by George Don, F.L.S.; and 800 new Figures of Plants, on Wood, from Drawings by J. D. C. Sowerby, F.L.S. 1 very large vol. 8vo. with nearly 10,000 Wood Engravings, 3*l.* 13*s.* 6*d.* cloth.

*. * The last Supplement, *separately*, 8vo. 15*s.* cloth.

LOUDON.—AN ENCYCLOPÆDIA OF COTTAGE, FARM, AND VILLA

ARCHITECTURE and FURNITURE. Containing Designs for Cottages, Villas, Farm Houses, Farmhouses, Country Inns, Public Houses, Parochial Schools, etc.; with the requisite Fittings-up, Fixtures, and Furniture, and appropriate Offices, Gardens, and Garden Scenery: each Design accompanied by Analytical and Critical Remarks illustrative of the Principles of Architectural Science and Taste on which it is composed, and General Estimates of the Expense. By J. C. Loudon, F.L.S. etc. New Edition, corrected, with a Supplement containing 160 additional pages of letter-press, and nearly 300 new engravings. 1 very thick vol. 8vo. with more than 2000 Engravings on Wood, 63*s.* cloth.

*. * The Supplement, *separately*, 8vo. 7*s.* 6*d.* sewed.

LOUDON.—HORTUS BRITANNICUS:

A Catalogue of all the Plants indigenous to or introduced into Britain. The 3d Edition (1832), with a New Supplement, prepared, under the direction of J. C. Loudon, by W. H. Baxter, and revised by George Don, F.L.S. 8vo. 3*l.* 6*d.* cloth.

The Supplement *separately*, 8vo. 2*s.* 6*d.* sewed.

The later Supplement *separately*, 8vo. 8*s.* sewed.

LOUDON.—THE SUBURBAN GARDENER AND VILLA COMPANION:

Comprising the Choice of a Villa or Suburban Residence, or of a situation on which to form one; the Arrangement and Furnishing of the House; and the Laying-out, Planting, and general Management of the Garden and Grounds; the whole adapted for Grounds from one perch to fifty acres and upwards in extent; intended for the instruction of those who know little of Gardening or Rural Affairs, and more particularly for the use of Ladies. By J. C. Loudon, F.L.S., etc. 8vo. with above 300 Wood Engravings, 20*s.* cloth.

LOUDON.—HORTUS LIGNOSUS LONDINENSIS;

Or, a Catalogue of all the Ligneous Plants cultivated in the neighbourhood of London. To which are added their usual Prices in Nurseries. By J. C. Loudon, F.L.S. etc. 8vo. 7*s.* 6*d.*

LOW.—ON LANDED PROPERTY, AND THE MANACEMENT OF ESTATES;

Comprehending the Relations between Landlord and Tenant, and the Principles and Forms of Leases; of Farm-buildings, Enclosures, Drains, Embankments, Roads, and other Rural Works, Minerals, and Woods. By David Low, Esq. F.R.S.E. etc., author of "Elements of Practical Agriculture," etc. 8vo. with numerous Engravings, 2*l.* cloth.

LOW.—ON THE DOMESTICATED ANIMALS OF GREAT BRITAIN,

comprehending the Natural and Economical History of the Species and Breeds; Illustrations of the Properties of External Form; and Observations on the Principles and Practice of Breeding. By David Low, Esq., F.R.S.E. Professor of Agriculture in the University of Edinburgh; Member of the Royal Academy of Agriculture of Sweden; Corresponding Member of the Conseil Royal d'Agriculture de France, of the Société Royal et Centrale, etc.; author of "Elements of Practical Agriculture," "Illustrations of the Breeds of the Domesticated Animals of the British Islands," "On Landed Property and the Economy of Estates," etc. 8vo. with Engravings on Wood. [In October.]

LOW.—THE BREEDS OF THE DOMESTICATED ANIMALS OF GREAT

BRITAIN described. By David Low, Esq. F.R.S.E., Professor of Agriculture in the University of Edinburgh; Member of the Royal Academy of Agriculture of Sweden; Corresponding Member of the Conseil Royal d'Agriculture de France, of the Société Royal et Centrale, etc. etc. The Plates from drawings by W. Nicholson, R.S.A., reduced from a Series of Oil Paintings, executed for the Agricultural Museum of the University of Edinburgh by W. Shiels, R.S.A. In 2 vols. atlas quarto, with 56 plates of Animals, beautifully coloured after Nature, 16*l.* 16*s.* half-bound in morocco.

Or in four separate portions, as follow:—

The OX. 1 vol. atlas quarto, with 22 Plates, price 6*l.* 16*s.* 6*d.* half-bound morocco.

The HORSE. 1 vol. atlas quarto, with 8 Plates, price 3*l.* half-bound morocco.

The SHEEP. 1 vol. atlas quarto, with 21 Plates, price 6*l.* 16*s.* 6*d.* half-bound morocco.

The HOG. 1 vol. atlas quarto, with 5 Plates, price 2*l.* 2*s.* half-bound morocco.

LOW.—ELEMENTS OF PRACTICAL AGRICULTURE;

Comprehending the Cultivation of Plants, the Husbandry of the Domestic Animals, and the Economy of the Farm. By David Low, Esq. F.R.S.E., Professor of Agriculture in the University of Edinburgh. 4th Edition, with Alterations and Additions, and above 200 Woodcuts. 8vo. 2*l.* cloth.

MACAULAY.—CRITICAL AND HISTORICAL ESSAYS CONTRIBUTED TO THE EDINBURGH REVIEW. By the Right Hon. Thomas Babington Macaulay, M.P. 3d Edition. 3 vols. 8vo. 36s. cloth.

MACAULAY.—LAYS OF ANCIENT ROME.

By the Right Honorable Thomas Babington Macaulay, M.P. 6th Edition. Crown 8vo. 10s. 6d. cloth.

MACKENZIE.—THE PHYSIOLOGY OF VISION.

By W. Mackenzie, M.D., Lecturer on the Eye in the University of Glasgow. 8vo. with Woodcuts, 10s. 6d. boards.

MACKINNON.—THE HISTORY OF CIVILISATION.

By Wm. Alexander Mackinnon, F.R.S. M.P. for Lymington. 2 vols. 8vo. [In October.

MACKINTOSH (SIR JAMES).—THE LIFE OF SIR THOMAS MORE.

By the Right Hon. Sir James Mackintosh. Reprinted from the Cabinet Cyclopædia; and intended for a Present-Book or School Prize. Folscap 8vo. with Portrait, 5s. cloth; or bound in vellum gilt (*old style*), 8s.

MACKINTOSH'S (SIR JAMES) MISCELLANEOUS WORKS;

Including his Contributions to The EDINBURGH REVIEW. Collected and Edited by his Son. 3 vols. 8vo.—*In the press.*

MACKINTOSH, ETC.—THE HISTORY OF ENGLAND.

By Sir James Mackintosh; W. Wallace, Esq.; and Robert Bell, Esq. 10 vols. folscap 8vo. with Vignette Titles, 3l. cloth.

M'CULLOCH.—THE LITERATURE OF POLITICAL ECONOMY;

Being a Classified Catalogue of the principal Works in the different departments of Political Economy, interspersed with Historical, Critical, and Biographical Notices. By J. R. M'Culloch, Esq. 8vo. 14s. cloth.

M'CULLOCH.—A TREATISE ON THE PRINCIPLES AND PRACTICAL INFLUENCE OF TAXATION AND THE FUNDING SYSTEM. By J. R. M'Culloch, Esq. 8vo. 15s. cloth.

M'CULLOCH.—A DICTIONARY, GEOGRAPHICAL, STATISTICAL, AND HISTORICAL, of the various Countries, Places, and Principal Natural Objects in the World. By J. R. M'Culloch, Esq. A new Edition, 2 thick vols. 8vo. with Six large Maps, 4l. cloth.

* * *The new Articles on the British Empire, England, Ireland, and Scotland, will be printed separately as a supplement to the former Edition. They comprise a pretty full account of the present state of the British Empire.*

M'CULLOCH.—A DICTIONARY, PRACTICAL, THEORETICAL, AND HISTORICAL, OF COMMERCE, AND COMMERCIAL NAVIGATION. By J. R. M'Culloch, Esq. An entirely New Edition, corrected throughout, enlarged, and improved. 1 very thick vol. 8vo., illustrated with Maps and Plans, 50s. cloth; or 55s. strongly half-bound in Russia, with flexible back.

"Mr. M'Culloch's Commercial Dictionary has for several years been a vade mecum for merchants, traders, ship-owners, and ship-masters, to guide and assist them in conducting the details of their respective occupations, we need not therefore expatiate upon the general merits of this well-known work, in announcing to the mercantile world a new, enlarged, and improved edition. The subjects handled in a commercial dictionary are not of a stationary but a progressive character, and those who mostly use such repertoires are not curious about historical notices or theoretical discussions, but concern themselves solely with practical details immediately connected with the present moment. The changes made in our commercial policy by the Tariff Act of 1842, and the late acts for regulating the corn and colonial trades, are so multiform, so important, and affect so many articles and interests, that Mr. M'Culloch despaired of introducing them into a supplement of a less size than the original work, he has therefore reconstructed his dictionary altogether. We have carefully examined this vast work, and are of opinion that the indefatigable author has produced a digest of the most useful and authentic information respecting the past and present state of the commerce of Europe and the world at large, and the laws and regulations under which commercial operations are carried on. We have not space, nor would it be useful if we had, to enumerate the new subjects treated in the edition before us; it will suffice practical men to be assured that in the course of their business scarcely any commercial question can arise upon which they will not find useful information in Mr. M'Culloch's well-stored pages."

From an article on Mr. M'Culloch's Dictionary in the "Times" newspaper.

MALTE-BRUN.—A SYSTEM OF UNIVERSAL GEOGRAPHY,

Founded on the Works of Malte-Brun and Balbi, embracing an Historical Sketch of the Progress of Geographical Discovery, the Principles of Mathematical and Physical Geography, and a complete Description, from the most recent sources, of the Political and Social Condition of all the Countries in the World: with numerous Statistical Tables, and an Alphabetical Index of 12,000 Names. 8vo. 30s. cloth.

MARCET (MRS.)—CONVERSATIONS ON THE HISTORY OF ENGLAND.

For the Use of Children. By Mrs. Marcet, author of "Conversations on Chemistry," etc. 2d Edition. 18mo. 5s. cloth.

MARCET.—CONVERSATIONS ON CHEMISTRY;

In which the Elements of that Science are familiarly Explained and Illustrated by Experiments. 14th Edition, enlarged and corrected. 2 vols. foolscap 8vo. 14s. cloth.

MARCET.—CONVERSATIONS ON NATURAL PHILOSOPHY;

In which the Elements of that Science are familiarly explained, and adapted to the comprehension of Young Persons. 10th Edition, enlarged and corrected by the Author. Fcap. 8vo. with 23 Plates, 10s. 6d. cloth.

MARCET.—CONVERSATIONS ON POLITICAL ECONOMY;

In which the Elements of that Science are familiarly explained. 7th Edition, revised and enlarged. Foolscap 8vo. 7s. 6d. cloth.

MARCET.—CONVERSATIONS ON VEGETABLE PHYSIOLOGY;

Comprehending the Elements of Botany, with their application to Agriculture. 3d Edition. Foolscap 8vo. with Four Plates, 9s. cloth.

MARCET.—CONVERSATIONS FOR CHILDREN;

On Land and Water. 2d Edition revised and corrected. Foolscap 8vo., with coloured Maps, shewing the comparative Altitude of Mountains, 5s. 6d. cloth.

MARCET.—CONVERSATIONS ON LANGUAGE,

For Children. By Mrs. Marcet, author of "Mary's Grammar," etc. 18mo. 4s. 6d. cloth.

MARCET.—THE GAME OF GRAMMAR,

With a Book of Conversations, shewing the Rules of the Game, and affording Examples of the manner of playing at it. In a varnished box, or done up as a post 8vo. volume, 8s.

MARCET.—WILLY'S GRAMMAR;

Interspersed with Stories, and intended for the Use of Boys. By Mrs. Marcet, author of "Mary's Grammar," etc. New edition. 18mo. 2s. 6d. cloth.

"A sound and simple work for the earliest ages."—Quarterly Review.

MARCET.—LESSONS ON ANIMALS, VEGETABLES, AND MINERALS.

By Mrs. Marcet, author of "Conversations on Chemistry," etc. 12mo. 2s. cloth.

"One of Mrs. Marcet's carefully written books of instruction, in which natural history is made pleasant and intelligible for the young."—Athenæum.

MARRIAGE GIFT.

By a Mother. A Legacy to her Children. Post 8vo. 5s. cloth, gilt edges.

MARRYAT (CAPT).—THE MISSION;

Or, Scenes in Africa. Written for Young People. By Captain Marryat, C.B., author of "Peter Simple," "Masterman Ready," "The Settlers in Canada," etc. 2 vols. fcap. 8vo. 12s. cloth.

"A delightful book for young people, written with great truth and point, and abounding in the natural but exciting adventure, that Cape emigrants are sure to encounter. The descriptions of natural objects—of the scenes of earth, air, and sea—of plants and animals—of the Caffres and their mode of life—are all filled in with the happiest effect. It is one of the author's most successful efforts in his more recent and best style, written with the tact, simplicity, and obvious moral bearing of Miss Edgeworth's stories for the young, but animated by a higher purpose than the Irishwoman's, and dashed with that romance of travel which creates an interest that survives the taste for every other kind of fictitious narrative."—Britannia.

MARRYAT.—THE SETTLERS IN CANADA.

Written for Young People. By Captain Marryat, C.B. author of "Peter Simple," "Masterman Ready," etc. 2 vols. fcap. 8vo. 12s. cloth.

MARRYAT.—MASTERMAN READY;

Or, the Wreck of the Pacific. Written for Young People. By Captain Marryat. 3 vols. foolscap 8vo. with numerous Engravings on Wood, 22s. 6d. cloth.

. The volumes, separately, 7s. 6d. each, cloth.

"The best of Robinson Crusoe's numerous descendants, and one of the most captivating of modern children's books. The only danger is, lest parents should dispute with their children the possession of it."—Quarterly Review.

MARX AND WILLIS.—ON THE DECREASE OF DISEASE EFFECTED BY THE PROGRESS OF CIVILIZATION.

By C. F. H. Marx, M.D. Professor of Medicine in the University of Gottingen, etc.; and R. Willis, M.D. Member of the Royal College of Physicians, etc. Foolscap 8vo. 4s. cloth.

MAUNDER.—THE TREASURY OF KNOWLEDGE,

And LIBRARY OF REFERENCE. By Samuel Maunder. 16th Edition, revised throughout and enlarged. 1 thick vol. foolscap 8vo., with two engraved Frontispieces, 10s. cloth; bound in roan, 12s.

. The principal contents of this new and thoroughly revised edition of "The Treasury of Knowledge," are—a new and enlarged English Dictionary, with a Grammar, Verbal Distinctions, and Exercises; a new Universal Gazetteer; a compendious Classical Dictionary; an Analysis of History and Chronology; a Dictionary of Law Terms; a new Synopsis of the British Peerage; and various useful tabular addenda.

MAUNDER.—THE BIOGRAPHICAL TREASURY:

Consisting of Memoirs, Sketches, and brief Notices of above 12,000 Eminent Persons of all Ages and Nations, from the Earliest Period of History; forming a new and complete Dictionary of Universal Biography. 5th Edition, revised throughout, and containing a copious Supplement, brought down to December 1844. Foolsap 8vo. with engraved Frontispiece, 10s. cloth; bound in roan, 12s.

MAUNDER.—THE SCIENTIFIC AND LITERARY TREASURY:

A New and Popular Encyclopædia of Science and the Belles Lettres; including all Branches of Science, and every Subject connected with Literature and Art. The whole written in a familiar style, adapted to the comprehension of all persons desirous of acquiring information on the subjects comprised in the work, and also adapted for a Manual of convenient Reference to the more instructed. By Samuel Maunder. 3d Edition. 1 thick vol. fcap. 8vo. with an engraved Frontispiece, 10s. cloth; bound in roan, 12s.

MAUNDER.—THE TREASURY OF HISTORY;

Comprising a General Introductory Outline of Universal History, Ancient and Modern, and a Series of separate Histories of every principal Nation that exists; developing their Rise, Progress, and Present Condition, the Moral and Social Character of their respective Inhabitants, their Religion, Manners, and Customs, etc. etc. By Samuel Maunder. 2d Edit. 1 thick vol. fcap. 8vo. 10s. cloth; bound in roan, 12s.

MAUNDER.—THE UNIVERSAL CLASS-BOOK:

A new Series of Reading Lessons (original and selected) for Every Day in the Year; each Lesson recording some important Event in General History, Biography, etc., which happened on the day of the month under which it is placed, or detailing, in familiar language, interesting facts in Science; also a variety of Descriptive and Narrative Pieces, interspersed with Poetical Gleanings: Questions for Examination being appended to each day's Lesson, and the whole carefully adapted to Practical Tuition. By Samuel Maunder, author of "The Treasury of Knowledge." 2d Edition, revised. 12mo. 5s. bound.

MICHELET (J.).—PRIESTS, WOMEN, AND FAMILIES.

By J. Michelet. Translated from the French (third edition), with the Author's permission, by C. Cocks, Bachelier-ès-Lettres, and Professor (breveté) of the Living Languages in the Royal Colleges of France. Post 8vo. 9s. cloth.

"A book uniting many excellences; the interest of the memoir, the fervency of a theological enquiry, and the pungency and force of a dissection of human nature. We recommend it most earnestly to our readers, as not only powerful and profound, but as written so clearly and agreeably that the most volatile and inattentive will comprehend and enjoy the remarkable disclosures made in its pages."—Jerrold's Magazine.

MILNER (REVS. J. AND I.)—THE HISTORY OF THE CHURCH OF

CHRIST. By the Rev. Joseph Milner, A.M. With Additions and Corrections by the late Rev. Isaac Milner, D.D. F.R.S., Dean of Carlisle, and President of Queen's College, Cambridge. A New Edition. 4 vols. 8vo. 2l. 8s. boards.

MONTGOMERY'S (JAMES) POETICAL WORKS.

New and only complete Edition. With some additional Poems and Autobiographical Prefaces. Collected and edited by Mr. Montgomery. 4 vols. foolscap 8vo. with Portrait, and 7 other beautifully engraved Plates, 20s. cloth; or bound in morocco, 1l. 16s.

MOORE'S POETICAL WORKS;

Containing the Author's recent Introduction and Notes. Complete in one volume, uniform with Lord Byron's Poems. With a New Portrait, by George Richmond, engraved in the line manner, and a View of Sloperton Cottage, the Residence of the Poet, by Thomas Creswick, A.R.A. Medium 8vo. 1l. 1s. cloth; or 42s. bound in morocco, in the best manner, by Hayday.

* Also, an Edition in 10 vols. foolscap 8vo. with Portrait, and 19 Plates, 2l. 10s. cloth; morocco, 4l. 10s.

MOORE'S LALLA ROOKH.

Twentieth Edition. Medium 8vo. illustrated with 13 Engravings, finished in the highest style of art, 2ls. cloth; morocco, 35s.; or 42s. with India Proof Plates, cloth.

MOORE'S LALLA ROOKH.

Twenty-first Edition. Foolsap 8vo. with 4 Engravings, from Paintings by Westall, 10s. 6d. cloth; or 14s. bound in morocco.

MOORE'S IRISH MELODIES.

Illustrated by D. MacIlae, R.A. Imp. 8vo. with 160 Designs, engraved on Steel, 3l. 3s. boards; proof Impressions, 6l. 6s. bound. [In October.]

* * * This work has been some years in preparation, and will be ready for publication in October. The text, with an ornamental border to each page, as well as the other Designs, are all engraved on steel; and it is believed that the novelty of the mode of production, combined with the care bestowed in the execution of every part of this elaborate work, will render it one of the most interesting volumes that have ever appeared.

MOORE'S IRISH MELODIES.

Fifteenth Edition. Fcap. 8vo. with Engraved Title and Vignette, 10s. cloth; or 13s. 6d. bound in morocco.

MOORE.—THE HISTORY OF IRELAND.

By Thomas Moore, Esq. Vols. 1 to 3, with Vignette Titles, 18s. cloth.

* * The third and concluding volume, which also completes "The Cabinet Cyclopædia," is nearly ready.

MORAL OF FLOWERS.

3d Edition. Royal 8vo. with 24 beautifully coloured Engravings, 1l. 10s. half-bound.

MORTON.—A VETERINARY TOXICOLOGICAL CHART,

Containing those Agents known to cause Death in the Horse; with the Symptoms, Antidotes, Action on the Tissues, and Tests. By W. J. T. Morton. 12mo. 6s. in case; 8s. 6d. on rollers.

MORTON.—A MANUAL OF PHARMACY,

For the Student in Veterinary Medicine; containing the Substances employed at the Royal Veterinary College, with an Attempt at their Classification, and the Pharmacopœia of that Institution. By W. J. T. Morton. 3d Edition. 12mo. 10s. cloth.

MOSELEY.—ILLUSTRATIONS OF PRACTICAL MECHANICS.

By the Rev. H. Moseley, M.A., Professor of Natural Philosophy and Astronomy in King's College, London; being the First Volume of the Illustrations of Science by the Professors of King's College. Fcap. 8vo. with numerous Woodcuts, 8s. cloth.

MOSELEY.—THE MECHANICAL PRINCIPLES OF ENGINEERING AND ARCHITECTURE.

By the Rev. H. Moseley, M.A. F.R.S., Professor of Natural Philosophy and Astronomy in King's College, London; and author of "Illustrations of Mechanics," etc. 8vo. with Woodcuts and Diagrams, 1l. 4s. cloth.

MÜLLER.—INTRODUCTION TO A SCIENTIFIC SYSTEM OF MYTHOLOGY.

By C. O. Müller, author of "The History and Antiquities of the Doric Race," etc. Translated from the German by John Leitch. 8vo. uniform with "Müller's Dorians," 12s. cloth.

MURRAY.—ENCYCLOPÆDIA OF GEOGRAPHY;

Comprising a complete Description of the Earth: exhibiting its Relation to the Heavenly Bodies, its Physical Structure, the Natural History of each Country, and the Industry, Commerce, Political Institutions, and Civil and Social State of all Nations. By Hugh Murray, F.R.S.E.: assisted in Astronomy, etc. by Professor Wallace; Geology, etc. by Professor Jameson; Botany, etc. by Sir W. J. Hooker; Zoology, etc. by W. Swainson, Esq. New Edition, with Supplement, bringing down the Statistical Information contained in the Work, to December 1843: with 82 Maps, drawn by Sidney Hall, and upwards of 1000 other Engravings on Wood, from Drawings by Swainson, T. Landseer, Sowerby, Strutt, etc. representing the most remarkable Objects of Nature and Art in every Region of the Globe. 1 very large vol. 8vo. containing upwards of 1500 pages, 3l. cloth.

NEWELL (REV. R. H.)—THE ZOOLOGY OF THE ENGLISH POETS.

Corrected by the Writings of Modern Naturalists. By the Rev. R. H. Newell, Rector of Little Hormead. 8vo. [In October.]

NICOLAS—THE CHRONOLOGY OF HISTORY,

Containing Tables, Calculations, and Statements indispensable for ascertaining the Dates of Historical Events, and of Public and Private Documents, from the Earliest Period to the Present Time. By Sir Harris Nicolas, K.C. M.G. Second Edition, corrected throughout. Folscap 8vo. with Vignette Title, 6s. cloth.

NISBET (JAMES).—THE FRENCH IN RHEINSTADT:

A Romance of the Day. A Friendly Voice from the Avon's Banks to the Nations of Germany, and other Poems. By James Nisbet. Post 8vo. 7s. 6d. cloth.

OWEN.—LECTURES ON THE COMPARATIVE ANATOMY AND PHYSIOLOGY OF THE INVERTEBRATE ANIMALS,

delivered at the Royal College of Surgeons in 1843. By Richard Owen, F.R.S. Hunterian Professor to the College. From Notes taken by William White Cooper, M.R.C.S. and revised by Professor Owen. With Glossary and Index. 8vo. with nearly 140 Illustrations on Wood, 14s. cloth.

•• A Second and concluding Volume, being the Lectures on Vertebrata delivered by Professor Owen during the last session, is preparing for publication.

PARABLES (THE).

The Parables of Our Lord, richly Illuminated with appropriate Borders, printed in Colours, and in Black and Gold; with a Design from one of the early German engravers. Square Folscap 4to., uniform in size with the "Sermon on the Mount," 2ls., in a new and characteristic binding; or 30s. bound in morocco, by Hayday. [In October.]

PARKES.—DOMESTIC DUTIES;

Or, Instructions to Young Married Ladies on the Management of their Households and the Regulation of their Conduct in the various Relations and Duties of Married Life. By Mrs. W. Parkes. 5th Edition. Folscap 8vo. 9s. cloth.

PARNELL.—A TREATISE ON ROADS;

Wherein the Principles on which Roads should be made are explained and illustrated by the Plans, Specifications, and Contracts made use of by Thomas Telford, Esq. on the Holyhead Road. By the Right Hon. Sir Henry Parnell, Bart., Hon. Memb. Inst. Civ. Eng. London. Second Edition, greatly enlarged. 8vo. with 9 large Plates, 1l. 1s. cloth.

PATON (A. A.)—SERVIA, THE YOUNGEST MEMBER OF THE EUROPEAN

FAMILY; or, a Residence in Belgrade, and Travels through the Highlands and Woodlands of the Interior, during the years 1843 and 1844. By Andrew Archibald Paton, Esq. Post 8vo. with portrait and plate, 12s. cloth.

By the same Author.

THE MODERN SYRIANS; or, Native Society in Damascus, Aleppo, and the Mountains of the Druses. Post 8vo. 10s. 6d. cloth.

PEARSON.—AN INTRODUCTION TO PRACTICAL ASTRONOMY:

By the Rev. W. Pearson, LL.D. F.R.S. etc. Rector of South Kilworth, Leicestershire, and Treasurer to the Astronomical Society of London. 2 vols. 4to. with Plates, 7l. 7s. boards.

Vol. 1 contains Tables, recently computed, for facilitating the Reduction of Celestial Observations; and a popular Explanation of their Construction and Use.

Vol. 2 contains Descriptions of the various Instruments that have been usefully employed in determining the Places of the Heavenly Bodies, with an Account of the Methods of Adjusting and Using them.

PERCIVALL.—THE ANATOMY OF THE HORSE:

Embracing the Structure of the Foot. By W. Percivall, M.R.C.S. 8vo. 1l. cloth.

PERCIVALL.—HIPPOPATHOLOGY:

A Systematic Treatise on the Disorders and Lameness of the Horse; with their Modern and most approved Methods of Cure; embracing the Doctrines of the English and French Veterinary Schools. By W. Percivall, M.R.C.S., Veterinary Surgeon in the First Life Guards. Vols. 1, 2, and 3, 8vo.; Vol. 1, 10s. 6d.; vol. 2, 14s.; vol. 3, 14s. boards.

PEREIRA.—A TREATISE ON FOOD AND DIET:

With Observations on the Dietetical Regimen suited for Disordered States of the Digestive Organs; and an Account of the Dietaries of some of the principal Metropolitan and other Establishments for Paupers, Lunatics, Criminals, Children, the Sick, etc. By Jon. Pereira, M.D. F.R.S., author of "Elements of Materia Medica." 8vo. 16s. cloth.

PESCHEL (C. F.)—ELEMENTS OF PHYSICS,

Part I. Ponderable Bodies. By C. F. Peschel, Principal of the Royal Military College, Dresden, etc. etc. Translated from the German, with Notes, by E. West. Foolscap 8vo. with Diagrams and Woodcuts. 7s. 6d. cloth.

An useful and well-digested 'Elementary Treatise on Physics.' Its plan is intermediate between the mere popular enunciation of physical facts, and the rigorous mathematical demonstrations of more scientific writers. Thus it is well calculated to meet the wants of those by whom a sound general knowledge of the elementary principles of natural philosophy is desired. The book contains more matter than is found in many ponderous volumes; the style is throughout neat, close, concise, and perspicuous, and the sense everywhere clearly and even elegantly expressed.—Eclectic Review.

PHILLIPS.—AN ELEMENTARY INTRODUCTION TO MINERALOGY:

Comprising a Notice of the Characters and Elements of Minerals; with Accounts of the Places and Circumstances in which they are found. By William Phillips, F.L.S. M.G.S. etc. 4th Edition, considerably augmented by R. Allan, F.R.S.E. 8vo. numerous Cuts, 12s. cloth.

PHILLIPS.—FIGURES & DESCRIPTIONS OF THE PALÆOZOIC FOSSILS OF

CORNWALL, DEVON, and WEST SOMERSET; observed in the course of the Ordnance Geological Survey of that District. By John Phillips, F.R.S. F.G.S. etc. Published by Order of the Lords Commissioners of H. M. Treasury. 8vo. with 60 Plates, comprising very numerous Figures, 9s. cloth.

PHILLIPS.—A GUIDE TO GEOLOGY.

By John Phillips, F.R.S. G.S. etc. Foolscap 8vo. with Plates, 5s. cloth.

PHILLIPS.—A TREATISE ON GEOLOGY.

By John Phillips, F.R.S. G.S. etc. 2 vols. foolscap 8vo. with Vignette Titles and Woodcuts. 12s. cloth.

PORTER.—A TREATISE ON THE MANUFACTURE OF SILK.

By G. R. Porter, Esq. F.R.S., author of "The Progress of the Nation," etc. Feap. 8vo. with Vignette Title, and 39 Engravings on Wood, 6s. cloth.

PORTER.—A TREATISE ON THE MANUFACTURES OF PORCELAIN AND

GLASS. By G. R. Porter, Esq. F.R.S. Foolscap 8vo. with Vignette Title and 50 Woodcuts, 6s. cloth.

PORTLOCK.—REPORT ON THE GEOLOGY OF THE COUNTY OF

LONDONDERRY, and of Parts of Tyrone and Fermanagh, examined and described under the Authority of the Master-General and Board of Ordnance. By J. E. Portlock, F.R.S. etc. 8vo. with 48 Plates, 24s. cloth.

POSTAN'S (CAPTAIN).—PERSONAL OBSERVATIONS ON SINDH,

The Manners and Customs of its Inhabitants, and its Productive Capabilities: with a Narrative of the Recent Events. By Captain Postans, Bombay Army, late Assistant to the Political Agent, Sindh. 8vo. with Map, col'd Frontispiece, and Illustrations on Wood. 18s. cloth.

POWELL.—THE HISTORY OF NATURAL PHILOSOPHY.

From the Earliest Periods to the Present Time. By Baden Powell, M. A., Savilian Professor of Mathematics in the University of Oxford. Feap. 8vo. Vignette Title, 6s. cloth.

PROCEEDINGS OF THE ZOOLOGICAL SOCIETY OF LONDON.

The last Part published is Part 11 for 1843. 8vo. 6s. cloth.

PYCROFT.—A COURSE OF ENGLISH READING;

Adapted to every Taste and Capacity. With Anecdotes of Men of Genius. By the Rev. James Pycroft, B.A., Trinity College, Oxford, author of "Greek Grammar Practice," "Latin Grammar Practice," etc. Foolscap 8vo., 6s. 6d. cloth.

"This course is admirably adapted to promote a really intellectual study of history, philosophy, and the belles lettres, as distinguished from that mere accumulation of words and dates in the memory which passes for education. We would recommend to every idle and inattentive reader, whether old or young, the author's sound and judicious advice, 'How to remember what we read.'"—John Bull.

QUARTERLY JOURNAL OF THE GEOLOGICAL SOCIETY OF LONDON.

Edited by David Thomas Ansted, M.A. F.R.S., Fellow of Jesus College, Cambridge; Professor of Geology in King's College, London; Vice-Secretary of the Geological Society Nos. 1. and 11. 8vo. 4s. each, sewed. [Published Quarterly.]

RANKE (PROFESSOR).—RANKE'S HISTORY OF THE REFORMATION.

Translated by Sarah Austin, translator of Ranke's "History of the Popes." Vols. 1 and 2, 8vo. 30s. cloth.

REECE.—THE MEDICAL GUIDE:

For the use of the Clergy, Heads of Families, Seminaries, and Junior Practitioners in Medicine; comprising a complete Modern Dispensatory, and a Practical Treatise on the distinguishing Symptoms, Causes, Prevention, Cure, and Palliation of the Diseases incident to the Human Frame. By R. Reece, M.D., late Fellow of the Royal College of Surgeons of London, etc. 16th Edition. 8vo. 12s. boards.

REID (DR.).—ILLUSTRATIONS OF THE THEORY AND PRACTICE OF VENTILATION:

with Remarks on Warming, Exclusive Lighting, and the Communication of Sound. By D. B. Reid, M.D. F.R.S.E. etc. 8vo. with Diagrams, and 320 Engravings on Wood, 16s. cloth.

"A complete development of the theory and practice of ventilation, made known to the public for the first time. There is not a chapter that does not offer a great number of novel and important suggestions, well worthy of the careful consideration alike of the public and the professions. Dr. Reid's work is, besides, full of curious illustrations; the descriptions and application of the 'principles' being interspersed throughout with a variety of amusing anecdotes bearing upon the general subject."—Morning Chronicle.

REPTON.—THE LANDSCAPE GARDENING & LANDSCAPE ARCHITECTURE

of the late Humphrey Repton, Esq.; being his entire Works on these subjects. New Edition, with an Historical and Scientific Introduction, a systematic Analysis, a Biographical Notice, Notes, and a copious Alphabetical Index. By J. C. Loudon, F.L.S., etc. Originally published in one folio and three quarto volumes, and now comprised in 1 vol. 8vo. illustrated by upwards of 250 Engravings, and Portrait, 30s. cloth; with coloured Plates, 3l. 6s. cloth.

REYNARD THE FOX;

A renowned Apology of the Middle Age. Reproduced in Rhyme. Embellished throughout with Scroll Capitals, in Colours, from Wood-block Letters made expressly for this work, after Designs of the 12th and 13th Centuries. With an Introduction by Samuel Naylor, late of Queen's College, Oxford. Large square 8vo. 18s. vellum cloth.

RIDDLE—A COMPLETE ENGLISH-LATIN AND LATIN-ENGLISH DICTIONARY,

compiled from the best sources, chiefly German. By the Rev. J. E. Riddle, M.A. 4th Edition. 8vo. 31s. 6d. cloth.

*. Separately—The English-Latin part, 10s. 6d. cloth; the Latin-English part, 21s. cloth.

RIDDLE.—A DIAMOND LATIN ENGLISH DICTIONARY.

For the waistcoat-pocket. A Guide to the Meaning, Quality, and right Accentuation of Latin Classical Words. By the Rev. J. E. Riddle, M.A. Royal 32mo. 4s. bound.

RIDDLE.—LETTERS FROM AN ABSENT GODFATHER;

Or, a Compendium of Religious Instruction for Young Persons. By the Rev. J. E. Riddle, M.A. Foolscep 8vo. 6s. cloth.

RIDDLE.—ECCLESIASTICAL CHRONOLOGY;

Or, Annals of the Christian Church, from its Foundation to the present Time. Containing a View of General Church History, and the Course of Secular Events; the Limits of the Church and its Relations to the State; Controversies; Sects and Parties; Rites, Institutions, and Discipline; Ecclesiastical Writers. The whole arranged according to the order of Dates, and divided into Seven Periods. To which are added, Lists of Councils and Popes, Patriarchs, and Archbishops of Canterbury. By the Rev. J. E. Riddle, M.A. 8vo. 15s. cloth.

RITCHIE (ROBERT).—RAILWAYS: THEIR RISE AND PROGRESS, AND

CONSTRUCTION, with Remarks on Railway Accidents, and Proposals for their prosecution. By Robert Ritchie, Esq. Fcap. 8vo. [In October.]

RIVERS.—THE ROSE AMATEUR'S GUIDE:

Containing ample Descriptions of all the fine leading varieties of Roses, regularly classed in their respective Families; their History and mode of Culture. By T. Rivers, Jun. Third Edition, corrected and improved. Foolscep 8vo. 6s. cloth.

ROBERTS (GEORGE).—THE LIFE, PROGRESSES, AND REBELLION OF

JAMES DUKE OF MONMOUTH, to his Capture and Execution: with a full Account of the Bloody Assizes, and copious Biographical Notices. By George Roberts, author of "The History of Lyme Regis," etc. etc. 2 vols. post 8vo. with Portrait, Maps, and other Illustrations, 24s. cloth.

ROBERTS.—A COMPREHENSIVE VIEW OF THE CULTURE OF THE VINE

under GLASS. By James Roberts, Gardener to M. Wilson, Esq., Eshton Hall, Yorkshire. 12mo. 5s. 6d. cloth.

ROBERTS.—AN ETYMOLOGICAL AND EXPLANATORY DICTIONARY OF the TERMS and LANGUAGE of GEOLOGY; designed for the early Student, and those who have not made great progress in the Science. By G. Roberts. Foolscap 8vo. 6s. cloth.

ROBINSON—GREEK AND ENGLISH LEXICON TO THE NEW TESTAMENT. By E. Robinson, D.D., author of "Biblical Researches." Edited, with careful revision, corrections, etc., by the Rev. Dr. Bloomfield. 8vo. 18s. cloth.

ROGERS.—THE VEGETABLE CULTIVATOR;

Containing a plain and accurate Description of all the different Species of Culinary Vegetables, with the most approved Method of Cultivating them by Natural and Artificial Means, and the best Modes of Cooking them; alphabetically arranged. Together with a Description of the Physical Herbs in General Use. Also, some Recollections of the Life of Philip Miller, F.A.S., Gardener to the Worshipful Company of Apothecaries at Chelsea. By John Rogers, author of "The Fruit Cultivator." 2d Edition. Foolscap 8vo. 7s. cloth.

ROME.—THE HISTORY OF ROME, IN LARDNER'S CYCLOPÆDIA. 2 vols. foolscap 8vo. with Vignette Titles, 12s. cloth.

ROSCOE.—LIVES OF EMINENT BRITISH LAWYERS.

By Henry Roscoe, Esq. Foolscap 8vo. with Vignette Title, 6s. cloth.

SANDBY (REV. G.)—MESMERISM AND ITS OPPONENTS:

With a Narrative of Cases. By the Rev. George Sandby, Jun., Vicar of Flixton, and Rector of All Saints with St. Nicholas, South Elmham, Suffolk; Domestic Chaplain to the Right Hon. the Earl of Abergavenny. Foolscap 8vo. 6s. cloth.

SANDFORD (REV. JOHN).—PAROCHIALIA,

or Church, School, and Parish. By the Rev. John Sandford, M.A. Vicar of Dunchurch, Chaplain to the Lord Bishop of Worcester, Hon. Canon of Worcester, and Rural Dean. 8vo. with numerous Woodcuts. 16s. cloth.

"The title of Mr. Sandford's volume expresses what it is meant to be—a manual for the clergyman in making the most suitable arrangements for the worship of God, and the education of the young, and the supervision of all; and the book is what its title imports. It is a book which we should advise every clergyman to consult, who wishes for full information on any of these points. And we can also recommend it as containing practical information, the result of experience and going into all the necessary details, and accompanied with plans and illustrations, and tables and estimates."—Church of England Quarterly Review.

SANDFORD.—WOMAN IN HER SOCIAL AND DOMESTIC CHARACTER.

By Mrs. John Sandford. 6th Edition. Foolscap 8vo. 6s. cloth.

SANDFORD.—FEMALE IMPROVEMENT.

By Mrs. John Sandford. 2d Edition. Foolscap 8vo. 7s. 6d. cloth.

SANDHURST COLLEGE MATHEMATICAL COURSE.

ELEMENTS of ARITHMETIC and ALGEBRA. By W. Scott, Esq., A.M. and F.R.A.S. Second Mathematical Professor at the Royal Military College, Sandhurst. Being the 1st Volume of the Sandhurst Course of Mathematics. 8vo. 16s. bound.

ELEMENTS of GEOMETRY; consisting of the first Four and Sixth Books of Euclid, chiefly from the Text of Dr. Robert Simson; with the principal Theorems in Proportion, and a Course of Practical Geometry on the Ground; also, Four Tracts relating to Circles, Planes, and Solids, with one on Spherical Geometry. By John Narrien, Professor of Mathematics in the Royal Military College, Sandhurst. Being the 2d Volume of the Sandhurst Course of Mathematics. 8vo. with many Diagrams, 10s. 6d. bound.

PLAIN TRIGONOMETRY AND MENSURATION; for the use of the Royal Military College, Sandhurst. By W. Scott, Esq. A.M. and F.R.A.S., Second Mathematical Master in the Institution. Being the 3d Volume of the Sandhurst Course of Mathematics. 8vo. 9s. 6d. bound.

PRACTICAL ASTRONOMY and GEODESY; Including the Projections of the Sphere, and Spherical Trigonometry. By John Narrien, F.R.S. and R.A.S. Professor of Mathematics in the Royal Military College, Sandhurst. Being the 5th Volume of the Sandhurst Mathematical Course. 8vo. 14s. bound.

SCHLEIDEN (PROF.)—PRINCIPLES OF SCIENTIFIC BOTANY.

By M. J. Schleiden, Professor of Botany at Jena. Translated by E. Lankester, M.D. F.L.S. 8vo. with numerous wood engravings. [In October.]

SCORESBY.—MAGNETICAL INVESTIGATIONS.

By the Rev. William Scoresby, D.D. F.R.S.L. and E. etc. etc. Comprising Investigations concerning the Laws or Principles affecting the Power of Magnetic Steel Plates or Bars, in combination as well as singly, under various conditions as to Mass, Hardness, Quality, Form, etc. as also concerning the comparative Powers of Cast Iron. Part 1, 8vo. with Plates, 5s. cloth; Part 2, 10s. 6d.

SCOTT.—THE HISTORY OF SCOTLAND.

By Sir Walter Scott, Bart. New Edition, 2 vols. foolscap 8vo. with Vignette Titles, 12s. cloth.

SEAWARD.—SIR EDWARD SEAWARD'S NARRATIVE OF HIS SHIPWRECK, and consequent Discovery of certain Islands in the Caribbean Sea: with a Detail of many extraordinary and highly interesting Events in his Life, from 1733 to 1749, as written in his own Diary. Edited by Miss Jane Porter. 3d Edition, with a New Nautical and Geographical Introduction, containing Extracts from a Paper by Mr C. F. Collett, of the Royal Navy, identifying the Islands described by Sir Edward Seaward. 2 vols. post 8vo. 21s. cloth.

SELECT WORKS OF THE BRITISH POETS:

From Chaucer to Withers. With Biographical Sketches, by R. Southey, LL.D. Medium 8vo. 30s. cloth; or 31s. 6d. with gilt edges.

SELECT WORKS OF THE BRITISH POETS:

With Biographical and Critical Prefaces by Dr. Alkin.

A New Edition, with Supplement by Lucy Aikin, consisting of Selections from the Works of Crabbe, Scott, Coleridge, Pringle, Charlotte Smith, and Mrs. Barbauld. Medium 8vo. 18s. cl.

* * *The peculiar feature of these two works is, that the Poems included are printed entire, without mutilation or abridgment; a feature not possessed by any similar work, and adding obviously to their interest and utility.*

SERMON ON THE MOUNT (THE).

[St. Matthew v. vi. vii.] Intended for a Birthday-Present, or Gift-Book for all Seasons. Printed in Gold and Colours, in the Missal Style, with Ornamental Borders by Owen Jones, Architect, and an Illuminated Frontispiece by W. Boxall, Esq. A new edition. Foolscap 4to. in a rich brocaded silk cover, manufactured expressly, 21s.; or bound in morocco, in the Missal style, by Hayday, 26s.

SHAKESPEARE, BY BOWDLER.

THE FAMILY SHAKESPEARE, in which nothing is added to the Original Text; but those Words and Expressions are omitted which cannot with propriety be read aloud. By T. Bowdler, Esq. F.R.S. Seventh Edition, 1 large vol. 8vo. with 36 Illustrations after Smirke, etc. 30s. cloth; or 31s. 6d. gilt edges.

* * * A LIBRARY EDITION, without Illustrations, 8 vols. 8vo. 4l. 14s. 6d. boards.

SHELLEY, ETC.—LIVES OF THE MOST EMINENT LITERARY MEN OF ITALY, SPAIN and PORTUGAL. By Mrs. Shelley, Sir D. Brewster, J. Montgomery, etc. 3 vols. foolscap 8vo. with Vignette Titles, 18s. cloth.

SHELLEY.—LIVES OF THE MOST EMINENT FRENCH WRITERS.

By Mrs. Shelley and others. 2 vols. foolscap 8vo. with Vignette Titles, 12s. cloth.

SHORT WHIST:

Its Rise, Progress, and Laws; with Observations to make any one a Whist Player; containing also the Laws of Piquet, Cassino, Ecarte, Cribbage, Backgammon. By Major A * * * * *. 9th Edition. To which are added, Precepts for Tyros. By Mrs. B * * * * *. Foolscap 8vo. 3s. cloth, gilt edges.

SISMONDI.—THE HISTORY OF THE ITALIAN REPUBLICS;

Or, of the Origin, Progress, and Fall of Freedom in Italy, from A.D. 476 to 1805. By J. C. L. De Sismondi. Foolscap 8vo. with Vignette Title, 6s. cloth.

SISMONDI.—THE HISTORY OF THE FALL OF THE ROMAN EMPIRE.

Comprising a View of the Invasion and Settlement of the Barbarians. By J. C. L. De Sismondi. 2 vols. foolscap 8vo. with Vignette Titles, 12s. cloth.

SMITH (MRS. H.)—THE FEMALE DISCIPLE OF THE FIRST THREE CENTURIES OF THE CHRISTIAN ERA: Her Trials and Her Mission. By Mrs. Henry Smith. Foolscap 8vo. 6s. cloth.

"Mrs. Smith's little book possesses the rare merit of presenting a subject of general interest, which nevertheless has hitherto excited but little attention out of the schools, in an attractive shape, and of concisely embodying the subject-matter of many volumes of patristic writings, which in their original form could never be consulted by the great majority of readers."

Atlas.

SMITH.—AN INTRODUCTION TO THE STUDY OF BOTANY.

By Sir J. E. Smith, late President of the Linnæan Society. 7th Edition, corrected; in which the object of Smith's "Grammar of Botany" is combined with that of the "Introduction." By Sir William Jackson Hooker, K.H. LL.D. etc. 8vo. with 36 Steel Plates, 16s. cloth; with coloured Plates, 2l. 12s. 6d. cloth.

SMITH.—COMPENDIUM OF THE ENGLISH FLORA.

By Sir J. E. Smith. 2d Edition, with Additions and Corrections. By Sir W. J. Hooker. 12mo. 7s. 6d. cloth. THE SAME IN LATIN. 5th Edition, 12mo. 7s. 6d.

SMITH.—THE ENGLISH FLORA.

By Sir James Edward Smith, M.D. F.R.S., late President of the Linnæan Society, etc. 6 vols. 8vo. 3l. 12s. boards.

CONTENTS:

Vols. I. to IV. the Flowering Plants and the Ferns, 2l. 8s.
Vol. V. Part 1, 12s.—Cryptogamia; comprising the Mosses, Hepaticæ, Lichens, Characeæ, and Algae. By Sir W. J. Hooker. | Vol. V. Part 2, 12s.—The Fungi—completing the work, by Sir W. J. Hooker, and the Rev. M. J. Berkeley, F.L.S. etc.

SMITH.—THE WORKS OF THE REV. SYDNEY SMITH.

3d Edition. 3 vols. 8vo. with Portrait, 36s. cloth.

* * *This collection consists of the Author's contributions to the "Edinburgh Review," Peter Plymley's "Letters on the Catholics," and other miscellaneous works; to which are now first added—"Three Letters on Railways;" "Letter to Mr. Horner;" "Two Letters on American Debts;" "A Prayer;" "Changes" (never before published); "A Fragment on the Irish Roman Catholic Church."*

SOUTHEY, ETC.—LIVES OF THE BRITISH ADMIRALS;

With an Introductory View of the Naval History of England. By R. Southey, Esq. and R. Bell, Esq. 5 vols. foolscap 8vo., with Vignette Titles, 1*l.* 10*s.* cloth.

SOUTHEY'S (ROBERT) COMPLETE POETICAL WORKS:

Containing all the Author's last Introductions and Notes. Complete in one volume, medium 8vo. with Portrait and Vignette, uniform with Byron's Poems and Thomas Moore's Poetical Works, 2*l.*s.; or 4*2s.* bound in morocco, in the best manner, by Hayday.

"The present (collective) edition, consisting of one volume only, includes the contents of the former ten, autobiographical prefaces, as well as poems. It is got up in an exceedingly beautiful style, with a clear though small type, and is adorned with a portrait of the author, and a vignette engraving of his residence at Keswick. Altogether, it forms a handsome drawing-room, or library book, whilst its reduced price, as compared with the ten volume edition, will render it highly acceptable to a large class. No lover of elegant literature will now content himself without possessing a copy of works which, however various the opinions entertained respecting some of them, have long taken their place amongst the enduring productions of our age."—*Eclectic Review*.

Also, an Edition in 10 vols. foolscap 8vo. with Portrait and 19 Plates, 2*l.* 10*s.*; morocco, 4*l.* 10*s.*

The following Works separately:—

JOAN OF ARC - - -	Fcap. 8vo. 5 <i>s.</i> cloth.	THALABA - - -	Fcap. 8vo. 5 <i>s.</i> cloth.
MADOC - - - - -	" 5 <i>s.</i> "	BALLADS, etc. 2 vols.	" 10 <i>s.</i> "
CURSE OF KEHAMA - -	" 5 <i>s.</i> "	RODERICK - - - -	" 5 <i>s.</i> "

SPALDING—THE PHILOSOPHY OF CHRISTIAN MORALS.

By Samuel Spalding, M.A. of the London University. 8vo. 10*s.* 6*d.* cloth.

SPIRIT OF THE WOODS.

By the author of "The Moral of Flowers." 2d Edition. Royal 8vo. with 23 beautifully coloured Engravings of the Forest Trees of Great Britain, 1*l.* 11*s.* 6*d.* cloth.

SPOONER.—A TREATISE ON THE STRUCTURE, FUNCTIONS, AND

DISEASES of the FOOT and LEG of the HORSE; comprehending the Comparative Anatomy of these Parts in other Animals; embracing the subject of Shoeing and the proper Treatment of the Foot; with the Rationale and Effects of various Important Operations, and the best Methods of performing them. By W. C. Spooner, M.R.V.C. 12mo. 7*s.* 6*d.* cloth.

STABLE TALK AND TABLE TALK; OR, SPECTACLES FOR YOUNG

SPORTSMEN. By Harry Hieover. 8vo.

[In October.]

STEAM ENGINE, BY THE ARTIZAN CLUB.

A Treatise on the Steam Engine. By the Artizan Club. Nos. 1 to 16, 4to. 1*s.* each, sewed. To be completed in 24 Monthly Parts, each illustrated by a Steel Plate and several Woodcuts.

"The object of this elaborate work is to supply practical machinists and engineers, with a complete and easily-accessible treatise on the steam engine. The labour expended upon it is clearly very great. It is published under the highest auspices, and cannot fail to become the standard authority on the subject; not a merely popular production, but full, explicit, and scientific."—*Railway Chronicle*.

STEBBING (REV. H.)—THE HISTORY OF THE CHURCH OF CHRIST,

From the Diet of Augsburg, 1530, to the Eighteenth Century; originally designed as a Continuation of Milner's "History of the Church of Christ." By the Rev. Henry Stebbing, D.D. 3 vols. 8vo. 36*s.* cloth.

STEBBING.—THE HISTORY OF THE CHRISTIAN CHURCH,

From its Foundation to A.D. 1492. By the Rev. H. Stebbing, M.A., etc. 2 vols. foolscap 8vo. with Vignette Titles, 12*s.* cloth.

STEBBING.—THE HISTORY OF THE REFORMATION.

By the Rev. H. Stebbing. 2 vols. foolscap 8vo. with Vignette Titles, 12*s.* cloth.

STEEL'S SHIPMASTER'S ASSISTANT,

And OWNER'S MANUAL; containing information necessary for persons connected with Mercantile Affairs; consisting of the Regulation Acts of the Customs for the United Kingdom, and British Possessions abroad; Navigation Laws; Registry Acts; Duties of Customs of the United Kingdom, the British Plantations in America, Canada, and Isle of Man; in the East Indies, Cape of Good Hope, New South Wales, and Van Diemen's Land; Smuggling Acts; Pilotage throughout England and Scotland; Insurances; Commercial Treaties; Dock Charges on Shipping, etc. An entirely new Edition, corrected and revised throughout, and brought down to the present Time. 8vo.

[In the press.]

STEPHENS.—A MANUAL OF BRITISH COLEOPTERA;

Or, BEETLES: containing a Description of all the Species of Beetles hitherto ascertained to inhabit Great Britain and Ireland, etc. With a complete Index of the Genera. By J. F. Stephens, F.L.S., author of "Illustrations of Entomology." Post 8vo. 14*s.* cloth.

STRONG—GREECE AS A KINGDOM:

A Statistical Description of that Country—its Laws, Commerce, Resources, Public Institutions, Army, Navy, etc.—from the Arrival of King Otho, in 1833, down to the present time. From Official Documents and Authentic Sources. By Frederick Strong, Esq., Consul at Athens for the Kingdoms of Bavaria and Hanover. 8vo. 15*s.* cloth.

SUMMERLY (MRS. FELIX).—THE MOTHER'S PRIMER:

A Little Child's First Steps in many Ways. By Mrs. Felix Summerly. Fcap. 8vo. printed in Colours, with a Frontispiece drawn on zinc by William Mulready, R.A. 1*s.* sewed.

SUNDAY LIBRARY:

Containing nearly one hundred Sermons by eminent Divines. With Notes, etc. by the Rev. T. F. Dibdin, D.D. 6 vols. foolscap 8vo. with 6 Portraits, 30s. cloth; neatly half-bound in morocco, with gilt edges, 2l. 12s. 6d.

SWAINSON.—A PRELIMINARY DISCOURSE ON THE STUDY OF NATURAL HISTORY. By W. Swainson, Esq. Foolscap 8vo. 6s. cloth.

A TREATISE ON THE NATURAL HISTORY & CLASSIFICATION OF ANIMALS. By W. Swainson, Esq. Fcap. 8vo. 6s. cloth.

NATURAL HISTORY AND CLASSIFICATION OF QUADRUPEDS. By W. Swainson, Esq. Fcap. 8vo. with Vignette Title, and 176 Woodcuts, 6s. cloth.

NATURAL HISTORY AND CLASSIFICATION OF BIRDS. By W. Swainson, Esq. 2 vols. fcap. 8vo. Vignette Titles, and above 300 Woodcuts, 12s. cloth.

A TREATISE ON MALACOLOGY; Or, the Natural Classification of Shells and Shell-Fish. By W. Swainson, Esq. Fcap. 8vo. with Vignette Title and numerous Woodcuts, 6s. cloth.

NATURAL HISTORY AND CLASSIFICATION OF FISH, AMPHIBIANS, AND REPTILES. By W. Swainson, Esq. 2 vols. fcap. 8vo. with Vignette Titles and Woodcuts 12s. cloth.

HABITS AND INSTINCT OF ANIMALS. By W. Swainson, Esq. Fcap. 8vo. with Vignette and Woodcuts, 6s. cloth.

ANIMALS IN MENAGERIES. By W. Swainson, Esq. Fcap. 8vo. Vignette Title and numerous Woodcuts, 6s. cloth.

HISTORY AND NATURAL ARRANGEMENT OF INSECTS. By W. Swainson, Esq. and W. E. Shuckard, Esq. Fcap. 8vo. with Vignette Title and Woodcuts, 6s. cloth.

A TREATISE ON TAXIDERMY; with the Biography of Zoologists, and Notices of their Works. Fcap. 8vo. with Portrait of the Author, 6s. cloth.

SWITZERLAND.—THE HISTORY OF SWITZERLAND.

Foolscap 8vo. with Vignette Title, 6s. cloth.

TATE.—HORATIUS RESTITUTUS;

Or, the Books of Horace arranged in Chronological Order, according to the Scheme of Dr. Bentley, from the Text of Gesner, corrected and improved. With a Preliminary Dissertation, very much enlarged, on the Chronology of the Works, on the Localities, and on the Life and Character of that Poet. By James Tate, M.A. Second edition, to which is now added, an original Treatise on the Metres of Horace. 8vo. 12s. cloth.

"Mr. Tate's *Horatius Restitutus* should find a place in the library of the mature scholar, of the youthful student, and of the accomplished man of the world."—Quarterly Review.

TATE.—THE CONTINUOUS HISTORY OF THE LIFE AND WRITINGS OF ST. PAUL, on the basis of the Acts; with Intercalary Matter of Sacred Narrative, supplied from the Epistles, and elucidated in occasional Dissertations; with the *Horæ Paulinæ* of Dr. Paley, in a more correct edition, subjoined. By James Tate, M.A., Canon Residentiary of St. Paul's. 8vo. with Map, 13s. cloth.

TAYLER (REV. CHARLES B.)—MARGARET;

Or, the Pearl. By the Rev. Charles B. Tayler, M.A. Rector of St. Peter's, Chester, author of "Lady Mary; or, Not of the World;" "Tractarianism not of God," etc. Foolscap 8vo. 6s. cloth.

"The design of this very pleasing and attractive story is to shew the errors of Tractarianism, and the happy effects that flow from a strict discharge of our duties, religious and moral. The book is written in a style extremely graceful, and is altogether free from that narrow-minded spirit which has but too frequently appeared of late in works of the same description."—Atlas.

TAYLER (REV. CHARLES B.)—LADY MARY; OR, NOT OF THE WORLD.

By the Rev. Charles B. Tayler, Rector of St. Peter's, Chester; author of "Margaret, or the Pearl," etc. Foolscap 8vo. 6s. 6d. cloth.

TAYLER (REV. CHARLES B.)—TRACTARIANISM NOT OF GOD.

Sermons. By the Rev. C. B. Tayler, Rector of St. Peter's, and Evening Lecturer at St. Mary's, Chester; author of "Lady Mary; or, Not of the World," etc. Fcap. 8vo. 6s. cloth.

TAYLER (REV. CHARLES B.)—DORA MELDER:

A Story of Alsace. By Meta Sander. A Translation. Edited by the Rev. C. B. Tayler, author of "Margaret; or, the Pearl," etc. Fcap. 8vo., with 2 Illustrations, 7s. cloth.

TAYLOR.—THE STATESMAN.

By Henry Taylor, Esq., author of "Philip Van Artevelde." 12mo. 6s. 6d. boards.

THACKER.—POCKET COMPENDIUM OF COURSING RULES & BYE-LAWS, For Use in the Field. By Thomas Thacker. 12mo. 1s. 6d. sewed.

THACKER.—THE COURSER'S ANNUAL REMEMBRANCER, AND STUD-BOOK; being an Alphabetical Return of the Running at all the Public Coursing Clubs in England, Ireland, and Scotland, for the Season 1841-42; with the Pedigrees (as far as received) of the Dogs that won, and the Dogs that ran up second for each Prize; also, a Return of all Single Matches run at those Meetings; with a Preliminary Essay on the Decision of Short Courses. By T. Thacker. 8vo. 10s. cloth.

THOMSON.—THE DOMESTIC MANAGEMENT OF THE SICK ROOM,

Necessary, in Aid of Medical Treatment, for the Cure of Diseases. By Anthony Todd Thomson, M.D. F.L.S. etc. 2d Edition. Post 8vo. 10s. 6d. cloth.

THIRLWALL.—THE HISTORY OF GREECE.

By the Right Rev. the Lord Bishop of St. David's. A new Edition, revised; with Notes. Vol. 1. demy 8vo. with two Maps, 12s. cloth. To be completed in 8 volumes. [Vol. 2, is in the press.]
"A history of Greece, written with profound and well-digested learning, free from all party bias, executed on an extensive scale, and with no small measure of enthusiastic love for the subject: finally, at so moderate a price, as to be accessible to most students. This elaborate work will long be a standard of reference. Its characteristic excellences are to be looked for in erudition, sound judgment, wise political remark, and philosophic perspicuity; and there are throughout a flow and grace in the narrative which make the reading pleasing to an English reader. There can be little doubt, considering the general suffrage which has been given in favour of the work alike in England and in Germany, that no history of Greece now exists in German, or in any language, which can be compared with Thirlwall's."—Eclectic Review.

*. * Also, an Edition in 8 vols. fcp. 8vo. with Vignette Titles, 2l. 8s. cloth.

THOMSON'S SEASONS.

Edited by Bolton Corney, Esq. Illustrated with Seventy-seven Designs drawn on Wood by the following Members of the Etching Club:—

J. Bell, Sculptor.	J. C. Horsley.	Frank Stone.	H. J. Townsend,
C. W. Cope.	J. P. Knight.	C. Stonhouse.	T. Webster, A.R.A.
Thomas Creswick.	R. Redgrave, A.R.A.	F. Tayler.	

Engraved by Thompson and other eminent Engravers.

Square crown 8vo. 21s. cloth; in morocco in the best manner by Hayday, 36s.

THOMSON.—AN ELEMENTARY TREATISE ON ALGEBRA,

Theoretical and Practical. By James Thomson, LL.D. Professor of Mathematics in the University of Glasgow. 12mo. 5s. cloth.

THOMSON (JOHN).—TABLES OF INTEREST,

At Three, Four, Four-and-a-half, and Five per Cent., from One Pound to Ten Thousand, and from One to Three Hundred and Sixty-five Days, in a regular progression of Single Days; with Interest at all the above Rates, from One to Twelve Months, and from One to Ten Years. Also, Tables shewing the Exchange on Bills, or Commission on Goods, etc. from One-eighth to Five per Cent.; and Tables shewing the Amount of any Salary, Income, Expense, etc. by the Day, Month, or Year. To which are prefixed, a Table of Discount on Bills at a certain Number of Days or Months; and a Table shewing the exact Number of Days, from any Day throughout the Year, to the 31st of December, the usual Period to which Interest is calculated. By John Thomson, Accountant in Edinburgh. 12mo. 8s. bound.

TOMLINE (BISHOP).—ELEMENTS OF CHRISTIAN THEOLOGY;

Containing Proofs of the Authenticity and Inspiration of the Holy Scriptures; a Summary of the History of the Jews; a Brief Statement of the Contents of the several Books of the Old and New Testaments; a Short Account of the English Translations of the Bible, and of the Liturgy of the Church of England; and a Scriptural Exposition of the Thirty-Nine Articles of Religion. By George Tomline, D.D. F.R.S., Lord Bishop of Winchester. Designed principally for the use of Young Students in Divinity. 14th Edition. With Additional Notes, and a Summary of Ecclesiastical History. By Henry Stebbing, D.D., author of "A History of the Church of Christ, from the Confession of Augsburg," etc. etc. 2 vols. 8vo. 21s. cloth.

TOMLINS.—A POPULAR LAW DICTIONARY;

Familiarly explaining the Terms and Nature of English Law; adapted to the comprehension of Persons not educated for the Legal Profession, and affording Information peculiarly useful to Magistrates, Merchants, Parochial Officers, and others. By Thomas Edlyne Tomlins, Attorney and Solicitor. 1thick vol. post 8vo. 18s. cloth.

The whole work has been revised by a Barrister.

TOOKE.—A HISTORY OF PRICES;

With reference to the Causes of their principal Variations, from 1792 to the Present Time. Preceded by a Sketch of the History of the Corn Trade in the last Two Centuries. By Thomas Tooke, Esq. F.R.S. 2 vols. 8vo. 1l. 16s. cloth.

A continuation of the above.

AN ACCOUNT OF PRICES and of the State of the CIRCULATION in 1838 and 1839; with Remarks on the Corn Laws, and on proposed Alterations in our Banking System. 8vo. 12s. cloth.

TRANSACTIONS OF THE GEOLOGICAL SOCIETY OF LONDON.

Second Series. Vol. 7. Part I, 4to. with coloured maps, 4s. 6d.; Vol. 7. Part 2, 4s.

TRANSACTIONS OF THE ENTOMOLOGICAL SOCIETY.

The last Part published is Part I of Vol. IV. 8vo. with Plates, 6s.

TRANSACTIONS OF THE ZOOLOGICAL SOCIETY OF LONDON.

The last Part published is Part 3, Vol. III. 4to. with Plates, 19s. 6d. coloured, and 12s. plain.

TRANSACTIONS OF THE ROYAL INSTITUTE OF BRITISH ARCHITECTS OF

LONDON: consisting of a Series of Papers on "Antiquities," and "Construction." By R. Willis, M.A. F.R.S. etc.; Ambrose Poynter; Herr Hallmann, of Hanover; Dr. Faraday; Mr. Bracebridge; Herr Beuth, of Berlin; Joseph Gwilt, F.S.A. F.A.S.; Mr. C. H. Smith; Mr. C. Fowler, Hon. Sec.; Mr. W. A. Nicholson, of Lincoln; and Mr. J. P. Papworth. Vol. I. Part 2, 4to. with numerous lithographic and woodcut Illustrations, 24s. cloth.

*. * Part I. Vol. I. uniform with the above, 16s. cloth.

TRANSACTIONS OF THE LINNEAN SOCIETY OF LONDON.

The last Part published is Part 3, Vol. XIX. 4to. with Plates, 40s.

TRANSACTIONS OF THE INSTITUTION OF CIVIL ENGINEERS.

Vol. II. 4to. with Twenty-three finely engraved Plates, 28s. cloth.

Vol. III., with Nineteen finely engraved Plates. 2l. 12s. 6d. cloth.

TURNER.—THE SACRED HISTORY OF THE WORLD.

Philosophically considered. By S. Turner, F.S.A. R.A.S.L. New Edit. 3 vols. 8vo. 42s. bds.
Vol. 1 considers the Creation and System of the Earth, and its Vegetable and Animal Races and Material Laws, and Formation of Mankind.
Vol. 2, the Divine Economy in its special Relation to Mankind, and in the Deluge, and the History of Human Affairs.
Vol. 3, the Provisions for the Perpetuation and Support of the Human Race, the Divine System of our Social Combinations, and the Supernatural History of the World.

TURNER (SHARON).—RICHARD III.:

A Poem. By Sharon Turner, Esq., F.S.A. and R.A.S.L., author of "The History of the Anglo-Saxons," "The Sacred History of the World," etc. Foolsap 8vo. 7s. 6d., cloth.

TURNER.—THE HISTORY OF ENGLAND.

From the Earliest Period to the Death of Elizabeth. By Sharon Turner, Esq. F.A.S. R.A.S.L. 12 vols. 8vo. 8l. 3s. cloth.

Or four separate portions, as follows:—

THE HISTORY of the ANGLO-SAXONS; comprising the History of England from the Earliest Period to the Norman Conquest. 6th Edition, 3 vols. 8vo. 27. 5s. boards.

THE HISTORY of ENGLAND during the MIDDLE AGES; comprising the Reigns from William the Conqueror to the Accession of Henry VIII., and also the History of the Literature, Religion, Poetry, and Progress of the Reformation, and of the Language during that period. 3d Edition, 6 vols. 8vo. 3l. boards.

THE HISTORY of the REIGN of HENRY VIII.; comprising the Political History of the commencement of the English Reformation: being the First Part of the Modern History of England. 3d Edition, 2 vols. 8vo. 26s. boards.

THE HISTORY of the REIGNS of EDWARD VI., MARY, and ELIZABETH; being the Second Part of the Modern History of England. 3d Edition, 2 vols. 8vo. 32s. boards.

TURNER.—A TREATISE ON THE FOOT OF THE HORSE.

And a New System of Shoeing, by One-sided Nailing; and on the Nature, Origin, and Symptoms of the Navicular Joint Lameness, with Preventive and Curative Treatment. By James Turner, M.R.V.C. Royal 8vo. 7s. 6d. boards.

TURTON'S (DR.) MANUAL OF THE LAND AND FRESHWATER SHELLS OF

THE BRITISH ISLANDS. A new Edition, thoroughly revised, and with considerable Additions. By John Edward Gray, Keeper of the Zoological Collection in the British Museum. Post 8vo. with Woodcuts, and 12 coloured Plates, 15s. cloth.

UNCLE PETER.—UNCLE PETER'S FAIRY TALES.

The First Story, containing the History and Adventures of Little Mary, Queen of the Great Island of Brakarakakaka. By Uncle Peter, F.R.L. M.M. T.T. F.A.S. Q.Q. X.Y.Z., etc. Foolsap 8vo. 5s. 6d. cloth.

URE.—DICTIONARY OF ARTS, MANUFACTURES, AND MINES;

Containing a clear Exposition of their Principles and Practice. By Andrew Ure, M.D. F.R.S. M.G.S. M.A.S. Lond.; M. Acad. N.S. Philad.; S. Ph. Soc. N. Germ. Hanov.; Mullii. etc. etc. Third Edition, corrected. 8vo. illustrated with 1240 Woodcuts, 50s. cloth.

URE (DR.)—RECENT IMPROVEMENTS IN ARTS, MANUFACTURES,

AND MINES; being the 2d Edition of a Supplement to the 3d Edition of his "Dictionary." By Andrew Ure, M.D. F.R.S. etc. 8vo. with numerous Wood Engravings, 14s. cloth.

VON ORLICH (CAP.)—TRAVELS IN INDIA,

And the adjacent Countries, in 1842 and 1843. By Capt. Leopold Von Orlich. Translated from the German by H. Evans Lloyd, Esq. 2 vols. 8vo. with coloured Frontispieces, and numerous Illustrations on Wood. 25s. cloth.

WALKER (GEO.)—CHESS STUDIES:

Comprising One Thousand Games actually Played during the last Half Century; presenting a unique Collection of Classical and Brilliant Specimens of Chess Skill in every stage of the Game, and thus forming a complete Encyclopedia of Reference. By George Walker. Medium 8vo. 10s. 6d. sewed.

WATERTON.—ESSAYS ON NATURAL HISTORY,

Chiefly Ornithology. By Charles Waterton, Esq., author of "Wanderings in South America." With an Autobiography of the Author, and a View of Walton Hall. 5th Edition, foolsap 8vo. 8s. cloth.

SECOND SERIES. With Continuation of Mr. Waterton's Autobiography. 2d Edition, fcap.

8vo. with Vignette by T. Creswick, A.R.A. 6s. 6d. cloth

WATTS (A. A.)—LYRICS OF THE HEART,

And other Poems. By Alaric A. Watts. Illustrated by a Series of Engravings from the most celebrated works of modern Painters, executed in the most finished style of Art. Square crown 8vo. 1l. 1s.; proof impressions, 3l. 3s. [In November.

WEBSTER.—AN ENCYCLOPÆDIA OF DOMESTIC ECONOMY;

Comprising such subjects as are most immediately connected with Housekeeping; as, The Construction of Domestic Edifices, with the modes of Warming, Ventilating, and Lighting them—A description of the various articles of Furniture, with the nature of their Materials—Duties of Servants—A general account of the Animal and Vegetable Substances used as Food, and the methods of preserving and preparing them by Cooking—Making Bread—The Chemical Nature and the Preparation of all kinds of Fermented Liquors used as Beverage—Materials employed in Dress and the Toilet—Business of the Laundry—Description of the various Wheel Carriages—Preservation of Health—Domestic Medicine, etc. etc. etc. By Thomas Webster, F.G.S., etc.; assisted by The late Mrs. Parkes, author of "Domestic Duties." 1 thick vol. 8vo. illustrated with nearly 1000 Woodcuts. 50s. cl.

WARDLAW.—DISCOURSES ON THE PRINCIPAL POINTS OF THE SOCINIAN CONTROVERSY—the Unity of God, and the Trinity of Persons in the Godhead—the Supreme Divinity of Jesus Christ—the Doctrine of the Atonement—the Christian Character, etc. By Ralph Wardlaw, D.D. 5th Edition, 8vo. 15s. cloth.

WEIL (DR.).—THE BIBLE, THE KORAN, AND THE TALMUD; Or, Biblical Legends of the Mahometans and Hebrews, from Arabic and Hebrew Sources. By Dr. Weil, of Heidelberg. Translated, with Notes, by the Rev. H. Douglas, A.M. Fcap 8vo. [Just ready.]

WELSFORD (HENRY).—ON THE ORIGIN AND RAMIFICATIONS OF THE ENGLISH LANGUAGE; preceded by an Investigation into the Primitive Seats, Early Migrations, and Final Settlements, of the principal European Nations. By Henry Welsford. 8vo. 10s. 6d. cloth.

WHITE'S COMPENDIUM OF THE VETERINARY ART; Containing Plain and Concise Observations on the Construction and Management of the Stable; a brief and popular Outline of the Structure and Economy of the Horse; the Nature, Symptoms, and Treatment of the Diseases and Accidents to which the Horse is liable: the best Method of performing various important Operations; with Advice to the Purchasers of Horses; and a copious *Materia Medica* and *Pharmacopœia*. 17th edition, entirely reconstructed, with considerable Additions and Alterations, bringing the work up to the present state of Veterinary Science. By W. C. Spooner, 8vo. with coloured Plate, 16s. cloth.

WHITE'S COMPENDIUM OF CATTLE MEDICINE; Or, Practical Observations on the Disorders of Cattle and other Domestic Animals, except the Horse. 6th Edition, re-arranged, with copious Additions and Notes. By W. C. Spooner, Vet. Surgeon, author of a "Treatise on the Influenza," and a "Treatise on the Foot and Leg of the Horse," etc. 8vo. 9s. cloth.

WIGAN (DR. A. L.).—THE DUALITY OF THE MIND, Proved by the Structure, Functions, and Diseases of the Brain, and by the Phenomena of Mental Derangement; and shewn to be essential to Moral Responsibility. With an Appendix. 1. On the Influence of Religion on Insanity; 2. Conjectures on the Nature of the Mental Operations; 3. On the Management of Lunatic Asylums. By A. L. Wigan, M.D. 8vo. 12s. cloth.

WILBERFORCE (W.).—A PRACTICAL VIEW OF THE PREVAILING RELIGIOUS SYSTEMS OF PROFESSED CHRISTIANS, in the Higher and Middle Classes in this Country, contrasted with Real Christianity. By William Wilberforce, Esq. M.P. for the County of York. 17th Edition. 8vo. 8s. boards.
* 19th Edition. 12mo. 4s. boards.

WILKINSON.—THE ENGINES OF WAR, ETC. Being a History of Ancient and Modern Projectile Instruments and Engines of Warfare and Sporting; including the Manufacture of Fire Arms, the History and Manufacture of Gunpowder, of Swords, and of the cause of the Damascus Figure in Sword Blades, with some Observations on Bronze: to which are added, Remarks on some Peculiarities of Iron, and on the extraordinary Effect produced by the Action of Sea Water on Cast Iron; with Details of the various miscellaneous Experiments. By H. Wilkinson, M.R.A.S. 8vo. 9s. cloth.

WILLIS (N. P.).—DASHES AT LIFE WITH A FREE PENCIL. By N. P. Willis, Esq., author of "Pencilings by the Way," "Inklings of Adventure," etc. 8 vols. post 8vo. 31s. 6d. boards.

"An exceedingly amusing book, dashed off with the freest of pencils."—Bell's Messenger.

WILLOUGHBY (LADY).—A DIARY. Purporting to be by the LADY WILLOUGHBY of the Reign of Charles I., embracing some Passages of her Domestic History from 1635 to 1648. 3d edition. Square foolscap 8vo. 8s. boards, or 18s. bound in morocco (old style.)

* This volume is printed and bound in the style of the period to which the Diary refers.


"The great charm of the book, which makes it almost impossible to lay it aside until wholly perused, is its beautiful simplicity, united to the most touching pathos, ever and anon relieved by little notices of household cares, and sweet pictures of domestic felicity."


Scotsman.

ZUMPT (PROF.).—A GRAMMAR OF THE LATIN LANGUAGE.

By C. G. Zumpt, Ph. D. Professor in the University, and Member of the Royal Academy of Berlin. Translated from the 9th Edition of the original, and adapted to the use of English Students, by Leonhard Schmitz, Ph. D., late of the University of Bonn; with numerous Additions and Corrections by the Author. 8vo. 14s. cloth.

"It would seem by a preface to this translation, from the pen of Professor Zumpt himself, that the translation so long in use in our English schools and colleges has not been satisfactory to the learned author. He describes it as imperfect, inaccurate, and translated from an early edition: 'while the German original, by continued labour on my part, had in its details become quite a different work.' To the new translation, here published, he has given every assistance. Though the edition from which it is translated has appeared within a year, his unceasing philological labours had already collected a large number of corrections and additions for future use; these he has sent to Dr. Schmitz for incorporation in the English text, and they make the volume before us more perfect than the German original. The Professor expresses the greatest confidence in his translator's learning and accuracy; and intimates that it will henceforth be their 'united endeavour' to remedy what deficiencies may be found. Thus, beyond all question, is the work of Dr. Schmitz henceforward the authorized version of Zumpt's Grammar: a book which well deserves its great celebrity and the high esteem by which it is held by the best scholars."





BOUND BY
WESTLEYS &
CLARK.

